

The proton radius: From a puzzle to precision

Hans-Werner Hammer^{a,b}, Ulf-G. Meißner^{c,d,e}

^a*Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany*

^b*ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany*

^c*Helmholtz Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

^d*Institute for Advanced Simulation and Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany*

^e*Tbilisi State University, 0186 Tbilisi, Georgia*

Abstract

We comment on the status and history of the proton charge radius determinations.

The proton charge radius r_p is a fundamental quantity in particle physics, as it challenges our understanding of the so successful Standard Model in the non-perturbative regime of the strong interactions. It is defined by the slope of the proton charge form factor $G_p^E(t)$ at zero momentum transfer,

$$r_p^2 = 6 \frac{dG_p^E(t)}{dt} \Big|_{t=0}, \quad (1)$$

with t the invariant four-momentum transfer squared. The proton charge radius was first indirectly measured in the Nobel prize winning electron scattering experiments by Hofstadter et al. [1, 2], who fitted the form factor data with a dipole form and extracted the radius from the slope of the dipole. While electron scattering was the method of choice to refine the measurements of the proton radius in the decades following these pioneering experiments, the Lamb shift in electronic hydrogen and muonic hydrogen is also sensitive to the proton radius [3]. These are electromagnetic bound states of an electron (a muon) with a proton, where the finite size of the proton leaves a small imprint in the energy spectrum. Such determinations, however, require precision experiments and precision theory and thus came into the game only later, with a much higher sensitivity to r_p for muonic hydrogen. This is due to the larger muon mass, $m_\mu/m_e \simeq 200$, so that the corresponding Bohr radius is smaller and the effect of the proton radius much enhanced. Most electron scattering experiments gave the so-called **large** radius, $r_p \simeq 0.88$ fm, which was also the value given by CODATA [4]. This was also consistent with the determination from the electronic Lamb shift. It then came as a true surprise to most researchers (but not all, see below) when the first measurement of the muonic hydrogen Lamb shift led to

the so-called **small** radius, $r_p = 0.84184(67)$ fm, differing by 5σ from the CODATA value [5]. The plot further thickened when high-precision electron-proton scattering data from the Mainz Microtron (MAMI) reinforced the large radius [6], which was also consistent with the average value from electronic Lamb shift measurements, see, e.g., [4]. Another measurement of muonic hydrogen, however, supported the small value [7]. This glaring discrepancy in such a fundamental quantity, which was believed to be understood since long, became known as the “proton radius puzzle”, that featured prominently in many print and online media. For a review see, e.g., [8].

However, while this led to a large number of publications scrutinizing the experimental and theoretical approaches, or even questioning the lepton universality underlying the Standard Model (cf. Ref. [8]), this is not the whole story. Electron scattering data leading to the proton and neutron charge and magnetic form factors are best analyzed using dispersion relations, as these embody the general principles of unitarity, crossing and analyticity. In particular, the contribution from the closest singularity in the momentum transfer t , the two-pion continuum, can be included in a model-independent fashion and is of utmost importance for a proper extraction of the proton radius [9]. This approach was pioneered and utilized first by the Karlsruhe group [10], and further developed and made consistent with symmetries and constraints from Quantum Chromodynamics (QCD) by the Bonn-Mainz group [11, 12]. In fact, the proton radius puzzle was anticipated in the 2007 paper by Belushkin and the authors based on a thorough dispersion-theoretical analysis of the world data base of nucleon form factor data [13]. Again,

Table 1: Modern extractions of the proton charge radius from the electronic Lamb shift and electron-proton scattering.

r_p [fm]	year	method	Ref.
0.8335(95)	2017	el. Lamb shift	[16]
0.877(13)	2018	el. Lamb shift	[17]
0.833(10)	2019	el. Lamb shift	[18]
0.831(7)(12)	2019	$e - p$ scattering	[19]

a small radius in the range $r_p = 0.82 \dots 0.85$ fm was found and it was further shown that a large proton radius $r_p = 0.88 \dots 0.90$ fm could hardly be accommodated by the form factor data if the constraints from unitarity and analyticity are taken into account. The reanalysis of the exquisite MAMI data from 2010 using the same dispersion-theoretical framework also led to a small radius of $r_p = 0.84(1)$ fm [14], similar to all other such analyses before. This was later refined including effects from the two-photon exchange and performing an improved error analysis, leading to $r_p = 0.840$ (0.828 – 0.855) fm [15].

Still, the situation remained unsatisfactory as r_p from the electronic Lamb shift was on the large side and there has been on-going debate about the extraction of the radius from electron scattering experiments. The situation changed, however, dramatically when three new experiments on the electronic Lamb shift [16, 17, 18], a novel measurement of electron-proton scattering at unprecedented small momentum transfer [19], and another dispersion-theoretical inspired analysis of electron scattering data [20] became available in the last few years, with the latter one just reinforcing the claims made by the Bonn-Mainz group since the mid 1990ties. With the exception of the Paris electronic Lamb shift measurement [17], all of these new determinations of r_p consistently give a small proton radius. Consequently, the newest addition of the CODATA compilation lists the proton charge radius as $r_p = 0.8414(19)$ fm [21], completely consistent with the value from muonic hydrogen and electron scattering data analysed using dispersion theory. In light of these results, an improved dispersion-theoretical analysis including the most recent high-precision determination of the two-pion continuum contribution [22] based on the tremendously successful Roy-Steiner analysis of pion-nucleon scattering [23] should be performed. Complementary scattering experiments using muons are also planned [24, 25].

In summary, in view of the new extractions of the proton charge radius from electronic Lamb shift measurements and very low-energy electron-proton scatter-

ing as well as the on-going activities to analyze electron scattering data using dispersive methods, we can now consider the proton radius puzzle solved and look forward to an increased precision in the determination of this fundamental quantity.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We thank Hans Ströher for pertinent comments and Bastian Kubis for a careful reading of the manuscript. This work was supported in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 279384907 – CRC 1245 and the DFG and the NSFC through funds provided to the Sino-German CRC 110 “Symmetries and the Emergence of Structure in QCD”. The work of UGM was also supported by the Chinese Academy of Sciences (CAS) President’s International Fellowship Initiative (PIFI) (Grant No. 2018DM0034) and by VolkswagenStiftung (Grant. No. 93562). The work of HWH was also supported by the German Federal Ministry of Education and Research (BMBF) (Grant no. 05P18RDFN1).

References

- [1] R.W. McAllister and R. Hofstadter, Phys. Rev. **102**, 851 (1956).
- [2] R. Hofstadter, Ann. Rev. Nucl. Part. Sci. **7**, 231 (1957).
- [3] R. Karplus, A. Klein and J. Schwinger, Phys. Rev. **86**, 288 (1952).
- [4] P. J. Mohr, B. N. Taylor and D. B. Newell, Rev. Mod. Phys. **80**, 633 (2008) [arXiv:0801.0028 [physics.atom-ph]].
- [5] R. Pohl *et al.*, Nature **466**, 213 (2010).
- [6] J. C. Bernauer *et al.* [A1 Collaboration], Phys. Rev. Lett. **105**, 242001 (2010) [arXiv:1007.5076 [nucl-ex]].
- [7] A. Antognini *et al.*, Science **339**, 417 (2013).
- [8] R. Pohl, R. Gilman, G. A. Miller and K. Pachucki, Ann. Rev. Nucl. Part. Sci. **63**, 175 (2013) [arXiv:1301.0905 [physics.atom-ph]].
- [9] G. Höhler and E. Pietarinen, Phys. Lett. **53B**, 471 (1975).
- [10] G. Höhler, E. Pietarinen, I. Sabba Stefanescu, F. Borkowski, G. G. Simon, V. H. Walther and R. D. Wendling, Nucl. Phys. B **114**, 505 (1976).
- [11] P. Mergell, U.-G. Meißner and D. Drechsel, Nucl. Phys. A **596**, 367 (1996). [hep-ph/9506375].
- [12] H. W. Hammer, U.-G. Meißner and D. Drechsel, Phys. Lett. B **385**, 343 (1996) [hep-ph/9604294].
- [13] M. A. Belushkin, H.-W. Hammer and U.-G. Meißner, Phys. Rev. C **75**, 035202 (2007). [hep-ph/0608337].
- [14] I. T. Lorenz, H.-W. Hammer and U.-G. Meißner, Eur. Phys. J. A **48**, 151 (2012) [arXiv:1205.6628 [hep-ph]].
- [15] I. T. Lorenz, U.-G. Meißner, H.-W. Hammer and Y.-B. Dong, Phys. Rev. D **91**, 014023 (2015) [arXiv:1411.1704 [hep-ph]].
- [16] A. Beyer *et al.*, Science **358**, 79 (2017).

- [17] H. Fleurbaey *et al.*, Phys. Rev. Lett. **120**, 183001 (2018) [arXiv:1801.08816 [physics.atom-ph]].
- [18] N. Bezginov, T. Valdez, M. Horbatsch, A. Marsman, A. C. Vutha and E. A. Hessels, Science **365**, 1007 (2019).
- [19] W. Xiong *et al.*, Nature **575**, 147 (2019).
- [20] J. M. Alarcon, D. W. Higinbotham, C. Weiss and Z. Ye, Phys. Rev. C **99**, 044303 (2019) [arXiv:1809.06373 [hep-ph]].
- [21] <https://physics.nist.gov/cgi-bin/cuu/Value?rp>
- [22] M. Hoferichter, B. Kubis, J. Ruiz de Elvira, H.-W. Hammer and U.-G. Meißner, Eur. Phys. J. A **52**, 331 (2016) [arXiv:1609.06722 [hep-ph]].
- [23] M. Hoferichter, J. Ruiz de Elvira, B. Kubis and U.-G. Meißner, Phys. Rept. **625**, 1 (2016) [arXiv:1510.06039 [hep-ph]].
- [24] R. Gilman *et al.* [MUSE Collaboration], arXiv:1303.2160 [nucl-ex].
- [25] C. Dreisbach *et al.* [COMPASS++/AMBER working group], PoS DIS **2019**, 222 (2019).