

Subspace Power Method for Symmetric Tensor Decomposition and Generalized PCA

Joe Kileel · João M. Pereira

Abstract We introduce the *Subspace Power Method* (SPM) for calculating the CP decomposition of low-rank even-order real symmetric tensors. This algorithm applies the tensor power method of Kolda-Mayo to a certain modified tensor, constructed from a matrix flattening of the original tensor, and then uses deflation steps. Numerical simulations indicate SPM is roughly one order of magnitude faster than state-of-the-art algorithms, while performing robustly for low-rank tensors subjected to additive noise. We obtain rigorous guarantees for SPM regarding convergence and global optima, for tensors of rank up to roughly the square root of the number of tensor entries, by drawing on results from classical algebraic geometry and dynamical systems. In a second contribution, we extend SPM to compute De Lathauwer’s symmetric block term tensor decompositions. As an application of the latter decomposition, we provide a method-of-moments for generalized principal component analysis.

Keywords CP symmetric tensor decomposition · generalized PCA · power method · trisecant lemma · Łojasiewicz inequality · center-stable manifold

1 Introduction

A tensor is a multi-dimensional array [46], and a symmetric tensor is an array unchanged by permutation of indices, i.e., a tensor T of order m is symmetric if for each index (j_1, \dots, j_m) and permutation σ , we have $T_{j_1 \dots j_m} = T_{j_{\sigma(1)} \dots j_{\sigma(m)}}$.

Joe Kileel

Mathematics Department and Oden Institute, University of Texas at Austin
E-mail: jkileel@math.utexas.edu

João M. Pereira

Oden Institute, University of Texas at Austin
E-mail: joao.pereira@utexas.edu

*The authors contributed equally.

Symmetric tensors arise naturally in several data processing tasks as higher order moments, generalizing the mean and covariance of a dataset or random vector; or as derivatives of multivariate real-valued functions, generalizing the gradient and Hessian of a function. Being able to decompose symmetric tensors is important in applications such as blind source separation (BSS) [13, 26], independent component analysis (ICA) [41, 66], antenna array processing [32, 18], telecommunications [76, 17], psychometrics [15], chemometrics [9], magnetic resonance imaging [5] and latent variable estimation for Gaussian mixture models, topic models and hidden Markov models [2].

In this paper, we focus on decomposing real symmetric tensors of even-order. Although our results may be modified to apply to real odd-order tensors (up to a suitable rank, see Remark 3.4), we focus on tensors with order $2n$, to simplify notation. Also, we restrict to real rather than complex tensors.

– **Unless otherwise stated, in this paper $n \geq 2$ is always assumed.**

Our goal is to compute *real symmetric rank decompositions of tensors*:

$$T = \sum_{i=1}^R \lambda_i a_i^{\otimes 2n}. \quad (1.1)$$

Such an expression is also known as a *CP decomposition* of T . In (1.1), the left-hand side T is a real symmetric tensor of dimension $L \times \dots \times L$ ($2n$ times). Meanwhile on the right-hand side, R is an integer, $\lambda_i \in \mathbb{R}$ are scalars, $a_i \in \mathbb{R}^L$ are unit norm vectors, and $a_i^{\otimes 2n}$ denotes the $2n$ -th tensor power (or outer product) of a_i . The tensor power of a vector is given by

$$(a_i^{\otimes 2n})_{j_1 \dots j_{2n}} = (a_i)_{j_1} \dots (a_i)_{j_{2n}}.$$

Symmetric CP tensor decompositions enjoy excellent existence and uniqueness properties, which is one reason for their use in inferential applications. Firstly, given any real symmetric tensor T , there always exists a decomposition of type (1.1). This is because the set $\{a^{\otimes 2n} : \|a\| = 1\}$ spans the vector space of all symmetric tensors [48]. On the other hand, suppose we consider a decomposition (1.1) in which R is minimal (what we shall mean by rank decomposition). The minimal R is called the rank of T . It is a crucial theoretical fact that, generically, rank decompositions are *unique* for rank-deficient tensors. If T satisfies (1.1) with $R < \frac{1}{L} \binom{L+2n-1}{2n} = O(L^{2n-1})$ and $L > 6$, then for Zariski-generic (λ_i, a_i) , the rank of T is R and the minimal decomposition (1.1) is unique (up to permutation and sign flips of a_i). This is a result in algebraic geometry about secant varieties of Veronese varieties [20, 3, 19].

While low-rank symmetric CP rank decompositions are generically unique, computing the decomposition is another matter. Hillar-Lim showed tensor decomposition is NP-hard in general [39]. Nonetheless, a number of works have sought efficient algorithms for sufficiently low-rank tensors, e.g., [37, 26, 2, 45, 35, 54, 40, 60]. Indeed, in theoretical computer science, a picture has emerged that efficient algorithms should exist that succeed in decomposing random tensors of rank on the order of the square root of the number of




Fig. 1.1: *Schematic of the Subspace Power Method.* The input is the symmetric tensor on the left (the low-rank decomposition of T is unknown). The output is $(\lambda_i, a_i)_{i=1}^R$. SPM has three main steps: (A) EXTRACT SUBSPACE; (B) POWER METHOD; and (C) DEFLATE.

tensor entries with high probability (but not so for ranks substantially more). Thus, tensor decomposition has a conjectural computational-statistical gap [4]. Producing a *practically efficient* method (even with restricted rank) is a further challenge.

1.1 Our contributions

In the present paper, we derive an algorithm that accepts T as input, and aims to output the minimal decomposition (1.1), up to trivial ambiguities, provided $R \leq \binom{L+n-1}{n} - L = O(L^n)$. Numerical simulations indicate our algorithm successfully computes the decomposition for $R = O(L^n)$, with no random restarts. Most notably, the method outperforms existing state-of-art methods by roughly one order of magnitude in terms of speed. Also it matches existing methods in terms of robustness to perturbations in the entries of T (including perturbations that give an input only approximately low-rank).

We call the method the *Subspace Power Method* (SPM). To give a glimpse, it consists of three parts.

- A. EXTRACT SUBSPACE: We flatten T to a square matrix, and compute a matrix eigendecomposition. From this, we extract the subspace of order- n tensors spanned by $a_i^{\otimes n}$, denoted \mathcal{A} .
- B. POWER METHOD: We try to find one of the rank-1 points $a_i^{\otimes n}$ in the subspace \mathcal{A} . To this end, we apply SS-HOPM [46] for computing tensor eigenvectors to an appropriately modified tensor, constructed from the subspace \mathcal{A} .
- C. DEFLATION: Finally, we solve for the corresponding scalar λ_i , and remove the obtained (λ_i, a_i) from T .

The pipeline repeats R times, until all (λ_i, a_i) are recovered. Fig. 1.1 shows a schematic of SPM (see Algorithm 1 in Section 3 for details). Step C (deflation) is optimized so as to avoid re-computation of a large eigendecomposition.

Beyond introducing a fast algorithm, we prove various theoretical guarantees for SPM. Firstly, we characterize the *global optima* in our reformulation of tensor decomposition. That is, in Propositions 3.1 and 3.2, we use the trisecant lemma from classical algebraic geometry to show the only rank-1 points

in \mathcal{A} are $a_i^{\otimes n}$ (up to scale). Secondly, we establish *convergence guarantees* for the power method iteration (Step B in Figure 1.1). The analysis is summarized by Theorem 4.1. In particular, using the Łojasiewicz inequality for real analytic functions, we show the power method indeed always converges (i.e., does not merely have accumulation points on the unit sphere \mathbb{S}^{L-1}). This is in fact an important technical contribution: while proving this, we also settle an outstanding conjecture of Kolda-Mayo that their SS-HOPM method for computing eigenvectors of symmetric tensors always converges (see page 1107 of [44]). We also get a qualitative bound on the rate of convergence. Our analysis further yields that the power method converges to a *second-order optimal point* for almost all initializations, and each component vector $\pm a_i$ is an *attractive* fixed point of the power method. Although this stops of justifying our empirical finding that SPM converges in practice always to $\pm a_i$ (the *global* optima), it does give another key explanation for the success of SPM. The technical tool here is the center-stable manifold theorem from dynamical systems.

Our last main contribution is in extending SPM to compute a certain generalization of the standard CP decomposition (1.1), De Lathauwer’s *symmetric block term tensor decomposition* [24, 25, 28]:

$$T = \sum_{i=1}^R (A_i; \dots; A_i) \cdot \mathbf{A}_i. \quad (1.2)$$

Here, T is a real symmetric tensor of size $L^{\times 2n}$, decomposed as a sum of *symmetric Tucker factorizations* (rather than rank-1 tensors). See Definitions 2.5 and 2.6. While decomposition (1.2) includes decomposition (1.1) as a strictly special case, SPM extends gracefully to compute symmetric block term tensor decompositions. We supply the details in Algorithm 2. The global optima characterization and local attractiveness are now based on the algebraic geometry of *subspace arrangements*; see Proposition 4.11 and Lemma 5.6.

To give a concrete application, we propose an SPM-based method-of-moments for *generalized principal component analysis* (GPCA). GPCA is the task of estimating a union of subspaces from noisy samples [79, 55, 80]. In simulations, we demonstrate that SPM for symmetric block term tensor decomposition (1.2) produces a competitive method for GPCA.

1.2 Comparison to prior art

SPM (Algorithm 1) integrates various ideas in the literature into a single natural algorithm for symmetric tensor decomposition, with innovations in computational efficiency and convergence theory.

Step A of SPM is a variation of the classical Catalecticant Method [42]. This goes back to Sylvester of the 19th century [74]. Although Sylvester’s work is well-known to experts, SPM is, to our knowledge, the first scalable numerical method for tensor decomposition based on this reformulation of tensor decomposition. The work [7] proposed using symbolic techniques with

Sylvester’s Catalecticant Method to decompose symmetric tensors, but the algorithm seems slow already on modest-sized examples. Other works are [42] and [61]. Perhaps most related to ours is the latter by Oeding-Ottaviani. Importantly however, we differ in that [61] proposes using standard polynomial-solving techniques, e.g., Gröbner bases, to find (λ_i, a_i) after reformulating tensor decomposition via Sylvester’s Catalecticant Method. Unfortunately, a Gröbner basis approach is impractical already for small-sized problem dimensions, since Gröbner bases have doubly exponential running time (and require exact arithmetic). By contrast, Steps B and C of SPM consist of fast numerical iterations (based on the functional $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$ in Theorem 4.1) and optimized numerical linear algebra (see Appendix B), respectively.

Next, Step B of SPM connects with Kolda-Mayo’s SS-HOPM method for computing tensor eigenvectors [44]. Indeed, Step B is the higher-order power method applied to there modified symmetric tensor \tilde{T} of the same size as T , where \tilde{T} corresponds to the functional f in Theorem 4.1 (see Subsection 4.3). Here, the important point is that SS-HOPM is not designed to compute *CP tensor components* as in Decomposition 1.1, rather *tensor Z-eigenvectors* (see the definition in Section 2). These vectors are not the same for T . However, SPM identifies the Z-eigenvectors of \tilde{T} (computable by SS-HOPM) with the CP tensor components a_i of T . Also, in analyzing the convergence of Step B, we improve on the theory for SS-HOPM in general, settling a conjecture of [44].

Finally, Step C of SPM is related to the matrix identities in [22] and Wedderburn’s rank reduction formula [21]. For maximal efficiency, we derive an optimized implementation of this procedure using Householder reflections, to avoid as much recomputation as possible during deflation (see Appendix B).

In other respects, Algorithm 1 bears resemblances to De Lathauwer *et al.*’s Fourth-Order-Only Blind Identification (FOOBI) algorithm for fourth-order symmetric tensor decomposition [26], the methods for finding low-rank elements in subspaces of matrices in [66, 34, 57], and the asymmetric decomposition method in [65] repeatedly reducing the tensor length in given directions.

Most prior practical decomposition methods rely on direct non-convex optimization, essentially minimizing the squared residual

$$\left\| T - \sum_{i=1}^R \lambda_i a_i^{\otimes 2n} \right\|^2.$$

This may be done, e.g., by gradient descent or symmetric variants of alternating least squares [45]. In Section 6, we compare SPM to state-of-the-art numerical methods, in particular De Lathauwer and collaborators’ MATLAB package Tensorlab [77]. The comparison is done using standard random ensembles for symmetric tensors, as well as “worse-conditioned” tensors with correlated components. We also compare to quite different but state-of-the-art theoretical methods, namely Nie’s method [59, 60] and FOOBI [26, 12].

As for Algorithm 2 (SPM to compute (1.2)), this method has less precedence in the literature. Block term tensor decompositions were introduced in [24, 25, 28], with a focus on the asymmetric case. The individual blocks in the

symmetric block term decomposition (1.2), symmetric Tucker products, are studied in [11, 68, 43]. However, to our knowledge, Algorithm 2 is the first to specifically compute the symmetric tensor decomposition (1.2).

In Lemma 5.1, we derive a relationship between Decomposition (1.2) and subspace arrangements. The case $n = 2$ connects with [14, Section 5]. For a concrete application, we propose using SPM (Algorithm 2) for generalized principal component analysis. While GPCA is a popular learning problem, e.g., [80, 51, 75], we are not aware of any method-of-moments based approach similar to our proposal. Simulations in Section 6 show that SPM yields a competitive method for GPCA.

1.3 Organization of the paper

The rest of this paper is organized as follows. Section 2 establishes notation and basic definitions. Section 3 states the algorithm for CP symmetric tensor rank decomposition. Section 4 analyzes the convergence of the power iteration (Step B). Section 5 extends SPM to compute the symmetric block term decompositions. Section 6 presents numerics, with comparisons of runtime and noise sensitivity to the state-of-the-art. Also, we discuss the application to GPCA. Section 7 concludes, and the appendices contain certain technical proofs.

MATLAB code for the Subspace Power Method is publicly available at:

<https://www.github.com/joaompereira/SPM>

This may be used to run SPM and compute the decompositions (1.1) or (1.2) for a user-specified symmetric tensor, or to reproduce the results in Section 6.

2 Definitions and notation

We fix notation regarding tensors. We use $\mathcal{T}_L^m = (\mathbb{R}^L)^{\otimes m} = \mathbb{R}^{L^m}$ to denote the vector space of real (not necessarily symmetric) tensors of *order* m and *length* L in each *mode*. \mathcal{T}_L^m is a Euclidean space, with Frobenius inner product and norm. If $T \in \mathcal{T}_L^m$, then $T_{j_1 \dots j_m}$ is the entry indexed by $(j_1 \dots j_m) \in [L]^m$, where $[L] = \{1, \dots, L\}$. For tensors $T \in \mathcal{T}_L^m$ and $U \in \mathcal{T}_L^{m'}$, their *tensor product* in $\mathcal{T}_L^{m+m'}$ is defined by

$$(T \otimes U)_{j_1 \dots j_{m+m'}} = T_{j_1 \dots j_m} U_{j_{m+1} \dots j_{m+m'}} \quad \forall (j_1, \dots, j_{m+m'}) \in [L]^{m+m'}. \quad (2.1)$$

The *tensor power* $T^{\otimes d} \in \mathcal{T}_L^{dm}$ is the tensor product of T with itself d times. The *tensor dot product* (or tensor contraction) between $T \in \mathcal{T}_L^m$ and $U \in \mathcal{T}_L^{m'}$, with $m \geq m'$, is the tensor in $\mathcal{T}_L^{m-m'}$ defined by

$$(T \cdot U)_{j_{m'+1} \dots j_m} = \sum_{j_1=1}^L \dots \sum_{j_{m'}=1}^L T_{j_1 \dots j_m} U_{j_1 \dots j_{m'}}. \quad (2.2)$$

If $m = m'$, contraction coincides with the inner product, i.e., $T \cdot U = \langle T, U \rangle$. For $T \in \mathcal{T}_L^m$, a (real normalized) *Z-eigenvector/eigenvalue pair* $(v, \lambda) \in \mathbb{R}^m \times \mathbb{R}$ is a vector/scalar pair satisfying $T \cdot v^{\otimes(m-1)} = \lambda v$ and $\|v\|_2 = 1$, see [52, 67].

Definition 2.1 A tensor $T \in \mathcal{T}_L^m$ is *symmetric* if it is unchanged by any permutation of indices, that is,

$$T_{j_1 \dots j_m} = T_{j_{\sigma(1)} \dots j_{\sigma(m)}} \quad \forall (j_1, \dots, j_m) \in [L]^m \text{ and } \sigma \in \Pi^m, \quad (2.3)$$

where Π^m is the permutation group on $[m]$. We denote by $\text{Sym } \mathcal{T}_L^m$ the vector space of real symmetric tensors of order m and length L . A tensor $T \in \mathcal{T}_L^m$ may be *symmetrized* by applying the *symmetrizing operator*, Sym , defined as

$$\text{Sym}(T)_{j_1 \dots j_m} = \frac{1}{m!} \sum_{\sigma \in \Pi^m} T_{j_{\sigma(1)} \dots j_{\sigma(m)}} \quad \forall (j_1, \dots, j_m) \in [L]^m. \quad (2.4)$$

A nice conceptual fact is the existence of a one-to-one map between symmetric tensors and homogeneous polynomials, which we formalize as follows. Here $\mathbb{R}[X_1, \dots, X_L]$ stands for the ring of real polynomials in n variables X , and $\mathbb{R}[X_1, \dots, X_L]_m$ denotes the subspace of homogeneous degree m forms.

Lemma 2.2 *There is a function Φ from $\bigcup_{m=0}^{\infty} \mathcal{T}_L^m$ to $\bigcup_{m=0}^{\infty} \mathbb{R}[X_1, \dots, X_L]_m$ (the set of homogeneous polynomials in L variables X) such that:*

- For every integer m , Φ is a linear map from \mathcal{T}_L^m to $\mathbb{R}[X_1, \dots, X_L]_m$ and a one-to-one map from $\text{Sym } \mathcal{T}_L^m$ to $\mathbb{R}[X_1, \dots, X_L]_m$.
- For any vector $v \in \mathbb{R}^L$, symmetric matrix $M \in \mathbb{R}^{L \times L}$ and symmetric tensor $T \in \mathcal{T}_L^m$, the homogeneous polynomials $\Phi(v)$, $\Phi(M)$, $\Phi(T)$ are such that

$$\Phi(v)(X) = v^T X, \quad \Phi(M)(X) = X^T M X, \quad \Phi(T)(X) = \langle T, X^{\otimes m} \rangle. \quad (2.5)$$

- For symmetric tensors $T_1 \in \mathcal{T}_L^m$ and $T_2 \in \mathcal{T}_L^{m'}$, we have

$$\Phi(T_1)\Phi(T_2) = \Phi(\text{Sym}(T_1 \otimes T_2)). \quad (2.6)$$

The proof of Lemma 2.2 is standard, and available in Appendix A. We can now give two viewpoints on symmetric tensor decomposition.

Definition 2.3 For a symmetric tensor $T \in \text{Sym } \mathcal{T}_L^m$, a *real symmetric rank decomposition* (or *real symmetric CP decomposition*) is an expression

$$T = \sum_{i=1}^R \lambda_i a_i^{\otimes m}, \quad (2.7)$$

where $R \in \mathbb{Z}$ is smallest possible, $\lambda_i \in \mathbb{R}$, and $a_i \in \mathbb{R}^L$ (without loss of generality, $\|a_i\|_2 = 1$). The minimal R is the *real symmetric rank* of T .

Equivalently, via the one-to-one correspondence¹ Φ , a *real Waring decomposition* of a homogeneous polynomial $F \in \mathbb{R}[X_1, \dots, X_L]_m$ is an expression

$$F = \sum_{i=1}^R \lambda_i (a_i^T X)^m, \quad (2.8)$$

¹ That is, $\Phi(a_i^{\otimes m})(X) = (a_i^T X)^m$.

where $R \in \mathbb{Z}$ is smallest possible, $\lambda_i \in \mathbb{R}$, and $a_i^T X \in \mathbb{R}[X_1, \dots, X_L]_1$ are linear forms (wlog normalized). The minimal R is the *real Waring rank* of F .

In (2.8), each term $\lambda_i(a_i^T X)^m$ is a homogeneous polynomial in one linear form, or “latent variable”, namely $a_i^T X$. We consider a generalization: write F as a sum of homogeneous polynomials each in a small number of latent variables (linear forms). More formally, for a matrix $A_i \in \mathbb{R}^{L \times \ell_i}$, with $\ell_i \leq L$, we consider homogeneous polynomials in the ℓ_i linear forms $A_i^T X \in \mathbb{R}^{\ell_i}$.

Definition 2.4 For a homogeneous polynomial $F \in \mathbb{R}[X_1, \dots, X_L]_m$, consider an expression

$$F = \sum_{i=1}^R f_i(A_i^T X). \quad (2.9)$$

where $R, \ell_i \in \mathbb{Z}$ are positive, $A_i \in \mathbb{R}^{L \times \ell_i}$ and $f_i \in \mathbb{R}[a_1, \dots, a_{\ell_i}]_m$ is a homogeneous polynomial in the ℓ_i latent variables $A_i^T X \in \mathbb{R}^{\ell_i}$. We call (2.9) an (A_1, \dots, A_R) -*block term Waring decomposition* of F .

Similarly to (2.7) and (2.8), we can recast (2.9) as a type of symmetric tensor decomposition. First we need the definition of symmetric Tucker product.

Definition 2.5 For a symmetric tensor $T \in \text{Sym } \mathcal{T}_L^m$, a *real symmetric Tucker product* is an expression

$$T = (A; \dots; A) \cdot \mathbf{A}, \quad (2.10)$$

where $\mathbf{A} \in \text{Sym } \mathcal{T}_\ell^m$ (some ℓ) is the “core tensor”, $A \in \mathbb{R}^{L \times \ell}$ is the “factor matrix”, and the (j_1, \dots, j_m) -entry of the middle/RHS in (2.10) is defined by:

$$((A; \dots; A) \cdot \mathbf{A})_{j_1 \dots j_m} = \sum_{k_1=1}^{\ell} \dots \sum_{k_m=1}^{\ell} \mathbf{A}_{k_1 \dots k_m} A_{j_1 k_1} \dots A_{j_m k_m}. \quad (2.11)$$

The notation of [29] is used for Tucker products, and a total of m matrices A are meant by $(A; \dots; A)$. Speaking roughly, Definition 2.5 encapsulates a symmetric tensor T that is supported in an ℓ^m subtensor or “block”, after a symmetric change of basis. When $\ell = 1$, symmetric Tucker products are symmetric rank-1 tensors. So, sums of symmetric Tucker products give a generalization of rank decompositions.

Definition 2.6 For a symmetric tensor $T \in \text{Sym } \mathcal{T}_L^m$, consider an expression

$$T = \sum_{i=1}^R (A_i; \dots; A_i) \cdot \mathbf{A}_i \quad (2.12)$$

for $\mathbf{A}_i \in \text{Sym } \mathcal{T}_{\ell_i}^m$ (some ℓ_i) and $A_i \in \mathbb{R}^{L \times \ell_i}$. We call (2.12) an (A_1, \dots, A_R) -*symmetric block term tensor decomposition*.

As with rank and Waring decompositions, Definitions 2.4 and 2.6 are equivalent in the following sense. Since f_i is a homogeneous polynomial, there exists $\mathbf{A}_i \in \mathcal{T}_{\ell_i}^m$ such that $\Phi(\mathbf{A}_i) = f_i$. We then have²

$$\Phi((A_i; \dots; A_i) \cdot \mathbf{A}_i)(X) = f_i(A_i^T X). \quad (2.13)$$

Definition 2.6 is the symmetric variant of De Lathauwer's *block term decomposition*, introduced in [24, 25, 28]. Definitions 2.4 and 2.5 specialize to Definition 2.3 when $\ell_1 = \dots = \ell_R = 1$ and R is minimal. We return to these generalizations of usual rank decompositions in Section 5, in connection to GPCA.

Next, we set conventions for unfolding tensors into matrices and vectors.

Definition 2.7 For an even-order tensor $T \in \mathcal{T}_L^{2n}$, the *square matrix flattening* of T denoted $\text{mat}(T) \in \mathbb{R}^{L^n \times L^n}$ is defined by

$$\text{mat}(T)_{j_1 \dots j_n, j_{n+1} \dots j_{2n}} = T_{j_1 \dots j_{2n}} \quad \forall (j_1, \dots, j_n), (j_{n+1}, \dots, j_{2n}) \in [L]^n, \quad (2.14)$$

where lexicographic order is used on $[L]^n$. For $U \in \mathcal{T}_L^n$, the *vectorization* of U denoted $\text{vec}(U) \in \mathbb{R}^{L^n}$ is also with respect to the lexicographic order on $[L]^n$.

Definition 2.8 Let $A \in \mathbb{R}^{L \times R}$ be a matrix with columns $a_1, \dots, a_R \in \mathbb{R}^L$. Then the n -th *Khatri-Rao power* of A , denoted $A^{\bullet n} \in \mathbb{R}^{L^n \times R^n}$, is the matrix with columns $\text{vec}(a_1^{\otimes n}), \dots, \text{vec}(a_R^{\otimes n}) \in \mathbb{R}^{L^n}$.

Remark 2.9 In this paper, some of our guarantees regarding decomposition (2.7) hold for *generic* a_i, λ_i . By that, we mean that there exists a polynomial p (often unspecified) in the entries of a_i and λ_i such that the guarantee is valid whenever $p(a_1, \dots, \lambda_R) \neq 0$, and moreover, $p(a_1, \dots, \lambda_R) \neq 0$ holds for some choice of unit-norm $a_i \in \mathbb{R}^L$ and $\lambda_i \in \mathbb{R}$. This is called *Zariski-generic* in algebraic geometry [38]. In particular, it implies that the guarantees hold with probability 1 if a_i and λ_i are drawn from *any* absolutely continuous probability distributions on the sphere and real line. Similar remarks apply to decomposition (2.12) and A_i, \mathbf{A}_i .

3 Algorithm description

In this section, we describe SPM (Algorithm 1). The input is a real even-order symmetric tensor $T \in \text{Sym } \mathcal{T}_L^{2n}$ ($n \geq 2$) assumed to obey a low-rank decomposition (1.1), where $R = O(L^n)$. The output is approximations to $(\lambda_i, \pm a_i)$ for $i = 1, \dots, R$. It is convenient to divide the description into three main steps: EXTRACT SUBSPACE, POWER METHOD and DEFLATE.

² We include a proof of (2.13) in Appendix A.

To start, observe the tensor decomposition (1.1) of T is equivalent to a certain matrix factorization of the flattening of T (Definition 2.7).

$$\begin{aligned} \text{mat}(T) &= \sum_{i=1}^R \lambda_i \text{mat}(a_i^{\otimes 2n}), \\ &= \sum_{i=1}^R \lambda_i \text{vec}(a_i^{\otimes n}) \text{vec}(a_i^{\otimes n})^T. \end{aligned} \quad (3.1)$$

Letting $A \in \mathbb{R}^{L \times R}$ denote the matrix with columns $a_1, \dots, a_R \in \mathbb{R}^L$, and $\Lambda \in \mathbb{R}^{R \times R}$ be the diagonal matrix with entries $\lambda_1, \dots, \lambda_R$, then (3.1) reads:

$$\text{mat}(T) = A^{\bullet n} \Lambda (A^{\bullet n})^T. \quad (3.2)$$

Here $A^{\bullet n} \in \mathbb{R}^{L^n \times R}$ is the Khatri-Rao power of A (Definition 2.8). Define the subspace $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ as the column space of A (after un-vectorization), i.e.,

$$\mathcal{A} = \text{Span}\{a_1^{\otimes n}, \dots, a_R^{\otimes n}\} \subset \text{Sym } \mathcal{T}_L^n. \quad (3.3)$$

In light of 3.2, EXTRACT SUBSPACE obtains an orthonormal basis for \mathcal{A} using $\text{mat}(T)$. Note that if the columns of $A^{\bullet n}$ were orthogonal, (3.2) would be an eigendecomposition of $\text{mat}(T)$; however $\langle \text{vec}(a_i^{\otimes n}), \text{vec}(a_j^{\otimes n}) \rangle = \langle a_i, a_j \rangle^n$, so the columns of $A^{\bullet n}$ are generally not orthogonal (and cannot be so if $R > L$). Nevertheless, the eigendecomposition of $\text{mat}(T)$ does provide *some* orthonormal basis for \mathcal{A} (not $a_1^{\otimes n}, \dots, a_R^{\otimes n}$), so long as $A^{\bullet n}$ is full-rank.

Proposition 3.1 – Assume $A^{\bullet n}$ has rank R . Then $\text{mat}(T)$ has rank R .

Moreover, if (V, D) is a thin eigendecomposition of $\text{mat}(T)$, i.e., $\text{mat}(T) = VDV^T$ where $V \in \mathbb{R}^{L^n \times R}$ has orthonormal columns and $D \in \mathbb{R}^{R \times R}$ is full-rank diagonal, then the columns of V give an orthonormal basis of \mathcal{A} .

– If $R \leq \binom{L+n-1}{n}$ and a_1, \dots, a_R are generic, then $A^{\bullet n}$ has rank R .

Proof First assume $A^{\bullet n}$ has rank R . Let (Q, W) be a thin QR-factorization of $A^{\bullet n}$, i.e., $A^{\bullet n} = QW$ where $Q \in \mathbb{R}^{L^n \times R}$ is a matrix with orthogonal columns and $W \in \mathbb{R}^{R \times R}$ is upper-triangular. Let (O, D) be an eigendecomposition of $W\Lambda W^T$, i.e., $W\Lambda W^T = ODO^T$ where $O \in \mathbb{R}^{R \times R}$ is orthogonal and $D \in \mathbb{R}^{R \times R}$ is diagonal. We have:

$$\text{mat}(T) = A^{\bullet n} \Lambda (A^{\bullet n})^T = (QW) \Lambda (QW)^T = Q(ODO^T)Q^T = (QO)D(QO)^T. \quad (3.4)$$

Since $A^{\bullet n}$ has full column rank, W is invertible, and so D is invertible. Thus the above shows $\text{rank}(\text{mat}(T)) = R$, and (QO, D) is a thin eigendecomposition of $\text{mat}(T)$. In particular, $\text{colspan}(QO) = \text{colspan}(Q) = \text{colspan}(A^{\bullet n})$, and the columns of QO give an orthonormal basis of \mathcal{A} (after un-vectorization).

Next assume $R \leq \binom{L+n-1}{n}$. Full column rank of $A^{\bullet n}$ imposes a Zariski-open condition on a_1, \dots, a_R , described by the non-vanishing of the maximal minors of $A^{\bullet n}$, i.e., certain polynomials in the entries of a_i . The condition holds generically because it holds for some a_1, \dots, a_R , since rank-1 symmetric tensors span $\text{Sym } \mathcal{T}_L^n$ and $\dim(\text{Sym } \mathcal{T}_L^n) = \binom{L+n-1}{n}$. \square

Thus SPM extracts $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ from an eigendecomposition of $\text{mat}(T)$.

POWER METHOD, the next step of SPM, seeks to find a rank-1 point in \mathcal{A} . As in [61], the following fact is essential underpinning. For the proof and discussion, let $\mathcal{V}_{L,n}$ denote the set of rank ≤ 1 tensors in $\text{Sym } \mathcal{T}_L^n$.

Proposition 3.2 *If $R \leq \binom{L+n-1}{n} - L$, then for generic a_1, \dots, a_R , the only rank-1 tensors in $\mathcal{A} = \text{Span}\{a_1^{\otimes n}, \dots, a_R^{\otimes n}\} \subset \text{Sym } \mathcal{T}_L^n$ are $a_1^{\otimes n}, \dots, a_R^{\otimes n}$ (up to nonzero scale).*

Proof This is a special case of the generalized trisecant lemma in algebraic geometry, see [19, Prop. 2.6] or [38, Exer. IV-3.10]. The set of rank ≤ 1 tensors forms an irreducible algebraic cone of dimension L linearly spanning its ambient space, $\text{Sym } \mathcal{T}_L^n$. This is the Veronese variety, $\mathcal{V}_{L,n}$. Meanwhile, \mathcal{A} is a secant plane through R general points on $\mathcal{V}_{L,n}$. The dimensions of $\mathcal{V}_{L,n}$ and \mathcal{A} are *sub-complimentary*, i.e.,

$$\dim(\mathcal{V}_{L,n}) + \dim(\mathcal{A}) = L + R \leq \binom{L+n-1}{n} = \dim(\text{Sym } \mathcal{T}_L^n). \quad (3.5)$$

So the generalized trisecant lemma applies and tells us that we have no unexpected intersection points, i.e., $\mathcal{V}_{L,n} \cap \mathcal{A} = \text{Span}\{a_1^{\otimes n}\} \cup \dots \cup \text{Span}\{a_R^{\otimes n}\}$. \square

Continuing, the second step of SPM seeks a rank-1 element in \mathcal{A} by iterating the tensor power method on a suitably modified tensor, such that a_i are eigenvectors of the modified tensor. Our power iteration may be motivated as an approximate alternating projection scheme to compute (unit-norm) points in $\mathcal{V}_{L,n} \cap \mathcal{A}$, as follows. Let $P_{\mathcal{A}}$ be orthogonal projection from symmetric order- n tensors onto \mathcal{A} , and $P_{\text{rank-1}}$ orthogonal projection onto the set of unit-norm elements of $\mathcal{V}_{L,n}$. So $P_{\mathcal{A}}$ is a linear map, while $P_{\text{rank-1}}$ is a nonlinear map well-defined outside an algebraic hypersurface in $\text{Sym } \mathcal{T}_L^n$ [27]. To compute $\mathcal{V}_{L,n} \cap \mathcal{A}$, a naive alternating projection method could initialize unit-norm $x \in \mathbb{R}^L$ and iterate until convergence:

$$x^{\otimes n} \leftarrow P_{\text{rank-1}}(P_{\mathcal{A}}(x^{\otimes n})). \quad (3.6)$$

In order to calculate $P_{\mathcal{A}}(x^{\otimes n})$, one could use that the columns of V (known from the first step of SPM) form an orthonormal basis of \mathcal{A} , and compute

$$\text{vec}(P_{\mathcal{A}}(x^{\otimes n})) = VV^T \text{vec}(x^{\otimes n}). \quad (3.7)$$

Unfortunately, evaluating $P_{\text{rank-1}}$ is NP-hard as soon as $n \geq 3$. Nevertheless, one could try the tensor power method for an approximation [27, 44]. Given a tensor U and unit-norm vector z , the tensor power method [27, 44] iterates

$$z \leftarrow \frac{U \cdot z^{\otimes n-1}}{\|U \cdot z^{\otimes n-1}\|}. \quad (3.8)$$

However performing this until convergence inside each iteration of alternating projections seems costly, and it is unclear if having a very good rank-1

approximation of $P_{\mathcal{A}}(x^{\otimes n})$ at each step improves the convergence of the alternating projection method. Instead, one could simply apply (3.8) once with $U = P_{\mathcal{A}}(x^{\otimes n})$ and $z = x$.

Such informal reasoning, in the previous paragraph, suggests replacing alternating projections (3.6) by the cheaper iteration

$$x \leftarrow \frac{P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes(n-1)}}{\|P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes(n-1)}\|}, \quad (3.9)$$

where (3.7) is used to evaluate $P_{\mathcal{A}}$. As in [44], we will actually use a shifted variant for its better convergence theory:

$$x \leftarrow \frac{P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes(n-1)} + \gamma x}{\|P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes(n-1)} + \gamma x\|}, \quad (3.10)$$

where $\gamma > 0$ is a constant depending just on n (see Lemma 4.4). Thus the POWER METHOD step of SPM is to iterate (3.10) until convergence. The reader may verify that the iteration (3.10) is equivalent to projected gradient ascent applied to the function $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$ with constant step-size $\frac{1}{\gamma}$.

We defer analysis of (3.10) to the Section 4, and instead finish explaining Algorithm 1. The last step, DEFLATE, is a procedure to remove rank-1 components from T . Recall that (V, D) is a thin eigendecomposition of $\text{mat}(T)$ (Proposition 3.1). Suppose that (3.10) with random initialization converged to $a \in \mathbb{R}^L$. We check whether $\text{vec}(a^{\otimes n})$ lies in the column space of V (in practice, up to some numerical threshold). If it does not, a is discarded and we repeat POWER METHOD with a fresh random initialization. Otherwise, we assume $a = a_1$ (after relabeling), and then determine λ_1 Wedderburn rank reduction [21], as follows. Let $\alpha \in \mathbb{R}^R$ be such that $V\alpha = \text{vec}(a_1^{\otimes n})$, i.e., $\alpha = V^T \text{vec}(a_1^{\otimes n})$. Consider rank-1 updates, $T - t a_1^{\otimes 2n}$, for scalars $t \in \mathbb{R}$. On one hand, these flatten to

$$\begin{aligned} \text{mat}(T) - t \text{vec}(a_1^{\otimes n}) \text{vec}(a_1^{\otimes n})^T = \\ (\lambda_1 - t) \text{vec}(a_1^{\otimes n}) \text{vec}(a_1^{\otimes n})^T + \sum_{i=2}^R \lambda_i \text{vec}(a_i^{\otimes n}) \text{vec}(a_i^{\otimes n})^T, \end{aligned} \quad (3.11)$$

while on the other hand,

$$\text{mat}(T) - t \text{vec}(a_i^{\otimes n}) \text{vec}(a_i^{\otimes n})^T = V(D - t\alpha\alpha^T)V^T. \quad (3.12)$$

Assuming $A^{\bullet n}$ has full column rank (the hypothesis in Proposition 3.1), then (3.11) shows $\text{mat}(T - t a_1^{\otimes 2n})$ has rank $R - 1$ if $t = \lambda_1$, and rank R otherwise. By (3.12), $D - t\alpha\alpha^T$ is singular if $t = \lambda_1$, and is non-singular otherwise. As shown in Lemma B.1, this implies $\lambda_1 = (\alpha^T D^{-1} \alpha)^{-1}$.

We proceed with the deflation. For faster computation, instead of updating $T \leftarrow T - \lambda_1 a_1^{\otimes 2n}$ and recalculating the eigendecomposition of $\text{mat}(T)$, DEFLATE updates the eigendecomposition of T directly.

Let (O, \tilde{D}) be a thin eigendecomposition of $D - \lambda_1 \alpha \alpha^T$, thus O is $R \times (R-1)$ with orthonormal columns, \tilde{D} is $(R-1) \times (R-1)$ diagonal, and $D - \lambda_1 \alpha \alpha^T = O \tilde{D} O^T$. Then,

$$\text{mat}(T - \lambda_1 a_1^{\otimes 2n}) = V(D - \lambda_1 \alpha \alpha^T) V^T = VO \tilde{D} (VO)^T. \quad (3.13)$$

Since $VO \in \mathbb{R}^{R \times (R-1)}$ has orthonormal columns and $\tilde{D} \in \mathbb{R}^{(R-1) \times (R-1)}$ is diagonal, (VO, \tilde{D}) is a thin eigendecomposition of $\text{mat}(\tilde{T})$, and so we deflate:

$$(V, D) \leftarrow (VO, \tilde{D}). \quad (3.14)$$

POWER METHOD and DEFLATE repeat as subroutines, such that each tensor component $(\lambda_i, \pm a_i)$ is removed one at a time, until all components have been found. The entire algorithm is specified by the following pseudocode.

Algorithm 1 Subspace Power Method

Input: Generic $T \in \text{Sym } \mathcal{T}_L^{2n}$ of rank $\leq \binom{L+n-1}{n} - L$
Output: Rank R and tensor decomposition $\{(\lambda_i, a_i)\}_{i=1}^R$

$(V, D) \leftarrow \text{eig}(\text{mat}(T))$ EXTRACT SUBSPACE
 $R \leftarrow \text{rank}(\text{mat}(T))$
for $i = 1$ **to** R **do**
 $x \leftarrow \text{random}(\mathbb{S}^{L-1})$ POWER METHOD
 repeat
 $x \leftarrow \frac{P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes n-1} + \gamma x}{\|P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes n-1} + \gamma x\|}$
 until convergence
 $a_i \leftarrow x$

 $\alpha \leftarrow V^T \text{vec}(a_i^{\otimes n})$ DEFLATE
 $\lambda_i \leftarrow (\alpha^T D^{-1} \alpha)^{-1}$
 $(O, \tilde{D}) \leftarrow \text{eig}(D - \lambda_i \alpha \alpha^T)$
 $V \leftarrow VO$
 $D \leftarrow \tilde{D}$
return R and $\{(\lambda_i, a_i)\}_{i=1}^R$

The computational costs of Algorithm 1 are upper-bounded as follows. In EXTRACT SUBSPACE, computing $\text{eig}(\text{mat}(T))$ upfront costs $O(L^{3n})$. Suppose $r \leq R$ components (λ_i, a_i) are yet to be found. In POWER METHOD, each iteration is $O(rL^n)$, the price of applying $P_{\mathcal{A}}$. In DEFLATE, computing $V^T \alpha$ is $O(rL^n)$, computing $\lambda_i = (\alpha^T D^{-1} \alpha)^{-1}$ is $O(r)$, computing an eigendecomposition of $D - \lambda_i \alpha \alpha^T$ naively is $O(r^3)$ and computing VO is $O(r^2 L^n)$.

Remark 3.3 For the simulations in Section 6, we slightly modified DEFLATE. This gave a modest speedup to SPM overall, in some cases $\approx 20\%$. The modified deflation is slightly more involved, and we present it in Appendix B.

Remark 3.4 An odd-order variant of SPM is straightforward to describe. Given $T \in \text{Sym } \mathcal{T}_L^{2n-1}$ ($n \geq 2$), one flattens T to an $L^n \times L^{n-1}$ matrix and extracts the

column space $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$. The power method proceeds as before, while deflation receives minor adjustment. Unfortunately, odd-order SPM has no chance of working up rank roughly the square root of the number of tensor entries, as believed tractable for efficient algorithms [54]. It is limited by the maximal rank of the rectangular flattening, which is $\dim(\text{Sym } \mathcal{T}_L^{n-1}) = \binom{L+n-2}{n-1}$ since each row of the flattened matrix lies in $\text{Sym } \mathcal{T}_L^{n-1}$ (after un-vectorization).

4 Power Method Analysis

The object of this section is to prove the following guarantees regarding POWER METHOD in Algorithm 1. We denote by \mathbb{S}^{L-1} the unit sphere in \mathbb{R}^L .

Theorem 4.1 *Given a tensor $T \in \text{Sym } \mathcal{T}_L^{2n}$ with decomposition as in (1.1), where $\lambda_i \in \mathbb{R}$ and $a_i \in \mathbb{S}^{L-1}$, set $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$, and consider the constrained optimization problem:*

$$\max_{x \in \mathbb{S}^{L-1}} f(x). \quad (4.1)$$

Define the sequence:

$$x_{k+1} = \frac{P_{\mathcal{A}}(x_k^{\otimes n}) \cdot x_k^{\otimes(n-1)} + \gamma x_k}{\|P_{\mathcal{A}}(x_k^{\otimes n}) \cdot x_k^{\otimes(n-1)} + \gamma x_k\|}, \quad (4.2)$$

where $x_1 \in \mathbb{S}^{L-1}$ is some initialization, and $\gamma \in \mathbb{R}_{>0}$ is any fixed shift such that $F(x) = f(x) + \frac{\gamma}{2n}(x^T x)^n$ is a strictly convex function on \mathbb{R}^n . For example, the following is a sufficiently large shift:

$$\gamma = \begin{cases} \sqrt{\frac{(n-1)}{2n}} & \text{if } n \leq 4 \\ \frac{2-\sqrt{2}}{2}\sqrt{n} & \text{if } n > 4. \end{cases} \quad (4.3)$$

- For all initializations x_1 , (4.2) is well-defined and converges monotonically to a constrained critical point x_* of (4.1) at a power rate. (That is, $f(x_{k+1}) \geq f(x_k)$ for all k , and there exist positive constants $\tau = \tau(\mathcal{A}, \gamma, x_1)$ and $C = C(\mathcal{A}, \gamma, x_1)$ such that $\|x_k - x_*\| < C \left(\frac{1}{k}\right)^\tau$ for all k .)
- For a full Lebesgue measure subset of initializations x_1 , (4.2) converges to a constrained second-order critical point of (4.1).
- For $R \leq \binom{L+n-1}{n} - L$ and generic $\{a_i\}_{i=1}^R$, the global optima of (4.1) are precisely $\pm a_i$. Moreover, if $L \geq n - 1$, each global maxima is attractive: for all initializations x_1 sufficiently close to $\pm a_i$, (4.2) converges to $\pm a_i$ at an exponential rate. (That is, there exist positive constants $\delta = \delta(\mathcal{A}, \gamma, i)$, $\tau = \tau(\mathcal{A}, \gamma, i)$ and $C = C(\mathcal{A}, \gamma, i)$ such that $\|x_1 - \pm a_i\| < \delta$ implies $\|x_k - \pm a_i\| < Ce^{-k\tau}$ for all k .)

Remark 4.2 It is clear that if $\gamma > 0$, then the sequence (4.2) is well-defined. Indeed, $\langle x_k, P_{\mathcal{A}}(x_k^{\otimes n}) \cdot x_k^{\otimes(n-1)} + \gamma x_k \rangle = \|P_{\mathcal{A}}(x_k^{\otimes n})\|^2 + \gamma > 0$, so that the denominator in (4.2) does not vanish.

Remark 4.3 Critical points are meant in the usual Lagrange multiplier sense. Setting $\mathcal{L}(x, \mu) = f(x) + \frac{\mu}{2}(x^T x - 1)$, one says $x_* \in \mathbb{S}^{L-1}$ is a *constrained critical point* of (4.1) if there exists $\mu_* \in \mathbb{R}$ such that $\nabla_x \mathcal{L}(x_*, \mu_*) = 0$, while x_* is *second-order critical* if in addition $(I - xx^T) \nabla_{xx}^2 \mathcal{L}(x_*, \mu_*) (I - xx^T) \preceq 0$.

Our proof strategy for Theorem 4.1 is as follows. First, we will regard (4.2) as the shifted power method (SS-HOPM) [44] applied to a certain modified tensor $\tilde{T} \in \text{Sym } \mathcal{T}_L^{2n}$. Then, we will obtain the first two parts of theorem by sharpening the analysis of SS-HOPM in general. Finally, for the third statement, we will exploit special geometric properties of the modified tensor \tilde{T} .

4.1 POWER METHOD as SS-HOPM applied to a modified tensor $\tilde{T} \in \text{Sym } \mathcal{T}_L^{2n}$

In Algorithm 1, let $V_1, \dots, V_R \in \text{Sym } \mathcal{T}_L^n$ be the orthonormal basis of \mathcal{A} given by the columns of V . Then

$$P_{\mathcal{A}}(x^{\otimes n}) = \sum_{i=1}^R \langle V_i, x^{\otimes n} \rangle V_i. \quad (4.4)$$

Now define

$$\tilde{T} = \sum_{i=1}^R \text{Sym}(V_i \otimes V_i) \in \text{Sym } \mathcal{T}_L^{2n}, \quad (4.5)$$

and notice that

$$\tilde{T} \cdot x^{\otimes 2n} = \sum_{i=1}^R \langle V_i, x^{\otimes n} \rangle^2 = \|P_{\mathcal{A}}(x^{\otimes n})\|^2, \quad (4.6)$$

and

$$\tilde{T} \cdot x^{\otimes 2n-1} = \sum_{i=1}^R \langle V_i, x^{\otimes n} \rangle V_i \cdot x^{\otimes n-1} = P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes n-1}. \quad (4.7)$$

Hence (3.9) is the vanilla power method (3.8) for the tensor \tilde{T} . Recalling Lemma 2.2, we also have $\tilde{T} = \Phi^{-1}(f(X))$, with f defined in (4.1). Shifting by γx , (3.10) and (4.2) are SS-HOPM applied to \tilde{T} , or equivalently, the vanilla power method (3.8) applied to $\Phi^{-1}(F(X))$ where $F(X) = f(X) + \frac{\gamma}{2n}(X^T X)^n$:

$$x_{k+1} = \frac{\nabla F(x_k)}{\|\nabla F(x_k)\|}. \quad (4.8)$$

Usually in SS-HOPM, the shift γ is selected so the homogeneous polynomial F becomes convex on \mathbb{R}^L . For functions of the form $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$, where $P_{\mathcal{A}}$ is orthogonal projection onto *any* linear subspace $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ (spanned by rank-1 points, or not), we have the following estimate for γ .

Lemma 4.4 *Let $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ be any linear subspace, $P_{\mathcal{A}} : \text{Sym } \mathcal{T}_L^n \rightarrow \mathcal{A}$ be orthogonal projection, and $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$. If $\|x\| = 1$ and $f(x) \geq \nu$, then*

$$C_{\nu} C_n \geq -\frac{1}{2n} \min_{y \in \mathbb{S}^{L-1}} y^T \nabla^2 f(x) y, \quad (4.9)$$

where

$$C_{\nu} = \begin{cases} \sqrt{\nu(1-\nu)} & \text{if } \nu > \frac{1}{2} \\ \frac{1}{2} & \text{if } \nu \leq \frac{1}{2}, \end{cases} \quad C_n = \begin{cases} \sqrt{\frac{2(n-1)}{n}} & \text{if } n \leq 4 \\ (2-\sqrt{2})\sqrt{n} & \text{if } n > 4. \end{cases} \quad (4.10)$$

In particular, $F(x) = f(x) + \frac{\gamma}{2n}(x^T x)^n$ is convex on \mathbb{R}^L when $\gamma \geq \frac{1}{2}C_n$.

The proof of Lemma 4.4 is a lengthy calculation given in Appendix C.1. We have provided this bound, instead of an easier one, since (4.9) and (4.10) suggest interesting *adaptive* shifts for SPM as in [47] (see Remark 4.13).

4.2 Sharpening the global convergence analysis of SS-HOPM in general

We now prove the first two parts of Theorem 4.1 by sharpening the analysis of SS-HOPM in general. For this subsection, $f(x)$ is **any** homogeneous polynomial function on \mathbb{R}^L of degree $2n$. We consider the optimization problem

$$\max_{x \in \mathbb{S}^{L-1}} f(x), \quad (4.11)$$

and define the sequence

$$x_{k+1} = \frac{\nabla f(x_k) + \gamma x_k}{\|\nabla f(x_k) + \gamma x_k\|}, \quad (4.12)$$

where $x_1 \in \mathbb{S}^{L-1}$ is some initialization, and $\gamma \in \mathbb{R}$ is any shift such that $F(x) = f(x) + \frac{\gamma}{2n}(x^T x)^n$ is convex on \mathbb{R}^n and strictly positive on $\mathbb{R}^L \setminus \{0\}$.

The next result resolves Kolda-Mayo's conjecture stated on [44, p. 1107]. This proves that SS-HOPM converges unconditionally. Previously, convergence was guaranteed only for tensors with finitely many real eigenvectors. While all symmetric tensors have finitely many eigenvalues, not all have finitely many real eigenvectors (see [16, Theorem 5.5]). Further, given a low-dimensional family of “structured” tensors, we do not know how to efficiently verify if a generic member of that family has only finitely many real eigenvectors. In particular, we could not verify the condition for the modified low-rank tensors arising in SPM³, i.e., \tilde{T} in (4.5), so we focused on proving the conjecture. The main technical tool is the Łojasiewicz inequality for real analytic functions [53], which was first used in convergence analysis in [1].

³ For the generalization of SPM in Section 5 to compute symmetric block term decompositions (2.12), the corresponding tensors *do have infinitely many real eigenvectors*, namely $\mathcal{S} \cap \mathbb{S}^{L-1}$.

Theorem 4.5 (Unconditional convergence of SS-HOPM) *As above, let f be any homogeneous polynomial function on \mathbb{R}^L of degree $2n$. Let $\gamma \in \mathbb{R}$ be any shift such that $F(x) = f(x) + \frac{\gamma}{2n}(x^T x)^n$ is convex on \mathbb{R}^L and strictly positive on $\mathbb{R}^L \setminus \{0\}$. Then for all initializations $x_1 \in \mathbb{S}^{L-1}$, the iterates (4.12) are well-defined and converge monotonically to a constrained critical point x_* of (4.11) at a power rate.*

Proof Note that the denominator in the RHS of (4.12) does not vanish, as $\nabla F(x_k) = \nabla f(x_k) + \gamma(x_k^T x_k)^{n-1}x_k = \nabla f(x_k) + \gamma x_k$ implies $\langle x_k, \nabla f(x_k) + \gamma x_k \rangle = \langle x_k, \nabla F(x_k) \rangle = 2nF(x_k) > 0$. Hence (x_k) is well-defined. Also, $F(x) = f(x) + \frac{\gamma}{2n}$ for $\|x\| = 1$, so constrained critical points of (4.11) are the same as constrained critical points of

$$\max_{x \in \mathbb{S}^{L-1}} F(x), \quad (4.13)$$

monotonic convergence with respect to f is the same as with respect to F , the iteration (4.12) is the same as

$$x_{k+1} = \frac{\nabla F(x_k)}{\|\nabla F(x_k)\|}, \quad (4.14)$$

and we may work just with the convex function $F(x)$ and (4.14).

By convexity of F ,

$$F(x_{k+1}) - F(x_k) \geq \nabla F(x_k)^T (x_{k+1} - x_k). \quad (4.15)$$

By definition of x_{k+1} and Cauchy-Schwarz,

$$\nabla F(x_k)^T (x_{k+1} - x_k) \geq 0, \quad (4.16)$$

so $(F(x_k))$ monotonically increases.

Now suppose $\lim_{k \rightarrow \infty} x_k = x_*$. Taking the limit of both sides of (4.14), continuity gives:

$$\frac{\nabla F(x_*)}{\|\nabla F(x_*)\|} = x_* \Rightarrow \exists \mu_* \in \mathbb{R} \text{ s.t. } \nabla F(x_*) = -\mu_* x_*, \quad (4.17)$$

so x_* is a constrained critical point.

Since $(x_k) \subset \mathbb{S}^{L-1}$, the sequence (x_k) has an accumulation point by compactness. To prove x_k in fact converges,⁴ we shall apply the powerful convergence result of Schneider-USchmajew [70, Theorem 2.3]. This is based on the Łojasiewicz inequality for real analytic functions. For completeness, we have provided a precise statement of the we need result in Appendix C.2 (for convenience, slightly simplified as compared to Schneider-USchmajew's own statement). We take $\mathcal{M}, \mathcal{D}, f$ in Theorem C.1 to be \mathbb{S}^{L-1} , an open neighborhood of \mathcal{M} and F . It remains to verify the hypotheses of Theorem C.1.

⁴ Intuitively, we must rule out that (x_k) follows pathological dynamics as in [1, Fig. 2.1].

- Firstly, here $\mathcal{M} \subseteq \mathbb{R}^N$ of Theorem C.1 is the unit sphere $\mathbb{S}^{L-1} \subseteq \mathbb{R}^L$. Indeed, the sphere is locally the image of a real analytic map out of Euclidean space. Also, the function f of Theorem C.1 is F , which is a polynomial, certainly real-analytic.
- Here the tangent cones $T_x \mathcal{M}$ are given by $T_x(\mathbb{S}^{L-1}) = \{\xi \in \mathbb{R}^{L-1} : \langle \xi, x \rangle = 0\} \subseteq \mathbb{R}^L$, the perpendicular space to x . Therefore, in the notation of (C.21), $(\nabla F(x))^\perp = \|\text{grad } F(x)\|$, where $\text{grad } F(x) := (I - xx^\top) \nabla F(x)$ (the Riemannian gradient of F).
- To check **(A1)** in Theorem C.1, we must verify there exists $\sigma > 0$ such that for large enough k it holds

$$F(x_{k+1}) - F(x_k) \geq \sigma \|\text{grad } F(x_k)\| \|x_{k+1} - x_k\|. \quad (4.18)$$

We shall prove that $\sigma = \frac{1}{2}$ works. Since F is convex:

$$\begin{aligned} F(x_{k+1}) - F(x_k) &\geq \nabla F(x_k)^T (x_{k+1} - x_k) \\ &= \|\nabla F(x_k)\| x_{k+1}^T (x_{k+1} - x_k) \\ &= \|\nabla F(x_k)\| (1 - \langle x_{k+1}, x_k \rangle) \\ &= \frac{1}{2} \|\nabla F(x_k)\| \|x_{k+1} - x_k\|^2. \end{aligned} \quad (4.19)$$

We also have:

$$\begin{aligned} \|\text{grad } F(x_k)\|^2 &= \|\nabla F(x_k) - \langle \nabla F(x_k), x_k \rangle x_k\|^2 \\ &= \|\nabla F(x_k)\|^2 \|x_{k+1} - \langle x_{k+1}, x_k \rangle x_k\|^2 \\ &= \|\nabla F(x_k)\|^2 (1 - \langle x_{k+1}, x_k \rangle)^2 \\ &= \|\nabla F(x_k)\|^2 (1 - \langle x_{k+1}, x_k \rangle)(1 + \langle x_{k+1}, x_k \rangle) \\ &\leq 2 \|\nabla F(x_k)\|^2 (1 - \langle x_{k+1}, x_k \rangle) \\ &\leq \|\nabla F(x_k)\|^2 \|x_{k+1} - x_k\|^2. \end{aligned} \quad (4.20)$$

Substituting the square root of (4.20) into (4.19) gives (4.18) with $\sigma = \frac{1}{2}$.

- To check **(A2)** in Theorem C.1, we must verify for large enough k it holds

$$\text{grad } F(x_k) = 0 \implies x_{k+1} = x_k. \quad (4.21)$$

But this is clear from (4.14).

- To check **(A3)** in Theorem C.1, we must verify there exists a constant $\rho > 0$ such that for large enough k it holds

$$\|x_{k+1} - x_k\| \geq \rho \|\text{grad } F(x_k)\|. \quad (4.22)$$

By (4.20), we may take $\rho = (\max_{\|x\|=1} \|\nabla F(x)\|)^{-1}$.

- As noted, by compactness of \mathbb{S}^{L-1} , the sequence (x_k) has a cluster point.

Theorem 4.5 now follows from Theorem C.1. \square

Next we prove that SS-HOPM in general converges to second-order critical points for almost all initializations. In the language of [44], SS-HOPM converges to *stable eigenvectors*. While intuitive and essentially implicit in [44], this does require proof. We adapt the arguments in [62, 50] based on the center-stable manifold theorem from dynamical systems. It is convenient to set

$$G(x) = \frac{\nabla F(x)}{\|\nabla F(x)\|}, \quad (4.23)$$

and record a straightforward computation.

Lemma 4.6 *Let $x_* \in \mathbb{S}^{L-1}$ be a critical point of (4.11), with $\mu_* \in \mathbb{R}$ so that $\nabla f(x_*) + \mu_* x_* = 0$. Then the Jacobian $DG(x_*) \in \mathbb{R}^{L \times L}$ is symmetric, has the same eigenvectors as the Hessian $\nabla^2 f(x_*)$, and their eigenvalues are related as follows. If $y \in \mathbb{R}^L$ with $\langle y, x_* \rangle = 0$ satisfies $\nabla^2 f(x_*)y = \lambda y$, then $DG(x_*)y = \frac{\lambda + \gamma}{|-\mu_* + \gamma|} y$.*

Proof See Appendix C.3. \square

The following lemmas are crucial for proving SS-HOPM almost always converges to second-order critical point. The first one is a direct application of the center-stable manifold theorem, and the second one is technical lemma regarding the inverse image of a measure zero set.

Lemma 4.7 *Suppose that $x_* \in \mathbb{S}^{L-1}$ is a constrained critical point of (4.11) that fails the second-order criticality conditions. Then there exists an open neighborhood $B_{x_*} \subset \mathbb{S}^{L-1}$ of x_* , and a smoothly embedded disc $D_{x_*} \subset B_{x_*}$ of dimension strictly less than $L - 1$ such that:*

$$(x \in B_{x_*} \text{ and } G^k(x) \in B_{x_*} \forall k \geq 1) \Rightarrow x \in D_{x_*}. \quad (4.24)$$

Proof See Appendix C.5. \square

Lemma 4.8 *The inverse image of a measure zero subset of \mathbb{S}^{L-1} under $G|_{\mathbb{S}^{L-1}}$ is also measure zero.*

Proof See Appendix C.6. \square

We finally show that SS-HOPM converges to a second-order critical point almost always.

Theorem 4.9 (Almost always convergence of SS-HOPM to second-order critical point) *As above, let f be any homogeneous polynomial function on \mathbb{R}^L of degree $2n$. Let $\gamma \in \mathbb{R}$ be any shift such that $F(x) = f(x) + \frac{\gamma}{2n}(x^T x)^n$ is convex on \mathbb{R}^n and strictly positive on $\mathbb{R}^L \setminus \{0\}$. Then there exists a full-measure subset $\Omega \subset \mathbb{S}^{L-1}$ such that for all initializations $x_1 \in \Omega$, the iterates (4.12) converge to a second-order critical point x_* of (4.11).*

Proof By Theorem 4.5, for any initialization $x_1 \in \mathbb{S}^{L-1}$, the iterates $G^k(x_1)$ converge to some constrained critical point x_* of (4.11). Let $\mu_* \in \mathbb{R}$ be the corresponding Lagrange multiplier, so $\nabla f(x_*) + \mu_* x_* = 0$ (4.25).

Now suppose x_* *fails* the second-order criticality conditions. Then by Lemma 4.7, there exists an open neighborhood $B_{x_*} \subset \mathbb{S}^{L-1}$ of x_* , and a smoothly embedded disc $D_{x_*} \subset B_{x_*}$ of dimension strictly less than $L-1$ such that:

$$(x \in B_{x_*} \text{ and } G^k(x) \in B_{x_*} \forall k \geq 1) \Rightarrow x \in D_{x_*}. \quad (4.26)$$

Since $x_k \rightarrow x_*$, there is K such that $x_k \in B_{x_*}$ for all $k \geq K$. So $x_K \in D_{x_*}$ and $x_1 \in G|^{-K}(D_{x_*}) \subset \mathbb{S}^{L-1}$ where $G|^{-K}$ denotes K -fold inverse image under $G| = G|_{\mathbb{S}^{L-1}}$ (G restricted to the unit sphere). In particular,

$$x_1 \in \bigcup_{k \geq 1} G|^{-k}(D_{x_*}). \quad (4.27)$$

Since D_{x_*} is a manifold of dimension $< L-1$, it has measure zero (with respect to the usual surface measure on \mathbb{S}^{L-1}). Furthermore, Lemma 4.8 implies that the inverse image of a measure zero subset of \mathbb{S}^{L-1} under G is also measure zero. So the RHS of (4.27) is a countable union of measure zero subsets, so has measure zero. We have shown the set of initializations that converge to a *particular* critical point, failing second-order conditions, is a null set of \mathbb{S}^{L-1} .

We finish the argument with a union bound. Let \mathcal{Z} be the set of all “bad” critical points, that is, first-order critical points but not second-order critical points (this is possibly an uncountable set). For each such x_* , let B_{x_*} and D_{x_*} be as above. Consider the following open subset of \mathbb{S}^{L-1} :

$$\mathcal{B} = \bigcup_{x_* \in \mathcal{Z}} B_{x_*} \quad (4.28)$$

where the union is over all “bad” critical points $x_* \in \mathcal{Z}$. This means that $\{B_{x_*}\}_{x_* \in \mathcal{Z}}$ is an open cover of \mathcal{B} . Since \mathbb{S}^{L-1} is second-countable, there exists a countable subcover of \mathcal{B} , that is, there exists a *countable* set $\mathcal{C} \subset \mathbb{S}^{L-1}$ of “bad” critical points such that (4.28) equals the union just over \mathcal{C} :

$$\mathcal{B} = \bigcup_{x_* \in \mathcal{C}} B_{x_*}. \quad (4.29)$$

If $x_k \rightarrow x_{\dagger}$ for some point $x_{\dagger} \in \mathcal{Z}$, then there is a point $x_* \in \mathcal{C}$ such that $x_{\dagger} \in B_{x_*}$. Since $x_k \rightarrow x_{\dagger}$ and $x_{\dagger} \in B_{x_*}$, there exists K such that $x_k \in B_{x_*}$ for all $k \geq K$ and

$$x_1 \in \bigcup_{k \geq 1} G|^{-k}(D_{x_*}). \quad (4.30)$$

We conclude that any initialization x_1 such that (4.12) does not converge to a second-order critical point must lie in

$$\bigcup_{x_* \in \mathcal{C}} \bigcup_{k \geq 1} G|^{-k}(D_{x_*}), \quad (4.31)$$

which is a countable union of null sets. This completes the proof. \square

4.3 Local linear convergence of POWER METHOD to $\pm a_i$

Here we show the third part of Theorem 4.1, starting first with an easy lemma and then an essentially geometric proposition.

Lemma 4.10 *Under the assumptions of Theorem 4.9, suppose $x_* \in \mathbb{S}^{L-1}$ is a strict second-order critical point of (4.11), that is, where $\mu_* \in \mathbb{R}$ is as in (4.25), we have*

$$P_{(x_*)^\perp} (\nabla^2 f(x_*) + \mu_* I) P_{(x_*)^\perp}^T \prec 0, \quad (4.32)$$

where $P_{(x_*)^\perp} \in \mathbb{R}^{(L-1) \times L}$ represents orthogonal projection onto the perpendicular space $(x_*)^\perp \subset \mathbb{R}^L$. Then, (4.12) has local linear convergence to x_* .

Proof See Appendix C.7. \square

Proposition 4.11 *Let $R \leq \binom{L+n-1}{n} - n + 1$. For generic $a_1, \dots, a_R \in \mathbb{S}^{L-1}$ with $\mathcal{A} = \text{Span}\{a_1^{\otimes n}, \dots, a_R^{\otimes n}\} \subset \text{Sym } \mathcal{T}_L^n$, the following holds: whenever $x \in \mathbb{R}^L$ is nonzero and perpendicular to a_i , then $\text{Sym}(a_i^{\otimes n-1} \otimes x) \notin \mathcal{A}$.*

Proposition 4.11 may be interpreted as a statement in projective algebraic geometry about Zariski tangent spaces. We give the proof in Appendix C.8.

Theorem 4.12 *Let $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$, where $\mathcal{A} = \text{Span}\{a_1^{\otimes n}, \dots, a_R^{\otimes n}\}$, $a_1, \dots, a_R \in \mathbb{S}^{L-1}$ and $P_{\mathcal{A}} : \text{Sym } \mathcal{T}_L^n \rightarrow \mathcal{A}$ is orthogonal projection, as in Theorem 4.1. Fix $\gamma \in \mathbb{R}$ satisfying (4.3). Assume a_1, \dots, a_R are generic. For $R \leq \binom{L+n-1}{n} - L$, the only constrained global maxima of (4.1) are $\pm a_i$. For $R \leq \binom{L+n-1}{n} - n + 1$, the iteration (4.12) has local linear convergence to $\pm a_i$.*

Proof Since $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2 \leq \|x^{\otimes n}\|^2 = \|x\|^{2n}$ and $f(\pm a_i) = 1$, evidently $\pm a_i$ are global maxima of (4.1). By Proposition 3.2, generically there are no other global maxima provided $R \leq \binom{L+n-1}{n} - L$.

For local linear convergence, firstly $\pm a_i$ satisfy the first and second-order conditions in Remark 4.3 by global optimality, where $\mu_* = -2nf(\pm a_i) = -2n$. By Lemma 4.10, it suffices to show strictness of the second-order condition (4.32). Thus for each fixed a_i , it is enough to verify:

$$x^T \left(\frac{1}{2n} \nabla^2 f(a_i) - I \right) x < 0 \quad \text{for all nonzero } x \perp a_i. \quad (4.33)$$

(The result for $-a_i$ will follow by evenness of f .) As in (C.2), we compute:

$$\begin{aligned} \frac{1}{2n} x^T \nabla^2 f(a_i) x &= n \|P_{\mathcal{A}}(a_i^{\otimes n-1} \otimes x)\|^2 + (n-1) \langle a_i^{\otimes n}, a_i^{\otimes n-2} \otimes x^{\otimes 2} \rangle \\ &= n \|P_{\mathcal{A}}(a_i^{\otimes n-1} \otimes x)\|^2 \\ &= n \|P_{\mathcal{A}}(\text{Sym}(a_i^{\otimes n-1} \otimes x))\|^2. \end{aligned} \quad (4.34)$$

On the other hand, $\|\text{Sym}(a_i^{\otimes n-1} \otimes x)\| = \frac{1}{n} \|x\|^2$. To see this, we note

$$\text{Sym}(a_i^{\otimes n-1} \otimes x) = \frac{1}{n} \sum_{n_1+n_2=n-1} a_i^{\otimes n_1} \otimes x \otimes a_i^{\otimes n_2}, \quad (4.35)$$

where the summands are pairwise orthogonal in $\text{Sym } \mathcal{T}_L^n$ (using $x \perp a_i$) and each has norm $\|x\|$ (using $\|a_i\| = 1$). From this and (4.34), we obtain:

$$\begin{aligned} x^T \left(\frac{1}{2n} \nabla^2 f(a_i) - I \right) x &= n \left(\|P_{\mathcal{A}}(\text{Sym}(a_i^{\otimes n-1} \otimes x))\|^2 - \frac{\|x\|^2}{n} \right) \\ &= n \left(\|P_{\mathcal{A}}(\text{Sym}(a_i^{\otimes n-1} \otimes x))\|^2 - \|\text{Sym}(a_i^{\otimes n-1} \otimes x)\|^2 \right) \\ &= -n \|P_{\mathcal{A}^\perp}(\text{Sym}(a_i^{\otimes n-1} \otimes x))\|^2. \end{aligned} \quad (4.36)$$

Here $P_{\mathcal{A}^\perp} : \text{Sym } \mathcal{T}_L^n \rightarrow \mathcal{A}^\perp$ denotes the orthogonal projection onto the orthogonal complement $\mathcal{A}^\perp \subset \text{Sym } \mathcal{T}_L^n$ to $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$. However, by Proposition 4.11 below, we have that the RHS in Eqn (4.36) is strictly negative for all $x \perp a_i$, if $R \leq \binom{L+n-1}{n} - n + 1$ and a_1, \dots, a_R are generic. That completes the proof. \square

4.4 Putting it all together

To sum up, Theorem 4.1 is now proven. The first part follows from Lemma 4.4 and Theorem 4.5 applied to $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$. The second part similarly follows from Theorem 4.9, noting that positivity obviously holds for $F(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2 + \frac{\gamma}{2n}(x^T x)^n$. Finally, the third part exactly is Theorem 4.12.

Remark 4.13 As an aside, we discuss the possibility of using *adaptive* shifts in SPM. Similarly to [47], the idea would be to use smaller shifts at x_k according to how close $f(x)$ is to being locally convex around x_k . To this end, Lemma 4.4 suggests making POWER METHOD, governed by (4.2), adaptive as follows:

$$x_{k+1} = \frac{P_{\mathcal{A}}(x_k^{\otimes n}) \cdot x_k^{\otimes(n-1)} + \gamma_k x_k}{\|P_{\mathcal{A}}(x_k^{\otimes n}) \cdot x_k^{\otimes(n-1)} + \gamma_k x_k\|}, \quad \text{where } \gamma_k = C_{f(x_k)} C_n. \quad (4.37)$$

For this adaptive POWER METHOD, we are able to easily modify the proofs of Theorem 4.5 about unconditional global convergence and also Theorem 4.12 about linear local convergence to $\pm a_i$. However, we have been unable to prove avoidance of saddles, that is, an analog to Theorem 4.9. Time-inhomogeneous variants of the center-stable manifold theorem might resolve this, as in [63].

4.5 Optimization landscape

Now that we have established that POWER METHOD converges to a local maximum of (4.1), it is important to understand if there are local maxima which are not global maxima, and when the power method iterations converge to these critical points. Note that when the algorithm converges to a local maximum $x_* \in \mathbb{S}^{L-1}$, we know it is a global maximum if and only if $f(x_*) = 1$. By Proposition 3.2, we know up to which rank, it holds that the global maxima of (4.1) are generically precisely the desired tensor components (up to sign).

To understand the *optimization landscape*, we have devised the following experiment, depicted in Figure 4.1. We considered fourth-order tensors ($n =$

2), fixed the length $L = 20$ and varied the rank R from 120 to 200. Note that here the threshold given by Proposition 3.2 is $R = \binom{21}{2} = 190$. For each rank, we sample R i.i.d. standard Gaussian vectors $\{a_i\}_{i=1}^R$ and calculate the corresponding $\mathcal{A} = \text{span}\{a_i, i = 1, \dots, R\}$. Then we sample another vector x_0 uniformly in \mathbb{S}^{L-1} and perform the power method, starting at x_0 , until convergence. We repeat this experiment 10000 times, register when the power method converged (up to sign) to one of the components in $\{a_i\}_{i=1}^R$, and report the relative frequency of success. A value of 1 means that the power method with a random starting point converged always to one of the components $\{a_i\}_{i=1}^R$, and a value of 0 means that either the power method did not converge, got stuck on a saddle point or converged to other local maxima.

In the experiment, for all values of R smaller than 140, the algorithm always converged to one of the components. Between 140 to 160 the frequency is at least 99.8%, and decreases to 98.5% at $R = 170$. Then, we observe a sharp transition when the rank varies between 170 and 190. The width of the transition is $\approx L$. This suggests if the rank scales like $c_n L^n$, for a fixed constant $c_n < \frac{1}{n!}$ (since $\binom{L+n-1}{n} \approx \frac{1}{n!} L^n$), then the power method converges to a global maximum with high probability.

Conjecture 4.14 Let $\{a_i\}_{i=1}^R$ be R i.i.d. standard Gaussian vectors in \mathbb{R}^L , $c_n < \frac{1}{n!}$ a constant and consider the power method iterations given by (4.2), with x_1 drawn uniformly from \mathbb{S}^{L-1} . Then if $R < c_n L^n$, the power method iterations converge to $\pm a_j$ for some j in $\{1, \dots, R\}$ with high probability, i.e., with probability converging to 1 as L diverges to infinity.

We have repeated the experiment drawing $\{a_i\}_{i=1}^R$ from distributions other than the isotropic Gaussian, and the results were identical.

We also understand the optimization landscape for some values of L and R . We have found an example of a tensor of dimension 4, order 4 and rank 5 for which the optimization landscape contains a local maximum which is not global. We can easily extend this example to show that there can be bad local




Fig. 4.1: Relative frequency of convergence to one of the components $\{a_i\}_{i=1}^R$. The dashed line is the threshold given by Proposition 3.2.

maxima if the order is 4 and $R > L$. On the other hand, we conjecture that if $R \leq L$, there are no bad local maxima. We note that this conjecture can be reduced to the case $R = L$ by projecting the tensor to the space spanned by the $\{a_i\}_{i=1}^R$. The proof of this conjecture for $R = 1$ is trivial, and we sketch the details of the proof for $R = 2$ followingly, but the proof remains elusive for $R > 2$.

If $R = 2$, we can rescale and rotate a_1 and a_2 , without changing the optimization landscape, so that $a_1 = (\cos \theta, \sin \theta)$ and $a_2 = (\cos \theta, -\sin \theta)$ for some value $\theta \in (0, \frac{\pi}{2})$. One first checks the critical points are $\pm(\cos \theta, \sin \theta)$, $\pm(\cos \theta, -\sin \theta)$, $\pm(1, 0)$ and $\pm(0, 1)$, and by computing the Riemannian Hessian, one concludes the first four are global maxima and the last four are local minimum.

5 Symmetric block term decomposition

In this section, we describe how to adapt Algorithm 1 to compute the symmetric block term tensor decomposition in Definition 2.10. We begin with a lemma linking this tensor decomposition to generalized PCA.

Lemma 5.1 *Consider a subspace arrangement $\mathcal{S} = \cup_{i=1}^R S_i \subset \mathbb{R}^L$ where S_i is a linear subspace of dimension ℓ_i given by $S_i = \text{colspan}(A_i)$ for $A_i \in \mathbb{R}^{L \times \ell_i}$. Let $Y \in \mathbb{R}^L$ be a random variable supported on \mathcal{S} . For each m , the moment tensor $\mathbb{E}[Y^{\otimes m}] \in \text{Sym } \mathcal{T}_L^m$ admits a symmetric block term decomposition:*

$$\mathbb{E}[Y^{\otimes m}] = \sum_{i=1}^R (A_i; \dots; A_i) \cdot \mathbf{A}_i, \quad (5.1)$$

for some $\mathbf{A}_i \in \text{Sym } \mathcal{T}_{\ell_i}^m$.

Proof We may write $Y = A_\kappa y_\kappa$ where $\kappa \in [R]$ is a random discrete variable with $\text{Prob}(\kappa = i) = p_i$ for (p_1, \dots, p_R) in the probability simplex, and where $y_i \in \mathbb{R}^{\ell_i}$ is a random variable for each $i \in [R]$. Then, by linearity of expectation:

$$\mathbb{E}[Y^{\otimes m}] = \sum_{i=1}^R p_i \mathbb{E}[(A_i y_i)^{\otimes m}] = \sum_{i=1}^R p_i (A_i; \dots; A_i) \cdot \mathbb{E}[y_i^{\otimes m}]. \quad (5.2)$$

So we may set $\mathbf{A}_i = p_i \mathbb{E}[y_i^{\otimes m}]$. \square

Lemma 5.1 is one reason to develop a method for computing symmetric block term decompositions. If in the setting of GPCA, we apply the method to the empirical moments of a dataset of i.i.d. draws of Y , then we can obtain an estimate to A_i (up to $\text{GL}(\ell_i, \mathbb{R})$), and whence $S_i = \text{colspan}(A_i)$. If in GPCA, the data points are subject to additive noise, upon debiasing the empirical moments, we can still reduce to the symmetric block term decomposition in Lemma 5.1 (see Subsection 6.5 and Appendix E). Other applications of (2.12) exist, e.g., variations on blind source separation, but we omit these here.

As for usual symmetric rank decomposition in Section 3, it is easier to describe SPM for the symmetric block term decomposition in the case of even order $m = 2n$. Thus, assume we have a tensor $T \in \text{Sym } \mathcal{T}_L^{2n}$ satisfying:

$$T = \sum_{i=1}^R (A_i; \dots; A_i) \cdot \mathbf{A}_i, \quad (5.3)$$

where as above $\mathbf{A}_i \in \text{Sym } \mathcal{T}_{\ell_i}^{2n}$ and $A_i \in \mathbb{R}^{L \times \ell_i}$. We assume A_i is full-rank, set $S_i = \text{colspan}(A_i) \subset \mathbb{R}^L$ and $\mathcal{S} = \cup_{i=1}^R S_i$, however, we do not assume knowledge of R or ℓ_i . Given T , the goal is to recover (\mathbf{A}_i, A_i) for $i = 1, \dots, R$ up to permutation of summands and change of basis in \mathbb{R}^{ℓ_i} , that is, the equivalence:

$$(\mathbf{A}_i, A_i) \sim (\mathbf{A}_i \times Q_i^{\times 2n}, A_i Q_i^{-1}), \quad \text{for } Q_i \in \text{GL}(\ell_i, \mathbb{R}). \quad (5.4)$$

To this end, we generalize SPM. We begin by adapting (3.2), to generalize EXTRACT SUBSPACE. Let $A_i^{\star n}$ denote the $L^n \times \binom{\ell_i+n-1}{n}$ matrix whose columns are given by all combinations $\text{vec}(\text{Sym}(a_{ij_1} \otimes \dots \otimes a_{ij_n}))$ where $1 \leq j_1 \leq \dots \leq j_n \leq \ell_i$ and $a_{ij} \in \mathbb{R}^L$ are the columns of A_i (and these combinations appear lexicographically in $A_i^{\star n}$). From the symmetric tensor \mathbf{A}_i in $\mathcal{T}_{\ell_i}^{2n}$ there is an injective linear map to a $\binom{\ell_i+n-1}{n} \times \binom{\ell_i+n-1}{n}$ symmetric matrix Λ_i such that

$$\text{mat}((A_i; \dots; A_i) \cdot \mathbf{A}_i) = A_i^{\star n} \Lambda_i A_i^{\star n}, \quad (5.5)$$

where mat is the flattening operator in Definition 2.7. Then (5.3) gives:

$$\text{mat}(T) = [A_1^{\star n} \dots A_R^{\star n}] \begin{bmatrix} \Lambda_1 & & & \\ & \Lambda_2 & & \\ & & \ddots & \\ & & & \Lambda_R \end{bmatrix} [A_1^{\star n} \dots A_R^{\star n}]^T. \quad (5.6)$$

This is a block matrix factorization of $\text{mat}(T)$ corresponding to (3.2). It is convenient to have shorthand for the left factor, of size $L^n \times \sum_{i=1}^R \binom{\ell_i+n-1}{n}$:

$$\mathbf{A} = [A_1^{\star n} \dots A_R^{\star n}]. \quad (5.7)$$

Generalizing (3.3), we set $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ as the column space of \mathbf{A} after unvectorization. We may also describe \mathcal{A} using the Veronese embedding [79].

Lemma 5.2 *We have $\mathcal{A} = \text{Span}\left(\cup_{i=1}^R \{a^{\otimes n} : a \in S_i\}\right) \subset \text{Sym } \mathcal{T}_L^n$. In equivalent geometric language, \mathcal{A} is the linear span of the subspace arrangement $\mathcal{S} = \cup_{i=1}^R S_i$ following re-embedding by the Veronese map $\nu_n : \mathbb{R}^L \rightarrow \text{Sym } \mathcal{T}_L^n$.*

Proof The second sentence is a restatement of the first because the Veronese map ν_n sends $a \mapsto a^{\otimes n}$. For the first sentence, it suffices to fix i and check

$\text{unvec}(\text{colspan}(A_i^{\star n})) = \text{Span}\{a^{\otimes n} : a \in S_i\}$. For ‘ \supset ’, take $a \in S_i$ and write $a = \sum_{j=1}^{\ell_i} \alpha_j a_{ij}$ where $a_{ij} \in \mathbb{R}^L$ are the columns of A_i and $\alpha_j \in \mathbb{R}$. Then

$$\begin{aligned} a^{\otimes n} &= \text{Sym}(a^{\otimes n}) = \text{Sym}\left(\sum_{j=1}^{\ell_i} \alpha_j a_{ij}\right)^{\otimes n} = \\ &\sum_{j_1, \dots, j_n=1}^{\ell_i} \alpha_{j_1} \dots \alpha_{j_n} \text{Sym}(a_{j_1} \otimes \dots \otimes a_{j_n}) \in \text{unvec}(\text{colspan}(A_i^{\star n})). \end{aligned} \quad (5.8)$$

The reverse inclusion is seen using homogeneous polynomials (Lemma 2.2). Via Φ , the LHS corresponds to $\mathbb{R}[a_{i1}^T X, \dots, a_{i\ell_i}^T X]_n \subset \mathbb{R}[X_1, \dots, X_L]_n$, i.e., degree- n polynomials in the “latent variables” $a_{i1}^T X, \dots, a_{i\ell_i}^T X$. Meanwhile, the RHS corresponds to $\text{Span}\{(a^T X)^n : a \in \text{colspan}(A)\}$. So, ‘ \subset ’ reduces to the fact that any polynomial may be written as a sum of powers of linear forms. \square

SPM for the block term decomposition relies on the subspace $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ given in Lemma 5.2. We proceed, similarly to Algorithm 1, by obtaining a thin eigendecomposition of $\text{mat}(T)$ to extract \mathcal{A} . The next result is analogous to Proposition 3.1, except for the more general case of symmetric block term decompositions rather than symmetric rank decompositions.

Proposition 5.3 *Let \mathbf{A} have rank \mathbf{R} , so $\mathbf{R} \leq \sum_{i=1}^R \binom{\ell_i+n-1}{n}$. Assume $\mathbf{A}_i \in \text{Sym } \mathcal{T}_{\ell_i}^{2n}$ are generic. Then the rank of $\text{mat}(T)$ is also \mathbf{R} . Furthermore, if $\text{mat}(T) = V D V^T$ is a thin eigendecomposition of $\text{mat}(T)$, that is, $V \in \mathbb{R}^{L^n \times \mathbf{R}}$ has orthonormal columns and D is full-rank diagonal, then the columns of V give an orthonormal basis for \mathcal{A} (after un-vectorization).*

Proof Certainly, $\mathbf{R} \leq \sum_{i=1}^R \binom{\ell_i+n-1}{n}$ as the RHS is the number of columns in \mathbf{A} . For each i , fix generic vectors $b_1, \dots, b_{\binom{\ell_i+n-1}{n}}$ in $\text{colspan}(A_i) \subset \mathbb{R}^L$ and let B_i be the matrix with these as columns. Consider the Khatri-Rao $B_i^{\bullet n} \in \mathbb{R}^{L^n \times \binom{\ell_i+n-1}{n}}$ (Definition 2.8). Similarly to Lemma 5.2, we can argue that $B_i^{\bullet n}$ and $A_i^{\star n}$ have the same column space. So there exists $\mathbf{M}_i \in \text{GL}(\binom{\ell_i+n-1}{n}, \mathbb{R})$ such that $A_i^{\star n} = B_i^{\bullet n} \mathbf{M}_i$. Substituting into (5.6) gives:

$$\text{mat}(T) = [B_1^{\bullet n} \dots B_R^{\bullet n}] \begin{bmatrix} \mathbf{M}_1 A_1 \mathbf{M}_1^T & & & \\ & \mathbf{M}_2 A_2 \mathbf{M}_2^T & & \\ & & \ddots & \\ & & & \mathbf{M}_R A_R \mathbf{M}_R^T \end{bmatrix} \begin{bmatrix} (B_1^{\bullet n})^T \\ \vdots \\ (B_R^{\bullet n})^T \end{bmatrix} \quad (5.9)$$

We want to show that $\text{mat}(T)$ and $[B_1^{\bullet n} \dots B_R^{\bullet n}]$ have the same column space, for generic \mathbf{A}_i . Since the condition is Zariski-open (characterized by polynomials), it is enough to see there exists some such \mathbf{A}_i . To this end, consider \mathbf{A}_i so that $\mathbf{M}_i A_i \mathbf{M}_i^T$ are generic diagonal matrices. Then the argument of Proposition 3.1 applies nearly identically (genericity, instead of nonzeroness, of the diagonal matrices is needed in the case $\mathbf{R} < \sum_{i=1}^R \binom{\ell_i+n-1}{n}$). That gives $\text{rank}(\text{mat}(T)) = \mathbf{R}$ as well as the statement about the columns of V and \mathcal{A} . \square

Remark 5.4 Proposition 5.3 is precisely where our GPCA method relies on the assumption that the ground-truth probability distribution supported on the subspace arrangement \mathcal{S} is *generic*. Specifically, we require that the moment tensors \mathbf{A}_i be Zariski-generic in $\text{Sym } \mathcal{T}_{\ell_i}^{2n}$ in order to be able to correctly extract the subspace \mathcal{A} corresponding to \mathcal{S} . While it is true that several distinct subspace arrangements could give rise to the same moment tensors if their probability distributions are chosen appropriately, our results prove that only one subspace arrangement \mathcal{S} equipped with a generic probability distribution (subject to being supported on \mathcal{S}) can be consistent with the moment tensors.

Remark 5.5 For the symmetric block term decomposition, we call \mathbf{R} the *square flattening rank* of T , and the quantity $\min\left(\binom{L+n-1}{n}, \sum_{i=1}^R \binom{\ell_i+n-1}{n}\right)$ the *expected square flattening rank*. The latter is unknown to us, except in synthetic experiments, as we do not assume knowledge of R and ℓ_1, \dots, ℓ_R .

Thus SPM again extracts $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ from an eigendecomposition of $\text{mat}(T)$.

The next lemma is a simple equivalence, related to Proposition 3.2. It re-expresses the condition that $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$ has no extraneous rank-1 points in terms of algebraic geometry language, connecting with existing algebraic literature on subspace arrangements and GPCA, for example, [30, 14].

Lemma 5.6 *Suppose the subspace arrangement $\mathcal{S} = \cup_{i=1}^R S_i \subset \mathbb{R}^L$ is set-theoretically defined by degree- n polynomial equations. This means that if we let $I_n(\mathcal{S}) = \{g \in \mathbb{R}[X_1, \dots, X_L] : g(x) = 0 \text{ for all } x \in \mathcal{S}\}$ be set of degree- n equations of \mathcal{S} , then these equations exactly characterize \mathcal{S} , i.e., the set of common zeroes $Z(I_d(\mathcal{S})) = \{x \in \mathbb{R}^L : g(x) = 0 \text{ for all } g \in I_n(\mathcal{S})\}$ equals \mathcal{S} . In this case, the only rank-1 tensors in \mathcal{A} are $\cup_{i=1}^R \{a^{\otimes n} : a \in S_i\}$ (up to sign if n is even). Moreover, the converse holds.*

Proof This follows from the identification $\Phi(I_n(\mathcal{S})) = \mathcal{A}^\perp \subset \text{Sym } \mathcal{T}_n^L$, where \mathcal{A}^\perp is the orthogonal complement to \mathcal{A} and Φ is as in Lemma 2.2. \square

Remark 5.7 Given $\mathcal{S} = \cup_{i=1}^R S_i \subset \mathbb{R}^L$ and a degree n , it is subtle to decide if the condition in Lemma 5.6 holds, i.e., if \mathcal{A} contains any extraneous rank-1 tensors. The challenge persists even if $S_i \subset \mathbb{R}^L$ are generic subspaces, subject to $\dim(S_i) = \ell_i$ for fixed ℓ_1, \dots, ℓ_R . An exception is the case $\ell_i = 1$ for all $i = 1, \dots, R$, which corresponds to usual tensor decomposition; then the trisecant lemma implies extraneous rank-1 tensors generically do not lie in \mathcal{A} up to an explicit R (Proposition 3.2). Generally, for any fixed arrangement \mathcal{S} , the condition in Lemma 5.6 holds for all sufficiently large n . As a pessimistic general bound, if $n \geq R$ then \mathcal{A} cannot have extraneous rank-1 points by [31].

Practically speaking, in light of Lemma 5.6, Remark 5.7 and Proposition 5.10 below, when applying generalized SPM to GPCA where the subspace arrangement \mathcal{S} is fixed but unknown, the sufficient conditions on \mathcal{S} and n for our guarantees cannot be verified a priori. So, it is reasonable to try a *degree-increasing approach*. That is, we apply SPM to the degree- $2n$ moments of the dataset for $n = 2, 3, \dots$, until the method does not fail and so \mathcal{S} is recovered.

Continuing the algorithm description for generalized SPM, now that we have an orthonormal basis for \mathcal{A} , we seek a rank-1 element in \mathcal{A} . To that end, as in Algorithm 1, we apply POWER METHOD, that is, we iterate (3.10) where γ is given by (4.3) (recall that Lemma 4.4 applied to *any* linear subspace $\mathcal{A} \subset \text{Sym } \mathcal{T}_L^n$). Here the first and second items in Theorem 4.1 apply, as Subsection 4.2 pertained to SS-HOPM and symmetric tensors in general. So POWER METHOD converges to a critical point $x_* \in \mathbb{S}^{L-1}$, and almost surely a local maximum, of $\max_{\|x\|=1} \|P_{\mathcal{A}}(x^{\otimes n})\|^2$. To see if x_* is a global maximum, we check if the objective function value at x_* is 1 (up to a numerical tolerance); if not, we discard x_* and repeat POWER METHOD with a fresh initialization.

The next step of generalized SPM departs from Algorithm 1. We assume x_* lies on exactly one S_i (the current algorithm may fail if the constituent subspaces in \mathcal{S} have nontrivial pairwise intersection and x_* lies on two of them). The next step, LOCAL COMPONENT, determines the subspace S_i containing x_* . For Algorithm 1, this is immediate, as usual tensor decomposition corresponds to (5.3) when each S_i is a line, so simply the span of $x_* \in S_i$. For the generalized case, we note that an open neighborhood of x_* in \mathcal{S} is an neighborhood of x_* in S_i . So, S_i is recovered by linearizing equations for \mathcal{S} around x_* , that is, by computing the tangent space to \mathcal{S} at x_* . To this end, we let $Q_1, \dots, Q_{\mathbf{R}^\perp} \in \text{Sym } \mathcal{T}_L^n$, where $\mathbf{R}^\perp = \binom{L+n-1}{n} - \mathbf{R}$, denote an orthonormal basis for \mathcal{A}^\perp (the orthogonal complement of \mathcal{A} in $\text{Sym } \mathcal{T}_L^n$). Then the equations

$$\langle Q_j, x^{\otimes n} \rangle = 0, \quad \text{for all } 1 \leq j \leq \mathbf{R}^\perp \quad (5.10)$$

span the vector space of all degree- n equations for \mathcal{S} . We try to determine the column span of A_i by looking at the null space of the Jacobian of these equations around $x = x^*$. Explicitly, we set

$$A_i \leftarrow \text{nullspace}(\mathbf{Q}), \quad (5.11)$$

where \mathbf{Q} is a $\mathbf{R}^\perp \times L$ matrix with rows given by

$$\mathbf{Q}_j = \nabla \langle Q_j, x^{\otimes n} \rangle_{x=x_*} = n \text{Sym}(Q_j) \cdot x_*^{\otimes n-1}, \quad (5.12)$$

and $\text{nullspace}(\mathbf{Q})$ is a matrix whose columns form a basis for the kernel of \mathbf{Q} .

Remark 5.8 Equations (5.11) and (5.12) correctly recover the column space of A_i provided the tangent space to \mathcal{S} at x_* is determined by differentiating *just* degree- n equations. See (5.16) in Proposition 5.10 below. Similarly to Remark 5.7, for any fixed arrangement \mathcal{S} , this condition holds at all points x_* in \mathcal{S} for sufficiently large degrees n . Again by [31], $n \geq R$ is sufficient but pessimistic.

Remark 5.9 Computationally, we calculate the null space of \mathbf{Q} using the matrix $V \in \mathbb{R}^{L^n \times \mathbf{R}}$ in Proposition 5.3, whose columns are vectorizations of an orthonormal basis of tensors spanning \mathcal{A} . This procedure is particularly efficient as \mathbf{R} decreases, and \mathbf{R}^\perp increases. We explain the details in Appendix D.

Algorithm 2 Generalized Subspace Power Method

Input: $T \in \text{Sym } \mathcal{T}_L^{2n}$ satisfying (2.12) and conditions of Lemma 5.6, Proposition 5.10
Output: Dimensions (ℓ_1, \dots, ℓ_R) and symmetric block term decomposition $\{(A_i, \mathbf{A}_i)\}_{i=1}^R$.

$(V, D) \leftarrow \text{eig}(\text{mat}(T))$ EXTRACT SUBSPACE
 $\mathbf{R} \leftarrow \text{rank}(\text{mat}(T))$
 $\mathbf{R}^\perp \leftarrow \binom{L+n-1}{n} - \mathbf{R}$
 $i \leftarrow 0$
while $\mathbf{R} > 0$ **do**
 $i \leftarrow i + 1$
 $x \leftarrow \text{random}(\mathbb{S}^{L-1})$ POWER METHOD
 repeat
 $x \leftarrow \frac{P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes n-1} + \gamma x}{\|P_{\mathcal{A}}(x^{\otimes n}) \cdot x^{\otimes n-1} + \gamma x\|}$
 until convergence
 $x^* \leftarrow x$

 $Q_1, \dots, Q_{\mathbf{R}^\perp} \leftarrow$ orthonormal basis of \mathcal{A}^\perp (obtained from V) LOCAL COMPONENT
 for $j = 1$ **to** \mathbf{R}^\perp **do**
 $\mathbf{Q}_j \leftarrow \text{Sym}(Q_j) \cdot x_*^{\otimes n-1}$
 $A_i \leftarrow \text{nullspace}(\mathbf{Q})$
 $\ell_i \leftarrow \# \text{columns of } A_i$

 construct A_i^{*n} from A_i DEFLATE
 $\alpha \leftarrow V^T A_i^{*n}$
 $\Lambda_i \leftarrow (\alpha^T D^{-1} \alpha)^{-1}$
 construct \mathbf{A}_i from Λ_i
 $(O, \tilde{D}) \leftarrow \text{eig}(D - \alpha A_i \alpha^T)$
 $V \leftarrow VO$
 $D \leftarrow \tilde{D}$
 $\mathbf{R} \leftarrow \mathbf{R} - \binom{\ell_i+n-1}{n}$

 $R \leftarrow i$
return $\{(A_i, \mathbf{A}_i)\}_{i=1}^R$

Finally comes DEFLATE, in which the summand of (2.12) corresponding to the subspace S_i is removed. With A_i in hand, we construct the matrix $A_i^{*n} \in \mathbb{R}^{L^n \times \binom{\ell_i+n-1}{n}}$, with columns given by the combinations $\text{Sym}(a_{ij_1} \otimes \dots \otimes a_{ij_n})$, where $1 \leq j_1 \leq \dots \leq j_n \leq \ell_i$ and $a_{ij} \in \mathbb{R}^L$ are the columns of A_i . Similarly to Algorithm 1, we consider updates of the form $\text{mat}(T) - A_i^{*n} W (A_i^{*n})^T$, where W is a $\binom{\ell_i+n-1}{n} \times \binom{\ell_i+n-1}{n}$ matrix. On one hand, by (5.6),

$$\text{mat}(T) - A_i^{*n} W (A_i^{*n})^T = A_i^{*n} (\Lambda_i - W) (A_i^{*n})^T + \sum_{j \neq i} A_j^{*n} \Lambda_j (A_j^{*n})^T, \quad (5.13)$$

while on the other hand,

$$\text{mat}(T) - A_i^{*n} W (A_i^{*n})^T = V(D - \alpha W \alpha^T) V^T, \quad (5.14)$$

where $\alpha = V^T A_i^{*n}$. As before, we proceed by applying Lemma B.1, now with $k = \binom{\ell_i+n-1}{n}$, to set $\Lambda_i \leftarrow (\alpha^T D^{-1} \alpha)^{-1}$. We then obtain \mathbf{A}_i from Λ_i by inverting the map used in (5.5). Finally, we update $V \leftarrow VO$ and $D \leftarrow \tilde{D}$, where (O, \tilde{D}) is the thin eigendecomposition of $D - \alpha A_i \alpha^T$. POWER METHOD, LOCAL

COMPONENT and DEFLATE repeat as subroutines, such that each block term component (A_i, A_i) is removed one at a time, until all components have been found. An overview of SPM for symmetric block term tensor decomposition is presented in Algorithm 2.

The last result this section is an analog to Theorem 4.12. For simplicity, we assume the subspace arrangement $\mathcal{S} = \cup_{i=1}^R S_i$ satisfies $S_i \cap S_j = 0$ for all $i \neq j$. We show, under certain additional geometric conditions on \mathcal{S} , each $S_i \cap \mathbb{S}^{L-1}$ is an attractive set for POWER METHOD enjoying local linear convergence.

Proposition 5.10 *Suppose $\mathcal{S} = \cup_{i=1}^R S_i \subset \mathbb{R}^L$ satisfies $S_i \cap S_j = 0$ for all $i \neq j$. For each i , let $P_{S_i^\perp} \in \mathbb{R}^{(L-\ell_i) \times L}$ represent orthogonal projection onto the orthogonal complement $S_i^\perp \subset \mathbb{R}^L$. As above, set $f(x) = \|P_{\mathcal{A}}(x^{\otimes n})\|^2$ where \mathcal{A} is given by the column space of (5.7). Assume for all $a \in S_i \cap \mathbb{S}^{L-1}$:*

$$P_{S_i^\perp} \left(\frac{1}{2n} \nabla^2 f(a) - I \right) \left(P_{S_i^\perp} \right)^T \prec 0. \quad (5.15)$$

Then, the iteration (4.12) has local linear convergence to the set $S_i \cap \mathbb{S}^{L-1}$. That is, there exist positive constants $\delta = \delta(\mathcal{S}, \gamma, i)$, $\tau = \tau(\mathcal{S}, \gamma, i)$ and $C = C(\mathcal{S}, \gamma, i)$ such that $\text{dist}(x_1, S_i \cap \mathbb{S}^{L-1}) < \delta$ implies $\text{dist}(x_k, S_i \cap \mathbb{S}^{L-1}) < Ce^{-k\tau}$ for all k , where $\text{dist}(x_1, S_i \cap \mathbb{S}^{L-1}) = \min_{a \in S_i \cap \mathbb{S}^{L-1}} \|x_1 - a\|$.

Here, (5.15) is equivalent to the condition that the degree- n equations of \mathcal{S} correctly determine the tangent space to \mathcal{S} at $a \in S_i \cap \mathbb{S}^{L-1}$. More precisely, the gradients of the degree- n equations of \mathcal{S} evaluated at a define S_i , i.e.,

$$S_i = \text{Span}\{\nabla g(a) \in \mathbb{R}^L : g \in \mathbb{R}[X_1, \dots, X_L]_n, g(x) = 0 \forall x \in \mathcal{S}\}^\perp. \quad (5.16)$$

Proof By Taylor's theorem and compactness, there exists a constant $M = M(\mathcal{S}, \gamma) > 0$ such that:

$$\|G(x) - G(y) - DG(y)(x - y)\| \leq M\|x - y\|^2 \quad \text{for all } x, y \in \mathbb{S}^{L-1}, \quad (5.17)$$

where G is the power iteration in (4.23).

Now fix i , let $x \in \mathbb{S}^{L-1}$, and write $x = x_\parallel + x_\perp$, where $x_\parallel \in S_i$ and $x_\perp \perp S_i$. Equivalently, we have $x_\parallel = \text{argmin}_{z \in S_i} \|x - z\|$. Moreover, let

$$y = \text{argmin}_{z \in S_i \cap \mathbb{S}^{L-1}} \|x - z\|, \quad (5.18)$$

thus $\text{dist}(x, S_i \cap \mathbb{S}^{L-1}) = \|x - y\|$. Since $y \in S_i$, we have

$$\|x_\perp\| = \|x - x_\parallel\| \leq \|x - y\|. \quad (5.19)$$

On the other hand, since G fixes $S_i \cap \mathbb{S}^{L-1}$ pointwise, $DG(y)(z) = 0$ for all $z \in S_i$. In particular, since DG is a linear map,

$$DG(y)(x - y) = DG(y)(x_\perp). \quad (5.20)$$

Using Lemma 4.6 and (5.15), similarly to (4.32) and the proof of Lemma 4.10, we get that S_i^\perp decomposes as a sum of eigenspaces of $DG(y)$ with eigenvalues all strictly less than 1 in absolute value. So by continuity of eigenvalues and

compactness, there exists a constant $\lambda_{\max} = \lambda_{\max}(\mathcal{S}, \gamma, i) \in (0, 1)$ such that for all $z \in S_i \cap \mathbb{S}^{L-1}$ each of eigenvalue of $P_{S_i^\perp} DG(z) \left(P_{S_i^\perp}\right)^T$ is at most λ_{\max} in absolute value.

We proceed by substituting (5.20) into (5.17), using $G(y) = y$, (5.19) and the triangle inequality:

$$\begin{aligned} \text{dist}(G(x), S_i \cap \mathbb{S}^{L-1}) &\leq \|G(x) - y\| \\ &\leq \|DG(y)(x_\perp)\| + M\|x - y\|^2 \\ &\leq \lambda_{\max}\|x_\perp\| + M\|x - y\|^2 \\ &\leq (\lambda_{\max} + M\|x - y\|)\|x - y\|. \end{aligned} \quad (5.21)$$

If $\|x - y\| = \text{dist}(x, S_i \cap \mathbb{S}^{L-1})$ is sufficiently small, then the bracketed quantity in (5.21) is less than and bounded away from 1. For the local linear convergence statement, we can take $\delta = (1 - \lambda_{\max}) / (2M)$ (and related expressions for τ and C).

For the next statement, as in the proof of Theorem 4.12, we have

$$x^T \left(\frac{1}{2n} \nabla^2 f(a) - I \right) x = -n \|P_{\mathcal{A}^\perp}(\text{Sym}(a^{\otimes n-1} \otimes x))\|^2, \quad (5.22)$$

for $a \in L_i$ and $x \perp S_i$, where \mathcal{A}^\perp is the orthogonal complement of \mathcal{A} in $\text{Sym } \mathcal{T}_L^n$. So (5.15) is equivalent to $P_{\mathcal{A}^\perp}(\text{Sym}(a^{\otimes n-1} \otimes x)) \neq 0$ for all $x \perp S_i$. On the other hand, consider the RHS of (5.16). As above in the explanation of LOCAL COMPONENT, this equals

$$\text{Span}\{Q \cdot a^{\otimes n-1} : Q \in \mathcal{A}^\perp\}^\perp, \quad (5.23)$$

which always contains S_i . So, (5.16) is equivalent to the intersection of (5.23) and S_i^\perp being zero, i.e., for all $x \perp S_i$, there exists $Q \in \mathcal{A}^\perp$ such that

$$0 \neq x \cdot (Q \cdot a^{\otimes n-1}) = \langle Q, a^{\otimes n-1} \otimes x \rangle = \langle Q, \text{Sym}(a^{\otimes n-1} \otimes x) \rangle. \quad (5.24)$$

Equivalently, $P_{\mathcal{A}^\perp}(\text{Sym}(a^{\otimes n-1} \otimes x)) \neq 0$ for all $x \perp S_i$. \square

In general, verifying the sufficient conditions in Proposition 5.10 is a subtle matter⁵, though they hold for any fixed \mathcal{S} and large enough degree n , pessimistically for $n \geq R$ by [31]. For the special case of usual tensor decompositions ($\ell_1 = \dots = \ell_R = 1$), we have better control over n by Proposition 4.11.

⁵ Interestingly, the question of whether quadratic equations ($n = 2$) correctly determine Zariski tangent spaces recently arose in a different context, see [6].

6 Numerical simulations

In this section, we perform various numerical tests of SPM and provide comparisons of runtime, accuracy and noise stability against existing state-of-the-art algorithms. For symmetric tensor rank decompositions, we compare against FOABI [26] (state-of-the-art among provable algorithms), Tensorlab [77] (state-of-the-art among heuristic algorithms) and the method described in the paper titled “Low rank symmetric tensor approximations” [60], which we abbreviate to LRSTA. For GPCA, we compare against PDA [79] and GPCA-Voting [82, 55], algebraic methods somewhat related to ours.

All of the numerical experiments were performed on a personal laptop with a Intel® Core™ i7-7700HQ CPU and 16.0GB of RAM. On our implementation of SPM, we have set the following procedures and hyper-parameters. The maximum number of power method iterations per component is 5000. We note that, by Theorem 4.1, if $R \leq \binom{L+n-1}{n} - L$, $f(x^*) = 1$ if and only if x^* is one of the components a_i ’s, thus we can use the function value as a criteria for repeating/restarting the power method. Therefore, in the current implementation of SPM, if the function value of x^* (obtained by the POWER METHOD) is smaller than $1 - 10^{-6}$, we repeat POWER METHOD step with a new random initialization. If after repeating 3 times the POWER METHOD step, the function value is still smaller than $1 - 10^{-6}$, we continue the algorithm by choosing which of the 3 points had higher function value.

6.1 Comparing SPM for different tensor decompositions

In this subsection, we test the performance of SPM for different tensors. The results are displayed in Table 6.1. We applied SPM to ten symmetric tensors $T_1, \dots, T_{10} \in \text{Sym } \mathcal{T}_L^m$ with order m and length L , generated as follows:

- For tensors T_1, \dots, T_5 and T_9 , we planted a low-rank rank decomposition (1.1) with L, m, R as in Table 6.1, by drawing $\{a_i\}_{i=1}^R$ independently and from the uniform distribution in the unit sphere in \mathbb{R}^L , and by drawing $\{\lambda_i\}_{i=1}^R$ independently from the standard Gaussian distribution. Notice T_9 is order-6. We then used SPM for computing rank decompositions (Algorithm 1).
 - For tensors T_6 and T_7 , we planted a low-rank rank decomposition (1.1) with correlated components. We drew $\{\lambda_i\}_{i=1}^R$ as above, but the vectors $\{a_i\}_{i=1}^R$ were drawn independently from a different distribution.
 - For T_6 , we sample a vector v from a Gaussian distribution with unit variance and mean $\mathbf{1}$ (the all-ones vector), and then let $a_i = v/\|v\|$. This implies that the average correlation between the vectors will be $1/2$.
 - For T_7 , we sample each $\{a_i\}_{i=1}^R$ from the uniform distribution in the positive orthant of the unit sphere in \mathbb{R}^L .
- Again, Algorithm 1 was used to recover $\{(\lambda_i, a_i)\}_{i=1}^R$.

T	m	L	R/\mathbf{R}	EXTRACT SUBSPACE time (s)	POWER METHOD time (s)	DEFLATE time (s)	Total time (s)	Avg. n° iter. / Restarts	Error
T_1	4	40	200	0.10	0.27	0.22	0.59	45 / 0	1.67×10^{-13}
T_2	4	40	400	0.09	1.52	0.92	2.53	71 / 0	5.13×10^{-13}
T_3	4	40	600	0.12	9.12	2.38	11.62	118 / 0	1.68×10^{-12}
T_4	4	60	400	0.67	3.09	1.68	5.44	41 / 0	2.93×10^{-13}
T_5	4	80	400	3.51	5.42	2.93	11.86	33 / 0	1.77×10^{-13}
T_6	4	40	200^C	0.11	1.23	0.21	1.55	214 / 0	1.14×10^{-12}
T_7	4	40	200^+	0.09	2.83	0.22	3.14	618 / 9	1.66×10^{-12}
T_8	4	40	180^G	0.10	0.04	0.08	0.22	42 / 0	1.43×10^{-12}
T_9	6	16	400	0.13	6.71	1.99	8.83	68 / 0	5.68×10^{-13}
T_{10}	6	16	112^G	0.13	0.02	0.06	0.22	33 / 0	7.14×10^{-12}

Table 6.1: SPM’s performance for different low-rank tensors. The symmetric tensors T_1, \dots, T_{10} are described in Section 6.1. Here, the exponent C indicates that the rank-1 components have positive correlation, $+$ indicates that the rank-1 components were drawn from the positive orthant, and G indicates that the tensors were planted with symmetric block term decompositions.

- For tensors T_8 and T_{10} , we planted symmetric block term decompositions (2.6), with L, m, \mathbf{R} as in Table 6.1, drawing each \mathbf{A}_i to be a symmetric tensor with standard Gaussian entries, and each matrix A_i with orthonormal columns so that their span is a subspace in \mathbb{R}^L that is uniformly random in the corresponding Grassmannian. Specifically, $T_8 \in \text{Sym } \mathcal{T}_{40}^4$ is a linear combination of $R = 40$ Tucker decompositions, 20 of dimension 3 and 20 of dimension 2, that is, $\ell_1 = \dots = \ell_{20} = 3$ and $\ell_{21} = \dots = \ell_{40} = 2$. The expected square flattening rank of T_8 is $20 \binom{3+1}{2} + 20 \binom{2+1}{2}$, and indeed we found this to be its square flattening rank, $\mathbf{R} = 180$ (Remark 5.5). Meanwhile, $T_{10} \in \text{Sym } \mathcal{T}_{16}^6$ is a linear combination of $R = 16$ Tucker decompositions, 8 of dimension 3 and 8 of dimension 2. The expected square flattening rank of T_* is $8 \binom{3+2}{3} + 8 \binom{2+2}{3}$, and we found this to be its square flattening rank, $\mathbf{R} = 112$. For tensors T_8 and T_{10} , Algorithm 2 was applied.

For each example tensor T_1, \dots, T_{10} , we have recorded the time per each step of SPM as well as the total time in Table 6.1 (we joined DEFLATE and LOCAL COMPONENT together in our accounting of Algorithm 2). We also recorded the average number of iterations of the power method and number of times the POWER METHOD step was restarted because the function value was less than $1 - 10^{-6}$. Finally, we report the error of the decomposition of the tensor T obtained by SPM, $(\hat{\mathbf{A}}_i, \hat{A}_i)_{i=1}^R$, calculated using the following formula:

$$\text{error} := \|T - \hat{T}\| \quad \text{where} \quad \hat{T} = \sum_{i=1}^R (\hat{A}_i; \dots; \hat{A}_i) \cdot \hat{\mathbf{A}}_i. \quad (6.1)$$

We followingly provide an analysis of these numerical results:

General comments: We first note that in most cases EXTRACT SUBSPACE is the fastest step in SPM (except for T_5 , T_8 and T_{10}). We believe this is due to only performing this operation once upfront, whereas POWER METHOD and DEFLATE are performed multiple times throughout SPM; also, the eigenvalue decomposition algorithm, on which EXTRACT SUBSPACE relies, is a highly optimized algorithm in MATLAB. On the other hand, the POWER METHOD takes the most time in most cases.

Increasing rank: By comparing the average number of iterations in T_1 , T_2 and T_3 , one observes that it increases with the rank. Interestingly, while the ranks of the tensors T_2 and T_4 and T_5 are the same, the average number of iterations for T_2 which has smaller dimension, is the highest. Based on empirical observations, the average number of iterations appears to be an increasing function of R/L^2 for tensors of order 4, where R is the rank and L is the length of the tensor. For tensor of even order $m = 2n$, we conjecture the average number of iterations depends on the rank and length through R/L^n .

Increasing dimension: When comparing T_2 , T_4 and T_5 , one observes that the time of the EXTRACT SUBSPACE step is increasing dramatically with the length of the vector. This happens because in this step we calculate a full eigenvector decomposition of a $L^n \times L^n$ matrix, which takes time $O(L^{3n})$. As L increases, calculating this decomposition becomes the major bottleneck of the algorithm for low rank tensors (the computational times for the POWER METHOD and DEFLATION steps are $O(R^2 L^n)$, respectively), and algorithms that calculate only the top eigenvalues (such as the routine `eigs` in MATLAB), should be used.

Correlated components: Table 7.1 shows SPM taking more iterations for random tensors with correlated entries than for those with uncorrelated entries (specifically, see T_1 vs T_6 and T_7). This is possibly due to a higher *condition number* for the symmetric tensor decomposition problem in the case of tensor correlated entries. See [8] for more on the appropriate notion of condition number. We expect this corresponds to the optimization landscape for $\|P_A(x^{\otimes n})\|^2$ being “more flat” in the vicinity of $\pm a_i$ when the components are correlated. Further work on the rate of convergence for SPM is warranted.

Higher-order tensors and symmetric block term decomposition: The tensors T_8 , T_9 and T_{10} give examples of SPM’s performance on higher-order tensors (sixth order), and on the symmetric block term decompositions. Note that for the tensors with planted block term decompositions (T_8 and T_{10}) the timings for the POWER METHOD and DEFLATE steps are much smaller than their counterparts in the other tensors. This is due to two reasons: the tensors have a smaller flattening rank \mathbf{R} ; and the number of times Algorithm 2 repeats the POWER METHOD and DEFLATE subroutines is equal to the number of subspaces (40 for T_8 and 16 for T_{10}), which is also considerably less than the number of repeats for the other tensors and Algorithm 1.

Number of iterations and restarts: While the current implementation of SPM performs at most 5000 power iterations per component, and may also restart, in most cases the number of iterations before convergence is much smaller, and it only restarts for T_7 . This means that, most of the time, the




Fig. 6.1: Frequency of SPM successes as a function of length L and rank R for tensors of order 4.

power method converges to a global maxima at the first try and in fewer iterations.

Error: We also note that the error shown here is for decomposition of exact low rank tensors, therefore it isn't surprising that the error is so low. We show it mostly to affirm that SPM indeed succeeds to decompose these tensors and to show that it is numerically stable.

6.2 SPM maximal rank

In this subsection, we test what is the maximal rank for which Algorithm 1 still works. We generated low-rank tensors of order 4, similarly to T_1, \dots, T_5 in Subsection 6.1, with lengths ranging from 10 to 40 and ranks ranging from 8 to $\frac{L^2}{2}$, where L is the length of the tensor. For each length/rank pair, we generated 100 random tensors and registered the frequency of times SPM returns a decomposition with error, calculated by formula (6.1), at most 10^{-6} . Figure 6.1 plots this frequency as a function of the length L and rank R .

We observe that the maximal rank is slightly below the red line, which is the threshold provided by Proposition 3.2. However there seems to be a sharp threshold near the dashed blue line, below which SPM works, and above which it does not. This furnishes further evidence that SPM will work if $R \leq c_2 L^2$, for some constant $c_2 \leq \frac{1}{2}$, as stated in Conjecture 4.14.

6.3 Computation time comparison

We compare computation times for rank decompositions amongst SPM (Algorithm 1), FOOBI [26], the Tensorlab package [77] and LRSTA [60].

While in [26], there are two variants of FOOBI, we consider only the fastest method (FOOBI-1), and use our own optimized implementation of the algorithm. Tensorlab has two algorithms specialized for symmetric tensor decompositions, `ccpd_nls` and `ccpd_minf`. We compare our method only with `ccpd_nls`, since it always performed better than `ccpd_minf` in our comparisons. The method employs second-order methods to solve the non-convex least-squares optimization problem:

$$\operatorname{argmin}_{A \in \mathbb{R}^{L \times R}} \left\| T - \sum_{i=1}^R a_i^{\otimes m} \right\|^2, \quad (6.2)$$

where a_i are the columns of A . We set the maximum number of iterations at 1000, and the gradient and function tolerances (as stop criterions) to 1×10^{-12} and 1×10^{-24} , respectively. In the comparison below, `ccpd_nls` converged in every run. Finally, for LRSTA we used the MATLAB code available in Jiawang Nie's webpage, with default parameters.

In Figure 6.2, we plot the computation time (in seconds) of the rank decomposition of T for SPM, FOOBI and Tensorlab, as a function of L . For that comparison, we set $m = 4$, since FOOBI only applies to fourth-order tensors. For several values of L , ranging from 10 to 55, we generated T as follows. We set the rank $R = \left\lfloor \frac{L^2}{3} \right\rfloor$ and generated $\{a_i\}_{i=1}^R$ as R i.i.d. standard Gaussian random vectors in \mathbb{R}^L , letting $\lambda_i = 1$ for $i = 1, \dots, R$, and then formed T as in (1.1)⁶. To obtain more accurate computation times, we have computed the rank decomposition of each tensor 2 times with SPM and FOOBI, and 20 times with Tensorlab⁷, and report the average for each method. The Tensorlab method fails sometimes to produce the correct tensor decomposition, when that happens we just disregard that run, restart the clock and run the algorithm again.

In Figure 6.3, we plot a similar comparison for SPM and LRSTA. For this comparison, we set $m = 6$, L ranges from 4 to 30 and we generated T as follows. We set the rank $R = \left\lfloor \frac{L^2}{3} \right\rfloor$, generated $\{a_i\}_{i=1}^R$ as R i.i.d. random vectors sampled uniformly from the unit sphere in \mathbb{R}^L and let $\lambda_i = 729^{\frac{i}{R}}$ for $i = 1, \dots, R$, and then formed T as in (1.1). As in the previous comparison, we computed the rank decomposition of each tensor 2 times with SPM and with LRSTA, and report the average for each method.

The main takeaway from Figure 6.2 is that SPM considerably outperforms the other algorithms in time (note the logarithmic scale). SPM is, on aver-

⁶ While SPM and Tensorlab allow for negative scalars, the FOOBI algorithm does not deal with this case.

⁷ The running time for Tensorlab crucially depends on the initialization, thus we run it more times to get a more accurate estimate of its average computation time.




Fig. 6.2: Computation time comparison between SPM, FOOBI and Tensorlab.




Fig. 6.3: Computation time comparison between SPM and LRSTA.

age, 20 times faster than Tensorlab, which in terms of empirical performance, is a state-of-the-art package for tensor decomposition in MATLAB. We also observe that, while FOOBI's complexity is polynomial in L , the exponent is considerably high (around $O(R^4 L^2)$), which makes this algorithm considerably slower than alternatives, especially for large tensors.

We note however that the computation times may vary, depending on different hardware, software and implementation tweaks that change between different computers, servers or operative systems. Nevertheless, even after taking these differences into account, SPM should still considerably out-perform the other methods in terms of computation speed.

6.4 Noise stability

In this subsection, we compare the sensitivity to noise of SPM, FOOBI and Tensorlab. We fixed $m = 4$, $L = 10$, $R = 20$, generated T as in Section 6.3 and added independent centered Gaussian noise to the entries of T , with standard deviation ranging from 10^{-8} to 10^{-2} , in a symmetric manner. It is important to note that the resulting noisy tensors are no longer exactly low-rank, rather they are only approximately so. Nevertheless, the algorithms return estimates for $\{a_i\}_{i=1}^R$. In real-data applications, there are no exact low rank tensors, and therefore knowing that SPM works for approximately low rank tensors is of utmost importance. We measured the error using formula (6.1), and show the results in Table 6.2, where the last row is the average ratio between the noise standard deviation and the estimation error.

As can be observed in Table 6.2, Tensorlab is the most stable to noise, possibly due to its minimization of the L^2 norm. The sensitivity of SPM to noise is slightly larger than that of Tensorlab, comparable to LRSTA and slightly smaller than the sensitivity of FOOBI.

Finally, we note that to obtain the best low rank approximation, a combination of SPM and Tensorlab can be used: SPM can be used to determine

Noise Level	FOOBI	SPM	Tensorlab	LRSTA
1×10^{-08}	1.624×10^{-07}	1.224×10^{-07}	1.098×10^{-07}	1.093×10^{-07}
1×10^{-06}	1.484×10^{-05}	1.023×10^{-05}	9.270×10^{-06}	1.015×10^{-05}
1×10^{-04}	1.247×10^{-03}	7.726×10^{-04}	6.144×10^{-04}	6.790×10^{-04}
1×10^{-02}	1.133×10^{-01}	6.338×10^{-02}	5.220×10^{-02}	8.866×10^{-02}
Ratio	13.72	9.13	7.90	9.18

Table 6.2: Noise stability

the rank and obtain a fairly good estimate of the solution, and Tensorlab can be used to refine the solution of SPM. This procedure should be faster than just using Tensorlab, since that algorithm converges very rapidly when close to the local maximum.

6.5 Generalized principal component analysis

In this subsection, we measure Algorithm 2’s performance on GPCA, using simulated data. As a first experiment, we drew 20 three-dimensional subspaces of \mathbb{R}^{20} uniformly at random (with respect to the orthogonally-invariant probability measure on the set of subspaces, i.e., the Grassmannian $\text{Gr}(\mathbb{R}^3, \mathbb{R}^{20})$). We drew N sample points from this subspace arrangement, with N ranging from 10^2 to 10^5 , from an equally-weighted mixture of standard Gaussian distributions supported on each subspace. We then added centered Gaussian noise to the data, independently drawn from $\mathcal{N}(0, \sigma^2 I_{20})$, where $\sigma = 0.1$. Note that the noisy dataset is no longer an exact subset of the ground truth subspace arrangement. Nevertheless, we calculated the sample 4-th order moment and debiased it using [64, Lemma 3.3.2]. We then applied SPM (Algorithm 2) to compute the symmetric block term decomposition of this tensor. From the symmetric block term tensor decomposition, we obtained subspace estimates via $\hat{S}_i = \text{colspan}(\hat{A}_i) \subset \mathbb{R}^L$. Finally, the error in the subspace estimates was measured using the formula:

$$\text{error} := \sqrt{\min_{\pi \in \Pi^{20}} \sum_{i=1}^{20} \|P_{S_i} - P_{\hat{S}_i}\|^2}. \quad (6.3)$$

Here $P_{S_i} \in \mathbb{R}^{20 \times 3}$ is the orthogonal projection operator onto the subspace S_i and likewise for $P_{\hat{S}_i}$ (both with respect to the standard bases), the norm is the Frobenius norm, $\{S_i\}_{i=1}^{20}$ are the original subspaces and $\{\hat{S}_i\}_{i=1}^{20}$ are the ones obtained by SPM. In Figure 6.4, we plotted the error and times for estimating the 4-th moment and computing the subspaces via SPM, as functions of N .

We see in Figure (6.4a) that the error scales as $O(N^{-\frac{1}{2}})$, which is expected since the variance of the 4-th moment scales as $\frac{1}{N}$. On the other hand, in Figure 6.4b we observe that computing the decomposition using SPM always




Fig. 6.4: SPM for GPCA with 20 subspaces of \mathbb{R}^{20} and dimension 3.

takes approximately the same time independently of N , since the length, rank and order of the tensor do not change, however estimating and debiasing the moment scales proportionally to the number of points.

Followingly, we compare SPM's performance with Polynomial Differentiation (PDA) [79] and GPCA-Voting (GPCA-V) [82, 55], two existing algorithms for GPCA (somewhat close to SPM's approach). To this end, we drew 2 three-dimensional subspaces of \mathbb{R}^6 uniformly at random and drew 300 points from the standard Gaussian distribution in each subspace (600 points at total). We then add centered Gaussian noise, with variance σ^2 , to all points, with σ varying from 10^{-2} to 10^0 . Note that PDA is an unsupervised subspace clustering algorithm, so, to compare the performance of the three algorithms, we measured the misclassification error, using the formula:

$$\text{misclassification error} := \min \left\{ \frac{1}{600} \sum_{i=1}^{600} \mathbb{1}_{C(x_i) \neq \hat{C}(x_i)}, \frac{1}{600} \sum_{i=1}^{600} \mathbb{1}_{C(x_i) \neq -\hat{C}(x_i)} \right\}, \quad (6.4)$$

where $\{x_i\}_{i=1}^{600}$ are the drawn points, $C(x_i) \in \{1, -1\}$ are their true subspace labels and $\hat{C}(x_i)$ are the labels estimated by each algorithm. We plotted the average misclassification error, over 100 runs, as a function of σ in Figure 6.5.

In order to use SPM, we calculated the sample second and fourth-order moments of the data points, used the heuristic described in Appendix E to estimate σ from the dataset, and then debiased the fourth moment using [64, Lemma 3.3.2]. Subspace estimates were obtained using SPM (Algorithm 2) to compute the block term tensor decomposition of the debiased fourth moment. We then assigned each data point to the closest subspace obtained by SPM. As can be observed in Figure 6.5, SPM outperforms the other algorithms and is able to do classification at higher noise levels.




Fig. 6.5: Comparing missclassification error in GPCA.

We remark that for subspace estimation (rather than point classification), SPM should work at arbitrary SNR, assuming the noise statistics are known (or can be reliably estimated) and assuming the number of sample points is sufficiently large compared to the noise level. This is because Algorithm 2 interacts with the dataset only through sample moments; and upon debiasing, these converge to clean population moments (as the number of samples grows).

7 Discussion

This paper introduced a new algorithm, called SPM, for symmetric even-order tensor decomposition. While this algorithm significantly outperforms state-of-the-art algorithms for tensor decomposition in terms of computational speed, we were also able to establish a rich mathematical foundation for its properties.

We provided various guarantees for SPM. For tensors of order $2n$, with generic rank-1 components $a_i \in \mathbb{R}^L$, $i = 1, \dots, R$, provided the rank R is below a certain $O(L^n)$ threshold, we showed the rank of the matrix flattening is R , the subspace $\mathcal{A} = \{a_i^{\otimes n} : i = 1, \dots, R\} \subset \text{Sym } \mathcal{T}_L^n$ can be obtained from a matrix eigendecomposition and the rank-1 points in the subspace \mathcal{A} are precisely $a_i^{\otimes n}$ up to scale (Propositions 3.1 and 3.2). We proved that SS-HOPM converges unconditionally, which is important for the analysis of structured tensors (Theorem 4.5). Under the conditions mentioned above, we also proved that, for almost all initializations, SPM's power method converges to a local maximum of a certain constrained polynomial optimization problem, such that $\pm a_i$ in (2.7) are the only global maximizers (Theorem 4.1) and $\pm a_i$ attractive fixed points of the power method (Theorem 4.12). Empirically, we have observed that for most ranks there are no bad local maxima (Figure 4.1), and

we conjecture that these do not exist with high probability as the length of the tensor goes to infinity (Conjecture 4.14).

We also introduced symmetric block term tensor decompositions (Definition 2.6) and extended SPM to compute these. Using these decompositions, we were able to use SPM for generalized principal component analysis, outperforming state-of-the-art algorithms, especially in the presence of noise. We made connections to degree bounds in the algebra of subspace arrangements.

However, there are aspects of SPM that need further analysis. One is its performance for low-rank tensors perturbed with noise. While the rank of the flattening of a tensor is equal to its symmetric rank in the absence of noise, in the presence of noise the flattening matrix is full rank, and determining the correct singular value cut-off is crucial for SPM to work. On the other hand, the added noise might shift the eigenvalues, similarly to the spiked covariance model [33], and determining the optimal shrinkage for SPM is currently an open problem. Another question of interest is the effect of noise in the obtained subspace \mathcal{A} and in the analysis of the power method. Empirically, we observe that, as long as the noise perturbation is reasonable, the global maxima are only slightly perturbed. We intend to investigate how perturbations affect the second-order optimality conditions of these maxima, and if local maxima appear as the noise increases. It seems that this effort may require the development of novel perturbation bounds for tensor eigenvectors.

Another aspect we would like to study is the possible extension of SPM to *implicit* settings, as in [71]. Here a low-rank symmetric tensor arises as the population moment of a latent variable model (for example, a mixture of Gaussians). The idea is that to reduce storage costs, one wishes to recover the latent variables, still essentially by tensor decomposition, but without ever explicitly forming the moment tensor. It will be interesting to see whether our subspace-based algorithm can be made to work in the implicit framework.

We are also interested in extending SPM to compute other tensor decompositions. While we can use a rectangular matrix flattening in SPM to decompose symmetric tensors of order $2n + 1$ (Remark 3.4), the rank to which this approach applies is merely $O(L^n)$; however, rank $R = O(L^{n+0.5})$ is believed to be tractable. We have tried to extend SPM to decompose symmetric odd-order tensors of such rank by decomposing $\text{Sym}(T^{\otimes 2})$, where T is the odd-order tensor, but were unable to obtain meaningful results. Possibly, *Young flattenings*, used in [49] to determine defining equations for the set of low-rank tensors, may provide algorithmic clues for an improved odd-order SPM. We also intend to extend SPM to calculate non-symmetric tensor decompositions. Another extension of the SPM algorithm of interest arises when comparing SPM with the power method for determining eigenvalues of a matrix. We can think of the QR algorithm as a generalization of the power method, that performs n power iterations all at once (where the matrix is $n \times n$) and that is more stable to noise. Therefore, we are interested in an “all-at-once” analog for SPM where all power iterations and deflation steps are performed at the same time. Hopefully, this modification would further improve SPM’s numerical stability.

Finally, we are interested in the applications of SPM in statistics and analysis of datasets. A first application arises from applying SPM to GPCA datasets, but to harness the full potential of SPM, it should be further investigated how the noise affects the moments and how to best debias these moments. We suspect that high noise and big data sets in GPCA may be a setting in which SPM is most helpful, as SPM interacts with noisy datasets only via averages. To test this, we wish to apply SPM to cryo-electron microscopy (cryo-EM), a molecular imaging technique known to produce massive and extremely noisy data sets. Specifically, we will investigate whether GPCA can help with parameter estimation in the rigid subunits model for molecular variability [56], since rigid motion segmentation is an established application of GPCA [78]. Lastly, SPM can also recover the underlying polynomials in the latent variables, as in (2.9). It would be interesting to find applications where recovering the underlying polynomials is of most relevance (rather than the subspaces).

Acknowledgements The authors thank Emmanuel Abbe, Amir Ali Ahmadi, Tamir Bendory, Greg Blekherman, Nicolas Boumal, Justin Chen, Tamara Kolda, János Kollár, Pablo Parrilo, Kristian Ranestad and Amit Singer for fruitful discussions. The authors also thank Jiawang Nie for sharing code and the anonymous referees for helpful comments, including further pointers to the literature. JK was supported in part by the Simons Collaboration on Algorithms and Geometry. JP was supported in part by NSF BIGDATA Award IIS-1837992. Much of this research was carried out while JK and JP were with the Program in Applied and Computational Mathematics, Princeton University.

References

1. Absil, P.A., Mahony, R., Andrews, B.: Convergence of the iterates of descent methods for analytic cost functions. *SIAM Journal on Optimization* **16**(2), 531–547 (2005)
2. Anandkumar, A., Ge, R., Hsu, D.J., Kakade, S.M., Telgarsky, M.: Tensor decompositions for learning latent variable models. *Journal of Machine Learning Research* **15**(1), 2773–2832 (2014)
3. Ballico, E.: On the weak non-defectivity of veronese embeddings of projective spaces. *Central European Journal of Mathematics* **3**(2), 183–187 (2005)
4. Bandeira, A.S., Perry, A., Wein, A.S.: Notes on computational-to-statistical gaps: predictions using statistical physics. *Portugaliae Mathematica* **75**(2), 159–186 (2018)
5. Basser, P.J., Pajevic, S.: Spectral decomposition of a 4th-order covariance tensor: Applications to diffusion tensor MRI. *Signal Processing* **87**(2), 220–236 (2007)
6. Blekherman, G., Sinn, R., Smith, G.G., Velasco, M.: Sums of squares and quadratic persistence on real projective varieties. *arXiv preprint arXiv:1902.02754* (2019)

7. Brachat, J., Comon, P., Mourrain, B., Tsigaridas, E.: Symmetric tensor decomposition. *Linear Algebra and its Applications* **433**(11-12), 1851–1872 (2010)
8. Breiding, P., Vannieuwenhoven, N.: The condition number of join decompositions. *SIAM Journal on Matrix Analysis and Applications* **39**(1), 287–309 (2018)
9. Bro, R.: Parafac. tutorial and applications. *Chemometrics and intelligent laboratory systems* **38**(2), 149–171 (1997)
10. Bunch, J.R., Nielsen, C.P., Sorensen, D.C.: Rank-one modification of the symmetric eigenproblem. *Numerische Mathematik* **31**(1), 31–48 (1978)
11. Cambré, J., De Lathauwer, L., De Moor, B.: Best rank (r, r, r) super-symmetric tensor approximation – a continuous-time approach. In: *Proceedings of the IEEE Signal Processing Workshop on Higher-Order Statistics. SPW-HOS'99*, pp. 242–246. IEEE (1999)
12. Cardoso, J.F.: Super-symmetric decomposition of the fourth-order cumulant tensor. blind identification of more sources than sensors. In: *ICASSP 91: 1991 International Conference on Acoustics, Speech, and Signal Processing*, pp. 3109–3112. IEEE (1991)
13. Cardoso, J.F.: Blind signal separation: statistical principles. *Proceedings of the IEEE* **86**(10), 2009–2025 (1998)
14. Carlini, E., Catalisano, M.V., Geramita, A.V.: Subspace arrangements, configurations of linear spaces and the quadrics containing them. *Journal of Algebra* **362**, 70–83 (2012)
15. Carroll, J.D., Chang, J.J.: Analysis of individual differences in multidimensional scaling via an N -way generalization of “Eckart-Young” decomposition. *Psychometrika* **35**(3), 283–319 (1970)
16. Cartwright, D., Sturmfels, B.: The number of eigenvalues of a tensor. *Linear Algebra and its Applications* **438**(2), 942–952 (2013)
17. Chevalier, P.: Optimal separation of independent narrow-band sources: Concept and performance. *Signal Processing* **73**(1-2), 27–47 (1999)
18. Chevalier, P., Albera, L., Ferréol, A., Comon, P.: On the virtual array concept for higher order array processing. *IEEE Transactions on Signal Processing* **53**(4), 1254–1271 (2005)
19. Chiantini, L., Ciliberto, C.: Weakly defective varieties. *Transactions of the American Mathematical Society* **354**(1), 151–178 (2002)
20. Chiantini, L., Ottaviani, G., Vannieuwenhoven, N.: On generic identifiability of symmetric tensors of subgeneric rank. *Transactions of the American Mathematical Society* **369**(6), 4021–4042 (2017)
21. Chu, M.T., Funderlic, R.E., Golub, G.H.: A rank-one reduction formula and its applications to matrix factorizations. *SIAM Review* **37**(4), 512–530 (1995)
22. Cline, R.E., Funderlic, R.: The rank of a difference of matrices and associated generalized inverses. *Linear Algebra and its Applications* **24**, 185–215 (1979)
23. Comon, P., Golub, G., Lim, L.H., Mourrain, B.: Symmetric tensors and symmetric tensor rank. *SIAM Journal on Matrix Analysis and Applica-*

- tions **30**(3), 1254–1279 (2008)
24. De Lathauwer, L.: Decompositions of a higher-order tensor in block terms—part i: Lemmas for partitioned matrices. *SIAM Journal on Matrix Analysis and Applications* **30**(3), 1022–1032 (2008)
 25. De Lathauwer, L.: Decompositions of a higher-order tensor in block terms—part ii: Definitions and uniqueness. *SIAM Journal on Matrix Analysis and Applications* **30**(3), 1033–1066 (2008)
 26. De Lathauwer, L., Castaing, J., Cardoso, J.F.: Fourth-order cumulant-based blind identification of underdetermined mixtures. *IEEE Transactions on Signal Processing* **55**(6), 2965–2973 (2007)
 27. De Lathauwer, L., De Moor, B., Vandewalle, J.: On the best rank-1 and rank- (R_1, R_2, \dots, R_n) approximation of higher-order tensors. *SIAM Journal on Matrix Analysis and Applications* **21**(4), 1324–1342 (2000)
 28. De Lathauwer, L., Nion, D.: Decompositions of a higher-order tensor in block terms—part iii: Alternating least squares algorithms. *SIAM Journal on Matrix Analysis and Applications* **30**(3), 1067–1083 (2008)
 29. De Silva, V., Lim, L.H.: Tensor rank and the ill-posedness of the best low-rank approximation problem. *SIAM Journal on Matrix Analysis and Applications* **30**(3), 1084–1127 (2008)
 30. Derksen, H.: Hilbert series of subspace arrangements. *Journal of Pure and Applied Algebra* **209**(1), 91–98 (2007)
 31. Derksen, H., Sidman, J.: A sharp bound for the Castelnuovo-Mumford regularity of subspace arrangements. *Advances in Mathematics* **172**(2), 151–157 (2002)
 32. Dogan, M.C., Mendel, J.M.: Applications of cumulants to array processing. I. aperture extension and array calibration. *IEEE Transactions on Signal Processing* **43**(5), 1200–1216 (1995)
 33. Donoho, D.L., Gavish, M., Johnstone, I.M.: Optimal shrinkage of eigenvalues in the spiked covariance model. *Annals of Statistics* **46**(4), 1742–1778 (2018)
 34. Fornasier, M., Vybíral, J., Daubechies, I.: Identification of shallow neural networks by fewest samples. *arXiv preprint arXiv:1804.01592* (2018)
 35. Ge, R., Ma, T.: On the optimization landscape of tensor decompositions. In: *Advances in Neural Information Processing Systems*, pp. 3653–3663 (2017)
 36. Hackbusch, W.: *Tensor Spaces and Numerical Tensor Calculus*, vol. 42. Springer (2012)
 37. Harshman, R.A.: Foundations of the PARAFAC procedure: models and conditions for an “explanatory” multi-model factor analysis. *UCLA Working Papers in Phonetics* **16**, 1–84 (1970)
 38. Hartshorne, R.: *Algebraic Geometry*, vol. 52. Springer Science & Business Media (2013)
 39. Hillar, C.J., Lim, L.H.: Most tensor problems are NP-hard. *Journal of the ACM (JACM)* **60**(6), 45:1–45:39 (2013)
 40. Hopkins, S.B., Schramm, T., Shi, J., Steurer, D.: Fast spectral algorithms from sum-of-squares proofs: tensor decomposition and planted sparse vec-

- tors. In: *Proceedings of the Forty-Eighth Annual ACM Symposium on Theory of Computing*, pp. 178–191. ACM (2016)
41. Hyvärinen, A., Karhunen, J., Oja, E.: *Independent Component Analysis*, vol. 46. John Wiley & Sons (2004)
 42. Iarrobino, A., Kanev, V.: *Power Sums, Gorenstein Algebras, and Determinantal Loci*, vol. 1721. Springer Science & Business Media (1999)
 43. Ishteva, M., Absil, P.A., Van Dooren, P.: Jacobi algorithm for the best low multilinear rank approximation of symmetric tensors. *SIAM Journal on Matrix Analysis and Applications* **34**(2), 651–672 (2013)
 44. Kolda, T., Mayo, J.: Shifted power method for computing tensor eigenpairs. *SIAM Journal on Matrix Analysis and Applications* **32**(4), 1095–1124 (2011)
 45. Kolda, T.G.: Numerical optimization for symmetric tensor decomposition. *Mathematical Programming* **151**(1), 225–248 (2015)
 46. Kolda, T.G., Bader, B.W.: Tensor decompositions and applications. *SIAM Review* **51**(3), 455–500 (2009)
 47. Kolda, T.G., Mayo, J.R.: An adaptive shifted power method for computing generalized tensor eigenpairs. *SIAM Journal on Matrix Analysis and Applications* **35**(4), 1563–1581 (2014)
 48. Landsberg, J.M.: *Tensors: Geometry and Applications*, vol. 128. AMS Graduate Studies in Mathematics (2012)
 49. Landsberg, J.M., Ottaviani, G.: Equations for secant varieties of veronese and other varieties. *Annali di Matematica Pura ed Applicata* **192**(4), 569–606 (2013)
 50. Lee, J.D., Simchowitz, M., Jordan, M.I., Recht, B.: Gradient descent only converges to minimizers. In: *Conference on Learning Theory (COLT)*, pp. 1246–1257 (2016)
 51. Lerman, G., Zhang, T., et al.: Robust recovery of multiple subspaces by geometric l_p minimization. *Annals of Statistics* **39**(5), 2686–2715 (2011)
 52. Lim, L.H.: Singular values and eigenvalues of tensors: a variational approach. In: *1st IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing*, pp. 129–132. IEEE (2005)
 53. Łojasiewicz, S.: Ensembles semi-analytiques. *Institute Des Hautes Études Scientifiques (IHES) Notes*, Bures-sur-Yvette, France (1965)
 54. Ma, T., Shi, J., Steurer, D.: Polynomial-time tensor decompositions with sum-of-squares. In: *2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS)*, pp. 438–446. IEEE (2016)
 55. Ma, Y., Yang, A.Y., Derksen, H., Fossum, R.: Estimation of subspace arrangements with applications in modeling and segmenting mixed data. *SIAM Review* **50**(3), 413–458 (2008)
 56. Nakane, T., Kimanius, D., Lindahl, E., Scheres, S.: Characterisation of molecular motions in cryo-EM single-particle data by multi-body refinement in RELION. *eLife* **7**, e36861 (2018)
 57. Nakatsukasa, Y., Soma, T., Uschmajew, A.: Finding a low-rank basis in a matrix subspace. *Mathematical Programming* **162**(1-2), 325–361 (2017)

58. Nelson, R.B.: Simplified calculation of eigenvector derivatives. *AIAA Journal* **14**(9), 1201–1205 (1976)
59. Nie, J.: Generating polynomials and symmetric tensor decompositions. *Foundations of Computational Mathematics* **17**(2), 423–465 (2017)
60. Nie, J.: Low rank symmetric tensor approximations. *SIAM Journal on Matrix Analysis and Applications* **38**(4), 1517–1540 (2017)
61. Oeding, L., Ottaviani, G.: Eigenvectors of tensors and algorithms for waring decomposition. *Journal of Symbolic Computation* **54**, 9–35 (2013)
62. Panageas, I., Piliouras, G.: Gradient descent only converges to minimizers: Non-isolated critical points and invariant regions. In: *8th Innovations in Theoretical Computer Science (ITCS)* (2017)
63. Panageas, I., Piliouras, G., Wang, X.: First-order methods almost always avoid saddle points: The case of vanishing step-sizes. In: *Advances in Neural Information Processing Systems* (2019)
64. Pereira, J.M.: *Information Theoretic Aspects of Cryo-Electron Microscopy*. Ph.D. thesis, Princeton University (2019)
65. Phan, A.H., Tichavský, P., Cichocki, A.: Tensor deflation for candecomp/parafac—part i: Alternating subspace update algorithm. *IEEE Transactions on Signal Processing* **63**(22), 5924–5938 (2015)
66. Podosinnikova, A., Perry, A., Wein, A., Bach, F., d’Aspremont, A., Sontag, D.: Overcomplete independent component analysis via SDP. In: *22nd International Conference on Artificial Intelligence and Statistics (AISTATS)* (2019)
67. Qi, L.: Eigenvalues of a real supersymmetric tensor. *Journal of Symbolic Computation* **40**(6), 1302–1324 (2005)
68. Regalia, P.A.: Monotonically convergent algorithms for symmetric tensor approximation. *Linear Algebra and its Applications* **438**(2), 875–890 (2013)
69. Rheinboldt, W.C.: *Methods for Solving Systems of Nonlinear Equations*, vol. 70. SIAM Regional Conference Series in Applied Mathematics (1974)
70. Schneider, R., Uschmajew, A.: Convergence results for projected line-search methods on varieties of low-rank matrices via Łojasiewicz inequality. *SIAM Journal on Optimization* **25**(1), 622–646 (2015)
71. Sherman, S., Kolda, T.G.: Estimating higher-order moments using symmetric tensor decomposition. *arXiv preprint* arXiv:1911.03813 (2019)
72. Shub, M.: *Global Stability of Dynamical Systems*. Springer Science & Business Media (2013)
73. Stor, N.J., Slapničar, I., Barlow, J.L.: Forward stable eigenvalue decomposition of rank-one modifications of diagonal matrices. *Linear Algebra and its Applications* **487**, 301–315 (2015)
74. Sylvester, J.J.: Lx. on a remarkable discovery in the theory of canonical forms and of hyperdeterminants. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **2**(12), 391–410 (1851)
75. Tsakiris, M.C., Vidal, R.: Filtrated algebraic subspace clustering. *SIAM Journal on Imaging Sciences* **10**(1), 372–415 (2017)

76. Van der Veen, A.J., Paulraj, A.: An analytical constant modulus algorithm. *IEEE Transactions on Signal Processing* **44**(5), 1136–1155 (1996)
77. Vervliet, N., Debals, O., Sorber, L., Van Barel, M., De Lathauwer, L.: Tensorlab 3.0. Available online. URL: <http://www.tensorlab.net> (2016)
78. Vidal, R., Hartley, R.: Motion segmentation with missing data using PowerFactorization and GPCA. In: *2004 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR)*. IEEE (2004)
79. Vidal, R., Ma, Y., Sastry, S.: Generalized principal component analysis (GPCA). *IEEE Transactions on Pattern Analysis and Machine Intelligence* **27**(12), 1945–1959 (2005)
80. Vidal, R., Ma, Y., Sastry, S.: *Generalized Principal Component Analysis*, vol. 40. Springer Interdisciplinary Applied Mathematics (2016)
81. Woodbury, M.A.: Inverting modified matrices. *Memorandum Report, Institute of Advanced Studies*, Princeton, NJ **42** (1950)
82. Yang, A.Y., Rao, S., Wagner, A., Ma, Y., Fossum, R.M.: Hilbert functions and applications to the estimation of subspace arrangements. In: *Tenth IEEE International Conference on Computer Vision (ICCV'05)*, vol. 1, pp. 158–165. IEEE (2005)

A Inner product and symmetric tensors

In this appendix, we present various results on symmetric tensors and inner product between tensors which will be useful in several proofs throughout this paper.

Lemma A.1 *For two tensors $T, T' \in \mathcal{T}_L^m$, such that T is a symmetric tensor, we have*

$$\langle T, T' \rangle = \langle T, \text{Sym}(T') \rangle. \quad (\text{A.1})$$

Moreover if $U, U' \in \mathcal{T}_L^{m'}$, we have

$$\langle T \otimes U, T' \otimes U' \rangle = \langle T, T' \rangle \langle U, U' \rangle. \quad (\text{A.2})$$

Proof Since Sym is a self-adjoint operator [23]:

$$\langle T, \text{Sym}(T') \rangle = \langle \text{Sym}(T), T' \rangle = \langle T, T' \rangle. \quad (\text{A.3})$$

and (A.1) follows. A proof of (A.2) is available in [36].

We now prove Lemma 2.2.

Proof (Lemma 2.2) We define Φ as in (2.5), that is, for $T \in \mathcal{T}_L^n$:

$$\Phi(T)(X) = \langle T, X^{\otimes n} \rangle, \quad (\text{A.4})$$

with $X \in \mathbb{R}^L$. The linearity of Φ follows from the bilinearity of the inner product. Since all the entries of the tensor $X^{\otimes n}$ are in $\mathbb{R}[X_1, \dots, X_L]_n$, $\Phi(T)$ is also in $\mathbb{R}[X_1, \dots, X_L]_n$. Moreover, the tensor $X^{\otimes n}$ contains all the monomials of degree n in L variables, thus Φ is surjective over all tensors in \mathcal{T}_L^n . Additionally, since $X^{\otimes n} = \text{Sym}(X^{\otimes n})$, (A.1) implies that $\langle T, X^{\otimes n} \rangle = \langle \text{Sym}(T), X^{\otimes n} \rangle$, thus Φ is surjective over all symmetric tensors in \mathcal{T}_L^n . Since

the vector space dimensions of $\text{Sym}(\mathcal{T}_L^n)$ and $\mathbb{R}[X_1, \dots, X_L]_n$ coincide, Φ is a one-to-one map between these spaces. Finally, (2.6) follows from (A.2):

$$\begin{aligned} \Phi(\text{Sym}(T_1 \otimes T_2)) &= \langle \text{Sym}(T_1 \otimes T_2), X^{\otimes m+n} \rangle \\ &= \langle T_1 \otimes T_2, X^{\otimes m+n} \rangle \\ &= \langle T_1, X^{\otimes m} \rangle \langle T_2, X^{\otimes n} \rangle \\ &= \Phi(T_1)\Phi(T_2). \end{aligned} \quad \square$$

We now prove (2.13).

Proof (2.13) A useful property of symmetric Tucker products is that the adjoint of a symmetric Tucker product is the symmetric Tucker product associated with the matrix transposed. Let $T \in \text{Sym } \mathcal{T}_n^d$, $U \in \text{Sym } \mathcal{T}_m^d$ and $A \in \mathbb{R}^{m \times n}$. Then (2.11) implies:

$$\begin{aligned} \langle U, (A; \dots; A) \cdot T \rangle &= \sum_{j_1=1}^m \dots \sum_{j_d=1}^m \sum_{k_1=1}^n \dots \sum_{k_d=1}^n U_{j_1 \dots j_d} T_{k_1 \dots k_d} A_{j_1 k_1} \dots A_{j_d k_d} \\ &= \langle (A^T; \dots; A^T) \cdot U \times T \rangle. \end{aligned} \quad (\text{A.5})$$

Recalling now (2.5), we have:

$$\begin{aligned} \Phi((A_i; \dots; A_i) \cdot \mathbf{A}_i)(X) &= \langle (A_i; \dots; A_i) \cdot \mathbf{A}_i, X^{\otimes m} \rangle \\ &= \langle \mathbf{A}_i, (A_i^T; \dots; A_i^T) \cdot X^{\otimes m} \rangle \\ &= \langle \mathbf{A}_i, (A_i^T X)^{\otimes m} \rangle \\ &= \Phi(\mathbf{A}_i)(A_i^T X) \\ &= f_i(A_i^T X). \end{aligned} \quad (\text{A.6})$$

B Speeding up deflation using Householder reflections

In this appendix, we present some results involving the deflation step in SPM and its variant for the symmetric block term decomposition. The following lemma states a formula for determining the λ_i (or \mathbf{A}_i for generalized variant) in the deflation step of SPM. This formula has similarities with the Woodbury matrix identity [81], the Bunch-Nielsen-Sorensen formula [10], and follows from matrix identities in [22]. However, we present the proof here for completeness and brevity.

Lemma B.1 *Let $A_{n \times n}$, $U_{n \times k}$ and $V_{n \times k}$ be full rank matrices with $k \leq n$, let $C^* = (V^T A^{-1} U)^\dagger$, where \dagger denotes Moore-Penrose pseudo-inverse, and $d = \text{rank}(C^*)$. Then*

$$\text{rank}(A - U C V^T) \geq n - d, \quad (\text{B.1})$$

with equality if and only if $C = C^$. Moreover, the null space of $A - U C^* V^T$ is spanned by the columns of $A^{-1} U C^*$.*

Proof Since U is full-rank, there exists an SVD decomposition

$$U = W \begin{bmatrix} D \\ 0_{(n-k) \times k} \end{bmatrix} B^T, \quad (\text{B.2})$$

where W and B are $n \times n$ and $k \times k$ orthogonal matrices respectively, and D is a $k \times k$ a diagonal and full-rank matrix. Let

$$U^\dagger = \begin{bmatrix} B D^{-1} & 0_{k \times (n-k)} \\ 0_{(n-k) \times k} & I_{n-k} \end{bmatrix} W^T \quad \text{and} \quad U^\perp = W \begin{bmatrix} 0_{k \times (n-k)} \\ I_{n-k} \end{bmatrix}. \quad (\text{B.3})$$

Thus we have

$$U^\dagger[U \ U^\perp] = \begin{bmatrix} I_k & 0_{k \times (n-k)} \\ 0_{(n-k) \times k} & I_{n-k} \end{bmatrix}. \quad (\text{B.4})$$

We now calculate $U^\dagger(A - UCV^T)A^{-1}[U \ U^\perp]$, since multiplying by these full rank matrices do not change the rank of $A - UCV^T$. We have $U^\dagger AA^{-1}[U \ U^\perp] = I_n$. On the other hand,

$$\begin{aligned} U^\dagger UCV^T A^{-1}[U \ U^\perp] &= \begin{bmatrix} I_k \\ 0_{n-k \times k} \end{bmatrix} CV^T A^{-1}[U \ U^\perp] \\ &= \begin{bmatrix} CV^T A^{-1}U & CV^T A^{-1}U^\perp \\ 0_{(n-k) \times k} & 0_{(n-k) \times (n-k)} \end{bmatrix}, \end{aligned} \quad (\text{B.5})$$

thus

$$U^\dagger(A - UCV^T)A^{-1}[U \ U^\perp] = \begin{bmatrix} I_k - CV^T A^{-1}U & -CV^T A^{-1}U^\perp \\ 0_{(n-k) \times k} & I_{n-k} \end{bmatrix}. \quad (\text{B.6})$$

Since this is a block upper triangular matrix, $\text{rank}(A - UCV^T) = \text{rank}(I_{n-k}) + \text{rank}(I_k - CV^T A^{-1}U)$, and $\text{rank}(I_k - CV^T A^{-1}U)$ is minimized when $C = (V^T A^{-1}U)^\dagger = C^*$. Finally, we note that $C^* V^T A^{-1}U$ is an orthogonal projection matrix, therefore the null space of $I_k - C^* V^T A^{-1}U$ is spanned by the columns of C^* , and the result follows. \square

We now explain the modified deflation step in Algorithm 1. Supposing $r \leq R$ components (λ_i, a_i) are yet to be found, then the modified deflation step takes time $O(rL^n)$, which is less than the time $O(r^2 L^n)$ it takes to calculate VO . The key observation is noticing that (V, D) does not in fact need to be an eigendecomposition, as long as V is an $L^n \times r$ matrix with orthonormal columns. In general, $\text{vec}(P_{\mathcal{A}}(x^{\otimes n})) = VV^\dagger \text{vec}(x^{\otimes n})$, and therefore it is convenient if V has orthonormal columns, for then $V^\dagger = V^T$. As a starting point, we still obtain (V, D) from the thin eigendecomposition of $\text{mat}(T)$; this way it is easier to estimate the rank by selecting a eigenvalue cut-off, and after each deflation step, we maintain V as a matrix with orthonormal columns by updating $V \leftarrow VO$ where O is a $r \times (r-1)$ matrix also with orthonormal columns. However, we do not need D to be diagonal. The iterations which have higher computational cost in the deflation step of Algorithm 1 are as follows:

- (a) Calculating $D^{-1}\alpha$, as an intermediate step to calculate $\lambda = (\alpha D^{-1}\alpha)^{-1}$. If D is a diagonal matrix this takes $O(r)$ operations, but for a general $r \times r$ matrix D this might take up to $O(r^3)$ operations.
- (b) Determining $O_{r \times (r-1)}$ and $\tilde{D}_{(r-1) \times (r-1)}$ such that O has orthogonal columns and

$$O\tilde{D}O^T = D - \lambda\alpha\alpha^T. \quad (\text{B.7})$$

We need this for the updates $D \leftarrow \tilde{D}$ and $V \leftarrow VO$. We note that if \tilde{D} is diagonal, (B.7) is an eigendecomposition and takes time⁸ $O(r^3)$, therefore not requiring \tilde{D} to be diagonal might speed up this step.

- (c) Calculating VO . Since V is a $L^n \times r$ matrix and O is a $r \times r-1$ matrix, this operation might take time up to $O(r^2 L^n)$, if not done intelligently.

We first focus on speeding up (c). To this end, we define O as a submatrix of a Householder reflection. Householder reflections are matrices that can be written as $H_x = I - 2xx^T$ for some unit norm vector $x \in \mathbb{R}^n$. These matrices are orthogonal, symmetric, and since they are rank-1 updates of the identity matrix, multiplying an $m \times n$ matrix by H_x takes time $O(mn)$, rather than $O(mn^2)$. In our case, we will define O such that

$$[O \ y] = H_x, \quad (\text{B.8})$$

for some unit-norm vector x , and for y in the null space of $D - \lambda\alpha\alpha^T$. By Lemma B.1, the null space of $D - \lambda\alpha\alpha^T$ is $\text{Span}(D^{-1}\alpha)$, so we set $y = D^{-1}\alpha/\|D^{-1}\alpha\|$. Note that $D^{-1}\alpha$

⁸ In fact, the eigendecomposition of a rank-1 update of a diagonal matrix costs only $O(r^2)$ operations by a specialized method [73]. However, using this would not reduce the overall computation time of the deflation step because of the update $V \leftarrow VO$.

was previously calculated as an intermediate step to calculate λ . Equation (B.8) implies $y = e_n - 2x_n x$, where e_n is the standard basis vector $(e_n)_j = \delta_{jn}$. Finally, we may solve for x to obtain $x_n = \sqrt{(1 - y_n)/2}$ and $x_i = -y_i/x_n$ for $i = 1, \dots, n-1$. Since the obtained matrix O is a submatrix of a Householder reflection, multiplying V by O takes time $O(L^n r)$, instead of $O(L^n r^2)$. Once O is calculated, we have:

$$\begin{aligned} D - \lambda \alpha \alpha^T &= (D - \lambda \alpha \alpha^T) H_x H_x^T \\ &= (D - \lambda \alpha \alpha^T) [O y] [O y]^T \\ &= (D - \lambda \alpha \alpha^T) (O O^T + y y^T) \\ &= (D - \lambda \alpha \alpha^T) O O^T \end{aligned} \tag{B.9}$$

$$= O O^T (D - \lambda \alpha \alpha^T) O O^T, \tag{B.10}$$

where (B.9) follows from y being in the null space of $D - \lambda \alpha \alpha^T$, and (B.10) follows from $D - \lambda \alpha \alpha^T$ being symmetric. Thus, if we set $\tilde{D} = O^T (D - \lambda \alpha \alpha^T) O$, we have $D - \lambda \alpha \alpha^T = O \tilde{D} O^T$. To calculate \tilde{D} we merely need to multiply an $r \times r$ matrix by O , which takes $O(r^2)$ time. While steps (b) and (c) now take $O(r^2)$, \tilde{D} is not diagonal anymore, so (a) might now take up to $O(r^3)$ time. In order to speed up (a), we change our algorithm to store $C' = D^{-1}$ instead of D . By doing so, we can now calculate $D^{-1} \alpha = C' \alpha$ in $O(r^2)$ time. However, to update C , we now need to update $C \leftarrow \tilde{C} = \tilde{D}^{-1}$, where $\tilde{D} = O^T (D - \lambda \alpha \alpha^T) O$. It turns out that $\tilde{D}^{-1} = O^T D^{-1} O = O^T C O$, which we may verify as follows:

$$\begin{aligned} (O^T D^{-1} O) \tilde{D} &= O^T D^{-1} O O^T (D - \lambda \alpha \alpha^T) O \\ &= O^T D^{-1} (D - \lambda \alpha \alpha^T) O \\ &= O^T (O - \lambda \tilde{D}^{-1} \alpha \alpha^T O) \\ &= I. \end{aligned} \tag{B.11}$$

Here we used (B.9), the symmetry of $D - \lambda \alpha \alpha^T$, $O^T y = 0 = O^T D^{-1} \alpha$ and $O^T O = I$. We then update $C \leftarrow O^T C O$, which takes $O(r^2)$ time.

We can also apply this improved deflation procedure to the block term decomposition (Algorithm 2), although with some modifications. Here α is not a column vector, but is instead an $r \times k$ matrix, where $k = \binom{\ell_i + n - 1}{n}$. Nevertheless, Lemma B.1 still applies, with $\Lambda = (\alpha^T D \alpha)^{-1}$ now being a $k \times k$ matrix, and the null space of $D - \alpha \Lambda \alpha^T$ being $\text{colspan}(D^{-1} \alpha)$. Accordingly, $[O y]$ is now a product of k Householder reflections, obtained by calculating the QR decomposition of $D^{-1} \alpha$. Here, y is a matrix with orthogonal columns and U is an upper triangular $k \times k$ matrix such that $y U = D^{-1} \alpha$. For our purposes, U is a byproduct of the QR decomposition, and our main concern is setting the columns of y to be an orthonormal basis of the null space of $D - \alpha \Lambda \alpha^T$. Since $[O y]$ is a product of k Householder reflections, multiplying V by O costs $O(L^n k r)$. Finally, we keep storing $C = D^{-1}$, and similarly update $C \leftarrow O^T C O$, which now takes $O(k r^2)$ time.

C Supporting proofs for Section 4

This appendix contains proofs of several supporting lemmas and propositions in Section 4. We have placed these arguments here in order to streamline the reading of Section 4.

C.1 Proof of Lemma 4.4

The Hessian of f at x is given by

$$\nabla^2 f(x) = 2n^2 \sum_{i=1}^R (V_i \cdot x^{\otimes n-1}) (V_i \cdot x^{\otimes n-1})^T + 2n(n-1) \sum_{i=1}^R \langle V_i, x^{\otimes n} \rangle V_i \cdot x^{\otimes n-2}. \tag{C.1}$$

In order to avoid repetition of the tensor product notation and make the proof more readable, for the rest of this proof we use the notation $xy := x \otimes y$ and $x^n := x^{\otimes n}$. Let $x, y \in \mathbb{R}^L$ such that $\|x\| = \|y\| = 1$. We have:

$$\begin{aligned} \frac{1}{2n} y^T \nabla^2 f(x) y &= n \sum_{i=1}^R \langle V_i, x^{n-1} y \rangle^2 + (n-1) \sum_{i=1}^R \langle V_i, x^n \rangle \langle V_i, x^{n-2} y^2 \rangle \\ &= n \|P_{\mathcal{A}}(x^{n-1} y)\|^2 + (n-1) \langle P_{\mathcal{A}}(x^n), x^{n-2} y^2 \rangle, \end{aligned} \quad (\text{C.2})$$

where (C.2) follows from (4.4). Define $\bar{y} \in \mathbb{R}^L$ and $\alpha, \beta \in \mathbb{R}$ such that $\bar{y} \perp x$, $\|\bar{y}\| = 1$, $\alpha^2 + \beta^2 = 1$ and $y = \alpha \bar{y} + \beta x$. Substituting this expression for y into (C.2), and rearranging using $P_{\mathcal{A}}$ is self-adjoint and $P_{\mathcal{A}}^2 = P_{\mathcal{A}}$,

$$\begin{aligned} \frac{1}{2n} y^T \nabla^2 f(x) y &= n \alpha^2 \|P_{\mathcal{A}}(x^{n-1} \bar{y})\|^2 + (n-1) \alpha^2 \langle P_{\mathcal{A}}(x^n), x^{n-2} \bar{y}^2 \rangle \\ &\quad + (2n-1) \beta \langle P_{\mathcal{A}}(x^n), x^{n-1} (\beta x + 2\alpha \bar{y}) \rangle. \end{aligned} \quad (\text{C.3})$$

On the other hand,

$$\begin{aligned} 0 &\leq \|P_{\mathcal{A}}(x^{n-1}(\sqrt{n}\alpha\bar{y} + \sqrt{2n-1}\beta x))\|^2 \\ &= \langle P_{\mathcal{A}}(x^{n-1}(\sqrt{n}\alpha\bar{y} + \sqrt{2n-1}\beta x)), x^{n-1}(\sqrt{n}\alpha\bar{y} + \sqrt{2n-1}\beta x) \rangle \\ &= n \alpha^2 \|P_{\mathcal{A}}(x^{n-1} \bar{y})\|^2 + (2n-1) \beta^2 \|P_{\mathcal{A}}(x^n)\|^2 \\ &\quad + 2\alpha\beta\sqrt{n(2n-1)} \langle P_{\mathcal{A}}(x^n), x^{n-1} \bar{y} \rangle. \end{aligned} \quad (\text{C.4})$$

Subtracting (C.4) from (C.3), we obtain

$$\frac{1}{2n} y^T \nabla^2 f(x) y \geq \langle P_{\mathcal{A}}(x^n), \alpha\beta b_n x^{n-1} \bar{y} + \alpha^2(n-1)x^{n-2} \bar{y}^2 \rangle, \quad (\text{C.5})$$

where

$$b_n = 2 \left(2n-1 - \sqrt{n(2n-1)} \right). \quad (\text{C.6})$$

Letting $Y = \alpha\beta b_n x^{n-1} \bar{y} + \alpha^2(n-1)x^{n-2} \bar{y}^2$, we have

$$\begin{aligned} \frac{1}{2n} y^T \nabla^2 f(x) y &\geq \langle P_{\mathcal{A}}(x^n), Y \rangle \\ &= \langle P_{\mathcal{A}}(x^n), \text{Sym}(Y) \rangle, \end{aligned} \quad (\text{C.7})$$

where the last equation follows from $P_{\mathcal{A}}(x^n)$ being a symmetric tensor and (A.1). From (A.1), (A.2) and $\bar{y} \perp x$, we see $\langle x^n, \text{Sym}(Y) \rangle = \langle x^n, Y \rangle = 0$. Since $P_{\mathcal{A}}$ is a self-adjoint projection operator,

$$\langle x^n, P_{\mathcal{A}}(x^n) \rangle = \|P_{\mathcal{A}}(x^n)\|^2. \quad (\text{C.8})$$

Now by Bessel's inequality,

$$\|P_{\mathcal{A}}(x^n)\|^2 \geq \langle P_{\mathcal{A}}(x^n), x^n \rangle^2 + \left\langle P_{\mathcal{A}}(x^n), \frac{\text{Sym}(Y)}{\|\text{Sym}(Y)\|} \right\rangle^2. \quad (\text{C.9})$$

Rearranging (C.9) and using (C.8),

$$\begin{aligned} |\langle P_{\mathcal{A}}(x^n), \text{Sym}(Y) \rangle| &\leq (\|P_{\mathcal{A}}(x^n)\|^2 - \|P_{\mathcal{A}}(x^n)\|^4)^{1/2} \|\text{Sym}(Y)\| \\ &\leq C_\nu \|\text{Sym}(Y)\|. \end{aligned} \quad (\text{C.10})$$

Here (C.10) is due to $\|P_{\mathcal{A}}(x^n)\|^2 = f(x) \leq 1$, the assumption $f(x) \geq \nu$ and the definition

$$C_\nu = \begin{cases} \sqrt{\nu(1-\nu)} & \text{if } \nu > \frac{1}{2} \\ \frac{1}{2} & \text{if } \nu \leq \frac{1}{2}. \end{cases}$$

Substituting (C.10) into (C.7),

$$\frac{1}{2n} y^T \nabla^2 f(x) y \geq -C_\nu \|\text{Sym}(Y)\|. \quad (\text{C.11})$$

The proof is complete if we show $\|\text{Sym}(Y)\| \leq C_n$. We first calculate $\|\text{Sym}(x^{n-2}\bar{y}^2)\|^2$. Denoting by $\binom{[n]}{2}$ the set of subsets of $\{1, \dots, n\}$ of cardinality two, we have

$$\text{Sym}(x^{n-2}\bar{y}^2) = \binom{n}{2}^{-1} \sum_{S \in \binom{[n]}{2}} \pi_S(x^{n-2}\bar{y}^2). \quad (\text{C.12})$$

By this notation, we mean the following: if we let $S = \{s_1, s_2\} \subset \{1, \dots, n\}$, then $\pi_S(x^{n-2}\bar{y}^2)$ permutes the tensor product $x^{n-2}\bar{y}^2$ such that y appears in positions s_1 and s_2 . Now from (A.2), $\bar{y} \perp x$ and $\|x\| = \|y\| = 1$, we obtain

$$\langle \pi_{S_1}(x^{n-2}\bar{y}^2), \pi_{S_2}(x^{n-2}\bar{y}^2) \rangle = \mathbb{1}_{S_1=S_2}. \quad (\text{C.13})$$

Thus

$$\|\text{Sym}(x^{n-2}\bar{y}^2)\|^2 = \binom{n}{2}^{-1} = \frac{2}{n(n-1)}. \quad (\text{C.14})$$

Analogously, $\|\text{Sym}(x^{n-1}\bar{y})\|^2 = \frac{1}{n}$ and $\langle \text{Sym}(x^{n-2}\bar{y}^2), \text{Sym}(x^{n-1}\bar{y}) \rangle = 0$, therefore

$$\|\text{Sym}(Y)\|^2 = \frac{\alpha^2}{n} (\beta^2 b_n^2 + 2\alpha^2(n-1)). \quad (\text{C.15})$$

Letting $t = \alpha^2$, and noting that $\beta^2 = 1 - t$,

$$\begin{aligned} \|\text{Sym}(Y)\|^2 &= \frac{b_n^2}{n} t - \left(\frac{b_n^2 - 2(n-1)}{n} \right) t^2 \\ &\leq \max_{t \in [0,1]} \frac{b_n^2}{n} t - \left(\frac{b_n^2 - 2(n-1)}{n} \right) t^2 \end{aligned} \quad (\text{C.16})$$

The unique critical point of this quadratic function in $t \in \mathbb{R}$ is

$$t^* = \frac{b_n^2}{2b_n^2 - 4(n-1)}. \quad (\text{C.17})$$

On the other hand, for $n \geq 2$, the coefficient of t^2 in (C.16) is negative, i.e., $-\frac{b_n^2 - 2(n-1)}{n} < 0$ (this may be verified by substituting (C.6), rearranging, squaring and estimating roots of a quartic polynomial in n , e.g., with the help of computer algebra software). Therefore the maximum in (C.16) is attained at $t = \min(t^*, 1)$. Further elementary calculations verify $t^* > 1$ if $n \leq 4$ and $t^* < 1$ if $n > 4$. So (C.16) becomes

$$\|\text{Sym}(Y)\|^2 \leq \begin{cases} \frac{2(n-1)}{n} & \text{if } n \leq 4 \\ \frac{b_n^4}{4nb_n^2 - 8n(n-1)} & \text{if } n > 4. \end{cases} \quad (\text{C.18})$$

Equation (4.10) now follows from:

$$\begin{aligned} \frac{b_n^4}{4nb_n^2 - 8n(n-1)} &= \frac{\sqrt{2}(2n-1)(3n-1-2\sqrt{n(2n-1)})}{\sqrt{n(3-11n+12n^2-(8n-4)\sqrt{n(2n-1)})}} \\ &\leq (2-\sqrt{2})\sqrt{n} \end{aligned} \quad (\text{C.19})$$

when $n > 4$. This completes the proof of Lemma 4.4. \square

C.2 Statement of convergence result [70, Theorem 2.3]

Here we record the “meta convergence result” of Schneider–Uschmajew. To prove that SS-HOPM converges unconditionally, we will verify the conditions in this theorem. For convenience, we will indicate a corollary and slightly weaker result than [70, Theorem 2.3].

Within this appendix, $\mathcal{M} \subseteq \mathbb{R}^N$ is a closed subset. For $x \in \mathcal{M}$, the *tangent cone* to \mathcal{M} at x is defined to be

$$T_x \mathcal{M} = \{\xi \in \mathbb{R}^N : \exists (x_n) \subseteq \mathcal{M}, \exists (a_n) \subseteq \mathbb{R}_{>0} \text{ s.t. } x_n \rightarrow x, a_n(x_n - x) \rightarrow \xi\} \subseteq \mathbb{R}^N. \quad (\text{C.20})$$

This is a closed cone, in general not convex (although when \mathcal{M} is an embedded submanifold, this coincides with the usual embedded tangent space to \mathcal{M} at x).

Given a vector $g \in \mathbb{R}^N$, introduce the notation

$$g^- := \max_{\substack{\xi \in T_x \mathcal{M} \\ \|\xi\| \leq 1}} -g^\top \xi \in \mathbb{R}. \quad (\text{C.21})$$

Thus $g^- \in \mathbb{R}$ is the norm of any orthogonal projection of $-g$ to $T_x \mathcal{M}$.

Theorem C.1 [70, Theorem 2.3] *Let $\mathcal{M} \subseteq \mathbb{R}^N$ be a closed set that is locally the image of a real analytic map out of Euclidean space. Let $\mathcal{D} \subseteq \mathbb{R}^N$ be an open neighborhood of \mathcal{M} . Suppose $f : \mathcal{D} \rightarrow \mathbb{R}$ is a real analytic function that is bounded below. Consider the problem*

$$\max_{x \in \mathcal{M}} f(x). \quad (\text{C.22})$$

Consider a sequence $(x_k) \subseteq \mathcal{M}$ seeking to solve (C.22), and set

$$f_k = f(x_k), \quad \nabla f_k = \nabla f(x_k), \quad (\nabla f_k)^- \text{ as in (C.21),} \quad \text{and} \quad T_k \mathcal{M} = T_{x_k} \mathcal{M}. \quad (\text{C.23})$$

Consider the following three assumptions:

(A1) *There exists $\sigma > 0$ such that for k large enough it holds*

$$f_{k+1} - f_k \geq \sigma (\nabla f_k)^- \|x_{k+1} - x_k\|.$$

(A2) *For large enough k it holds*

$$(\nabla f_k)^- = 0 \implies x_{k+1} = x_k.$$

(A3) *There exists $\rho > 0$ such that for large enough k it holds*

$$\|x_{k+1} - x_k\| \geq \rho (\nabla f_k)^-.$$

*If (x_k) satisfies the assumptions **(A1)** and **(A2)**, then any cluster point $x_* \in \mathcal{M}$ of (x_k) must actually be the limit point of (x_k) . Further if **(A3)** holds, then there exists constants $C > 0$ and $\tau > 1$ such that*

$$\|x_k - x_*\| < Ck^{-\tau}. \quad (\text{C.24})$$

Moreover, $(\nabla f_k)^- \rightarrow 0$ as $k \rightarrow \infty$.

The proof in [70] uses the real analytic hypotheses on \mathcal{M} and f by invoking the Łojasiewicz inequality for real analytic functions. We note that typically knowledge of τ is not accessible.

C.3 Proof of Lemma 4.6

Simple calculations show

$$DG(x_*) = \frac{\nabla^2 F(x_*)}{\|\nabla F(x_*)\|} \left(I - \left(\frac{\nabla F(x_*)}{\|\nabla F(x_*)\|} \right) \left(\frac{\nabla F(x_*)}{\|\nabla F(x_*)\|} \right)^T \right) = \frac{\nabla^2 F(x_*)}{\|\nabla F(x_*)\|} (I - x_* x_*^T). \quad (\text{C.25})$$

We may also verify

$$\nabla F(x_*) = \nabla f(x_*) + \gamma(x_*^T x_*)^{n-1} x_* = \nabla f(x_*) + \gamma x_*, \quad (\text{C.26})$$

and likewise

$$\begin{aligned} \nabla^2 F(x_*) &= \nabla^2 f(x_*) + (2n-2)\gamma(x_*^T x_*)^{n-2} x_* x_*^T + \gamma(x_*^T x_*)^{n-1} I \\ &= \nabla^2 f(x_*) + (2n-2)\gamma x_* x_*^T + \gamma I. \end{aligned} \quad (\text{C.27})$$

Substituting (C.26) and (C.27) into (C.25) implies the result. \square

C.4 Statement of center-stable manifold theorem

Theorem C.2 (Central Stable Manifold Theorem, [72, Theorem III.7(2)]) *Let 0 be a fixed point for a C^r local diffeomorphism $\phi : U \rightarrow \mathbb{R}^n$ where U is a neighborhood of zero in \mathbb{R}^n ($r \geq 1$). Let $E^s \oplus E^c \oplus E^u$ be the invariant splitting of \mathbb{R}^n into the generalized eigenspaces of $D\phi(0)$ corresponding to eigenvalues of absolute values less than one, equal to one, and greater than one. Then there exists a neighborhood B of 0, and a C^r embedded disk $W_{\text{loc}}^{\text{cs}} \subset B$ that is tangent to $E^s \oplus E^c$ at 0, such that $\phi(W_{\text{loc}}^{\text{cs}}) \cap B \subset W_{\text{loc}}^{\text{cs}}$ and $\cap_{k=0}^{\infty} \phi^{-k}(B) \subset W_{\text{loc}}^{\text{cs}}$. One calls $W_{\text{loc}}^{\text{cs}}$ the local stable center manifold of ϕ at 0.*

C.5 Proof of Lemma 4.7

Suppose x_* fails the second-order optimality conditions. Remark 4.3 then implies that:

$$(I - x_* x_*^T) (\nabla^2 f(x_*) + \mu_* I) (I - x_* x_*^T) \not\leq 0. \quad (\text{C.28})$$

In other words, there exists $y \in \mathbb{R}^L \setminus \{0\}$ lying in the tangent space $T_{x_*}(\mathbb{S}^{L-1})$, i.e., $\langle y, x_* \rangle = 0$, satisfying $\nabla^2 f(x_*)y = \lambda y$ and $\lambda + \mu_* > 0$. By Lemma 4.6, y is also a eigenvector of $DG(x_*)$ with eigenvalue $\frac{\lambda + \gamma}{-\mu_* + \gamma}$. Multiplying (C.26) and (4.25) on the left by x_*^T shows $-\mu_* + \gamma = F(x_*)$, which is positive by assumption on γ . So the eigenvalue of $DG(x_*)$ associated with y is $\frac{\lambda + \gamma}{-\mu_* + \gamma}$, and crucially, $\frac{\lambda + \gamma}{-\mu_* + \gamma} > 1$ since $\lambda + \mu_* > 0$.

Now, we wish to apply the center-stable manifold theorem to G (the statement is given above in Appendix C.4). To apply this theorem, we need to map locally \mathbb{S}^{L-1} to \mathbb{R}^{L-1} . Let O be a $L \times L-1$ matrix with orthogonal columns, such that $O^T x_* = 0$. Define $\Gamma : \mathbb{R}^{L-1} \rightarrow \mathbb{S}^{L-1}$ such that $\Gamma(y) = \frac{x_* + Oy}{\|x_* + Oy\|}$, and note that $\Gamma(0) = x_*$ and $D\Gamma(0) = O(I - x_* x_*^T) = O$. Now let $\tilde{G} = \Gamma^{-1} \circ G \circ \Gamma$, and note that $D\tilde{G}(0) = O^T DG(x_*) O$. Since O is orthogonal to x_* , and $DG(x_*)x_* = (\mu_* + \gamma)x_*$, thus x_* is an eigenvector of $DG(x_*)$, we have that if $y \perp x_*$ is an eigenvector of $DG(x_*)$, with associated eigenvalue λ , then Oy is an eigenvector of $D\tilde{G}(0)$ with the same eigenvalue. Finally, the argument from the previous paragraph implies that $D\tilde{G}(0)$ has at least one eigenvalue greater than 1. Letting W be the subspace spanned by all eigenvectors of $D\tilde{G}(0)$ with associated eigenvalue at most 1, we then conclude $\dim(W) < L-1$.

By our choice of γ , all eigenvalues of $D\tilde{G}(0)$ are positive, which implies that \tilde{G} is locally a C^∞ diffeomorphism. Therefore, the center-stable manifold theorem (stated in Theorem C.2)

implies that there exists an open neighborhood $B_0 \subset \mathbb{R}^{L-1}$ of 0, and a smoothly embedded disc $D_0 \subset B$, tangent to W , and therefore of *dimension strictly less than* $L - 1$ such that:

$$(x \in B_0 \text{ and } \tilde{G}^k(x) \in B_0 \forall k \geq 1) \Rightarrow x \in D_0. \quad (\text{C.29})$$

The result now follows from taking $B_{x_*} = \Gamma(B_0)$, $D_{x_*} = \Gamma(D_0)$ and noticing that

$$\Gamma\left(\left\{x \in B_0 : \tilde{G}^k(x) \in B_0 \forall k \geq 1\right\}\right) = \left\{x \in B_{x_*} : G^k(x_*) \in B_0 \forall k \geq 1\right\}.$$

□

C.6 Proof of Lemma 4.8

We first prove G is a local diffeomorphism everywhere. Fix $x_* \in \mathbb{S}^{L-1}$ (which might not be a local maximum), and define \tilde{G} as in the proof of Lemma 4.7. We then have (C.25) together with the gradient implies that

$$D\tilde{G}(0) = O^T \frac{\nabla^2 F(x_*)}{\|\nabla F(x_*)\|} \left(I - x_* x_*^T\right) O = O^T \frac{\nabla^2 F(x_*)}{\|\nabla F(x_*)\|} O,$$

where we used that $x_*^T O = 0$. Therefore $D\tilde{G}(0)$, and since F is strictly convex, the eigenvalues of $D\tilde{G}(0)$ are positive which implies that \tilde{G} is locally a diffeomorphism around 0, and G is locally a diffeomorphism around x_* . Now identify $\tilde{G} = \tilde{G}_{x_*}$ and $\Gamma = \Gamma_{x_*}$. Then for all x^* there exist neighborhoods B_{x_*} such that G^{-1} is well defined at these neighborhoods, and if $A \subset G^{-1}(A)$

$$\mu(G^{-1}(A)) = \int_{G^{-1}(A)} 1 d\mu = \int_{\tilde{G}_{x_*}^{-1}(\Gamma_{x_*}^{-1}(A))} 1 d\mu = \int_{\Gamma_{x_*}^{-1}(A)} |D\tilde{G}| d\mu,$$

by the change of measure theorem. By compactness of \mathbb{S}^{L-1} , and since $D\tilde{G}$ is continuous, there exists $M \in \mathbb{R}$ such that $|D\tilde{G}(x)| \leq M$ for all $x \in \mathbb{S}^{L-1}$, and since \mathbb{S}^{L-1} is second-countable and $\bigcup_{x^* \in \mathbb{S}^{L-1}} B_{x^*}$ is a cover of \mathbb{S}^{L-1} , there is a countable sub-cover, that is, a countable set $\mathcal{C} \subset \mathbb{S}^{L-1}$, such that $\bigcup_{x^* \in \mathcal{C}} B_{x^*} = \mathbb{S}^{L-1}$. This then implies that

$$\mu(G^{-1}(A)) \leq \sum_{x^* \in \mathcal{C}} \mu(G^{-1}(A \cap B_{x^*})) \leq M \sum_{x^* \in \mathcal{C}} \mu(A \cap B_{x^*}) = 0,$$

which completes the proof. □

C.7 Proof of Lemma 4.10

By first-order optimality condition (4.25), we have $G(x_*) = x_*$, for G in (4.23) where (4.12) is $x_{k+1} = G(x_k)$. Now consider the Jacobian $DG|(x_*)$, for $G|$ the restriction of G to \mathbb{S}^{L-1} . Here $DG|(x_*)$ is a linear map $T_{x_*}(\mathbb{S}^{L-1}) \rightarrow T_{x_*}(\mathbb{S}^{L-1})$, and $T_{x_*}(\mathbb{S}^{L-1}) = x_*^\perp$. As in the proof of Theorem 4.9, the eigenvalues of $DG|(x_*)$ are $\frac{\lambda + \gamma}{-\mu_* + \gamma}$, where λ ranges over the eigenvalues of the Hessian matrix $\nabla^2 f(x_*)$ (restricted to x_*^\perp). By assumptions, we have $\lambda + \gamma > 0$ and $-\mu_* + \gamma > 0$. So, all eigenvalues of $DG(x_*)$ are strictly less than 1 in magnitude, if and only if $\lambda + \mu_* < 0$ for each eigenvalue of $\nabla^2 f(x_*)$. However, this holds by the second-order strictness condition (4.32). Therefore, $DG|(x_*)$ is a contractive linear mapping, and $G|$ is locally contractive around x_* . As $G(x_*) = x_*$, this implies local linear convergence to x_* as desired, see [69, p. 18]. □

C.8 Proof of Proposition 4.11

Proof Relabeling if necessary, we assume $i = 1$. Applying a suitable orthogonal transformation to \mathbb{R}^L , we may further assume $a_1 = e_1$ (the first standard basis vector). Recalling Lemma 2.2, we set $l_j = \Phi(a_j)$ and p for the linear forms in $\mathbb{R}[X_1, \dots, X_L]$ corresponding to the vectors a_j and x , and so in particular $l_1 = X_1$. The condition $x \perp a_1$ corresponds to $p \in \mathbb{R}[X_2, \dots, X_L]$ (that is, the variable X_1 does not appear in p). Supposing this, it suffices to assume

$$X_1^{n-1}p = \alpha_1 l_1^n + \dots + \alpha_R l_R^n \quad (\text{C.30})$$

for some scalars $\alpha_1, \dots, \alpha_R \in \mathbb{R}$, and then to deduce $p = 0$. Note (C.30) is an equality between degree n forms. By genericity, the variable X_1 appears in each l_2, \dots, l_R (this is equivalent to $\langle a_1, a_j \rangle \neq 0$ for each j). Thus dividing by appropriate nonzero coefficients, and then scaling α_j , we may assume the coefficient of X_1 in l_j exactly equals 1, for all j .

Now plug in $X_1 = 0$ into (C.30). The LHS is identically 0, while the RHS is an expression in the other variables, X_2, \dots, X_L :

$$0 = \alpha_2 \tilde{l}_2^n + \dots + \alpha_R \tilde{l}_R^n \quad (\text{C.31})$$

where $\tilde{l}_j := l_j(0, X_2, \dots, X_L) \in \mathbb{R}[X_2, \dots, X_L]$.

If $n > 2$, we differentiate (C.30) with respect to X_1 , obtaining

$$(n-1)X_1^{n-2}p = n\alpha_1 l_1^{n-1} + \dots + n\alpha_R l_R^{n-1}. \quad (\text{C.32})$$

Plugging in $X_1 = 0$ into (C.32) and dividing by n yields

$$0 = \alpha_2 \tilde{l}_2^{n-1} + \dots + \alpha_R \tilde{l}_R^{n-1} \quad (\text{C.33})$$

with \tilde{l}_j as above.

Continuing in this way, differentiating (C.30) repeatedly with respect to X_1 and subsequently plugging in $X_1 = 0$, it follows that

$$0 = \alpha_2 \begin{bmatrix} \tilde{l}_2^n \\ \tilde{l}_2^{n-1} \\ \vdots \\ \tilde{l}_2^2 \end{bmatrix} + \dots + \alpha_R \begin{bmatrix} \tilde{l}_R^n \\ \tilde{l}_R^{n-1} \\ \vdots \\ \tilde{l}_R^2 \end{bmatrix}. \quad (\text{C.34})$$

Here (C.34) is a linear dependence between elements in the vector space $\mathbb{R}[X_2, \dots, X_L]_n \oplus \dots \oplus \mathbb{R}[X_2, \dots, X_L]_2$, that is, between vectors of homogeneous polynomials in $L-1$ variables in which the first entries are degree n , the second entries degree $n-1$ and so on until the last entries of degree 2. By homogenizing, we regard this as a linear dependence of polynomials in L variables. Specifically, using $l_j = \tilde{l}_j + X_1$ and the binomial theorem, it is equivalent that

$$0 = \alpha_2 \mathcal{P}_{\mathcal{B}}(l_2^n) + \dots + \alpha_R \mathcal{P}_{\mathcal{B}}(l_R^n), \quad (\text{C.35})$$

where $\mathcal{B} = \langle X_1^n, X_1^{n-1}X_2, \dots, X_1^{n-1}X_R \rangle^\perp \subset \mathbb{R}[X_1, \dots, X_L]$ is the orthogonal complement to the space of polynomials divisible by X_1^{n-1} and where the mapping $\mathcal{P}_{\mathcal{B}} : \mathbb{R}[X_1, \dots, X_L]_n \rightarrow \mathcal{B}$ is orthogonal projection (orthogonality is with respect to the inner product on polynomials induced by Φ in Lemma 2.2).

To finish, it is enough to know that, owing to Zariski genericity of l_2, \dots, l_R , the polynomials $\mathcal{P}_{\mathcal{B}}(l_2^n), \dots, \mathcal{P}_{\mathcal{B}}(l_R^n)$ are linearly independent. For then, (C.35) implies $\alpha_2 = \dots = \alpha_R = 0$, and whence $p = 0$ from (C.30), as desired. However this is true by elementary facts from classical algebraic geometry, namely that the Veronese variety is non-degenerate and that the linear projection of any non-degenerate variety is also non-degenerate. \square

D Determining the null space of \mathbf{Q}

In this appendix, we explain how we calculate the null space of \mathbf{Q} in the LOCAL COMPONENT step of Algorithm 2 (as sketched in Remark 5.9). The calculation is based on the eigendecomposition of $\mathbf{Q}^T \mathbf{Q}$. Since, by definition, $\text{Sym}(\mathcal{T}_L^n) = \mathcal{A} \oplus \mathcal{A}^\perp$, we have:

$$\begin{aligned} y^T \mathbf{Q}^T \mathbf{Q} y &= \sum_{j=1}^K \langle Q_j, x_*^{\otimes n-1} \otimes y \rangle^2 \\ &= \|P_{\mathcal{A}^\perp}(x^{\otimes n-1} \otimes y)\|^2 \\ &= \|P_{\text{Sym}}(x^{\otimes n-1} \otimes y)\|^2 - \|P_{\mathcal{A}}(x^{\otimes n-1} \otimes y)\|^2. \end{aligned} \quad (\text{D.1})$$

Using an argument similar to the one for (C.12), we see

$$\begin{aligned} \|P_{\text{Sym}}(x^{\otimes n-1} \otimes y)\|^2 &= \frac{1}{n} \|y\|^2 + \frac{n-1}{n} (x^T y)^2 \\ &= y^T \left(\frac{1}{n} I + \frac{n-1}{n} x x^T \right) y, \end{aligned} \quad (\text{D.2})$$

and therefore

$$\mathbf{Q}^T \mathbf{Q} = \frac{1}{n} I + \frac{n-1}{n} x x^T - \mathbf{V}^T \mathbf{V}. \quad (\text{D.3})$$

Here the rows of \mathbf{V} are defined by

$$\mathbf{V}_j = V_j \cdot x_*^{\otimes n-1}, \quad j = 1, \dots, \mathbf{R} \quad (\text{D.4})$$

where V_j for $j = 1, \dots, \mathbf{R}$ form an orthonormal basis for \mathcal{A} .

E A heuristic for determining the noise level in GPCA

Suppose we have N points Y_1, \dots, Y_N in \mathbb{R}^L which are obtained by adding Gaussian noise with variance σ^2 to Z_1, \dots, Z_N , that is,

$$Y_i = Z_i + \xi_i, \quad i = 1, \dots, N, \quad (\text{E.1})$$

where $\xi_i \sim \mathcal{N}(0, \sigma^2 I_L)$. Letting

$$M_n^Z := \sum_{i=1}^N Z_i^{\otimes n} \quad \text{and} \quad M_n^Y := \mathbb{E} \left[\sum_{i=1}^N Y_i^{\otimes n} \right], \quad (\text{E.2})$$

we have by [64, Lemma 3.3.2]

$$M_2^Y = M_2^Z + \sigma^2 I_L, \quad (\text{E.3})$$

where I_L is the identity matrix, and

$$M_4^Y = M_4^Z + 6\sigma^2 \text{Sym}(M_2^Z \otimes I_L) + 3\sigma^4 \text{Sym}(I_L \otimes I_L), \quad (\text{E.4})$$

or equivalently

$$M_4^Y = M_4^Z + 6\sigma^2 \text{Sym}(M_2^Y \otimes I_L) - 3\sigma^4 \text{Sym}(I_L \otimes I_L). \quad (\text{E.5})$$

If the points Z_1, \dots, Z_N live on a union of subspaces, such that the corresponding square flattening rank is less than $\binom{L+1}{2}$, then the smallest eigenvector of $\text{mat}(M_4^Z)$ should be 0. Therefore, we will estimate σ as the minimum $t \in \mathbb{R}$ such that the smallest eigenvalue of

$$\text{mat} \left(M_4^Y - 6t^2 \text{Sym}(M_2^Y \otimes I_L) + 3t^4 \text{Sym}(I_L \otimes I_L) \right) \quad (\text{E.6})$$

is 0. Letting λ and v be the smallest eigenvalue and corresponding normalized eigenvector of $\text{mat}(M_4^Y)$ and μ the smallest eigenvalue of $\text{mat}(M_4^Z)$, we have, by matrix eigenvalue perturbation theory [58]:

$$\mu \approx \lambda - 6t^2 a_1 + 3t^4 a_2, \quad (\text{E.7})$$

where

$$a_1 = v^T \text{mat}(\text{Sym}(M_2^Y \otimes I_L))v \quad \text{and} \quad a_2 = v^T \text{mat}(\text{Sym}(I_L \otimes I_L))v. \quad (\text{E.8})$$

We now set $\mu = 0$ in (E.7) and solve for t . This yields the following estimator:

$$\hat{\sigma} = \sqrt{\frac{a_1 - \sqrt{a_1^2 - a_2 \lambda / 3}}{a_2}}. \quad (\text{E.9})$$