

ON THE FAST ROTATION ASYMPTOTICS OF A NON-HOMOGENEOUS INCOMPRESSIBLE MHD SYSTEM

*Dimitri Cobb*¹ and *Francesco Fanelli*²

^{1, 2} UNIVERSITÉ DE LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1

Institut Camille Jordan – UMR 5208

43 blvd. du 11 novembre 1918, F-69622 Villeurbanne cedex, FRANCE

¹ `cobb@math.univ-lyon1.fr`

² `fanelli@math.univ-lyon1.fr`

December 24, 2021

Abstract

This paper is devoted to the analysis of a singular perturbation problem for a 2-D incompressible MHD system with density variations and Coriolis force, in the limit of small Rossby numbers. Two regimes are considered. The first one is the quasi-homogeneous regime, where the densities are small perturbations around a constant state. The limit dynamics is identified as an incompressible homogeneous MHD system, coupled with an additional transport equation for the limit of the density variations. The second case is the fully non-homogeneous regime, where the densities vary around a general non-constant profile. In this case, in the limit, the equation for the magnetic field combines with an underdetermined linear equation, which links the limit density variation function with the limit velocity field. The proof is based on a compensated compactness argument, which enables us to consider general ill-prepared initial data. An application of Di Perna-Lions theory for transport equations allows to treat the case of density-dependent viscosity and resistivity coefficients.

2010 Mathematics Subject Classification: 35Q35 (primary); 35B25, 76U05, 35B40, 76W05 (secondary).

Keywords: incompressible MHD; density variations; variable viscosity and resistivity; Coriolis force; low Rossby number; singular perturbation problem.

1 Introduction

In this article, we study the $\epsilon \rightarrow 0^+$ asymptotics of the following non-homogeneous incompressible MHD system with Coriolis force:

$$(1) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \frac{1}{\epsilon} \nabla \pi + \frac{1}{\epsilon} \rho u^\perp = \operatorname{div}(\nu(\rho) \nabla u) + \operatorname{div}(b \otimes b) - \frac{1}{2} \nabla |b|^2 \\ \partial_t b + \operatorname{div}(u \otimes b) - \operatorname{div}(b \otimes u) = \nabla^\perp(\mu(\rho) \operatorname{curl}(b)) \\ \operatorname{div} u = \operatorname{div} b = 0. \end{cases}$$

These equations are set in a two dimensional domain Ω , which is either the plane \mathbb{R}^2 or the torus \mathbb{T}^2 . The vector fields u and b are the velocity and the magnetic fields, the scalar fields

$\rho \geq 0$ and π represent the density and the pressure fields and ν and μ are two functions defined on \mathbb{R}_+ . The notation u^\perp refers the rotation of angle $\pi/2$ of the vector u : in other words, if $u = (u_1, u_2)$, then $u^\perp = (-u_2, u_1)$. Analogously, we have set $\nabla^\perp = (-\partial_2, \partial_1)$. We have also defined $\text{curl } u = \partial_1 u_2 - \partial_2 u_1$ to be the curl of the 2-D vector u . As we will see, since the first equation is written up to a gradient field, the term $\nabla|b|^2/2$ does not appear in the weak form of the equations.

The main goal of this article is to show that solutions $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$ to system (1) converge (in some way) to some functions (ρ, u, b) , and to describe the limit dynamics by proving that the limit (ρ, u, b) solves an evolution PDE system.

1.1 General physical remarks

Magnetohydrodynamic models are used whenever describing a fluid which is subject to the magnetic field it generates through its own motion. Examples of such fluids range from the industrial scale, with plasma confinement in fusion research or some types of electrolytics, to the geophysical or astrophysical scale, with atmospheric plasmas, planetary mantle convection or the solar interior. Their mathematical study thus combines the Navier-Stokes and the Maxwell equations.

We focus on fluids on which the Coriolis force $\epsilon^{-1}\rho u^\perp$ has a major influence compared to the kinematics of the said fluid, such as large-scale fluids evolving on a celestial body. The importance of this effect is measured by the Rossby number of the fluid $Ro = \epsilon$, the condition $\epsilon \ll 1$ defining the regime of large-scale planetary or stellar fluid dynamics. At the mathematical level, in the limit $\epsilon \rightarrow 0^+$, the Coriolis force can only be balanced by the pressure term, which is reflected by the ϵ^{-1} factor in front of $\nabla\pi$.

1.1.1 Derivation of the equations

Let us give a few additional details concerning the derivation of the MHD equations¹. We conduct these computations in the physically relevant three-dimensional setting $\Omega = \mathbb{R}^3$.

First of all, the Laplace force exerted on the fluid is $f_L = j \times b$, where j is the electric current density. We further assume the fluid viscosity $\nu(\rho)$ to depend on the density ρ , the precise nature of the function ν depending on the exact composition of the fluid (alternate models include anisotropic scaling in the viscosity to take into account the joint effects of turbulence and the asymmetry induced by the rotation, see [6], [10] for more on this topic). Writing Newton's law for a density-dependent fluid, we get

$$\rho \partial_t u + \rho(u \cdot \nabla)u + \frac{1}{\epsilon} \nabla \pi + \frac{1}{\epsilon} \mathfrak{C}[\rho, u] - \text{div}(\nu(\rho) \nabla u) = j \times b,$$

where $\mathfrak{C}[\rho, u] = e^3 \times \rho u = (-\rho u_2, \rho u_1, 0)$ is the (three-dimensional) Coriolis force. The rotation axis is taken constant, parallel to the vertical unit vector $e^3 = (0, 0, 1)$. We will comment more on this in the next paragraph.

The fluid is assumed to be *non-relativistic*, with negligible velocities $|u| \ll c$ when compared to the speed of light. This justifies the use of an electrostatic approximation in the Maxwell equations, which are simplified by omitting the time derivative of the electric field. Obviously, this is not a wild assumption since we intend to work on planetary or stellar fluids subject to the body's rotation (see also [10], [27]). Thus, Ampère's circuital law reads

$$\text{curl}(b) = j + \partial_t e \approx j,$$

where e is the electric field. The electrical resistivity is described by Ohm's law, which links the electrical current j to the electrical field e and the other physical quantities:

$$j = \sigma(\rho)(e + u \times b).$$

¹In order to simplify notations, we assume the electrical permittivity and the magnetic permeability to be of unit value, which can be done by working in an appropriate set of units.

Note that we have assumed that the conductivity $\sigma(\rho)$ depends on the density, the precise nature of the function σ depending on the exact composition of the fluid. Combining the above relation with the Maxwell-Faraday equation $\partial_t b = -\text{curl}(e)$ gives a relation describing the evolution of the magnetic field:

$$\partial_t b = -\text{curl} \left(\frac{1}{\sigma(\rho)} \text{curl}(b) - u \times b \right).$$

By noting $\mu(\rho) = 1/\sigma(\rho)$ the electrical resistivity, we thus get

$$\partial_t b + (u \cdot \nabla) b - (b \cdot \nabla) u = -\text{curl}(\mu(\rho) \text{curl}(b)).$$

Next, Gauss's law for magnetism, which rules out the possibility of magnetic monopoles, gives the divergence-free condition $\text{div}(b) = 0$.

Finally, we assume the fluid to be incompressible, so that $\text{div}(u) = 0$. Therefore, the mass conservation equation reads

$$\partial_t \rho + \text{div}(\rho u) = \partial_t \rho + u \cdot \nabla \rho = 0.$$

Putting everything together, and using the conservative form of the equations, we obtain the following three-dimensionnal MHD system,

$$(2) \quad \begin{cases} \partial_t \rho + \text{div}(\rho u) = 0 \\ \partial_t(\rho u) + \text{div}(\rho u \otimes u) + \nabla \pi + \frac{1}{\epsilon} \mathfrak{C}[\rho, u] = \text{div}(\nu(\rho) \nabla u) + \text{div}(b \otimes b) - \frac{1}{2} \nabla(|b|^2) \\ \partial_t b + \text{div}(u \otimes b - b \otimes u) = -\text{curl}(\mu(\rho) \text{curl}(b)) \\ \text{div}(b) = \text{div}(u) = 0, \end{cases}$$

of which system (1) is the 2-D equivalent.

1.1.2 Physical relevance of the system

This paragraph is devoted to a few critical remarks concerning system (1), and the physical setting that led to its derivation.

First of all, the model neglects any effect due to temperature variations, which is a debatable simplification, even in the case of non-conducting fluids. For instance, ocean water density is an intricate function of the pressure, salinity and temperature, and the temperature of air masses plays a major role in weather evolution. In those cases, dependence on the pressure is often neglected, and both temperature and salinity are assumed to evolve through a diffusion process (see [7], Chapter 3). For conducting fluids, which are generally heated magma or plasmas, the temperature is expected to play an even greater role. However, the equations, as they are, already provide interesting challenges and are widely used by physicists for practical purposes.

Secondly, we spend a few words concerning the two-dimensional setting. Our main motivation for restricting to 2-D domains is purely technical: in our analysis, we face similar difficulties as in [16], devoted to the fast rotation asymptotics for density-dependent incompressible Navier-Stokes equations in dimension $d = 2$. In particular, the fast rotation limit for incompressible non-homogeneous fluids in 3-D is a widely open problem (see more details here below). However, let us notice that one of the common features of highly rotating fluids is to be, in a first approximation, planar: the fluid is devoid of vertical motion and the particles move in columns. This property is known as the Taylor-Proudman theorem (see [7], [28] for useful insight). Therefore, the 2-D setting is in itself a relevant approximation for geophysical fluids.

At this point, note that equations (1) *per se* do not describe a conducting fluid confined to a quasi-planar domain. If that were the case, the magnetic field would circulate *around* the current lines, hence being orthogonal to the plane of the fluid, assuming the form $b = b_3(t, x)e^3$ for some

scalar function b_3 . Our problem, which involves a 2-D magnetic field $b = (b_1, b_2)$ is a projection of the full three-dimensional MHD system (2). Taking this step away from the physical problem brings us closer to the actual form of the physically relevant 3-D problem, and we hope it will provide a step towards its understanding.

Next, we remark that we have taken a quite simple form for the Coriolis force: namely, $\mathfrak{C}[\rho, u] = \rho u^\perp$. This means that the rotation axis is constant and normal to the plane where the fluid moves. Of course, more complicated choices are possible. However, on the one hand this form for the rotation term is physically consistent with a fluid evolving at mid-latitudes, in a region small enough compared to the radius of the planetary or stellar body. On the other hand, this choice is quite common in mathematical studies, and the obtained model is already able to explain several physical phenomena.

Finally, we point out that the incompressibility assumption $\operatorname{div}(u) = 0$ is a valid approximation for flows in the ocean and in the atmosphere, and we will assume it. On the other hand, by our choice of considering domains with a very simple geometry, we completely avoid boundary effects.

1.2 Previous mathematical results on fast rotating fluids

The mathematical study of rotating fluids is by no means new in the mathematical literature. It has started in the 1990s with the pioneering works [1]-[2]-[3] of Babin, Mahalov and Nikolaenko, and has since been deeply investigated, above all for models of homogeneous incompressible fluids. We refer to book [6] for a complete treatment of the incompressible Navier-Stokes equations with Coriolis force, and for further references on this subject.

The study of fast rotation asymptotics for non-homogeneous fluids has a much more recent history. However, efforts have mainly been focused on compressible fluid models: see e.g. [18], [17], [20], [19], [13], [24]. We refer to [15] for additional details and further references, as well as for recent developments. On the contrary, not so many results are available for density-dependent incompressible fluids. To the best of our knowledge, the only work in this direction is [16], treating the case of the non-homogeneous Navier-Stokes equations in two-dimensional domains. The reason for such a gap between the compressible and the (non-homogeneous) incompressible cases is that the coupling between the mass and the momentum equation is weaker in the latter situation than in the former one. As a consequence, less information is available on the limit points of the sequences of solutions, and taking the limit in the equations becomes a harder task. This explains also the lack of results for 3-D incompressible flows with variable densities.

The case of rotating MHD equations has also received some attention in the past years. Once again, most of the available results concern the case of homogeneous flows: for instance, we mention papers [10] and [30], concerning the stability of boundary layers in homogenous rotating MHD, and [27], about the stabilising effect the rotation has on solution lifespan. See also references therein, as well as Chapter 10 of [6], for further references. On the density-dependent side, fast rotating asymptotics has recently been conducted in [23] for compressible flows, in two space dimensions. We point out that the approach of [23] is based on relative entropy estimates; if on the one hand this method enables to consider also a vanishing viscosity and resistivity regime, on the other hand it requires to assume well-prepared initial data.

1.3 Overview of the main results of the paper

Our main motivation here is to extend the results of [16] to the case of the MHD equations. Therefore, we choose to work with incompressible density-dependent fluids, see system (1). More precisely, we study the fast rotation asymptotics in two different regimes: the *quasi-homogeneous* regime (meaning that the initial densities are small variations of a constant state) and the *fully non-homogeneous* regime (when the initial densities are perturbations of a fixed *non-constant* profile, in the sense of relation (11) below). In the former case, the limit dynamics is identified as a homogeneous incompressible MHD system, coupled (via a lower order term) with a pure

transport equation for the limit density variation function. In the latter case, we show convergence to an underdetermined equation, expressed in terms of the vorticity of the limit velocity field and the limit density oscillation function. The fact that the limit system is underdetermined can be viewed as an expression of the weaker coupling we mentioned above, between the mass and momentum equations. In order to prove our results, we will resort to the techniques of [16], based on a compensated compactness argument, which allows us to consider general *ill-prepared* initial data. Roughly speaking, compensated compactness consists in exploiting the structure of the equations (written in the form of a wave system governing oscillations, which propagate in the form of Poincaré-Rossby waves), in order to find special algebraic cancellations and relations which allow to pass to the limit in the non-linear terms, even in absence of strong convergence. That technique goes back to the pioneering work [26] by P.-L. Lions and Masmoudi, where the authors dealt with the incompressible limit of the compressible Navier-Stokes equations; it was later adapted by Gallagher and Saint-Raymond in [21] to the context of fast rotating fluids, and then broadly exploited in similar studies (see e.g. [17], [14] and [15]).

We remark that, in the fully non-homogeneous regime, the limit equation (combining the mass and momentum equations of the primitive system) is *linear* in the unknowns. This is a remarkable property, which is however now quite well-understood (see e.g. [21], [17]): let us give an insight of it. We will be able to show that, if the initial densities are small perturbations around a non-constant state (say) ρ_0 , then, at any later time, the solutions stay close (in a suitable topology, but quantitatively, in powers of ϵ) to the same state ρ_0 . Notice that this property is not obvious at all, as the densities satisfy a pure transport equation by the velocity fields. Anyhow, as a consequence the limit density profile is exactly the initial reference state ρ_0 . Roughly speaking, this fact restricts much more the limit motion than in the case when the target density is constant, since the kernel of the singular perturbation operator is smaller. The additional constraint implies that the average process (convergence in the weak formulation of the equations) tends to kill the convective term in the limit $\epsilon \rightarrow 0^+$.

As a last comment, let us point out that system (1) differs from the Navier-Stokes-Coriolis equations of [16] in a crucial way: namely, the density-dependent viscosity and resistivity terms, respectively $\operatorname{div}(\nu(\rho)\nabla u)$ and $\nabla^\perp(\mu(\rho)\operatorname{curl}(b))$, introduce new difficulties in the analysis. Indeed, the methods of [16], which rely on compactness of the densities in spaces of negative index of regularity, are insufficient to take the limit in those non-linear terms: overcoming this obstacle requires almost everywhere convergence of the densities. Now, the sought almost everywhere convergence is implied by strong convergence in suitable Lebesgue spaces, which we achieve by using well-posedness results on linear transport equations proved by Di Perna and P.-L. Lions [12]. Besides, we point out that strong convergence of the densities has the additional advantage of providing simpler proofs: where the analysis of [16] relies on paradifferential calculus to obtain convergence of some quadratic terms, we can often replace it by plain Hölder inequalities.

We conclude this introduction by giving a short overview of the paper.

In the next section we fix our assumptions and state our main results, both for the quasi-homogeneous and the fully non-homogeneous regimes. Section 3 is devoted to the derivation of uniform bounds for the sequence of weak solutions, which enable us to infer first convergence properties and to identify weak-limit points. There, we will also derive constraints those limit points have to satisfy, and establish strong convergence of the density functions. In Section 4 we complete the proof of the convergence; the main part of the analysis will be devoted to passing to the limit in the convective term. Section 5 focuses on the well-posedness of the limit system obtained in the quasi-homogeneous regime. An appendix about Littlewood-Paley theory and paradifferential calculus will end the manuscript.

Notation and conventions

Before starting, let us introduce some useful notation we use throughout this text.

The space domain will be denoted by $\Omega \subset \mathbb{R}^d$: throughout the text, we will always work in the case $d = 2$. All derivatives are (weak) derivatives, and the symbols ∇ , div and Δ are, unless specified otherwise, relative to the space variables. Given a subset $U \subset \Omega$ or $U \subset \mathbb{R}_+ \times \Omega$, we note $\mathcal{D}(U)$ the space of compactly supported C^∞ functions on U . If f is a tempered distribution, we note $\mathcal{F}[f] = \widehat{f}$ the Fourier transform of f with respect to the space variables.

For $1 \leq p \leq +\infty$, we will note $L^p(\Omega) = L^p$ when there is no ambiguity regarding the domain of definition of the functions. Likewise, we omit the dependency on Ω in functional spaces when no mistake can be made. If X is a Fréchet space of functions, we note $L^p(X) = L^p(\mathbb{R}_+; X)$. For any finite $T > 0$, we note $L_T^p(X) = L^p([0, T]; X)$ and $L_T^p = L^p[0, T]$.

Let $(f_\epsilon)_{\epsilon>0}$ be a sequence of functions in a normed space X . If this sequence is bounded in X , we use the notation $(f_\epsilon)_{\epsilon>0} \subset X$. If X is a topological linear space, whose (topological) dual is X' , we note $\langle \cdot | \cdot \rangle_{X' \times X}$ the duality brackets.

Any constant will be generically noted C , and, whenever deemed useful, we will specify the dependencies by writing $C = C(a_1, a_2, a_3, \dots)$. In all the text, $M_p(t) \in L^p(\mathbb{R}_+)$ will be a generic globally L^p function; on the other hand, we will use the notation $N_p(t)$ to denote a generic function in $L_{\text{loc}}^p(\mathbb{R}_+)$.

Acknowledgements

The work of the second author has been partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissement d’Avenir” (ANR-11-IDEX-0007), and by the projects BORDS (ANR-16-CE40-0027-01) and SingFlows (ANR-18-CE40-0027), all operated by the French National Research Agency (ANR).

The second author is deeply grateful to M. Paicu for interesting discussions on the MHD system.

2 Main assumptions and results

Let us fix the initial domain Ω to be either the whole space \mathbb{R}^2 or the torus \mathbb{T}^2 . In $\mathbb{R}_+ \times \Omega$, we consider the following *non-homogeneous incompressible MHD system*:

$$(3) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \frac{1}{\epsilon} \nabla \pi + \frac{1}{\epsilon} \rho u^\perp = \operatorname{div}(\nu(\rho) \nabla u) + \operatorname{div}(b \otimes b) - \frac{1}{2} \nabla |b|^2 \\ \partial_t b + \operatorname{div}(u \otimes b) - \operatorname{div}(b \otimes u) = \nabla^\perp(\mu(\rho) \operatorname{curl}(b)) \\ \operatorname{div} u = \operatorname{div} b = 0. \end{cases}$$

In the previous system, the viscosity coefficient ν and the resistivity coefficient μ are assumed to be continuous and non-degenerate: more precisely, they satisfy

$$\nu, \mu \in C^0(\mathbb{R}_+), \quad \text{with, } \forall \rho \geq 0, \quad \nu(\rho) \geq \nu_* > 0 \quad \text{and} \quad \mu(\rho) \geq \mu_* > 0,$$

for some positive real numbers ν_* and μ_* .

Our main goal here is to perform the limit for $\epsilon \rightarrow 0^+$ in equations (3) for general *ill-prepared* initial data. Let us then specify the assumptions on the initial density, velocity field and magnetic field.

2.1 Initial data

We supplement system (3) with general ill-prepared initial data. Let us be more precise, and start by considering the density function: for any $0 < \epsilon \leq 1$, we take

$$\rho_{0,\epsilon} = \rho_0 + \epsilon r_{0,\epsilon}, \quad \text{with} \quad \rho_0 \in C_b^2(\Omega) \quad \text{and} \quad (r_{0,\epsilon})_{\epsilon>0} \subset (L^2 \cap L^\infty)(\Omega).$$

Here above, we have denoted $C_b^2 := C^2 \cap W^{2,\infty}$. We also assume that there is a constant $\rho^* > 0$ such that, for any $\epsilon > 0$, one has

$$0 \leq \rho_0 \leq \rho^* \quad \text{and} \quad 0 \leq \rho_{0,\epsilon} \leq 2\rho^*.$$

In the case $\Omega = \mathbb{R}^2$, we require the initial densities $\rho_{0,\epsilon}$ to fulfill an extra integrability assumption. Namely, we suppose that one of the two following (non-equivalent) conditions is satisfied: either

$$(4) \quad \exists \delta > 0 \quad | \quad \left(\frac{1}{\rho_{0,\epsilon}} \mathbb{1}_{\{\rho_{0,\epsilon} < \delta\}} \right)_{\epsilon > 0} \subset L^1(\Omega),$$

where the symbol $\mathbb{1}_A$ stands for the characteristic function of a set $A \subset \Omega$, or

$$(5) \quad \exists p_0 \in]1, +\infty[, \quad \exists \bar{\rho} > 0 \quad | \quad \left((\bar{\rho} - \rho_{0,\epsilon})^+ \right)_{\epsilon > 0} \subset L^{p_0}(\Omega).$$

These two conditions allow for a low frequency control on the fluid velocity, uniformly with respect to ϵ . We will comment more about them at the end of the next subsection, and refer to Chapter 2 of [25] (see conditions (2.8), (2.9) and (2.10) therein) for more details.

For the velocity field, in order to avoid the trouble of defining the speed of the fluid in a vacuum zone $\{\rho = 0\}$, we work instead on the momentum $m = \rho u$. For any $\epsilon > 0$, we take an initial momentum $m_{0,\epsilon}$ such that

$$(m_{0,\epsilon})_{\epsilon > 0} \subset L^2(\Omega), \quad \left(\frac{|m_{0,\epsilon}|^2}{\rho_{0,\epsilon}} \right)_{\epsilon > 0} \subset L^1(\Omega),$$

where we agree that $m_{0,\epsilon} = 0$ and $|m_{0,\epsilon}|^2 / \rho_{0,\epsilon} = 0$ wherever $\rho_{0,\epsilon} = 0$.

Finally, let us now consider the magnetic field: we choose initial data

$$(b_{0,\epsilon})_{\epsilon > 0} \subset L^2(\Omega), \quad \text{with, } \forall \epsilon > 0, \quad \operatorname{div} b_{0,\epsilon} = 0.$$

Because of the previous uniform bounds, we deduce that, up to an extraction, one has the weak convergence properties

$$(6) \quad m_{0,\epsilon} \rightharpoonup m_0 \quad \text{in } L^2(\Omega), \quad r_{0,\epsilon} \xrightarrow{*} r_0 \quad \text{in } (L^2 \cap L^\infty)(\Omega), \quad b_{0,\epsilon} \rightharpoonup b_0 \quad \text{in } L^2(\Omega),$$

for suitable functions m_0 , r_0 and b_0 belonging to the respective functional spaces. Notice that we obviously have the strong convergence property $\rho_{0,\epsilon} - \rho_0 \rightarrow 0$ in $L^2 \cap L^\infty$ for $\epsilon \rightarrow 0^+$.

2.2 Finite energy weak solutions

For smooth solutions of (3) related to the initial data (ρ_0, m_0, b_0) , we can multiply the momentum equation by u and the magnetic field equation by b and integrate both equations. We have, after integration by parts,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |u|^2 dx + \int_{\Omega} \nu(\rho) |\nabla u|^2 dx &= \int_{\Omega} (b \cdot \nabla) b \cdot u dx \\ \frac{1}{2} \frac{d}{dt} \int_{\Omega} |b|^2 dx + \int_{\Omega} \mu(\rho) |\operatorname{curl}(b)|^2 dx &= \int_{\Omega} (b \cdot \nabla) u \cdot b dx. \end{aligned}$$

One more integration by parts show that the right-hand side of both equations are opposite. So, by summing the equations and integrating over $t \in [0, T]$ (for any fixed $T > 0$) and using the non-degeneracy hypothesis on the viscosity and resistivity coefficients, we get

$$(7) \quad \int_{\Omega} \left(\rho(T) |u(T)|^2 + |b(T)|^2 \right) dx + \int_0^T \int_{\Omega} \left(\nu_* |\nabla u|^2 + c\mu_* |\nabla b|^2 \right) dx dt \leq \int_{\Omega} \left(\frac{|m_0|^2}{\rho_0} + |b_0|^2 \right) dx.$$

In the above inequality, we have used the fact that b is divergence free, which implies that for almost all times $t \geq 0$ the norms $\|\operatorname{curl} b(t)\|_{L^2}$ and $\|\nabla b(t)\|_{L^2}$ are equivalent².

On the other hand, ρ is simply transported by the divergence-free velocity field u . Hence, for all $t \geq 0$, one formally has

$$(8) \quad \forall p \in [1, +\infty), \quad \|\rho(t)\|_{L^p} = \|\rho_0\|_{L^p}.$$

The previous inequalities give us grounds to define the notion of finite energy weak solution.

Definition 2.1. Let $T > 0$ and let (ρ_0, m_0, b_0) be initial data fulfilling the assumptions described in Section 2.1 above. We say that (ρ, u, b) is a *finite energy weak solution* to system (3) in $[0, T] \times \Omega$ related to the previous initial data if the following conditions are verified:

- (i) $\rho \in L^\infty([0, T] \times \Omega)$ and $\rho \in C^0([0, T]; L^q_{\text{loc}}(\Omega))$ for all $1 \leq q < +\infty$;
- (ii) $\rho|u|^2 \in L^\infty([0, T]; L^1(\Omega))$, with $u \in L^2([0, T]; H^1(\Omega)) \cap C_w^0([0, T]; L^2(\Omega))$, where the index w refers to continuity with respect to the weak topology;
- (iii) $b \in L^\infty([0, T]; L^2(\Omega)) \cap C_w^0([0, T]; L^2(\Omega))$, with $\nabla b \in L^2([0, T]; L^2(\Omega))$;
- (iv) the mass equation is satisfied in the weak sense: for any $\psi \in \mathcal{D}([0, T[\times \Omega])$, one has

$$\int_0^T \int_\Omega \left\{ \rho \partial_t \psi + \rho u \cdot \nabla \psi \right\} dx dt = - \int_\Omega \rho_0 \psi|_{t=0} dx;$$

- (v) the divergence-free conditions $\operatorname{div}(u) = \operatorname{div}(b) = 0$ are satisfied in $\mathcal{D}'([0, T[\times \Omega])$;
- (vi) the momentum equation is satisfied in the weak sense: for any $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2])$ such that $\operatorname{div}(\phi) = 0$, one has

$$\int_0^T \int_\Omega \left\{ \rho u \cdot \partial_t \phi + (\rho u \otimes u - b \otimes b) : \nabla \phi - \frac{1}{\epsilon} \rho u^\perp \cdot \phi - \nu(\rho) \nabla u : \nabla \phi \right\} dx dt = - \int_\Omega m_0 \phi|_{t=0} dx;$$

- (vii) the equation for the magnetic field is satisfied in the weak sense: for all $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2])$, one has

$$\int_0^T \int_\Omega \left\{ b \cdot \partial_t \phi + (u \otimes b - b \otimes u) : \nabla \phi - \mu(\rho) \operatorname{curl}(b) \operatorname{curl}(\phi) \right\} dx dt = - \int_\Omega b_0 \cdot \phi|_{t=0} dx;$$

- (viii) for almost every $t \in [0, T]$, the energy inequality (7) is satisfied.

The solution (ρ, u, b) is said to be *global* if the above conditions hold for all $T > 0$.

Existence of such finite energy weak solutions has been shown for fluids with density dependent viscosities in the case where there is no magnetic field (namely $b \equiv 0$ in system (3) above) by P.-L. Lions. His result even allows the initial density to vanish, under conditions (4) and (5). We refer to Chapter 2 of [25] for more details and references.

Concerning conductive fluids, more limited results are available. Gerbeau and Le Bris prove in [22] existence of finite energy weak solutions in a bounded domain of \mathbb{R}^3 (their proof can be extended to \mathbb{R}^2 or \mathbb{T}^2 with standard modifications), but only for fluids with non-vanishing initial densities. Desjardins and Le Bris do so in [11] for cylindrical or toroidal domains based on bounded subsets of \mathbb{R}^2 , and for flows with translation invariance.

Even if we do not have a full existence result for flows presenting vacuum patches, as described above, our arguments are robust enough to consider possible (mild) vanishing of the density, in the same spirit of (4) and (5). Therefore, we will work under those conditions, in order to accommodate possible future existence results.

²In fact, in dimension 2 and for the L^2 norm, we have the exact equality: namely, $\|\operatorname{curl} b\|_{L^2} = \|\nabla b\|_{L^2}$. Indeed, since $\xi \cdot \widehat{b}(\xi) = 0$, by Cauchy-Schwarz one has that $|\xi^\perp \cdot \widehat{b}(\xi)| = |\xi| |\widehat{b}(\xi)|$.

2.3 Statement of the results

We consider a sequence of initial data $(\rho_{0,\epsilon}, m_{0,\epsilon}, b_{0,\epsilon})_\epsilon$ satisfying all the assumptions and uniform bounds described in Section 2.1 above. We further consider a sequence $(\rho_\epsilon, u_\epsilon, b_\epsilon)_\epsilon$ of finite energy weak solutions (in the sense of Definition 2.1) related to those initial data. We aim at proving some kind of convergence of the solutions $(\rho_\epsilon, u_\epsilon, b_\epsilon)_\epsilon$ and identify the limit dynamics for $\epsilon \rightarrow 0^+$ in the form of a PDE solved by the limit points of the sequence. We consider two cases.

Firstly, we consider the case of a *quasi-homogeneous* density, meaning that the initial density $\rho_{0,\epsilon}$ is supposed to be a perturbation of a constant density state, say 1 for simplicity. We then write $\rho_{0,\epsilon} = 1 + \epsilon r_{0,\epsilon}$: this assumption simplifies the equations very much. Indeed, on the one hand, at any later time we still have $\rho_\epsilon = 1 + \epsilon r_\epsilon$, with r_ϵ solving a linear transport equation

$$(9) \quad \begin{cases} \partial_t r_\epsilon + \operatorname{div}(r_\epsilon u_\epsilon) = 0 \\ (r_\epsilon)|_{t=0} = r_{0,\epsilon}, \end{cases}$$

thanks to the divergence-free condition $\operatorname{div}(u_\epsilon) = 0$. On the other hand, the momentum equation can be written (in a suitable sense)

$$\begin{aligned} & \partial_t(\rho_\epsilon u_\epsilon) + \operatorname{div}(\rho_\epsilon u_\epsilon \otimes u_\epsilon - b_\epsilon \otimes b_\epsilon) + \frac{1}{\epsilon} \nabla \pi_\epsilon + \frac{1}{2} \nabla(|b_\epsilon|^2) + \frac{1}{\epsilon} \rho_\epsilon u_\epsilon^\perp - \operatorname{div}(\nu(\rho_\epsilon) \nabla u_\epsilon) \\ &= \partial_t u_\epsilon + \operatorname{div}(u_\epsilon \otimes u_\epsilon - b_\epsilon \otimes b_\epsilon) - \nu(1) \Delta u_\epsilon + r_\epsilon u_\epsilon^\perp + \left\{ \frac{1}{\epsilon} \nabla \pi_\epsilon + \frac{1}{2} \nabla(|b_\epsilon|^2) + \frac{1}{\epsilon} u_\epsilon^\perp \right\} + O(\epsilon), \end{aligned}$$

where the terms in the brackets (which are singular in ϵ) are gradient terms, hence do not appear in the weak form of the equations. Therefore, taking the limit in this case will not be too complicated. In the end, we can prove the following result.

Theorem 2.2. *Suppose that $\rho_0 = 1$ and consider a sequence $(\rho_{0,\epsilon}, m_{0,\epsilon}, b_{0,\epsilon})_{\epsilon>0}$ of initial data satisfying the assumptions fixed in Section 2.1. Let $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$ be a sequence of corresponding finite energy weak solutions to (3). Finally, let m_0, b_0 and r_0 be as in (6) and define $r_\epsilon = (\rho_\epsilon - 1)/\epsilon$.*

Then, there exists a triplet (r, u, b) in the space $L^\infty(\mathbb{R}_+; L^2(\Omega) \cap L^\infty(\Omega)) \times L^\infty(\mathbb{R}_+; L^2(\Omega)) \times L^\infty(\mathbb{R}_+; L^2(\Omega))$, with ∇u and ∇b belonging to $L^2(\mathbb{R}_+; L^2(\Omega))$ and $\operatorname{div} u = \operatorname{div} b = 0$, such that, up to the extraction of a subsequence, the following convergence properties hold: for any $T > 0$, we have

- (1) $r_\epsilon \xrightarrow{*} r$ in $L_T^\infty(L^2 \cap L^\infty)$, and $r_\epsilon \rightarrow r$ in $L_T^2(L_{\text{loc}}^2)$;
- (2) $u_\epsilon \xrightarrow{*} u$ in $L^\infty(L^2) \cap L_T^2(H^1)$;
- (3) $b_\epsilon \xrightarrow{*} b$ in $L^\infty(L^2) \cap L_T^2(H^1)$, and $b_\epsilon \rightarrow b$ in $L_T^2(H_{\text{loc}}^s)$ for any $s < 1$.

The limit dynamics is described by a homogeneous MHD-type system, which the triplet (r, u, b) solves in the weak sense: namely,

$$(10) \quad \begin{cases} \partial_t r + \operatorname{div}(r u) = 0 \\ \partial_t u + \operatorname{div}(u \otimes u) + \nabla \pi + \frac{1}{2} \nabla(|b|^2) + r u^\perp = \nu(1) \Delta u + \operatorname{div}(b \otimes b) \\ \partial_t b + \operatorname{div}(u \otimes b - b \otimes u) = \mu(1) \Delta b \\ \operatorname{div}(u) = \operatorname{div}(b) = 0, \end{cases}$$

for some pressure function π and with initial data (r_0, m_0, b_0) .

In addition, if $(r_0, u_0, b_0) \in H^{1+\beta} \times H^1 \times H^1$, for some $\beta \in]0, 1[$, then the solution (r, u, b) to system (10) is unique. As a consequence, the whole sequence $(r_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$ converges.

We will study the limit system (10) in Section 5; there, we will also specify better in which functional class the uniqueness of solutions holds.

The second case we consider is the case of a *fully non-homogeneous* density, in the sense that the reference density profile ρ_0 is non-constant. Moreover, we need an extra technical assumption on ρ_0 : we suppose that

$$(11) \quad \forall K \subset \Omega, \quad K \text{ compact}, \quad \text{meas} \left\{ x \in K \mid |\nabla \rho_0(x)| \leq \delta \right\} \xrightarrow{\delta \rightarrow 0^+} 0.$$

Roughly speaking, we are requiring that, for any fixed compact $K \subset \Omega$, the set of critical points of ρ_0 in K is of zero measure. We remark that this condition is a relaxed version of the condition imposed in [16] (see also [21], [14]), which involves no difficulties in the proof, and has the advantage of allowing for more general reference densities (for instance, profiles which exponentially decay to some positive constant at $|x| \sim +\infty$).

This case is understandably more difficult, since none of the two previous simplifications can be made. However, we will see that, by using the structure of system (3), we can find an analogous decomposition $\rho_\epsilon = \rho_0 + \epsilon \sigma_\epsilon$, where $(\sigma_\epsilon)_\epsilon$ is bounded in a low regularity space. Unfortunately, this does not simplify much the singular term, as $\frac{1}{\epsilon} \rho_0 u_\epsilon^\perp$ is not a gradient term. Another problem is that the bounds we will find on $(\sigma_\epsilon)_\epsilon$ are in such a low regularity space ($H^{-3-\delta}$ in fact) that taking the limit $\epsilon \rightarrow 0^+$ directly in the momentum equation is impossible. We will need to, instead, work on the vorticity and take the curl of the momentum equation.

The result in the fully non-homogeneous case is contained in the next statement.

Theorem 2.3. *Assume that ρ_0 satisfies condition (11) and (when $\Omega = \mathbb{R}^2$) either (4) or (5). Consider a sequence $(\rho_{0,\epsilon}, m_{0,\epsilon}, b_{0,\epsilon})_{\epsilon>0}$ of initial data satisfying the assumptions fixed in Section 2.1, and let $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$ be a sequence of corresponding weak solutions to (3). Finally, let m_0, b_0 and r_0 be as in (6) and define $\sigma_\epsilon := (\rho_\epsilon - 1)/\epsilon$.*

Then, there exist $\sigma \in L_{\text{loc}}^\infty(\mathbb{R}_+; H^{-3-\delta}(\Omega))$ for any $\delta > 0$ arbitrarily small, $u \in L^\infty(\mathbb{R}_+; L^2(\Omega))$ and $b \in L^\infty(\mathbb{R}_+; L^2(\Omega))$, with ∇u and ∇b in $L^2(\mathbb{R}_+; L^2(\Omega))$ and $\text{div}(\rho_0 u) = \text{div} u = \text{div} b = 0$, such that, up to the extraction of a subsequence, the following convergence properties hold true: for any $T > 0$, one has

- (1) $\rho_\epsilon \rightarrow \rho_0$ in $L_T^2(L_{\text{loc}}^2)$;
- (2) $\sigma_\epsilon \xrightarrow{*} \sigma$ in $L_T^\infty(H^{-3-\delta})$, for all arbitrarily small $\delta > 0$;
- (3) $u_\epsilon \xrightarrow{*} u$ in $L^\infty(L^2) \cap L_T^2(H^1)$;
- (4) $b_\epsilon \xrightarrow{*} b$ in $L^\infty(L^2) \cap L_T^2(H^1)$, and $b_\epsilon \rightarrow b$ in $L_T^2(H_{\text{loc}}^s)$ for any $s < 1$.

Moreover, there exists a distribution $\Gamma \in \mathcal{D}'(\mathbb{R}_+ \times \Omega)$ of order at most 1 such that

$$\begin{cases} \partial_t \left(\text{curl}(\rho_0 u) - \sigma \right) - \text{curl} \left(\text{div}(\nu(\rho_0) \nabla u) \right) + \text{curl} \left(\rho_0 \nabla \Gamma - \text{div}(b \otimes b) \right) = 0 \\ \partial_t b + \text{div}(u \otimes b - b \otimes u) = \nabla^\perp(\mu(\rho_0) \text{curl}(b)) \\ \text{div}(\rho_0 u) = 0 \\ \text{div}(u) = \text{div}(b) = 0, \end{cases}$$

with initial data $\left(\text{curl}(\rho_0 u) - \sigma \right)_{|t=0} = \text{curl}(m_0) - r_0$ and $b_{|t=0} = b_0$.

Remark 2.4. The summand $\rho_0 \nabla \Gamma$ can be interpreted as a Lagrange multiplier associated to the constraint $\text{div}(\rho_0 u) = 0$, just as the pressure term $\nabla \pi$ in (3) and (10) can be seen as a Lagrange multiplier for the incompressibility constraint $\text{div}(u) = 0$.

3 Uniform bounds and convergence properties

The next three sections are devoted to the proofs of Theorems 2.2 and 2.3. In all that follows, $(\rho_{0,\epsilon}, m_{0,\epsilon}, b_{0,\epsilon})_\epsilon$ is a sequence of initial data satisfying all the assumptions and uniform bounds described in Section 2.1 above, and $(\rho_\epsilon, u_\epsilon, b_\epsilon)_\epsilon$ is an associated sequence of finite energy weak solutions related to those initial data, in the sense of Definition 2.1 above.

In this section, we first use uniform bounds on the solutions to prove their weak convergence. Then, we focus on convergence results for the density, which we will need later.

3.1 Uniform bounds

In this section, we establish uniform bounds (i.e. bounds independent of ϵ) on the sequence of solutions $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$, thus enabling us to extract weakly converging subsequences.

First of all, we notice that the solutions satisfy the energy inequality (7) (this is point (viii) of Definition 2.1): for almost every $t > 0$ fixed, we have

$$\int_{\Omega} \left(\rho_\epsilon(t) |u_\epsilon(t)|^2 + |b_\epsilon(t)|^2 \right) dx + \int_0^t \int_{\Omega} \left(\nu_* |\nabla u_\epsilon|^2 + \mu |\nabla b_\epsilon|^2 \right) dx ds \leq \int_{\Omega} \left(\frac{|m_{0,\epsilon}|^2}{\rho_{0,\epsilon}} + |b_{0,\epsilon}|^2 \right) dx.$$

In view of our assumptions on the initial data, the right-hand side of the previous inequality is uniformly bounded. Thus we get

$$(12) \quad (\sqrt{\rho_\epsilon} u_\epsilon)_{\epsilon>0}, (b_\epsilon)_{\epsilon>0} \subset L^\infty(\mathbb{R}_+; L^2(\Omega)),$$

$$(13) \quad (\nabla u_\epsilon)_{\epsilon>0}, (\nabla b_\epsilon)_{\epsilon>0} \subset L^2(\mathbb{R}_+; L^2(\Omega)).$$

Secondly, because both ρ_ϵ and (in the quasi-homogeneous case) $r_\epsilon := (\rho_\epsilon - 1)/\epsilon$ solve a pure transport equation by the divergence-free vector field u_ϵ (keep in mind (9) above), we see that, for all $\epsilon > 0$, one has

$$\forall 0 \leq \alpha \leq \beta < +\infty, \quad \text{meas}\{\alpha \leq \rho_\epsilon \leq \beta\} = \text{meas}\{\alpha \leq \rho_{0,\epsilon} \leq \beta\},$$

and the same holds for r_ϵ (this is the same property as in Theorem 2.1, Chapter 2, p. 23 of [25]). On the one hand, this of course implies that $(\rho_\epsilon)_{\epsilon>0} \subset L^\infty(L^\infty)$ and $(r_\epsilon)_{\epsilon>0} \subset L^\infty(L^2 \cap L^\infty)$, together with the bounds

$$(14) \quad 0 \leq \rho_\epsilon \leq 2\rho^* \quad \text{almost everywhere in } \mathbb{R}_+ \times \Omega,$$

in view of the assumptions on the initial datum $\rho_{0,\epsilon}$. Therefore, up to extracting a subsequence, we gather the convergences

$$(15) \quad \rho_\epsilon \xrightarrow{*} \rho \quad \text{in } L^\infty(L^\infty) \quad \text{and} \quad r_\epsilon \xrightarrow{*} r \quad \text{in } L^\infty(L^2 \cap L^\infty),$$

for some $\rho \in L^\infty(L^\infty)$ and $r \in L^\infty(L^2 \cap L^\infty)$. On the other hand, the same property also shows that, for almost every $t > 0$, the density $\rho_\epsilon(t)$ satisfies the extra regularity properties which we had required if $\Omega = \mathbb{R}^2$, and it does so independently of $t > 0$ and uniformly with respect to $\epsilon > 0$:

$$(16) \quad \left(\frac{1}{\rho_\epsilon(t)} \mathbf{1}_{\{\rho_\epsilon(t) < \delta\}} \right)_{\epsilon>0} \subset L^1(\Omega) \quad \text{or} \quad \left((\bar{\rho} - \rho_\epsilon(t))^+ \right)_{\epsilon>0} \subset L^{p_0}(\Omega),$$

where $\delta > 0$, $p_0 \in]1, +\infty[$ and $\bar{\rho} > 0$ are the same as in conditions (4) and (5).

Finally, we also see that $(u_\epsilon)_{\epsilon>0}$ is in fact uniformly bounded in $L_T^2(L^2)$ for any finite time $T > 0$. Indeed, if $\Omega = \mathbb{R}^2$, this is a consequence of either one of the two previous conditions in (16) (see also [25], point 8 in Remark 2.1 pp. 24-25). If $\Omega = \mathbb{T}^2$ instead, the same can be shown

without the extra assumptions, by means of the Poincaré-Wirtinger inequality (see again [25], Subsection 2.3 p. 37). Therefore, up to an extraction, we deduce that

$$(17) \quad u_\epsilon \rightharpoonup u \quad \text{in } L^2_{\text{loc}}(\mathbb{R}_+; H^1(\Omega)) \quad \text{and} \quad b_\epsilon \rightharpoonup b \quad \text{in } L^2_{\text{loc}}(\mathbb{R}_+; H^1(\Omega)),$$

for suitable functions u and b belonging to $L^2_{\text{loc}}(\mathbb{R}_+; H^1(\Omega))$.

Remark that, in fact, we have a more precise convergence property for the magnetic fields: in view of (12)-(13), we know that $b \in L^\infty(\mathbb{R}_+; L^2(\Omega))$, with $\nabla b \in L^2(\mathbb{R}_+; L^2(\Omega))$ and, up to an extraction, we have the convergences $b_\epsilon \xrightarrow{*} b$ in $L^\infty(L^2)$ and $\nabla b_\epsilon \rightharpoonup \nabla b$ in $L^2(L^2)$. We will resort to those precise features when needed.

3.2 Strong convergence of the densities

This section is dedicated to the quest of pointwise convergence for the ρ_ϵ . This will be useful for two reasons. Firstly, strong convergence makes the proofs simpler: on many occasions, the use of paradifferential calculus in [16] can be replaced by more elementary arguments. Secondly, pointwise convergence is necessary to deal with the viscosity and resistivity terms: since we only have weak convergence of the velocity fields, strong convergence of both $\nu(\rho_\epsilon)$ and $\mu(\rho_\epsilon)$ is required to achieve convergence of the product terms $\text{div}(\nu(\rho_\epsilon)\nabla u_\epsilon)$ and $\nabla^\perp(\mu(\rho_\epsilon)\text{curl } u_\epsilon)$.

However, the uniform bounds alone are insufficient to prove the strong convergence we seek. So far, we have only obtained mere weak convergence $\rho_\epsilon \xrightarrow{*} \rho$ in $L^\infty(L^\infty)$ (recall uniform bound (15) above). We now resort to the arguments of Di Perna and P.-L. Lions [12]: if somehow we proved that

$$(18) \quad \rho_\epsilon^2 \xrightarrow{*} \rho^2 \quad \text{in } L^\infty(\mathbb{R}_+; L^\infty(\Omega)),$$

then, by using the characteristic function $\mathbb{1}_K$ of a compact subset $K \subset \Omega$ as a test function, we would recover convergence of the L^2 norms. Using the euclidean structure of $L^2_T(L^2(K))$, we would then deduce local strong convergence, hence pointwise convergence, after extraction.

Therefore, the argument boils down to proving (18). The quadratic non-linearity is the main challenge as, by the uniform bounds $(\rho_\epsilon)_{\epsilon>0} \subset L^\infty(L^\infty)$, we only know that there exists some function $g \in L^\infty(L^\infty)$ such that $\rho_\epsilon^2 \rightharpoonup^* g$ in $L^\infty(L^\infty)$, and this function g need not be ρ^2 . The trick is that both g and ρ^2 are (weak) solutions of the transport PDE

$$(19) \quad \begin{cases} \partial_t a + u \cdot \nabla a = 0 \\ a|_{t=0} = a_0, \end{cases}$$

with same initial datum and divergence-free velocity field. After the work [12], problem (19) is well-posed, so g and ρ^2 must be equal. To sum up, we get the following statement.

Proposition 3.1. *The convergence property (18) holds true. In particular, in the limit $\epsilon \rightarrow 0^+$, we have the strong convergence*

$$\rho_\epsilon \longrightarrow \rho \quad \text{in } L^2_{\text{loc}}(\mathbb{R}_+ \times \Omega),$$

and, up to the extraction of a suitable subsequence, the convergence holds also almost everywhere in $\mathbb{R}_+ \times \Omega$.

Before proving the previous proposition, some preliminary lemmas are in order. First of all, we establish that all the ρ_ϵ^2 are solutions to the continuity equation. Since this fact is to be shown for all $\epsilon > 0$ independently, we drop the ϵ indices for more clarity.

Lemma 3.2. *Let $\rho_0 \in L^\infty$ and $u \in L^2_{\text{loc}}(\mathbb{R}_+; H^1)$ be a divergence-free vector field. Let $\rho \in L^\infty(L^\infty)$ be a weak solution to the Cauchy problem (19), with initial datum ρ_0 .*

Then ρ^2 is also a weak solution of the same equation, related to the initial datum ρ_0^2 .

Proof of Lemma 3.2. We wish to prove that ρ^2 is a weak solution of (19), with initial datum ρ_0^2 : this means that, for all $T > 0$ and all $\psi \in \mathcal{D}([0, T[\times \Omega)$, one has

$$\int_0^T \int_{\Omega} \rho^2 (\partial_t \psi + u \cdot \nabla \psi) dx dt + \int_{\Omega} \rho_0^2 \psi|_{t=0} dx = 0.$$

We consider a smoothing kernel $(\mu_{\alpha})_{\alpha>0}$ such that, if $\Omega = \mathbb{R}^2$, we have $\mu_{\alpha}(x) = \mu(x/\alpha)/\alpha^d$, where $\mu \in C^\infty(\mathbb{R}^2)$ is such that $\text{Supp } \mu \subset B(0, 1)$ and $\mu(x) = \mu(-x)$. For $\Omega = \mathbb{T}^2$, instead, we set $\mu'_{\alpha}(x) = \sum_{k \in \mathbb{Z}^2} \mu_{\alpha}(x + k)$.

For all $\alpha > 0$, we define $\rho_{\alpha} = \mu_{\alpha} * \rho$ (use μ'_{α} instead of μ_{α} if $\Omega = \mathbb{T}^2$). Then, using also the divergence-free condition for u and the evenness of μ_{α} , we deduce that ρ_{α} solves (in the weak sense) the following approximate equation:

$$\partial_t \rho_{\alpha} + u \cdot \nabla \rho_{\alpha} = [u \cdot \nabla, \mu_{\alpha} *] \rho, \quad (\rho_{\alpha})|_{t=0} = \mu_{\alpha} * \rho_0,$$

where we have denoted $[u \cdot \nabla, \mu_{\alpha} *]$ the commutator between $u \cdot \nabla$ and the convolution by μ_{α} . Multiplying this equation by $2\rho_{\alpha}$ shows that

$$(20) \quad \partial_t (\rho_{\alpha}^2) + u \cdot \nabla (\rho_{\alpha}^2) = 2\rho_{\alpha} [u \cdot \nabla, \mu_{\alpha} *] \rho, \quad (\rho_{\alpha}^2)|_{t=0} = (\mu_{\alpha} * \rho_0)^2.$$

The space differentiation $2\rho_{\alpha} \nabla \rho_{\alpha} = \nabla (\rho_{\alpha}^2)$ is justified because $\rho_{\alpha}(t) \in C^\infty$ for almost all times $0 \leq t \leq T$ (for any fixed $T > 0$), and the time differentiation $2\rho_{\alpha} \partial_t \rho_{\alpha} = \partial_t (\rho_{\alpha}^2)$ is justified because $\rho_{\alpha} \in W_T^{1,2}(H_{\text{loc}}^s)$ for every $\alpha > 0$ and $s \geq 0$. This comes from the property $\rho u \in L_T^2(L^2)$ and the relation $\partial_t \rho_{\alpha} = -\text{div}(\mu_{\alpha} * (\rho u))$ (which follows from (19) and $\text{div } u = 0$), which implies $\partial_t \rho_{\alpha} \in L_T^2(H_{\text{loc}}^s)$.

Our next goal is to take the limit $\alpha \rightarrow 0^+$ in the weak formulation of (20), namely in the relation

$$(21) \quad \int_0^T \int_{\Omega} \rho_{\alpha}^2 \left\{ \partial_t \psi + u \cdot \nabla \psi \right\} dx dt + \int_{\Omega} (\mu_{\alpha} * \rho_0)^2 \psi|_{t=0} dx + \int_0^T \int_{\Omega} 2\psi \rho_{\alpha} [u \cdot \nabla, \mu_{\alpha} *] \rho dx dt = 0,$$

for any arbitrary test function $\psi \in \mathcal{D}([0, T[\times \Omega)$. On the one hand, we remark that, since $u \in L_T^2(H^1)$ and $\rho \in L^\infty(L^\infty)$, we can apply Lemma II.1 in [12] to get

$$[u \cdot \nabla, \mu_{\alpha} *] \rho \xrightarrow{\alpha \rightarrow 0^+} 0 \quad \text{in } L_T^1(L_{\text{loc}}^2).$$

Using this, we find that the commutator term in (20) cancels in the limit $\alpha \rightarrow 0^+$, as, for every compact $K \subset \Omega$, we have

$$\|2\rho_{\alpha} [u \cdot \nabla, \mu_{\alpha} *] \rho\|_{L_T^1(L^2(K))} \leq 2\|\rho\|_{L_{t,x}^\infty} \| [u \cdot \nabla, \mu_{\alpha} *] \rho \|_{L_T^1(L^2(K))} \xrightarrow{\alpha \rightarrow 0^+} 0.$$

On the other hand, since $\rho \in L^\infty(\mathbb{R}_+ \times \Omega)$, by standard properties of mollification kernels we gather the strong convergence $\rho_{\alpha} \rightarrow \rho$ in e.g. $L_T^2(L_{\text{loc}}^2)$ when $\alpha \rightarrow 0^+$, for any fixed $T > 0$. By the same token, we also have $\mu_{\alpha} * \rho_0 \rightarrow \rho_0$ in $L_{\text{loc}}^2(\Omega)$, in the limit $\alpha \rightarrow 0^+$. Thanks to those properties, it is easy to take the limit in the first and second term in (21).

The proof of the lemma is now completed. \square

We also need the following result.

Lemma 3.3. *Let $g \in L^\infty(\mathbb{R}_+ \times \Omega)$ be any weak-* limit point of the sequence $(\rho_\epsilon^2)_{\epsilon>0}$ with respect to the $L^\infty(\mathbb{R}_+ \times \Omega)$ topology. Let ρ_0 be the limit density profile and u the limit velocity field identified in (17).*

Then g is a solution of the linear transport equation

$$(22) \quad \partial_t g + u \cdot \nabla g = 0, \quad \text{with} \quad g|_{t=0} = \rho_0^2.$$

Proof of Lemma 3.3. By Lemma 3.2 above, we know that the ρ_ϵ^2 solve the transport equation

$$(23) \quad \partial_t (\rho_\epsilon^2) + u_\epsilon \cdot \nabla (\rho_\epsilon^2) = 0,$$

related to the initial datum $(\rho_\epsilon^2)|_{t=0} = \rho_{0,\epsilon}^2$. Therefore, for proving our claim, it is enough to take the limit in the weak formulation of the previous equation. The main issue is showing that $\rho_\epsilon^2 u_\epsilon$ converges in (say) \mathcal{D}' to gu : for doing so, we resort to some arguments of [16] (see Paragraph 3.1.2. therein). The basic idea is to use the transport equation solved by the ρ_ϵ^2 to trade space regularity for time compactness.

We start by remarking that, in view of (23), one has $\partial_t(\rho_\epsilon^2) = -\operatorname{div}(\rho_\epsilon^2 u_\epsilon)$. Since $|\rho_\epsilon^2 u_\epsilon| \leq (\rho^*)^{3/2} |\sqrt{\rho_\epsilon} u_\epsilon|$ by use of (14) and (12), we infer that $(\partial_t(\rho_\epsilon^2))_{\epsilon>0} \subset L^\infty(H^{-1})$, hence

$$(\rho_\epsilon^2)_{\epsilon>0} \subset W_T^{1,\infty}(H_{\text{loc}}^{-1}),$$

where the localisation in space comes from the fact that the initial data ρ_0^2 is just $L^\infty(\Omega)$. Let now $\theta \in]0, 1[$: standard Sobolev interpolation gives, for almost all $0 \leq s, t \leq T$, the estimate

$$\|(\rho_\epsilon(t) - \rho_\epsilon(s))\chi\|_{H^{-\theta}} \leq \|(\rho_\epsilon(t) - \rho_\epsilon(s))\chi\|_{H^{-1}}^\theta \|(\rho_\epsilon(t) - \rho_\epsilon(s))\chi\|_{L^2}^{1-\theta},$$

where $\chi \in \mathcal{D}(\Omega)$ is an arbitrary compactly supported function. This shows that $(\rho_\epsilon^2)_{\epsilon>0}$ is bounded in every space $C_T^{0,\theta}(H_{\text{loc}}^{-\theta})$. Therefore, by the Ascoli-Arzelà theorem, we gather the strong convergence

$$\rho_\epsilon^2 \longrightarrow g \quad \text{in } C^{0,\theta}([0, T]; H_{\text{loc}}^{-\theta}(\Omega)),$$

for all $0 < \theta < 1$ and all fixed $T > 0$. Combining this property with the Lemma A.5, which provides continuity of the function product $(a, b) \mapsto ab$ in the $H^{-\theta} \times H^1 \rightarrow H^{-\theta-\delta}$ topology (for $\delta > 0$ arbitrarily small), we get

$$\rho_\epsilon^2 u_\epsilon \rightharpoonup gu \quad \text{in } \mathcal{D}'([0, T] \times \Omega).$$

It is now possible to take the limit $\epsilon \rightarrow 0^+$ in the weak form of equation (23), thus recovering equation (22). \square

We can now complete the proof to Proposition 3.1.

Proof of Proposition 3.1. In view of Lemmas 3.3 and 3.2, it follows that both ρ^2 and g are weak solutions to the initial value problem (22), and they both belong to $L^\infty(\mathbb{R}_+ \times \Omega)$. But problem (22) is in fact well-posed in the previous space, as a consequence of Di Perna and P.-L. Lions theory. More precisely, in order to apply their uniqueness result (see Theorem II.2 of [12]), we have to make sure that the limit velocity field $u \in L_{\text{loc}}^2(\mathbb{R}_+; H^1(\Omega))$ fulfills the following condition: for any fixed $T > 0$,

$$(24) \quad \frac{u(t, x)}{1 + |x|} \in L_T^1(L^1) + L_T^1(L^\infty).$$

To see this, let us set an arbitrary $R > 0$ and decompose according to whether $|u| < R$ or not:

$$\frac{|u(t, x)|}{1 + |x|} = \mathbf{1}_{\{|u| < R\}} \frac{|u(t, x)|}{1 + |x|} + \mathbf{1}_{\{|u| \geq R\}} \frac{|u(t, x)|}{1 + |x|}.$$

On the one hand, the measure of the set $A_R(t) := \{|u(t)| \geq R\}$ is bounded by the Bienaymé-Chebyshev inequality, as

$$\operatorname{meas} A_R(t) \leq \frac{1}{R^2} \int_\Omega |u(t, x)|^2 dx.$$

Observe that the term on the right-hand side of the previous estimate belongs to L_T^1 for all fixed $T > 0$. Therefore, Hölder's inequality yields

$$\int_0^T \int_{A_R} \frac{|u(t, x)|}{1 + |x|} dx dt \leq \int_0^T \|u(t)\|_{L^1(A_R)} dt \leq \int_0^T \|u(t)\|_{L^2} (\text{meas } A_R(t))^{1/2} dt < +\infty,$$

implying that $\mathbb{1}_{A_R(t)} u(t, x) (1 + |x|)^{-1} \in L_T^1(L^1)$. On the other hand, we obviously have

$$\mathbb{1}_{\{|u(t)| < R\}} \frac{|u(t, x)|}{1 + |x|} \leq R \in L_T^1(L^\infty).$$

Therefore, (24) is indeed satisfied by u , so we can apply Theorem II.2 of [12]. This result implies that we do have $\rho^2 \equiv g$ almost everywhere. In particular, we also deduce the weak convergence (18). As already remarked above, that property in turn yields local strong convergence of the ρ_ϵ to ρ . Indeed, let $K \subset \Omega$ be a compact set; then, using (18) we get

$$\langle \rho_\epsilon^2, \mathbb{1}_K \rangle_{L_{t,x}^\infty \times L_{t,x}^1} = \|\rho_\epsilon\|_{L_T^2 L^2(K)}^2 \longrightarrow \langle \rho^2, \mathbb{1}_K \rangle_{L_{t,x}^\infty \times L_{t,x}^1} = \|\rho\|_{L_T^2 L^2(K)}^2.$$

At this point, the fact that $L_T^2(L^2(K))$ has a Euclidean structure gives strong convergence: because of the weak-* convergence (15) of the ρ_ϵ , we infer that $\langle \rho_\epsilon, \rho \rangle_{L^\infty([0, T[\times K) \times L^1([0, T[\times K)}$ tends to $\|\rho\|_{L_T^2(L^2(K))}^2$ and hence

$$\|\rho_\epsilon - \rho\|_{L_T^2(L^2(K))}^2 = \|\rho_\epsilon\|_{L_T^2(L^2(K))}^2 + \|\rho\|_{L_T^2(L^2(K))}^2 - 2 \int_0^T \int_K \rho_\epsilon \rho dx dt \xrightarrow{\epsilon \rightarrow 0^+} 0.$$

In particular, after extracting one more time, we deduce the pointwise convergence $\rho_\epsilon \longrightarrow \rho$, in the limit $\epsilon \rightarrow 0^+$. \square

3.3 The singular part of the equations

In this part, we focus our attention on the singular part of system (3), namely on the term $\epsilon^{-1}(\nabla \pi_\epsilon + \rho_\epsilon u_\epsilon^\perp)$ in the momentum equation. Note that any singular gradient term disappears in its weak formulation, due to the divergence-free condition on the test functions.

Proposition 3.4. *Let $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon > 0}$ be a sequence of weak solutions to system (3), associated with the sequence of initial data $(\rho_{0,\epsilon}, u_{0,\epsilon}, b_{0,\epsilon})_{\epsilon > 0}$ satisfying the assumptions fixed in Subsection 2.1. Let (ρ, u, b) be a limit point of the sequence $(\rho_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon > 0}$, as identified in Subsection 3.1.*

1) *In the case of a quasi-homogeneous density, for all test function $\phi \in \mathcal{D}(\mathbb{R}_+ \times \Omega; \mathbb{R}^2)$ such that $\text{div } \phi = 0$, we have*

$$\frac{1}{\epsilon} \int_0^{+\infty} \int_\Omega \rho_\epsilon u_\epsilon^\perp \cdot \phi dx dt \xrightarrow{\epsilon \rightarrow 0^+} \int_0^{+\infty} \int_\Omega r u^\perp \cdot \phi dx dt.$$

2) *In the fully non-homogeneous case, the limit density satisfies $\rho(t, x) = \rho_0(x)$ for almost every $(t, x) \in \mathbb{R}_+ \times \Omega$. Moreover, we have the relations $\text{div}(\rho_0 u) = \text{div } u = 0$ almost everywhere in $\mathbb{R}_+ \times \Omega$. In particular, $\nabla \rho_0 \cdot u = 0$ almost everywhere in $\mathbb{R}_+ \times \Omega$.*

Proof. We start by attending to the quasi-homogeneous setting, where the singularity *de facto* disappears. Indeed, we can write

$$\frac{1}{\epsilon} (\nabla \pi_\epsilon + \rho_\epsilon u_\epsilon^\perp) = \frac{1}{\epsilon} (\nabla \pi_\epsilon + u_\epsilon^\perp) + r_\epsilon u_\epsilon,$$

and the terms in the brackets are perfect gradients. Therefore, if $T > 0$ and $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2])$ is a divergence-free test function, one gets

$$\frac{1}{\epsilon} \int_0^T \int_{\Omega} \rho_{\epsilon} u_{\epsilon}^{\perp} \cdot \phi \, dx \, dt = \int_0^T \int_{\Omega} r_{\epsilon} u_{\epsilon}^{\perp} \cdot \phi \, dx \, dt.$$

To take the limit $\epsilon \rightarrow 0^+$ in this last integral, we observe that the functions r_{ϵ} solve the linear transport equation (9); hence, Proposition 3.1 applies to the sequence $(r_{\epsilon})_{\epsilon > 0}$, yielding the strong convergence

$$r_{\epsilon} \xrightarrow{\epsilon \rightarrow 0^+} r \quad \text{in} \quad L_T^2(L_{\text{loc}}^2),$$

for any fixed $T > 0$. Using the weak convergence (17) for $(u_{\epsilon})_{\epsilon}$, we finally infer that

$$r_{\epsilon} u_{\epsilon} \longrightarrow r u \quad \text{in} \quad \mathcal{D}'([0, T[\times \Omega).$$

This completes the proof of the first property of the proposition.

The study of the fully non-homogeneous case is very much similar, with the exception that the singularity does not disappear. We start by remarking that, exactly as done above, as a consequence of Proposition 3.1 we deduce that $\rho_{\epsilon} u_{\epsilon} \longrightarrow \rho u$ in the sense of distributions. Now, multiplying the momentum equation in its weak form by ϵ , we see that, for any divergence-free $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2])$, one has

$$\int_0^T \int_{\Omega} \rho_{\epsilon} u_{\epsilon}^{\perp} \cdot \phi \, dx \, dt = O(\epsilon).$$

Indeed, the uniform bounds established in Section 3.1 show that, for any $T > 0$, one gets $(\rho_{\epsilon} u_{\epsilon})_{\epsilon > 0} \subset L_T^2(L^2)$, $(\rho_{\epsilon} u_{\epsilon} \otimes u_{\epsilon})_{\epsilon > 0}$ and $(b_{\epsilon} \otimes b_{\epsilon})_{\epsilon > 0}$ uniformly bounded in $L_T^{\infty}(L^1)$, and $(\nu(\rho_{\epsilon}) \nabla u_{\epsilon})_{\epsilon > 0} \subset L_T^2(L^2)$, while by assumption $(m_{0,\epsilon})_{\epsilon > 0}$ is bounded in L^2 . Therefore, we can take the limit $\epsilon \rightarrow 0^+$ and get, for any test function ϕ as above, that

$$\int_0^T \int_{\Omega} \rho u^{\perp} \cdot \phi \, dx \, dt = 0.$$

This means that $\rho u^{\perp} = \nabla p$ for some suitable function p , which implies, after taking the curl, the constraint $\text{div}(\rho u) = 0$. With this latter relation at hand, we look at the mass equation: using again that $\rho_{\epsilon} u_{\epsilon} \rightarrow \rho u$ in \mathcal{D}' , we have no trouble in taking the limit $\epsilon \rightarrow 0^+$, and we obtain

$$\partial_t \rho + \text{div}(\rho u) = 0, \quad \text{which implies} \quad \partial_t \rho = 0.$$

The consequence is that $\rho(t, x) = \rho_0(x)$ for almost every (t, x) . In particular, the relation $\text{div}(\rho_0 u) = \nabla \rho_0 \cdot u = 0$ is satisfied almost everywhere in $\mathbb{R}_+ \times \Omega$. \square

3.4 Quantitative convergence properties for the density

This paragraph centers on the density functions in the fully non-homogeneous case. In Subsection 3.2 above, we have shown strong convergence of the densities in $L_{\text{loc}}^2(\mathbb{R}_+ \times \Omega)$. However, this convergence is not enough for the convergence in the fully non-homogeneous case, since neither quantitative nor uniform with respect to time.

As we have seen, there is no obvious way to write $\rho_{\epsilon} = \rho_0 + \epsilon \sigma_{\epsilon}$, with $(\sigma_{\epsilon})_{\epsilon}$ being bounded in some Banach space. Nonetheless, it turns out that the previous decomposition holds true thanks to the structure of the system, but will yield uniform bounds for $(\sigma_{\epsilon})_{\epsilon}$ in the very low regularity space $H^{-3-\delta}$. We refer to Subsection 3.3 of [16] for the original proofs (the presence of the magnetic field introduces only minor modifications).

Proposition 3.5. *Define the functions $\sigma_\epsilon := (\rho_\epsilon - \rho_0)/\epsilon$. Then the sequence $(\sigma_\epsilon)_{\epsilon>0}$ is uniformly bounded in $L_{\text{loc}}^\infty(\mathbb{R}_+; H^{-3-\delta}(\Omega))$ for all $\delta > 0$. In particular, up to an extraction, it weakly-* converges to some σ in that space.*

Proof. For notational convenience, set

$$(25) \quad V_\epsilon = \rho_\epsilon u_\epsilon \quad \text{and} \quad f_\epsilon = \operatorname{div}(\nu(\rho_\epsilon) \nabla u_\epsilon) + \operatorname{div}(b_\epsilon \otimes b_\epsilon - \rho_\epsilon u_\epsilon \otimes u_\epsilon).$$

Because of the Sobolev embedding $H^{1+\delta} \subset L^\infty$ (see the note following Proposition A.3), which holds for any $\delta > 0$, we see that $L^1 \subset H^{-1-\delta}$. Hence, arguing as in the proof of Proposition 3.4, we see that, for any $T > 0$ and any arbitrarily small $\delta > 0$, one has

$$(26) \quad (f_\epsilon)_\epsilon \subset L_T^2(H^{-1}) + L_T^\infty(H^{-2-\delta}) \subset L_T^2(H^{-2-\delta}).$$

Now, because ρ_0 is time-independent, we can write the mass and momentum equations as

$$(27) \quad \begin{cases} \epsilon \partial_t \sigma_\epsilon + \operatorname{div} V_\epsilon = 0 \\ \epsilon \partial_t V_\epsilon + \nabla \pi_\epsilon + V_\epsilon^\perp = \epsilon f_\epsilon. \end{cases}$$

Taking the curl of the second equation and computing the difference with the first one leads to

$$(28) \quad (\partial_t(\eta_\epsilon - \sigma_\epsilon))_{\epsilon>0} = (\operatorname{curl}(f_\epsilon))_{\epsilon>0} \subset L_T^2(H^{-3-\delta}),$$

where we have set $\eta_\epsilon = \operatorname{curl} V_\epsilon$. Notice that $(\eta_\epsilon)_\epsilon \subset L_T^\infty(H^{-1})$ for all $T > 0$ and all $\delta > 0$, in view of (12) and (14), whence the uniform bound $(\sigma_\epsilon)_\epsilon \subset L_T^\infty(H^{-3-\delta})$. In particular, there exists $\sigma \in L_T^\infty(H^{-3-\delta})$ such that $\sigma_\epsilon \xrightarrow{*} \sigma$ in this space. \square

Next, for any $\epsilon > 0$ let us set

$$s_\epsilon := \rho_\epsilon - \rho_0 = \epsilon \sigma_\epsilon.$$

By interpolating between the regularity of s_ϵ and the one of σ_ϵ , we want to show convergence of the densities in better spaces, at the cost of losing the linear convergence speed $O(\epsilon)$, which we will have to replace by $O(\epsilon^\theta)$, for some $0 < \theta < 1$.

Proposition 3.6. *Given $0 < \gamma < 1$, there exists $(\theta, \beta, k) \in \mathbb{R}^3$, satisfying*

$$0 < \beta < \gamma < k < 1 \quad \text{and} \quad 0 < \theta < 1,$$

such that the following uniform embeddings,

$$(\epsilon^{-\theta} s_\epsilon)_{\epsilon>0} \subset C^{0,\beta}([0, T]; H^{-k}(\Omega)) \quad \text{and} \quad (\epsilon^{-\theta} s_\epsilon u_\epsilon)_{\epsilon>0} \subset L^2([0, T]; H^{-k-\delta}(\Omega)),$$

hold true for any $T > 0$ and any arbitrarily small $\delta > 0$. In addition,

$$\epsilon^{-\theta} s_\epsilon \longrightarrow 0 \quad \text{in } L^\infty([0, T]; H_{\text{loc}}^{-k-\delta}(\Omega)) \quad \text{and} \quad \epsilon^{-\theta} s_\epsilon u_\epsilon \rightharpoonup 0 \quad \text{in } L^2([0, T]; H_{\text{loc}}^{-k-\delta}(\Omega))$$

for any $T > 0$ and any small $\delta > 0$.

Proof. The function s_ϵ solves a transport equation with a second member:

$$\partial_t s_\epsilon + \operatorname{div}(s_\epsilon u_\epsilon) = -u_\epsilon \cdot \nabla \rho_0.$$

with initial datum $(s_\epsilon)|_{t=0} = \epsilon r_{0,\epsilon}$. Because we have assumed that $\rho_0 \in C_b^2$, Sobolev embeddings show that, for any fixed $T > 0$, the sequence $(u_\epsilon \cdot \nabla \rho_0)_\epsilon \subset L_T^2(H^1)$ is bounded in every space

$L_T^2(L^q)$, for $2 \leq q < +\infty$. In addition, $s_\epsilon = \rho_\epsilon - \rho_0$ is trivially uniformly bounded in $L^\infty(L^\infty)$; therefore, we finally infer, for all fixed $T > 0$, the uniform bounds

$$(s_\epsilon)_{\epsilon>0} \subset L_T^\infty(L^2 \cap L^\infty).$$

Furthermore, writing $\partial_t s_\epsilon = -\operatorname{div}(\rho_\epsilon u_\epsilon)$ and reasoning as in the proof of Lemma 3.3, we see that $(s_\epsilon)_{\epsilon>0} \subset W_T^{1,\infty}(H^{-1})$ for all $T > 0$; after interpolation between Sobolev spaces, we get, for every $0 \leq \gamma \leq 1$, the embedding

$$(s_\epsilon)_{\epsilon>0} \subset C_T^{0,\gamma}(H^{-\gamma}).$$

On the other hand, from Proposition 3.6 we know that $(\sigma_\epsilon)_\epsilon \subset L_T^\infty(H^{-3-\delta})$ for $T > 0$ and arbitrarily small $\delta > 0$. Therefore, for $0 \leq t_1, t_2 \leq T$, and for $0 < \theta < 1$ such that $k = \gamma(1 - \theta) + (3 + \delta)\theta$, we have

$$\begin{aligned} \|s_\epsilon(t_2) - s_\epsilon(t_1)\|_{H^{-k}} &\leq \|s_\epsilon(t_2) - s_\epsilon(t_1)\|_{H^{-\gamma}}^{1-\theta} \|s_\epsilon(t_2) - s_\epsilon(t_1)\|_{H^{-3-\delta}}^\theta \\ &\leq 2 \|s_\epsilon\|_{C_T^{0,\gamma}(H^{-\gamma})}^{1-\theta} |t_2 - t_1|^{\gamma(1-\theta)} \epsilon^\theta \|\sigma_\epsilon\|_{L_T^\infty(H^{-3-\delta})}^\theta. \end{aligned}$$

By setting $\beta = (1 - \theta)\gamma$, we get $(\epsilon^{-\theta} s_\epsilon)_\epsilon \subset C_T^{0,\beta}(H^{-k})$, as claimed. We deduce from the Ascoli-Arzelà theorem that the sequence $(\epsilon^{-\theta} s_\epsilon)_\epsilon$ is compact in $L_T^\infty(H_{\text{loc}}^{-k-\delta})$, for arbitrarily small $\delta > 0$, so that it converges strongly to some s in that space. Finally, we remark that we must have $s = 0$, because $\epsilon^{-\theta} s_\epsilon = \epsilon^{1-\theta} \sigma_\epsilon \rightarrow 0$ in \mathcal{D}' .

Next, Corollary A.6 gives continuity of the function product in the $H^{-k} \times H^1 \rightarrow H^{-k-\delta}$ topology, for arbitrarily small $\delta > 0$, whence the uniform bound

$$(\epsilon^{-\theta} s_\epsilon u_\epsilon)_{\epsilon>0} \subset L_T^2(H^{-k-\delta}),$$

for all fixed $T > 0$. In addition, using the strong convergence $\epsilon^{-\theta} s_\epsilon \rightarrow 0$ in $L_T^\infty(H_{\text{loc}}^{-k-\delta})$, we get the weak convergence $\epsilon^{-\theta} s_\epsilon u_\epsilon \rightharpoonup 0$ in $L_T^2(H_{\text{loc}}^{-k-\delta})$. \square

4 Convergence

In this section we complete the proof of Theorems 2.2 and 2.3, up to the uniqueness part of the former statement (which will be considered in Section 5 below). Subsections 4.1 and 4.2 are common to both the quasi-homogeneous and the fully non-homogeneous case. In Subsection 4.3 we take care of the convective term in the quasi-homogeneous case. Subsections 4.4 and 4.5 are dedicated to the fully non-homogeneous case. Discussion first bears on the convergence of the convective term, which is the most involved part of the proof; lastly, we handle the Coriolis term. In both parts, we thoroughly exploit the decomposition $\rho_\epsilon = \rho_0 + \epsilon \sigma_\epsilon$ and the vorticity form of the momentum equation.

4.1 The magnetic field

In this section, we take care of all the terms containing the magnetic fields b_ϵ , except for the resistivity term $\nabla^\perp(\mu(\rho_\epsilon) \operatorname{curl} b_\epsilon)$. As in the previous section with the continuity equation, we use the magnetic field equation to trade space regularity against time compactness.

Proposition 4.1. *We have the following strong convergence for the magnetic fields: up to the extraction of a suitable subsequence, for any $0 < s < 1$ we have*

$$b_\epsilon \xrightarrow{\epsilon \rightarrow 0^+} b \quad \text{in} \quad L_{\text{loc}}^2(\mathbb{R}_+; H_{\text{loc}}^s(\Omega)).$$

In particular, we deduce the convergence of all bilinear terms involving the magnetic field:

$$\operatorname{div}(b_\epsilon \otimes b_\epsilon) \rightarrow \operatorname{div}(b \otimes b) \quad \text{in} \quad \mathcal{D}'(\mathbb{R}_+ \times \Omega),$$

and analogous convergence properties hold for $\operatorname{div}(u_\epsilon \otimes b_\epsilon)$ and $\operatorname{div}(b_\epsilon \otimes u_\epsilon)$.

Proof. Let $T > 0$ be a fixed positive time. Recall that we have $(b_\epsilon)_\epsilon \subset L^\infty(L^2)$, as well as $(u_\epsilon)_\epsilon, (b_\epsilon)_\epsilon \subset L_T^2(H^1)$. The magnetic field equation reads

$$\partial_t b_\epsilon = \operatorname{div} (b_\epsilon \otimes u_\epsilon - u_\epsilon \otimes b_\epsilon) + \nabla^\perp (\mu(\rho_\epsilon) \operatorname{curl} (b_\epsilon)).$$

Gagliardo-Nirenberg inequality of Lemma A.7 gives $(b_\epsilon)_\epsilon \subset L_T^4(L^4)$ and $(u_\epsilon)_\epsilon \subset L_T^2(L^4)$. Hence $(u_\epsilon \otimes b_\epsilon)_\epsilon \subset L_T^{4/3}(L^2)$ and $(\partial_t b_\epsilon)_\epsilon \subset L_T^{4/3}(H^{-1})$. As a result, we get the Hölder bound $(b_\epsilon)_\epsilon \subset C_T^{0,3/4}(H^{-1})$, and the Ascoli-Arzelà theorem gives compactness of $(b_\epsilon)_\epsilon$ in e.g. $C_T^0(H_{\operatorname{loc}}^{-1-\delta})$, for any small $\delta > 0$. Using the uniform bound in $L_T^2(H^1)$ and Sobolev interpolation gives strong convergence of the magnetic fields $(b_\epsilon)_\epsilon$ in $L_T^2(H_{\operatorname{loc}}^s)$, for any $0 \leq s < 1$.

As a consequence, we gather strong convergence of the tensor products $(b_\epsilon \otimes b_\epsilon)_\epsilon$ in, say, $L_T^1(L_{\operatorname{loc}}^1)$, which implies in particular that $\operatorname{div} (b_\epsilon \otimes b_\epsilon) \rightarrow \operatorname{div} (b \otimes b)$ in $\mathcal{D}'([0, T] \times \Omega)$. In an analogous way, we can achieve weak convergence of the mixed tensor products $b_\epsilon \otimes u_\epsilon$ and $u_\epsilon \otimes b_\epsilon$: for the sake of brevity, we omit the details here. \square

Note that this proposition does not complete the study of the magnetic field equation: the convergence of the resistivity term $\nabla^\perp (\mu(\rho_\epsilon) \operatorname{curl} (b_\epsilon))$ still remains. This is the goal of the next paragraph.

4.2 The viscosity and resistivity terms

In this section, we take care of the convergence of the viscosity and resistivity terms, namely $\operatorname{div} (\nu(\rho_\epsilon) \nabla u_\epsilon)$ and $\nabla^\perp (\mu(\rho_\epsilon) \operatorname{curl} (b_\epsilon))$ respectively. Remember that we have taken ν and μ to be continuous on \mathbb{R}_+ .

Proposition 4.2. *The following convergence of the viscosity and resistivity terms holds true, in the sense of $\mathcal{D}'(\mathbb{R}_+ \times \Omega)$:*

$$\operatorname{div} (\nu(\rho_\epsilon) \nabla u_\epsilon) \rightarrow \operatorname{div} (\nu(\rho_0) \nabla u) \quad \text{and} \quad \nabla^\perp (\mu(\rho_\epsilon) \operatorname{curl} (b_\epsilon)) \rightarrow \nabla^\perp (\mu(\rho_0) \operatorname{curl} (b)),$$

where ρ_0 is either 1 (in the quasi-homogeneous case) or the truly variable profile satisfying the assumptions of Theorem 2.3 (in the fully-non-homogeneous case).

Proof. We only prove the convergence of the viscosity term, the one of the resistivity term being, in all that matters, identical.

In the case where the density is quasi-homogeneous, we already have strong convergence $\rho_\epsilon \rightarrow 1$ in $L^\infty(L^\infty)$, because $(r_\epsilon)_\epsilon$ is bounded in $L^\infty(L^2 \cap L^\infty)$. This makes the viscosity term easy to handle: let $T > 0$ and $\phi \in \mathcal{D}([0, T] \times \Omega; \mathbb{R}^2)$ be a test function, then

$$\begin{aligned} & \int_0^T \int_\Omega \nu(\rho_\epsilon) \nabla u_\epsilon : \nabla \phi \, dx \, dt - \int_0^T \int_\Omega \nu(1) \nabla u : \nabla \phi \, dx \, dt \\ &= \int_0^T \int_\Omega (\nu(\rho_\epsilon) - \nu(1)) \nabla u_\epsilon : \nabla \phi \, dx \, dt + \int_0^T \int_\Omega \nu(1) (\nabla u_\epsilon - \nabla u) : \nabla \phi \, dx \, dt. \end{aligned}$$

The second integral has limit zero, because of the weak convergence of the u_ϵ in $L_T^2(H^1)$. As for the first integral, uniform convergence of the ρ_ϵ and continuity of ν gives

$$\left| \int_0^T \int_\Omega (\nu(\rho_\epsilon) - \nu(1)) \nabla u_\epsilon : \nabla \phi \right| \leq \|\nu(\rho_\epsilon) - \nu(1)\|_{L^\infty(L^\infty)} \|\nabla u_\epsilon\|_{L_T^2(L^2)} \|\nabla \phi\|_{L_T^2(L^2)} \xrightarrow{\epsilon \rightarrow 0^+} 0.$$

Obviously, this does not work as well in the fully non-homogeneous case. Hence, we will need to use the strong convergence result of Proposition 3.1: namely, in the limit $\epsilon \rightarrow 0^+$, one has $\rho_\epsilon \rightarrow \rho_0$ in $L_{\operatorname{loc}}^2(\mathbb{R}_+ \times \Omega)$ and almost everywhere in $\mathbb{R}_+ \times \Omega$, where we have used also the information coming from the second item of Proposition 3.4.

The uniform bounds $(\rho_\epsilon)_{\epsilon > 0} \subset L^\infty(L^\infty)$ and the continuity of ν show that the $\nu(\rho_\epsilon)$ are also uniformly bounded in $L^\infty(L^\infty)$: namely, there is a constant $\nu^* > 0$ such that $\nu(\rho_\epsilon) \leq \nu^*$ for all $\epsilon > 0$. The dominated convergence theorem then gives strong convergence of the viscosities:

$$(29) \quad \nu(\rho_\epsilon) \xrightarrow{\epsilon \rightarrow 0^+} \nu(\rho_0) \quad \text{in} \quad L^2_{\text{loc}}(\mathbb{R}_+ \times \Omega).$$

Now, let $T > 0$ and $K \subset \Omega$ be a compact set. For any $\phi \in \mathcal{D}([0, T[\times K)$, we can estimate

$$\begin{aligned} & \left| \int_0^T \int_\Omega \nu(\rho_\epsilon) \nabla u_\epsilon \cdot \nabla \phi \, dx \, dt - \int_0^T \int_\Omega \nu(\rho_0) \nabla u \cdot \nabla \phi \, dx \, dt \right| \\ & \leq \left| \int_0^T \int_\Omega \nu(\rho_0) (\nabla u_\epsilon - \nabla u) \cdot \nabla \phi \, dx \, dt \right| + \left| \int_0^T \int_\Omega (\nu(\rho_\epsilon) - \nu(\rho_0)) \nabla u_\epsilon \cdot \nabla \phi \, dx \, dt \right|. \end{aligned}$$

The first integral obviously tends to zero as $\epsilon \rightarrow 0^+$, because of weak convergence $\nabla u_\epsilon \rightharpoonup \nabla u$ in $L^2_T(L^2)$. The second integral, instead, can be bounded as follows:

$$\left| \int_0^T \int_\Omega (\nu(\rho_\epsilon) - \nu(\rho_0)) \nabla u_\epsilon \cdot \nabla \phi \, dx \, dt \right| \leq \|\nabla \phi\|_{L^\infty(L^\infty)} \|\nu(\rho_\epsilon) - \nu(\rho_0)\|_{L^2_T(L^2(K))} \|\nabla u_\epsilon\|_{L^2(L^2)},$$

where the quantity on the right-hand side goes to 0 in view of (29).

The proposition is now proved. \square

4.3 The convective term: the quasi-homogeneous case

In the slightly non-homogeneous case, the convective term $\text{div}(\rho_\epsilon u_\epsilon \otimes u_\epsilon)$ is the last one we have to study. The argument is in three steps. First of all, we reduce the problem to the study of $\text{div}(u_\epsilon \otimes u_\epsilon)$, taking advantage of the approximation $\rho_\epsilon \approx 1$. Then, we use the uniform H^1 regularity to find an approximation of u_ϵ by smooth functions, which we will need for the last step, a compensated compactness argument.

Proposition 4.3. *For all divergence-free $\phi \in \mathcal{D}(\mathbb{R}_+ \times \Omega; \mathbb{R}^2)$, one has*

$$\int_0^{+\infty} \int_\Omega \rho_\epsilon u_\epsilon \otimes u_\epsilon : \nabla \phi \, dx \, dt \xrightarrow{\epsilon \rightarrow 0^+} \int_0^{+\infty} \int_\Omega u \otimes u : \nabla \phi \, dx \, dt.$$

Proof. Let $T > 0$ be a fixed positive time. To start the proof, recall that we can write $\rho_\epsilon = 1 + \epsilon r_\epsilon$, with $(r_\epsilon)_\epsilon \subset L^\infty(L^2 \cap L^\infty)$. Then, by virtue of the $L^2_T(L^2)$ uniform boundedness of $(u_\epsilon)_\epsilon$, for all divergence-free $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2)$ we have

$$(30) \quad \left| \int_0^T \int_\Omega (\rho_\epsilon u_\epsilon \otimes u_\epsilon - u_\epsilon \otimes u_\epsilon) : \nabla \phi \, dx \, dt \right| \xrightarrow{\epsilon \rightarrow 0^+} 0.$$

Next, we seek a uniform approximation of u_ϵ by a smooth function (that is, smooth in the space variable). Let S_j be the low-frequency cut-off operator from the Littlewood-Paley decomposition given by (56) in the appendix. Then, using the uniform bound $(u_\epsilon)_\epsilon \subset L^2_T(H^1)$ and the characterisation (57) of Sobolev spaces, it immediately follows that

$$\|(I - S_j)u_\epsilon\|_{L^2_T(L^2)} \leq C 2^{-j},$$

for some constant $C > 0$ possibly depending on T , but independent of both j and ϵ . Using the previous bound, we can thus estimate

$$(31) \quad \left| \int_0^T \int_\Omega \left\{ u_\epsilon \otimes u_\epsilon - S_j u_\epsilon \otimes S_j u_\epsilon \right\} : \nabla \phi \, dx \, dt \right|$$

$$\leq \|\nabla\phi\|_{L^\infty(L^\infty)} \left(\|(I - S_j)u_\epsilon \otimes u_\epsilon\|_{L_T^1(L^1)} + \|S_j u_\epsilon \otimes (I - S_j)u_\epsilon\|_{L_T^1(L^1)} \right) \leq C 2^{-j}.$$

Therefore, the problem is now to prove convergence of the term

$$\int_0^T \int_\Omega S_j u_\epsilon \otimes S_j u_\epsilon : \nabla \phi \, dx \, dt = - \int_0^T \int_\Omega \operatorname{div} (S_j u_\epsilon \otimes S_j u_\epsilon) \cdot \phi \, dx \, dt$$

when $\epsilon \rightarrow 0^+$, for any fixed $j \geq -1$. Notice that the integration by parts is justified, since all the quantities are now smooth with respect to the space variable. For convenience purposes, we henceforth note $u_{\epsilon,j} = S_j u_\epsilon$. Because the operator S_j is a Fourier-multiplier, it commutes with all the partial derivatives: in particular, $\operatorname{div}(u_{\epsilon,j}) = 0$ and therefore, after denoting $\omega_\epsilon := \operatorname{curl} u_\epsilon$ and $\omega_{\epsilon,j} := S_j \omega_\epsilon = \operatorname{curl} u_{\epsilon,j}$, we deduce

$$(32) \quad \operatorname{div} (u_{\epsilon,j} \otimes u_{\epsilon,j}) = \frac{1}{2} \nabla |u_{\epsilon,j}|^2 + \omega_{\epsilon,j} u_{\epsilon,j}^\perp.$$

The first term in the right-hand side disappears when tested against a divergence-free function, hence we deduce

$$\int_0^T \int_\Omega u_{\epsilon,j} \otimes u_{\epsilon,j} : \nabla \phi \, dx \, dt = - \int_0^T \int_\Omega \omega_{\epsilon,j} u_{\epsilon,j}^\perp \cdot \phi \, dx \, dt.$$

As for the vorticity term, we resort to the reformulation (27) of the momentum equation, which can be rewritten, in the quasi-homogeneous case, in the following way:

$$\epsilon \partial_t V_\epsilon + \nabla \pi_\epsilon + \frac{1}{2} \epsilon \nabla |b_\epsilon|^2 + u_\epsilon^\perp = \epsilon (f_\epsilon - r_\epsilon u_\epsilon^\perp),$$

where V_ϵ and f_ϵ have been defined in (25). Then, applying the operator S_j to the previous equation and taking the curl gives

$$\partial_t \eta_{\epsilon,j} = \operatorname{curl} \left(f_{\epsilon,j} - S_j (r_\epsilon u_\epsilon^\perp) \right),$$

where we have set $\eta_{\epsilon,j} := S_j \operatorname{curl} (V_\epsilon)$ and $f_{\epsilon,j} = S_j f_\epsilon$. From (26), we know that $(f_\epsilon)_\epsilon$ is bounded in $L_T^2(H^{-2-\delta})$ for any $\delta > 0$, and so, for any fixed j , the sequence $(f_{\epsilon,j})_\epsilon$ is bounded in every $L_T^2(H^m)$, with $m \in \mathbb{R}$. Likewise, because the sequence $(r_\epsilon u_\epsilon)_\epsilon$ is bounded in $L_T^2(L^2)$, we also see that $(r_\epsilon u_\epsilon^\perp)_\epsilon \subset L_T^2(H^m)$ for every $m \in \mathbb{R}$. We deduce a uniform bound for the $\eta_{\epsilon,j}$: for $m \in \mathbb{R}$,

$$(33) \quad (\eta_{\epsilon,j})_\epsilon \subset C_T^{0,1/2}(H^m).$$

Hence, the Ascoli-Arzelà theorem provides strong convergence (up to the extraction of a subsequence) to some $\eta_j \in L_{\operatorname{loc}}^\infty(H_{\operatorname{loc}}^m)$: more precisely, for all fixed $T > 0$ and $m \in \mathbb{R}$,

$$\forall j \geq 1, \quad \eta_{\epsilon,j} \xrightarrow{\epsilon \rightarrow 0^+} \eta_j \quad \text{in } L_T^\infty(H_{\operatorname{loc}}^m).$$

But since we already know that $V_\epsilon \rightharpoonup u$ in $L_T^2(L^2)$, it follows that $\eta_j = \omega_j$. Thanks to the previous strong convergence property, for fixed $j \geq 1$ and for every $m \in \mathbb{R}$, we get also the strong convergence of $(\omega_{\epsilon,j})$: namely,

$$\omega_{\epsilon,j} = \eta_{\epsilon,j} - \epsilon S_j \operatorname{curl} (r_\epsilon u_\epsilon) \xrightarrow{\epsilon \rightarrow 0^+} \omega_j \quad \text{in } L_T^\infty(H_{\operatorname{loc}}^m).$$

Combining this information with the weak convergence $u_\epsilon \rightharpoonup u$ in $L_T^2(H^1)$ finally yields

$$\omega_{\epsilon,j} u_{\epsilon,j}^\perp \xrightarrow{\epsilon \rightarrow 0^+} \omega_j u_j^\perp \quad \text{in } \mathcal{D}'([0, T] \times \Omega).$$

Thus, we have proved that, for any divergence-free test function $\phi \in \mathcal{D}([0, T] \times \Omega)$, we have

$$\langle \operatorname{div} (u_{\epsilon,j} \otimes u_{\epsilon,j}), \phi \rangle_{\mathcal{D}' \times \mathcal{D}} = \langle \omega_{\epsilon,j} u_{\epsilon,j}^\perp, \phi \rangle_{\mathcal{D}' \times \mathcal{D}} \xrightarrow{\epsilon \rightarrow 0^+} \langle \omega_j u_j^\perp, \phi \rangle_{\mathcal{D}' \times \mathcal{D}} = \langle \operatorname{div} (u_j \otimes u_j), \phi \rangle_{\mathcal{D}' \times \mathcal{D}}.$$

Now, keeping in mind (30), using the uniform approximation property (31) yields the claimed convergence result. \square

4.4 The convective term: the fully non-homogeneous case

The main ideas for handling the convective term in the fully non-homogeneous case are very similar to those used for the quasi-homogeneous case, although many complications occur. Because ρ_0 is not constant, the equivalent of decomposition (32) will instead be (omitting for the time being the regularisation argument)

$$\operatorname{div}(\rho_\epsilon u_\epsilon \otimes u_\epsilon) \approx \operatorname{div}(\rho_0 u_\epsilon \otimes u_\epsilon) = \frac{1}{2} \rho_0 \nabla |u_\epsilon|^2 + \rho_0 \omega_\epsilon u_\epsilon^\perp + (u_\epsilon \cdot \nabla \rho_0) u_\epsilon.$$

To simplify those terms, we can no longer rely on the fact that we use divergence-free test functions: $\langle \rho_0 \nabla |u_\epsilon|^2, \phi \rangle \neq 0$ even when $\operatorname{div}(\phi) = 0$. However, any term of the form $\rho_0 \nabla \Lambda_\epsilon$ or $\Lambda_\epsilon \nabla \rho_0$ will give rise to a term of the form $\rho_0 \nabla \Gamma$ in the limit (see below), which can be considered as a “pressure” term associated to the constraint $\operatorname{div}(\rho_0 u) = 0$.

In the end, we can prove the next statement.

Proposition 4.4. *There is a distribution Γ (of order at most one) such that, for all $\phi \in \mathcal{D}(\mathbb{R}_+ \times \Omega; \mathbb{R}^2)$ such that $\operatorname{div}(\phi) = 0$, one has*

$$\int_0^{+\infty} \int_\Omega \rho_\epsilon u_\epsilon \otimes u_\epsilon : \nabla \phi \, dx \, dt \xrightarrow{\epsilon \rightarrow 0^+} \langle \rho_0 \nabla \Gamma, \phi \rangle_{\mathcal{D}' \times \mathcal{D}}.$$

The rest of this subsection is devoted to the proof of the previous proposition. Our argument, mainly borrowed from [16], consists of several steps. We start by taking a positive time $T > 0$, a compact set $K \subset \Omega$ and a divergence-free test function $\phi \in \mathcal{D}([0, T[\times K; \mathbb{R}^2)$, which we keep fixed throughout all the proof.

Step 1: approximation of the densities. First of all, we justify the approximation $\rho_\epsilon u_\epsilon \otimes u_\epsilon \approx \rho_0 u_\epsilon \otimes u_\epsilon$. Note that, because we have accounted for the presence of vacuum, the best uniform bound we have for the velocity field is $(u_\epsilon)_\epsilon \subset L_T^2(H^1)$. This means that, when estimating the difference $(\rho_\epsilon - \rho_0)u_\epsilon \otimes u_\epsilon$, we must use strong convergence of the densities *uniformly with respect to time*. In particular, we cannot benefit of the convergence properties proved in Proposition 4.2, which only provides strong convergence $\rho_\epsilon \rightarrow \rho_0$ in the spaces $L_T^p(L_{\text{loc}}^q)$ for $1 \leq p, q < +\infty$ (thanks to the uniform bound $0 \leq \rho_\epsilon \leq \rho^*$ and dominated convergence). Instead, from Proposition 3.6 we know that $\rho_\epsilon = \rho_0 + s_\epsilon$, with $s_\epsilon \rightarrow 0$ in $C_T^{0,\beta}(H_{\text{loc}}^{-k})$, where $\beta, k \in]0, 1[$ are as in that proposition (relatively to some $\gamma \in]0, 1[$).

Firstly, using Corollary A.6, we see that, for every $\delta > 0$, the function product is continuous in the $H^1 \times H^1 \rightarrow H^{1-\delta}$ topology. This implies that the tensor product $(u_\epsilon \otimes u_\epsilon)_\epsilon$ is bounded in the space $L_T^1(H^{1-\delta})$, for every $\delta > 0$. Next, using Lemma A.5, we get continuity of the product in the $H^{-k} \times H^{1-\delta} \rightarrow H^{-k-\delta}$ topology provided that $\delta > 0$ be small enough (i.e. small enough for $1 - k - \delta$ to be positive). We therefore gather that

$$\|s_\epsilon u_\epsilon \otimes u_\epsilon\|_{L_T^1(H^{-k-\delta})} \leq \|s_\epsilon\|_{L_T^\infty(H^{-k})} \|u_\epsilon \otimes u_\epsilon\|_{L_T^1(H^{1-\delta})} \xrightarrow{\epsilon \rightarrow 0^+} 0$$

and are reconducted to taking the limit $\epsilon \rightarrow 0^+$ in the integral

$$\int_0^T \int_\Omega \rho_0 u_\epsilon \otimes u_\epsilon : \nabla \phi \, dx \, dt.$$

At this point, the idea is to resort once again to a compensated compactness argument: for this, we need first to smooth out the velocity fields.

Step 2: regularisation. We use the same regularisation procedure than in the quasi-homogeneous case: here S_j still is the Littlewood-Paley operator defined by (56) below. We continue to note $S_j g_\epsilon = g_{\epsilon,j}$ for any sequence of functions $(g_\epsilon)_{\epsilon>0}$, whenever we feel it make things more legible.

We set as above $\eta_\epsilon = \text{curl } V_\epsilon = \text{curl } (\rho_\epsilon u_\epsilon)$. We now state a simple approximation lemma: its proof is simple, hence omitted. It is enough to recall that $(\eta_\epsilon)_\epsilon$ is bounded in $L_T^\infty(H^{-1})$ and that $(\sigma_\epsilon)_\epsilon$ is bounded in every $L_T^\infty(H^{-3-\delta})$, with $\delta > 0$.

Lemma 4.5. *The following uniform properties hold, in the limit for $j \rightarrow +\infty$:*

1) for all $s > 3$, we have $\sup_{\epsilon > 0} \|\sigma_\epsilon - S_j \sigma_\epsilon\|_{L_T^\infty(H^{-s})} \rightarrow 0$;

2) for all $s > 1$, we have $\sup_{\epsilon > 0} \|\eta_\epsilon - S_j \eta_\epsilon\|_{L_T^\infty(H^{-s})} \rightarrow 0$.

The first new problem we face, after introducing S_j , is that $S_j(\rho_\epsilon u_\epsilon) \neq \rho_0 S_j u_\epsilon$, because ρ_0 is no longer constant. With the same notation introduced in Section 3.4, we write

$$(34) \quad S_j(\rho_\epsilon u_\epsilon) = S_j(\rho_0 u_\epsilon) + \epsilon^\theta S_j(\epsilon^{-\theta} s_\epsilon u_\epsilon) = \rho_0 u_{\epsilon,j} + [S_j, \rho_0] u_\epsilon + \epsilon^\theta S_j(\epsilon^{-\theta} s_\epsilon u_\epsilon).$$

In the above, $[S_j, \rho_0]$ is the commutator between S_j and the multiplication by ρ_0 operator. By Proposition 3.6, we already know that $(S_j(\epsilon^{-\theta} s_\epsilon u_\epsilon))_\epsilon \subset L_T^2(H^s)$ for any $s \geq 0$. To deal with the second summand, we use Lemma A.9 on $[S_j, \rho_0] u_\epsilon$. On the one hand, we have

$$\|[S_j, \rho_0] u_\epsilon\|_{L_T^2(L^2)} \leq \frac{C}{2^j} \|\nabla \rho_0\|_{L^\infty} \|u_\epsilon\|_{L_T^2(L^2)} \leq C 2^{-j}.$$

On the other hand, by differentiating the commutator, we get, for $i \in \{1, 2\}$,

$$\partial_i [S_j, \rho_0] u_\epsilon = [S_j, \partial_i \rho_0] u_\epsilon + [S_j, \rho_0] \partial_i u_\epsilon.$$

Therefore, from Lemma A.9 we infer

$$\|\partial_i [S_j, \rho_0] u_\epsilon\|_{L_T^2(L^2)} \leq \frac{C}{2^j} \left\{ \|\nabla^2 \rho_0\|_{L^\infty} \|u_\epsilon\|_{L_T^2(L^2)} + \|\nabla \rho_0\|_{L^\infty} \|\nabla u_\epsilon\|_{L_T^2(L^2)} \right\} \leq C 2^{-j}.$$

Thus, from (34), we have obtained the following decomposition of $S_j(\rho_\epsilon u_\epsilon)$:

$$(35) \quad S_j(\rho_\epsilon u_\epsilon) = \rho_0 u_{\epsilon,j} + \epsilon^\theta \zeta_{\epsilon,j} + h_{\epsilon,j},$$

where we have defined $\zeta_{\epsilon,j} := S_j(\epsilon^{-\theta} s_\epsilon u_\epsilon)$ and $h_{\epsilon,j} := [S_j, \rho_0] u_\epsilon$. Notice that

$$(36) \quad \forall j \geq -1, \forall s \geq 0, \quad (\zeta_{\epsilon,j})_\epsilon \subset L_T^2(H^s) \quad \text{and} \quad \sup_{\epsilon > 0} \|h_{\epsilon,j}\|_{L_T^2(H^1)} \leq C 2^{-j}.$$

We make a couple of remarks before going onwards. Firstly, we have seen in Proposition 3.6 that $(\epsilon^{-\theta} s_\epsilon u_\epsilon)_\epsilon$ is bounded in $L_T^2(H^{-k-\delta})$, where $0 < k < 1$ has been fixed in that proposition and $\delta > 0$ is arbitrarily small. Therefore, we can further write

$$(37) \quad \eta_{\epsilon,j} = S_j \text{curl} \left[\rho_0 u_\epsilon + \epsilon^\theta (\epsilon^{-\theta} s_\epsilon u_\epsilon) \right] = \eta_{\epsilon,j}^{(1)} + \epsilon^\theta \eta_{\epsilon,j}^{(2)},$$

with the uniform bounds (with respect to ϵ)

$$\left\| \eta_{\epsilon,j}^{(1)} \right\|_{L_T^2(L^2)} \leq C_1 \quad \text{and} \quad \left\| \eta_{\epsilon,j}^{(2)} \right\|_{L_T^2(H^s)} \leq C(s, j),$$

for any given $s \geq 0$. Note that the constant C_1 does not depend on ϵ nor on j .

Finally, exactly as in the quasi-homogeneous case (keep in mind estimate (31) above), thanks to the uniform approximation properties of S_j , we note that it is enough to prove the convergence of the term

$$\int_0^T \int_\Omega \rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j} : \nabla \phi \, dx \, dt = - \int_0^T \int_\Omega \text{div} (\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) \cdot \phi \, dx \, dt,$$

where the integration by parts is now well-justified, since all the quantities in the integral are smooth in the space variable.

Step 3: reformulation. In view of the previous discussion, we are left with $\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j})$. Since all functions are smooth, we can write

$$\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = (u_{\epsilon,j} \cdot \nabla \rho_0) u_{\epsilon,j} + \rho_0 \omega_{\epsilon,j} u_{\epsilon,j}^\perp + \frac{1}{2} \rho_0 \nabla |u_{\epsilon,j}|^2,$$

with $\omega_{\epsilon,j} = S_j \operatorname{curl}(u_\epsilon)$. We remark that the last term in the righthand side of this equation contributes $\rho_0 \nabla \Gamma$ in the limit, for some distribution Γ of order at most one. In the same way, any term of the form $\langle \Lambda_{\epsilon,j} \nabla \rho_0, \phi \rangle$ has a limit of the same form $\langle \rho_0 \nabla \Gamma, \phi \rangle$: since $\operatorname{div} \phi = 0$, an integration by parts gives

$$\int_0^T \int_\Omega \Lambda_{\epsilon,j} \nabla \rho_0 \cdot \phi \, dx \, dt = \int_0^T \int_\Omega \Lambda_{\epsilon,j} \operatorname{div}(\rho_0 \phi) \, dx \, dt = - \int_0^T \int_\Omega \rho_0 \nabla \Lambda_{\epsilon,j} \cdot \phi \, dx \, dt.$$

Since all terms of the form $\rho_0 \nabla \Lambda_{\epsilon,j}^{(1)} + \Lambda_{\epsilon,j}^{(2)} \nabla \rho_0$ can be treated in this way, we will generically note any of them by $\Gamma_{\epsilon,j}$. Likewise, we note $R_{\epsilon,j}$ any remainder term, that is any term such that

$$\lim_{j \rightarrow +\infty} \limsup_{\epsilon \rightarrow 0^+} \left| \int_0^T \int_\Omega R_{\epsilon,j} \cdot \phi \, dx \, dt \right| = 0.$$

With that notation, we can write

$$\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = (u_{\epsilon,j} \cdot \nabla \rho_0) u_{\epsilon,j} + \rho_0 \omega_{\epsilon,j} u_{\epsilon,j}^\perp + \Gamma_{\epsilon,j}.$$

We are now going to deal with the vorticity term, namely the second term in the right-hand side of the previous equation.

Step 4: the vorticity term. By use of (35), we get

$$\begin{aligned} \eta_{\epsilon,j} &= \operatorname{curl}(\rho_0 u_{\epsilon,j}) + \epsilon^\theta \operatorname{curl} \zeta_{\epsilon,j} + \operatorname{curl} h_{\epsilon,j} \\ &= \rho_0 \omega_{\epsilon,j} + \nabla^\perp \rho_0 \cdot u_{\epsilon,j} + \epsilon^\theta \operatorname{curl} \zeta_{\epsilon,j} + \operatorname{curl} h_{\epsilon,j}. \end{aligned}$$

Therefore, by virtue of (36), we gather that $\rho_0 \omega_{\epsilon,j} u_{\epsilon,j}^\perp = \eta_{\epsilon,j} u_{\epsilon,j}^\perp - (u_{\epsilon,j} \cdot \nabla^\perp \rho_0) u_{\epsilon,j}^\perp + R_{\epsilon,j}$, which in turn gives

$$\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = \eta_{\epsilon,j} u_{\epsilon,j}^\perp + (u_{\epsilon,j} \cdot \nabla \rho_0) u_{\epsilon,j} - (u_{\epsilon,j} \cdot \nabla^\perp \rho_0) u_{\epsilon,j}^\perp + \Gamma_{\epsilon,j} + R_{\epsilon,j}.$$

Step 5: a geometric property. Now we focus on the term

$$X_{\epsilon,j} := (u_{\epsilon,j} \cdot \nabla \rho_0) u_{\epsilon,j} - (u_{\epsilon,j} \cdot \nabla^\perp \rho_0) u_{\epsilon,j}^\perp.$$

The main idea to treat this term is to decompose $u_{\epsilon,j}(t, x)$ in the orthonormal basis of \mathbb{R}^2 given by $\left(\frac{\nabla \rho_0(x)}{|\nabla \rho_0(x)|}, \frac{\nabla^\perp \rho_0(x)}{|\nabla \rho_0(x)|} \right)$. However, to avoid complications, we have first to deal with those $x \in \Omega$ for which $|\nabla \rho_0(x)|$ is small. More precisely, let $B \in \mathcal{D}(\mathbb{R}^2)$ be such that

$$0 \leq B \leq 1 \quad \text{and} \quad \begin{cases} B(y) = 1 & \text{for } |y| \leq 1 \\ B(y) = 0 & \text{for } |y| \geq 2, \end{cases}$$

and let $B_j(x) = B(2^{j/2} \nabla \rho_0(x))$. The function B_j is so chosen that $|\nabla \rho_0| \geq 2^{-j/2}$ on $\operatorname{Supp}(1 - B_j)$. Recall that $\operatorname{Supp} \phi \subset [0, T] \times K$, where ϕ is the divergence-free test function we have fixed at the beginning of the argument. Then, for any $2 < q < +\infty$, Hölder's inequality with $1 = (2/q) + (q-2)/q$ yields

$$\|B_j X_{\epsilon,j}\|_{L_T^1(L^1(K))} \leq C \|X_{\epsilon,j}\|_{L_T^1(L^{q/2})} \operatorname{meas} \left\{ x \in K \mid |\nabla \rho_0(x)| \leq 2^{1-j/2} \right\}^{(q-2)/q}.$$

Using the Sobolev embedding $H^1 \subset L^q$, we get

$$\|X_{\epsilon,j}\|_{L_T^1(L^{q/2})} \leq C \|\nabla \rho_0\|_{L^\infty} \|u_{\epsilon,j}\|_{L_T^2(H^1)}^2 \leq C.$$

Hence we see that $B_j X_{\epsilon,j} = R_{\epsilon,j}$ is a remainder term, thanks to the assumption (11) on ρ_0 .

Next, we look at $(1 - B_j) X_{\epsilon,j}$: following the idea explained here above, we get

$$\begin{aligned} (1 - B_j) u_{\epsilon,j} &= \frac{1 - B_j}{|\nabla \rho_0|^2} \left\{ (u_{\epsilon,j} \cdot \nabla \rho_0) \nabla \rho_0 + (u_{\epsilon,j} \cdot \nabla^\perp \rho_0) \nabla^\perp \rho_0 \right\} \\ (1 - B_j) u_{\epsilon,j}^\perp &= \frac{1 - B_j}{|\nabla \rho_0|^2} \left\{ - (u_{\epsilon,j} \cdot \nabla^\perp \rho_0) \nabla \rho_0 + (u_{\epsilon,j} \cdot \nabla \rho_0) \nabla^\perp \rho_0 \right\}. \end{aligned}$$

Putting this in $(1 - B_j) X_{\epsilon,j}$, we see that the two terms parallel to $\nabla^\perp \rho_0$ cancel out. All that remains is

$$(1 - B_j) X_{\epsilon,j} = \frac{1 - B_j}{|\nabla \rho_0|^2} \left\{ (u_{\epsilon,j} \cdot \nabla \rho_0)^2 + (u_{\epsilon,j} \cdot \nabla^\perp \rho_0)^2 \right\} \nabla \rho_0 = \Gamma_{\epsilon,j}.$$

We have thus proved that $X_{\epsilon,j} = \Gamma_{\epsilon,j} + R_{\epsilon,j}$, therefore

$$\operatorname{div} (\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = \eta_{\epsilon,j} u_{\epsilon,j}^\perp + \Gamma_{\epsilon,j} + R_{\epsilon,j}.$$

Step 6: the new vorticity term. It remains us to deal with the new vorticity term $\eta_{\epsilon,j} u_{\epsilon,j}^\perp$. First we prove that $B_j \eta_{\epsilon,j} u_{\epsilon,j}^\perp$ is a remainder term $R_{\epsilon,j}$. Writing $\eta_{\epsilon,j} = \eta_{\epsilon,j}^{(1)} + \epsilon^\theta \eta_{\epsilon,j}^{(2)}$ as in (37), we see that, for $2 < q < +\infty$, one has

$$\begin{aligned} \|B_j \eta_{\epsilon,j} u_{\epsilon,j}^\perp\|_{L_T^1(L^1(K))} &\leq C \|\eta_{\epsilon,j}^{(1)}\|_{L_T^2(L^2)} \|u_{\epsilon,j}\|_{L_T^2(L^q)} \operatorname{meas} \left\{ x \in K \mid |\nabla \rho_0(x)| \leq 2^{1-j/2} \right\}^{(q-2)/(2q)} \\ &\quad + C \epsilon^\theta \|\eta_{\epsilon,j}^{(2)}\|_{L_T^2(L^2)} \|u_{\epsilon,j}\|_{L_T^2(L^2)}, \end{aligned}$$

which implies the bound

$$\|B_j \eta_{\epsilon,j} u_{\epsilon,j}^\perp\|_{L_T^1(L^1(K))} \leq C \operatorname{meas} \left\{ x \in K \mid |\nabla \rho_0(x)| \leq 2^{1-j/2} \right\}^{(q-2)/(2q)} + C(j) \epsilon^\theta.$$

Therefore, we do in fact see that $B_j \eta_{\epsilon,j} u_{\epsilon,j}^\perp = R_{\epsilon,j}$, as claimed.

Next, we use one last time the decomposition of $u_{\epsilon,j}$ on the basis $\left(\frac{\nabla \rho_0}{|\nabla \rho_0|}, \frac{\nabla^\perp \rho_0}{|\nabla \rho_0|} \right)$ to get

$$\begin{aligned} (1 - B_j) \eta_{\epsilon,j} u_{\epsilon,j}^\perp &= \frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} \left\{ (u_{\epsilon,j}^\perp \cdot \nabla \rho_0) \nabla \rho_0 + (u_{\epsilon,j} \cdot \nabla \rho_0) \nabla^\perp \rho_0 \right\} \\ &= \frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} (u_{\epsilon,j} \cdot \nabla \rho_0) \nabla^\perp \rho_0 + \Gamma_{\epsilon,j}. \end{aligned}$$

At this point, observe that, by taking the divergence of equation (35), we can write $u_{\epsilon,j} \cdot \nabla \rho_0 = \operatorname{div} (\rho_0 u_{\epsilon,j}) = \operatorname{div} V_{\epsilon,j} - \operatorname{div} (\epsilon^\theta \zeta_{\epsilon,j} + h_{\epsilon,j})$, so

$$(1 - B_j) \eta_{\epsilon,j} u_{\epsilon,j}^\perp = \frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} \operatorname{div} V_{\epsilon,j} \nabla^\perp \rho_0 + \Gamma_{\epsilon,j} + R_{\epsilon,j}.$$

Indeed, on the one hand $\epsilon^\theta \operatorname{div} \zeta_{\epsilon,j} \eta_{\epsilon,j} = R_{\epsilon,j}$; on the other, we can estimate

$$\left\| \frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} \operatorname{div} h_{\epsilon,j} \right\|_{L_T^1(L^1)} \leq C \left(\|\eta_{\epsilon,j}^{(1)}\|_{L_T^2(L^2)} \|h_{\epsilon,j}\|_{L_T^2(H^1)} + \epsilon^\theta \|\eta_{\epsilon,j}^{(2)}\|_{L_T^2(L^2)} \|h_{\epsilon,j}\|_{L_T^2(H^1)} \right),$$

so also that term is a remainder, thanks to the uniform bound (36) on $(h_{\epsilon,j})_\epsilon$.

So far, we have thus obtained

$$\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = \frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} \operatorname{div} V_{\epsilon,j} \nabla^\perp \rho_0 + \Gamma_{\epsilon,j} + R_{\epsilon,j}.$$

Step 7: end of the proof. Recalling equation (27), we compute

$$\begin{aligned} \eta_{\epsilon,j} \operatorname{div} V_{\epsilon,j} &= -\epsilon \eta_{\epsilon,j} \partial_t \sigma_{\epsilon,j} = -\frac{1}{2} \epsilon \partial_t (|\sigma_{\epsilon,j}|^2) - \epsilon (\eta_{\epsilon,j} - \sigma_{\epsilon,j}) \partial_t \sigma_{\epsilon,j} \\ &= -\frac{1}{2} \epsilon \partial_t (|\sigma_{\epsilon,j}|^2) - \epsilon \partial_t \left\{ (\eta_{\epsilon,j} - \sigma_{\epsilon,j}) \sigma_{\epsilon,j} \right\} + \epsilon \operatorname{curl} f_{\epsilon,j} \sigma_{\epsilon,j}, \end{aligned}$$

where we have used also equation (28) in passing from the first to the second line. By virtue of the uniform bounds we have already obtained on $(\sigma_{\epsilon,j})_\epsilon$, $(\eta_{\epsilon,j})_\epsilon$ and $(f_{\epsilon,j})_\epsilon$, we see that

$$\frac{1 - B_j}{|\nabla \rho_0|^2} \eta_{\epsilon,j} \operatorname{div} V_{\epsilon,j} \nabla^\perp \rho_0 = R_{\epsilon,j}.$$

In the end, we have shown that

$$\operatorname{div}(\rho_0 u_{\epsilon,j} \otimes u_{\epsilon,j}) = \Gamma_{\epsilon,j} + R_{\epsilon,j},$$

which finally achieves the proof of Proposition 4.4.

4.5 Conclusion: taking the limit

For the quasi-homogeneous case, the results of Sections 3.3, 4.1, 4.2 and 4.3 combine to show convergence of the sequence $(r_\epsilon, u_\epsilon, b_\epsilon)_{\epsilon>0}$ to a solution (r, u, b) of system (10). Notice that we still have to prove the uniqueness part of Theorem 2.2.

In the fully non-homogeneous case, it remains to take care of the singular Coriolis force term: for this, we use the linear convergence properties for the density of Proposition 3.5.

Proposition 4.6. *The Coriolis force term satisfies the following convergence property: for any divergence-free $\phi \in \mathcal{D}(\mathbb{R}_+ \times \Omega; \mathbb{R}^2)$, write $\phi = \nabla^\perp \psi$; then*

$$\frac{1}{\epsilon} \int_0^{+\infty} \int_\Omega \rho_\epsilon u_\epsilon^\perp \cdot \phi \, dx \, dt \xrightarrow{\epsilon \rightarrow 0^+} - \int_0^\infty \int_\Omega \sigma \partial_t \psi \, dx \, dt + \int_\Omega r_0 \psi|_{t=0} \, dx,$$

where σ is the limit density oscillation function identified in Proposition 3.5, and we have improperly written the integral in space for the duality product in $H^{-3-\delta} \times H^{3+\delta}$.

Proof. Let $T > 0$ be a fixed positive time and let $\phi = \nabla^\perp \psi$ be a divergence-free test function, with $\psi \in \mathcal{D}([0, T[\times \Omega)$. We use the decomposition $\rho_\epsilon = \rho_0 + \epsilon \sigma_\epsilon$ to write

$$\begin{aligned} \frac{1}{\epsilon} \int_0^T \int_\Omega \rho_\epsilon u_\epsilon^\perp \cdot \phi \, dx \, dt &= -\frac{1}{\epsilon} \langle \operatorname{div}(\rho_\epsilon u_\epsilon), \psi \rangle_{\mathcal{D}' \times \mathcal{D}} = \frac{1}{\epsilon} \langle \partial_t \rho_\epsilon, \psi \rangle_{\mathcal{D}' \times \mathcal{D}} \\ &= \langle \partial_t \sigma_\epsilon, \psi \rangle_{\mathcal{D}' \times \mathcal{D}} = - \int_0^T \int_\Omega \sigma_\epsilon \partial_t \psi \, dx \, dt - \int_\Omega r_{0,\epsilon} \psi|_{t=0} \, dx. \end{aligned}$$

It is now matter of applying the convergence properties of (6) and Proposition 3.5. \square

This last trick of using ψ as a test function rather than the divergence-free ϕ allows us to get rid of the singularity in the case of non-constant densities. However, it forces us to take the curl of the whole equation. Propositions 3.4 and 4.1 to 4.6 show that, for any divergence-free $\phi \in \mathcal{D}(\mathbb{R}_+ \times \Omega; \mathbb{R}^2)$, which we can write $\phi = \nabla^\perp \psi$ with $\psi \in \mathcal{D}(\mathbb{R}_+ \times \Omega)$, one has

$$\begin{aligned} \int_0^{+\infty} \int_{\Omega} \sigma \partial_t \psi \, dx \, dt + \int_0^{+\infty} \int_{\Omega} \left(\rho_0 u \cdot \partial_t \nabla^\perp \psi - b \otimes b : \nabla \nabla^\perp \psi - \nu(\rho_0) \nabla u : \nabla \nabla^\perp \psi \right) dx \, dt \\ + \langle \Gamma, \operatorname{div}(\rho_0 \nabla^\perp \psi) \rangle_{\mathcal{D}' \times \mathcal{D}} = \int_{\Omega} \left(r_0 \psi|_{t=0} - m_0 \nabla^\perp \psi|_{t=0} \right) dx \, dt. \end{aligned}$$

Integration by parts show that we have indeed the weak form of the sought equation. The proof of Theorem 2.3 is then completed.

5 The quasi-homogeneous case: study of the limit system

In this section, we focus our attention on the limit system for the quasi-homogeneous case, which we recall for the reader's convenience:

$$(38) \quad \begin{cases} \partial_t r + \operatorname{div}(r u) = 0 \\ \partial_t u + \operatorname{div}(u \otimes u) + \nabla \pi + \frac{1}{2} \nabla(|b|^2) + r u^\perp = \nu(1) \Delta u + \operatorname{div}(b \otimes b) \\ \partial_t b + \operatorname{div}(u \otimes b - b \otimes u) = \mu(1) \Delta b \\ \operatorname{div}(u) = \operatorname{div}(b) = 0, \end{cases}$$

for some pressure function π . For simplicity, from now on we set $\nu(1) = \mu(1) = 1$.

We shall prove that the solutions to system (38) are unique in the energy space, given regular enough initial data. In particular, this completes the proof of Theorem 2.2, showing that the whole sequence of solutions $(r_\epsilon, u_\epsilon, b_\epsilon)_\epsilon$ weakly converges to the limit point (r, u, b) , without the need to extract a subsequence. We proceed in four steps. First, we find energy estimates for (38) at the order of regularity suited to prove uniqueness with stability estimates. Then, we show rigorously the existence of solutions at this level of regularity. Finally, we prove uniqueness for system (38). The end result of this section is a well-posedness theorem for the limit system (38).

Theorem 5.1. *Let $0 < \beta < 1$ and let $(r_0, u_0, b_0) \in H^1 \times H^1 \times H^{1+\beta}$ be a set of initial data. There exists a (unique) solution (r, u, b) of (38) related to those initial data such that*

(i) *for all $0 < \gamma < \beta$, we have $r \in C^0(\mathbb{R}_+; H^{1+\gamma})$;*

(ii) *we have $u, b \in L_{\text{loc}}^\infty(H^1) \cap L_{\text{loc}}^2(H^2)$.*

Moreover, such a solutions is unique in the energy space $L_{\text{loc}}^\infty(L^2 \cap L^\infty) \times L_{\text{loc}}^\infty(L^2) \times L_{\text{loc}}^\infty(L^2)$.

Remark 5.2. (i) It goes without saying that the previous regularity properties for the solution (r, u, b) have also a quantitative counterpart, for which we refer to the estimates of Proposition 5.3.

(ii) The uniqueness of solutions is a consequence of a stability estimate and a *weak-strong* uniqueness result, which are stated in Proposition 5.5 below.

In what follows, we will make extensive use of the Gagliardo-Nirenberg inequality (GN inequality for short, see Lemma A.7) as well as the Young inequality in the following form: if $1/p + 1/q = 1$ then, for any $\eta > 0$ and $a, b \geq 0$, we have $ab \leq \eta a^p + \eta^{-q/p} b^q$. From now on, $\eta > 0$ will always note a small positive constant to be fixed in the later parts of the proofs. In addition, we try to find inequalities that are as precise as (reasonably) possible in order to highlight which terms have the most impact on the final estimates.

5.1 Order 2 energy estimates

In this section, we focus on finding order 2 *a priori* estimates for the limit system; one way to do this (see e.g. Section 4.4.1 of [16]) is to use Δu and Δb as test functions in (38). In addition, we attempt to optimise the estimates as far as growth in time is concerned: we show that the higher order energy grows slower than any polynomial function of $T \geq 0$ of positive degree $h > 0$.

Proposition 5.3. *Let (r, u, b) be a regular enough solution of (38) related to the (regular) initial datum (r_0, u_0, b_0) . Then, we have the following properties:*

- (i) *we have $u, b \in L^\infty(L^2)$ and $\nabla u, \nabla b \in L^2(L^2)$, with the standard energy estimate (41) below;*
- (ii) *for any fixed $T > 0$, we have $\nabla u, \nabla b \in L_T^\infty(L^2)$ and $\Delta u, \Delta b, \partial_t u, \partial_t b \in L_T^2(L^2)$, with explicit bounds: for all $h > 0$, there is a constant $C = C(\|r_0\|_{L^2 \cap L^\infty}, \|(u_0, b_0)\|_{H^1}, h) > 0$ such that*

$$(39) \quad \|\nabla u\|_{L_T^\infty(L^2)} + \|(\Delta u, \partial_t u)\|_{L_T^2(L^2)} + \|\nabla b\|_{L_T^\infty(L^2)} + \|(\Delta b, \partial_t b)\|_{L_T^2(L^2)} \leq C(1 + T^h);$$

- (iii) *for all $2 \leq p \leq +\infty$ and all $t \geq 0$, $\|r(t)\|_{L^p} = \|r_0\|_{L^p}$, and for all $0 < \gamma < \beta < 1$ and any fixed $T > 0$, we also have*

$$(40) \quad \|r\|_{L_T^\infty(H^{1+\gamma})} \leq C(\beta, \gamma) \exp \left\{ C(\beta, \gamma) \left(\int_0^T \|\nabla u\|_{H^1} dt \right)^2 \right\} \|r_0\|_{H^{1+\beta}}.$$

Proof. First, testing the momentum equation with u and the magnetic field equation with b gives a basic energy estimate similar to (7): namely,

$$(41) \quad \frac{1}{2} \left(\|u(t)\|^2 + \|b(t)\|^2 \right) + \int_0^t \left(\|\nabla u(s)\|^2 + \|\nabla b(s)\|^2 \right) ds \leq \frac{1}{2} \left(\|u_0\|^2 + \|b_0\|^2 \right).$$

Next, we use the fact that r solves a pure transport equation with a divergence-free flow u to see that the L^p norms of $r(t)$ are preserved: $\|r(t)\|_{L^p} = \|r_0\|_{L^p}$ for all $p \in [2, +\infty]$ and all $t \geq 0$.

Now, we consider order 2 energy estimates. We test the momentum equation with $-\Delta u$ and the magnetic field equation with $-\Delta b$; summing and integrating by parts gives

$$(42) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|\nabla u\|^2 + \|\nabla b\|^2 \right) + \left(\|\Delta u\|^2 + \|\Delta b\|^2 \right) + \int_\Omega (b \cdot \nabla) b \cdot \Delta u \, dx + \int_\Omega (b \cdot \nabla) u \cdot \Delta b \, dx \\ = \int_\Omega (u \cdot \nabla) u \cdot \Delta u \, dx + \int_\Omega r u^\perp \Delta u \, dx + \int_\Omega (u \cdot \nabla) b \cdot \Delta b \, dx. \end{aligned}$$

We start by handling the two integrals which do not involve the magnetic field. On the one hand, using Hölder's inequality and then Proposition A.8 yields

$$\left| \int_\Omega (u \cdot \nabla) u \cdot \Delta u \, dx \right| \leq \|\Delta u\|_{L^2} \|u\|_{L^\infty} \|\nabla u\|_{L^2} \leq C \|\Delta u\|_{L^2}^{3/2} \|u\|_{L^2}^{1/2} \|\nabla u\|_{L^2}.$$

Young's inequality with exponents $\frac{3}{4} + \frac{1}{4} = 1$ gives in turn

$$\left| \int_\Omega (u \cdot \nabla) u \cdot \Delta u \, dx \right| \leq \eta \|\Delta u\|_{L^2}^2 + C(\eta) \|u\|_{L^2}^2 \|\nabla u\|_{L^2}^4 = \eta \|\Delta u\|_{L^2}^2 + M_1(t) \|\nabla u\|_{L^2}^2,$$

where $M_1(t) := C(\eta) \|u(t)\|_{L^2}^2 \|\nabla u(t)\|_{L^2}^2$. Remark that, by (41), $M_1 \in L^1(\mathbb{R}_+)$ is a globally integrable function, with $\|M_1\|_{L^1(\mathbb{R}_+)}$ depending only on η , $\|u_0\|_{L^2}$ and $\|b_0\|_{L^2}$.

On the other hand, we use Hölder's inequality with exponents $\frac{1}{p} + \frac{1}{q} + \frac{1}{2} = 1$ which we will choose later, followed by the GN inequality: for all $\eta > 0$,

$$\left| \int_\Omega r u^\perp \cdot \Delta u \, dx \right| \leq \|r\|_{L^p} \|u\|_{L^q} \|\Delta u\|_{L^2} \leq C(q) \|r\|_{L^2}^{2/p} \|r\|_{L^\infty}^{1-2/p} \|u\|_{L^2}^{2/q} \|\nabla u\|_{L^2}^{1-2/q} \|\Delta u\|_{L^2}$$

$$\leq \eta \|\Delta u\|_{L^2}^2 + C(\eta, \|r_0\|_{L^2 \cap L^\infty}, q) \|u\|_{L^2}^{4/q} \|\nabla u\|_{L^2}^{2(1-2/q)}.$$

Since $\|u\|_{L^2} \in L^\infty(\mathbb{R}_+)$ and $\|\nabla u\|_{L^2}^2 \in L^1(\mathbb{R}_+)$ by the energy inequality (41), we see that, for any arbitrary $h > 0$, we can chose q so large that the function $M_{1+h}(t) := C\|u(t)\|_{L^2}^{4/q} \|\nabla u(t)\|_{L^2}^{2(1-2/q)}$ belongs to the space $L^{1+h}(\mathbb{R}_+)$. Therefore

$$(43) \quad \left| \int_{\Omega} r u^\perp \cdot \Delta u \, dx \right| \leq \eta \|\Delta u\|_{L^2}^2 + M_{1+h}(t),$$

with the norm $\|M_{1+h}\|_{L^{1+h}(\mathbb{R}_+)}$ only depending on the quantities $(\eta, \|r_0\|_{L^2 \cap L^\infty}, \|(u_0, b_0)\|_{L^2}, h)$.

Now we take care of the three remaining integrals in (42) involving the magnetic field. As with the first integral, Proposition A.8 yields

$$\begin{aligned} \left| \int_{\Omega} (b \cdot \nabla) b \cdot u \, dx \right| &\leq \|\Delta u\|_{L^2} \|b\|_{L^\infty} \|\nabla b\|_{L^2} \leq C \|\Delta u\|_{L^2} \|\Delta b\|_{L^2}^{1/2} \|b\|_{L^2}^{1/2} \|\nabla b\|_{L^2} \\ &\leq \eta \|\Delta u\|_{L^2}^2 + C(\eta) \|\Delta b\|_{L^2} \|b\|_{L^2} \|\nabla b\|_{L^2}^2 \\ &\leq \eta \left\{ \|\Delta u\|_{L^2}^2 + \|\Delta b\|_{L^2}^2 \right\} + M_1(t) \|\nabla b\|_{L^2}^2, \end{aligned}$$

where we have set $M_1(t) := C(\eta) \|b(t)\|_{L^2}^2 \|\nabla b(t)\|_{L^2}^2$. Notice that $M_1 \in L^1(\mathbb{R}_+)$, with $\|M_1\|_{L^1(\mathbb{R}_+)}$ depending only on η , $\|u_0\|_{L^2}$ and $\|b_0\|_{L^2}$.

Next, similarly as above, we use Hölder's inequality, the GN inequality and then Young's inequality twice, to obtain

$$\begin{aligned} \left| \int_{\Omega} (b \cdot \nabla) u \cdot \Delta b \, dx \right| &\leq \|b\|_{L^4} \|\nabla u\|_{L^4} \|\Delta b\|_{L^2} \leq C \|b\|_{L^2}^{1/2} \|\nabla b\|_{L^2}^{1/2} \|\nabla u\|_{L^2}^{1/2} \|\Delta u\|_{L^2}^{1/2} \|\Delta b\|_{L^2} \\ &\leq \eta \|\Delta b\|_{L^2}^2 + C(\eta) \|b\|_{L^2} \|\nabla b\|_{L^2} \|\nabla u\|_{L^2} \|\Delta u\|_{L^2} \\ &\leq \eta \left\{ \|\Delta b\|_{L^2}^2 + \|\Delta u\|_{L^2}^2 \right\} + M_1(t) \|\nabla b\|_{L^2}^2, \end{aligned}$$

where this time $M_1 = C(\eta) \|b\|_{L^2}^2 \|\nabla u\|_{L^2}^2 \in L^1(\mathbb{R}_+)$, with $\|M_1\|_{L^1(\mathbb{R}_+)} = C(\eta, \|u_0\|_{L^2}, \|b_0\|_{L^2})$. Exactly the same computations *mutatis mutandi* yield

$$\left| \int_{\Omega} (u \cdot \nabla) b \cdot \Delta b \, dx \right| \leq \eta \left\{ \|\Delta b\|_{L^2}^2 + \|\Delta u\|_{L^2}^2 \right\} + M_1(t) \|\nabla u\|_{L^2}^2,$$

with $M_1 = C(\eta) \|u\|_{L^2}^2 \|\nabla b\|_{L^2}^2 \in L^1(\mathbb{R}_+)$ and $\|M_1\|_{L^1(\mathbb{R}_+)} = C(\eta, \|u_0\|_{L^2}, \|b_0\|_{L^2})$.

In the end, putting all these estimates together and choosing $\eta > 0$ small enough, we obtain the differential inequality

$$\frac{1}{2} \frac{d}{dt} (\|\nabla u\|^2 + \|\nabla b\|^2) + \frac{1}{2} (\|\Delta u\|^2 + \|\Delta b\|^2) \leq M_{1+h}(t) + M_1(t) (\|\nabla u\|^2 + \|\nabla b\|^2).$$

An application of Grönwall's lemma finally yields, for all fixed $T > 0$, the inequality

$$(44) \quad \sup_{t \in [0, T]} (\|\nabla u(t)\|^2 + \|\nabla b(t)\|^2) + \int_0^T (\|\Delta u(t)\|^2 + \|\Delta b(t)\|^2) \, dt \\ \leq 2 \left(\|\nabla u_0\|_{L^2}^2 + \|\nabla b_0\|_{L^2}^2 + \int_0^T M_{1+h}(t) \, dt \right) \exp \left\{ \int_0^{+\infty} M_1(t) \, dt \right\}.$$

Observe that

$$\int_0^T M_{1+h}(t) \, dt = \|M_{1+h}\|_{L^{1+h}(\mathbb{R}_+)} T^{h/(1+h)} \leq C (1 + T^h),$$

which obviously implies estimate (39), up to the time derivative terms, which will be dealt with in a while. For the moment, remark that, since we have shown $L_T^2(H^2)$ bounds on the velocity field, we can apply Proposition 5.2. of [8] (which is also expressed in a much more thorough form in Theorem 3.33 of [4]) to the transport equation for r : for all $0 < \gamma < \beta < 1$, we get propagation of the $L_T^\infty(H^{1+\gamma})$ regularity of r , together with estimate (40).

Let us come back to the proof of (39): it remains us to bound the time derivative terms. We start by looking at $\partial_t u$. In order to get rid of the pressure term, we apply the Leray projector \mathbb{P} , which is the L^2 -orthogonal projector on the subspace of divergence-free functions. It can be defined as a Fourier multiplier:

$$\forall f \in L^2, \quad \widehat{\mathbb{P}f}(\xi) = \widehat{f}(\xi) - \left(\frac{\xi}{|\xi|^2} \cdot \widehat{f}(\xi) \right) \xi = \left(\frac{\xi^\perp}{|\xi|^2} \cdot \widehat{f}(\xi) \right) \xi^\perp.$$

In particular, it commutes with differential operators (both on time and space variables). Applying the Leray projector to the momentum equation in (38) gives

$$(45) \quad \partial_t u + \mathbb{P}[\operatorname{div}(u \otimes u - b \otimes b) + ru^\perp] = \Delta u,$$

since $\operatorname{div} u = 0$. In fact, since \mathbb{P} is a Fourier multiplier associated to a homogeneous function of degree zero, it is a continuous operator over L^2 ; so we need only estimates on Δu , ru and $\operatorname{div}(u \otimes u - b \otimes b)$ to conclude (39).

First of all, for $1/p + 1/q = 1/2$, Hölder's and the GN inequalities, combined with (41), give

$$\|ru\|_{L^2} \leq \|r\|_{L^p} \|u\|_{L^q} \leq C(q, \|r_0\|_{L^2 \cap L^\infty}, \|u_0\|_{L^2}, \|b_0\|_{L^2}) \|\nabla u\|_{L^2}^{1-2/q} = C M_{1+h}^{1/2},$$

where M_{1+h} has been defined in (43). Hence, by taking q large enough, we see that $ru \in L^{2(1+h)}(L^2)$. Secondly, using that $\operatorname{div}(u \otimes u) = (u \cdot \nabla)u$, an application of Proposition A.8 gives

$$(46) \quad \|\operatorname{div}(u \otimes u)\|_{L^2}^2 \leq C \|u\|_{L^\infty}^2 \|\nabla u\|_{L^2}^2 \leq C \|u\|_{L^2} \|\Delta u\|_{L^2} \|\nabla u\|_{L^2}^2.$$

Integrating over $t \in [0, T]$ and using estimate (41), after an application of Young inequality we obtain, for all $h > 0$, the bound

$$\begin{aligned} \|\operatorname{div}(u \otimes u)\|_{L_T^2(L^2)}^2 &\leq C \int_0^T (\|u\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + \|\Delta u\|_{L^2}^2) dt \\ &\leq C(\|(u_0, b_0)\|_{L^2}) \left(\sup_{[0, T]} (\|\nabla u\|_{L^2}^2) + \|\Delta u\|_{L_T^2(L^2)}^2 \right). \end{aligned}$$

In view of (44), this term can be finally bounded by the right-hand side of that inequality. The same computations with the magnetic field yield

$$\|\operatorname{div}(b \otimes b)\|_{L_T^2(L^2)}^2 \leq C(\|(u_0, b_0)\|_{L^2}) \left(\sup_{[0, T]} (\|\nabla b\|_{L^2}^2) + \|\Delta b\|_{L_T^2(L^2)}^2 \right),$$

which again can be bounded by the right-hand side of (41). In the end, the combination of all those estimates implies that also $\|\partial_t u\|_{L_T^2(L^2)}$ satisfies (39).

It only remains to find the estimate on $\|\partial_t b\|_{L_T^2(L^2)}$, for which we use the magnetic field equation, namely the third equation in (38). Estimate (44) already provides us with a bound for Δb in the sought space. On the other hand, the quadratic terms can be estimated exactly as the corresponding ones appearing in the momentum equation. We finally deduce that also $\|\partial_t b\|_{L_T^2(L^2)}$ fulfills (39), and this completes the proof of the proposition. \square

5.2 Existence result

In this section, we explain how solutions of (38), whether they be weak solution in the energy space or with the level of regularity described in Proposition 5.3, can be constructed.

Proposition 5.4. *Assume that $r_0 \in L^2 \cap L^\infty$ and that $u_0, b_0 \in L^2$ are two divergence-free functions. Then there exists a weak solution (r, u, b) of system (38) related to those initial data, such that $r \in L^\infty(L^2 \cap L^\infty)$ and³ $u, b \in C_b^0(L^2)$, with ∇u and ∇b belonging to $L^2(L^2)$. Moreover this solution satisfies the basic energy inequality (41) and the conservation of the L^p norms of r in time, for any $p \in [2, +\infty]$.*

Suppose now that $r_0 \in H^{1+\beta}$, for some $\beta > 0$, and that $u_0, b_0 \in H^1$. Then there exists a weak solution (r, u, b) of (38) such that, in addition to the previous properties, one also has, for all $T > 0$, $r \in C_T^0(H^{1+\gamma})$ for all $0 < \gamma < \min\{1, \beta\}$, and $u, b \in C_T^0(H^1) \cap L_T^2(H^2)$. Moreover, this solution satisfies inequalities (39) and (40).

The proof of the previous proposition is standard (see e.g. [6] or [9]); we give here most of the details for reader's convenience. First of all, we implement Friedrichs scheme and construct smooth solutions to an approximate system. After deriving uniform bounds (of first and second order), we show that those solutions tend (weakly) to a solution of (38).

Step 1: approximate system. Let $j \geq 2$ and A_j be the spectral projection operator defined in the following way:

$$\forall f \in L^2, \quad \mathcal{F}[A_j f](\xi) = \mathbf{1}_{|\xi| \leq j} \widehat{f}(\xi).$$

Recall the Leray projector \mathbb{P} from (45), and that we have set $\mu(1) = \nu(1) = 1$. Set $r_1(t, x) = S_1 r_0(x)$, where S_1 is the low frequency cut-off operator defined in (56). For $j \geq 2$, we consider the sequence of approximate systems

$$(47) \quad \begin{cases} \partial_t u_j + \mathbb{P} A_j \operatorname{div} (u_j \otimes u_j - b_j \otimes b_j) + \mathbb{P} A_j [r_{j-1} u_j^\perp] = \Delta u_j \\ \partial_t b_j + A_j \operatorname{div} (u_j \otimes b_j - b_j \otimes u_j) = \Delta b_j \\ \operatorname{div}(u_j) = \operatorname{div}(b_j) = 0, \end{cases}$$

which we equip with the initial datum $(u_j, b_j)|_{t=0} = (A_j u_0, A_j b_0)$. Given r_{j-1} , applying the Cauchy-Lipschitz theorem in the Banach space

$$\mathcal{X}_j = \left\{ f \in L^2 \mid \operatorname{Supp} \widehat{f} \subset B(0, j) \right\}$$

gives the existence of a unique $C^\infty([T_-(j), T_+(j)]; \mathcal{X}_j)$ maximal solution. Then, we compute the function r_j by solving the linear transport equation

$$\begin{cases} \partial_t r_j + \operatorname{div} (r_j u_j) = 0 \\ (r_j)|_{t=0} = S_j r_0. \end{cases}$$

We now look for uniform bounds for $(r_j, u_j, b_j)_j$ in suitable spaces. First of all, using the fact that r_j solves a pure transport equation by a divergence-free velocity field u_j , one gathers that

$$\|r_j(t)\|_{L^2} + \|r_j(t)\|_{L^\infty} = \|S_j r_0\|_{L^2} + \|S_j r_0\|_{L^\infty} \leq C \|r_0\|_{L^2 \cap L^\infty}.$$

for all $t \in [0, T_+(j)[$. This implies that $(r_j)_j \subset L^\infty(L^2 \cap L^\infty)$. Next, we easily see that the maximal solution (u_j, b_j) satisfies the basic energy estimate (41); it also satisfies the order 2

³Given a Banach space X , we note $C_b^0(X)$ the space of continuous and globally bounded functions on \mathbb{R}_+ with values in X .

estimates (39). Indeed, testing the momentum equation in (47) with $-\Delta A_j u$ (for example), which is both in \mathcal{X}_j and divergence-free, gives, for all $0 \leq t < T_+(j)$, the equality

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla u_j|^2 dx + \int_{\Omega} |\Delta u_j| dx = \int_{\Omega} \operatorname{div} (u_j \otimes u_j - b_j \otimes b_j) \cdot \Delta u_j dx + \int_{\Omega} r_{j-1} u_j^\perp \cdot \Delta u_j dx.$$

An analogous relation holds for the magnetic field b_j . Then, one can repeat the same computations as in Subsection 5.1 to finally get (39), as claimed.

We now show that the approximate solutions do not blow-up in finite time. For this, fix $j \geq 2$: the basic energy estimates state that $\|u_j(t)\|_{L^2}$ and $\|b_j(t)\|_{L^2}$ are bounded for $0 \leq t < T_+(j)$, therefore (using the bounds on $(r_j)_j$ and the spectral localisation) so are the norms of the time derivatives $\|\partial_t u_j(t)\|_{L^2}$ and $\|\partial_t b_j(t)\|_{L^2}$. Hence, the solution of the ODE system (47) satisfies the Cauchy criterion for $t < T_+(j)$, and necessarily $T_+(j) = +\infty$.

Step 2: convergence. For the sake of generality, we prove convergence of the sequence of approximate solutions relying only on the basic energy estimates (41) and the conservation of L^p norms for the r_j . Specifically, we will be using only the following uniform bounds:

$$(u_j)_{j \geq 2}, (b_j)_{j \geq 2} \subset L^\infty(L^2) \cap L_{\text{loc}}^2(H^1) \quad \text{and} \quad (r_j)_{j \geq 2} \subset L^\infty(L^2 \cap L^\infty).$$

Those bounds yield the following weak convergence properties (up to an extraction): there exists a triplet $(r, u, b) \in L^\infty(L^2 \cap L^\infty) \times L_{\text{loc}}^2(H^1) \times L_{\text{loc}}^2(H^1)$ such that, for all fixed $T > 0$, one has

$$(48) \quad (u_j, b_j) \rightharpoonup (u, b) \quad \text{in } L_T^2(H^1) \quad \text{and} \quad r_j \xrightarrow{*} r \quad \text{in } L^\infty(L^2 \cap L^\infty).$$

In order to achieve convergence in the non-linear terms, we are going to prove that both $(\partial_t u_j)_j$ and $(\partial_t b_j)_j$ are uniformly bounded in $L_{\text{loc}}^2(H^{-1})$. Notice that both \mathbb{P} and A_j are continuous for all the H^s topologies (with $s \in \mathbb{R}$). Therefore, by using (47), we get

$$\begin{aligned} \|\partial_t u_j\|_{L_T^2(H^{-1})} + \|\partial_t b_j\|_{L_T^2(H^{-1})} &\leq \|u_j\|_{L_T^2(H^1)} + \|b_j(t)\|_{L_T^2(H^1)} \\ &\quad + \sum_{f, g \in \{u_j, b_j\}} \left\| \operatorname{div} (f \otimes g) \right\|_{L_T^2(H^{-1})} + \|r_{j-1} u_j\|_{L_T^2(H^{-1})}. \end{aligned}$$

The last term is bounded by $\|r_{j-1} u_j\|_{L_T^2(H^{-1})} \leq C_T \|r_{j-1}\|_{L_T^\infty(L^\infty)} \|u_j\|_{L_T^\infty(L^2)}$, so we only have to worry about the quadratic terms. If $f, g \in L_T^2(H^1)$, then, for all $0 \leq t \leq T$, using the Sobolev embedding $H^1 \subset L^4$ followed by the Cauchy-Schwarz inequality in L_T^2 , we get

$$\left\| \operatorname{div} (f \otimes g) \right\|_{L_T^2(H^{-1})}^2 \leq \int_0^T \|f(t)\|_{L^4}^2 \|g(t)\|_{L^4}^2 dt \leq \|f\|_{L_T^2(H^1)}^2 \|g\|_{L_T^2(H^1)}^2 < +\infty.$$

These computations show that $(\partial_t u_j)_j$ and $(\partial_t b_j)_j$ are indeed bounded in $L_T^2(H^{-1})$, from which we infer the uniform boundness of $(u_j)_j$ and $(b_j)_j$ in $C_T^{0,1/2}(H^{-1})$. On the other hand, thanks to the compact embedding $L^2(K) \subset H^{-1}(K)$ for all compact $K \subset \Omega$, we deduce that, for almost all $0 \leq t \leq T$, the sequences $(u_j(t))_j$ and $(b_j(t))_j$ are relatively compact in H_{loc}^{-1} . An application of the Ascoli-Arzelà theorem gives then, for all $T > 0$, the strong convergence

$$(u_j, b_j) \longrightarrow (u, b) \quad \text{in } L_T^\infty(H_{\text{loc}}^{-1}).$$

Because $(u_j)_j$ and $(b_j)_j$ are also bounded in $L_T^2(H^1)$, interpolation between Sobolev spaces gives strong L^2 convergence

$$(u_j, b_j) \longrightarrow (u, b) \quad \text{in } L_T^2(L_{\text{loc}}^2).$$

Next, using the fact that the r_j solve a linear transport equation and arguing exactly as in Subsection 3.2, we get strong convergence

$$r_j \longrightarrow r \quad \text{in} \quad L_T^2(L_{\text{loc}}^2),$$

which in turn gives convergence of the products $(r_j u_{j-1})_j$ and $(r_j u_j)_j$ in the sense of distributions in $\mathbb{R}_+ \times \Omega$.

Step 3: weak solutions. We aim to prove that the triplet (r, u, b) , identified in (48), is in fact a weak solution of (38). The only terms whose convergence is not completely obvious at this point are the quadratic terms in u_j and b_j . Let $\phi \in \mathcal{D}([0, T[\times \Omega; \mathbb{R}^2)$ be a divergence-free test function. We will only prove the convergence of

$$\int_0^T \int_{\Omega} A_j \mathbb{P}(u_j \otimes b_j) : \nabla \phi \, dx \, dt = \int_0^T \int_{\Omega} (u_j \otimes b_j) : A_j \nabla \phi \, dx \, dt,$$

all other quadratic terms being similar. Taking the difference between the previous integral and the one we desire, we get

$$\begin{aligned} & \left| \int_0^T \int_{\Omega} \left\{ (u_j \otimes b_j) : A_j \nabla \phi - (u \otimes b) : \nabla \phi \right\} dx \, dt \right| \\ & \leq \left| \int_0^T \int_{\Omega} (u_j \otimes b_j) : (A_j - I) \nabla \phi \, dx \, dt \right| + \left| \int_0^T \int_{\Omega} (u_j \otimes b_j - u \otimes b) : \nabla \phi \, dx \, dt \right|. \end{aligned}$$

Using the Sobolev embedding $H^1 \subset L^4$, we see that the first integral on the right-hand side is bounded by

$$\left| \int_0^T \int_{\Omega} (u_j \otimes b_j) : (A_j - I) \nabla \phi \, dx \, dt \right| \leq \|u_j\|_{L_T^2(L^4)} \|b_j\|_{L_T^2(L^4)} \|(A_j - I) \nabla \phi\|_{L_T^\infty(L^2)},$$

which converges to 0 for $j \rightarrow +\infty$. This comes from the uniform bound (with respect to time)

$$\|(A_j - I) \nabla \phi(t)\|_{L^2}^2 \leq \frac{1}{j^2} \|\nabla^2 \phi(t)\|_{L^2}^2 \leq \frac{C}{j^2}.$$

For the other integral, we simply recall that $(u_j, b_j) \rightarrow (u, b)$ in $L_T^2(L_{\text{loc}}^2)$, so that we have strong convergence of the tensor products

$$(49) \quad u_j \otimes b_j \longrightarrow u \otimes b \quad \text{in} \quad L_T^1(L_{\text{loc}}^1).$$

Thus, we have proved convergence for the quadratic term:

$$\int_0^T \int_{\Omega} A_j \mathbb{P}(u_j \otimes b_j) : \nabla \phi \, dx \, dt \xrightarrow{j \rightarrow +\infty} \int_0^T \int_{\Omega} (u \otimes b) : \nabla \phi \, dx \, dt.$$

In the end, we have shown that (r, u, b) is indeed a weak solution of (38). Finally, the Banach-Steinhaus theorem makes sure that the inequalities of Proposition 5.3 are carried from the approximate solutions to the limit point (r, u, b) . This completes the proof of Proposition 5.4.

5.3 Uniqueness for the limit system

The proof of the uniqueness for system (38) is based on stability estimates in the energy space. Hence, we require enough regularity to perform those energy estimates, without any attempt of sharpness in our statement (very likely, a well-posedness result like the one in [29] may be proved also for our system).

Proposition 5.5. Fix $0 < \beta < 1$. Let $(r_{0,1}, u_{0,1}, b_{0,1}) \in H^{1+\beta} \times H^1 \times H^1$ and $(r_{0,2}, u_{0,2}, b_{0,2}) \in (L^2 \cap L^\infty) \times L^2 \times L^2$ be two sets of initial data, and for $i = 1, 2$, consider (r_i, u_i, b_i) two associated solutions of (38) such that:

- (1) $r_i \in L^\infty(L^2 \cap L^\infty)$ and $u_i, b_i \in L^\infty(L^2)$, with $\nabla u_i, \nabla b_i \in L^2(L^2)$;
- (2) for all $T > 0$, $u_1, b_1 \in L_T^\infty(H^1) \cap L_T^2(H^2) \cap W_T^{1,2}(L^2)$ and $r_1 \in L_T^\infty(H^{1+\gamma})$ for all $0 < \gamma < \beta$.

Define $\delta r = r_2 - r_1$, $\delta u = u_2 - u_1$ and $\delta b = b_2 - b_1$, and note by $\delta r_0, \delta u_0, \delta b_0$ the same quantities computed on the initial data.

Then, we have the following stability estimate: for every $0 \leq t \leq T$,

$$(50) \quad \|\delta u(t)\|_{L^2}^2 + \|\delta b(t)\|_{L^2}^2 + \|\delta r(t)\|_{L^2}^2 + \int_0^t \left\{ \|\nabla \delta u\|_{L^2}^2 + \|\nabla \delta b\|_{L^2}^2 \right\} ds \\ \leq C \left(\|\delta u_0\|_{L^2}^2 + \|\delta b_0\|_{L^2}^2 + \|\delta r_0\|_{L^2}^2 \right),$$

where the constant $C > 0$ depends on T , $\mu(1)$, $\nu(1)$ and $(\|r_{0,1}\|_{H^{1+\beta}}, \|u_{0,1}\|_{H^1}, \|b_{0,1}\|_{H^1})$.

Our statement is pretty much a *weak-strong uniqueness* result. In particular, we deduce uniqueness of the solutions in the energy space, given regular initial data.

Corollary 5.6. Consider $0 < \beta < 1$ and $(r_0, u_0, b_0) \in H^{1+\beta} \times H^1 \times H^1$. For that initial datum, there is exactly one solution (r, u, b) of (38) in the energy space, that is such that $r \in L^\infty(L^2 \cap L^\infty)$ and $u, b \in L^\infty(L^2)$, with $\nabla u, \nabla b \in L^2(H^1)$.

Let us now prove the previous proposition.

Proof of Proposition 5.5. We start by remarking that the existence of the two sets of solutions $(r_i, u_i, b_i)_{i=1,2}$, with the claimed level of regularity, is a consequence of Proposition 5.4.

In order to prove inequality (50), we take the difference between the equation solved by (r_2, u_2, b_2) and the one solved by (r_1, u_1, b_1) : this gives

$$\begin{cases} \partial_t \delta u + (u_2 \cdot \nabla) \delta u + (\delta u \cdot \nabla) u_1 + \nabla \Pi + r_2 \delta u^\perp + \delta r u_1^\perp = \Delta \delta u + (\delta b \cdot \nabla) b_1 + (b_2 \cdot \nabla) \delta b \\ \partial_t \delta b + (\delta u \cdot \nabla) b_1 + (u_2 \cdot \nabla) \delta b = (\delta b \cdot \nabla) u_1 + (b_2 \cdot \nabla) \delta u + \Delta \delta b \\ \partial_t \delta r + (u_2 \cdot \nabla) \delta r = -\delta u \cdot \nabla r_1 \\ \operatorname{div}(\delta u) = \operatorname{div}(\delta b) = 0, \end{cases}$$

where we have set $\Pi := \pi_2 - \pi_1 + \frac{1}{2}|b_2|^2 - \frac{1}{2}|b_1|^2$ in the first equation. Recall that we are taking $\nu(1) = \mu(1) = 1$ for simplicity of presentation.

Omitting (for the sake of brevity) a standard regularisation process, let us perform energy estimates directly on the previous system. So, test the first equation with δu , the second one with δb and the third one with δr : one gets

$$(51) \quad \frac{1}{2} \frac{d}{dt} \int_\Omega |\delta u|^2 dx + \int_\Omega (u_2 \cdot \nabla) \delta u \cdot \delta u dx + \int_\Omega (\delta u \cdot \nabla) u_1 \cdot \delta u dx + \int_\Omega \delta r u_1^\perp \cdot \delta u dx \\ + \int_\Omega |\nabla \delta u|^2 dx = \int_\Omega (\delta b \cdot \nabla) b_1 \cdot \delta u dx + \int_\Omega (b_2 \cdot \nabla) \delta b \cdot \delta u dx$$

$$(52) \quad \frac{1}{2} \frac{d}{dt} \int_\Omega |\delta b|^2 dx + \int_\Omega (\delta u \cdot \nabla) b_1 \cdot \delta b dx + \int_\Omega (u_2 \cdot \nabla) \delta b \cdot \delta b dx + \int_\Omega |\nabla \delta b|^2 dx \\ = \int_\Omega (\delta b \cdot \nabla) u_1 \cdot \delta b dx + \int_\Omega (b_2 \cdot \nabla) \delta u \cdot \delta b dx$$

$$(53) \quad \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\delta r|^2 dx + \int_{\Omega} (u_2 \cdot \nabla \delta r) \delta r dx = - \int_{\Omega} (\delta u \cdot \nabla r_1) \delta r dx.$$

An integration by parts shows that the second integral in (51), the third in (52) and the second in (53) are all equal to zero. Next, note that the last integrals in (51) and (52) are opposite. Therefore, by adding the three equations together, we gather

$$(54) \quad \frac{1}{2} \frac{d}{dt} \int_{\Omega} \left\{ |\delta u|^2 + |\delta b|^2 + |\delta r|^2 \right\} dx + \int_{\Omega} \left\{ |\nabla \delta u|^2 + |\nabla \delta b|^2 \right\} dx \\ \leq \left| \int_{\Omega} (\delta u \cdot \nabla) u_1 \cdot \delta u dx \right| + \left| \int_{\Omega} \delta r u_1^\perp \cdot \delta u dx \right| + \left| \int_{\Omega} (\delta u \cdot \nabla r_1) \delta r dx \right| \\ + \left| \int_{\Omega} (\delta u \cdot \nabla) b_1 \cdot \delta b dx \right| + \left| \int_{\Omega} (\delta b \cdot \nabla) u_1 \cdot \delta b dx \right| + \left| \int_{\Omega} (\delta b \cdot \nabla) b_1 \cdot \delta u dx \right|.$$

The first three integrals, which do not involve the magnetic field, can be dealt with as in [16] (see Paragraph 4.4.2 therein). We briefly summarise the computations. Firstly, using in turn the Hölder, the GN and Young's inequalities with exponents $1/4 + 3/4 = 1$, we infer

$$\left| \int_{\Omega} (\delta u \cdot \nabla) u_1 \cdot \delta u dx \right| \leq \|\delta u\|_{L^4} \|u_1\|_{L^4} \|\nabla \delta u\|_{L^2} \leq C \|\delta u\|_{L^2}^{1/2} \|\nabla \delta u\|_{L^2}^{3/2} \|u_1\|_{L^4} \\ \leq \eta \|\nabla \delta u\|_{L^2}^2 + C(\eta) \|u_1\|_{L^4}^4 \|\delta u\|_{L^2}^2 = \eta \|\nabla \delta u\|_{L^2}^2 + C(\eta) M_1 \|\delta u\|_{L^2}^2.$$

Notice that the GN inequality again gives that $M_1(t) = \|u_1(t)\|_{L^4}^4 \leq C \|u_1(t)\|_{L^2}^2 \|\nabla u_1(t)\|_{L^2}^2$, hence $M_1 \in L^1(\mathbb{R}_+)$ is an integrable function.

Next, making use of Proposition A.8, we have

$$\left| \int_{\Omega} \delta r u_1^\perp \cdot \delta u dx \right| \leq \|u_1\|_{L^\infty} (\|\delta u\|_{L^2}^2 + \|\delta r\|_{L^2}^2) = N_4 (\|\delta u\|_{L^2}^2 + \|\delta r\|_{L^2}^2),$$

with $N_4(t) = \|u_1(t)\|_{L^\infty} \leq C \|u_1(t)\|_{L^2}^{1/2} \|\Delta u_1(t)\|_{L^2}^{1/2} \in L_T^4$ for any fixed $T > 0$.

As for the third integral, we use the fact that $\nabla r_1 \in L_T^\infty(H^\gamma)$ for some $\gamma > 0$. By fractional Sobolev embedding (see equation (58) in the appendix), we know that $\nabla r_1 \in L_T^\infty(L^p)$ for some $p > 2$. Let q be an exponent such that $1/p + 1/q = \frac{1}{2}$. Applying the GN inequality first, and then Young's inequality with exponents $1/(1 - 2/p) + 1/(2/p) = 1$ gives

$$\left| \int_{\Omega} \delta u \cdot \nabla r_1 \delta r dx \right| \leq \|\nabla r_1 \cdot \delta u\|_{L^2}^2 + \|\delta r\|_{L^2}^2 \leq \|\nabla r_1\|_{L^p}^2 \|\delta u\|_{L^q}^2 + \|\delta r\|_{L^2}^2 \\ \leq C(p, \|r_{0,1}\|_{H^{1+\beta}}, \|u_{0,1}\|_{H^1}, \|b_{0,1}\|_{H^1}) \|\delta u\|_{L^2}^{4/p} \|\nabla \delta u\|_{L^2}^{2(1-2/p)} + \|\delta r\|_{L^2}^2 \\ \leq \eta \|\nabla \delta u\|_{L^2}^2 + C(\eta, \beta, \|r_{0,1}\|_{H^{1+\beta}}, \|u_{0,1}\|_{H^1}, \|b_{0,1}\|_{H^1}) \|\delta u\|_{L^2}^2 + \|\delta r\|_{L^2}^2.$$

We still have to handle three integrals, which involve the magnetic field. Firstly, integration by parts gives

$$\left| \int_{\Omega} (\delta u \cdot \nabla) b_1 \cdot \delta b dx \right| = \left| \int_{\Omega} (\delta u \cdot \nabla) \delta b \cdot b_1 dx \right| \leq \|\delta u\|_{L^4} \|\nabla \delta b\|_{L^2} \|b_1\|_{L^4} \\ \leq \eta \|\nabla \delta b\|_{L^2}^2 + C(\eta) \|\delta u\|_{L^4}^2 \|b_1\|_{L^4}^2 \leq \eta \|\nabla \delta b\|_{L^2}^2 + \|\delta u\|_{L^2} \|\nabla \delta u\|_{L^2} \|b_1\|_{L^4}^2 \\ \leq \eta (\|\nabla u\|_{L^2}^2 + \|\nabla b\|_{L^2}^2) + M_1(t) \|\delta b\|_{L^2}^2,$$

where we have also used the GN inequality in the second line, and Young's inequality in order to get the last inequality, and where we have set $M_1(t) = C(\eta) \|b_1(t)\|_{L^4}^4 \in L^1(\mathbb{R}_+)$, because of the same bounds exhibited above for $\|u_1\|_{L^4}^4$. The last integral in (54) can be treated in the same way: after integration by parts, we get

$$\left| \int_{\Omega} (\delta b \cdot \nabla) b_1 \cdot \delta u dx \right| = \left| \int_{\Omega} (\delta b \cdot \nabla) \delta u \cdot b_1 dx \right| \leq \|\delta u\|_{L^4} \|\nabla \delta b\|_{L^2} \|b_1\|_{L^4} \leq M_1(t) \|\delta b\|_{L^2}^2.$$

Finally, for the remaining term we can apply one last time the GN inequality: we infer

$$\begin{aligned} \left| \int_{\Omega} (\delta b \cdot \nabla) u_1 \cdot \delta b \, dx \right| &\leq \|\nabla u_1\|_{L^2} \|\delta b\|_{L^4}^2 \leq C \|\nabla u_1\|_{L^2} \|\delta b\|_{L^2} \|\nabla \delta b\|_{L^2} \\ &\leq \eta \|\nabla \delta b\|_{L^2}^2 + C(\eta) \|\nabla u_1\|_{L^2}^2 \|\delta b\|_{L^2}^2 \leq \eta \|\nabla \delta b\|_{L^2}^2 + M_1(t) \|\delta b\|_{L^2}^2, \end{aligned}$$

where $M_1(t) = C(\eta) \|\nabla u_1(t)\|_{L^2}^2 \in L^1(\mathbb{R}_+)$.

Define now

$$E(t) := \|\delta u(t)\|_{L^2}^2 + \|\delta b(t)\|_{L^2}^2 + \|\delta r(t)\|_{L^2}^2 \quad \text{and} \quad E_0 := \|\delta u_0\|_{L^2}^2 + \|\delta b_0\|_{L^2}^2 + \|\delta r_0\|_{L^2}^2.$$

Putting all the previous bounds together and choosing η so small that the gradient terms can be absorbed in the left-hand side, from (54) we arrive at the differential inequality

$$(55) \quad \frac{1}{2} \frac{d}{dt} E(t) + \frac{1}{2} \int_{\Omega} \left\{ |\nabla \delta u|^2 + |\nabla \delta b|^2 \right\} dx \leq N_1(t) E(t),$$

where $N_1(t) \in L_T^1$ is the sum of all the functions $M_1 \in L^1(\mathbb{R}_+)$, $N_4(t) \in L_T^4$ in the previous inequalities. Therefore, Grönwall's lemma implies that, for all $t \geq 0$, one has

$$E(t) \leq C(t) E_0.$$

Coming back to (55), we finally get (50). The proposition is then proved. \square

A Appendix – Fourier and harmonic analysis toolbox

We recall here the main ideas of Littlewood-Paley theory, which we exploited in the previous analysis. We refer e.g. to Chapter 2 of [4] for details. For simplicity of exposition, let us deal with the \mathbb{R}^d case; however, the whole construction can be adapted also to the d -dimensional torus \mathbb{T}^d .

First of all, let us introduce the so called “Littlewood-Paley decomposition”, based on a non-homogeneous dyadic partition of unity with respect to the Fourier variable. We fix a smooth radial function χ supported in the ball $B(0, 2)$, equal to 1 in a neighborhood of $B(0, 1)$ and such that $r \mapsto \chi(re)$ is nonincreasing over \mathbb{R}_+ for all unitary vectors $e \in \mathbb{R}^d$. Set $\varphi(\xi) = \chi(\xi) - \chi(2\xi)$ and $\varphi_j(\xi) := \varphi(2^{-j}\xi)$ for all $j \geq 0$.

The dyadic blocks $(\Delta_j)_{j \in \mathbb{Z}}$ are defined by⁴

$$\Delta_j := 0 \quad \text{if } j \leq -2, \quad \Delta_{-1} := \chi(D) \quad \text{and} \quad \Delta_j := \varphi(2^{-j}D) \quad \text{if } j \geq 0.$$

We also introduce the following low frequency cut-off operator:

$$(56) \quad S_j u := \chi(2^{-j}D) = \sum_{k \leq j-1} \Delta_k \quad \text{for } j \geq 0.$$

Note that the operator S_j is a convolution operator with a function $K_j(x) = 2^{dj} K_1(2^j x) = \mathcal{F}^{-1}[\chi(2^{-j}\xi)](x)$ of constant L^1 norm, and hence defines a continuous operator for the $L^p \rightarrow L^p$ topologies, for any $1 \leq p \leq +\infty$.

The following classical property holds true: for any $u \in \mathcal{S}'$, then one has the equality $u = \sum_j \Delta_j u$ in the sense of \mathcal{S}' . Let us also mention the so-called *Bernstein inequalities*, which explain the way derivatives act on spectrally localized functions.

Lemma A.1. *Let $0 < r < R$. A constant C exists so that, for any nonnegative integer k , any couple (p, q) in $[1, +\infty]^2$, with $p \leq q$, and any function $u \in L^p$, we have, for all $\lambda > 0$,*

$$\begin{aligned} \text{Supp } \widehat{u} \subset B(0, \lambda R) &\implies \|\nabla^k u\|_{L^q} \leq C^{k+1} \lambda^{k+d(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p}; \\ \text{Supp } \widehat{u} \subset \{\xi \in \mathbb{R}^d \mid r\lambda \leq |\xi| \leq R\lambda\} &\implies C^{-k-1} \lambda^k \|u\|_{L^p} \leq \|\nabla^k u\|_{L^p} \leq C^{k+1} \lambda^k \|u\|_{L^p}. \end{aligned}$$

⁴Throughout we agree that $f(D)$ stands for the pseudo-differential operator $u \mapsto \mathcal{F}^{-1}[f(\xi)\widehat{u}(\xi)]$.

By use of Littlewood-Paley decomposition, we can define the class of Besov spaces.

Definition A.2. Let $s \in \mathbb{R}$ and $1 \leq p, r \leq +\infty$. The *non-homogeneous Besov space* $B_{p,r}^s$ is defined as the subset of tempered distributions u for which

$$\|u\|_{B_{p,r}^s} := \left\| \left(2^{js} \|\Delta_j u\|_{L^p} \right)_{j \geq -1} \right\|_{\ell^r} < +\infty.$$

Besov spaces are interpolation spaces between Sobolev spaces. In fact, for any $k \in \mathbb{N}$ and $p \in [1, +\infty]$ we have the following chain of continuous embeddings: $B_{p,1}^k \hookrightarrow W^{k,p} \hookrightarrow B_{p,\infty}^k$, where $W^{k,p}$ denotes the classical Sobolev space of L^p functions with all the derivatives up to the order k in L^p . When $1 < p < +\infty$, we can refine the previous result (this is the non-homogeneous version of Theorems 2.40 and 2.41 in [4]): we have

$$B_{p,\min(p,2)}^k \hookrightarrow W^{k,p} \hookrightarrow B_{p,\max(p,2)}^k.$$

In particular, for all $s \in \mathbb{R}$ we deduce the equivalence $B_{2,2}^s \equiv H^s$, with equivalence of norms:

$$(57) \quad \|f\|_{H^s} \sim \left(\sum_{j \geq -1} 2^{2js} \|\Delta_j f\|_{L^2}^2 \right)^{1/2}.$$

As an immediate consequence of the first Bernstein inequality, one gets the following embedding result.

Proposition A.3. *The space $B_{p_1,r_1}^{s_1}$ is continuously embedded in the space $B_{p_2,r_2}^{s_2}$ for all indices satisfying $p_1 \leq p_2$ and*

$$s_2 < s_1 - d \left(\frac{1}{p_1} - \frac{1}{p_2} \right) \quad \text{or} \quad s_2 = s_1 - d \left(\frac{1}{p_1} - \frac{1}{p_2} \right) \quad \text{and} \quad r_1 \leq r_2.$$

In particular, in dimension $d = 2$, we get the regular Sobolev embeddings

$$(58) \quad H^s \simeq B_{2,2}^s \hookrightarrow B_{p,2}^{s-2\left(\frac{1}{2}-\frac{1}{p}\right)} \hookrightarrow B_{p,2}^0 = B_{p,\min(p,2)}^0 \hookrightarrow L^p$$

as long as $0 \leq s < 1$ and $2 \leq p \leq 2/(1-s)$. Also note that, still for $d = 2$, one has the embedding $H^s \hookrightarrow L^\infty \cap C^0$ for all $s > 1$.

Let us now introduce the paraproduct operator (after J.-M. Bony, see [5]). Constructing the paraproduct operator relies on the observation that, formally, any product of two tempered distributions u and v , may be decomposed into

$$(59) \quad uv = T_u(v) + T_v(u) + R(u, v),$$

where we have defined

$$T_u(v) := \sum_j S_{j-1} u \Delta_j v, \quad \text{and} \quad R(u, v) := \sum_j \sum_{|j'-j| \leq 1} \Delta_j u \Delta_{j'} v.$$

The above operator T is called “paraproduct” whereas R is called “remainder”. The paraproduct and remainder operators have many nice continuity properties. The following ones have been of constant use in this paper (see the proof in e.g. Chapter 2 of [4]).

Proposition A.4. *For any $(s, p, r) \in \mathbb{R} \times [1, \infty]^2$ and $t > 0$, the paraproduct operator T maps continuously $L^\infty \times B_{p,r}^s$ in $B_{p,r}^s$ and $B_{\infty,\infty}^{-t} \times B_{p,r}^s$ in $B_{p,r}^{s-t}$. Moreover, the following estimates hold:*

$$\|T_u(v)\|_{B_{p,r}^s} \leq C \|u\|_{L^\infty} \|\nabla v\|_{B_{p,r}^{s-1}} \quad \text{and} \quad \|T_u(v)\|_{B_{p,r}^{s-t}} \leq C \|u\|_{B_{\infty,\infty}^{-t}} \|\nabla v\|_{B_{p,r}^{s-1}}.$$

For any (s_1, p_1, r_1) and (s_2, p_2, r_2) in $\mathbb{R} \times [1, \infty]^2$ such that $s_1 + s_2 > 0$, $1/p := 1/p_1 + 1/p_2 \leq 1$ and $1/r := 1/r_1 + 1/r_2 \leq 1$, the remainder operator R maps continuously $B_{p_1,r_1}^{s_1} \times B_{p_2,r_2}^{s_2}$ into $B_{p,r}^{s_1+s_2}$. In the case $s_1 + s_2 = 0$, provided $r = 1$, operator R is continuous from $B_{p_1,r_1}^{s_1} \times B_{p_2,r_2}^{s_2}$ with values in $B_{p,\infty}^0$.

As a corollary of the previous proposition, we deduce the following continuity properties of the product in Sobolev spaces, which have been used in the course of the analysis. In the statement, we limit ourselves to the case of space dimension $d = 2$, the only relevant one for this study.

Lemma A.5. *Take the space dimension to be $d = 2$. For appropriate f and g , one has the following properties:*

- (1) for $s \in \mathbb{R}$ and $t > 0$, $\|T_f(g)\|_{H^{s-t}} \leq C\|f\|_{H^{1-t}}\|g\|_{H^s}$.
- (2) for $s \in \mathbb{R}$, $\|T_f(g)\|_{H^s} \leq C\|f\|_{L^\infty}\|g\|_{H^s}$;
- (3) for $s_1, s_2 \in \mathbb{R}$ such that $s_1 + s_2 > 0$, $\|R(f, g)\|_{H^{s_1+s_2-1}} \leq C\|f\|_{H^{s_1}}\|g\|_{H^{s_2}}$.

Proof. We start by proving the first point. From the second inequality in Proposition A.4, we get

$$\|T_f(g)\|_{H^{s-t}} = \|T_f(g)\|_{B_{2,2}^{s-t}} \leq C\|f\|_{B_{\infty,\infty}^{-t}}\|\nabla g\|_{B_{2,2}^{s-1}} = C\|f\|_{B_{\infty,\infty}^{-t}}\|\nabla g\|_{H^{s-1}}.$$

Because $d = 2$, Proposition A.3 gives the embedding $H^{1-t} = B_{2,2}^{1-t} \hookrightarrow B_{\infty,\infty}^{-t}$, from which we infer the first inequality.

Next, using the first inequality in Proposition A.4, we gather

$$\|T_f(g)\|_{H^s} = \|T_f(g)\|_{B_{2,2}^{s-t}} \leq C\|f\|_{L^\infty}\|\nabla g\|_{B_{2,2}^{s-1}} \leq C\|f\|_{L^\infty}\|g\|_{H^s},$$

which proves the second point.

Finally, using Proposition A.4 to estimate the remainder term, because we have assumed that $s_1 + s_2 > 0$, we get

$$\|R(f, g)\|_{B_{1,1}^{s_1+s_2}} \leq C\|f\|_{H^{s_1}}\|g\|_{H^{s_2}}.$$

Proposition A.3 provides the embedding $B_{1,1}^{s_1+s_2} \hookrightarrow B_{2,2}^{s_1+s_2-1} = H^{s_1+s_2-1}$, which gives the last inequality, thus completing the proof of the lemma. \square

Corollary A.6. *As a consequence of the previous lemma, we see that*

- (i) for any $\delta > 0$, the space $H^{1+\delta}$ is a Banach algebra;
- (ii) for all $s > -1$ and all $(f, g) \in H^1 \times H^s$, we have $fg \in H^{s-\delta}$ for any $\delta > 0$, with

$$\|fg\|_{H^{s-\delta}} \leq C\|f\|_{H^1}\|g\|_{H^s}.$$

The next two propositions are functional inequalities which we use repeatedly in this article. The first one is the classical Gagliardo-Nirenberg inequality, whose proof can be found e.g. in Corollary 1.2 of [6].

Lemma A.7. *Let $2 \leq p < +\infty$ such that $1/p > 1/2 - 1/d$. Then, for all $u \in H^1$, one has*

$$\|u\|_{L^p} \leq C(p)\|u\|_{L^2}^\lambda \|\nabla u\|_{L^2}^{1-\lambda}, \quad \text{with} \quad \lambda = \frac{d(p-2)}{2p}.$$

In particular, in dimension $d = 2$, we have $\|u\|_{L^p} \leq \|u\|_{L^2}^{2/p} \|\nabla u\|_{L^2}^{1-2/p}$ for any $p \in [2, +\infty[$.

The following proposition, which is in the same spirit of the Gagliardo-Nirenberg inequality, gives a bound for the endpoint case $p = +\infty$. It is proved by resorting to Littlewood-Paley decomposition and the Bernstein inequalities.

Proposition A.8. *Let $f \in H^s$, for some $s > d/2$. Then there exists a constant $C = C(d, s) > 0$ and an exponent $\alpha = \alpha(d, s) = d/(2s)$ such that*

$$\|f\|_{L^\infty} \leq C \|f\|_{L^2}^{1-\alpha} \left\| (-\Delta)^{s/2} f \right\|_{L^2}^\alpha.$$

In particular, when $d = 2$ and $s = 2$, then $\alpha = 1/2$ and $\|f\|_{L^\infty} \leq C \|f\|_{L^2}^{1/2} \|\Delta f\|_{L^2}^{1/2}$.

Proof. The main idea of the proof is to look separately at the high and low frequencies. Let $N \geq 1$ be an integer to be fixed later on. Thanks to the Littlewood-Paley decomposition, we can write

$$\|f\|_{L^\infty} \leq \sum_{j < N} \|\Delta_j f\|_{L^\infty} + \sum_{j \geq N} \|\Delta_j f\|_{L^\infty}.$$

Using the first Bernstein inequality in the first sum gives $\|\Delta_j f\|_{L^\infty} \leq C 2^{jd/2} \|\Delta_j f\|_{L^2}$. On the other hand, the two Bernstein inequalities applied to the high frequency term yield $\|\Delta_j f\|_{L^2} \leq C 2^{jd/2} 2^{-sj} \|\Delta_j (-\Delta)^{s/2} f\|_{L^2}$. Therefore

$$\begin{aligned} \|f\|_{L^\infty} &\leq C \|f\|_{L^2} \sum_{j < N} 2^{jd/2} + C \left\| (-\Delta)^{s/2} f \right\|_{L^2} \sum_{j \geq N} 2^{-j(s-d/2)} \\ &\leq C \left(\|f\|_{L^2} 2^{Nd/2} + \left\| (-\Delta)^{s/2} f \right\|_{L^2} 2^{-N(s-d/2)} \right). \end{aligned}$$

By choosing N so that $2^{Ns} \approx \left\| (-\Delta)^{s/2} f \right\|_{L^2} \|f\|_{L^2}^{-1}$ (say that N is the largest integer such that 2^{Ns} is smaller than $\left\| (-\Delta)^{s/2} f \right\|_{L^2} \|f\|_{L^2}^{-1}$), we deduce the desired inequality. \square

Finally, we recall a classical commutator estimate, which we have needed in our analysis (see Lemma 2.97 of [4] for the proof).

Lemma A.9. *Let $\chi \in C^1(\mathbb{R}^d)$ be such that $H(\xi) := (1 + |\xi|)\widehat{\chi}(\xi) \in L^1$. There exists a constant $C > 0$ depending only on $\|H\|_{L^1}$ such that*

$$\forall f \in W^{1,\infty}, \forall g \in L^2, \forall \lambda > 0, \quad \left\| \left[\chi \left(\frac{1}{\lambda} D \right), f \right] g \right\|_{L^2} \leq C \frac{1}{\lambda} \|\nabla f\|_{L^\infty} \|g\|_{L^2}.$$

References

- [1] A. Babin, A. Mahalov, B. Nicolaenko: *Global splitting, integrability and regularity of 3D Euler and Navier-Stokes equations for uniformly rotating fluids*. European J. Mech. B Fluids 15 (1996), n. 3, pp. 291–300.
- [2] A. Babin, A. Mahalov, B. Nicolaenko: *Regularity and integrability of 3D Euler and Navier-Stokes equations for rotating fluids*. Asymptot. Anal. 15 (1997), n. 2, pp. 103–150.
- [3] A. Babin, A. Mahalov, B. Nicolaenko: *Global regularity of 3D rotating Navier-Stokes equations for resonant domains*. Indiana Univ. Math. J. 48 (1999), n. 3, pp. 1133–1176.
- [4] H. Bahouri, J.-Y. Chemin and R. Danchin: *“Fourier analysis and nonlinear partial differential equations”*. Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), 343. Springer, Heidelberg, 2011.
- [5] J.-M. Bony: *Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires*. (in french) Ann. Sci. École Norm. Sup. 14 (1981), n. 2, pp. 209–246.
- [6] J. -Y. Chemin, B. Desjardins, I. Gallagher and E. Grenier: *“Mathematical geophysics. An introduction to rotating fluids and the Navier-Stokes equations”*. Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 2006.
- [7] B. Cushman-Roisin: *“Introduction to geophysical fluid dynamics”*. Prentice-Hall, Englewood Cliffs, 1994.

- [8] R. Danchin: *Local and global well-posedness results for flows of inhomogeneous viscous fluids*. Adv. Differential Equations 9 (2004), n. 3–4, pp. 353–386.
- [9] F. De Anna, F. Fanelli: *Global well-posedness and long-time dynamics for a higher order quasi-geostrophic type equation*. J. Funct. Anal. 274 (2018), n. 8, pp. 2291–2355.
- [10] B. Desjardins, E. Dormy and E. Grenier: *Stability of mixed Ekman-Hartmann boundary layers*. Nonlinearity 12 (1999), n. 2, pp. 181–199.
- [11] B. Desjardins and C. Le Bris: *Remarks on a nonhomogeneous model of magnetohydrodynamics*. Differential Integral Equations 11 (1998), n. 3, pp. 377–394.
- [12] R. J. Di Perna and P.-L. Lions: *Ordinary differential equations, transport theory and Sobolev spaces*. Invent. Math. 98 (1989), n. 3, pp. 511–547.
- [13] F. Fanelli: *Highly rotating viscous compressible fluids in presence of capillarity effects*. Math. Ann. 366 (2016), n. 3–4, pp. 981–1033.
- [14] F. Fanelli: *A singular limit problem for rotating capillary fluids with variable rotation axis*. J. Math. Fluid Mech. 18 (2016), n. 4, pp. 625–658.
- [15] F. Fanelli: *Incompressible and fast rotation limit for barotropic Navier-Stokes equations at large Mach numbers*. Submitted (2019), arxiv preprint arxiv.org/abs/1909.09894.
- [16] F. Fanelli and I. Gallagher: *Asymptotics of fast rotating density-dependent incompressible fluids in two space dimensions*. Rev. Mat. Iberoam. 35 (2019), n. 6, pp. 1763–1807.
- [17] E. Feireisl, I. Gallagher, D. Gerard-Varet and A. Novotný: *Multi-scale analysis of compressible viscous and rotating fluids*. Comm. Math. Phys. 314 (2012), n. 3, pp. 641–670.
- [18] E. Feireisl, I. Gallagher and A. Novotný: *A singular limit for compressible rotating fluids*. SIAM J. Math. Anal. 44 (2012), n. 1, pp. 192–205.
- [19] E. Feireisl, Y. Lu, A. Novotný: *Rotating compressible fluids under strong stratification*. Nonlinear Anal. Real World Appl. 19 (2014), pp. 11–18.
- [20] E. Feireisl, A. Novotný: *Multiple scales and singular limits for compressible rotating fluids with general initial data*. Comm. Partial Differential Equations 39 (2014), n. 6, pp. 1104–1127.
- [21] I. Gallagher and L. Saint-Raymond: *Weak convergence results for inhomogeneous rotating fluid equations*. J. Anal. Math. 99 (2006), pp. 1–34.
- [22] J.-F. Gerbeau and C. Le Bris: *Existence of solution for a density-dependent magnetohydrodynamic equation*. Adv. Differential Equations 2 (1997), n. 3, pp. 427–452.
- [23] Y.-S. Kwon, Y.-C. Lin and C.-F. Su: *Derivation of inviscid quasi-geostrophic equation from rotational compressible magnetohydrodynamic flows*, J. Nonlinear Sci. 28 (2018), n. 2, pp. 599–620.
- [24] Y.-S. Kwon, D. Maltese, A. Novotný: *Multiscale analysis in the compressible rotating and heat conducting fluids*. J. Math. Fluid Mech. 20 (2018), n. 2, pp. 421–444.
- [25] P.-L. Lions: *“Mathematical topics in fluid dynamics. Vol. 1: incompressible models”*. Oxford University Press, Oxford, 1996.
- [26] P.-L. Lions, N. Masmoudi: *Incompressible limit for a viscous compressible fluid*. J. Math. Pures Appl. 77 (1998), n. 6, pp. 585–627.
- [27] V.-S. Ngo: *A Global existence result for the anisotropic rotating magnetohydrodynamical system*. Acta Appl. Math. 150 (2017), n. 1, pp. 1–42.
- [28] J. Pedlosky: *“Geophysical fluid dynamics”*. Springer-Verlag, New-York, 1987.
- [29] M. Paicu, P. Zhang, Z. Zhang: *Global unique solvability of inhomogeneous Navier-Stokes equations with bounded density*. Comm. Partial Differential Equations 38 (2013), n. 7, pp. 1208–1234.
- [30] F. Rousset: *Stability of large amplitude Ekman-Hartmann boundary layers in MHD: the case of ill-prepared data*. Comm. Math. Phys. 259 (2005), n. 1, pp. 223–256.