

Lie Polynomials and a Twistorial Correspondence for Amplitudes

Hadleigh Frost and Lionel Mason

**The Mathematical Institute, University of Oxford,
AWB, ROQ, Oxford OX2 6GG, United Kingdom**

ABSTRACT: We review Lie polynomials as a mathematical framework that underpins the structure of the so-called double copy relationship between gauge and gravity theories (and a network of other theories besides). We explain how Lie polynomials naturally arise in the geometry and cohomology of $\mathcal{M}_{0,n}$, the moduli space of n points on the Riemann sphere up to Möbius transformation. We introduce a twistorial correspondence between the cotangent bundle $T_D^*\mathcal{M}_{0,n}$, the bundle of forms with logarithmic singularities on the divisor D as the twistor space, and \mathcal{K}_n the space of momentum invariants of n massless particles subject to momentum conservation as the analogue of space-time. This gives a natural framework for Cachazo He and Yuan (CHY) and ambitwistor-string formulae for scattering amplitudes of gauge and gravity theories as being the corresponding Penrose transform. In particular we show that it gives a natural correspondence between CHY half-integrands and scattering forms, certain $n-3$ -forms on \mathcal{K}_n , introduced by Arkani-Hamed, Bai, He and Yan (ABHY). We also give a generalization and more invariant description of the associahedral $n-3$ -planes in \mathcal{K}_n introduced by ABHY.

KEYWORDS: Scattering amplitudes, Lie Polynomials, twistor theory

Contents

1	Introduction	1
2	The double copy and Lie polynomials	2
2.1	A review of words, Lie polynomials and trees	3
2.2	The geometry of \mathcal{K}_n	5
2.3	The double copy from biadjoint scalars to gauge and gravity theories	6
2.4	A note on the Kinematic Algebra	8
3	Trees and words in $\mathcal{M}_{0,n}$	8
3.1	The cohomology of $\mathcal{M}_{0,n}$ and Parke-Taylor	10
3.2	Homology of $\mathcal{M}_{0,n}$	11
4	A Penrose transform for amplitudes	13
4.1	The double fibration and the CHY formulae	14
4.2	The geometry of the correspondence	15
4.3	The symplectic form and the holomorphic volume form	15
4.4	Associahedral $(n - 3)$ -planes in \mathcal{K}_n and forms on $\mathcal{M}_{0,n}$.	17
4.5	Scattering forms and CHY	20
5	Discussion	21
6	Acknowledgements	22

1 Introduction

Color-kinematics duality and the double copy [1, 2] have had a powerful influence on recent developments in scattering amplitudes. They stem from the KLT relations in string theory [3] between gravity and Yang-Mills tree-amplitudes and have been developed as a tool for the study of multiloop gravity amplitudes and more recently for applications to perturbative classical gravity calculations in connection with gravitational waves [4]. Notwithstanding the physical applications, the underlying mathematical framework is perhaps rather surprising even at tree level. There is little hint of such a double copy structure in standard approaches to perturbation theory of the classical nonlinear theories involved. The purpose of this article is to develop some of the underpinning mathematical structures. We build on observations by Kapranov in an after dinner talk [5] concerning the relevance of Lie polynomials, both in the double copy and in the Parke-Taylor expressions that pervade the subject. We also

build on the recent work by Arkani-Hamed, Bai, He and Yan [6] that introduces differential forms in the space of kinematic invariants, \mathcal{K}_n . We tie them together by means of a double fibration correspondence that leads to a Penrose-like transform for the formulae of Cachazo He and Yuan (CHY) arising from the scattering equations [7, 8].

The first section provides an elementary review of the theory of Lie polynomials as relevant to this topic, and expresses standard facts about the double copy in this language. In particular, the trivalent diagrams of BCJ are a representation of elements $\Gamma \in \text{Lie}(n-1)$, the space of Lie polynomials of degree $n-1$, and BCJ numerators N_Γ are homomorphisms $N : \text{Lie}(n-1) \rightarrow V$ where V is some vector space of polynomials in the polarization data and momenta.

We next review the role played by Lie polynomials in the geometry of the moduli space $\mathcal{M}_{0,n}$ of n points σ_i in \mathbb{CP}^1 , both in describing the compact cycles in the homology $H_{n-3}(\mathcal{M}_{0,n} - D)$, which is isomorphic to $\text{Lie}(n-1)$, and dually the relative cocycles in $H^{n-3}(\mathcal{M}_{0,n}, D)$, represented by the top degree holomorphic *Parke-Taylor forms*.

With these preliminaries in hand, we study a double fibration between the space of Mandelstam variables, \mathcal{K}_n , and $T_D^* \mathcal{M}_{0,n}$

$$\begin{array}{ccc} \mathcal{Y}_n = \mathcal{K}_n \times \mathcal{M}_{0,n}, (s_{ij}, \sigma_j) & & \\ p \swarrow & \searrow q & \\ (s_{ij}), \mathcal{K}_n & & \mathbb{T} = T_D^* \mathcal{M}_{0,n}, (\tau_i, \sigma_i). \end{array} \quad (1.1)$$

where p forgets the second factor and q is defined by the incidence relations

$$\tau_i = E_i(s_{kl}, \sigma_m) := \sum_j \frac{s_{ij}}{\sigma_{ij}}, \quad (1.2)$$

which give the left hand side of the scattering equations. In the language of this correspondence, the CHY formulae are a Penrose transform, being simply the push down of the pullback of certain forms on \mathbb{T} . We investigate other more geometrical aspects of the Penrose transform. In particular we show that the top power of the symplectic form ω^{n-3} provides a correspondence between certain $(n-3)$ -forms w_Γ on \mathcal{K}_n that were introduced by ABHY and homology classes in $\mathcal{M}_{0,n}$. ABHY use the w_Γ as numerators so that given a set of conventional numerators N one can associate a *scattering* $(n-3)$ -form Ω_N . These arise from our double fibration via a Penrose transform also. Dually, ABHY introduce associahedral $(n-3)$ -planes in \mathcal{K}_n that can be used to convert a scattering form into a conventional amplitude. We give an improved and extended definition of these, and show how they tie into the geometry of the correspondence.

2 The double copy and Lie polynomials

Colour structures for n -point amplitudes are degree n invariant polynomials of weight one in each of the n Lie algebra ‘colours’ of the external particles. These naturally arise in

Feynman rules as trivalent Feynman diagrams whose vertices are the structure constants of some unspecified Lie algebra. If we fix the n^{th} particle, and an invariant inner product on the Lie algebra, at tree-level, such a polynomial can be realized as the inner product of the n^{th} colour with the Lie algebra element with a *Lie polynomial* formed by successive commutators of the $n-1$ other colours working through the diagram back from the n^{th} particle. This section reviews material concerning such colour structures in the language of free Lie algebras and Lie polynomials together with their duality with words formed from permutations of the $n-1$ labels of the first $n-1$ external particles. A classic text on free Lie algebras is [9].

2.1 A review of words, Lie polynomials and trees

The space of words, $W(n-1)$, is the $(n-1)!$ -dimensional linear span of words

$$a = x_{a(1)}x_{a(2)} \cdots x_{a(n-1)},$$


where the letters $x_{a(i)} \in \{x_1, \dots, x_{n-1}\}$ are all distinct, so that the a 's define permutations on $n-1$ letters. There is a natural bilinear inner product on $W(n-1)$ that is defined on monomials a and b by

$$(a, b) := \prod_{i=1}^{n-1} \delta_{a(i)b(i)},$$

i.e., (a, b) is 1 if a and b are the same word, and 0 otherwise. A ‘Lie polynomial’ in $W(n-1)$ is any expression formed by taking $n-2$ iterated commutators of the x_i . An example is

$$\Gamma = [x_1, [\dots, [x_{n-1}, x_n] \dots]] + [x_n, [\dots, [x_{n-2}, x_{n-1}] \dots]],$$

where $[x_i, x_j]$ is the commutator, $x_i x_j - x_j x_i$. Let $Lie(n-1)$ be the linear subspace of $W(n-1)$ generated by all Lie polynomials Γ of weight $n-1$ in the $n-1$ variables, x_1, \dots, x_{n-1} , with weight one in each. Every Lie monomial Γ defines a rooted trivalent tree decorated by an orientation. We denote this tree also by Γ . An orientation of the tree can be presented as a planar embedding, where two planar embeddings have the same orientation if they differ from one another by an even number of flips. Thus, for example, the monomial $\Gamma = [2, [[1, 3], [5, 4]]]$ is associated with the following two planar embeddings, among others.



The Jacobi identity implies the vanishing of the sum of the three four-point oriented trees corresponding to an s, t and u -channel exchange graph.

$$\begin{array}{c} 2 \quad 3 \\ \diagdown \quad \diagup \\ 1 \text{---} \text{---} 4 \end{array} - \begin{array}{c} 2 \quad 3 \\ | \quad | \\ 1 \text{---} \text{---} 4 \end{array} + \begin{array}{c} 3 \quad 2 \\ | \quad | \\ 1 \text{---} \text{---} 4 \end{array} = 0$$

We will denote three Lie polynomials or corresponding graphs that differ only on such a four point subgraph by Γ_s, Γ_t and Γ_u , and we will consequently have

$$\Gamma_s + \Gamma_t + \Gamma_u = 0. \quad (2.1)$$

The Lie monomial notation is useful for keeping track of the orientations of trees. Recall the inner product $(\ , \)$ defined on words. For a Lie monomial Γ , (Γ, a) is the coefficient of a in the expansion of its Lie monomial Γ . When Γ is not planar for the ordering a , $(\Gamma, a) = 0$. When Γ is planar for the ordering a , (Γ, a) is the orientation of that planar embedding, either +1 or -1. As an application of the notation we can write

$$\Gamma = \sum_a (\Gamma, a) a, \quad (2.2)$$

which is simply the expansion of the commutators in Γ .

There are many characterizations of $Lie(n-1)$ as a subspace of $W(n-1)$ [9], and some of these have long been known in the physics literature as relations among gauge theory tree amplitudes. For instance, the U(1) decoupling identity is a consequence of Ree's theorem.

Proposition 2.1 (Ree [10]) *A polynomial $w \in W[n-1]$ is a Lie polynomial iff $(w, a \sqcup b) = 0$ for all nontrivial shuffles $a \sqcup b$.¹*

This proposition implies that $Lie(n-1)^*$, the dual vector space of $Lie(n-1)$, can be understood as the quotient vector space $W(n-1)/\text{Sh}$, where Sh is the subspace generated by all nontrivial shuffles.

Lemma 2.2 (Radford [11]) *The $(n-2)!$ words of the form $1a$ are a basis for $Lie(n-1)^*$.*

The direct expansion of Γ_{1a} into words is

$$\Gamma_{1a} = \sum_{a \in \bar{u} \sqcup v} (-1)^{|u|} u 1 v, \quad (2.3)$$

where $|u|$ is the length of u . It follows that

$$(\Gamma_{1a}, 1b) = (a, b).$$

In other words, we have the following.

Lemma 2.3 *The Kleiss-Kuijff (KK) basis of $Lie(n-1)^*$ given by words $1a$ is dual to the DDM basis of $Lie(n-1)$ given by combs Γ_{1a} .*

¹The shuffle product of a and b , $a \sqcup b$, is the sum over ordered permutations of the letters of a and b that preserve the orderings of the letters in a and b .

An immediate consequence of this Lemma is that any $b + \text{Sh} \in W(n-1)/\text{Sh}$ may be expanded in this basis as

$$b + \text{Sh} = \sum_a (\Gamma_{1a}, b) 1a + \text{Sh}. \quad (2.4)$$

Dually, given that the combs are a basis for $\text{Lie}(n-1)$, a polynomial w is a Lie polynomial iff there is an expansion of w in the combs Γ_{1a} . Using Lemma 2.3, we find that

$$w = \sum_a (w, 1a) \Gamma_{1a}. \quad (2.5)$$

By equation (2.3), we find that

$$w = \sum (w, 1(\bar{u} \sqcup v)) (-1)^{|u|} u 1v,$$

where \bar{u} is the reversal of u . This identity implies the Kleiss-Kuijf relations, which we can restate as a theorem about Lie polynomials.

Proposition 2.4 (*Kleiss-Kuijf*) *A polynomial $w \in W[n-1]$ is a Lie polynomial iff the Kleiss-Kuijf relations [12] hold:*

$$(w, a1b) = (-1)^{|a|} (w, 1\bar{a} \sqcup b), \quad (2.6)$$

.

2.2 The geometry of \mathcal{K}_n

The space of Mandelstam variables is $\mathcal{K}_n \simeq \mathbb{R}^{n(n-3)/2}$. In coordinates s_{ij} , $i, j = 1, \dots, n$ (with $s_{ij} = s_{ji}$, $s_{ii} = 0$), \mathcal{K}_n is the hyperplane given by the equations

$$\sum_{j=1}^n s_{ij} = 0. \quad (2.7)$$

For amplitudes, the key geometric structure in \mathcal{K}_n are the factorization hyperplanes given by $s_I = 0$, where $I \subset \{1, 2, \dots, n\}$ and

$$s_I := \sum_{i,j \in I} s_{ij} = \left(\sum_i k_i \right)^2. \quad (2.8)$$

Let \bar{I} be the complement of I , so that $s_{\bar{I}} = -s_I$ by (2.7). Locality states that the only singularities of tree amplitudes are simple poles on these hyperplanes. A further requirement is that a double pole on the intersection of $s_I = 0$ and $s_J = 0$ occurs in an amplitude only if $I \subset J$ or $\bar{I} \subset \bar{J}$. It follows that the allowed pole structure of a contribution to an n -point amplitude has poles along at most $n-3$ factorization hyperplanes $s_{I_p} = 0$ for $p = 1, \dots, n-3$. Such choices are in one-to-one correspondence with trivalent (and unoriented) trees.

2.3 The double copy from biadjoint scalars to gauge and gravity theories

The double copy principle is that massless n -point tree amplitudes for a large web of important theories, including many gauge and gravity theories, can be expressed as a double copy in the form

$$\mathcal{M} = \sum_{\Gamma} \frac{N_{\Gamma} \tilde{N}_{\Gamma}}{d_{\Gamma}}. \quad (2.9)$$

Here the denominators

$$d_{\Gamma} = \prod_{r=1}^{n-3} s_{I_r} \quad (2.10)$$

are the propagator factors associated to the graph Γ thought of as a Feynman graph. Further, each trivalent diagram Γ has a pair of *numerator* factors N_{Γ} and \tilde{N}_{Γ} that are functions of momenta, polarization data, flavour and colour. Such factors are said to be *local* if they are polynomial, i.e., admit no spurious singularities.

The key additional feature required to be a BCJ numerator is that N_{Γ} and \tilde{N}_{Γ} should represent homomorphisms from Lie polynomials to some vector space V of functions,

$$N : \text{Lie}(n-1) \rightarrow V. \quad (2.11)$$

Thus for any three graphs Γ_s , Γ_t and Γ_u satisfying $\Gamma_s + \Gamma_t + \Gamma_u = 0$ as Lie polynomials, we must also have that

$$N_{\Gamma_s} + N_{\Gamma_t} + N_{\Gamma_u} = 0. \quad (2.12)$$

For this reason, the numerators are not uniquely determined: given a triple Γ_s , Γ_t and Γ_u we can perform the shift $\delta(N_{\Gamma_s}, N_{\Gamma_t}, N_{\Gamma_u}) = (s, t, u)A$ for any $A \in V$. It also follows that BCJ numerators can be determined from their values on a comb basis $N_{\Gamma_{1a}}$ by

$$N_{\Gamma} = \sum_a (\Gamma, 1a) N_{\Gamma_{1a}}. \quad (2.13)$$

In the case of Yang mills, the claim of BCJ [1] is that

$$\mathcal{A} = \sum_{\Gamma} \frac{N_{\Gamma}^{k,\epsilon} c_{\Gamma}}{d_{\Gamma}} \quad (2.14)$$

for some *kinematic numerators* $N_{\Gamma}^{k,\epsilon}$ depending linearly on each polarization vector ϵ_i and rationally (or even polynomially) on the momenta satisfying (2.12). The key nontrivial output of the double copy is that gravity amplitudes are obtained when $\tilde{N}_{\Gamma} = N_{\Gamma}^{k,\epsilon}$. The same numerators determine both the colour-ordered Yang-Mills amplitude with order a is then

$$\mathcal{A}_a = \sum_{\Gamma} \frac{N_{\Gamma}(\Gamma, a)}{d_{\Gamma}} \quad (2.15)$$

and gravity amplitudes by

$$\mathcal{M} = \sum_{\Gamma} \frac{N_{\Gamma} N_{\Gamma}}{d_{\Gamma}}. \quad (2.16)$$

The most basic theory in this framework is the bi-adjoint scalar theory whose colour ordered amplitudes are given by

$$m(a, b) := \sum_{\Gamma} \frac{(\Gamma, a)(\Gamma, b)}{d_{\Gamma}}. \quad (2.17)$$

and we can introduce two underlying abstract amplitudes for these theories given by

$$m = \sum_{\Gamma} \frac{\Gamma \otimes \Gamma}{d_{\Gamma}} \in Lie(n-1) \otimes Lie(n-1), \quad m(a) = \sum_{\Gamma} \frac{(\Gamma, a)\Gamma}{d_{\Gamma}} \in Lie(n-1). \quad (2.18)$$

Substituting (2.13) into (2.15) we obtain Yang-Mills amplitudes in terms of numerators and $m(a, b)$ by

$$\mathcal{A}_a = (N, m(a)) = \sum_b m(a, 1b) N_{\Gamma_{1b}}^{k, \epsilon}, \quad (2.19)$$

with a similar form for gravity

$$\mathcal{M} = (N \otimes \tilde{N}, m) = \sum_{a, b} m(1a, 1b) N_{1a}^{k, \epsilon} \tilde{N}_{1b}^{k, \tilde{\epsilon}}. \quad (2.20)$$

We briefly remark that the basic kinematic numerators $N_{\Gamma}^{k, \epsilon}$ for Yang Mills were obtained in [13] and other related ones such as $N_{\Gamma}^{k, k}$ can be created by setting $k_i \cdot \epsilon_j = 0$, and $\epsilon_i \cdot \epsilon_j = k_i \cdot k_j$ or $N_{\Gamma}^{\epsilon, k, m}$ by taking some components of the polarization vectors to be in a higher dimension to the momenta. We thereby obtain, amplitudes for a variety of theories as

$$\text{Amplitudes:} \quad \mathcal{M} = \sum_{\Gamma} \frac{N_{\Gamma}^l N_{\Gamma}^r}{d_{\Gamma}}$$

with N_{Γ} 's from in table 1. See [4] for an up-to-date list of available numerators and their details.

$N^l \backslash N^r$	$N_{\Gamma}^{k, \epsilon}$	$N_{\Gamma}^{k, k}$	$N_{\Gamma}^{k, \epsilon, m}$	c_{Γ} or (Γ, a)
$N_{\Gamma}^{k, \epsilon}$	E			
$N_{\Gamma}^{k, k}$	BI	Galileon		
$N_{\Gamma}^{k, \epsilon, m}$	$EM_{U(1)^m}$	DBI	$EMS_{U(1)^m \times U(1)^{\tilde{m}}}$	
c_{Γ} or (Γ, a)	YM	Nonlinear σ	$EYMS_{SU(N) \times U(1)^{\tilde{m}}}$	Biadjoint Scalar $SU(N) \times SU(\tilde{N})$

Table 1. Theories arising from the different choices of numerators.

We cannot simply invert (2.19) to obtain the N_{1b} as $m(1a, 1b)$ is not invertible. Indeed, all theories that can be expressed in this double-copy format with one explicit Lie polynomial factor satisfy the fundamental BCJ relations [1]. There are many forms of the relations, one version being, for a word² $a \in W(n-2)$

$$\sum_{bc=a} s_{i, b|_b} m((i \sqcup b)c) = 0, \text{ where } s_{b,k} := \sum_{i=1}^{|b|} s_{b_i k}. \quad (2.21)$$

Thus (2.19) determines the N_{1a} only up to the addition of multiples of the BCJ relations. This freedom can be used to set all but $(n-3)!$ of the N_{1a} to zero, but this is at the expense of requiring numerators that are rational rather than polynomial in the momenta, so that the remaining numerators will then have spurious poles.

2.4 A note on the Kinematic Algebra

Given a Lie algebra, g , with an inner product, a Lie monomial $\Gamma \in Lie(n-1)$ gives rise to a ‘colour factor,’ c_Γ , for every n -tuple of Lie algebra elements, T_1, \dots, T_n ,

$$c_\Gamma := \text{tr}(\Gamma[T_1, \dots, T_{n-1}]T_n).$$

This means we have a map

$$c : Lie(n-1) \rightarrow \otimes^n g^*,$$


and it is a homomorphism, since the c_Γ clearly satisfy the Jacobi identity. The BCJ kinematic numerators for Yang-Mills, $N_\Gamma^{k,\epsilon}$, likewise satisfy the Jacobi identity leading to a suggestion that they might arise from some *kinematic algebra*, an as yet unidentified Lie algebra. If that were the case, the double copy would be replacing the Yang-Mills Lie algebra numerator c_Γ with kinematic numerators $N_\Gamma^{k,\epsilon}$. However (2.12) does not imply that there is a Lie algebra, g , such that N_Γ is the colour factor for that Lie algebra. This is clear for example in the case of (Γ, a) (and the forms w_Γ below). Nevertheless, there has been some interesting work to identify such a Lie algebra associated to the kinematic numerators, [15–17], (2.12).

In general, a homomorphism from $Lie(n-1)$ to a vector space of functions can be given by choosing any $(n-2)!$ such functions, as in equation (2.13). These have been identified in the case of Yang-Mills by various authors by recursion and in particular for Yang Mills in [13].

3 Trees and words in $\mathcal{M}_{0,n}$

In this section we consider the homology and cohomology of $\mathcal{M}_{0,n}$, the Deligne-Mumford compactification of the space of n distinct points on the Riemann sphere \mathbb{CP}^1 up to Möbius transformations. We first recall the basic properties of $\mathcal{M}_{0,n}$. $\mathcal{M}_{0,n}$ has a normal crossing divisor D , whose top dimensional strata are the codimension one components D_I , for $I \subset \{1, \dots, n\}$, where the points in I bubble off onto a new \mathbb{CP}^1 , attached to the first by a node.

²This is eq 3.8 of [14] together with the use of the $U(1)$ -decoupling identity in the bracketed term.



Two such components, D_I and D_J , intersect iff $I \subset J$ or $I \subset \bar{J}$. Then $D_I \cap D_J$ corresponds to nodal curves with 3 components; for example when $I \subset J$, one containing the points I , the second $J - I$ and the third \bar{J} . It follows that the maximal intersections of these D_I are points, $D_\Gamma \in \mathcal{M}_{0,n}$, given by the intersection of $n - 3$ compatible D_{I_p} . Each such point correspond to a nodal curve with $n - 3$ components, each with three points that are either nodes or marked points. Such a tuple of compatible sets I_p defines a trivalent tree, Γ , with the components corresponding to the vertices and nodes to propagators. The 0-dimensional strata of D are thus in one-to-one correspondence with trivalent (and unoriented) trees.

The complement of the divisor in $\mathcal{M}_{0,n}$ is an open top cell, $\mathcal{M}_{0,n}^\# := \mathcal{M}_{0,n} - D$, on which we can use simplicial coordinates σ_i , $i = 1, \dots, n$, with the gauge fixing $(\sigma_1, \sigma_{n-1}, \sigma_n) = (0, 1, \infty)$. However, in order to study $\mathcal{M}_{0,n}$ in the neighborhood of the divisor, it is useful to introduce dihedral coordinates, which are a set of cross-ratios of the points. There is one such set of coordinates for every dihedral structure [18]. Given an ordering, a , the associated dihedral coordinates are the cross-ratios

$$u_{ij} := (a_i a_{j-1} | a_{i-1} a_j) = \frac{\sigma_{a_i a_{j-1}} \sigma_{a_{i-1} a_j}}{\sigma_{a_i a_j} \sigma_{a_{i-1} a_{j-1}}}, \quad (3.1)$$

for $1 < i + 1 < j$ and $\sigma_{ij} = \sigma_i - \sigma_j$. Each such cross-ratio u_{ij} is associated to the cord, (a_i, a_{j+1}) , of the n -gon labelled by the ordering, a . We will also denote u_{ij} by u_I , where $I \subset \{1, \dots, n\}$ is the subset $\{a_i, \dots, a_{j-1}\}$ or its complement.

If the ordering a is compatible with the set $I \subset \{1, \dots, n\}$, then the divisor component D_I can be seen in these coordinates as the locus of $u_I = 0$ [18]. The points D_Γ in the divisor are, in dihedral coordinates, given as follows. Choose any ordering a such that Γ is planar for a (i.e. $(\Gamma, a) \neq 0$), with propagators given by the subsets $I_p \subset \{1, \dots, n - 1\}$. In the dihedral coordinates associated to a , there are $n - 3$ cross-ratios u_{I_p} corresponding to the propagators of Γ . The point D_Γ is then given, in these coordinates, by $u_{I_p} = 0$ for all $p = 1, \dots, n - 3$.

The u_{I_p} form a good set of coordinates near Γ . Relations between such coordinate systems near different such points are obtained from the non-crossing identity

$$u_{ij} = 1 - \prod_{(k,l) \in (i,j)^c} u_{kl}, \quad (3.2)$$

where for $k < l$, $(k, l) \in (i, j)^c$ means that the diagonal (k, l) of the polygon with vertices $\{1, \dots, n\}$ crosses the diagonal (i, j) .

3.1 The cohomology of $\mathcal{M}_{0,n}$ and Parke-Taylor

The dimensions of the cohomology groups of $\mathcal{M}_{0,n}$ are given by the Poincaré polynomial³

$$P(t) := \sum_i \dim(H^i(\mathcal{M}_{0,n}^\#)) t^i = \prod_{k=2}^{n-2} (1 + kt). \quad (3.3)$$

The cohomology ring is generated by the $d \log \sigma_{ij}$ in the standard gauge fixing, subject to the quadratic relations

$$\frac{d\sigma_{ij}}{\sigma_{ij}} \wedge \frac{d\sigma_{jk}}{\sigma_{jk}} + \frac{d\sigma_{jk}}{\sigma_{jk}} \wedge \frac{d\sigma_{ki}}{\sigma_{ki}} + \frac{d\sigma_{ki}}{\sigma_{ki}} \wedge \frac{d\sigma_{ij}}{\sigma_{ij}} = 0. \quad (3.4)$$

This gives the dimension of $\Gamma(\Omega_D^1)$ as $\sum_{k=2}^{n-2} = n(n-3)/2$ as claimed earlier. It also follows that the top cohomology $H^{n-3}(\mathcal{M}_{0,n}, D) \simeq \Gamma(\mathcal{M}_{0,n}, \Omega_D^{n-3})$ has dimension $(n-2)!$. A natural spanning set for $\Gamma(\Omega_D^{n-3})$ is provided by the Parke-Taylor forms

$$PT(123 \dots n) = \frac{1}{\text{Vol}SL(2)} \bigwedge_{i=1}^n \frac{d \log \sigma_{i i+1}}{2\pi i}. \quad (3.5)$$

In our gauge fixing

$$\frac{1}{\text{Vol}SL(2)} = \frac{(2\pi i)^3 \sigma_{1n-1} \sigma_{n-1n} \sigma_{n1}}{d\sigma_1 d\sigma_{n-1} d\sigma_n} \quad (3.6)$$

yielding now for a general choice of permutation a of $1, \dots, n-1$

$$PT_a := PT(an) = \frac{d^{n-3} \sigma}{\prod_{i=1}^{n-2} \sigma_{a(i) a(i+1)}}, \quad d^{n-3} \sigma := \frac{1}{(2\pi i)^{n-3}} \bigwedge_{i=2}^{n-2} d\sigma_i. \quad (3.7)$$

The $(n-1)!$ Parke-Taylor forms defined in this way are not linearly independent, because they satisfy the shuffle relations of 2.4,

$$PT_{b \sqcup c} = 0, \quad (3.8)$$

for b, c nontrivial, identically⁴ [21]. Thus, following Proposition 2.4, we deduce that

$$H^{n-3}(\mathcal{M}_{0,n}, D) \simeq \Gamma(\mathcal{M}_{0,n}, \Omega_D^{n-3}) \simeq \text{Lie}(n-1)^*. \quad (3.9)$$

Moreover, by Lemma 2.2, one can take a KK basis for $H^{n-3}(\mathcal{M}_{0,n}, D)$, given by the PT_{1a} , for all $(n-2)!$ permutations a . It follows from equation (2.4) that we have the following identity.

³Arnol'd [19] computes the Poincaré polynomial of the cohomology of the configuration space M_{n-1} of $n-1$ points in \mathbb{C} . This can be obtained inductively via the fibration $M_k \rightarrow M_{k-1}$ with fibre $\mathbb{C} - \{\sigma_1, \dots, \sigma_{k-1}\}$ giving factors of $(1 + (k-1)t)$. To get $\mathcal{M}_{0,n}^\#$, one needs to take the quotient of M_{n-1} by $\mathbb{C}^* \ltimes \mathbb{C}$, the stabilizer in PSL_2 of the point at infinity. Arnol'd's formula for the Poincaré polynomial of M_{n-1} must therefore be divided by $1+t$ to obtain the formula here.

⁴We thank Carlos Mafra for this reference and also refer the reader to sections 4.1. and 4.2 of [20] for further discussion.

Lemma 3.1 *The forms PT_a satisfy*

$$PT_a = \sum_b (\Gamma_{1b}, a) PT_{1b}.$$

For future reference, observe that we can write PT_{1a} in dihedral coordinates as

$$PT_{1a} = \text{sgn}(a) \bigwedge_{i=3}^{n-1} \frac{1}{2\pi i} d \log \frac{u_{1a_i}}{1 - u_{1a_i}}, \quad (3.10)$$

since, in the standard gauge fixing,

$$d \log \frac{u_{1a_i}}{1 - u_{1a_i}} = d\sigma_{a_{i-1}} \frac{\sigma_{1a_i}}{\sigma_{1a_{i-1}} \sigma_{a_{i-1}a_i}} + d\sigma_{a_i} \frac{1}{\sigma_{a_{i-1}a_i}}.$$

In fact, for any tree Γ compatible with the ordering a we have an associated top form,

$$PT_\Gamma := \bigwedge_{i=1}^{n-3} \frac{1}{2\pi i} d \log \frac{u_{I_p}}{1 - u_{I_p}}, \quad (3.11)$$

where the I_p are (some ordering of) the subsets defining the propagators of Γ . In particular, $PT_{\Gamma_{1a}} = PT_{1a}$. PT_Γ is not always equal to a standard Parke-Taylor for the given dihedral structure. For the ordering $1a$, an example of a tree, Γ , such that $PT_\Gamma \neq PT_{1a}$ is the ‘snowflake’, $\Gamma = [[[1, 2], [3, 4]], 5]$, for which one finds

$$PT_\Gamma = PT_{12345} \times (13|24).$$

However, for a tree Γ in which every vertex is attached to at least one external particle, PT_Γ does give rise to the ordinary Parke-Taylor for that dihedral structure, and these expressions for PT_Γ , in dihedral coordinates, were first considered and used by Koba and Nielsen in [22].

Note also that $\Gamma \mapsto PT_\Gamma$ is not a homomorphism from $Lie(n-1)$ to $H^{n-3}(\mathcal{M}_{0,n}, D)$, since the PT_Γ do not satisfy the Jacobi identity, which in particular implies that equation (2.5) cannot be used to expand PT_Γ in a basis of PT_a ’s.

3.2 Homology of $\mathcal{M}_{0,n}$

We saw, in equation (3.9), that there is an isomorphism of $H^{n-3}(\mathcal{M}_{0,n}, D)$ with $Lie(n-1)^*$. Integration gives a perfect pairing between relative cohomology, $H^{n-3}(\mathcal{M}_{0,n}, D)$, and the homology of the complement of the divisor, $H_{n-3}(\mathcal{M}_{0,n} - D)$ [23]. It follows that the homology $H_{n-3}(\mathcal{M}_{0,n} - D)$ can be identified (as a vector space) with $Lie(n-1)$.⁵ In this section, we will review the description of the homology cycles of $\mathcal{M}_{0,n} - D$.

A point D_Γ naturally gives rise to a class in $H_{n-3}(\mathcal{M}_{0,n}^\#)$ represented by a real half-dimensional torus that surrounds D_Γ . Explicitly, the cycle can be defined as the locus

$$C_\Gamma : |u_{I_p}| = \epsilon_p, \quad p = 1, \dots, n-3 \quad (3.12)$$

⁵There is a free action of the symmetric group S_n on $H_{n-3}(\mathcal{M}_{0,n} - D)$. As a module for S_n , $H_{n-3}(\mathcal{M}_{0,n} - D)$ is not isomorphic to $Lie(n-1)$, but $Lie(n-1)$ twisted by the sign representation of S_n .

for some small ϵ_p and choice of orientation. These cycles were first described in [24], and they generate the homology, but are not independent.⁶ Integrating a holomorphic top-form with C_Γ evaluates the residue at D_Γ [26]. Lemma 7.1 of [18] states that PT_a can only have a pole at those D_Γ which are compatible with the ordering a . Moreover, it is clear that we can orient the C_{Γ_a} so that

$$\int_{C_{\Gamma_a}} PT_a = +1,$$

which leads to the following,

Lemma 3.2 *The cycles $C_{\Gamma_{1a}}$ represent a basis for $H_{n-3}(\mathcal{M}_{0,n}^\#)$, dual to the KK basis of $H^{n-3}(\mathcal{M}_{0,n}, D)$:*


$$\int_{C_{\Gamma_{1a}}} PT_{1b} = (1a, 1b),$$

where we have chosen to orient the $C_{\Gamma_{1a}}$ using the PT_{1a} forms so that the residues are +1.

The relations among the top cycles are given by Jacobi-type relations.

Lemma 3.3 (Cohen⁷) *For three trees related by Jacobi, there exists a contraction of the sum of the corresponding cycles, $C_{\Gamma_s} + C_{\Gamma_t} + C_{\Gamma_u}$.*

An explicit homotopy contracting $C_{\Gamma_s} + C_{\Gamma_t} + C_{\Gamma_u}$ is easy to visualize for $n = 4$, since $\mathcal{M}_{0,4}^\# = \mathbb{CP}^1 - \{0, 1, \infty\}$, with boundary points $D_s = 0$, $D_t = 1$, and $D_u = \infty$. Appropriately oriented, the three small circles around these points add up to zero in homology.



More generally, for $n > 4$, we can exhibit a contraction of $C_{\Gamma_s} + C_{\Gamma_t} + C_{\Gamma_u}$ by making small all ϵ 's in definition of the cycle, equation (3.12), except for the ϵ corresponding to the propagator being exchanged. This restricts us to a \mathbb{CP}^1 component of D , where the same argument made for $n = 4$ can be applied.⁸

It follows from Lemma 3.3 and equation (2.5) that,

$$C_\Gamma = \sum_a (\Gamma, 1a) \Gamma_{1a}.$$

Combining this with Lemma 3.1 and Lemma 3.2 implies,

⁶The homology of configuration spaces was first computed by [25], but the cycles are described by Cohen [24]. Cohen was studying the homology of configuration spaces. To apply his results here one uses the isomorphism $\mathcal{M}_{0,n} - D \simeq \text{Conf}_{n-3}(\mathbb{R}^2 - \{0, 1\})$.

⁷This is a consequence of Theorem 1.2 in [24], as reviewed in [27].

⁸The maximum dimension tori appearing in Lemma 3.3 can be constructed by taking the wedge products of lower dimensional tori. This leads to a natural product in the full homology $H_*(\mathcal{M}_{0,n} - D)$, under which H_* can be identified with a graded Lie algebra, with graded Jacobi relations holding in all dimensions. [24] The (ungraded) Jacobi relation in Lemma 3.3 is a special case. [24, 27]

Lemma 3.4 *The integration pairing between C_Γ and PT_a is*

$$\int_{C_\Gamma} PT_a = (\Gamma, a).$$

4 A Penrose transform for amplitudes

Our starting point is the observation that \mathcal{K}_n can be identified with $H^1(\mathcal{M}_{0,n}, D)$. Our ‘twistor space’ for the Penrose transform will be $\mathbb{T} = T_D^* \mathcal{M}_{0,n}$, the total space of the bundle of holomorphic 1-forms on $\mathcal{M}_{0,n}$ with logarithmic singularities on D . The relationship with \mathcal{K}_n is given by the isomorphism

$$\mathcal{K}_n := H^1(\mathcal{M}_{0,n}, D) \simeq \Gamma(\mathcal{M}_{0,n}, T_D^*). \quad (4.1)$$

This correspondence can be expressed by considering $d \log$ of the Koba-Nielsen factor [22], which is, in the standard gauge fixing,

$$KN := \prod_{i < j} \sigma_{ij}^{s_{ij}}, \quad \sigma_{ij} = \sigma_i - \sigma_j. \quad (4.2)$$

This gives the general section of $\tau \in \Gamma(T_D^* \mathcal{M}_{0,n})$ as

$$\tau = \sum_i E_i d\sigma_i := \sum_{i < j} s_{ij} d \log \sigma_{ij}, \quad E_i = \sum_j \frac{s_{ij}}{\sigma_{ij}}. \quad (4.3)$$

The second equality shows that this is clearly invariant under translations and rescalings of the σ_i , but full Mobius invariance (i.e., vanishing when contracted with $\sum_i \sigma_i^2 \partial \sigma_i$) requires $\sum_i s_{ij} = 0$. Our normalizations $(\sigma_1, \sigma_{n-1}, \sigma_n) = (0, 1, \infty)$ gives the triviality of $d \log \sigma_{in}$ and $d \log \sigma_{1n-1}$ giving the correct dimensionality of the $d \log \sigma_{ij}$ basis of H^1 . Note that the equations $E_i = 0$ are the *scattering equations*.

To more clearly demonstrate the $d \log$ behaviour on D , given a choice of the standard ordering, we can also represent the Koba Nielsen factor as [28]

$$KN = \prod_{j > i+1} u_{ij}^{X_{ij}}, \quad X_{ij} = \sum_{i \leq l < m < j} s_{lm}. \quad (4.4)$$

This gives the useful representation of the general section in terms of the $n(n-3)/2$ basis $d \log u_{ij}$

$$\sum_i E_i d\sigma_i = \sum_{j > i+1} X_{ij} d \log u_{ij}. \quad (4.5)$$

This representation manifests the $d \log$ behaviour on the components of D compatible with this choice of ordering.

4.1 The double fibration and the CHY formulae

The twistor correspondence arises from the following double fibration:

$$\begin{array}{ccc} \mathcal{Y}_n = \mathcal{K}_n \times \mathcal{M}_{0,n}, & (s_{ij}, \sigma_j) & \\ p \swarrow & & \searrow q \\ (s_{ij}), \mathcal{K}_n & & \mathbb{T} = T_D^* \mathcal{M}_{0,n}, (\tau_i, \sigma_i). \end{array} \quad (4.6)$$

where p forgets the second factor and q is defined by the incidence relations

$$\tau_i = E_i(s_{kl}, \sigma_m) := \sum_j \frac{s_{ij}}{\sigma_{ij}}. \quad (4.7)$$

A point in \mathcal{K}_n therefore determines a section $\tau_i = E_i$ of $\mathbb{T} \rightarrow \mathcal{M}_{0,n}$.

A special role is played by the zero-section \mathbb{T}_0 of \mathbb{T} as it encodes the scattering equations; given generic s_{ij} , the section $\tau_i = E_i(\sigma)$ intersects \mathbb{T}_0 at the $(n-3)!$ solutions to the scattering equations. We therefore introduce $\bar{\delta}(\tau)^{n-3}$ to be the $(0, n-3)$ -form delta function supported on \mathbb{T}_0 . In the standard gauge fixing above it can be defined by

$$\bar{\delta}(\tau)^{n-3} := \prod_{i=2}^{n-2} \bar{\partial} \frac{1}{2\pi i \tau_i}, \quad \text{where} \quad \bar{\partial} \frac{1}{2\pi i z} = \delta(\Re z) \delta(\Im z) d\bar{z}. \quad (4.8)$$

More invariantly, this takes values in $(\Omega_D^{n-3} \mathcal{M}_{0,n})^*$ so that to use it in an integrand, we will need an extra factor with values in $(\Omega_D^{n-3} \mathcal{M}_{0,n})^2$.

A first observation is that the CHY formulae can be regarded as examples of a Penrose transform in the sense that the amplitudes are obtained as the pushdown to \mathcal{K}_n of a pullback of an object from \mathbb{T} . The generic CHY formula takes the form:

$$\mathcal{M}(s_{ij}, \dots) = \int_{\mathcal{M}_{0,n=p^{-1}(s_{ij})}} q^* (\mathcal{I}_l \mathcal{I}_r \bar{\delta}(\tau)^{n-3}) \quad (4.9)$$

Here $\mathcal{I}_l, \mathcal{I}_r \in \Omega_D^{n-3} \mathcal{M}_{0,n}$ are CHY half-integrands but also often depending also on polarization data, with the most basic example being $m(a, b)$ when $(\mathcal{I}_l, \mathcal{I}_r) = (PT_a, PT_b)$. There is an empirical direct correspondence between choices of $\mathcal{I}_{l/r}$ and numerators N_Γ with for example the CHY Pfaffian⁹ $\text{Pf}'(M)$ corresponding to the $N_\Gamma^{k, \epsilon}$ described earlier. See [29, 30] for details of half-integrands for other theories and their origins.

⁹For completeness, the CHY Pfaffian is the reduced Pfaffian of the matrix M

$$M = \begin{pmatrix} A & C \\ -C^t & B \end{pmatrix}, \quad A_{ij} = \frac{k_i \cdot k_j}{\sigma_i - \sigma_j}, \quad B_{ij} = \frac{\epsilon_i \cdot \epsilon_j}{\sigma_i - \sigma_j}, \quad C_{ij} = \frac{k_i \cdot \epsilon_j}{\sigma_i - \sigma_j}, \quad \text{for } i \neq j$$

constructed from polarization vectors ϵ_i with $k_i \cdot \epsilon_i = 0 \dots$, and $A_{ii} = B_{ii} = 0$, $C_{ii} = \epsilon_i \cdot \sum k_i / \sigma_{ij}$. This matrix is degenerate, but removing two rows and columns, say ij , taking the Pfaffian and dividing by σ_{ij} yields a well-defined half-integrand.

4.2 The geometry of the correspondence

A generic point (τ_i, σ_i) of \mathbb{T} corresponds to a codimension- $n-3$ plane in \mathcal{K}_n . This plane is the $(n-2)(n-3)/2$ dimensional space of sections that pass through the point. For a point lying in a top-stratum, D_I , of the divisor, these planes lie inside the factorisation hyperplane plane $s_I = 0$. This follows from the following combination [31] of the scattering equations

$$E_I = \frac{s_I}{2} + \sum_{i \in I, j \in \bar{I}} s_{ij} \frac{\sigma_{i1} \sigma_{jn}}{\sigma_{ij} \sigma_{1n}}. \quad (4.10)$$

and the fact that, assuming $1 \in I$ and $n \in \bar{I}$, the second term vanishes when restricted to D_I .

It follows that the point $(\tau_i, \sigma_i) = (0, D_\Gamma) \in \mathbb{T}$ corresponds to the codimension- $n-3$ plane in \mathcal{K}_n given by the intersection of the planes $s_{I_p} = 0$, where I_p are the subsets of $\{1, \dots, n-1\}$ corresponding to the momentum flowing through each propagator in Γ . We can characterize these planes as being those planes passing through the origin with normal $n-3$ -form

$$w_\Gamma = \pm \bigwedge_{p=1}^{n-3} ds_{I_p}. \quad (4.11)$$

In the next section, we will see how the w_Γ arise from the double fibration and show how their signs are fixed to make them satisfy the same relations as the $n-3$ -forms defined by [6].

4.3 The symplectic form and the holomorphic volume form

By studying the symplectic volume form ω^{n-3} on $T_D^* \mathcal{M}_{0,n}$, we find two elementary consequences of the double fibration. In this section, we describe how the symplectic volume gives rise to a transform between $n-3$ cycles in $\mathcal{M}_{0,n} - D$ and the $n-3$ -forms, equation (??), encountered above. In the next section we will describe the associated correspondence between $(n-3)$ -planes in \mathcal{K}_n and $(n-3)$ -cycles in $H^{n-3}(\mathcal{M}_{0,n}, D)$. Distinguished classes in H^{n-3} are seen to correspond to the planes defined by [6].

The symplectic form on $T_D^* \mathcal{M}_{0,n}$ can be written explicitly as

$$\omega = \sum_{i \neq 1, n-1, n} d\tau_i \wedge d\sigma_i,$$

where τ_i are the components of $\tau = \sum \tau_i d\sigma_i$ in these coordinates. Pulling back ω^{n-3} to \mathcal{Y}_n , we can decompose it into a sum over a basis of $\Gamma(\mathcal{M}_{0,n}, \Omega_D^{n-3})$ with coefficients given by $n-3$ -forms on \mathcal{K}_n . This gives rise to a correspondence between $n-3$ -forms on \mathcal{K}_n and $(n-3)$ -cycles in $\mathcal{M}_{0,n}$ which we explain in this section. In the next section, we describe the associated correspondence between $(n-3)$ -planes in \mathcal{K}_n ('ABHY planes') and $(n-3)$ -cocycles in $H^{n-3}(\mathcal{M}_{0,n}, D)$.

Every cycle C_Γ in $H_{n-3}(\mathcal{M}_{0,n} - D)$ defines an $n-3$ -form on \mathcal{K}_n ,

$$w_\Gamma := \int_{C_\Gamma} q^* \omega^{n-3} \in \Omega^{n-3}(\mathcal{K}_n). \quad (4.12)$$

It is clear that, for a Jacobi triple of trees,

$$w_{\Gamma_s} + w_{\Gamma_t} + w_{\Gamma_u} = 0. \quad (4.13)$$

In other words, the map $Lie(n-1) \hookrightarrow \wedge^{n-3} \mathcal{K}_n^*$ given by $\Gamma \mapsto w_\Gamma$ is a homomorphism.

To find an explicit expression for w_Γ , we use the representation (4.5) of $q^*\tau$ in a choice of dihedral coordinates u_{ij} for which Γ is planar. This gives

$$q^*\omega = d(q^*\tau) = \sum_{i+1 < j} dX_{ij} \wedge \frac{du_{ij}}{u_{ij}} \quad (4.14)$$

It is then easily seen that integration of $q^*\omega^{n-3}$ over C_Γ picks out only those poles that correspond to propagators of Γ . The residue thus gives a wedge product $\wedge ds_{I_p}$, with the overall sign determined by the orientation of C_Γ . Given this explicit form of w_Γ , the Jacobi relation, equation (4.13), can also be understood to follow from the momentum conservation relation, $ds + dt + du = 0$ [6].

It follows from equation (2.5) that w_Γ admits the expansion

$$w_\Gamma = \sum_{a \in S_{N-2}} w_{\Gamma_{1a}}(\Gamma, 1a). \quad (4.15)$$

Combining this with Lemmas 3.4 and 2.3, we find that we can write the pull-back of the symplectic volume form as

$$q^*\omega^{n-3} = \sum_{a \in S_{n-2}} w_{\Gamma_{1a}} PT_{1a}. \quad (4.16)$$

Although we have used the dual comb and KK bases, this relation follows in any dual basis, because w_Γ and PT_a furnish representations of $Lie(n-1)$ and $Lie(n-1)^*$ respectively so that (4.16) is an explicit of writing the Kronecker delta. For example, when $n = 4$, $q^*\omega$ can be written in any one of the three KK bases,

$$\begin{aligned} q^*\omega &= -ds_{12} \wedge PT_{213} - ds_{23} \wedge PT_{231} \\ &= ds_{12} \wedge PT_{123} + ds_{13} \wedge PT_{132} \\ &= -ds_{23} \wedge PT_{321} + ds_{13} \wedge PT_{312}. \end{aligned} \quad (4.17)$$

More generally, we can rewrite $q^*\omega^{n-3}$ using whatever bases we choose.

Lemma 4.1 *The pullback of the symplectic volume form to \mathcal{Y}_n can be written*

$$q^*\omega^{n-3} = \sum_{\Gamma \in H, a \in K} (\Gamma, a) w_\Gamma \wedge PT_a.$$

for a basis H of $Lie(n-1)$, and a basis K of $Lie(n-1)^*$.




Figure 1.

4.4 Associahedral $(n-3)$ -planes in \mathcal{K}_n and forms on $\mathcal{M}_{0,n}$.

An alternative way to study the correspondence is to restrict ω^{n-3} to different $(n-3)$ -planes in \mathcal{K}_n . This correspondence is related to the construction in [32], although they discuss different planes and polytopes. Distinguished classes in $H^{n-3}(\mathcal{M}_{0,n}, D)$ correspond to the associahedral planes found in [6].

An $n-3$ -plane in \mathcal{K}_n is defined up to translation by its tangent form $P \in \wedge^{n-3}\mathcal{K}_n$. For any such plane, P , the symplectic volume form gives rise to an $n-3$ -form in $\Gamma(\mathcal{M}_{0,n}, \Omega_D^{n-3})$,

$$P \lrcorner q^* \omega^{n-3}.$$

Recall the definition of $PT_\Gamma \in \Gamma(\mathcal{M}_{0,n}, \Omega_D^{n-3})$, given in equation (3.11), which is a top form associated to a dihedral structure, a , together with a tree Γ that is compatible with that ordering. PT_Γ corresponds to a plane P_Γ in \mathcal{K}_n defined as follows. Fix a dihedral structure a and let

$$D_I := \frac{\partial}{\partial X_I} - \sum_{J \in I^c} \frac{\partial}{\partial X_J},$$

such that we have

$$D_I \log KN = \log u_I - \sum_{J \in I^c} \log u_J = \log \frac{u_I}{1 - u_I}, \quad (4.18)$$

by the non-crossing identity, equation (3.2). It follows that, for a compatible set of propagators I_p , defining a tree Γ that is compatible with a , we can define a plane

$$P_\Gamma = \bigwedge_{p=1}^{n-3} D_{I_p},$$

for some ordering of the propagators chosen so that

$$P_\Gamma \lrcorner q^* \omega^{n-3} = q^* P T_{a,\Gamma}. \quad (4.19)$$

In particular, when we take $P_{a,\Gamma}$ to be the comb, $P_{1a} := P_{1a,\Gamma_{1a}}$, we recover the Parke-Taylor factors in the standard KK basis,

$$P_{1a} \lrcorner q^* \omega^{n-3} = q^* P T_{1a}.$$

For a general Γ , in dihedral structure a , the equations of the plane associated to P_Γ can be written as

$$X_J + \sum_{r \text{ s.t. } J \in I_r^c} X_{I_r} = \text{const.}, \quad (4.20)$$

for all J compatible with the ordering a and not corresponding to a propagator of Γ . These equations bear no obvious resemblance to those defined in [6], but we will see that our planes are the same as theirs.

First, notice that the P_a satisfy

$$P_a = \sum (\Gamma_{1b}, a) P_{1b},$$

by Lemma 3.1, and that, moreover,

$$(w_\Gamma, P_a) = \int_{C_\Gamma} P T_a = (\Gamma, a),$$

using Lemma 3.4.

In [6], the ABHY planes are defined, for the ordering $a = 1 \dots n-1$ and the comb Γ_a , to be defined by the $(n-2)(n-3)/2$ equations

$$s_{ij} = \text{const.},$$

for $1 \leq i < j-1 < n-2$. Using the identity, $s_{ij} = X_{ij} + X_{i+1j+1} - X_{ij+1} - X_{i+1j}$ we can verify that, for I_p in Γ_a ,

$$D_{I_p} s_{ij} = 0, \quad (4.21)$$

for all $1 \leq i < j-1 < n-2$, and each p . Fixing an s_{ij} , one can check this equation for each p . The chord I_p is the arc kn for $k = 2, \dots, n-2$. The five cases to check are (i) $k \leq i$, (ii) $k = i+1$, (iii) $i+1 < k < j$, (iv) $k = j$, and (v) $k > j$. That equation (4.21) vanishes in these five cases is easily seen from Figure 1. Conversely, we can define the ABHY plane for the ordering a and the comb Γ_a to be given by the tangent form

$$P_a = \bigwedge_{i=2}^{n-2} \left(\frac{\partial}{\partial s_{a_{i-1} a_i}} - \frac{\partial}{\partial s_{a_i a_{i+1}}} \right). \quad (4.22)$$




Figure 2.

For each factor in P_a we can check that it annihilates the expressions in equation (4.20). We have the following property

$$\left(\frac{\partial}{\partial s_{i-1i}} - \frac{\partial}{\partial s_{ii+1}} \right) X_{jk} = \begin{cases} 1, & i = k - 1, \\ -1, & i = j, \\ 0, & \text{otherwise.} \end{cases} \quad (4.23)$$

For fixed i , there are five cases to check, and these are shown in Figure 2. By these arguments, we conclude that

Lemma 4.2 *The planes $P_{a,\Gamma}$ defined by the correspondence, equation (4.19), are the ABHY planes when Γ_a is the comb for the ordering a .*

Moreover, the arguments that led to this Lemma can be applied to the case of general Γ , which leads to the following

Lemma 4.3 *The planes $P_{a,\Gamma}$ defined by the correspondence, equation (4.19), are given by the $(n-2)(n-3)/2$ equations*

$$s_{ij} = \text{const.},$$

for all chords (ij) compatible with the ordering that are not one of the $n-3$ propagators of Γ .

It is a consequence of the results in [6] that the planes in Lemma 4.3 are ABHY planes, in the sense that they cut out associahedra in the same way as the original planes in [33].

4.5 Scattering forms and CHY

The scattering forms of [6] are defined by the following sum over trees,

$$\Omega_a = \sum_{\Gamma} \frac{w_{\Gamma}(a, \Gamma)}{d_{\Gamma}}, \quad (4.24)$$

where $d_{\Gamma} = \prod_{p=1}^{n-3} s_{I_p}$ and I_p are the propagators of Γ . The (a, Γ) -factor reduces the sum to one over trees that are planar for the ordering a .

The symplectic form discussed above gives rise to the ABHY scattering forms on \mathcal{K}_n from the Dolbeault formula

$$\Omega_a = \int_{p^{-1}(s_{ij})} q^* [\bar{\delta}^{n-3}(\underline{\tau}) \wedge \omega^{n-3} PT_a], \quad (4.25)$$

where

$$\bar{\delta}^{n-3}(\underline{\tau}) = \bigwedge_{i=1}^{n-3} \bar{\delta}(\tau_i), \quad \bar{\delta}(z) = \frac{1}{2\pi i} \bar{\partial} \frac{1}{z}. \quad (4.26)$$

To see that equation (4.25) is equal to equation (4.24), we use Lemma 4.1 to rewrite equation (4.25) as

$$\Omega_a = \sum_b w_{\Gamma_{1b}} \int_{p^{-1}(s_{ij})} q^* [\bar{\delta}^{n-3}(\underline{\tau}) \wedge PT_{1b} PT_a].$$

We recognize that the integral is the CHY formula [7, 8] for $m(a, b)$, defined in equation (2.17). It follows that equation (4.25) can be expanded as

$$\Omega_a = \sum_b \frac{w_{\Gamma_{1b}}(\Gamma, 1b)(\Gamma, a)}{d_{\Gamma}},$$

which is equal to (4.24) by equation (4.15). We conclude that

Proposition 4.4 *The ABHY scattering form Ω_a is given by equation (4.25).*

More generally, given any CHY half-integrand \mathcal{I} , we can define an ABHY scattering form

$$\Omega_{\mathcal{I}} := \int_{p^{-1}(s_{ij})} q^* [\bar{\delta}^{n-3}(\underline{\tau}) \wedge \omega^{n-3} \mathcal{I}]. \quad (4.27)$$

This form admits an expansion

$$\Omega_{\mathcal{I}} = \sum_a w_{\Gamma_{1a}} \int_{p^{-1}(s_{ij})} q^* [\bar{\delta}^{n-3}(\underline{\tau}) PT_{1a} \mathcal{I}].$$

The CHY integral as given is not changed by adding to \mathcal{I} a term that vanishes on the support of the scattering equations. One can regard \mathcal{I} as a representative of a twisted cohomology class, as explained in [34].

All scattering forms, $\Omega_{\mathcal{I}}$, obtained in this way are projective. Let Υ be the Euler vector field on \mathcal{K}_n ,

$$\Upsilon = \sum_{i < j} s_{ij} \frac{\partial}{\partial s_{ij}}.$$

We can lift Υ trivially to \mathcal{Y}_n using the product structure. On objects pulled back from \mathbb{T} it then acts by

$$\Upsilon = \sum_i \Upsilon(\tau_i) \frac{\partial}{\partial \tau_i} = \sum_i \tau_i \frac{\partial}{\partial \tau_i}. \quad (4.28)$$

Contracting Υ into $\Omega(a)$ we find

$$\Upsilon \lrcorner \Omega_{\alpha}^{(n-3)} = \int q^* \left[\left(\sum_i \tau_i \frac{\partial}{\partial \tau_i} \right) \lrcorner \bar{\delta}^{n-3}(\underline{\tau}) \wedge \omega^{n-3} PT_a \right] = 0,$$

on account of the delta functions in the integrand.

Lemma 4.5 *The scattering form $\Omega_{\mathcal{I}}$ is a projective form on \mathbb{PK}_n .*

Finally, amplitudes are obtained by restricting the scattering forms $\Omega_{\mathcal{I}}$ to ABHY planes, as in [6]. In fact, the restriction of $\Omega_{\mathcal{I}}$ to P_a evaluates to give the CHY formula for the integrands PT_a and \mathcal{I} , as follows from the definition of P_a (equation (4.19) and subsequent lines). Moreover, for a general cohomology class

$$N := \sum_a N_{1a} PT_{1a} \in H^{n-3},$$

we have an associated plane, $P_N := \sum_a N_{1a} P_{1a}$. Pairing this with the general scattering form $\Omega_{\mathcal{I}}$ gives the CHY amplitude for integrands N and \mathcal{I} .

5 Discussion

We have seen that Lie polynomials underpin the color-kinematics and double copy framework of BCJ. We have reviewed the classical fact that the top homology of $\mathcal{M}_{0,n} - D$ is isomorphic to the Lie polynomials, $Lie(n-1)$, and shown that there is a natural correspondence between $T_D^* \mathcal{M}_{0,n}$ and \mathcal{K}_n under which the CHY integral formulae can be understood as a Penrose transform. This can be extended to a transform between the holomorphic Liouville form and the differential forms w_{Γ} introduced by [6], and also between CHY half-integrands and the scattering forms introduced by [6].

One underlying question in the subject is whether there is a kinematic algebra underpinning the kinematic numerators $N_{\Gamma}^{k,\epsilon}$. Although we have seen that the color factors of Lie algebras can provide such numerators, we have also seen many examples of numerators satisfying the Jacobi identity that do not arise as color factors for a Lie algebra: for example, (Γ, a) and w_{Γ} and so on. In other words, the existence of a homomorphism from $Lie(n-1)$ to some vector space does not of itself determine a Lie algebra.

The basic results in this paper can be taken further to yield natural recursions in field theory, which lead to both Lie polynomial and ABHY-form based proofs of the known properties of the field theory momentum kernel and of kinematic numerators. The momentum kernel can be studied also in $T^*\mathcal{M}_{0,n}$, where it arises in the CHY treatment of KLT orthogonality [35]. It also seems likely that the framework will naturally extend to loop integrands in the context of nodal spheres following the logic of [36–40].

The correspondence we have described is suggestive of the naive explicit formula

$$N_{\Gamma}^{\mathcal{I}_l} = \int_{C_{\Gamma}} \mathcal{I}_l, \quad (5.1)$$

for numerators in terms of CHY half-integrands. However, such formulae fail for the CHY Pfaffian that one would expect to give the basic Yang-Mills kinematic numerators; the formula is compromised by the interdependence between in particular reduced Pfaffians and scattering equations. This equation is shown to be invalid as written but a slightly different formulation in a similar spirit is shown to work when \mathcal{I}_l represents a cohomology class in the context of twisted cohomology by Mizera [41].

We remark that the twisted cycle formulation of string integrals in [42] naturally arises in the context of the holomorphic geometric quantization of $T_D^*\mathcal{M}_{0,n}$. To carry out geometric quantization one introduces the line bundle $\mathcal{L} \rightarrow T_D^*\mathcal{M}_{0,n}$ with connection $\nabla = d + \alpha'\tau$ where $\tau = \sum_i \tau_i d\sigma_i$ is the canonical 1-form (symplectic potential) and α' plays the role of Planck's constant. Polarized wave functions should be independent of τ_i . On pull back to the correspondence space, \mathcal{Y}_n , the connection ∇ becomes the standard twisted exterior derivative associated to the Koba-Nielsen factor. Such a quantization of $T^*\mathcal{M}_{0,n}$ perhaps most naturally arises from ambitwistor-string path-integral [43], where the Pfaffian half-integrand for kinematic numerators arises from the RNS spin field path-integral.

There are many further connections to be followed up; we briefly mention the delta algebras of [44] and the formulations of colour-kinematics duality in [45, 46].

6 Acknowledgements

It is a pleasure to acknowledge informative conversations and email exchanges with Nima Arkani-Hamed, Nick early, Yvonne Geyer, Carlos Mafra, Sebastian Mizera, Ricardo Monteiro, Michael Reiterer and Oliver Schlotterer and the hospitality of the CMSA at Harvard. HF would like to acknowledge support from ERC grant Galois theory of periods and applications 724638. While we were writing up this work we became aware of the parallel work of Sebastian Mizera [41] that has some overlap with this—we are grateful to him for letting us see a preview of his work before we finished this paper.

References

- [1] Z. Bern, J. Carrasco, and H. Johansson, *New Relations for Gauge-Theory Amplitudes*, *Phys.Rev.* **D78** (2008) 085011, [[arXiv:0805.3993](#)].

- [2] Z. Bern, J. J. M. Carrasco, and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys.Rev.Lett.* **105** (2010) 061602, [[arXiv:1004.0476](#)].
- [3] H. Kawai, D. C. Lewellen, and S. H. H. Tye, *A Relation Between Tree Amplitudes of Closed and Open Strings*, *Nucl. Phys.* **B269** (1986) 1.
- [4] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson, and R. Roiban, *The Duality Between Color and Kinematics and its Applications*, [arXiv:1909.0135](#).
- [5] M. Kapranov, *Lie operads*, in *The Geometry of Scattering Amplitudes workshop held at Banff in August of 2012*, 2012.
- [6] N. Arkani-Hamed, Y. Bai, S. He, and G. Yan, *Scattering Forms and the Positive Geometry of Kinematics, Color and the Worldsheet*, *JHEP* **05** (2018) 096, [[arXiv:1711.0910](#)].
- [7] F. Cachazo, S. He, and E. Y. Yuan, *Scattering of Massless Particles in Arbitrary Dimensions*, *Phys.Rev.Lett.* **113** (2014), no. 17 171601, [[arXiv:1307.2199](#)].
- [8] F. Cachazo, S. He, and E. Y. Yuan, *Scattering of Massless Particles: Scalars, Gluons and Gravitons*, *JHEP* **1407** (2014) 033, [[arXiv:1309.0885](#)].
- [9] C. Reutenauer, *Free Lie algebras*, vol. 7 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 1993. Oxford Science Publications.
- [10] R. Ree, *Lie elements and an algebra associated with shuffles*, *Annals of Mathematics* **68** (1958), no. 2 210–220.
- [11] D. E. Radford, *A natural ring basis for the shuffle algebra and an application to group schemes*, *Journal of Algebra* **58** (1979), no. 2 432 – 454.
- [12] R. Kleiss and H. Kuijf, *Multigluon cross sections and 5-jet production at hadron colliders*, *Nuclear Physics B* **312** (1989), no. 3 616 – 644.
- [13] C.-H. Fu, Y.-J. Du, R. Huang, and B. Feng, *Expansion of Einstein-Yang-Mills Amplitude*, *JHEP* **09** (2017) 021, [[arXiv:1702.0815](#)].
- [14] F. Cachazo, *Fundamental BCJ Relation in $N=4$ SYM From The Connected Formulation*, [arXiv:1206.5970](#).
- [15] R. Monteiro and D. O’Connell, *The Kinematic Algebra From the Self-Dual Sector*, *JHEP* **07** (2011) 007, [[arXiv:1105.2565](#)].
- [16] R. Monteiro and D. O’Connell, *The Kinematic Algebras from the Scattering Equations*, *JHEP* **03** (2014) 110, [[arXiv:1311.1151](#)].
- [17] G. Chen, H. Johansson, F. Teng, and T. Wang, *On the kinematic algebra for BCJ numerators beyond the MHV sector*, *arXiv:1906.10683 [hep-th]* (June, 2019). [arXiv: 1906.10683](#).
- [18] F. C. S. Brown, *Multiple zeta values and periods of moduli spaces $M_{0,n}(\mathbb{R})$* , *Annales Sci. Ecole Norm. Sup.* **42** (2009) 371, [[math/0606419](#)].
- [19] V. I. Arnol’d, *The cohomology ring of the group of dyed braids*, *Mat. Zametki* **5** (1969) 227–231.
- [20] C. R. Mafra and O. Schlotterer, *Berends-Giele recursions and the BCJ duality in superspace and components*, [arXiv:1510.0884](#).

- [21] J. Cresson, *Calcul moulien*, *Annales de la Faculté des sciences de Toulouse : Mathématiques* **6e série**, **18** (2009), no. 2 307–395.
- [22] Z. Koba and H. B. Nielsen, *Reaction amplitude for n mesons: A Generalization of the Veneziano-Bardakci-Ruegg-Virasoro model*, *Nucl. Phys.* **B10** (1969) 633–655.
- [23] R. Bott and L. W. Tu, *Differential forms in algebraic topology*, vol. 82 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [24] F. Cohen, *Homology of $\Omega^{(n+1)}\Sigma^{(n+1)}X$ and $C_{(n+1)}X$, $n > 0$* , *Bull. Amer. Math. Soc.* **79** (1973) 1236–1241 (1974).
- [25] E. Fadell and L. Neuwirth, *Configuration Spaces.*, *MATHEMATICA SCANDINAVICA* **10** (June, 1962) 111–118.
- [26] P. Griffiths and J. Harris, *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [27] F. R. Cohen, *On configuration spaces, their homology, and Lie algebras*, *J. Pure Appl. Algebra* **100** (1995), no. 1-3 19–42.
- [28] Z. Koba and H. B. Nielsen, *Manifestly crossing invariant parametrization of n meson amplitude*, *Nucl. Phys.* **B12** (1969) 517–536.
- [29] F. Cachazo, S. He, and E. Y. Yuan, *Scattering Equations and Matrices: From Einstein To Yang-Mills, DBI and NLSM*, *JHEP* **07** (2015) 149, [[arXiv:1412.3479](#)].
- [30] E. Casali, Y. Geyer, L. Mason, R. Monteiro, and K. A. Roehrig, *New Ambitwistor String Theories*, *JHEP* **11** (2015) 038, [[arXiv:1506.0877](#)].
- [31] L. Dolan and P. Goddard, *Proof of the Formula of Cachazo, He and Yuan for Yang-Mills Tree Amplitudes in Arbitrary Dimension*, *JHEP* **1405** (2014) 010, [[arXiv:1311.5200](#)].
- [32] S. He, G. Yan, C. Zhang, and Y. Zhang, *Scattering Forms, Worldsheet Forms and Amplitudes from Subspaces*, *JHEP* **08** (2018) 040, [[arXiv:1803.1130](#)].
- [33] V. Bazier-Matte, G. Douville, K. Mousavand, H. Thomas, and E. Yldrm, *ABHY Associahedra and Newton polytopes of \mathbb{F} -polynomials for finite type cluster algebras*, *arXiv:1808.09986 [math]* (Aug., 2018). arXiv: 1808.09986.
- [34] S. Mizera, *Scattering Amplitudes from Intersection Theory*, *Phys. Rev. Lett.* **120** (2018), no. 14 141602, [[arXiv:1711.0046](#)].
- [35] F. Cachazo, S. He, and E. Y. Yuan, *Scattering equations and Kawai-Lewellen-Tye orthogonality*, *Phys. Rev.* **D90** (2014), no. 6 065001, [[arXiv:1306.6575](#)].
- [36] Y. Geyer, L. Mason, R. Monteiro, and P. Tourkine, *Loop Integrands for Scattering Amplitudes from the Riemann Sphere*, *Phys. Rev. Lett.* **115** (2015), no. 12 121603, [[arXiv:1507.0032](#)].
- [37] Y. Geyer, L. Mason, R. Monteiro, and P. Tourkine, *One-loop amplitudes on the Riemann sphere*, *JHEP* **03** (2016) 114, [[arXiv:1511.0631](#)].
- [38] Y. Geyer, L. Mason, R. Monteiro, and P. Tourkine, *Two-Loop Scattering Amplitudes from the Riemann Sphere*, *Phys. Rev.* **D94** (2016), no. 12 125029, [[arXiv:1607.0888](#)].
- [39] Y. Geyer and R. Monteiro, *Gluons and gravitons at one loop from ambitwistor strings*, *JHEP* **03** (2018) 068, [[arXiv:1711.0992](#)].

- [40] S. He, O. Schlotterer, and Y. Zhang, *New BCJ representations for one-loop amplitudes in gauge theories and gravity*, *Nuclear Physics B* **930** (May, 2018) 328–383. arXiv: 1706.00640.
- [41] S. Mizera, *Kinematic Jacobi Identity is a Residue Theorem*, [arXiv:1912.0339](#).
- [42] S. Mizera, *Aspects of Scattering Amplitudes and Moduli Space Localization*. PhD thesis, Perimeter Inst. Theor. Phys., 2019. [arXiv:1906.0209](#).
- [43] L. Mason and D. Skinner, *Ambitwistor strings and the scattering equations*, *JHEP* **1407** (2014) 048, [[arXiv:1311.2564](#)].
- [44] F. Cachazo, N. Early, A. Guevara, and S. Mizera, *Δ -algebra and scattering amplitudes*, *JHEP* **02** (2019) 005, [[arXiv:1812.0116](#)].
- [45] G. Chen, H. Johansson, F. Teng, and T. Wang, *On the kinematic algebra for BCJ numerators beyond the MHV sector*, *JHEP* **11** (2019) 055, [[arXiv:1906.1068](#)].
- [46] M. Reiterer, *A homotopy BV algebra for Yang-Mills and color-kinematics*, [arXiv:1912.0311](#).