

On 2-dimensional expanding attractors of A-flows*

V. Medvedev

E. Zhuzhoma[†]

Abstract

We prove that given any closed n -manifold M^n , $n \geq 4$, there is an A-flow f^t on M^n such that the non-wandering set $NW(f^t)$ consists of 2-dimensional expanding attractor (the both, orientable and non-orientable) and trivial basic sets. For 3-manifolds, we prove that given any closed 3-manifold M^3 , there is an A-flow f^t on M^3 such that the non-wandering set $NW(f^t)$ consists of a non-orientable 2-dimensional expanding attractor and trivial basic sets. Moreover, there is a nonsingular A-flow f^t on a 3-sphere such that the non-wandering set $NW(f^t)$ consists of an orientable 2-dimensional expanding attractor and trivial basic sets (isolated periodic trajectories).

Introduction

A-flows was introduced by Smale [16]. This class of flows contains structurally stable flows including Morse-Smale flows and Anosov flows. Due to Smale's Spectral Theorem, the non-wandering set of A-flow is a disjoint union of closed transitive invariant pieces called basic sets. A basic set is called *trivial* if it is either an isolated fixed point or isolated periodic trajectory. Important type of basic sets are expanding attractors introduced by Williams [17]. A nontrivial basic set Λ is an *expanding attractor* provided Λ is an attractor and its topological dimension coincides with the dimension of unstable manifold at each point of Λ , see definitions at Section 1. Due to Williams [17], an expanding attractor consists of unstable manifolds of its points. Moreover, the unstable manifolds of points of expanding attractor form a lamination with leaves are planes and cylinders. In addition, this lamination locally homeomorphic to the product of Cantor set and Euclidean plane those dimension is more than one. Thus, the minimal topological dimension of expanding attractor for A-flows equals two, and the minimal dimension of supporting manifold equals three. It is natural to study the following question : what manifolds admit A-flows with 2-dimensional expanding attractors ?

In the paper, we solve completely this question for closed manifolds beginning with 3-manifolds. The main results of the paper are the following statements.

Theorem 1 *Given any closed 3-manifold M^3 , there is an A-flow f^t on M^3 such that the non-wandering set $NW(f^t)$ consists of a non-orientable 2-dimensional expanding attractor and trivial basic sets.*

*2000 *Mathematics Subject Classification*. Primary 37D05; Secondary 37B35, 34C40

[†]*Key words and phrases*: A-flow, expanding attractor, basic set

This result contrasts with the case for A-diffeomorphisms. To be precise, it follows from [9, 10, 13, 18] that if a closed 3-manifold M^3 admits an A-diffeomorphism with 2-dimensional expanding attractor, then $\pi_1(M^3) \neq 0$. Note that there are Anosov flows with 2-dimensional expanding attractors [3]. However, Margulis [12] proved that the fundamental group $\pi_1(M^3)$ of a supporting manifold M^3 has an exponential growth.

Theorem 2 *There is a nonsingular A-flow f^t on a 3-sphere such that the non-wandering set $NW(f^t)$ consists of an orientable 2-dimensional expanding attractor and trivial basic sets (isolated periodic trajectories).*

With the help of the results above, we get the final statement.

Theorem 3 *Given any closed n -manifold M^n , $n \geq 4$, there is an A-flow f^t on M^n such that the non-wandering set $NW(f^t)$ consists of 2-dimensional expanding attractor (the both, orientable and non-orientable) and trivial basic sets.*

Acknowledgments. The authors are partially supported by Laboratory of Topological Methods in Dynamics of National Research University Higher School of Economics, of the Ministry of science and higher education of the RF, grant ag. № 075-15-2019-1931.

1 Basic definitions

Basic definitions. Let f^t be a smooth flow on a closed n -manifold M^n , $n \geq 3$. A subset $\Lambda \subset M^n = M$ is *invariant* provided Λ consists of trajectories of f^t . An invariant nonsingular set $\Lambda \subset M$ is called *hyperbolic* if the sub-bundle $T_\Lambda M$ of the tangent bundle TM can be represented as a Df^t -invariant continuous splitting $E_\Lambda^{ss} \oplus E_\Lambda^t \oplus E_\Lambda^{uu}$ such that

- 1) $\dim E_\Lambda^{ss} + \dim E_\Lambda^t + \dim E_\Lambda^{uu} = n$;
- 2) E_Λ^t is the line bundle tangent to the trajectories of the flow f^t ;
- 3) there are $C_s > 0$, $C_u > 0$, $0 < \lambda < 1$ such that

$$\|df^t(v)\| \leq C_s \lambda^t \|v\|, \quad v \in E_\Lambda^{ss}, \quad t > 0, \quad \|df^{-t}(v)\| \leq C_u \lambda^t \|v\|, \quad v \in E_\Lambda^{uu}, \quad t > 0.$$

If $x \in \Lambda$ is a fixed point of hyperbolic Λ , then x is an isolated hyperbolic equilibrium state. The topological structure of flow near x is described by Grobman-Hartman theorem, see for example [14]. In this case $E_x^t = 0$ and $\dim E_\Lambda^{ss} + \dim E_\Lambda^{uu} = n$.

If hyperbolic Λ does not contain fixed points, then the bundles

$$E_\Lambda^{uu} \oplus E_\Lambda^1 = E_\Lambda^u, \quad E_\Lambda^{ss} \oplus E_\Lambda^1 = E_\Lambda^s, \quad E_\Lambda^{uu}, \quad E_\Lambda^{ss}$$

are uniquely integrable [7], [16]. The corresponding leaves

$$W^u(x), \quad W^s(x), \quad W^{uu}(x), \quad W^{ss}(x)$$

through a point $x \in \lambda$ are called *unstable*, *stable*, *strongly unstable*, and *strongly stable manifolds*.

Given a set $U \subset M^n$, denote by $f^{t_0}(U)$ the shift of A along the trajectories of f^t on the time t_0 . Recall that a point x is non-wandering if given any neighborhood U of x and a number T_0 ,

there is $t_0 \geq T_0$ such that $U \cap f^{t_0}(U) \neq \emptyset$. The *non-wandering set* $NW(f^t)$ of f^t is the union of all non-wandering point.

Denote by $Fix(f^t)$ the set of fixed points of flow f^t . Following Smale [16], we'll call f^t an *A-flow* provided its non-wandering set $NW(f^t)$ is hyperbolic and the periodic trajectories are dense in $NW(f^t) \setminus Fix(f^t)$. Well-knows [16, 11] that the non-wandering set $NW(f^t)$ of A-flow f^t is a disjoint union of closed, and invariant, and transitive sets called *basic sets*. Following Williams [17], we'll call a basic set Ω an *expanding attractor* provided Ω is an attractor and its topological dimension equals the dimension of unstable manifold $W^u(x)$ for every points $x \in \Omega$. A basic set Λ is called *orientable* provided the both fiber bundles E_Λ^{ss} and E_Λ^{uu} are orientable. Note that if E_Λ^{ss} and E_Λ^{uu} are one-dimensional, then the orientability of Λ means that the both E_Λ^{ss} and E_Λ^{uu} can be embedded in vector fields on M^n .

2 Proof of main results

We begin with previous results which are interesting itself. Recall that any closed manifold admits a Morse-Smale flow with a source that is a repelling fixed point. In particular, any closed manifold admits a gradient-like Morse-Smale flow having at least one source and sink [15]. The following result says that any closed 3-manifold admits a Morse-Smale flow with one-dimensional repelling periodic trajectory.

Lemma 1 *Given any closed 3-manifolds M^3 , there is a Morse-Smale flow f^t with repelling isolated periodic trajectory on M^3 .*

Proof. Take a gradient-like Morse-Smale flow f_0^t with a sink, say ω , on M^3 . Let $U(\omega)$ be a neighborhood of ω . Without loss of generality, we can suppose that $U(\omega)$ is a ball. Since ω is a hyperbolic fixed point, one can assume that the boundary $\partial U(\omega)$ is a smooth sphere that is transversal to the trajectories. This means that the vector field inducing the flow f_0^t is directed inside of $U(\omega)$. Let us introduce coordinates (x, y, z) and the corresponding cylinder coordinates (ρ, ϕ, z) in $U(\omega)$ smoothly connected with the original coordinates in $U(\omega)$. Consider the system

$$\dot{\rho} = \rho \cdot (1 - \rho), \quad \dot{\phi} = 1, \quad \dot{z} = -z.$$

It easy to check that this system has the attractive hyperbolic one-dimensional trajectory and the saddle fixed point at the origin. This complete the proof. \square

Lemma 2 *There is an A-flow on $S^2 \times S^1$ such that the spectral decomposition of f^t consists of two-dimensional (non-orientable) expanding attractor Λ_a and four isolated hyperbolic repelling trajectories. Moreover, there is a neighborhood P of Λ_a homeomorphic to a solid torus $S^1 \times D^2$ such that the boundary $\partial P = S^1 \times S^1$ is transversal to the trajectories which enter inside of P as a time parameter increases.*

Proof. Take an A-diffeomorphism $f : S^2 \rightarrow S^2$ the spectral decomposition of those consists of a Plykin attractor Λ_0 and four hyperbolic sources. Due to Plykin [13], such diffeomorphism exists. Moreover, well-known that Λ_0 is an expanding attractor. Without loss of generality, one can assume that f is a preserving orientation diffeomorphism (otherwise, one takes f^2).

Let $sus^t(f)$ be the dynamical suspension over f . Since f is an A-diffeomorphism, $sus^t(f)$ is an A-flow. Obviously, the spectral decomposition of f corresponds to the spectral decomposition of

$sus^t(f)$. Because of f preserves orientation, the supporting manifold for $sus^t(f)$ is homeomorphic to $S^2 \times S^1$. Since Λ_0 is a one-dimensional expanding attractor, $sus^t(f)$ has a two dimensional expanding attractor denoted by Λ_a .

Take an isolated hyperbolic repelling trajectory γ that corresponds to some source of f . Since γ is a repelling trajectory, there is a neighborhood $V(\gamma)$ homeomorphic to a solid torus such that the boundary $\partial V(\gamma)$ is transversal to the trajectories of $sus^t(f)$, so that the trajectories move outside of $V(\gamma)$ as a time parameter increases. This follows that the interior of $(S^2 \times S^1) \setminus V(\gamma)$ is the neighborhood, say P , of Λ_a homeomorphic to a solid torus $S^1 \times D^2$ such that the boundary $\partial P = S^1 \times S^1$ is transversal to the trajectories which enter inside of P as a time parameter increases. Hence, $sus^t(f) = f^t$ is a desired flow. \square

PROOF OF THEOREM 1. Let M^3 is a closed 3-manifold. According to Lemma 1, there is a Morse-Smale flow φ^t with repelling isolated periodic trajectory l on M^3 . Hence, there is a neighborhood $U(l)$ of l such that $U(l)$ is homeomorphic to the interior of the solid torus $S^1 \times D^2$, and the boundary $\partial U(l)$ is transversal to the trajectories of φ^t , so that the trajectories move outside of $U(l)$ as a time parameter increases. This follows that $U(l)$ can be replaced by the solid torus P satisfying Lemma 2. As a consequence, one gets the A-flow with two-dimensional (non-orientable) expanding attractor Λ_a . This completes the proof. \square

PROOF OF THEOREM 2. The matrix $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ defines the Anosov diffeomorphism $A : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ of the two-dimensional torus \mathbb{T}^2 . Let $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be a DA-diffeomorphism that is constructed by modifying A [14]. We know that the non-wandering set of f consists of an orientable one-dimensional expanding attractor and isolated source. Denote by $sus^t(f)$ the dynamical suspension over f . Then $sus^t(f)$ is an A-flow with the non-wandering set consisting of the two-dimensional orientable expanding attractor Λ and the isolated repelling periodic trajectory γ . The supporting manifold for $sus^t(f)$ is the mapping torus denoted by T_f^3 . Clearly, that T_f^3 is a total space of a locally trivial bundle over S^1 and a fiber 2-torus.

Take a tubular neighborhood $U(\gamma)$ of γ such that $U = U(\gamma)$ does not intersect Λ and the boundary $\partial U(\gamma) = \partial U$ is transversal to the trajectories of $sus^t(f)$. Due to [1, 2], there is the diffeomorphism $\vartheta : \partial U \rightarrow \partial U$ such that the manifold $(T_f^3 \setminus U) \cup_{\vartheta} U$ is a 3-sphere S^3 . Another words, one can take out $U(\gamma)$ from T_f^3 and after that return $U(\gamma)$ gluing together $(T_f^3 \setminus U)$ and $U(\gamma)$ along the boundary ∂U under the diffeomorphism ϑ . Since γ is a repelling set, the restriction $sus^t|_{T_f^3 \setminus U}$ continues to a flow f^t with an isolated repelling periodic trajectory. Since ∂U is transversal to sus^t , one can construct f^t to be an A-flow. It follows that the non-wandering set $NW(f^t)$ consists of an orientable 2-dimensional expanding attractor and trivial basic set (isolated periodic trajectory). This completes the proof. \square

PROOF OF THEOREM 3. First, we construct a flow desired on an n -sphere S^n for any $n \geq 4$. Let f^t be the A-flow on S^3 with a 2-dimensional expanding attractor Λ . Take S^3 to be smoothly embedded in S^4 such that $S^4 \setminus S^3$ is the disjoint union of 4-balls B_1^4, B_2^4 . One can continue f^t to S^4 to get A-flow, say f_4^t , such that f_4^t has a unique source at each B_i^4 , $i = 1, 2$, and S^3 is an attracting set for f_4^t . Clearly, Λ is the 2-dimensional expanding attractor of f_4^t . Continuing by similar way, one can construct an A-flow, say f_n^t , on S^n with the 2-dimensional expanding attractor Λ for any $n \geq 5$.

Now, let M^n be an arbitrary closed n -manifold, $n \geq 4$. Due to Smale [15], there is a gradient-like Morse-Smale flow g^t on M^n such that g^t has a sink s_0 . By construction, f_n^t has an isolated source, say r_n . Take out neighborhoods $U(s_0), U(r_n)$ of s_0, r_n such that the boundaries $\partial U(s_0,$

$\partial U(r_n)$ are transversal to g^t and f_n^t respectively. If one glue $M^n \setminus U(s_0)$, $S^n \setminus U(r_n)$ under a diffeomorphism $\partial U(s_0) \rightarrow \partial U(r_n)$, we get a connected sum $M^n \# S^n$ homeomorphic to M^n . Then the flows g^t and f_n^t define the A flow desired on M^n . This completes the proof. \square

Bibliography

- [1] **Christy J.** Branched surfaces and attractors I: dynamic branched surfaces. *Trans. Amer. Math. Soc.*, **336**(1993), 759-784.
- [2] **Francis D.** Drawing Seifert surfaces that fiber the figure 8 knot complement in \mathbb{S}^3 over \mathbb{S}^1 . *Amer. Math. Monthly*, **90**(1983), 589-599.
- [3] **Franks J., Williams B.** Anomalous Anosov flows. *Lect. Notes in Math.*, **819**(1980), 158-174.
- [4] **Grobman D.** On a homeomorphism of systems of differential equations. *Dokl. Akad. Nauk, USSR*, **128**(1959), 5, 880-881.
- [5] **Grobman D.** Topological classification of neighborhoods of singular point in n -dimensional space. *Matem. sbornik, USSR*, **56**(1962), 1, 77-94.
- [6] **Hartman P.** On the local linearization of differential equations. *Proc. AMS* **14**(1963), no 4, 568-573.
- [7] **Hirsch M., Pugh C., Shub M.** *Invariant Manifolds. Lect. Notes in Math.*, **583**(1977), Springer-Verlag.
- [8] **Hirsch M., Palis J., Pugh C., Shub M.** Neighborhoods of hyperbolic sets. *Invent. Math.*, **9**(1970), 121-134.
- [9] **Medvedev V., Zhuzhoma E.** On non-orientable two-dimensional basic sets on 3-manifolds, *Sbornik Math.*, **193**(2002), no 6, 869-888.
- [10] **Medvedev V., Zhuzhoma E.** On the existence of codimension one non-orientable expanding attractors. *Journ. Dyn. Contr. Syst.*, **11**(2005), no 3, 405-411.
- [11] **Pugh C., Shub M.** The Ω -stability theorem for flows. *Invent. Math.*, **11**(1970), 150-158.
- [12] **Margulis G.** U-flows on three-dimensional manifolds. Appendix of **Anosov D.V., Sinai Ya.G.** Some smooth ergodic systems, *Russian Math. Surveys*, **22**(1967), 103-168.
- [13] **Plykin R.V.** On the geometry of hyperbolic attractors of smooth cascades. *Russian Math. Surveys*, **39**(1984), no 6, 85-131.
- [14] **Robinson C.** *Dynamical Systems: stability, symbolic dynamics, and chaos*. Studies in Adv. Math., 2nd ed., CRC Press, Boca Raton, FL, **1999**.
- [15] **Smale S.** Morse inequalities for a dynamical system. *Bull. Amer. Math. Soc.*, **66**(1960), 43-49.

- [16] **Smale S.** Differentiable dynamical systems. *Bull. Amer. Math. Soc.*, **73**(1967), 747-817.
- [17] **Williams R.** Expanding attractors. *Publ. Math. IHES*, **43**(1974), 169-203.
- [18] **Zhuzhoma E.** Orientable basic sets of codimension 1. *Soviet Math. (Iz. VUZ)*, **26**(1082), no 5, 17–25.

National Research University Higher School of Economics, 25/12 Bolshaya Pecherskaya,
603005, Nizhni Novgorod, Russia

E-mail: medvedev-1942@mail.ru

E-mail: zhuzhoma@mail.ru