

How far can we trust published TESS periods?

Zdeněk Mikulášek¹ and Marek Skarka^{1,2}

¹ *Department of Theoretical Physics and Astrophysics, Masaryk University,
Kotlářská 2, 611 38 Brno, The Czech Republic (E-mail:
mikulas@physics.muni.cz)*

² *Astronomical Institute of the Czech Academy of Sciences
251 65 Ondřejov, The Czech Republic*

Received: May 1, 2005; Accepted: August 28, 2005

Abstract. Possible inaccuracies in the determination of periods from short-term time series caused by disregard of the real course of light curves and instrumental trends are documented on the example of the period analysis of simulated TESS-like light curve by notorious Lomb-Scargle method.

Key words: variable stars – period analysis – TESS data

1. Introduction

TESS data are now one of the most popular sources of information about variable stars, including their periods. However, TESS data suffers from two shortcomings that significantly corrupt the results of the period analysis with the standard tools. The data are strongly affected by instrumental trends of various kinds and are mostly obtained only in non-standard short time interval of 27 days (with a break) that is often comparable with the lengths of the periods themselves. The majority of such days-long periods and their uncertainties are mere artifacts of the method used to determine them and, at best, are just estimates.

The most widely used instruments of period analysis are frequency periodograms of various kinds, where astrophysically important frequencies are those in which the extremes of some suitably chosen characteristic of the studied data set occur. The reason that they generally give virtually identical results is that they are mostly consciously or covertly based on data interleaving by sine/cosine functions using the least squares method. The most commonly used is the historically oldest Lomb-Scargle method (Press & Rybicki, 1989), which after some algebra can be converted to a more informative version periodogram, where the frequency-dependent characteristic is the amplitude of light or other changes (Mikulášek et al., 2015).

The following demonstration, based on the simulations of the periodic light curve resembling a rotating chemically peculiar stars with a period of 2.7 days (see Fig. 1) shows possible pitfalls of standard processing of such type of data.




Figure 1. Simulated light curve without trends represented by 17 925 points.

1.1. Simulation of "TESS-like" data of a hypothetical CP star

Chemically peculiar (CP) stars are rotating variables with extensive photometric spots on their surface. The observed light changes of such objects are strictly periodic with the period of star rotation and can be described by a low degree harmonic polynomial. For our example, we chose a two-wave curve represented by a third-degree harmonic polynomial, described by five opted parameters (2) of the *phase function* ϑ (for details see in Mikulášek et al., 2008), $F(\vartheta, \mathbf{a})$, $P = 2.7$ d, with a maximum in the phase $\varphi = 0$.

$$\vartheta = \frac{t - M_0}{P} = E + \varphi, \quad M_0 = 1450 + P\Delta\varphi, \quad E = \text{IP}(\vartheta), \quad \varphi = \text{FP}(\vartheta), \quad (1)$$

$$F(\vartheta, \mathbf{a}) = a_1 \cos(2\pi\vartheta) + a_2 \cos(4\pi\vartheta) + a_3 \cos(6\pi\vartheta) + \quad (2)$$

$$+ a_4 [2 \sin(2\pi\vartheta) - \sin(4\pi\vartheta)] + a_5 [3 \sin(2\pi\vartheta) + 6 \sin(4\pi\vartheta) - 5 \sin(6\pi\vartheta)].$$

where t is the TESS BJD time of the observation ($t = \text{BJD} - 2\,457\,000$), M_0 is the TESS BJD moment of the initial light curve maximum, $\Delta\varphi$ is an optional *initial phase* parameter allowing horizontal shift of the simulated light curve, E is an integer epoch and φ is a common phase. IP and FP means the integer part and the fraction part of a number, $a_1 = -5$, $a_2 = -4.5$, $a_3 = -0.5$, $a_4 = -0.67$, and $a_5 = -0.17$ mmag.

The simulated TESS light curve is represented by 17 925 points with cadence of 2 minutes. The curve can, according to the assignment, show trends and Gaussian scatter (see Fig. 1).




Figure 2. The amplitude periodogram of the simulated light curve without scatter and trends for $\Delta\varphi = 0$.

2. The role of the initial phase $\Delta\varphi$

The amplitude frequency spectrum of the simulated light curve displays as expected three dominant, equidistantly spaced peaks with central frequencies f_1, f_2 , and f_3 (Fig. 2)), each carrying a period information: $1/f_1 = 2/f_2 = 3/f_3 = P$. However, this is fulfilled only approximately. If we limit ourselves to two peaks, then for $\Delta\varphi = 0$ we get: $P_1 = 1/f_1 = 2.7070(10)$ d, $P_2 = 2/f_2 = 2.6977(8)$ d. The deviation from the baseline period $P = 2.7$ d is thus evident and far exceeds the limits given by the uncertainty of the positioning of the frequency peaks. Why such a difference? The same results give other simple period finders.

The discrepancy would only disappear if the light curves were purely sinusoidal without higher harmonics. If they deviate from this ideal, the so-called 'periods' found are not real periods, but only parameters found by regression with an inadequate model that differ (sometimes fragrantly) from its pattern.

The following graph (Fig. 3) shows that the values of those 'periods' found in the periodograms are a complex periodic function of the initial phase $\Delta\varphi$, whose amplitude is simply shocking.

2.1. Influence of light curve trends

Both TESS and Kepler observations are strongly affected by aperiodic instrumental trends (Hümmerich et al., 2018; Mikulášek et al., 2019). Neglecting them has a completely devastating effect on period analysis (Fig. 4). Appropriate de-




Figure 3. $\Delta P_1 = P_1 - P$ (blue diamonds) and $\Delta P_2 = P_2 - P$ (red triangles) differences as function of the light curve shift $\Delta\varphi$.

trending of the observed light curves is highly desirable, because only in this way can we utilize the unprecedented accuracy of satellite photometry.

3. Modelling of light curves and trends

When applying the Lomb-Scargle method, we have to adjust the observation data by subtracting their mean value from them. Subsequently, these data are fitted by the simplest possible model of the periodic light curve, i.e. a linear combination of a pair of harmonic functions $F = a_1 \cos(2\pi f) + a_2 \sin(2\pi f)$, where the amplitude A , $A = \sqrt{a_1^2 + a_2^2}$, is a function of the frequency f and plotted on the $A(f)$ - amplitude periodogram, where the extremes of the plotted function are searched for and interpreted. It is even worse with estimating the uncertainty of determined frequencies with extreme amplitude values, because it is based on the apparatus of the least-squares method, but the conditions for its application are not met¹.

¹Observed deviations of the functional value model does not have a normal distribution, consecutive functional values are not independent, etc.




Figure 4. The comparison of ΔP_1 (blue diamonds) and ΔP_2 (red triangles) differences and true model ones (pink squares) for non-detrended increasing light curve as a function of the LC shift $\Delta\varphi$.

It is apparent from the foregoing examples that the use of conventional low parametric models is not sufficient for the description of the high-precision real light curves provided by modern instruments, in particular the satellites. It is, therefore, essential to use more advanced models to simultaneously fit a phase curve with a harmonic polynomial of at least third degree, while modeling trends by dividing the light curve into segments and describing them with a polynomial of the appropriate degree (Mikulášek et al., 2019; Hümmerich et al., 2018). Phase curve and trend models should be tailored to the actually observed light curves. The models should be functions of the parameters of the ephemeris of stellar periodicity, especially the time of the basic extremum, the mean period (or periods) and their time derivatives (Mikulášek, 2015, 2016).

Modern tools such as chi-square approach, robust regression, bootstrapping, etc. can be used to find parameters (including period/periods) and estimate their uncertainties. The results of this rigorous procedure can be compared with the results obtained by standard trivial period analysis procedures in Fig. 4. No further comment is needed.

4. Conclusions

- Short-term observational series corrupted by instrumental trends are not the easy observational material for accurate determination of periods.
- Using the Lomb-Scargle method we make serious errors both in the determination of the period and in the estimation of the uncertainty of this determination.
- In the periodic analysis of TESS data, the differences between the found period and the actual period are so huge that we are not able to correctly establish observations that are only a few months away.

A rigorous solution to the problem is to move to realistic models of light curves, including a true description of phase curve/curves as well as instrumental trends. Only then, we will fully use the information potential of short-term observation sets.

Acknowledgements. This work has been supported by grant GAČR 18-05665S and Operational Programme Research, Development and Education - Project "Post-doc@MUNI" (No. CZ.02.2.69/0.0/0.0/16_027/0008360).

References

- Hümmerich, S., Mikulášek, Z., Paunzen, E., et al., The Kepler view of magnetic chemically peculiar stars. 2018, *Astron. Astrophys.*, **619**, A98, DOI: 10.1051/0004-6361/201832938
- Mikulášek, Z., Phenomenological modelling of eclipsing system light curves. 2015, *Astron. Astrophys.*, **584**, A8, DOI: 10.1051/0004-6361/201425244
- Mikulášek, Z., Monitoring of rotational period variations in magnetic chemically peculiar stars. 2016, *Contributions of the Astronomical Observatory Skalnaté Pleso*, **46**, 95
- Mikulášek, Z., Krtička, J., Henry, G. W., et al., The extremely rapid rotational braking of the magnetic helium-strong star HD 37776. 2008, *Astron. Astrophys.*, **485**, 585, DOI: 10.1051/0004-6361:20077794
- Mikulášek, Z., Paunzen, E., Netopil, M., & Zejda, M., New Tools for Finding and Testing Weak Periodic Variability. 2015, in Astronomical Society of the Pacific Conference Series, Vol. **494**, *Physics and Evolution of Magnetic and Related Stars*, ed. Y. Y. Balega, I. I. Romanyuk, & D. O. Kudryavtsev, 320
- Mikulášek, Z., Zejda, M., Hümmerich, S., et al., Monitoring Period Variations of Variable Stars using Precise Photometric Surveys. 2019, in IAU Symposium, Vol. **339**, *Southern Horizons in Time-Domain Astronomy*, ed. R. E. Griffin, 110–113
- Press, W. H. & Rybicki, G. B., Fast Algorithm for Spectral Analysis of Unevenly Sampled Data. 1989, *Astrophys. J.*, **338**, 277, DOI: 10.1086/167197