

Oscillatory asymptotics for the Airy kernel determinant on two intervals

Elliot Blackstone, Christophe Charlier, Jonatan Lenells *

March 18, 2022

Abstract

We obtain asymptotics for the Airy kernel Fredholm determinant on two intervals. We give explicit formulas for all the terms up to and including the oscillations of order 1, which are expressed in terms of Jacobi θ -functions.

1 Introduction

The Airy point process is a well-known universal point process from the theory of random matrices and is used to model the behavior of the largest eigenvalue for a wide class of large Hermitian random matrices [6, 10, 11, 22, 30]. The Airy point process also appears in other related models. For example, it describes the transition between the solid and liquid regions in random tiling models [25] and the largest parts of Young diagrams with respect to the Plancherel measure [2, 4]. This is a determinantal point process on \mathbb{R} whose correlation kernel is given by

$$K^{\text{Ai}}(u, v) = \frac{\text{Ai}(u)\text{Ai}'(v) - \text{Ai}'(u)\text{Ai}(v)}{u - v}, \quad (1.1)$$

where Ai denotes the Airy function. Let $\vec{x} = (x_1, x_2, x_3)$ be such that $x_3 < x_2 < x_1 < 0$ and let $F(\vec{x}) = F(x_1, x_2, x_3)$ be the probability of finding no points on $(x_3, x_2) \cup (x_1, +\infty)$. In other words, F represents a gap probability on two disjoint intervals. It is well-known [31] that $F(\vec{x})$ can be written as a Fredholm determinant

$$F(\vec{x}) = \det \left(1 - \mathcal{K}^{\text{Ai}} \chi_{(x_3, x_2) \cup (x_1, +\infty)} \right), \quad (1.2)$$

where \mathcal{K}^{Ai} is the integral operator whose kernel is K^{Ai} . The purpose of this paper is to obtain asymptotics for $F(r\vec{x}) = F(rx_1, rx_2, rx_3)$ as $r \rightarrow +\infty$. Such asymptotics are often referred to as large gap asymptotics.

The history of gap expansions began with the work of Dyson [18] for the sine process. Many authors contributed over the years to the analysis of large gap asymptotics, including Widom [33], des Cloizeaux and Mehta [9], Krasovsky [26], Ehrhardt [19], etc., and we refer to [5] and the references therein for a historical review.

Large gap asymptotics on a single interval for the Airy kernel have been analysed in [1, 13, 32] and are now well-understood. The probability of a gap on $(-r, +\infty)$ enjoys the following asymptotics

$$\det \left(1 - \mathcal{K}^{\text{Ai}} \chi_{(-r, +\infty)} \right) = \exp \left(-\frac{|r|^3}{12} - \frac{1}{8} \log |r| + \zeta'(-1) + \frac{1}{24} \log 2 + \mathcal{O}(r^{-\frac{3}{2}}) \right), \quad \text{as } r \rightarrow +\infty, \quad (1.3)$$

where ζ' denotes the derivative of the Riemann zeta function. This expansion was first obtained in [32], but without rigorously proving the term of order 1. Note that the term of order 1 in (1.3) is

*Department of Mathematics, KTH Royal Institute of Technology, Lindstedtsvägen 25, SE-114 28 Stockholm, Sweden. We are grateful to the referees for several excellent suggestions. We are also grateful to B. Fahs, I. Krasovsky, and T. Maroudas for useful discussions. The work of all three authors is supported by the European Research Council, Grant Agreement No. 682537. Lenells also acknowledges support from the Göran Gustafsson Foundation, the Swedish Research Council, Grant No. 2015-05430, and the Ruth and Nils-Erik Stenbäck Foundation. E-mail: elliotb@kth.se, cchar@kth.se, jlenells@kth.se.

a constant (independent of r). In Theorem 1.1 below, we show that this is no longer the case when $x_3 < x_2 < x_1 < 0$; instead the term of order 1 presents some oscillations expressed in terms of a Jacobi θ -function.

We briefly introduce the necessary material to present our result. Let us define

$$\sqrt{\mathcal{Q}(z)} = \sqrt{(z - x_1)(z - x_2)(z - x_3)}, \quad (1.4)$$

where the principal branch is taken for each square root. Thus $\sqrt{\mathcal{Q}(z)}$ is analytic on $\mathbb{C} \setminus ((-\infty, x_3] \cup [x_2, x_1])$ and $\sqrt{\mathcal{Q}(z)} \sim z^{\frac{3}{2}}$ as $z \rightarrow \infty$. Let us also define

$$q(z) = -z^2 + \frac{x_1 + x_2 + x_3}{2}z - q_0, \quad (1.5)$$

where $q_0 \in \mathbb{R}$ is given by

$$q_0 = \left(\int_{x_3}^{x_2} \frac{ds}{\sqrt{\mathcal{Q}(s)}} \right)^{-1} \int_{x_3}^{x_2} \frac{-s^2 + \frac{x_1 + x_2 + x_3}{2}s}{\sqrt{\mathcal{Q}(s)}} ds. \quad (1.6)$$

The constant Ω , defined by

$$\Omega = \int_{x_2}^{x_1} \frac{2q(x)}{|\sqrt{\mathcal{Q}(x)}|} dx,$$

also appears in our final result. We show in Proposition 3.1 (see also Remark 5) that q has one simple root in (x_3, x_2) and another simple root in $(x_1, +\infty)$, so $q(x) > 0$ for $x \in (x_2, x_1)$, which implies that $\Omega > 0$. We define c_0 and τ by

$$c_0 = \left(\int_{x_3}^{x_2} \frac{2dx}{|\sqrt{\mathcal{Q}(x)}|} \right)^{-1} \in \mathbb{R}^+, \quad \tau = \int_{x_2}^{x_1} \frac{2ic_0 dx}{|\sqrt{\mathcal{Q}(x)}|} \in i\mathbb{R}^+. \quad (1.7)$$

The θ -function $\theta(z) = \theta(z; \tau)$ associated to the constant τ of (1.7) is given by (see e.g. [29])

$$\theta(z) = \sum_{m=-\infty}^{\infty} e^{2\pi i m z} e^{\pi i m^2 \tau}. \quad (1.8)$$

It is an entire function which satisfies

$$\theta(z + 1) = \theta(z), \quad \theta(z + \tau) = e^{-2\pi i z} e^{-\pi i \tau} \theta(z), \quad \theta(-z) = \theta(z), \quad \text{for all } z \in \mathbb{C}. \quad (1.9)$$

We are now ready to state our main result.

Theorem 1.1. *Let $x_3 < x_2 < x_1 < 0$ be fixed. As $r \rightarrow +\infty$, we have*

$$F(r\vec{x}) = \exp \left(c r^3 - \frac{1}{2} \log r + \log \theta(\nu) + C + \mathcal{O}(r^{-1}) \right), \quad (1.10)$$

$$c = \frac{x_1^3 + x_2^3 + x_3^3 - (x_1 + x_2)(x_1 + x_3)(x_2 + x_3)}{12} + \frac{q_0}{3}(x_1 + x_2 + x_3), \quad (1.11)$$

where $C = C(\vec{x})$ is independent of r and $\nu = \nu(r, \vec{x})$ is given by

$$\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}. \quad (1.12)$$

Remark 1. Theorem 1.1 has been verified numerically using the Linear Algebra package of Bornemann for Fredholm determinants [3].

Remark 2. The asymptotic formula (1.10) is not valid if $r \rightarrow +\infty$ and simultaneously $x_2 \uparrow x_1$ or $x_2 \downarrow x_3$. However, it is possible to naively expand our expression for the constant c as $x_2 \uparrow x_1$ or as

$x_2 \downarrow x_3$ and see if it agrees with the constants for the leading term of (1.3). The expression (1.11) for c is given in terms of q_0 defined in (1.6). We show in Lemma 3.3 that

$$q_0 = \frac{x_1 x_3}{2} + \mathcal{O}(x_2 - x_3), \quad \text{as } x_2 \downarrow x_3, \quad (1.13)$$

$$q_0 = \frac{x_1 x_3}{2} + \mathcal{O}\left(\frac{1}{\log(x_1 - x_2)}\right), \quad \text{as } x_2 \uparrow x_1. \quad (1.14)$$

Using these expansions, we obtain

$$\lim_{x_2 \uparrow x_1} c = -\frac{|x_3|^3}{12}, \quad \lim_{x_2 \downarrow x_3} c = -\frac{|x_1|^3}{12},$$

which is consistent with (1.3). Interestingly, the coefficient $-\frac{1}{2}$ of the $\log r$ term in (1.10) is independent of x_1, x_2, x_3 and different from the coefficient $-\frac{1}{8}$ of the $\log r$ term in (1.3).

Remark 3. In [16], Deift, Its and Zhou have obtained large gap asymptotics for the sine point process on disjoint intervals. Their final asymptotic formula is similar to our formula (1.10), in the sense that the subleading term in [16] is also logarithmic, and the third term presents oscillations also described in terms of Jacobi θ -functions. However, their constant associated to the logarithmic term is quite involved and expressed in terms of hyperelliptic integrals, while in our case the constant is simply $-\frac{1}{2}$. We were informed by B. Fahs at an early stage of our project that the prefactor of $\log s$ is also $-\frac{1}{2}$ for the sine kernel determinant on two large intervals [21].

1.1 Outline of the proof

We first follow the procedure developed by Its-Izergin-Korepin-Slavnov [24] to obtain an identity for $\partial_r \log F(r\vec{x})$ in terms of a certain resolvent operator. Next, we use a result of Claeys and Doeraene [7] to express this resolvent in terms of a model Riemann-Hilbert (RH) problem, whose solution is denoted Ψ , see (2.7) for the exact expression. We employ the Deift/Zhou [14, 17] steepest descent method to obtain asymptotics for Ψ as $r \rightarrow +\infty$. This method is well-established, but its application in the present situation leads to rather involved analysis. By integrating the differential identity $\partial_r \log F(r\vec{x})$, we have

$$\log F(r\vec{x}) = \log F(M\vec{x}) + \int_M^r \partial_{r'} \log F(r'\vec{x}) dr', \quad (1.15)$$

where $M > 0$ is a sufficiently large constant. We obtain our main result by substituting the large r' asymptotics for $\partial_{r'} \log F(r'\vec{x})$ and then performing integration with respect to r' .

Our analysis of the RH problem for Ψ involves several quantities related to a genus one Riemann surface, which is realized as two copies of the complex plane glued together along two disjoint intervals. In particular, Jacobi θ -functions, the Abel map, see (4.10), and the solution of a Jacobi inversion problem, all play an important role. The simple fact that the large r asymptotics for $F(r\vec{x})$ are of the form

$$F(r\vec{x}) = \exp\left(c r^3 + c_2 \log r + c_3 \log \theta(\nu) + C + \mathcal{O}(r^{-1})\right), \quad (1.16)$$

for certain constants c, c_2, c_3 and C requires a non-trivial amount of work. In particular, it is not straightforward to show that there are no terms proportional to $r^{\frac{3}{2}}$, no oscillations of order $\log r$, and that all oscillations of order 1 can be written in the form $c_3 \log \theta(\nu)$. An important part of this work is devoted to understanding the algebra related to the θ -functions and the Abel map (see [16] for pioneering work in this direction). We obtain very simple expressions for c_2 and c_3 only after considerable simplifications using many identities involving θ -functions and the Abel map. Furthermore, the identity $c_3 = 1$ requires the use of Riemann's bilinear identity, while $c_2 = -\frac{1}{2}$ requires the explicit evaluation of certain elliptic integrals, see Proposition 8.7. The fact that we manage to prove $c_2 = -\frac{1}{2}$ and $c_3 = 1$ can be viewed as one of the main contributions of this paper.

The integration constant $\log F(M\vec{x})$ in (1.15) is unknown and contributes to the expression C of (1.10), so our method does not allow for an explicit expression for C . Such constants are notoriously complicated to obtain (see [27, 20]), and we leave this for future works.

Remark 4. I. Krasovsky and T. Maroudas [28] have simultaneously and independently analyzed the large r asymptotics of the Fredholm determinant $F(r\vec{x})$. Their method is based on a different differential identity than the one considered here.

1.2 Organization of the paper

First, we present the model RH problem for Ψ and obtain the differential identity (2.7) in Section 2. We perform the first steps of the steepest descent analysis in Section 3. These steps are denoted by $\Psi \mapsto T \mapsto S$ and consists of constructing the so-called g -function and the opening of lenses. The second main part of the steepest descent method consists in constructing approximations to S in four distinct regions of the complex plane. The global parametrix $P^{(\infty)}$ approximates S everywhere in the complex plane except near the points $y_j := x_j - x_3$, and the local parametrix $P^{(y_j)}$ approximates S near the point y_j , $j = 1, 2, 3$. In Section 4, we construct $P^{(\infty)}$ explicitly in terms of Jacobi θ -functions. This construction is technically different from the construction in [15], but contains similar ideas [12]. In Section 5, we use some results from [16] to build local parametrices $P^{(y_j)}$, $j = 1, 2, 3$, in terms of Bessel functions. The last step $S \mapsto R$ of the steepest descent method is done in Section 6; in particular we show that $P^{(y_j)}$ approximates S in a small neighborhood of y_j , $j = 1, 2, 3$, for sufficiently large r . In Section 7, we summarize certain θ -function identities that are required to analyze the differential identity (2.7). In Section 8, we complete the proof of Theorem 1.1 by obtaining large r asymptotics for $\partial_r \log F(r\vec{x})$ via (2.7), and subsequently large r asymptotics for $\log F(r\vec{x})$ via (1.15).

2 Differential identity for F

In this section, we use the method of Its-Izergin-Korepin-Slavnov [24] to obtain a differential identity for $\partial_r \log F(r\vec{x})$ in terms of a model RH problem Ψ obtained in [7], which is a simple generalization of the problem in [8, p. 374] and the corresponding formula [8, eq (2.17)].

A kernel $K : \mathbb{R}^2 \rightarrow \mathbb{R}$ is said to be integrable if it can be written in the form $K(x, y) = \frac{f^T(x)g(y)}{x-y}$, where $f(x)$ and $g(y)$ are column vectors satisfying $f^T(x)g(x) = 0$. It is well-known and easy to see from (1.1) that the Airy kernel K^{Ai} is integrable. In our case, we need to consider the trace-class kernel

$$K_{\vec{x}}(x, y) = K^{\text{Ai}}(x, y)(\chi_{(x_3, x_2)}(y) + \chi_{(x_1, +\infty)}(y)). \quad (2.1)$$

This kernel is also integrable with f and g given by

$$f(x) = \begin{pmatrix} \text{Ai}(x) \\ \text{Ai}'(x) \end{pmatrix}, \quad g(y) = \begin{pmatrix} \text{Ai}'(y)(\chi_{(x_3, x_2)}(y) + \chi_{(x_1, +\infty)}(y)) \\ -\text{Ai}(y)(\chi_{(x_3, x_2)}(y) + \chi_{(x_1, +\infty)}(y)) \end{pmatrix}.$$

The associated integral operator $\mathcal{K}_{\vec{x}} : L^2((x_3, x_2) \cup (x_1, +\infty)) \rightarrow L^2((x_3, x_2) \cup (x_1, +\infty))$ is given by

$$\mathcal{K}_{\vec{x}}f(x) = \int_{-\infty}^{+\infty} K^{\text{Ai}}(x, y)(\chi_{(x_3, x_2)}(y) + \chi_{(x_1, +\infty)}(y))f(y)dy. \quad (2.2)$$

Let us now scale the size of the intervals with a parameter $r > 0$, i.e. we replace \vec{x} by $r\vec{x}$. Since $\partial_r \log F(r\vec{x}) = \partial_r \log \det(I - \mathcal{K}_{r\vec{x}})$, by using general theory of integrable kernel operators, we obtain

$$\begin{aligned} \partial_r \log F(r\vec{x}) &= -\text{Tr} \left[(I - \mathcal{K}_{r\vec{x}})^{-1} \partial_r \mathcal{K}_{r\vec{x}} \right] = \sum_{j=1}^3 (-1)^{j+1} x_j \text{Tr} \left[(I - \mathcal{K}_{r\vec{x}})^{-1} \mathcal{K}_{r\vec{x}} \delta_{rx_j} \right] \\ &= \sum_{j=1}^3 (-1)^{j+1} x_j \text{Tr} [\mathcal{R}_{r\vec{x}} \delta_{rx_j}] = \sum_{j=1}^3 (-1)^{j+1} x_j \lim_{u \rightarrow rx_j} R_{r\vec{x}}(u, u), \end{aligned} \quad (2.3)$$

where δ_{rx_j} is the Dirac delta operator, $\mathcal{R}_{r\vec{x}}$ is the resolvent operator given by

$$\mathcal{R}_{r\vec{x}} := (I - \mathcal{K}_{r\vec{x}})^{-1} \mathcal{K}_{r\vec{x}} = (I - \mathcal{K}_{r\vec{x}})^{-1} - I,$$

$R_{r\vec{x}}$ is the kernel of $\mathcal{R}_{r\vec{x}}$, and the limits $u \rightarrow rx_j$, $j = 1, 2, 3$ in (2.3) are taken from the interior of $(rx_3, rx_2) \cup (rx_1, +\infty)$.

Next, we use a result of [7] (based on [24]) which relates $\mathcal{R}_{r\vec{x}}$ to the model RH problem Ψ . For convenience, we define

$$y_j = x_j - x_3, \quad j = 1, 2, 3.$$




Figure 1: Jump contours for the model RH problem for Ψ .

Using the Its-Izergin-Korepin-Slavnov method, it was shown in [7, proof of Proposition 1] that the resolvent kernel $R_{r\vec{x}}(u; u)$ can be expressed in terms of the solution of a naturally associated RH problem. For $u \in (rx_3, rx_2) \cup (rx_1, +\infty)$, we have

$$R_{r\vec{x}}(u, u) = \frac{1}{2\pi i} (\Psi_+^{-1} \Psi'_+)_{21} (u - x_3; x_3, \vec{y}), \quad (2.4)$$

where $\vec{y} = (y_1, y_2, y_3)$ and $\Psi(z; x_3, \vec{y})$ is the solution of the following RH problem.

RH problem for Ψ

- (a) $\Psi : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{C}^{2 \times 2}$ is analytic, with

$$\Gamma = e^{\pm \frac{2\pi i}{3}}(0, +\infty) \cup (-\infty, 0] \cup [y_2, y_1] \quad (2.5)$$

and Γ oriented as in Figure 1.

- (b) $\Psi(z)$ has continuous boundary values as $z \in \Gamma \setminus \{y_1, y_2, y_3\}$ is approached from the left (+ side) or from the right (− side) and they are related by

$$\begin{aligned} \Psi_+(z) &= \Psi_-(z) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } z \in e^{\pm \frac{2\pi i}{3}}(0, +\infty), \\ \Psi_+(z) &= \Psi_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (-\infty, 0), \\ \Psi_+(z) &= \Psi_-(z) \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (y_2, y_1). \end{aligned}$$

- (c) As $z \rightarrow \infty$, we have

$$\Psi(z) = (I + \mathcal{O}(z^{-1})) z^{\frac{1}{4}\sigma_3} M^{-1} e^{-(\frac{2}{3}z^{\frac{3}{2}} + x_3 z^{\frac{1}{2}})\sigma_3}, \quad (2.6)$$

where $M = (I + i\sigma_1)/\sqrt{2}$, $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, and where principal branches of $z^{\frac{3}{2}}$ and $z^{\frac{1}{2}}$ are taken.

- (d) $\Psi(z) = \mathcal{O}(\log |z - y_j|)$ as $z \rightarrow y_j$, $j = 1, 2, 3$.

We conclude from the definition of F (1.2) together with (2.3) and (2.4) that

$$\partial_r \log F(r\vec{x}) = \sum_{j=1}^3 \frac{(-1)^{j+1} x_j}{2\pi i} \lim_{z \rightarrow y_j} (\Psi_+^{-1} \Psi'_+)_{21} (rz; rx_3, r\vec{y}), \quad (2.7)$$

where the limits as $z \rightarrow y_j$, $j = 1, 2, 3$, are taken such that $z \in (0, y_2) \cup (y_1, +\infty)$. The differential identity (2.7) expresses $\partial_r \log F(r\vec{x})$ in terms of Ψ , which is the solution to a RH problem. In Sections 3-6, we employ the Deift/Zhou steepest descent method [17] to obtain the asymptotics of $\Psi(z)$, $\Psi'(z)$ as $r \rightarrow +\infty$ for z in small neighborhoods of y_1, y_2, y_3 .

3 Steepest descent for Ψ : first steps

In this section, we perform the first steps of the steepest descent method. We obtain the so-called g -function in Subsection 3.1. This function is used to normalize the RH problem at ∞ via the $\Psi \rightarrow T$ transformation of Subsection 3.2. Finally, we proceed with the opening of the lenses $T \mapsto S$ in Subsection 3.3.

3.1 g -function

The first step of the analysis consists of normalizing the RH problem at ∞ by means of an appropriate g -function; see [16] for a similar construction. Let us define

$$\sqrt{\mathcal{R}(z)} = \sqrt{(z - y_1)(z - y_2)z}$$

where the principal branch is taken for each square root. More precisely, $\sqrt{\mathcal{R}(z)}$ is analytic on $\mathbb{C} \setminus ((-\infty, 0] \cup [y_2, y_1])$ and $\sqrt{\mathcal{R}(z)} \sim z^{\frac{3}{2}}$ as $z \rightarrow \infty$. In particular it satisfies the jumps

$$\sqrt{\mathcal{R}(z)}_+ + \sqrt{\mathcal{R}(z)}_- = 0 \quad \text{for } z \in (-\infty, 0) \cup (y_2, y_1). \quad (3.1)$$

We define the derivative of the g -function by

$$g'(z) = \frac{p(z)}{\sqrt{\mathcal{R}(z)}}, \quad (3.2)$$

where

$$p(z) := -z^2 + \frac{y_1 + y_2 - x_3}{2}z + \frac{x_3(y_1 + y_2)}{4} + \frac{(y_1 - y_2)^2}{8} - g_1 \quad (3.3)$$

and $g_1 \in \mathbb{R}$ is given by

$$g_1 = \left(\int_0^{y_2} \frac{ds}{\sqrt{\mathcal{R}(s)}} \right)^{-1} \left(\int_0^{y_2} \frac{-s^2 + \frac{y_1 + y_2 - x_3}{2}s + \frac{x_3(y_1 + y_2)}{4} + \frac{(y_1 - y_2)^2}{8}}{\sqrt{\mathcal{R}(s)}} ds \right). \quad (3.4)$$

By definition of g_1 , we have

$$\int_0^{y_2} g'(s) ds = 0. \quad (3.5)$$

Remark 5. The functions \mathcal{Q} and q defined in (1.4) and (1.5) have the relations with \mathcal{R} and p

$$\mathcal{Q}(z) = \mathcal{R}(z - x_3) \quad \text{and} \quad q(z) = p(z - x_3).$$

Proposition 3.1. *The degree two polynomial p has two simple real zeros z_{\pm} , with $z_- \in (0, y_2)$ and $z_+ \in (y_1, \infty)$. Moreover, we have*

$$p(y_3) < 0, \quad p(y_2) > 0, \quad p(y_1) > 0.$$

Proof. The polynomial p has real coefficients, thus $p(z) \in \mathbb{R}$ for $z \in \mathbb{R}$. Then we immediately conclude from (3.5) that $p(z)$ must have two real simple zeros and that at least one of them lies on $(0, y_2)$, because $\sqrt{\mathcal{R}(z)} < 0$ for all $z \in (0, y_2)$. However it is not straightforward to prove that p has one zero in $(y_1, +\infty)$, or equivalently that $p(y_1) > 0$ (since $p(z) \rightarrow -\infty$ as $z \rightarrow +\infty$). Note that $p(z)$ can be written as

$$p(z) = -z^2 + \frac{z}{2}(y_1 + y_2 - x_3) + \alpha, \quad \text{where} \quad \alpha := \left(\int_0^{y_2} \frac{ds}{\sqrt{\mathcal{R}(s)}} \right)^{-1} \left(\int_0^{y_2} \frac{s^2 - \frac{s}{2}(y_1 + y_2 - x_3)}{\sqrt{\mathcal{R}(s)}} ds \right).$$

The constant α can be written explicitly in terms of complete elliptic integrals of the first (**K**) and second (**E**) kind, which are defined as [23, eq 8.112]

$$\mathbf{K}(k) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \quad \mathbf{E}(k) = \int_0^1 \frac{\sqrt{1-k^2x^2}}{\sqrt{1-x^2}} dx.$$

Let us define $k_\star := \sqrt{\frac{y_2}{y_1}}$. Using [23, eq 3.131.4, 3.141.15 and 3.141.27], we obtain

$$\begin{aligned} \int_0^{y_2} \frac{ds}{\sqrt{\mathcal{R}(s)}} &= \frac{-2}{\sqrt{y_1}} \mathbf{K}(k_\star), \quad \int_0^{y_2} \frac{s ds}{\sqrt{\mathcal{R}(s)}} = -2\sqrt{y_1} (\mathbf{K}(k_\star) - \mathbf{E}(k_\star)), \\ \int_0^{y_2} \frac{s^2 ds}{\sqrt{\mathcal{R}(s)}} &= \frac{2}{3} \sqrt{y_1} (2(y_1 + y_2) \mathbf{E}(k_\star) - (2y_1 + y_2) \mathbf{K}(k_\star)), \end{aligned}$$

from which we deduce

$$\alpha = \frac{y_1}{2} \left[\frac{y_1 - y_2}{3} + x_3 - \frac{\mathbf{E}(k_\star)}{\mathbf{K}(k_\star)} \left(\frac{y_1 + y_2}{3} + x_3 \right) \right]. \quad (3.6)$$

From [29, eq 19.9.8], we have the lower bound

$$\frac{\mathbf{E}(k_\star)}{\mathbf{K}(k_\star)} > \sqrt{1 - k_\star^2} = \sqrt{1 - \frac{y_2}{y_1}} = \sqrt{\frac{x_1 - x_2}{x_1 - x_3}}.$$

We weaken this bound for our purpose. Notice

$$4(x_1 - x_2)(x_1 - x_3) = (x_1 + x_2 + x_3)^2 + 3x_1(x_1 - 2x_2 - 2x_3) - (x_2 - x_3)^2 < (x_1 + x_2 + x_3)^2,$$

because $x_1 - 2x_2 - 2x_3 > 0$ and $x_1 < 0$, so we have

$$\frac{x_1 - x_2}{x_1 - x_3} = \frac{4(x_1 - x_2)^2}{4(x_1 - x_2)(x_1 - x_3)} > \frac{4(x_1 - x_2)^2}{(x_1 + x_2 + x_3)^2}.$$

The new bound is

$$\frac{\mathbf{E}(k_\star)}{\mathbf{K}(k_\star)} > \frac{2(x_2 - x_1)}{x_1 + x_2 + x_3},$$

from which we have

$$p(y_1) = \frac{y_1}{2}(y_2 - y_1 - x_3) + \alpha = \frac{y_1}{6} \left[2(x_2 - x_1) - \frac{\mathbf{E}(k_\star)}{\mathbf{K}(k_\star)} (x_1 + x_2 + x_3) \right] > 0,$$

as desired. \square

We define the g -function by

$$g(z) = \int_{y_1}^z g'(s) ds, \quad (3.7)$$

where the path of integration does not cross $(-\infty, y_1]$.

Lemma 3.2. *Let $x_3 < x_2 < x_1 < 0$.*

1. *The g -function is analytic in $\mathbb{C} \setminus (-\infty, y_1]$ and satisfies $g(z) = \overline{g(\bar{z})}$.*
2. *The g -function satisfies the jump conditions*

$$g_+(z) + g_-(z) = 0, \quad z \in (-\infty, 0) \cup (y_2, y_1), \quad (3.8)$$

$$g_+(z) - g_-(z) = i\Omega, \quad z \in (0, y_2), \quad (3.9)$$

where $\Omega = 2i \int_{y_2}^{y_1} g'_+(s) ds = -2ig_+(y_2) \in \mathbb{R}_+$.

3. *As $z \rightarrow \infty$, $z \notin (-\infty, 0)$, we have*

$$g(z) = -\frac{2}{3}z^{\frac{3}{2}} - x_3z^{\frac{1}{2}} + 2g_1z^{-\frac{1}{2}} + \mathcal{O}(z^{-\frac{3}{2}}). \quad (3.10)$$

4. Let N_0 be an open neighborhood of $(-\infty, y_1]$, such that

$$\left\{ z \in \mathbb{C} : \arg z \in \left(-\pi, -\pi + \frac{\pi}{3} + \epsilon \right) \cup \left(\pi - \frac{\pi}{3} - \epsilon, \pi \right) \text{ and } |z| \geq M \right\} \subset N_0 \quad (3.11)$$

for some $\epsilon > 0$ and $M > 0$. We have

$$\Re[g(z)] \geq 0, \quad \text{for all } z \in N_0 \quad (3.12)$$

where equality holds only when $z \in (-\infty, 0] \cup [y_2, y_1]$.

Proof. 1. Analyticity follows from the definition (3.7). It is straightforward to verify from (3.2) that $g'(z) = \overline{g'(\bar{z})}$, from which we deduce

$$\overline{g(\bar{z})} = \int_{y_1}^{\bar{z}} g'(s) ds = \int_0^1 g'(y_1 + t(\bar{z} - y_1))(\bar{z} - y_1) dt = \int_0^1 g'(y_1 + t(z - y_1))(z - y_1) dt = g(z).$$

2. The jumps (3.8) follow from (3.1) and (3.5). Let γ be a closed curve with positive orientation surrounding $[y_2, y_1]$ and not intersecting $(-\infty, 0]$. For $z \in (0, y_2)$, we use (3.8) to write

$$g_+(z) - g_-(z) = \int_{\gamma} g'(s) ds = -2 \int_{y_2}^{y_1} g'_+(s) ds,$$

and this is (3.9). It follows from Proposition 3.1 that

$$\Omega = 2i \int_{y_2}^{y_1} g'_+(s) ds = 2 \int_{y_2}^{y_1} \frac{p(s) ds}{\sqrt{s} \sqrt{s - y_2} \sqrt{y_1 - s}} > 0.$$

3. From (3.2), a computation shows that

$$g'(z) = -\sqrt{z} - \frac{x_3}{2} z^{-\frac{1}{2}} - g_1 z^{-\frac{3}{2}} + \mathcal{O}(z^{-\frac{5}{2}}), \quad \text{as } z \rightarrow \infty,$$

and then after integration we have

$$g(z) = -\frac{2}{3} z^{\frac{3}{2}} - x_3 z^{\frac{1}{2}} + g_0 + 2g_1 z^{-\frac{1}{2}} + \mathcal{O}(z^{-\frac{3}{2}}), \quad \text{as } z \rightarrow \infty, \quad (3.13)$$

for a certain $g_0 \in \mathbb{C}$. We deduce from (3.13) that

$$g_+(z) + g_-(z) = 2g_0 + \mathcal{O}(z^{-\frac{1}{2}}), \quad \text{as } z \rightarrow -\infty, \quad z \in \mathbb{R}^-.$$

Comparing the above with (3.8), we conclude that $g_0 = 0$.

4. It follows from part 2 that $\Re[g(z)]$ is single valued on \mathbb{C} and furthermore $\Re[g(z)] \equiv 0$ for $z \in (-\infty, 0) \cup (y_2, y_1)$. Due to Proposition 3.1, we have

$$\Im[g'_+(x)] = \Im \left[\frac{p(x)}{\sqrt{\mathcal{R}(x)}_+} \right] < 0, \quad \text{for } x \in (-\infty, 0) \cup (y_2, y_1).$$

So $\Im[g_+(x)]$ decreases as $x \in (-\infty, 0) \cup (y_2, y_1)$ increases, and it follows from the Cauchy-Riemann equations that $\Re[g(x + i\epsilon)]$ increases (and is therefore positive) for sufficiently small $\epsilon > 0$. We deduce from part 1 that $\Re[g(x - i\epsilon)] > 0$ for sufficiently small $\epsilon > 0$. Another immediate consequence of Proposition 3.1 is that $\Re[g(z)] > 0$ for all $z \in (0, y_2)$. Furthermore, from (3.7) we have

$$\Re[g(z)] \sim c(z - y_1)^{\frac{1}{2}}, \quad c > 0,$$

(the exact value of c is unimportant here) as $z \rightarrow y_1$ so there is an open disk D_{y_1} around y_1 such that $\Re[g(z)] \geq 0$ for all $z \in D_{y_1}$ with equality $\Re[g(z)] = 0$ if and only if $z \in D_{y_1} \cap (y_2, y_1)$. We conclude similarly that there also exists open disks D_{y_2} and D_{y_3} around y_2 and y_3 , respectively, such that

$$\Re[g(z)] \geq 0, \quad \text{for all } z \in D_{y_2} \cup D_{y_3}$$

with equality only when $z \in (-\infty, 0] \cup [y_2, y_1]$. The fact that we can choose $\epsilon > 0$ and $M > 0$ such that (3.11) holds follows directly from (3.10). \square

In Lemma 3.3 below, we obtain asymptotics for g_1 as $y_2 \downarrow 0$ and as $y_2 \uparrow y_1$. This is useful to compare our results with [13], see Remark 2. Indeed, we deduce (1.13)-(1.14) from (3.15)-(3.16) by noting that

$$q_0 = g_1 + \frac{2x_1x_2 + 2x_1x_3 + 2x_2x_3 - x_1^2 - x_2^2}{8}. \quad (3.14)$$

Lemma 3.3. *Fix y_1 and x_3 . We have*

$$g_1 = \frac{y_1}{4} \left(x_3 + \frac{y_1}{2} \right) + \mathcal{O}(y_2^2), \quad \text{as } y_2 \downarrow 0 \quad (3.15)$$

$$g_1 = -\frac{y_1(2y_1 + 3x_3)}{3 \log(y_1 - y_2)} + \mathcal{O} \left(\frac{1}{\log^2(y_1 - y_2)} \right), \quad \text{as } y_2 \uparrow y_1. \quad (3.16)$$

Proof. As $y_2 \downarrow 0$, we have

$$\begin{aligned} \int_0^{y_2} \frac{ds}{\sqrt{\mathcal{R}(s)}} &= \int_0^{y_2} \frac{-ds}{\sqrt{s(y_2 - s)}} \frac{1}{\sqrt{y_1}} \left(1 + \frac{s}{2y_1} + \frac{3s^2}{8y_1^2} + \mathcal{O}\left(\frac{s^3}{y_1^3}\right) \right) \\ &= \frac{-\pi}{\sqrt{y_1}} \left(1 + \frac{y_2}{4y_1} + \frac{9y_2^2}{64y_1^2} + \mathcal{O}(y_2^3) \right). \end{aligned}$$

Similar computations show that

$$\int_0^{y_2} \frac{s ds}{\sqrt{\mathcal{R}(s)}} = \frac{-\pi y_2}{2\sqrt{y_1}} \left(1 + \frac{3y_2}{8y_1} + \mathcal{O}(y_2^2) \right), \quad \int_0^{y_2} \frac{s^2 ds}{\sqrt{\mathcal{R}(s)}} = \frac{-3\pi y_2^2}{8\sqrt{y_1}} + \mathcal{O}(y_2^3), \quad \text{as } y_2 \downarrow 0,$$

from which we obtain (3.15). Now we turn to the proof of (3.16). Using (3.6), we express the constant g_1 explicitly in terms of complete elliptic integrals:

$$g_1 = \frac{y_1}{2} \frac{\mathbf{E}(k_*)}{\mathbf{K}(k_*)} \left(\frac{y_1 + y_2}{3} + x_3 \right) - \frac{x_3}{4} (y_1 - y_2) - \frac{1}{24} (y_1 - y_2)(y_1 + 3y_2).$$

We then use [29, eq 19.12.1 and 19.12.2] for the asymptotics as $y_2 \uparrow y_1$ to obtain (3.16). \square

3.2 Rescaling of the RH problem

Define the function $T(z) = T(z; x_3, \vec{y})$ as follows:

$$T(z) = \begin{pmatrix} r^{-\frac{1}{4}} & 2ir^{\frac{7}{4}}g_1 \\ 0 & r^{\frac{1}{4}} \end{pmatrix} \Psi(rz; rx_3, r\vec{y}) e^{-r^{\frac{3}{2}}g(z)\sigma_3}, \quad z \in \mathbb{C} \setminus \Gamma. \quad (3.17)$$

The asymptotics (2.6) of Ψ then imply after a small calculation that T behaves as

$$T(z) = (I + \mathcal{O}(z^{-1})) z^{\frac{1}{4}\sigma_3} M^{-1}, \quad (3.18)$$

as $z \rightarrow \infty$, $z \notin (-\infty, 0)$, where the principal branches of the roots are chosen. Using (3.8), the jumps $T_-(z)^{-1}T_+(z)$ for $z \in (y_2, y_1)$ can be factorized as

$$\begin{pmatrix} e^{-2r^{\frac{3}{2}}g_+(z)} & 1 \\ 0 & e^{-2r^{\frac{3}{2}}g_-(z)} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ e^{-2r^{\frac{3}{2}}g_-(z)} & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ e^{-2r^{\frac{3}{2}}g_+(z)} & 1 \end{pmatrix}. \quad (3.19)$$

3.3 Opening of lenses

Around the interval (y_2, y_1) , we will split the jump contour in three parts using (3.19). This transformation is traditionally called opening of lenses. Let us consider lens-shaped contours $\gamma_{2,+}$ and $\gamma_{2,-}$ lying in the upper and lower half plane respectively, as shown in Figure 2. We define S by

$$S(z) = T(z) \begin{cases} \begin{pmatrix} 1 & 0 \\ -e^{-2r^{\frac{3}{2}}g(z)} & 1 \end{pmatrix}, & z \text{ is inside the lenses around } (y_2, y_1) \text{ and } \Im[z] > 0, \\ \begin{pmatrix} 1 & 0 \\ e^{-2r^{\frac{3}{2}}g(z)} & 1 \end{pmatrix}, & z \text{ is inside the lenses around } (y_2, y_1) \text{ and } \Im[z] < 0, \\ I, & \text{otherwise.} \end{cases} \quad (3.20)$$

We use Lemma 3.2, as well as the RH problem for Ψ and the definitions (3.17) of T and (3.20) of S to conclude that S satisfies the following RH problem.




Figure 2: Jump contours Σ_S for S .

RH problem for S

(a) $S : \mathbb{C} \setminus \Sigma_S \rightarrow \mathbb{C}^{2 \times 2}$ is analytic, with

$$\Sigma_S = (-\infty, y_1] \cup \gamma_+ \cup \gamma_-, \quad \gamma_{\pm} = \gamma_{2,\pm} \cup (e^{\pm \frac{2\pi i}{3}}(0, +\infty)) \quad (3.21)$$

and Σ_S oriented as in Figure 2.

(b) The jumps for S are given by

$$S_+(z) = S_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad z \in (-\infty, 0) \cup (y_2, y_1), \quad (3.22)$$

$$S_+(z) = S_-(z) \begin{pmatrix} 1 & 0 \\ e^{-2r^{\frac{2}{3}}g(z)} & 1 \end{pmatrix}, \quad z \in \gamma_+ \cup \gamma_-, \quad (3.23)$$

$$S_+(z) = S_-(z) e^{-i\Omega r^{\frac{3}{2}}\sigma_3}, \quad z \in (0, y_2). \quad (3.24)$$

(c) As $z \rightarrow \infty$, $z \notin (-\infty, 0)$, we have

$$S(z) = (I + \mathcal{O}(z^{-1})) z^{\frac{1}{4}\sigma_3} M^{-1}. \quad (3.25)$$

(d) As $z \rightarrow y_j$, $j = 1, 2, 3$, we have

$$S(z) = \mathcal{O}(\log |z - y_j|). \quad (3.26)$$

Deforming γ_+ and γ_- if necessary, we assume that $\gamma_+, \gamma_- \subset N_0$ where N_0 has the properties described in Lemma 3.2. In particular, $\Re[g(z)] > 0$ for all $z \in \gamma_+ \cup \gamma_-$, so that the jumps for S are exponentially close to the identity as $r \rightarrow +\infty$ on the lens boundaries. This convergence is uniform outside small fixed neighborhoods of y_1, y_2, y_3 , but is not uniform as $r \rightarrow +\infty$ and simultaneously $z \rightarrow y_j$, $j = 1, 2, 3$.

4 Global parametrix

In this section we construct the global parametrix $P^{(\infty)}$. We will show in Section 6 that $P^{(\infty)}$ approximates S outside of neighborhoods of y_1, y_2, y_3 .

RH problem for $P^{(\infty)}$

(a) $P^{(\infty)} : \mathbb{C} \setminus (-\infty, y_1] \rightarrow \mathbb{C}^{2 \times 2}$ is analytic.




Figure 3: The selection of a canonical homology basis for X . The solid parts are on the first sheet and the dashed parts are on the second sheet.

(b) The jumps for $P^{(\infty)}$ are given by

$$\begin{aligned} P_+^{(\infty)}(z) &= P_-^{(\infty)}(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, & z \in (-\infty, 0) \cup (y_2, y_1), \\ P_+^{(\infty)}(z) &= P_-^{(\infty)}(z) e^{-i\Omega r^{\frac{3}{2}} \sigma_3}, & z \in (0, y_2). \end{aligned} \quad (4.1)$$

(c) As $z \rightarrow \infty$, we have

$$P^{(\infty)}(z) = (I + \mathcal{O}(z^{-1})) z^{\frac{1}{4}\sigma_3} M^{-1}. \quad (4.2)$$

(d) As $z \rightarrow y_j$, we have $P^{(\infty)}(z) = \mathcal{O}((z - y_j)^{-\frac{1}{4}})$ for $j = 1, 2, 3$.

Conditions (a)-(c) for the RH problem for $P^{(\infty)}$ are obtained from the RH problem S by ignoring the (pointwise) exponentially small jumps of S . Condition (d) does not come from the RH problem for S and has been added to ensure uniqueness of the solution. The solution $P^{(\infty)}$ can be constructed in terms of Jacobi θ -functions [12]. This construction is technically different from the construction in [15], but contains similar ideas. Let X be the two-sheeted Riemann surface of genus one associated to $\sqrt{\mathcal{R}(z)}$, with

$$\sqrt{\mathcal{R}(z)} = \sqrt{z(z - y_2)(z - y_1)},$$

and we let $\sqrt{\mathcal{R}(z)} > 0$ for $z \in (y_1, +\infty)$ on the first sheet. We also define A and B cycles such that they form a canonical homology basis of X . The A cycle surrounds $(0, y_2)$ with counterclockwise orientation. The upper part of the A cycle (the solid line in Figure 3) lies on the first sheet, and the lower part (the dashed line in Figure 3) lies on the second sheet. The B cycle surrounds $(-\infty, 0)$ with clockwise orientation. The unique A -normalized holomorphic one-form ω on X is given by

$$\omega = \frac{c_0 dz}{\sqrt{\mathcal{R}(z)}}, \quad c_0 = \left(\int_A \frac{dz}{\sqrt{\mathcal{R}(z)}} \right)^{-1}. \quad (4.3)$$

By construction $\int_A \omega = 1$ and the lattice parameter is given by $\tau = \int_B \omega \in i\mathbb{R}^+$. The associated θ -function of the third kind $\theta(z) = \theta(z; \tau)$ is given by

$$\theta(z) = \sum_{m=-\infty}^{\infty} e^{2\pi i m z} e^{\pi i m^2 \tau}. \quad (4.4)$$

It is an entire function which satisfies (1.9). The zeros of $\theta(z)$ are the points $\frac{1}{2} + m_1 + \frac{\tau}{2} + m_2 \tau$, with $m_1, m_2 \in \mathbb{Z}$. Let X_1 and X_2 denote the upper and lower sheet of X , respectively, and consider the set

$$\Lambda := \{u \in \mathbb{C} : -\frac{1}{2} \leq \Re[u] \leq \frac{1}{2} \text{ and } -\frac{|\tau|}{2} \leq \Im[u] \leq \frac{|\tau|}{2}\},$$

together with the function

$$\varphi : (X_1 \setminus (-\infty, y_1]) \cup (X_2 \setminus (-\infty, y_1]) \rightarrow \Lambda, \quad z \mapsto \varphi(z) = \int_{y_1}^z \omega, \quad (4.5)$$




Figure 4: The image set of the function φ .

where the path does not cross $(-\infty, y_1]$ and lies in the same sheet as z . We denote $X_{j,+}$ and $X_{j,-}$ for the strict upper and lower half plane of the j -th sheet. The function φ maps $X_{1,+}, X_{1,-}, X_{2,+}, X_{2,-}$ into the four quadrants of Λ as shown in Figure 4. For z on the first sheet, $\varphi(z)$ has the following jumps

$$\varphi_+(z) + \varphi_-(z) = 0, \quad z \in (y_2, y_1), \quad (4.6)$$

$$\varphi_+(z) + \varphi_-(z) = 1, \quad z \in (-\infty, 0), \quad (4.7)$$

$$\varphi_+(z) - \varphi_-(z) = \tau, \quad z \in (0, y_2). \quad (4.8)$$

At the four branch points $y_1, y_2, 0, \infty$ on the first sheet, we have

$$\varphi(y_1) = 0, \quad \varphi_{\pm}(y_2) = \pm \frac{\tau}{2}, \quad \varphi_{\pm}(0) = \frac{1}{2} \pm \frac{\tau}{2}, \quad \varphi(\infty) = \frac{1}{2}, \quad (4.9)$$

see Figure 4. The Abel map is closely related to φ , so we denote it by φ_A , and is given by

$$\varphi_A : X \rightarrow \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), \quad z \mapsto \varphi_A(z) := \varphi(z) \mod (\mathbb{Z} + \tau\mathbb{Z}). \quad (4.10)$$

The Abel map is a bijection with the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ and is analytic on the full Riemann surface (as opposed to φ which presents discontinuities, as can be seen in (4.6)-(4.8)). There is an explicit expression for its inverse $\varphi_A^{-1}(u)$ in terms of the Jacobi elliptic function sn [29, eq 22.2.4]:

$$\varphi_A^{-1}(u) = y_1 + (y_2 - y_1) \text{sn}^2 \left(\frac{iu\sqrt{y_1}}{2c_0}, 1 - \frac{y_2}{y_1} \right).$$

We introduce the following ratio of θ -functions (cf. Baker-Akhiezer functions)

$$\mathcal{F}(z; \hat{d}, \nu) = \frac{\theta(z + \hat{d} + \nu)}{\theta(z + \hat{d})}, \quad z, \hat{d} \in \mathbb{C}, \quad \nu \in \mathbb{R}.$$

This function has a simple pole at $z = -\hat{d} + \frac{1}{2} + \frac{\tau}{2} \mod (\mathbb{Z} + \tau\mathbb{Z})$. From (1.9) and (4.6)-(4.8), for $z \in X_1$ we have

$$\begin{aligned} \mathcal{F}(\varphi_+(z); \hat{d}, \nu) &= \mathcal{F}(\varphi_-(z); -\hat{d}, -\nu), & z \in (-\infty, 0) \cup (y_2, y_1), \\ \mathcal{F}(\varphi_+(z); \hat{d}, \nu) &= e^{-2\pi i \nu} \mathcal{F}(\varphi_-(z); \hat{d}, \nu), & z \in (0, y_2). \end{aligned}$$

Let us consider the matrix

$$Q_1^{(\infty)}(z) := \begin{pmatrix} \mathcal{F}(\varphi(z); -\widehat{d}_1, -\nu) & \mathcal{F}(\varphi(z); \widehat{d}_1, \nu) \\ \mathcal{F}(\varphi(z); \widehat{d}_2, -\nu) & \mathcal{F}(\varphi(z); -\widehat{d}_2, \nu) \end{pmatrix}, \quad z \in \mathbb{C} \setminus (-\infty, y_1], \quad (4.11)$$

where $\varphi(z)$ is interpreted as the value of φ at the point z on the upper sheet X_1 . The matrix $Q_1^{(\infty)}(z)$ satisfies the jump conditions

$$\begin{aligned} Q_{1,+}^{(\infty)}(z) &= Q_{1,-}^{(\infty)}(z) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & z \in (-\infty, 0) \cup (y_2, y_1), \\ Q_{1,+}^{(\infty)}(z) &= Q_{1,-}^{(\infty)}(z) e^{2\pi i \nu \sigma_3}, & z \in (0, y_2). \end{aligned}$$

Note that $\det Q_1^{(\infty)}$ is not identically 1. To ensure that $Q_1^{(\infty)}$ has no pole at ∞ , we assume from now that $\widehat{d}_1, \widehat{d}_2 \neq \frac{\tau}{2} \pmod{(\mathbb{Z} + \tau\mathbb{Z})}$. We define

$$\beta(z) = \sqrt[4]{\frac{z(z-y_1)}{z-y_2}}, \quad z \in \mathbb{C} \setminus ((-\infty, 0) \cup [y_2, y_1]),$$

where the principal branch is chosen for each branch so $\beta(z) > 0$ for $z > y_1$. It can be verified that the matrix

$$Q_2^{(\infty)}(z) := \beta(z)^{\sigma_3} M^{-1} \quad (4.12)$$

is analytic in $\mathbb{C} \setminus ((-\infty, 0] \cup [y_2, y_1])$, satisfies the jumps

$$Q_{2,+}^{(\infty)}(z) = Q_{2,-}^{(\infty)}(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad z \in (-\infty, 0) \cup (y_2, y_1), \quad (4.13)$$

with asymptotic behavior at ∞ given by

$$Q_2^{(\infty)}(z) = (I + \mathcal{O}(z^{-1})) z^{\frac{\sigma_3}{4}} M^{-1}, \quad \text{as } z \rightarrow \infty. \quad (4.14)$$

Therefore, the matrix

$$Q_3^{(\infty)}(z) := \frac{\beta(z)^{\sigma_3}}{\sqrt{2}} \begin{pmatrix} \mathcal{F}(\varphi(z); -\widehat{d}_1, -\nu) & -i\mathcal{F}(\varphi(z); \widehat{d}_1, \nu) \\ -i\mathcal{F}(\varphi(z); \widehat{d}_2, -\nu) & \mathcal{F}(\varphi(z); -\widehat{d}_2, \nu) \end{pmatrix} \quad (4.15)$$

has the jumps

$$\begin{aligned} Q_{3,+}^{(\infty)}(z) &= Q_{3,-}^{(\infty)}(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, & z \in (-\infty, 0) \cup (y_2, y_1), \\ Q_{3,+}^{(\infty)}(z) &= Q_{3,-}^{(\infty)}(z) e^{2\pi i \nu \sigma_3}, & z \in (0, y_2). \end{aligned}$$

However, $Q_3^{(\infty)}(z)$ is not a solution to the RH problem for $P^{(\infty)}(z)$ because it does not have the required behavior at ∞ . Indeed, as $z \rightarrow \infty$, $z \notin (-\infty, 0]$, we have

$$\begin{aligned} Q_3^{(\infty)}(z) M z^{-\frac{\sigma_3}{4}} &= \frac{1}{2} \begin{pmatrix} [\mathcal{F}(\varphi(z); -\widehat{d}_1, -\nu) + \mathcal{F}(\varphi(z); \widehat{d}_1, \nu)] \beta(z) z^{-\frac{1}{4}} & \cdots \\ [\mathcal{F}(\varphi(z); -\widehat{d}_2, \nu) - \mathcal{F}(\varphi(z); \widehat{d}_2, -\nu)] i \beta(z)^{-1} z^{-\frac{1}{4}} & \cdots \\ \cdots & \cdots \\ [\mathcal{F}(\varphi(z); -\widehat{d}_1, -\nu) - \mathcal{F}(\varphi(z); \widehat{d}_1, \nu)] i \beta(z) z^{\frac{1}{4}} & \cdots \\ [\mathcal{F}(\varphi(z); \widehat{d}_2, -\nu) + \mathcal{F}(\varphi(z); -\widehat{d}_2, \nu)] \beta(z)^{-1} z^{\frac{1}{4}} & \cdots \end{pmatrix} = \tilde{Q} + \mathcal{O}(z^{-1}) \end{aligned} \quad (4.16)$$

where the leading coefficient \tilde{Q} is given by

$$\tilde{Q} = \begin{pmatrix} \mathcal{F}(\frac{1}{2}; \widehat{d}_1, \nu) & 2ic_0 \mathcal{F}'(\frac{1}{2}; \widehat{d}_1, \nu) \\ 0 & \mathcal{F}(\frac{1}{2}; -\widehat{d}_2, \nu) \end{pmatrix},$$

and where we have used the expansions

$$\begin{aligned}\varphi(z) &= \frac{1}{2} - 2c_0 z^{-\frac{1}{2}} + \mathcal{O}(z^{-\frac{3}{2}}), \\ \mathcal{F}(\varphi(z); \widehat{d}, \nu) &= \mathcal{F}(\tfrac{1}{2}; \widehat{d}, \nu) - 2c_0 \mathcal{F}'(\tfrac{1}{2}; \widehat{d}, \nu) z^{-\frac{1}{2}} + \mathcal{O}(z^{-1}), \\ \mathcal{F}(\varphi(z); -\widehat{d}, -\nu) &= \mathcal{F}(-\varphi(z) + 1; \widehat{d}, \nu) = \mathcal{F}(\tfrac{1}{2}; \widehat{d}, \nu) + 2c_0 \mathcal{F}'(\tfrac{1}{2}; \widehat{d}, \nu) z^{-\frac{1}{2}} + \mathcal{O}(z^{-1}),\end{aligned}$$

as $z \rightarrow \infty$, $z \notin (-\infty, 0]$. Since $\widehat{d}_1, \widehat{d}_2 \neq \frac{\tau}{2} \pmod{\mathbb{Z} + \tau\mathbb{Z}}$, the quantities $\mathcal{F}(\tfrac{1}{2}; \widehat{d}_1, \nu)$ and $\mathcal{F}(\tfrac{1}{2}; -\widehat{d}_2, \nu)$ are well-defined. We also assume that $\Im[\widehat{d}_1], \Im[\widehat{d}_2] \neq (\tfrac{1}{2} + n)\Im[\tau], n \in \mathbb{Z}$, so that $\mathcal{F}(\tfrac{1}{2}; \widehat{d}_1, \nu)$ and $\mathcal{F}(\tfrac{1}{2}; -\widehat{d}_2, \nu)$ are both different from 0 for all values of $\nu \in \mathbb{R}$. This implies that \tilde{Q} is invertible, therefore we can normalize $Q_3^{(\infty)}(z)$ at ∞ . Thus we define

$$P^{(\infty)}(z) := \tilde{Q}^{-1} Q_3^{(\infty)}(z) = \begin{pmatrix} \frac{1}{\mathcal{G}(\frac{1}{2})} & \frac{-ic_{\mathcal{F}}}{\mathcal{H}(\frac{1}{2})} \\ 0 & \frac{1}{\mathcal{H}(\frac{1}{2})} \end{pmatrix} \frac{\beta(z)^{\sigma_3}}{\sqrt{2}} \begin{pmatrix} \mathcal{G}(-\varphi(z)) & -i\mathcal{G}(\varphi(z)) \\ -i\mathcal{H}(-\varphi(z)) & \mathcal{H}(\varphi(z)) \end{pmatrix} \quad (4.17)$$

where

$$\mathcal{G}(z) = \mathcal{F}(z; \widehat{d}_1, \nu), \quad \mathcal{H}(z) = \mathcal{F}(z; -\widehat{d}_2, \nu), \quad c_{\mathcal{F}} = 2c_0(\log \mathcal{G})'(\tfrac{1}{2}).$$

The above matrix $P^{(\infty)}(z)$ has the same jumps as $Q_3^{(\infty)}(z)$ (since the multiplication by \tilde{Q}^{-1} is taken on the left), and has the prescribed asymptotic behavior as $z \rightarrow \infty$ given by (4.2). It remains to choose \widehat{d}_1 and \widehat{d}_2 appropriately so that $P^{(\infty)}(z)$ has no poles. Since β^{-1} vanishes at $z = y_2$, we choose $\widehat{d}_2 = \frac{1}{2}$ so that the pole of $\mathcal{H}(\pm\varphi(z))$ also lies at y_2 . Thus at the branch point y_2 , we have

$$\mathcal{H}(\pm\varphi(z))\beta(z)^{-1} = \mathcal{O}((z - y_2)^{-\frac{1}{4}}), \quad \text{as } z \rightarrow y_2. \quad (4.18)$$

Let us now inspect the first line of the right-most matrix in (4.17). Since β vanishes at 0 and y_1 , we have two choices for \widehat{d}_1 (see (4.9)):

$$\widehat{d}_1 = 0 \quad \text{or} \quad \widehat{d}_1 = \frac{1}{2} + \frac{\tau}{2}. \quad (4.19)$$

However, the parameter ν will be such that $\nu \rightarrow -\infty$ as $r \rightarrow +\infty$, so we need $\Im[\widehat{d}_1] \neq \frac{\Im[\tau]}{2}$ in order to ensure that \tilde{Q} is invertible, see the discussion above (4.17). Therefore we choose $\widehat{d}_1 = 0$. The choice for ν is imposed from the jumps on $(0, y_2)$, see (4.1), therefore we must have

$$\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}. \quad (4.20)$$

To summarize, the matrix-valued function $P^{(\infty)}(z)$ given by (4.17) with $\widehat{d}_1 = 0$, $\widehat{d}_2 = \frac{1}{2}$ and $\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}$ satisfies the RH problem for $P^{(\infty)}$. Furthermore, we have $\mathcal{H}(z) = \mathcal{G}(z - \frac{1}{2})$, so that (4.17) can be rewritten as

$$P^{(\infty)}(z) := \tilde{Q}^{-1} Q_3^{(\infty)}(z) = \begin{pmatrix} \frac{1}{\mathcal{G}(\frac{1}{2})} & \frac{-ic_{\mathcal{F}}}{\mathcal{G}(0)} \\ 0 & \frac{1}{\mathcal{G}(0)} \end{pmatrix} \frac{\beta(z)^{\sigma_3}}{\sqrt{2}} \begin{pmatrix} \mathcal{G}(-\varphi(z)) & -i\mathcal{G}(\varphi(z)) \\ -i\mathcal{G}(-\varphi(z) - \frac{1}{2}) & \mathcal{G}(\varphi(z) - \frac{1}{2}) \end{pmatrix} \quad (4.21)$$

where

$$\mathcal{G}(z) = \mathcal{F}(z; 0, \nu) = \frac{\theta(z + \nu)}{\theta(z)}.$$

The function \mathcal{G} is analytic in $\mathbb{C} \setminus \{\frac{1+\tau}{2} + n + m\tau : n, m \in \mathbb{Z}\}$ and satisfies

$$\mathcal{G}(z + 1) = \mathcal{G}(z), \quad \mathcal{G}(z + \tau) = e^{-2\pi i \nu} \mathcal{G}(z). \quad (4.22)$$

In the following subsections, we obtain expressions for the first four terms of the asymptotics of $P^{(\infty)}(z)$ as $z \rightarrow y_j$, $\Im[z] > 0$, $j = 1, 2, 3$. These coefficients will be needed in the computations of Section 6.

4.1 Asymptotics of $P^{(\infty)}(z)$ as $z \rightarrow y_1$

The asymptotics for $\beta(z)$ and $\varphi(z)$ as $z \rightarrow y_1$, $z \in \mathbb{C} \setminus (-\infty, y_1]$, are given by

$$\begin{aligned}\beta(z) &= \beta_{y_1}^{(\frac{1}{4})}(z - y_1)^{\frac{1}{4}} + \beta_{y_1}^{(\frac{5}{4})}(z - y_1)^{\frac{5}{4}} + \mathcal{O}((z - y_1)^{\frac{9}{4}}), \\ \beta_{y_1}^{(\frac{1}{4})} &= \frac{y_1^{\frac{1}{4}}}{(y_1 - y_2)^{\frac{1}{4}}}, \quad \beta_{y_1}^{(\frac{5}{4})} = \frac{-y_2}{4y_1^{\frac{3}{4}}(y_1 - y_2)^{\frac{5}{4}}}, \\ \varphi(z) &= \varphi_{y_1}^{(\frac{1}{2})}(z - y_1)^{\frac{1}{2}} + \varphi_{y_1}^{(\frac{3}{2})}(z - y_1)^{\frac{3}{2}} + \mathcal{O}((z - y_1)^{\frac{5}{2}}), \\ \varphi_{y_1}^{(\frac{1}{2})} &= \frac{2c_0}{\sqrt{y_1}\sqrt{y_1 - y_2}}, \quad \varphi_{y_1}^{(\frac{3}{2})} = \frac{-c_0(2y_1 - y_2)}{3y_1^{\frac{3}{2}}(y_1 - y_2)^{\frac{3}{2}}}.\end{aligned}\tag{4.23}$$

The asymptotics for $P^{(\infty)}(z)$ as $z \rightarrow y_1$ are given by

$$\begin{aligned}P^{(\infty)}(z) &= \sum_{j=0}^3 (P^{(\infty)})_{y_1}^{(-\frac{1}{4} + \frac{j}{2})} (z - y_1)^{-\frac{1}{4} + \frac{j}{2}} + \mathcal{O}((z - y_1)^{\frac{7}{4}}), \\ (P^{(\infty)})_{y_1}^{(-\frac{1}{4})} &= \frac{-\mathcal{G}(\frac{1}{2})}{\sqrt{2}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & ic_{\mathcal{F}} \\ i & -1 \end{pmatrix}, \\ (P^{(\infty)})_{y_1}^{(\frac{1}{4})} &= \frac{\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)}{\sqrt{2}\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & -i \\ 0 & 0 \end{pmatrix} + \frac{\varphi_{y_1}^{(\frac{1}{2})}\mathcal{G}'(\frac{1}{2})}{\sqrt{2}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & -ic_{\mathcal{F}} \\ i & 1 \end{pmatrix}, \\ (P^{(\infty)})_{y_1}^{(\frac{3}{4})} &= \frac{-\varphi_{y_1}^{(\frac{1}{2})}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}'(0)}{\sqrt{2}\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & i \\ 0 & 0 \end{pmatrix} + \frac{\frac{\beta_{y_1}^{(\frac{5}{4})}}{\beta_{y_1}^{(\frac{1}{4})}}\mathcal{G}(\frac{1}{2}) - \frac{1}{2}(\varphi_{y_1}^{(\frac{1}{2})})^2\mathcal{G}''(\frac{1}{2})}{\sqrt{2}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & ic_{\mathcal{F}} \\ i & -1 \end{pmatrix}, \\ (P^{(\infty)})_{y_1}^{(\frac{5}{4})} &= \frac{\beta_{y_1}^{(\frac{5}{4})}\mathcal{G}(0) + \frac{1}{2}\beta_{y_1}^{(\frac{1}{4})}(\varphi_{y_1}^{(\frac{1}{2})})^2\mathcal{G}''(0)}{\sqrt{2}\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & -i \\ 0 & 0 \end{pmatrix} \\ &+ \frac{\varphi_{y_1}^{(\frac{3}{2})}\mathcal{G}'(\frac{1}{2}) + \frac{1}{6}(\varphi_{y_1}^{(\frac{1}{2})})^3\mathcal{G}'''(\frac{1}{2}) - \frac{\beta_{y_1}^{(\frac{5}{4})}}{\beta_{y_1}^{(\frac{1}{4})}}\varphi_{y_1}^{(\frac{1}{2})}\mathcal{G}'(\frac{1}{2})}{\sqrt{2}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & -ic_{\mathcal{F}} \\ i & 1 \end{pmatrix}.\end{aligned}$$

4.2 Asymptotics of $P^{(\infty)}(z)$ as $z \rightarrow y_2$ from $\Im[z] > 0$

We recall that $\frac{1+\tau}{2}$ is a simple zero of $z \mapsto \theta(z)$. Thus the function $z \mapsto \mathcal{G}(z)$ has a simple pole at $z = \frac{1+\tau}{2}$. Since $\varphi_+(y_2) = \frac{\tau}{2}$, the entries on the second row of $P^{(\infty)}(z)$ blow up as $z \rightarrow y_2$, $\Im[z] > 0$. Let us define

$$z \mapsto \tilde{\mathcal{G}}(z) := \mathcal{G}(z)(z - \frac{1+\tau}{2}).\tag{4.24}$$

Using the relations (4.22) and (4.24), before expanding $P^{(\infty)}(z)$ as $z \rightarrow y_2$, $\Im[z] > 0$, we rewrite $P^{(\infty)}(z)$ as follows

$$P^{(\infty)}(z) = \begin{pmatrix} \frac{1}{\mathcal{G}(\frac{1}{2})} & \frac{-ic_{\mathcal{F}}}{\mathcal{G}(0)} \\ 0 & \frac{1}{\mathcal{G}(0)} \end{pmatrix} \frac{\beta(z)^{\sigma_3}}{\sqrt{2}} \begin{pmatrix} e^{2\pi i \nu} \mathcal{G}(\tau - \varphi(z)) & -i\mathcal{G}(\varphi(z)) \\ -ie^{2\pi i \nu} \frac{\tilde{\mathcal{G}}(-\varphi(z) + \frac{1}{2} + \tau)}{\frac{\tau}{2} - \varphi(z)} & \frac{\tilde{\mathcal{G}}(\varphi(z) + \frac{1}{2})}{\varphi(z) - \frac{\tau}{2}} \end{pmatrix}\tag{4.25}$$

The asymptotics for $\beta(z)$ and $\varphi(z)$ as $z \rightarrow y_2$, $\Im[z] > 0$, are of the form

$$\begin{aligned}\beta(z) &= \beta_{y_2}^{(-\frac{1}{4})}(z - y_2)^{-\frac{1}{4}} + \beta_{y_2}^{(\frac{3}{4})}(z - y_2)^{\frac{3}{4}} + \mathcal{O}((z - y_2)^{\frac{7}{4}}), \\ \varphi(z) &= \frac{\tau}{2} + \varphi_{y_2}^{(\frac{1}{2})}(z - y_2)^{\frac{1}{2}} + \varphi_{y_2}^{(\frac{3}{2})}(z - y_2)^{\frac{3}{2}} + \mathcal{O}((z - y_2)^{\frac{5}{2}}),\end{aligned}$$

and the asymptotics for $P^{(\infty)}(z)$ as $z \rightarrow y_2$, $\Im[z] > 0$, are of the form

$$P^{(\infty)}(z) = \sum_{j=0}^3 (P^{(\infty)})_{y_2}^{(-\frac{1}{4}+\frac{j}{2})} (z - y_2)^{-\frac{1}{4}+\frac{j}{2}} + \mathcal{O}((z - y_2)^{\frac{7}{4}}).$$

The explicit values for $(P^{(\infty)})_{y_2}^{(-\frac{1}{4}+\frac{j}{2})}$, $j = 0, \dots, 3$ are all needed to prove Theorem 1.1. They can be obtained after long but straightforward computations in terms of $\beta_{y_2}^{(-\frac{1}{4})}$, $\beta_{y_2}^{(\frac{3}{4})}$, $\varphi_{y_2}^{(\frac{1}{2})}$, $\varphi_{y_2}^{(\frac{3}{2})}$, $\mathcal{G}(0)$, $\mathcal{G}(\frac{1}{2})$, $\mathcal{G}^{(k)}(\frac{\tau}{2})$, $\tilde{\mathcal{G}}^{(k)}(\frac{1+\tau}{2})$, $k = 0, 1, 2, 3$, but for conciseness we do not write them down.

4.3 Asymptotics of $P^{(\infty)}(z)$ as $z \rightarrow y_3 = 0$ from $\Im[z] > 0$

Using the relations (4.22) and (4.24), before expanding $P^{(\infty)}(z)$ as $z \rightarrow y_3$, $\Im[z] > 0$, we rewrite $P^{(\infty)}(z)$ as follows

$$P^{(\infty)}(z) = \begin{pmatrix} \frac{1}{\mathcal{G}(\frac{1}{2})} & \frac{-i\mathcal{C}\mathcal{F}}{\mathcal{G}(0)} \\ 0 & \frac{1}{\mathcal{G}(0)} \end{pmatrix} \frac{\beta(z)^{\sigma_3}}{\sqrt{2}} \begin{pmatrix} e^{2\pi i\nu} \frac{\tilde{\mathcal{G}}(1+\tau-\varphi(z))}{\frac{1+\tau}{2}-\varphi(z)} & -i \frac{\tilde{\mathcal{G}}(\varphi(z))}{\varphi(z)-\frac{1+\tau}{2}} \\ -ie^{2\pi i\nu} \mathcal{G}(\frac{1}{2}+\tau-\varphi(z)) & \mathcal{G}(\varphi(z)-\frac{1}{2}) \end{pmatrix}$$

The asymptotics for $\beta(z)$ and $\varphi(z)$ as $z \rightarrow y_3 = 0$, $\Im[z] > 0$, are of the form

$$\begin{aligned} \beta(z) &= \beta_{y_3}^{(\frac{1}{4})} z^{\frac{1}{4}} + \beta_{y_3}^{(\frac{5}{4})} z^{\frac{5}{4}} + \mathcal{O}(z^{\frac{9}{4}}), \\ \varphi(z) &= \frac{1+\tau}{2} + \varphi_{y_3}^{(\frac{1}{2})} z^{\frac{1}{2}} + \varphi_{y_3}^{(\frac{3}{2})} z^{\frac{3}{2}} + \mathcal{O}(z^{\frac{5}{2}}), \end{aligned}$$

and the asymptotics for $P^{(\infty)}(z)$ as $z \rightarrow y_3 = 0$, $\Im[z] > 0$, are of the form

$$P^{(\infty)}(z) = \sum_{j=0}^3 (P^{(\infty)})_{y_3}^{(-\frac{1}{4}+\frac{j}{2})} z^{-\frac{1}{4}+\frac{j}{2}} + \mathcal{O}(z^{\frac{7}{4}}).$$

The explicit values for $(P^{(\infty)})_{y_3}^{(-\frac{1}{4}+\frac{j}{2})}$, $j = 0, \dots, 3$ are also all needed to prove Theorem 1.1. They can be expressed after long but straightforward computations in terms of $\beta_{y_3}^{(\frac{1}{4})}$, $\beta_{y_3}^{(\frac{5}{4})}$, $\varphi_{y_3}^{(\frac{1}{2})}$, $\varphi_{y_3}^{(\frac{3}{2})}$, $\mathcal{G}(0)$, $\mathcal{G}(\frac{1}{2})$, $\mathcal{G}^{(k)}(\frac{\tau}{2})$, $\tilde{\mathcal{G}}^{(k)}(\frac{1+\tau}{2})$, $k = 0, 1, 2, 3$, and we do not write them down for conciseness.

4.4 Asymptotics of $P^{(\infty)}(z)^{-1}$ as $z \rightarrow y_k$ from $\Im[z] > 0$, $k = 1, 2, 3$

Since $\det P^{(\infty)} \equiv 1$, we have

$$P^{(\infty)}(z)^{-1} = \begin{pmatrix} P_{22}^{(\infty)}(z) & -P_{12}^{(\infty)}(z) \\ -P_{21}^{(\infty)}(z) & P_{11}^{(\infty)}(z) \end{pmatrix},$$

and we deduce from Subsections 4.1, 4.2 and 4.3 the following asymptotic expansions

$$\begin{aligned} P^{(\infty)}(z)^{-1} &= \sum_{j=0}^3 (P_{\text{inv}}^{(\infty)})_{y_k}^{(-\frac{1}{4}+\frac{j}{2})} (z - y_k)^{-\frac{1}{4}+\frac{j}{2}} + \mathcal{O}((z - y_k)^{\frac{7}{4}}), \quad k = 1, 2, 3, \\ (P_{\text{inv}}^{(\infty)})_{y_k}^{(-\frac{1}{4}+\frac{j}{2})} &= \begin{pmatrix} (P^{(\infty)})_{y_k,22}^{(-\frac{1}{4}+\frac{j}{2})} & -(P^{(\infty)})_{y_k,12}^{(-\frac{1}{4}+\frac{j}{2})} \\ -(P^{(\infty)})_{y_k,21}^{(-\frac{1}{4}+\frac{j}{2})} & (P^{(\infty)})_{y_k,11}^{(-\frac{1}{4}+\frac{j}{2})} \end{pmatrix}, \quad k = 1, 2, 3, \quad j = 0, 1, 2, 3. \end{aligned}$$

5 Local parametrices

Let us choose $\delta > 0$ such that

$$\delta \leq \frac{\min\{-x_1, x_1 - x_2, x_2 - x_3\}}{3} \quad \text{and} \quad \mathbb{D}_{y_j} := \{z \in \mathbb{C} : |z - y_j| < \delta\} \subset N_0, \quad j = 1, 2, 3,$$

where N_0 is as described in Lemma 3.2. The local parametrix $P^{(y_j)}(z)$, $j = 1, 2, 3$, is defined for $z \in \mathbb{D}_{y_j}$ as the solution to a RH problem with the same jumps as S inside \mathbb{D}_{y_j} . Furthermore, we require that $P^{(y_j)}(z) = \mathcal{O}(\log|z - y_j|)$ as $z \rightarrow y_j$ and that $P^{(y_j)}$ matches with $P^{(\infty)}$ on the boundary of \mathbb{D}_{y_j} , in the sense that

$$P^{(y_j)}(z) = (I + o(1))P^{(\infty)}(z), \quad \text{as } r \rightarrow +\infty \quad (5.1)$$

uniformly for $z \in \partial\mathbb{D}_{y_j}$. These constructions are standard (so we give relatively short explanations) and given in terms of an explicitly solvable model RH problem Φ_{Be} which is expressed in terms of Bessel functions. This model RH problem was first derived in [16] and we present it in Appendix A for convenience.

5.1 Parametrix at y_1

Let f_{y_1} be the conformal map from \mathbb{D}_{y_1} to a neighborhood of 0 defined by

$$f_{y_1}(z) = \frac{g^2(z)}{4}. \quad (5.2)$$

The expansion of $f_{y_1}(z)$ as $z \rightarrow y_1$ is given by

$$\begin{aligned} f_{y_1}(z) &= c_{y_1}(z - y_1)(1 + c_{y_1}^{(2)}(z - y_1) + c_{y_1}^{(3)}(z - y_1)^2 + \mathcal{O}((z - y_1)^3)), \\ c_{y_1} &= \frac{p^2(y_1)}{y_1(y_1 - y_2)}, \quad c_{y_1}^{(2)} = \frac{2p'(y_1)}{3p(y_1)} - \frac{2y_1 - y_2}{3y_1(y_1 - y_2)}, \\ c_{y_1}^{(3)} &= \frac{1}{45} \left(\frac{23y_1^2 - 23y_1y_2 + 8y_2^2}{y_1^2(y_1 - y_2)^2} + \frac{5p'(y_1)^2}{p(y_1)^2} - \frac{14(2y_1 - y_2)p'(y_1)}{y_1(y_1 - y_2)p(y_1)} + \frac{9p''(y_1)}{p(y_1)} \right). \end{aligned} \quad (5.3)$$

We have freedom in the choice of $\gamma_{\pm} \subset N_0$. We choose the lenses such that $f_{y_1}(\gamma_+ \cap \mathbb{D}_{y_1}) \subset e^{\frac{2\pi i}{3}}\mathbb{R}^+$ and $f_{y_1}(\gamma_- \cap \mathbb{D}_{y_1}) \subset e^{\frac{2\pi i}{3}}\mathbb{R}^-$. We conclude from the RH problem for Φ_{Be} presented in Section A that the matrix

$$\Phi_{\text{Be}}(r^3 f_{y_1}(z)) e^{-r^{\frac{3}{2}}g(z)\sigma_3} \quad (5.4)$$

has the same jumps as $S(z)$ inside the disk $z \in \mathbb{D}_{y_1}$, see (3.22)-(3.24). Let us define $P^{(y_1)}$ by

$$P^{(y_1)}(z) := E_{y_1}(z) \Phi_{\text{Be}}(r^3 f_{y_1}(z)) e^{-r^{\frac{3}{2}}g(z)\sigma_3}, \quad (5.5)$$

$$E_{y_1}(z) = P^{(\infty)}(z) M^{-1} \left(2\pi r^{\frac{3}{2}} f_{y_1}(z)^{\frac{1}{2}} \right)^{\frac{\sigma_3}{2}}. \quad (5.6)$$

It can be verified from the jumps of $P^{(\infty)}(z)$ that the prefactor $E_{y_1}(z)$ is analytic for $z \in \mathbb{D}_{y_1}$. Thus, $P^{(y_1)}(z)$ has the (exact) same jumps as and endpoint behavior, see (A.3), as $S(z)$ for $z \in \mathbb{D}_{y_1}$. Furthermore, due to (5.3) and (A.2), we have

$$P^{(y_1)}(z) P^{(\infty)}(z)^{-1} = I + \frac{P^{(\infty)}(z) \Phi_{\text{Be},1} P^{(\infty)}(z)^{-1}}{r^{\frac{3}{2}} f_{y_1}(z)^{\frac{1}{2}}} + \mathcal{O}(r^{-3}) \quad (5.7)$$

as $r \rightarrow +\infty$ uniformly for $z \in \partial\mathbb{D}_{y_1}$. In Section 8 we will need the first columns of $E_{y_1}(y_1)$ and $E'_{y_1}(y_1)$. After some computations, we find

$$\begin{aligned} E_{y_1}(y_1) &= \frac{-\sqrt{2\pi} c_{y_1}^{\frac{1}{4}} \mathcal{G}(\frac{1}{2})}{\beta_{y_1}^{(\frac{1}{4})} \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} r^{\frac{3}{4}} + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}}, \\ E'_{y_1}(y_1) &= \sqrt{2\pi} c_{y_1}^{\frac{1}{4}} r^{\frac{3}{4}} \left\{ \frac{\left(\frac{\beta_{y_1}^{(\frac{5}{4})}}{\beta_{y_1}^{(\frac{1}{4})}} - \frac{1}{4} c_{y_1}^{(2)} \right) \mathcal{G}(\frac{1}{2}) - \frac{1}{2} (\varphi_{y_1}^{(\frac{1}{2})})^2 \mathcal{G}''(\frac{1}{2})}{\beta_{y_1}^{(\frac{1}{4})} \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} \right. \\ &\quad \left. - \frac{\varphi_{y_1}^{(\frac{1}{2})} \beta_{y_1}^{(\frac{1}{4})} \mathcal{G}'(0)}{\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\} + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}} \end{aligned} \quad (5.8)$$

where \star denotes unnecessary constants.

5.2 Parametrix at y_2

Let f_{y_2} be the conformal map from \mathbb{D}_{y_2} to a neighborhood of 0 defined by

$$f_{y_2}(z) = -\frac{1}{4} \left(g(z) \mp \frac{i\Omega}{2} \right)^2, \quad (5.9)$$

where we take $-/+$ when z is above/below the real axis. Its expansion as $z \rightarrow y_2$ is given by

$$\begin{aligned} f_{y_2}(z) &= c_{y_2}(z - y_2) \left(1 + c_{y_2}^{(2)}(z - y_2) + c_{y_2}^{(3)}(z - y_2)^2 + \mathcal{O}((z - y_2)^3) \right), \\ c_{y_2} &= \frac{p^2(y_2)}{y_2(y_1 - y_2)} > 0, \quad c_{y_2}^{(2)} = \frac{2p'(y_2)}{3p(y_2)} - \frac{y_1 - 2y_2}{3y_2(y_1 - y_2)}, \\ c_{y_2}^{(3)} &= \frac{1}{45} \left(\frac{8y_1^2 - 23y_1y_2 + 23y_2^2}{(y_1 - y_2)^2y_2^2} + \frac{5p'(y_2)^2}{p(y_2)^2} - \frac{14(y_1 - 2y_2)p'(y_2)}{(y_1 - y_2)y_2p(y_2)} + \frac{9p''(y_2)}{p(y_2)} \right). \end{aligned} \quad (5.10)$$

We choose the lenses $\gamma_+ \subset N_0$ and $\gamma_- \subset N_0$ such that $-f_{y_2}(\gamma_{\pm} \cap \mathbb{D}_{y_2}) \subset e^{\pm \frac{2\pi i}{3}} \mathbb{R}^+$, and we define

$$\begin{aligned} P^{(y_2)}(z) &= E_{y_2}(z) \sigma_3 \Phi_{\text{Be}}(-r^3 f_{y_2}(z)) \sigma_3 e^{-r^{\frac{3}{2}} g(z) \sigma_3}, \\ E_{y_2}(z) &= P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} M \left(2\pi r^{\frac{3}{2}} (-f_{y_2}(z))^{\frac{1}{2}} \right)^{\frac{\sigma_3}{2}}. \end{aligned} \quad (5.11)$$

It can be verified that $E_{y_2}(z)$ is analytic in the disk \mathbb{D}_{y_2} and that $P^{(y_2)}(z) = \mathcal{O}(\log |z - y_2|)$ as $z \rightarrow y_2$, see (A.3). From (5.9) and (A.2), we also conclude that

$$P^{(y_2)}(z) P^{(\infty)}(z)^{-1} = I + \frac{P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} \Phi_{\text{Be},1} \sigma_3 e^{\mp \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} P^{(\infty)}(z)^{-1}}{r^{\frac{3}{2}} (-f_{y_2}(z))^{\frac{1}{2}}} + \mathcal{O}(r^{-3}) \quad (5.12)$$

as $r \rightarrow +\infty$ uniformly for $z \in \partial \mathbb{D}_{y_2}$. Furthermore, we also verify that

$$\begin{aligned} E_{y_2}(y_2) &= e^{-\frac{\pi i}{4}} \sqrt{2\pi} c_{y_2}^{\frac{1}{4}} e^{\pi i \nu} r^{\frac{3}{4}} \left\{ \frac{\tilde{\mathcal{G}}(\frac{1+\tau}{2})}{\varphi_{y_2}^{(\frac{1}{2})} \beta_{y_2}^{(-\frac{1}{4})} \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} + \frac{\beta_{y_2}^{(-\frac{1}{4})} \mathcal{G}(\frac{\tau}{2})}{\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\} + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}}, \\ E'_{y_2}(y_2) &= e^{-\frac{\pi i}{4}} \sqrt{2\pi} c_{y_2}^{\frac{1}{4}} e^{\pi i \nu} r^{\frac{3}{4}} \left\{ \frac{(\beta_{y_2}^{(\frac{3}{4})} + \frac{1}{4} c_{y_2}^{(2)} \beta_{y_2}^{(-\frac{1}{4})}) \mathcal{G}(\frac{\tau}{2}) + \frac{1}{2} (\varphi_{y_2}^{(\frac{1}{2})})^2 \beta_{y_2}^{(-\frac{1}{4})} \mathcal{G}''(\frac{\tau}{2})}{\mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right. \\ &\quad \left. + \frac{(\frac{1}{4} c_{y_2}^{(2)} \varphi_{y_2}^{(\frac{1}{2})} \beta_{y_2}^{(-\frac{1}{4})} - \varphi_{y_2}^{(\frac{1}{2})} \beta_{y_2}^{(\frac{3}{4})} - \varphi_{y_2}^{(\frac{3}{2})} \beta_{y_2}^{(-\frac{1}{4})}) \tilde{\mathcal{G}}(\frac{1+\tau}{2}) + \frac{1}{2} (\varphi_{y_2}^{(\frac{1}{2})})^3 \beta_{y_2}^{(-\frac{1}{4})} \tilde{\mathcal{G}}''(\frac{1+\tau}{2})}{(\varphi_{y_2}^{(\frac{1}{2})})^2 (\beta_{y_2}^{(-\frac{1}{4})})^2 \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} \right\} \\ &\quad + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}}, \end{aligned} \quad (5.13)$$

as $r \rightarrow +\infty$, where \star denotes unnecessary constants.

5.3 Parametrix at $y_3 = 0$

Let f_{y_3} be the conformal map from \mathbb{D}_{y_3} to a neighborhood of 0 defined by

$$f_{y_3}(z) = \frac{1}{4} \left(g(z) \mp \frac{i\Omega}{2} \right)^2, \quad (5.14)$$

where we take the $-/+$ sign when z is above/below the real axis. The expansion of $f_{y_3}(z)$ as $z \rightarrow 0$ is given by

$$\begin{aligned} f_{y_3}(z) &= c_{y_3} z \left(1 + c_{y_3}^{(2)} z + c_{y_3}^{(3)} z^2 + \mathcal{O}(z^3) \right), \\ c_{y_3} &= \frac{p^2(0)}{y_1 y_2}, \quad c_{y_3}^{(2)} = \frac{2p'(0)}{3p(0)} + \frac{y_1 + y_2}{3y_1 y_2}, \\ c_{y_3}^{(3)} &= \frac{1}{45} \left(\frac{8y_1^2 + 7y_1 y_2 + 8y_2^2}{y_1^2 y_2^2} + \frac{5p'(0)^2}{p(0)^2} + \frac{14(y_1 + y_2)p'(0)}{y_1 y_2 p(0)} + \frac{9p''(0)}{p(0)} \right). \end{aligned} \quad (5.15)$$

We choose the lenses $\gamma_+ \subset N_0$ and $\gamma_- \subset N_0$ such that $f_{y_3}(\gamma_\pm \cap \mathbb{D}_{y_3}) \subset e^{\pm \frac{2\pi i}{3}} \mathbb{R}^+$, and we define

$$\begin{aligned} P^{(y_3)}(z) &= E_{y_3}(z) \Phi_{\text{Be}}(r^3 f_{y_3}(z)) e^{-r^{\frac{3}{2}} g(z) \sigma_3}, \\ E_{y_3}(z) &= P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} M^{-1} \left(2\pi r^{\frac{3}{2}} (f_{y_3}(z))^{\frac{1}{2}} \right)^{\frac{\sigma_3}{2}}. \end{aligned} \quad (5.16)$$

It can be verified that $E_{y_3}(z)$ is analytic inside the disk \mathbb{D}_{y_3} , $P^{(y_3)}(z) = \mathcal{O}(\log|z|)$ as $z \rightarrow 0$, see (A.3), and that

$$P^{(y_3)}(z) P^{(\infty)}(z)^{-1} = I + \frac{P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} \Phi_{\text{Be},1} e^{\mp \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} P^{(\infty)}(z)^{-1}}{r^{\frac{3}{2}} f_{y_3}(z)^{\frac{1}{2}}} + \mathcal{O}(r^{-3}) \quad (5.17)$$

as $r \rightarrow +\infty$ uniformly for $z \in \partial\mathbb{D}_{y_3}$. Furthermore, we verify that

$$\begin{aligned} E_{y_3}(y_3) &= -\sqrt{2\pi} c_{y_3}^{\frac{1}{4}} e^{\pi i \nu} r^{\frac{3}{4}} \left\{ \frac{\mathcal{G}(\frac{\tau}{2})}{\beta_{y_3}^{(\frac{1}{4})} \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} + \frac{\beta_{y_3}^{(\frac{1}{4})} \tilde{\mathcal{G}}(\frac{1+\tau}{2})}{\varphi_{y_3}^{(\frac{1}{2})} \mathcal{G}(\frac{1}{2})} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\} + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}}, \\ E'_{y_3}(y_3) &= -\sqrt{2\pi} c_{y_3}^{\frac{1}{4}} e^{\pi i \nu} r^{\frac{3}{4}} \left\{ \frac{\left(\frac{1}{4} c_{y_3}^{(2)} - \frac{\beta_{y_3}^{(\frac{5}{4})}}{\beta_{y_3}^{(\frac{1}{4})}} \right) \mathcal{G}(\frac{\tau}{2}) + \frac{1}{2} (\varphi_{y_3}^{(\frac{1}{2})})^2 \mathcal{G}''(\frac{\tau}{2})}{\beta_{y_3}^{(\frac{1}{4})} \mathcal{G}(0)} \begin{pmatrix} c_{\mathcal{F}} & 0 \\ i & 0 \end{pmatrix} \right. \\ &\quad \left. + \frac{\left(\frac{1}{4} c_{y_3}^{(2)} \beta_{y_3}^{(\frac{1}{4})} - \frac{\varphi_{y_3}^{(\frac{3}{2})}}{\varphi_{y_3}^{(\frac{1}{2})}} \beta_{y_3}^{(\frac{1}{4})} + \beta_{y_3}^{(\frac{5}{4})} \right) \tilde{\mathcal{G}}(\frac{1+\tau}{2}) + \frac{1}{2} (\varphi_{y_3}^{(\frac{1}{2})})^2 \beta_{y_3}^{(\frac{1}{4})} \tilde{\mathcal{G}}''(\frac{1+\tau}{2})}{\varphi_{y_3}^{(\frac{1}{2})} \mathcal{G}(\frac{1}{2})} \right\} + \begin{pmatrix} 0 & \star \\ 0 & \star \end{pmatrix} r^{-\frac{3}{4}} \end{aligned} \quad (5.18)$$

as $r \rightarrow +\infty$, where \star denotes unnecessary constants.

6 Small norm problem

In this section we show that for sufficiently large r , $P^{(\infty)}(z)$ approximates $S(z)$ for $z \in \mathbb{C} \setminus \cup_{j=1}^3 \mathbb{D}_{y_j}$ and $P^{(y_j)}(z)$ approximates $S(z)$ for $z \in \mathbb{D}_{y_j}$, $j = 1, 2, 3$. We define

$$R(z) = \begin{cases} S(z) P^{(y_j)}(z)^{-1}, & z \in \mathbb{D}_{y_j}, \quad j = 1, 2, 3, \\ S(z) P^{(\infty)}(z)^{-1}, & z \in \mathbb{C} \setminus \cup_{j=1}^3 \mathbb{D}_{y_j}. \end{cases} \quad (6.1)$$

Since $P^{(y_j)}(z)$, $j = 1, 2, 3$, has the exact same jumps as S inside the disks, $R(z)$ is analytic for $z \in \cup_{j=1}^3 \mathbb{D}_{y_j} \setminus \{y_j\}$. Furthermore, we verify from (3.26), (5.5), (5.11), (5.16) and (A.3) that $S(z) P^{(y_j)}(z)^{-1} = \mathcal{O}(\log(z - y_j))$ as $z \rightarrow y_j$, $j = 1, 2, 3$. This means that the singularities of R at y_1, y_2, y_3 are in fact removable and R is analytic in all three open disks. Let Σ_R denote the jump contour for R which is explicitly given by

$$\Sigma_R = \left(\gamma_+ \cup \gamma_- \cup \bigcup_{j=1}^3 \partial\mathbb{D}_{y_j} \right) \setminus \bigcup_{j=1}^3 \mathbb{D}_{y_j},$$

with orientation as shown in Figure 5. In particular, note that we orient the boundaries of the disks in the clockwise direction. The jumps for R are denoted by J_R ,

$$J_R : \Sigma_R \rightarrow \mathbb{C}^{2 \times 2}, \quad z \mapsto J_R(z) := R_-(z)^{-1} R_+(z),$$

and are explicitly given by

$$J_R(z) = \begin{cases} P^{(\infty)}(z) \begin{pmatrix} 1 & 0 \\ e^{-2r^{\frac{3}{2}} g(z)} & 1 \end{pmatrix} P^{(\infty)}(z)^{-1}, & z \in \gamma_+ \cup \gamma_- \setminus \cup_{j=1}^3 \mathbb{D}_{y_j} \\ P^{(y_j)}(z) P^{(\infty)}(z)^{-1}, & z \in \partial\mathbb{D}_{y_j}, \quad j = 1, 2, 3. \end{cases} \quad (6.2)$$




Figure 5: The jump contour Σ_R .

From Lemma 3.2 and (5.7), (5.12), (5.17), as $r \rightarrow \infty$ we have

$$J_R(z) = \begin{cases} I + \mathcal{O}(e^{-\tilde{c}|rz|^{\frac{3}{2}}}), & \text{uniformly for } z \in \Sigma_R \setminus \cup_{j=1}^3 \partial \mathbb{D}_{y_j}, \\ I + \frac{J_R^{(1)}(z)}{r^{\frac{3}{2}}} + \mathcal{O}(r^{-3}), & \text{uniformly for } z \in \cup_{j=1}^3 \partial \mathbb{D}_{y_j}, \end{cases} \quad (6.3)$$

where $\tilde{c} > 0$ is a sufficiently small constant, and $J_R^{(1)}(z) = \mathcal{O}(1)$ as $r \rightarrow +\infty$ uniformly for $z \in \cup_{j=1}^3 \partial \mathbb{D}_{y_j}$. It is important to recall that $P^{(\infty)}(z)$ has r dependence but this is of no consequence due to the periodicity of the θ -function in the real direction. The matrix $J_R^{(1)}(z)$ has been computed in (5.7), (5.12), (5.17), and is given by

$$J_R^{(1)}(z) = \begin{cases} \frac{P^{(\infty)}(z) \Phi_{\text{Be},1} P^{(\infty)}(z)^{-1}}{f_{y_1}(z)^{\frac{1}{2}}}, & \text{if } z \in \partial \mathbb{D}_{y_1}, \\ \frac{P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} \Phi_{\text{Be},1} \sigma_3 e^{\mp \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} P^{(\infty)}(z)^{-1}}{(-f_{y_2}(z))^{\frac{1}{2}}}, & \text{if } z \in \partial \mathbb{D}_{y_2}, \\ \frac{P^{(\infty)}(z) e^{\pm \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} \Phi_{\text{Be},1} e^{\mp \frac{i\Omega}{2} r^{\frac{3}{2}} \sigma_3} P^{(\infty)}(z)^{-1}}{f_{y_3}(z)^{\frac{1}{2}}}, & \text{if } z \in \partial \mathbb{D}_{y_3}. \end{cases} \quad (6.4)$$

By the standard theory for RH problems [17], $R(z)$ exists for sufficiently large r and satisfies

$$R(z) = I + \frac{R^{(1)}(z)}{r^{\frac{3}{2}}} + \mathcal{O}(r^{-3}), \quad (6.5)$$

as $r \rightarrow \infty$, uniformly for $z \in \mathbb{C} \setminus \Sigma_R$. The goal for the remainder of this section is to obtain explicit expressions for $R^{(1)'}(y_j)$, $j = 1, 2, 3$. Since $R(z)$ satisfies the equation

$$R(z) = I + \frac{1}{2\pi i} \int_{\Sigma_R} \frac{R_-(\xi) (J_R(\xi) - I)}{\xi - z} d\xi, \quad z \in \mathbb{C} \setminus \Sigma_R, \quad (6.6)$$

we deduce from (6.3) that

$$R^{(1)}(z) = \frac{1}{2\pi i} \int_{\bigcup_{j=1}^3 \partial \mathbb{D}_{y_j}} \frac{J_R^{(1)}(\xi)}{\xi - z} d\xi, \quad z \in \mathbb{C} \setminus \bigcup_{j=1}^3 \partial \mathbb{D}_{y_j}, \quad (6.7)$$

where we recall that the circles $\partial \mathbb{D}_{y_j}$, $j = 1, 2, 3$, have clockwise orientation. From (6.4), we note that the jumps $J_R^{(1)}$ can be analytically continued from $\cup_{j=1}^3 \partial \mathbb{D}_{y_j}$ to $\cup_{j=1}^3 (\overline{\mathbb{D}}_{y_j} \setminus \{y_j\})$, so we can evaluate (6.7) by a residue calculation. The asymptotic expansion of $J_R^{(1)}(z)$ as $z \rightarrow y_j$, $j = 1, 2, 3$, is of the form

$$J_R^{(1)}(z) = \sum_{k=-1}^N (J_R^{(1)})_{y_j}^{(k)} (z - y_j)^k + \mathcal{O}((z - y_j)^{N+1}),$$

for any $N \in \mathbb{N}_{>0}$, and therefore (6.7) can be rewritten as

$$R^{(1)}(z) = \sum_{j=1}^3 \frac{1}{z - y_j} (J_R^{(1)})_{y_j}^{(-1)}, \quad z \in \mathbb{C} \setminus \cup_{j=1}^3 \mathbb{D}_{y_j}, \quad (6.8)$$

$$R^{(1)}(z) = - \sum_{k=0}^{\infty} (J_R^{(1)})_{y_j}^{(k)} (z - y_j)^k + \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{z - y_{j'}} (J_R^{(1)})_{y_{j'}}^{(-1)}, \quad z \in \mathbb{D}_{y_j}, j = 1, 2, 3. \quad (6.9)$$

In particular, we have

$$R^{(1)'}(y_j) = -(J_R^{(1)})_{y_j}^{(1)} - \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{(y_j - y_{j'})^2} (J_R^{(1)})_{y_{j'}}^{(-1)}, \quad j = 1, 2, 3. \quad (6.10)$$

An explicit expression for $R^{(1)'}(y_j)$ requires the evaluation of the matrix-coefficients appearing in (6.10). The coefficients $(J_R^{(1)})_{y_j}^{(1)}$ are particularly hard to compute and require more coefficients in the expansions of $P^{(\infty)}(z)$ as $z \rightarrow y_j$ than those already computed in Subsections 4.1, 4.2 and 4.3. However, it turns out that what will be needed in Section 8 is to obtain explicit expressions for

$$\left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21}. \quad (6.11)$$

The quantities (6.11) are much easier to evaluate than the $R^{(1)'}(y_j)$ themselves, due to heavy cancellations between $R^{(1)'}(y_j)$ and the factors $E_{y_j}(y_j)^{\pm 1}$. In particular, (6.11) can be explicitly written in terms of the first four coefficients appearing in the expansions of $P^{(\infty)}(z)$ as $z \rightarrow y_j$ (these coefficients were computed in Subsections 4.1, 4.2 and 4.3). Let us define

$$\mathcal{J}_{y_j}(z) = \left[E_{y_j}(y_j)^{-1} J_R^{(1)}(z) E_{y_j}(y_j) \right]_{21}.$$

As $z \rightarrow y_{j'}$, $j' = 1, 2, 3$, we have

$$\begin{aligned} \mathcal{J}_{y_j}(z) &= \sum_{k=-1}^1 (\mathcal{J}_{y_j})_{y_{j'}}^{(k)} (z - y_{j'})^k + \mathcal{O}((z - y_{j'})^2), \\ (\mathcal{J}_{y_j})_{y_{j'}}^{(k)} &= \left[E_{y_j}(y_j)^{-1} (J_R^{(1)})_{y_{j'}}^{(k)} E_{y_j}(y_j) \right]_{21}, \end{aligned} \quad (6.12)$$

so that (6.11) can be written as

$$\left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21} = -(\mathcal{J}_{y_j})_{y_j}^{(1)} - \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{(y_j - y_{j'})^2} (\mathcal{J}_{y_j})_{y_{j'}}^{(-1)}.$$

After a standard computation using the expansions derived in Subsections 4.1, 4.2 and 4.3, and using the expressions for $E_{y_j}(y_j)$ given in (5.8), (5.13) and (5.18), we obtain

$$\begin{aligned} (\mathcal{J}_{y_1})_{y_2}^{(-1)} &= -\frac{\sqrt{c_{y_1}} e^{2\pi i \nu} \pi (\beta_{y_2}^{(-\frac{1}{4})})^2 \mathcal{G}(\frac{\tau}{2})^2}{8\sqrt{c_{y_2}} (\beta_{y_1}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2} r^{\frac{3}{2}}, & (\mathcal{J}_{y_1})_{y_3}^{(-1)} &= -\frac{i\sqrt{c_{y_1}} e^{2\pi i \nu} \pi (\beta_{y_3}^{(\frac{1}{4})})^2 \tilde{\mathcal{G}}(\frac{1+\tau}{2})^2}{8\sqrt{c_{y_3}} (\varphi_{y_3}^{(\frac{1}{2})})^2 (\beta_{y_1}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2} r^{\frac{3}{2}}, \\ (\mathcal{J}_{y_2})_{y_1}^{(-1)} &= -\frac{\sqrt{c_{y_2}} e^{2\pi i \nu} \pi (\beta_{y_2}^{(-\frac{1}{4})})^2 \mathcal{G}(\frac{\tau}{2})^2}{8\sqrt{c_{y_1}} (\beta_{y_1}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2} r^{\frac{3}{2}}, & (\mathcal{J}_{y_3})_{y_1}^{(-1)} &= -\frac{i\sqrt{c_{y_3}} e^{2\pi i \nu} \pi (\beta_{y_3}^{(\frac{1}{4})})^2 \tilde{\mathcal{G}}(\frac{1+\tau}{2})^2}{8\sqrt{c_{y_1}} (\varphi_{y_3}^{(\frac{1}{2})})^2 (\beta_{y_1}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2} r^{\frac{3}{2}}, \\ (\mathcal{J}_{y_2})_{y_3}^{(-1)} &= -\frac{\sqrt{c_{y_2}} e^{4\pi i \nu} \pi \left(\varphi_{y_2}^{(\frac{1}{2})} \varphi_{y_3}^{(\frac{1}{2})} (\beta_{y_2}^{(-\frac{1}{4})})^2 \mathcal{G}(\frac{\tau}{2})^2 - (\beta_{y_3}^{(\frac{1}{4})})^2 \tilde{\mathcal{G}}(\frac{1+\tau}{2})^2 \right)^2}{8\sqrt{c_{y_3}} (\varphi_{y_2}^{(\frac{1}{2})})^2 (\varphi_{y_3}^{(\frac{1}{2})})^2 (\beta_{y_2}^{(-\frac{1}{4})})^2 (\beta_{y_3}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2 \mathcal{G}(\frac{1}{2})^2} r^{\frac{3}{2}}, \\ (\mathcal{J}_{y_3})_{y_2}^{(-1)} &= -\frac{\sqrt{c_{y_3}} e^{4\pi i \nu} \pi \left(\varphi_{y_2}^{(\frac{1}{2})} \varphi_{y_3}^{(\frac{1}{2})} (\beta_{y_2}^{(-\frac{1}{4})})^2 \mathcal{G}(\frac{\tau}{2})^2 - (\beta_{y_3}^{(\frac{1}{4})})^2 \tilde{\mathcal{G}}(\frac{1+\tau}{2})^2 \right)^2}{8\sqrt{c_{y_2}} (\varphi_{y_2}^{(\frac{1}{2})})^2 (\varphi_{y_3}^{(\frac{1}{2})})^2 (\beta_{y_2}^{(-\frac{1}{4})})^2 (\beta_{y_3}^{(\frac{1}{4})})^2 \mathcal{G}(0)^2 \mathcal{G}(\frac{1}{2})^2} r^{\frac{3}{2}}. \end{aligned}$$

The coefficient $(\mathcal{J}_{y_1})_{y_1}^{(1)}$ requires more involved computations and is given by

$$(\mathcal{J}_{y_1})_{y_1}^{(1)} = \frac{i\pi r^{\frac{3}{2}}}{16\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)^2} \left\{ 3c_{y_1}^{(2)}\beta_{y_1}^{(\frac{1}{4})}\mathcal{G}(0)^2 - 2\left(6\beta_{y_1}^{(\frac{5}{4})}\mathcal{G}(0)^2 + (\varphi_{y_1}^{(\frac{1}{2})})^2\beta_{y_1}^{(\frac{1}{4})}\left[\mathcal{G}'(0)^2 + 3\mathcal{G}(0)\mathcal{G}''(0)\right]\right) \right\}.$$

The expressions for $(\mathcal{J}_{y_2})_{y_2}^{(1)}$ and $(\mathcal{J}_{y_3})_{y_3}^{(1)}$ are more complicated and significantly longer than $(\mathcal{J}_{y_1})_{y_1}^{(1)}$, so we do not write them down. In Section 7, we derive several identities involving the θ -function which allows one to simplify the formulas for $(\mathcal{J}_{y_j})_{y_j}^{(k)}$, see proof of Proposition 8.5.

7 Some θ -function identities

The coefficients $(P^{(\infty)})_{y_j}^{(-\frac{1}{4}+\frac{k}{2})}$ obtained in Subsections 4.1, 4.2, 4.3, the expressions for $E_{y_j}(y_j)$ and $E'_{y_j}(y_j)$ given by (5.8), (5.13), (5.18), and the matrices $(\mathcal{J}_{y_j})_{y_j}^{(k)}$ computed in Section 6, all involve the θ -function and its derivatives evaluated at the 8 points

$$0, \quad \frac{1}{2}, \quad \frac{\tau}{2}, \quad \frac{1+\tau}{2}, \quad \nu, \quad \nu + \frac{1}{2}, \quad \nu + \frac{\tau}{2}, \quad \nu + \frac{1+\tau}{2}.$$

In this section, we provide a systematic way of simplifying these expressions by showing that

$$\theta^{(j)}\left(\frac{1}{2}\right), \quad \theta^{(j)}\left(\frac{\tau}{2}\right), \quad \theta^{(j)}\left(\frac{1+\tau}{2}\right), \quad j \geq 0, \quad (7.1)$$

where $\theta^{(j)}(z) := \frac{d^j}{dz^j}\theta(z)$, can be expressed in terms of $\theta(0)$, $\theta^{(2)}(0)$, $\theta^{(4)}(0)$, ..., and similarly that

$$\theta^{(j)}\left(\nu + \frac{1}{2}\right), \quad \theta^{(j)}\left(\nu + \frac{\tau}{2}\right), \quad \theta^{(j)}\left(\nu + \frac{1+\tau}{2}\right), \quad j \geq 0, \quad (7.2)$$

can be expressed in terms of $\theta(\nu)$, $\theta'(\nu)$, $\theta^{(2)}(\nu)$, ... We start by recalling the basic symmetries of the θ -function, which can be found in (1.9). By differentiating the relations $\theta(-u) = \theta(u)$ and $\theta(\frac{1}{2} - u) = \theta(\frac{1}{2} + u)$, we immediately obtain

$$\theta'(0) = 0, \quad \theta^{(3)}(0) = 0, \quad \theta'\left(\frac{1}{2}\right) = 0, \quad \theta^{(3)}\left(\frac{1}{2}\right) = 0.$$

We also recall that $\theta(z)$ has a simple zero at $z = \frac{1+\tau}{2}$. Let us define $\tilde{\theta}$ by

$$\tilde{\theta}(z) = \frac{\theta(z)}{z - \frac{1+\tau}{2}}, \quad (7.3)$$

so that $\tilde{\mathcal{G}}$ defined in (4.24) is given by $\tilde{\mathcal{G}}(z) = \frac{\theta(z+\nu)}{\theta(z)}$. By differentiating (7.3), we obtain

$$\tilde{\theta}^{(j-1)}\left(\frac{1+\tau}{2}\right) = \frac{1}{j}\theta^{(j)}\left(\frac{1+\tau}{2}\right), \quad j = 1, 2, \dots \quad (7.4)$$

Remark 6. It can be shown by standard arguments that $\det P^{(\infty)}(z) \equiv 1$. This gives a non-trivial relation between the θ -function evaluated at several points. However, it turns out that this is not enough for our needs. Proposition 7.1 below states three fundamental relations between the Abel map and the θ -function. These identities are of central importance for us and will be used extensively in the remainder of this paper.

Proposition 7.1. *Let $z \mapsto \varphi_A(z)$ denote the Abel map (4.10). Then, for all z on the Riemann surface,*

$$\frac{e^{2i\pi\varphi_A(z)}\theta(\varphi_A(z) + \frac{\tau}{2})^2}{\theta(\varphi_A(z))^2} = \frac{D_1^2}{z}, \quad (7.5a)$$

$$\frac{e^{2i\pi\varphi_A(z)}\theta(\varphi_A(z) + \frac{1+\tau}{2})^2}{\theta(\varphi_A(z))^2} = \frac{D_2^2(z - y_1)}{z}, \quad (7.5b)$$

$$\frac{\theta(\varphi_A(z) + \frac{1}{2})^2}{\theta(\varphi_A(z))^2} = \frac{D_3^2(z - y_2)}{z}, \quad (7.5c)$$

where we have identified $z \in X$ with its projection onto the complex plane on the right-hand sides, and

$$D_1 = (y_1 y_2)^{\frac{1}{4}} e^{-\frac{\pi i \tau}{4}}, \quad D_2 = i \frac{y_2^{\frac{1}{4}} e^{-\frac{\pi i \tau}{4}}}{(y_1 - y_2)^{\frac{1}{4}}}, \quad D_3 = \frac{y_1^{\frac{1}{4}}}{(y_1 - y_2)^{\frac{1}{4}}}. \quad (7.6)$$

Proof. An easy computation using the transformation properties (1.9) of θ show that the three quotients

$$\frac{e^{2\pi i u} \theta(u + \frac{\tau}{2})^2}{\theta(u)^2}, \quad \frac{e^{2i\pi u} \theta(u + \frac{1+\tau}{2})^2}{\theta(u)^2}, \quad \frac{\theta(u + \frac{1}{2})^2}{\theta(u)^2}$$

are invariant under the shifts $u \mapsto u + 1$ and $u \mapsto u + \tau$, so that they are well-defined on the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$. Let us first prove (7.5a). Since $\theta(u)$ has a simple zero at $u = \frac{1+\tau}{2}$ and no other zeros, it follows that the quotient $\frac{e^{2\pi i u} \theta(u + \frac{\tau}{2})^2}{\theta(u)^2}$ defines a meromorphic function on $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ with a double zero at $u = \frac{1}{2}$ and a double pole at $u = \frac{1+\tau}{2}$ and no other zeros or poles. Recalling that

$$\varphi_A(y_1) = 0, \quad \varphi_A(y_2) = \frac{\tau}{2}, \quad \varphi_A(y_3) = \frac{1}{2} + \frac{\tau}{2}, \quad \varphi_A(\infty) = \frac{1}{2},$$

this implies that the left-hand side of (7.5a) is a meromorphic function on the Riemann surface with a double zero at $z = \infty$ and a double pole at $z = y_3$. Since the torus is of genus 1, it follows from the Riemann-Roch theorem that the identity (7.5a) holds for some constant $D_1^2 \in \mathbb{C}$. Analogous arguments apply to (7.5b) and (7.5c). By evaluating the identities in (7.5) at y_1, ∞ and ∞ , respectively, we obtain

$$D_1 = \sqrt{y_1} \frac{\theta(\frac{\tau}{2})}{\theta(0)} \in \mathbb{R}, \quad D_2 = i \frac{\theta(\frac{\tau}{2})}{\theta(\frac{1}{2})} \in i\mathbb{R}, \quad D_3 = \frac{\theta(0)}{\theta(\frac{1}{2})} \in \mathbb{R}. \quad (7.7)$$

On the other hand, by expanding (7.5a) as $z \rightarrow y_2$ and (7.5b), (7.5c) as $z \rightarrow y_1$, we obtain

$$D_1 = \sqrt{y_2} \frac{e^{-\frac{\pi i \tau}{2}} \theta(0)}{\theta(\frac{\tau}{2})}, \quad D_2 = \frac{\sqrt{y_1} \varphi_{y_1}(\frac{1}{2}) \theta'(\frac{\tau+1}{2})}{\theta(0)}, \quad D_3 = \frac{\sqrt{y_1} \theta(\frac{1}{2})}{\sqrt{y_1 - y_2} \theta(0)}. \quad (7.8)$$

Comparing (7.7) with (7.8), we obtain (7.6). \square

We immediately obtain the following Corollary.

Corollary 7.2. *Let $z \mapsto \varphi_A(z)$ be the Abel map defined in (4.10). Then*

$$\frac{e^{2i\pi\varphi_A(z)} \theta(\varphi_A(z) + \frac{\tau}{2}) \theta(\varphi_A(z) + \frac{1+\tau}{2})}{\theta(\varphi_A(z)) \theta(\varphi_A(z) + \frac{1}{2})} = \frac{D_1 D_2}{D_3} \sqrt{\frac{z - y_1}{(z - y_2)z}} \quad (7.9)$$

for all z on the Riemann surface, where the branch is such that $\sqrt{\frac{z - y_1}{(z - y_2)z}} > 0$ for $z > y_1$ on the upper sheet, and

$$\frac{D_1 D_2}{D_3} = i \sqrt{y_2} e^{-\frac{\pi i \tau}{2}}. \quad (7.10)$$

Proof. By multiplying (7.5a) with (7.5b), and then dividing by (7.5c), we obtain

$$\left(\frac{e^{2i\pi\varphi_A(z)} \theta(\varphi_A(z) + \frac{\tau}{2}) \theta(\varphi_A(z) + \frac{1+\tau}{2})}{\theta(\varphi_A(z)) \theta(\varphi_A(z) + \frac{1}{2})} \right)^2 = \frac{D_1^2 D_2^2}{D_3^2} \frac{z - y_1}{(z - y_2)z}. \quad (7.11)$$

By taking the square root of (7.11), we obtain (7.9), up to a multiplicative sign that can be determined from an expansion as $z \rightarrow y_1$, see [29, Chapter 20]. \square

By taking derivatives of (7.5a)-(7.5c), we can obtain expressions for (7.1) in terms of only $\theta(0), \theta''(0), \dots$. We summarize the relations that will be used in Section 8 in the following Corollary.

Corollary 7.3. *We have the relations*

$$\theta\left(\frac{\tau}{2}\right) = e^{-\frac{\pi i \tau}{4}} \frac{y_2^{\frac{1}{4}}}{y_1^{\frac{1}{4}}} \theta(0), \quad \theta\left(\frac{1}{2}\right) = \frac{(y_1 - y_2)^{\frac{1}{4}}}{y_1^{\frac{1}{4}}} \theta(0), \quad \theta'\left(\frac{\tau}{2}\right) = -i\pi e^{-\frac{\pi i \tau}{4}} \frac{y_2^{\frac{1}{4}}}{y_1^{\frac{1}{4}}} \theta(0),$$

as well as

$$\begin{aligned} \theta'\left(\frac{1+\tau}{2}\right) &= ie^{-\frac{\pi i \tau}{4}} \frac{(y_1 - y_2)^{\frac{1}{4}} y_2^{\frac{1}{4}}}{2c_0} \theta(0), \\ \theta''\left(\frac{\tau}{2}\right) &= e^{-\frac{\pi i \tau}{4}} \frac{y_2^{\frac{1}{4}}}{y_1^{\frac{1}{4}}} \left(\theta''(0) - \left[\pi^2 + \frac{y_1 - y_2}{4c_0^2} \right] \theta(0) \right), \\ \theta''\left(\frac{1+\tau}{2}\right) &= \pi e^{-\frac{\pi i \tau}{4}} \frac{(y_1 - y_2)^{\frac{1}{4}} y_2^{\frac{1}{4}}}{c_0} \theta(0), \\ \theta''\left(\frac{1}{2}\right) &= \frac{(y_1 - y_2)^{\frac{1}{4}}}{4c_0^2 y_1^{\frac{1}{4}}} \left(y_2 \theta(0) + 4c_0^2 \theta''(0) \right), \\ \theta^{(3)}\left(\frac{1+\tau}{2}\right) &= -ie^{-\frac{\pi i \tau}{4}} \frac{(y_1 - y_2)^{\frac{1}{4}} y_2^{\frac{1}{4}}}{8c_0^3} \left((y_1 - 2y_2) \theta(0) + 12c_0^2 (\pi^2 \theta(0) - \theta''(0)) \right). \end{aligned}$$

Finally, we note that the relations (7.5a)-(7.5c) allows also to express (7.2) in terms of $\theta(\nu)$, $\theta'(\nu)$, $\theta^{(2)}(\nu), \dots$, via the following identities

$$\theta\left(\nu + \frac{\tau}{2}\right)^2 = e^{-2i\pi\nu} \frac{D_1^2}{a} \theta(\nu)^2, \quad (7.12a)$$

$$\theta\left(\nu + \frac{1+\tau}{2}\right)^2 = e^{-2i\pi\nu} \frac{D_2^2(a - y_1)}{a} \theta(\nu)^2, \quad (7.12b)$$

$$\theta\left(\nu + \frac{1}{2}\right)^2 = \frac{D_3^2(a - y_2)}{a} \theta(\nu)^2, \quad (7.12c)$$

where $a = \varphi_A^{-1}(\nu)$.

8 Proof of Theorem 1.1

In Section 2 we obtained the following differential identity:

$$\partial_r \log F(r\vec{x}) = \sum_{j=1}^3 K_j, \quad \text{with} \quad K_j := \frac{(-1)^{j+1} x_j}{2\pi i} \lim_{z \rightarrow y_j} (\Psi_+^{-1} \Psi'_+)^{-1}_{21}(rz; rx_3, r\vec{y}), \quad (8.1)$$

where the limits as $z \rightarrow y_j$, $j = 1, 2, 3$, are taken such that $z \in (0, y_2) \cup (y_1, +\infty)$. In this section, we use the analysis from Sections 3-6 to obtain large r asymptotics for K_j , $j = 1, 2, 3$. These asymptotics can be simplified in two steps: first, we use the numerous identities involving the θ -function and the Abel map of Section 7, and second, we use Riemann's bilinear identity and some other identities for elliptic integrals. Then, we integrate these asymptotics as shown in (1.15) and prove the asymptotic formula (1.10) for $\log F(r\vec{x})$.

Proposition 8.1. *Let $\vec{x} = (x_1, x_2, x_3)$ be fixed and such that $x_3 < x_2 < x_1 < 0$. We have*

$$\begin{aligned} \partial_r \log F(r\vec{x}) &= r^2 \sum_{j=1}^3 (-1)^{j+1} x_j c_{y_j} + \frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21} \\ &\quad + \frac{1}{2\pi i r} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} E'_{y_j}(y_j) \right]_{21} + \mathcal{O}(r^{-\frac{5}{2}}) \end{aligned} \quad (8.2)$$

as $r \rightarrow +\infty$, where c_{y_j} , $j = 1, 2, 3$, are given by (5.3), (5.10) and (5.15).

Proof. Using the transformations $\Psi \rightarrow T \rightarrow S \rightarrow R$ given by (3.17), (3.20), (6.1), we see that, for z inside \mathbb{D}_j , $j = 1, 2, 3$, but outside the lenses, we have

$$\Psi(rz) = \begin{pmatrix} r^{\frac{1}{4}} & -2ir^{\frac{7}{4}}g_1 \\ 0 & r^{-\frac{1}{4}} \end{pmatrix} R(z)P^{(y_j)}(z)e^{r^{\frac{3}{2}}g(z)\sigma_3}.$$

Thus, for z inside \mathbb{D}_j but outside the lenses, we obtain

$$\begin{aligned} (\Psi^{-1}\Psi')(rz) &= \frac{1}{r}e^{-r^{\frac{3}{2}}g(z)\sigma_3} \left\{ r^{\frac{3}{2}}g'(z)\sigma_3 + P^{(y_j)}(z)^{-1}P^{(y_j)'}(z) \right. \\ &\quad \left. + P^{(y_j)}(z)^{-1}R(z)^{-1}R'(z)P^{(y_j)}(z) \right\} e^{r^{\frac{3}{2}}g(z)\sigma_3}. \end{aligned}$$

Consequently, for $j = 1, 2, 3$, we have

$$K_j = \frac{(-1)^{j+1}x_j}{2\pi ir} \lim_{z \rightarrow y_j} e^{2r^{\frac{3}{2}}g_+(z)} \left[P_+^{(y_j)}(z)^{-1}P_+^{(y_j)'}(z) + P_+^{(y_j)}(z)^{-1}R(z)^{-1}R'(z)P_+^{(y_j)}(z) \right]_{21}, \quad (8.3)$$

where again the limits are taken such that $z \in (0, y_2) \cup (y_1, +\infty)$, and we write $R(z)$ instead of $R_+(z)$ because $R(z)$ has no jumps on $(0, y_2) \cup (y_1, +\infty)$. The expressions (5.5), (5.11), (5.16) for $P^{(y_j)}(z)$, $j = 1, 2, 3$, are of the form

$$\begin{aligned} P^{(y_j)}(z) &= E_{y_j}(z)\Phi_{\text{Be}}(r^3f_{y_j}(z))e^{-r^{\frac{3}{2}}g(z)\sigma_3}, \quad j = 1, 3 \\ P^{(y_2)}(z) &= E_{y_2}(z)\sigma_3\Phi_{\text{Be}}(-r^3f_{y_2}(z))\sigma_3e^{-r^{\frac{3}{2}}g(z)\sigma_3}, \end{aligned}$$

so we can write (8.3) as

$$\begin{aligned} K_j &= \frac{x_j}{2\pi ir} \lim_{z \downarrow y_j} \left[\Phi_{\text{Be}}(r^3f_{y_j}(z))^{-1}E_{y_j}(z)^{-1}E'_{y_j}(z)\Phi_{\text{Be}}(r^3f_{y_j}(z)) \right. \\ &\quad \left. + \Phi_{\text{Be}}(r^3f_{y_j}(z))^{-1}\Phi'_{\text{Be}}(r^3f_{y_j}(z))r^3f'_{y_j}(z) - r^{\frac{3}{2}}g'(z)\sigma_3 \right. \\ &\quad \left. + \Phi_{\text{Be}}(r^3f_{y_j}(z))^{-1}E_{y_j}(z)^{-1}R(z)^{-1}R'(z)E_{y_j}(z)\Phi_{\text{Be}}(r^3f_{y_j}(z)) \right]_{21}, \quad j = 1, 3, \\ K_2 &= \frac{-x_2}{2\pi ir} \lim_{z \uparrow y_2} \left[\sigma_3\Phi_{\text{Be}}(-r^3f_{y_2}(z))^{-1}\sigma_3E_{y_2}(z)^{-1}E'_{y_2}(z)\sigma_3\Phi_{\text{Be}}(-r^3f_{y_2}(z))\sigma_3 \right. \\ &\quad \left. + \sigma_3\Phi_{\text{Be}}(-r^3f_{y_2}(z))^{-1}\Phi'_{\text{Be}}(-r^3f_{y_2}(z))\sigma_3(-r^3f'_{y_2}(z)) - r^{\frac{3}{2}}g'(z)\sigma_3 \right. \\ &\quad \left. + \sigma_3\Phi_{\text{Be}}(-r^3f_{y_2}(z))^{-1}\sigma_3E_{y_2}(z)^{-1}R(z)^{-1}R'(z)E_{y_2}(z)\sigma_3\Phi_{\text{Be}}(r^3f_{y_2}(z))\sigma_3 \right]_{21}. \end{aligned}$$

Note that the term $-r^{\frac{3}{2}}g'(z)\sigma_3$ does not contribute for K_j , since $(\sigma_3)_{21} = 0$. As $z \rightarrow y_j$ with z outside the lenses, we have $(-1)^{j+1}f_{y_j}(z) \rightarrow 0$ with $|\arg[(-1)^{j+1}f_{y_j}(z)]| < \frac{2\pi}{3}$ for each $j = 1, 2, 3$. Thus, using (A.7), we have

$$\lim_{z \rightarrow y_j} \left[\Phi_{\text{Be}}((-1)^{j+1}r^3f_{y_j}(z))^{-1}\Phi'_{\text{Be}}((-1)^{j+1}r^3f_{y_j}(z)) \right]_{21} = 2\pi i, \quad j = 1, 2, 3.$$

Moreover, $E_{y_j}(z)$ and $R(z)$ are analytic for $z \in \mathbb{D}_{y_j}$. Hence, using (A.5)-(A.7) we have

$$K_j = \frac{(-1)^{j+1}x_j}{2\pi ir} \left\{ \left[E_{y_j}(y_j)^{-1}E'_{y_j}(y_j) \right]_{21} + 2\pi ir^3f'_{y_j}(y_j) + \left[E_{y_j}(y_j)^{-1}R(y_j)^{-1}R'(y_j)E_{y_j}(y_j) \right]_{21} \right\}$$

for $j = 1, 2, 3$. We also note that $f'_{y_j}(y_j) = c_{y_j}$ for $j = 1, 2, 3$. Furthermore, by (6.5) and (6.6), we have

$$R^{-1}(z)R'(z) = \left(I + \mathcal{O}(r^{-\frac{3}{2}}) \right) \left(\frac{R^{(1)'}(z)}{r^{\frac{3}{2}}} + \mathcal{O}(r^{-3}) \right), \quad \text{as } r \rightarrow +\infty,$$

and by (5.8), (5.13) and (5.18), we have

$$\begin{aligned} E_{y_j}(y_j) &= \begin{pmatrix} \mathcal{O}(r^{\frac{3}{4}}) & \mathcal{O}(r^{-\frac{3}{4}}) \\ \mathcal{O}(r^{\frac{3}{4}}) & \mathcal{O}(r^{-\frac{3}{4}}) \end{pmatrix}, \quad r \rightarrow +\infty, \quad j = 1, 2, 3, \\ E_{y_j}(y_j)^{-1} &= \begin{pmatrix} \mathcal{O}(r^{-\frac{3}{4}}) & \mathcal{O}(r^{-\frac{3}{4}}) \\ \mathcal{O}(r^{\frac{3}{4}}) & \mathcal{O}(r^{\frac{3}{4}}) \end{pmatrix}, \quad r \rightarrow +\infty, \quad j = 1, 2, 3. \end{aligned}$$

We infer that, for $j = 1, 2, 3$,

$$K_j = \frac{(-1)^{j+1}x_j}{2\pi i r} \left\{ 2\pi i c_{y_j} r^3 + \left[E_{y_j}(y_j)^{-1} E'_{y_j}(y_j) \right]_{21} + r^{-\frac{3}{2}} \left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21} + \mathcal{O}(r^{-\frac{3}{2}}) \right\}.$$

The proposition follows by summing from $j = 1$ to $j = 3$. \square

8.1 Evaluation of the first term

Proposition 8.2. *The first term on the right-hand side of (8.2) is given by*

$$r^2 \sum_{j=1}^3 (-1)^{j+1} x_j c_{y_j} = c \partial_r r^3, \quad (8.4)$$

where c is defined in (1.11).

Proof. By direct computation using the explicit expressions

$$c_{y_1} = \frac{p(y_1)^2}{y_1(y_1 - y_2)}, \quad c_{y_2} = \frac{p(y_2)^2}{y_2(y_1 - y_2)}, \quad c_{y_3} = \frac{p(y_3)^2}{y_1 y_2},$$

and the definition of $p(z)$ given by (3.3), we obtain

$$\begin{aligned} r^2 \sum_{j=1}^3 (-1)^{j+1} x_j c_{y_j} &= \frac{1}{3} \left(\frac{x_1 p(y_1)^2}{y_1(y_1 - y_2)} - \frac{x_2 p(y_2)^2}{y_2(y_1 - y_2)} + \frac{x_3 p(y_3)^2}{y_1 y_2} \right) \partial_r r^3 \\ &= \frac{1}{24} \left((x_1 - x_2)^2 (x_1 + x_2 - x_3) + 2x_3^3 + 8g_1(x_1 + x_2 + x_3) \right) \partial_r r^3. \end{aligned} \quad (8.5)$$

We note that \mathcal{Q}, q defined in (1.4), (1.5) have the relations with \mathcal{R}, p

$$\mathcal{Q}(z) = \mathcal{R}(z - x_3) \quad \text{and} \quad q(z) = p(z - x_3).$$

Therefore, g_1 given by (3.4) and q_0 given by (1.6) are related by

$$g_1 = q_0 - \frac{2x_1 x_2 + 2x_1 x_3 + 2x_2 x_3 - x_1^2 - x_2^2}{8}. \quad (8.6)$$

We obtain (8.4) after substituting (8.6) into (8.5). \square

8.2 Evaluation of the second term

Proposition 8.3. *The second term on the right-hand side of (8.2) is given by*

$$\begin{aligned} &\frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21} \\ &= \frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^j x_j (\mathcal{J}_{y_j})_{y_j}^{(1)} + \frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^j x_j \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{(y_j - y_{j'})^2} (\mathcal{J}_{y_j})_{y_{j'}}^{(-1)}. \end{aligned} \quad (8.7)$$

Proof. The claim follows after substituting the expression (6.10) for $R^{(1)'}(y_j)$ into the left-hand side of (8.7) and by using the definition (6.12) of $(\mathcal{J}_{y_j})_{y_{j'}}^{(k)}$. \square

Proposition 8.4. *The first term on the right-hand side of (8.7) is given by*

$$\frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^j x_j (\mathcal{J}_{y_j})_{y_j}^{(1)} = -\frac{1}{16} \sum_{j=1}^3 \frac{x_j p'(y_j)}{p(y_j)} \partial_r \log r. \quad (8.8)$$

Proof. This follows from involved computations which are quite long and thus will be omitted here. We use the identities of Corollary 7.3 to express $\theta^{(j)}(\frac{1}{2})$, $\theta^{(j)}(\frac{\tau}{2})$ and $\theta^{(j)}(\frac{1+\tau}{2})$, $j = 0, 1, 2, 3$, in terms of only $\theta^{(j)}(0)$, $j = 0, 2$. We also use the j -th derivative with respect to ν of (7.12a)-(7.12c), $j = 0, 1, 2, 3$, to express $\theta^{(j)}(\nu + \frac{1}{2})$, $\theta^{(j)}(\nu + \frac{\tau}{2})$ and $\theta^{(j)}(\nu + \frac{1+\tau}{2})$, $j = 0, 1, 2, 3$, in terms of only $\theta^{(j)}(\nu)$, $j = 0, 1, 2, 3$. Remarkably, the final expression on the right-hand side of (8.8) does not contain the θ -function. \square

Definition 1. Let $\varphi_{\mathbb{C}}^{-1}(u)$ be the projection into the complex plane of $\varphi_A^{-1}(u)$. We define $a = a(r)$ by

$$a := \varphi_{\mathbb{C}}^{-1}(\nu) = \varphi_{\mathbb{C}}^{-1}\left(-\frac{\Omega}{2\pi}r^{\frac{3}{2}}\right). \quad (8.9)$$

From the relations $\varphi_{\mathbb{C}}^{-1}(u) = \varphi_{\mathbb{C}}^{-1}(-u)$, $\varphi_{\mathbb{C}}^{-1}(u+1) = \varphi_{\mathbb{C}}^{-1}(u)$ and $\Omega > 0$, we conclude that the function $r \mapsto a(r)$ is oscillatory as $r \rightarrow +\infty$. Furthermore, we see that $\varphi_A(x)$ is monotone for $x \in (y_1, +\infty)$, which follows from direct inspection of (4.5). Since $\varphi_A(y_1) = 0$ and $\varphi_A(\infty) = \frac{1}{2}$, we conclude that $a(r)$ is bigger than y_1 for any r and oscillates between y_1 and $+\infty$.

Proposition 8.5. *The second term on the right-hand side of (8.7) is given by*

$$\frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^j x_j \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{(y_j - y_{j'})^2} (\mathcal{J}_{y_j})_{y_{j'}}^{(-1)} = n^{(0)} \partial_r \log r + n^{(-1)} \partial_r \int_M^r \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})}, \quad (8.10)$$

where $a = a(r)$ is oscillatory and given by (8.9), $M > 0$ is a large constant, and $n^{(-1)}, n^{(0)}$ are given by

$$n^{(-1)} = -\frac{1}{32} \left(\frac{1}{p(y_1)} + \frac{1}{p(y_2)} + \frac{1}{p(y_3)} \right) (x_1 - x_3)(x_2 - x_3)(x_1 + x_2 + x_3), \quad (8.11)$$

$$n^{(0)} = -\frac{x_1 \frac{y_2}{y_1} p(y_1) - x_2 \frac{y_1}{y_2} p(y_2)}{16 p(y_3)(x_1 - x_2)} - \frac{x_3 p(y_3)}{16} \left(\frac{1}{y_1 p(y_1)} + \frac{1}{y_2 p(y_2)} \right). \quad (8.12)$$

Proof. By using Proposition 3.1, Corollary 7.3 together with the derivatives with respect to ν of (7.12a)-(7.12c), we can simplify the quantities obtained at the end of Section 6 for $(\mathcal{J}_{y_j})_{y_{j'}}^{(-1)}$, $1 \leq j \neq j' \leq 3$, as follows

$$\begin{aligned} (\mathcal{J}_{y_1})_{y_2}^{(-1)} &= -\frac{\pi i}{8} \frac{(y_1 - y_2) y_2 p(y_1)}{p(y_2)} \frac{r^{\frac{3}{2}}}{a}, & (\mathcal{J}_{y_2})_{y_1}^{(-1)} &= -\frac{\pi i}{8} \frac{(y_1 - y_2) y_1 p(y_2)}{p(y_1)} \frac{r^{\frac{3}{2}}}{a}, \\ (\mathcal{J}_{y_1})_{y_3}^{(-1)} &= \frac{\pi i}{8} \frac{y_1 y_2 p(y_1)}{(y_1 - y_2) p(y_3)} \frac{r^{\frac{3}{2}}(a - y_1)}{a}, & (\mathcal{J}_{y_3})_{y_1}^{(-1)} &= \frac{\pi i}{8} \frac{y_1 p(y_3)}{p(y_1)} \frac{r^{\frac{3}{2}}(a - y_1)}{a}, \\ (\mathcal{J}_{y_2})_{y_3}^{(-1)} &= \frac{\pi i}{8} \frac{y_1 y_2 p(y_2)}{(y_1 - y_2) p(y_3)} \frac{r^{\frac{3}{2}}(a - y_2)}{a}, & (\mathcal{J}_{y_3})_{y_2}^{(-1)} &= \frac{\pi i}{8} \frac{y_2 p(y_3)}{p(y_2)} \frac{r^{\frac{3}{2}}(a - y_2)}{a}, \end{aligned}$$

where $a = a(r)$ depends on r and is given by (8.9). Then, a long but straightforward computation gives

$$\frac{1}{2\pi i r^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^j x_j \sum_{\substack{j'=1 \\ j' \neq j}}^3 \frac{1}{(y_j - y_{j'})^2} (\mathcal{J}_{y_j})_{y_{j'}}^{(-1)} = n^{(0)} \partial_r \log r + n^{(-1)} \partial_r \int_M^r \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})},$$

where $n^{(0)}$ is given by (8.12) and where $n^{(-1)}$ is given by

$$n^{(-1)} = \frac{1}{16} \left(\frac{1}{p(y_1)} + \frac{1}{p(y_2)} + \frac{1}{p(y_3)} \right) \left(x_1 p(y_1) \frac{x_2 - x_3}{x_1 - x_2} - x_2 p(y_2) \frac{x_1 - x_3}{x_1 - x_2} + x_3 p(y_3) \right).$$

Substituting the explicit expression (3.3) for p leads to further simplifications and we find (8.11). \square

The next proposition shows that $\int_M^r \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})}$ is proportional to $\log r$ and contains no oscillations of order 1.

Proposition 8.6. *As $r \rightarrow +\infty$, we have*

$$\int_M^r \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} = 2 \int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx \log r + \tilde{C} + \mathcal{O}(r^{-1}), \quad (8.13)$$

where \tilde{C} is a constant independent of r .

Proof. Let $r_0 > M$ be a large constant smaller than r and such that $a(r_0) = y_1$. Since $\Omega > 0$, the values of $r \geq r_0$ for which $a(r) = +\infty$ or $a(r) = y_1$ are given by

$$r_j := \left(r_0^{\frac{3}{2}} + j \frac{\pi}{\Omega} \right)^{\frac{2}{3}}, \quad j \geq 0.$$

Let us write

$$r = \left(r_0^{\frac{3}{2}} + 2k \frac{\pi}{\Omega} + \lambda \frac{\pi}{\Omega} \right)^{\frac{2}{3}}, \quad \text{where } k \in \mathbb{N} \quad \text{and} \quad 0 \leq \lambda < 2,$$

so that

$$\int_M^r \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} = \int_M^{r_0} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} + \sum_{j=0}^{k-1} \left[\int_{r_{2j}}^{r_{2j+1}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} + \int_{r_{2j+1}}^{r_{2j+2}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} \right] + \int_{r_{2k}}^{r_{2k} + \lambda \frac{\pi}{\Omega}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})}.$$

From the change of variables $\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}$ and using the notation $\nu_j = -\frac{\Omega}{2\pi} r_j^{\frac{3}{2}}$, we obtain

$$\begin{aligned} \int_{r_{2j}}^{r_{2j+1}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} &= \frac{2}{3} \int_{\nu_{2j}}^{\nu_{2j+1}} \frac{d\nu}{\nu \varphi_{\mathbb{C}}^{-1}(\nu)} = \frac{2}{3} \int_{\nu_{2j}}^{\nu_{2j+1}} \frac{1}{\nu_{2j}} \left(1 + \mathcal{O} \left(\frac{\nu - \nu_{2j}}{\nu_{2j}} \right) \right) \frac{d\nu}{\varphi_{\mathbb{C}}^{-1}(\nu)} \\ &= -\frac{4\pi}{3\Omega} r_{2j}^{-\frac{3}{2}} \left(\int_{\nu_{2j}}^{\nu_{2j+1}} \frac{d\nu}{\varphi_{\mathbb{C}}^{-1}(\nu)} + \mathcal{O} \left(\frac{\nu_{2j+1} - \nu_{2j}}{\nu_{2j}} \right) \right) = -\frac{4\pi}{3\Omega} r_{2j}^{-\frac{3}{2}} \left(\int_{\nu_{2j}}^{\nu_{2j+1}} \frac{d\nu}{\varphi_{\mathbb{C}}^{-1}(\nu)} + \mathcal{O}(r_{2j}^{-1}) \right) \end{aligned}$$

as $r_{2j} \rightarrow +\infty$. Noting that $\nu_{2j+1} = \nu_{2j} - \frac{1}{2}$, $\varphi_{\mathbb{C}}^{-1}(\nu_{2j+1}) = +\infty$, and $\varphi_{\mathbb{C}}^{-1}(\nu)$ is strictly increasing for ν in the oriented segment (ν_{2j}, ν_{2j+1}) , we use the change of variables $x = \varphi_{\mathbb{C}}^{-1}(\nu)$ to obtain

$$\int_{r_{2j}}^{r_{2j+1}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} = \frac{4\pi}{3\Omega} r_{2j}^{-\frac{3}{2}} \left(\int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx + \mathcal{O}(r_{2j}^{-1}) \right), \quad j = 0, \dots, k-1.$$

Similarly,

$$\int_{r_{2j+1}}^{r_{2j+2}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} = \frac{4\pi}{3\Omega} r_{2j+1}^{-\frac{3}{2}} \left(\int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx + \mathcal{O}(r_{2j+1}^{-1}) \right), \quad j = 0, \dots, k-1.$$

As $r \rightarrow +\infty$,

$$\sum_{j=0}^{2k-1} r_j^{-\frac{3}{2}} = \frac{\Omega}{\pi} \log k + d_0 + \mathcal{O}(k^{-1}) = \frac{3\Omega}{2\pi} \log r + d_1 + \mathcal{O}(r^{-\frac{3}{2}}), \quad \sum_{j=0}^{2k-1} r_j^{-\frac{5}{2}} = d_2 + \mathcal{O}(r^{-1}),$$

for certain constants d_0, d_1, d_2 , so we have

$$\sum_{j=0}^{k-1} \left[\int_{r_{2j}}^{r_{2j+1}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} + \int_{r_{2j+1}}^{r_{2j+2}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} \right] = 2 \int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx + \mathcal{O}(r^{-1})$$

as $r \rightarrow +\infty$. Since $a(\tilde{r}) \geq y_1$ for all \tilde{r} ,

$$\int_M^{r_{2k} + \lambda \frac{\pi}{\Omega}} \frac{d\tilde{r}}{\tilde{r} a(\tilde{r})} = \mathcal{O}(r^{-1}), \quad \text{as } r \rightarrow +\infty,$$

and the proof is completed. \square

In Propositions 8.4 and 8.5, we were able to express the right-hand side of (8.7) as a perfect derivative and in Proposition 8.6 we obtained the leading order r behavior of the only non-explicit term appearing on the right-hand side of (8.7). Thus, by combining Propositions 8.3, 8.4, 8.5, 8.6, we have, as $r \rightarrow +\infty$,

$$\int_M^r \frac{1}{2\pi i \tilde{r}^{\frac{5}{2}}} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} R^{(1)'}(y_j) E_{y_j}(y_j) \right]_{21} d\tilde{r} = c_2 \log r + \tilde{C}_2 + \mathcal{O}(r^{-1}), \quad (8.14)$$

where M and \tilde{C}_2 are independent of r , and c_2 is given by

$$c_2 = -\frac{1}{16} \sum_{j=1}^3 \frac{x_j p'(y_j)}{p(y_j)} + n^{(0)} + 2n^{(-1)} \int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx. \quad (8.15)$$

This expression for c_2 could be obtained thanks to the considerable simplifications of Propositions 8.4-8.6, and does not involve the θ -function. Surprisingly, it is possible to simplify c_2 significantly more.

Proposition 8.7. *We have $c_2 = -\frac{1}{2}$.*

Proof. Using [23, eqs 3.131.4, 3.134.28], we obtain

$$\int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx = \frac{1}{2y_2} \left(1 - \frac{\mathbf{E}(k_*)}{\mathbf{K}(k_*)} \right), \quad (8.16)$$

where $k_* = \sqrt{\frac{y_2}{y_1}}$. Recall from the proof of Proposition 3.1 that $p(z)$ can be written

$$p(z) = -z^2 + \frac{z}{2}(y_1 + y_2 - x_3) + \alpha, \quad \alpha = \frac{y_1}{2} \left[\frac{y_1 - y_2}{3} + x_3 - \frac{\mathbf{E}(k_*)}{\mathbf{K}(k_*)} \left(\frac{y_1 + y_2}{3} + x_3 \right) \right]. \quad (8.17)$$

Since $y_3 = 0$, and hence $p(y_3) \equiv p(0) = \alpha$, we have the relation

$$1 - \frac{\mathbf{E}(k_*)}{\mathbf{K}(k_*)} = \frac{6p(y_3) + 2y_1 y_2}{y_1(x_1 + x_2 + x_3)}$$

which together with (8.16) and the definition of $n^{(-1)}$ (8.11) implies that

$$2n^{(-1)} \int_{y_1}^{+\infty} \frac{\varphi'(x)}{x} dx = -\frac{1}{16} \left(\frac{1}{p(y_1)} + \frac{1}{p(y_2)} + \frac{1}{p(y_3)} \right) (3p(y_3) + y_1 y_2). \quad (8.18)$$

Using (8.17), we find the following identities:

$$\begin{aligned} -\frac{x_3 p(y_3)}{16y_1 p(y_1)} - \frac{x_1 p'(y_1)}{16p(y_1)} &= \frac{\frac{x_1}{2}(3y_1 - y_2 + x_3) - \frac{x_3}{y_1} \alpha}{16p(y_1)}, \\ -\frac{x_2 p'(y_2)}{16p(y_2)} + \frac{x_3 p(y_3)}{16y_2 p(y_2)} &= \frac{\frac{x_2}{2}(-y_1 + 3y_2 + x_3) - \frac{x_3}{y_2} \alpha}{16p(y_2)}, \\ -\frac{x_1 \frac{y_2}{y_1} p(y_1) - x_2 \frac{y_1}{y_2} p(y_2)}{16p(y_3)(x_1 - x_2)} - \frac{x_3 p'(y_3)}{16p(y_3)} &= \frac{y_1 y_2}{16p(y_3)} - \frac{1}{16} \left(1 - \frac{x_1}{y_1} - \frac{x_2}{y_2} \right). \end{aligned}$$

Substituting the above identities and (8.18) into (8.15), we obtain

$$\begin{aligned} c_2 = -\frac{1}{16} \left(4 - \frac{x_1}{y_1} - \frac{x_2}{y_2} \right) &+ \frac{\frac{x_1}{2}(3y_1 - y_2 + x_3) - y_1 y_2 - \alpha(3 + \frac{x_3}{y_1})}{16p(y_1)} + \\ &+ \frac{\frac{x_2}{2}(-y_1 + 3y_2 + x_3) - y_1 y_2 - \alpha(3 + \frac{x_3}{y_2})}{16p(y_2)}. \end{aligned} \quad (8.19)$$

The two numerators in (8.19) can be simplified as follows:

$$\begin{aligned} \frac{x_1}{2}(3y_1 - y_2 + x_3) - y_1 y_2 - \alpha \left(3 + \frac{x_3}{y_1} \right) &= -p(y_1) \left(3 + \frac{x_3}{y_1} \right), \\ \frac{x_2}{2}(-y_1 + 3y_2 + x_3) - y_1 y_2 - \alpha \left(3 + \frac{x_3}{y_2} \right) &= -p(y_2) \left(3 + \frac{x_3}{y_2} \right), \end{aligned}$$

and hence

$$c_2 = -\frac{1}{16} \left(4 - \frac{x_1}{y_1} - \frac{x_2}{y_2} + 3 + \frac{x_3}{y_1} + 3 + \frac{x_3}{y_2} \right) = -\frac{1}{2},$$

as desired. \square

8.3 Evaluation of the third term

We now consider the evaluation of the third term in (8.2) given by

$$\frac{1}{2\pi ir} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} E'_{y_j}(y_j) \right]_{21}. \quad (8.20)$$

Proposition 8.8 (Contribution from y_1). *The first term in the sum (8.20) is given by*

$$\frac{x_1}{2\pi ir} \left[E_{y_1}(y_1)^{-1} E'_{y_1}(y_1) \right]_{21} = \frac{8\pi c_0 x_1 p(y_1)}{3\Omega y_1(y_1 - y_2)} \frac{\partial}{\partial r} \log(\theta(\nu)).$$

Proof. A long but direct computation using (5.8) and $\theta'(0) = 0$ gives

$$\frac{x_1}{2\pi ir} \left[E_{y_1}(y_1)^{-1} E'_{y_1}(y_1) \right]_{21} = -x_1 \sqrt{r} \sqrt{c_{y_1}} \varphi_{y_1}^{(\frac{1}{2})} \frac{\theta'(\nu)}{\theta(\nu)} = -\frac{2c_0 \sqrt{r} x_1 p(y_1)}{y_1(y_1 - y_2)} \frac{\theta'(\nu)}{\theta(\nu)}.$$

Recalling that $\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}$, the claim follows. \square

The second and third terms in the sum (8.20) are significantly more complicated to analyze than the first term computed in Proposition 8.8 for two reasons: 1) they require several identities derived in Section 7 (while the first term only uses $\theta'(0) = 0$), and 2) the related computations are significantly longer.

Proposition 8.9 (Contribution from y_2). *The second term in the sum (8.20) is given by*

$$\frac{-x_2}{2\pi ir} \left[E_{y_2}(y_2)^{-1} E'_{y_2}(y_2) \right]_{21} = -\frac{8\pi}{3\Omega} \frac{x_2 p(y_2) c_0}{y_2(y_1 - y_2)} \frac{\partial}{\partial r} \log \theta(\nu).$$

Proof. The proof uses Propositions 3.1, 7.1 and Corollaries 7.2, 7.3. After long computations and considerable simplifications, we obtain

$$\frac{-x_2}{2\pi ir} \left[E_{y_2}(y_2)^{-1} E'_{y_2}(y_2) \right]_{21} = \frac{2c_0 \sqrt{r} x_2 p(y_2)}{(y_1 - y_2) y_2} \frac{\theta'(\nu)}{\theta(\nu)}.$$

The claim follows as in Proposition 8.8 by using $\nu = -\frac{\Omega}{2\pi} r^{\frac{3}{2}}$. \square

Proposition 8.10 (Contribution from y_3). *The third term in the sum (8.20) is given by*

$$\frac{x_3}{2\pi ir} \left[E_{y_3}(y_3)^{-1} E'_{y_3}(y_3) \right]_{21} = \frac{8\pi}{3\Omega} \frac{x_3 p(y_3) c_0}{y_1 y_2} \partial_r \log \theta(\nu).$$

Proof. The proof is similar to the proof of Proposition 8.9, and uses also Propositions 3.1, 7.1 and Corollaries 7.2, 7.3. \square

It follows from Propositions 8.8, 8.9 and 8.10 that

$$\frac{1}{2\pi ir} \sum_{j=1}^3 (-1)^{j+1} x_j \left[E_{y_j}(y_j)^{-1} E'_{y_j}(y_j) \right]_{21} = c_3 \partial_r \log \theta(\nu), \quad (8.21)$$

where the constant c_3 is given by

$$c_3 = \frac{8\pi c_0}{3\Omega} \left(\frac{x_1 p(y_1)}{(x_1 - x_3)(x_1 - x_2)} - \frac{x_2 p(y_2)}{(x_2 - x_3)(x_1 - x_2)} + \frac{x_3 p(y_3)}{(x_1 - x_3)(x_2 - x_3)} \right). \quad (8.22)$$

The three terms of the sum appearing on the left-hand side of (8.21) have been simplified separately in Propositions 8.8, 8.9 and 8.10, so that the expression (8.22) for c_3 does not involve the θ -function. Remarkably, these three terms combine together and it is possible to further simplify c_3 . The tools needed to simplify c_3 are of a different nature than those needed to simplify c_2 in (8.14). In Proposition 8.11 below, we show that $c_3 = 1$ by using Riemann's bilinear identity.

Proposition 8.11. *We have $c_3 = 1$.*

Proof. First, we use the explicit expression (3.3) for p to note that (8.22) can be simplified into

$$c_3 = -\frac{4\pi c_0}{3\Omega}(x_1 + x_2 + x_3). \quad (8.23)$$

We consider the following one-forms on the Riemann surface X :

$$\omega_1 = \frac{dz}{\sqrt{\mathcal{R}(z)}}, \quad \omega_2 = \frac{(p(z) + g_1)dz}{\sqrt{\mathcal{R}(z)}}.$$

Let us deform the cycle B of Figure 3 such that it has finite length and surrounds the interval $[y_2, y_1]$ in the positive direction. Riemann's bilinear identity gives the relation

$$A_1 B_2 - A_2 B_1 = 2\pi i \sum_{p=y_3, y_2, y_1, \infty} \text{Res} \left(\omega_2 \int_{\infty}^P \omega_1; P=p \right),$$

where

$$\begin{aligned} A_1 &= \int_A \omega_1 = 2 \int_0^{y_2} \frac{dx}{|\sqrt{\mathcal{R}(x)}|}, & B_1 &= \int_B \omega_1 = 2i \int_{y_2}^{y_1} \frac{dx}{|\sqrt{\mathcal{R}(x)}|}, \\ A_2 &= \int_A \omega_2 = 2 \int_0^{y_2} \frac{(p(x) + g_1)dx}{|\sqrt{\mathcal{R}(x)}|}, & B_2 &= \int_B \omega_2 = 2i \int_{y_2}^{y_1} \frac{(p(x) + g_1)dx}{|\sqrt{\mathcal{R}(x)}|}. \end{aligned}$$

A simple computation shows that the Abelian differential

$$\omega_2 \int_{\infty}^P \omega_1$$

has no residue at y_3, y_2, y_1 , and has a residue at ∞ given by $-\frac{2}{3}(x_1 + x_2 + x_3)$ in the local coordinate. Therefore, we have

$$A_1 B_2 - A_2 B_1 = -\frac{4\pi i}{3}(x_1 + x_2 + x_3). \quad (8.24)$$

Using the definition of g_1 , Ω and c_0 , it is a direct computation to verify from (8.23) that the claim $c_3 = 1$ is equivalent to (8.24). \square

Theorem 1.1 follows directly by combining Propositions 8.1, 8.2, 8.7 and 8.11, together with equations (8.14) and (8.21).

A Bessel Model RH problem

In this section, we recall the Bessel model RH problem (see [16]) whose solution is denoted by Φ_{Be} .

- (a) $\Phi_{\text{Be}} : \mathbb{C} \setminus \Sigma_{\text{Be}} \rightarrow \mathbb{C}^{2 \times 2}$ is analytic, where Σ_{Be} is shown in Figure 6.
- (b) Φ_{Be} satisfies the jump conditions

$$\begin{aligned} \Phi_{\text{Be},+}(z) &= \Phi_{\text{Be},-}(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, & z &\in \mathbb{R}^-, \\ \Phi_{\text{Be},+}(z) &= \Phi_{\text{Be},-}(z) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, & z &\in e^{\frac{2\pi i}{3}} \mathbb{R}^+, \\ \Phi_{\text{Be},+}(z) &= \Phi_{\text{Be},-}(z) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, & z &\in e^{-\frac{2\pi i}{3}} \mathbb{R}^+. \end{aligned} \quad (\text{A.1})$$




Figure 6: The jump contour Σ_{Be} for Φ_{Be} .

(c) As $z \rightarrow \infty$, $z \notin \Sigma_{\text{Be}}$, we have

$$\Phi_{\text{Be}}(z) = (2\pi z^{\frac{1}{2}})^{-\frac{\sigma_3}{2}} M \left(I + \frac{\Phi_{\text{Be},1}}{z^{\frac{1}{2}}} + \mathcal{O}(z^{-1}) \right) e^{2z^{\frac{1}{2}}\sigma_3}, \quad (\text{A.2})$$

where $\Phi_{\text{Be},1} = \frac{1}{16} \begin{pmatrix} -1 & -2i \\ -2i & 1 \end{pmatrix}$ and $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}$.

(d) As z tends to 0, the behavior of $\Phi_{\text{Be}}(z)$ is

$$\Phi_{\text{Be}}(z) = \begin{cases} \begin{pmatrix} \mathcal{O}(1) & \mathcal{O}(|\log z|) \\ \mathcal{O}(z) & \mathcal{O}(|\log z|) \end{pmatrix}, & |\arg z| < \frac{2\pi}{3}, \\ \begin{pmatrix} \mathcal{O}(|\log z|) & \mathcal{O}(|\log z|) \\ \mathcal{O}(|\log z|) & \mathcal{O}(|\log z|) \end{pmatrix}, & \frac{2\pi}{3} < |\arg z| < \pi. \end{cases} \quad (\text{A.3})$$

The unique solution to this RH problem is given by

$$\Phi_{\text{Be}}(z) = \begin{cases} \begin{pmatrix} I_0(2z^{\frac{1}{2}}) & \frac{i}{\pi} K_0(2z^{\frac{1}{2}}) \\ 2\pi i z^{\frac{1}{2}} I_0'(2z^{\frac{1}{2}}) & -2z^{\frac{1}{2}} K_0'(2z^{\frac{1}{2}}) \end{pmatrix}, & |\arg z| < \frac{2\pi}{3}, \\ \begin{pmatrix} \frac{1}{2} H_0^{(1)}(2(-z)^{\frac{1}{2}}) & \frac{1}{2} H_0^{(2)}(2(-z)^{\frac{1}{2}}) \\ \pi z^{\frac{1}{2}} (H_0^{(1)})'(2(-z)^{\frac{1}{2}}) & \pi z^{\frac{1}{2}} (H_0^{(2)})'(2(-z)^{\frac{1}{2}}) \end{pmatrix}, & \frac{2\pi}{3} < \arg z < \pi, \\ \begin{pmatrix} \frac{1}{2} H_0^{(2)}(2(-z)^{\frac{1}{2}}) & -\frac{1}{2} H_0^{(1)}(2(-z)^{\frac{1}{2}}) \\ -\pi z^{\frac{1}{2}} (H_0^{(2)})'(2(-z)^{\frac{1}{2}}) & \pi z^{\frac{1}{2}} (H_0^{(1)})'(2(-z)^{\frac{1}{2}}) \end{pmatrix}, & -\pi < \arg z < -\frac{2\pi}{3}, \end{cases} \quad (\text{A.4})$$

where $H_0^{(1)}$ and $H_0^{(2)}$ are the Hankel functions of the first and second kind, and I_0 and K_0 are the modified Bessel functions of the first and second kind.

For certain computations, we will need a more precise expansion than (A.3). By [29, Section 10.30(i)], as $z \rightarrow 0$ we have

$$\Phi_{\text{Be}}(z) = \begin{pmatrix} 1 + \mathcal{O}(z) & \mathcal{O}(|\log z|) \\ \mathcal{O}(z) & \mathcal{O}(1) \end{pmatrix}, \quad z \rightarrow 0, \quad |\arg z| < \frac{2\pi}{3}, \quad (\text{A.5})$$

$$\Phi_{\text{Be}}(z)^{-1} = \begin{pmatrix} \mathcal{O}(1) & \mathcal{O}(|\log z|) \\ \mathcal{O}(z) & 1 + \mathcal{O}(z) \end{pmatrix}, \quad z \rightarrow 0, \quad |\arg z| < \frac{2\pi}{3}, \quad (\text{A.6})$$

$$\Phi_{\text{Be}}(z)^{-1} \Phi'_{\text{Be}}(z) = \begin{pmatrix} \mathcal{O}(|\log z|) & \mathcal{O}(z^{-1}) \\ 2\pi i + \mathcal{O}(z) & \mathcal{O}(|\log z|) \end{pmatrix}, \quad z \rightarrow 0, \quad |\arg z| < \frac{2\pi}{3}. \quad (\text{A.7})$$

References

- [1] J. Baik, R. Buckingham, and J. Di Franco, Asymptotics of Tracy-Widom distributions and the total integral of a Painlevé II function, *Comm. Math. Phys.* **280** (2008), 463–497.

- [2] J. Baik, P. Deift, and E. Rains, A Fredholm determinant identity and the convergence of moments for random Young tableaux, *Comm. Math. Phys.* **223** (2001), 627–672.
- [3] F. Bornemann, On the numerical evaluation of Fredholm determinants, *Math. Comp.* **79** (2010), 871–915.
- [4] A. Borodin, A. Okounkov, and G. Olshanski, Asymptotics of Plancherel measures for symmetric groups, *J. Amer. Math. Soc.* **13** (2000), 481–515.
- [5] T. Bothner, P. Deift, A. Its and I. Krasovsky, On the asymptotic behavior of a log gas in the bulk scaling limit in the presence of a varying external potential I, *Comm. Math. Phys.* **337** (2015), 1397–1463.
- [6] P. Bourgade, L. Erdős, and H.-T. Yau, Edge universality of beta ensembles, *Comm. Math. Phys.* **332** (2014), 261–353.
- [7] T. Claeys and A. Doeraene, The generating function for the Airy point process and a system of coupled Painlevé II equations, *Stud. Appl. Math.* **140** (2018), 403–437.
- [8] T. Claeys, A. Its, and I. Krasovsky, Higher order analogues of the Tracy-Widom distribution and the Painlevé II hierarchy, *Comm. Pure Appl. Math.* **63** (2010), 362–412.
- [9] J. des Cloizeaux, M.L. Mehta, Asymptotic behavior of spacing distributions for the eigenvalues of random matrices, *J. Math. Phys.* **14** (1973), 1648–1650.
- [10] P. Deift, Orthogonal polynomials and random matrices: a Riemann-Hilbert approach, *Courant Lecture Notes in Mathematics*, 3. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [11] P. Deift, D. Gioev, Universality at the edge of the spectrum for unitary, orthogonal and symplectic ensembles of random matrices, *Comm. Pure Appl. Math.* **60** (2007), 867–910.
- [12] P. Deift, A. Its, A. Kapaev, and X. Zhou, On the algebro-geometric integration of the Schlesinger equations, *Comm. Math. Phys.* **203** (1999), 613–633.
- [13] P. Deift, A. Its, and I. Krasovsky, Asymptotics for the Airy-kernel determinant, *Comm. Math. Phys.* **278** (2008), 643–678.
- [14] P. Deift, T. Kriecherbauer, K. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.* **52** (1999), 1491–1552.
- [15] P. Deift, T. Kriecherbauer, K.T.-R. McLaughlin, S. Venakides and X. Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, *Comm. Pure Appl. Math.* **52** (1999), 1335–1425.
- [16] P. Deift, A. Its and X. Zhou, A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics, *Ann. of Math.* **146** (1997), 149–235.
- [17] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation, *Ann. of Math.* **137** (1993), 295–368.
- [18] F.J. Dyson, Fredholm determinants and inverse scattering problems, *Comm. Math. Phys.* **47** (1976), 171–183.
- [19] T. Ehrhardt, Dyson’s constant in the asymptotics of the Fredholm determinant of the sine kernel, *Comm. Math. Phys.* **262** (2006), 317–341.
- [20] B. Fahs and I. Krasovsky, Splitting of a gap in the bulk of the spectrum of random matrices, *Duke Math. J.* **168** (2019), 3529–3590.
- [21] B. Fahs and I. Krasovsky, Sine-kernel determinant on two large intervals, preprint, arXiv:2003.08136.

- [22] P.J. Forrester, The spectrum edge of random matrix ensembles, *Nuclear Phys. B* **402** (1993), 709–728.
- [23] I.S. Gradshteyn and I.M. Ryzhik, Table of integrals, series, and products. Seventh edition. Elsevier Academic Press, Amsterdam, 2007.
- [24] A. Its, A.G. Izergin, V.E. Korepin, and N.A. Slavnov, Differential equations for quantum correlation functions, *Proceedings of the Conference on Yang-Baxter Equations, Conformal Invariance and Integrability in Statistical Mechanics and Field Theory*, Volume **4**, (1990) 1003–1037.
- [25] K. Johansson, The arctic circle boundary and the Airy process, *Ann. Probab.* **33** (2005), 1–30.
- [26] I. Krasovsky, Gap probability in the spectrum of random matrices and asymptotics of polynomials orthogonal on an arc of the unit circle, *Int. Math. Res. Not.* **2004** (2004), 1249–1272.
- [27] I. Krasovsky, Large gap asymptotics for random matrices, in “New Trends in Mathematical Physics”, XVth International Congress on Mathematical Physics, *Springer*, 2009.
- [28] I. Krasovsky and T. Maroudas, Airy-kernel determinant on two large intervals, preprint.
- [29] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark, NIST handbook of mathematical functions, (2010) *Cambridge University Press*.
- [30] A. Soshnikov, Universality at the edge of the spectrum in Wigner random matrices, *Comm. Math. Phys.* **207** (1999), 697–733.
- [31] A. Soshnikov, Determinantal random point fields, *Russian Math. Surveys* **55** (2000), 923–975.
- [32] C. Tracy and H. Widom, Level-spacing distributions and the Airy kernel, *Comm. Math. Phys.* **159** (1994), 151–174.
- [33] H. Widom, Asymptotics for the Fredholm determinant of the sine kernel on a union of intervals, *Comm. Math. Phys.* **171** (1995), 159–180.