

# Rates of convergence to the local time of Oscillating and Skew Brownian Motions

Sara Mazzonetto\*

## Abstract

In this paper a class of statistics based on high frequency observations of oscillating Brownian motions and skew Brownian motions is considered. Their convergence rate towards the local time of the underlying process is obtained in form of a Central Limit Theorem.

**Keywords:** Skew Brownian motion, Oscillating Brownian motion, Local time, Functional limit theorems, Central Limit Theorem.

**AMS 2010:** 60F17, 60J55, 60F05.

## 1 Introduction

It is well known that the normalized number of crossings of the level  $r \in \mathbb{R}$  of the time discretization of a Brownian motion provides an estimator for its local time at  $r$ . Roughly speaking the local time at the point  $r$  measures the time the process spends around  $r$  (see (1.2) below for a precise definition), so a rescaled number of crossings for high frequency data is a natural approximation of the local time also for more general processes. Indeed, this was first proven in [3, 4] for Brownian diffusions whose drift and diffusion coefficient  $\sigma$  are sufficiently regular, in particular  $\sigma$  should be continuous. More general functionals of discrete observations can also be considered: given a stochastic process  $(X_t)_{t \in [0, \infty)}$ , let the following statistics for high frequency observation:

$$\varepsilon_{n,t}^{(r,f,X)} := \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} f(\sqrt{n}(X_{k/n} - r), \sqrt{n}(X_{(k-1)/n} - r)) \quad (1.1)$$

where  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  is a measurable function satisfying suitable integrability conditions. In the case of Brownian motion convergence towards the local time and Central Limit Theorem (CLT) were obtained in [7, 8] for these kind of estimators. In the context of Brownian diffusions with regular coefficients mentioned above, convergence results are proven for specific functions  $f$  in [9] and for more general statistics of multivariate diffusions in [13]. In the latter article the author also proves the associated CLT.

In this paper we treat the case of two classes of one-dimensional stochastic processes not covered by the existing literature: oscillating Brownian motion (OBM) and skew Brownian motion (SBM), respectively solutions of SDEs (2.4) and (2.1) below. A peculiarity of these processes is that they change behavior when they reach a point, called barrier or *threshold*,

---

\*Institute of Mathematics, University of Potsdam, Germany; mazzonetto@uni-potsdam.de

which is a *discontinuity point* for the the process local time  $x \mapsto L_t^x(X)$ .

We study the asymptotic behavior of  $\varepsilon_n^{(r,f,X)}$  in case  $X$  is an OBM or a SBM and  $r \in \mathbb{R}$  is its threshold. The convergence towards the local time was proven in the case of SBM in [21, Proposition 2] (see Proposition 1 below). In this article we provide the *rate of convergence* in form of a CLT in Theorem 1. We also prove the convergence result and the associated CLT for an OBM (resp. in Proposition 2 and Theorem 2 below). These CLTs are the main contributions of this paper. For motivation purposes, let us have a look at the following simplified statement: for appropriate constants  $c, k \in \mathbb{R}$  it holds for all  $t \in [0, \infty)$  that

$$n^{\frac{1}{4}} \left( \frac{\varepsilon_{n,t}^{(r,f,X)} - cL_t^r(X)}{k^2 \sqrt{L_t^r(X)}} \right) \underset{n \rightarrow \infty}{\sim} \mathcal{N}(0, 1).$$

The results we propose are more general, they are *functional limit theorems*: we see the processes as random variables with values in the Skorokhod space of càdlàg functions. Although OBM and SBM behave differently with respect to Brownian motion at the threshold, the speed of convergence,  $n^{\frac{1}{4}}$ , is the same as for Brownian motion. Euristically the convergence is different than what one might expect (i.e.  $n^{\frac{1}{2}}$ ), because the local time in  $r$  and its estimator change only when the process is close to  $r$ . And as  $n \rightarrow \infty$ , among  $n$  observations of the process on a fixed interval, the number of those which are sufficiently close to  $r$  to matter is of order  $n^{\frac{1}{2}}$ .

As a direct application of Theorem 1 and Theorem 2, we consider classical estimators of the local time of Brownian motion such as the normalized number of crossings. Since standard Brownian motion (and by the way reflected Brownian motion as well) is a special case of OBM and SBM, we recover the classical results on the convergence (rates) for these estimators.

Other applications of our theoretical results have already been proposed in the literature, for instance:

- In [21, 22] the authors highlight the usefulness of the missing CLT for SBM (our Theorem 1). They study the asymptotic behavior of the maximum likelihood estimator (MLE) of the skewness parameter of a SBM.
- Theorem 2, in particular the example in Proposition 7, is a necessary step for analyzing for instance the convergence rate of a MLE estimator for the drift parameter of the drifted OBM considered in [20].

Remark that the CLTs of this paper hold also for drifted SBM and OBM as soon as the drift allows for Girsanov's transform (see e.g. [19, Theorem 4.19] for a proof that Girsanov's transform does not affect the stable convergence). Extensions to more general statistics of solutions to SDEs with more general discontinuous diffusion coefficient is object of further research.

## 1.1 Outline of the paper

The paper is organized as follows. In Section 2 we state the main result for SBM and OBM in two different sections: Section 2.1 and Section 2.2. In both, we first introduce the process as unique strong solution of a SDE, then we specify the setting and the framework of the result, finally we state existing and new results. The proofs are provided in Section 3. Of particular interest is Section 3.1 whose main focus is to discuss, in the context of the proofs of this article, the strong relationship between OBM and SBM. Moreover we reduce to prove a unique result demonstrated in Appendix B. Appendix A instead deals with useful properties of OBM relevant in this article. Finally, in Section 4, we apply our main results for OBM and SBM to two classical estimators of the local time of Brownian motion.

## 1.2 Notation and notions of convergence

Throughout this document for every measurable functions  $g: \mathbb{R} \rightarrow \mathbb{R}$  and measure  $\mu$  on the Borel space  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$  we denote by  $\langle \mu, g \rangle$  the integral of  $g$  with respect to the measure  $\mu$ :

$$\langle \mu, g \rangle := \int_{-\infty}^{\infty} g(x) \mu(dx).$$

For every  $\gamma \in [0, \infty)$  let  $\lambda^{(\gamma)}$  be the measure on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$  absolutely continuous with respect to the Lebesgue measure satisfying  $\lambda^{(\gamma)}(dx) = |x|^\gamma dx$  and let

$$(\mathbf{L}^1(\lambda^{(\gamma)}), \|\cdot\|_{1,\gamma})$$

the set of Borel measurable  $\lambda^{(\gamma)}$ -integrable functions and its norm. If  $\lambda = 0$ , we simply denote by  $(\mathbf{L}^1, \|\cdot\|_1) := (\mathbf{L}^1(\lambda^{(0)}), \|\cdot\|_{1,0})$  the normed space of Lebesgue integrable functions.

**Definition 1.** Let  $\gamma \in [0, \infty)$ . We denote by  $\mathbf{L}^{1,b}(\lambda^{(\gamma)})$  the following subspace of  $\mathbf{L}^1$ :

$$\mathbf{L}^{1,b}(\lambda^{(\gamma)}) = \{f: \mathbb{R} \rightarrow \mathbb{R}, \text{ measurable and bounded s.t. } f \in \mathbf{L}^1(\lambda^{(\gamma)})\}$$

We denote by  $\mathbf{L}_\gamma$  the following space of bi-variate functions

$$\mathbf{L}_\gamma = \{h: \mathbb{R}^2 \rightarrow \mathbb{R}, \exists \bar{h} \in \mathbf{L}^{1,b}(\lambda^{(\gamma)}), \exists a \in [0, \infty) \text{ s.t. } \forall x, y \in \mathbb{R} : |h(x, y)| \leq \bar{h}(x)e^{a|y-x|}\}.$$

Let us give a more rigorous definition of the local time process. Let  $t \in [0, \infty)$  and let  $(X_s)_{s \in [0, \infty)}$  be a one-dimensional semi-martingale. The *symmetric local time at the point  $r$*  accumulated on the time interval  $[0, t]$  by the semi-martingale  $X$  is

$$L_t^r(X) = \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_0^t \mathbf{1}_{\{-\epsilon \leq X_s - r \leq \epsilon\}} d\langle X \rangle_s \quad (1.2)$$

and if  $r = 0$  we denote  $L_t^0(X)$  by  $L_t(X)$ .

As already mentioned, the main aim of this article is studying, as  $n \rightarrow \infty$ , the convergence towards the local time together with its rate of the statistics  $\varepsilon_{n, \cdot}^{(r, f, X)}$ , with  $X$  being an OBM or a SBM and  $f$  suitable function. Let us recall the notions of convergence used for the results of this paper. The statement of the CLT involves the notion of *stable convergence* which was introduced and studied first in [24] and [2]. We now specify it in the case used in this document.

**Definition 2.** Let  $(D, d)$  be a metric space,  $(\Omega', \mathcal{F}', \mathbb{P}')$  be an extension of the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , let  $X_n: \Omega \rightarrow D$ ,  $n \in \mathbb{N}$ , be a sequence of random variables, and let  $X: \Omega' \rightarrow D$  be a random variable. Then we say that  $X_n$  converges stably in law to  $X$  if for all  $f: D \rightarrow \mathbb{R}$  continuous and bounded and all bounded random variable  $Y: \Omega \rightarrow \mathbb{R}$  it holds that

$$\lim_{n \rightarrow \infty} \mathbb{E}[f(X_n)Y] = \mathbb{E}'[f(X)Y].$$

Let  $t \in [0, \infty)$ , let  $\mathbb{D}_t$ , resp.  $\mathbb{D}_\infty$ , be the Skorokhod space of càdlàg functions from  $[0, t]$ , resp.  $[0, \infty)$ , to  $\mathbb{R}$  endowed with the Skorokhod topology. When  $D = \mathbb{D}_t$ ,  $t \in [0, \infty]$  the functional stable convergence in law is usually denoted by

$$X_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} X.$$

Finally we recall the notion used in the convergence results, i.e. the *convergence in probability locally uniformly in time* or convergence *uniform on compacts in probability* (u.c.p.): let  $X, X_n$ ,  $n \in \mathbb{N}$ , be random variables with values in  $\mathbb{D}_\infty$ , then

$$X_n \xrightarrow{\text{u.c.p.}} X,$$

if for all  $t \in [0, \infty)$  it holds that  $\sup_{s \in [0, t]} |X_n(s) - X(s)| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ .

## 2 Main results

In this section we introduce the processes SBM and OBM and we provide the main results of this article.

In the entire document let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, \infty)}, \mathbb{P})$  be a stochastic basis (i.e. a complete filtered probability space whose filtration satisfies the usual conditions) and  $W$  be an  $(\mathcal{F}_t)_{t \in [0, \infty)}$ -adapted Wiener process.

### 2.1 The case of a skew Brownian motion

Roughly speaking a SBM can be described trajectoryally as a standard Brownian motion transformed by flipping its excursions from the origin with a certain probability. In this document we refer to the characterization as solution of a SDE involving the local time, which was first considered by [10]. We refer the reader to the recent survey paper on SBM [18].

Let  $x_0 \in \mathbb{R}$ . The SBM with skewness parameter  $\beta \in [-1, 1]$  at the threshold  $r \in \mathbb{R}$  is the diffusion which is strong solution of the following SDE

$$X_t = x_0 + W_t + \beta L_t^r(X) \quad (2.1)$$

where  $L_t^r(X)$  is the symmetric local time of the process at  $r$ , and  $\beta X_0 \geq 0$  if  $|\beta| = 1$ . Some properties of the local time of SBM are object of the recent paper [6].

We call *standard SBM* a SBM with threshold  $r = 0$  starting at 0. In this paper a SBM with skewness parameter  $\beta \in (-1, 1)$  is also denoted by  $\beta$ -SBM. Note that a 0-SBM is a Brownian motion. Moreover the  $\pm 1$ -SBM is a positively/negatively reflected Brownian motion.

#### 2.1.1 Notation and framework

We introduce some notation and a recent result about the convergence towards the local time.

Let  $\mu_\beta$  be the stationary measure associated to the standard  $\beta$ -SBM, that is

$$\mu_\beta(dx) := ((1 + \beta)\mathbf{1}_{(0, \infty)}(x) + (1 - \beta)\mathbf{1}_{(-\infty, 0)}(x)) dx = (1 + \operatorname{sgn}(x)\beta) dx,$$

and let  $p_\beta(t, x, y)$  its transition density (first computed in [25]) which satisfies

$$p_\beta(t, x, y) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(x-y)^2}{2t}\right) + \beta \operatorname{sgn}(y) \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(|x| + |y|)^2}{2t}\right). \quad (2.2)$$

Given two measurable functions  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$ , let

$$\mathbf{F}_{f,g}(x) = \int_{-\infty}^{\infty} f(x, y)g(y)p_\beta(1, x, y) dy \quad \text{and} \quad \mathbf{F}_f := \mathbf{F}_{f,1} \quad (\text{i.e. } g \equiv 1). \quad (2.3)$$

*Hypothesis 1.* The measurable bi-variate function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies that  $\mathbf{F}_f, \mathbf{F}_{f^2} \in \mathbf{L}^{1,b}(\lambda^{(2)})$ .

**Proposition 1** (Convergence towards the scaled local time, cf. [22, Proposition 2]). *Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function satisfying Hypothesis 1 and for every  $\beta \in [-1, 1]$  let  $X^\beta$  be the  $\beta$ -SBM strong solution of (2.1). Then for all  $t \in (0, \infty)$  it holds for all  $\varepsilon \in (0, \infty)$  that*

$$\lim_{n \rightarrow \infty} \sup_{\beta \in [-1, 1]} \mathbb{P} \left( \sup_{s \in [0, t]} \left| \varepsilon_{n,s}^{(r, f, X^\beta)} - \langle \mu_\beta, \mathbf{F}_f \rangle L_s^r(X^\beta) \right| \geq \varepsilon \right) = 0.$$

Proposition 1 follows from [22, Proposition 2] (with  $T = 1$ ) and the scaling property. Remark that the multiplicative factor in front of the local time is

$$\langle \mu_\beta, \mathbf{F}_f \rangle = \mathbb{E} \left[ f(X_0^\beta, X_1^\beta) | X_0^\beta \sim \mu_\beta \right].$$

### 2.1.2 Rate of convergence to the local time of a SBM

We refine the above convergence showing that the speed of convergence is of order  $1/4$ .

**Theorem 1.** *Let  $f \in \mathbf{I}_\gamma$ ,  $\gamma > 3$ , let  $X$  be the  $\beta$ -SBM solution of (2.1). Then there exists a Brownian motion  $B$  independent of  $X$  (possibly on an extension of the probability space) such that*

$$n^{1/4} \left( \varepsilon_{n,\cdot}^{(r,f,X)} - \langle \mu_\beta, \mathbf{F}_f \rangle L^r(X) \right) \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_f} B_{L^r(X)},$$

where

$$\begin{aligned} K_f &= \langle \mu_\beta, \mathbf{F}_{f^2} + 2\mathbf{F}_{f,\mathcal{P}_\beta} \rangle + \frac{8}{3\sqrt{2\pi}} (\langle \mu_\beta, \mathbf{F}_f \rangle)^2 \\ &\quad - 2\sqrt{\frac{2}{\pi}} \langle \mu_\beta, \mathbf{F}_f \rangle \int_{-\infty}^{\infty} \left( e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y|\Phi(-|y|) \right) \mathcal{P}_\beta(y) \mu_\beta(dy) dt \\ &\quad - 2\langle \mu_\beta, \mathbf{F}_f \rangle \int_{-\infty}^{\infty} \int_0^1 \int_{-\infty}^{\infty} \frac{|x|e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} \sqrt{\frac{1}{t} - 1} \Phi(-|y|) f(x\sqrt{t}, y\sqrt{1-t}) \mu_\beta(dy) dt \mu_\beta(dx), \end{aligned}$$

$\Phi$  is the cumulative distribution function of a standard Gaussian random variable,

$$\mathcal{P}_\beta(x) = \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} p_\beta(j, x, y) (\mathbf{F}_f(y) - \langle \mu_\beta, \mathbf{F}_f \rangle \mathbf{F}_{g_\beta}(y)) dy,$$

and  $g_\beta(x, y) = \frac{1}{1+\text{sgn}(y)\beta} \left( |y| - \frac{1+\text{sgn}(y)\beta}{1+\text{sgn}(x)\beta} |x| \right)$ .

*Remark 1.* If  $\beta = 0$  we recover the known result for Brownian motion: e.g. [7, 8] and a special case of the already cited [12, Theorem 1.2]. The expression for the constant  $K_f$  we propose is slightly more explicit.

## 2.2 The case of an oscillating Brownian motion

Let  $y_0 \in \mathbb{R}$ . The strong solution to the SDE

$$Y_t = y_0 + \int_0^t \sigma(Y_s) dW_s, \quad t \geq 0, \quad (2.4)$$

is called OBM with threshold  $r \in \mathbb{R}$  when the diffusion coefficient  $\sigma$  is the positive two-valued function discontinuous at the threshold:

$$\sigma := \sigma_- \mathbf{1}_{(-\infty, r)} + \sigma_+ \mathbf{1}_{[r, +\infty)}. \quad (2.5)$$

The existence of a strong solution to (2.4) follows for instance from the results of [17]. This process has been first defined and studied in [16]. We call *standard OBM* an OBM  $Y$  with threshold  $r = 0$  and starting point  $Y_0 = 0$ .

We can allow either  $\sigma_-$  or  $\sigma_+$  to be infinity: If  $\sigma_+ = 1$ ,  $\sigma_- = +\infty$ ,  $Y_0 \geq 0$  (resp.  $\sigma_- = 1$ ,  $\sigma_+ = +\infty$ ,  $Y_0 \leq 0$ ) then it is a positively (resp. negatively) reflected Brownian motion.

### 2.2.1 Notation and framework

In this section we provide in Proposition 2 the counterpart for OBM of the convergence result for SBM stated above in Proposition 1. To the best of our knowledge, this is the first time that the convergence to the local time of OBM is formulated at this level of generality.

We first introduce some notation. The stationary measure for the standard OBM is

$$\lambda_\sigma(dx) := \frac{1}{(\sigma(x-r))^2} dx = \left( \frac{1}{\sigma_-^2} \mathbf{1}_{(-\infty, 0)} + \frac{1}{\sigma_+^2} \mathbf{1}_{[0, \infty)} \right) dx$$

and its transition density, here denoted by  $q_\sigma(t, x, y)$ , satisfies

$$q_\sigma(t, x, y) = \frac{1}{\sigma(y-r)} p_{\beta_\sigma} \left( t, \frac{x}{\sigma(x-r)}, \frac{y}{\sigma(y-r)} \right) \quad \text{with} \quad \beta_\sigma := \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}, \quad (2.6)$$

where  $p_{\beta_\sigma}$  is the density of the SBM recalled in (2.2) (see, e.g. equation (3) in [19] for an explicit expression). Given two measurable functions  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$ , let

$$\mathbf{H}_{f,g}(x) = \int_{-\infty}^{\infty} f(x, y) g(y) q_\sigma(1, x, y) dy \quad \text{and} \quad \mathbf{H}_f(x) := \mathbf{H}_{f,1}(x). \quad (2.7)$$

*Hypothesis 2.* The measurable bi-variate function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies that  $\mathbf{H}_f, \mathbf{H}_{f^2} \in \mathbf{L}^{1,b}(\lambda^{(2)})$ .

**Proposition 2** (Convergence towards the scaled local time). *Let  $h: \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfying Hypothesis 2 and let  $Y$  be the OBM solution to (2.4). Then*

$$\varepsilon_{n,\cdot}^{(r,h,Y)} \xrightarrow[n \rightarrow \infty]{\text{u.c.p.}} \langle \lambda_\sigma, \mathbf{H}_h \rangle L^r(Y).$$

Note that, similarly to Proposition 1, the convergence in Proposition 2 is uniform in the choice of the pair of parameters  $\sigma_-, \sigma_+ \in (0, \infty]$ .

Moreover observe that the constant  $\langle \lambda_\sigma, \mathbf{H}_h \rangle$  can be rewritten as

$$\langle \lambda_\sigma, \mathbf{H}_h \rangle = \mathbb{E}[h(Y_0, Y_1) | Y_0 \sim \lambda_\sigma].$$

### 2.2.2 Rate of convergence to the local time of an OBM

We refine the above convergence showing that the rate is of order 1/4 in the following CLT:

**Theorem 2.** *Let  $h \in \mathbf{I}_\gamma$ ,  $\gamma > 3$ , and let  $Y$  be the OBM solution to (2.4). Then there exists a Brownian motion  $B$  independent of  $Y$  (possibly on an extension of the probability space) such that*

$$n^{1/4} \left( \varepsilon_{n,\cdot}^{(r,h,Y)} - \langle \lambda_\sigma, \mathbf{H}_h \rangle L^r(Y) \right) \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_h} B_{L^r(Y)}$$

where

$$\begin{aligned} K_h &= \langle \lambda_\sigma, \mathbf{H}_{h^2} + 2\mathbf{H}_{h,\mathfrak{O}_{\sigma,h}} \rangle + \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \frac{8}{3\sqrt{2\pi}} (\langle \lambda_\sigma, \mathbf{H}_h \rangle)^2 \\ &\quad - 2\sqrt{\frac{2}{\pi}} \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \langle \lambda_\sigma, \mathbf{H}_h \rangle \int_{-\infty}^{\infty} \sigma(y) \left( e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y|\Phi(-|y|) \right) \mathfrak{O}_{\sigma,h}(\sigma(y)y) \lambda_\sigma(dy) \\ &\quad - 2\langle \lambda_\sigma, \mathbf{H}_h \rangle \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \\ &\quad \cdot \int_{-\infty}^{\infty} \int_0^1 \int_{-\infty}^{\infty} \frac{|x| e^{-\frac{x^2}{2}} \Phi(-|y|)}{\sqrt{2\pi}\sigma(x)\sigma(y)} \sqrt{\frac{1}{t} - 1} h(\sigma(x)x\sqrt{t}, \sigma(y)y\sqrt{1-t}) dy dt dx, \end{aligned} \quad (2.8)$$




Figure 1: Map of the proofs.

Proposition 1, which follows from an existing result, implies Proposition 2. Theorem 1 follows from Theorem 2 which follows from the combination of Proposition 4 and (the existing) Proposition 3. Proposition 4 is the cornerstone of this article.

$\Phi$  the cumulative distribution function of a standard normal random variable,

$$\mathfrak{G}_{\sigma,h}(x) = \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} q_{\sigma}(j, x, y) (\mathbf{H}_h(y) - \langle \lambda_{\sigma}, \mathbf{H}_h \rangle \mathbf{H}_g(y)) dy \quad (2.9)$$

and  $g(x, y) = |y| - |x|$ .

*Remark 2.* Theorem 2 implies a weaker version of Proposition 2. Proposition 2 requires Hypothesis 2 which is satisfied if for instance  $h \in \mathbf{I}_2$ . Theorem 2 instead assumes  $h \in \mathbf{I}_{\gamma}$ ,  $\gamma > 3$ , which is a stronger condition.

Let us comment on how to derive the u.c.p. convergence from Theorem 2: The notions of convergence in law/stably in law/probability coincide when the limit is constant and so  $\varepsilon_{n,\cdot}^{(r,h,Y)} - \langle \mu_{\beta}, \mathbf{H}_h \rangle L^r(Y) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$  in the Skorokhod topology. Since  $L^r(Y)$  is (a.s.) continuous and increasing it can be proven (splitting into positive and negative part of  $h$ , and so of  $\mathbf{H}_h$ ) that  $\varepsilon_{n,\cdot}^{(r,h,Y)} \xrightarrow[n \rightarrow \infty]{\text{u.c.p.}} \langle \mu_{\beta}, \mathbf{H}_h \rangle L^r(Y)$  (see e.g. (2.2.16) in [14]).

### 3 Proofs of the main results

In this section we comment the results and their proof: The convergence in probability to the local time in Proposition 2 and the rate of (stable) convergence in the case of SBM and OBM, respectively Theorem 1 and Theorem 2.

The relationship between SBM and OBM is crucial to obtain Proposition 2 from its already proven analogous for SBM Proposition 1 and the rate in the case of SBM in Theorem 1 from Theorem 2. This is the content of Section 3.1.

Finally Theorem 2 is proved in Section 3.3 relying on an a well known CLT stated in Section 3.2.

Figure 1 is a map of what we just described.

#### 3.1 The interplay between SBM and OBM

SBM and OBM are strongly related in the following sense: Let  $r \in \mathbb{R}$ , let  $\sigma$  be the function in (2.5), let  $Y$  the solution to (2.4) and  $X$  be a SBM solution to (2.1) with skewness parameter  $\beta_{\sigma} := \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}$  and suitable initial condition: Solution to the SDE

$$X_t = \frac{Y_0}{\sigma(Y_0)} + W_t + \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+} L_t^r(X). \quad (3.1)$$

It holds that  $Y_t = \sigma(X_t)X_t$ , or equivalently  $\frac{Y_t}{\sigma(Y_t)} = X_t$ , and the local times satisfy

$$L^r(X) = \frac{\sigma_+ + \sigma_-}{2\sigma_+\sigma_-} L^r(Y) \quad (3.2)$$

(see, e.g. [19, page 3573]).

Let us recall, given a measurable function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  the definition of the quantity  $\mathbf{H}_f$  in (2.7), and for  $\beta \in [-1, 1]$  the functions  $\mathbf{F}_f$  in (2.3).

We are now ready to approach the proof of Proposition 2 for the OBM  $Y$ .

*Proof of Proposition 2.* Let  $X$  be the SBM associated to the OBM  $Y$ , i.e.  $X$  is a  $\beta_\sigma$ -SBM with skewness parameter  $\beta_\sigma = \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}$  solution to (3.1). Let  $h_\sigma$  be the function satisfying  $h_\sigma(x) := h(\sigma(x)x)$ . Note that a function  $h$  satisfies Hypothesis 2 if and only if  $h_\sigma$  satisfies Hypothesis 1. Moreover it holds that  $\mathbf{H}_{h_\sigma}(\sigma(x)x) = \mathbf{F}_{(h_\sigma)\gamma}(x)$  and so  $\langle \mu_{\beta_\sigma}, \mathbf{F}_{(h_\sigma)\gamma} \rangle = \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \langle \lambda_\sigma, \mathbf{H}_{h_\sigma} \rangle$ . Applying Proposition 1 for the SBM  $X$  and the function  $f := h_\sigma$  and taking into account the latter equalities and the relationship between local times (3.2) complete the proof.  $\square$

Now, assume that Theorem 2 holds, we prove now the analogous for SBM.

*Proof of Theorem 1.* Let  $\beta \in [-1, 1]$ ,  $\gamma > 3$ ,  $X^\beta$  the  $\beta$ -SBM solution to (2.1), and  $f \in \mathbf{I}_\gamma$ .

Take  $\sigma_+^{(\beta)}, \sigma_-^{(\beta)} \in (0, \infty]$  such that  $\beta = \frac{\sigma_-^{(\beta)} - \sigma_+^{(\beta)}}{\sigma_-^{(\beta)} + \sigma_+^{(\beta)}}$  and construct the diffusion coefficient  $\sigma_\beta := \sigma_-^{(\beta)} \mathbf{1}_{(-\infty, 0)} + \sigma_+^{(\beta)} \mathbf{1}_{[0, \infty)}$  in (2.5). Let  $Y^\beta$  be the OBM with this diffusion coefficient  $\sigma_\beta$  and initial condition  $Y_0^\beta = \sigma_\beta(X_0)X_0$  and let  $h_\beta: \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function satisfying  $h_\beta(x) := f(x/\sigma_\beta(x))$ . Note that  $f \in \mathbf{I}_\gamma$  if and only if  $h_\beta \in \mathbf{I}_\gamma$ . Theorem 2 can be applied to the OBM  $Y^\beta$  and the function  $h_\beta$  to obtain that

$$n^{1/4} \left( \mathbb{E}_{n, \cdot}^{(r, h_\beta, Y^\beta)} - \langle \lambda_{\sigma_\beta}, \mathbf{H}_{h_\beta} \rangle L^r(Y^\beta) \right) \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_f} B_{L^r(Y^\beta)}$$

with  $K_f := K_{h_\beta}$  in (2.8). The relationship between  $X^\beta$  and  $Y^\beta$ , in particular between their local time (3.2) and their transition densities (2.6), and the fact that  $\langle \lambda_{\sigma_\beta}, \mathbf{H}_{h_\beta} \rangle = \frac{\sigma_-^{(\beta)} + \sigma_+^{(\beta)}}{2\sigma_+^{(\beta)}\sigma_-^{(\beta)}} \langle \mu_\beta, \mathbf{F}_f \rangle$  are used to rewrite the constant  $K_f$  and the limit in the desired form.  $\square$

## 3.2 A Central Limit Theorem

In this paragraph we reformulate a special case of Theorem 3.2 in [12].

**Proposition 3** (cf. Theorem 3.2 in [12]). *Let  $(Y_t)_{t \in [0, 1]}$  be an  $(\mathcal{F}_t)_{t \in [0, 1]}$ -local martingale on the stochastic basis  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, 1]}, \mathbb{P})$ . Let  $\mathcal{Z}^n = \sum_{k=1}^{\lfloor nt \rfloor} \chi_k^n$  where  $\chi_k^n$  are square integrable  $\mathcal{F}_{\frac{k}{n}}$ -measurable,  $n \in \mathbb{N}$ , and assume that there are  $E$  and  $F$  continuous processes on  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, 1]}, \mathbb{P})$  such that  $E$  has bounded variation and it holds*

$$i) \sup_{s \in [0, 1]} \left| \sum_{k=1}^{\lfloor ns \rfloor} \mathbb{E}[\chi_k^n | \mathcal{F}_{(k-1)/n}] - E_s \right| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0,$$

$$ii) \text{ for all } t \in [0, 1] \text{ that } \sum_{k=1}^{\lfloor nt \rfloor} \left( \mathbb{E}[(\chi_k^n)^2 | \mathcal{F}_{(k-1)/n}] - (\mathbb{E}[\chi_k^n | \mathcal{F}_{(k-1)/n}])^2 \right) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} F_t,$$

$$iii) \text{ for all } t \in [0, 1] \text{ that } \sum_{k=1}^{\lfloor nt \rfloor} \mathbb{E}[\chi_k^n (Y_{k/n} - Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0,$$

$$iv) \text{ for all } \varepsilon \in (0, \infty) \text{ that } \sum_{k=1}^n \mathbb{E}[|\chi_k^n|^2 \mathbf{1}_{\{|\chi_k^n| \geq \varepsilon\}} | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0, \text{ and}$$



v) for all  $t \in [0, 1]$  and  $M$  bounded  $(\mathcal{F}_t)_{t \in [0, 1]}$ -martingale such that for all  $s \in [0, 1]$  the cross variation satisfies  $\mathbb{P}(\langle M, Y \rangle_s = 0) = 1$  that

$$\sum_{k=1}^{\lfloor nt \rfloor} \mathbb{E}[\chi_k^n (M_{k/n} - M_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

Then there exists a Brownian motion  $B$ , possibly on an extension of the probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, 1]}, \mathbb{P})$ , such that  $B$  and  $Y$  are independent and

$$\mathcal{Z}_t^n \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-}s} E. + B_E.$$

### 3.3 Proof of the Central Limit Theorem for OBM

We introduce now Proposition 4 and explain its key role in the proof of the main Theorem 2.

**Proposition 4.** *Let  $(Y_t)_{t \in [0, 1]}$  be the OBM with threshold  $r = 0$  strong solution to (2.4) on the stochastic basis  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, 1]}, \mathbb{P})$ , let  $\gamma > 3$ , and let  $h \in \mathbf{I}_\gamma$ .*

*Then there exist sequences of stochastic process  $(\mathcal{V}_t^n)_{t \in [0, 1]}$ ,  $n \in \mathbb{N}$ , with*

$$\sup_{s \in [0, 1]} m^{\frac{1}{4}} |\mathcal{V}_s^m| \xrightarrow[m \rightarrow \infty]{\mathbb{P}} 0,$$

$(\mathcal{F}_{\lfloor nt \rfloor / n})_{t \in [0, 1]}$ -martingales  $(\mathcal{M}_t^n)_{t \in [0, 1]}$ ,  $n \in \mathbb{N}$ , and random variables  $(\chi_k^n)_{k \in \{1, \dots, \lfloor nt \rfloor\}}$ ,  $n \in \mathbb{N}$ , such that it holds for all  $t \in [0, 1]$ ,  $n \in \mathbb{N}$  it holds that  $\mathcal{M}_t^n = \sum_{k=1}^{\lfloor nt \rfloor} \chi_k^n$  and

$$n^{\frac{1}{4}} \left( \varepsilon_{n,t}^{(0,h,Y)} - \langle \lambda_\sigma, \mathbf{H}_h \rangle L_t(Y) \right) = \mathcal{M}_t^n + n^{\frac{1}{4}} \mathcal{V}_t^n, \quad (3.3)$$

and it holds

i) for all  $k \in \{1, \dots, \lfloor nt \rfloor\}$   $\chi_k^n$  is square integrable  $\mathcal{F}_{\frac{k}{n}}$ -measurable and  $\mathbb{E}[\chi_k^n | \mathcal{F}_{(k-1)/n}] = 0$ ,

ii) for all  $t \in [0, 1]$  that  $\sum_{k=1}^{\lfloor nt \rfloor} \mathbb{E}[(\chi_k^n)^2 | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} K_h L_t(Y)$  with  $K_h$  given by (2.8),

iii) for all  $t \in [0, 1]$  that  $\sum_{k=1}^{\lfloor nt \rfloor} \mathbb{E}[\chi_k^n (Y_{k/n} - Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ ,

iv) for all  $\varepsilon \in (0, \infty)$  that  $\sum_{k=1}^n \mathbb{E}[|\chi_k^n|^2 \mathbf{1}_{\{|\chi_k^n| \geq \varepsilon\}} | \mathcal{F}_{(k-1)/n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ .

The proof of this result is provided in Appendix B. It consists in generalizing to the case of OBM the fundamental procedure used in [12] for Brownian motion.

Let us now assume that Proposition 4 holds and let us prove Theorem 2.

*Proof of Theorem 2.* Let  $Y$  be the OBM with threshold  $r$  strong solution to (2.4).

Without loss of generality we can assume in the proof of Theorem 2 the threshold  $r$  to be 0, because  $Y - r$  is an OBM with threshold  $r = 0$ .

Moreover we now show that we can reduce ourselves to prove Theorem 2 on the interval  $[0, 1]$  for  $n \in \mathbb{N}$  tending to infinity.

The scaling property for the OBM and its local time (see (A.1)) yields the result for all non-negative times: as processes on  $\mathbb{D}_{[0,t]}$ . (The scaling property also ensures that in Theorem 1, and so in Theorem 2,  $n$  is not necessarily a natural number, but it can stay for a positive real number tending to infinity.) Since  $B_{L(Y)}$  has (a.s.) continuous trajectories, it follows, e.g. combining [5,

Theorem 16.2] and [14, Proposition 2.2.4] that  $\varepsilon_{n,\cdot}^{(0,h,Y)} - \langle \lambda_\sigma, \mathbf{H}_h \rangle L_\cdot(Y) \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_h} B_{L_\cdot(Y)}$  in  $\mathbb{D}_\infty$  if and only if for all  $t \in [0, \infty)$   $(\varepsilon_{n,\cdot}^{(0,h,Y)} - \langle \lambda_\sigma, \mathbf{H}_h \rangle L_\cdot(Y))|_{[0,t]} \xrightarrow[n \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_h} B_{L_\cdot(Y)}|_{[0,t]}$  in  $\mathbb{D}_t$ .

Proposition 4 implies that there exists a decomposition as in (3.3) and its desired stable limit as  $n \in \mathbb{N}$  goes to infinity coincides with the stable limit of the sequence  $\mathcal{M}^n$  of càdlàg  $(\mathcal{F}_{\lfloor ns \rfloor / n})_{s \in [0,1]}$ -martingales,  $n \in \mathbb{N}$ . Indeed the fact that  $\sup_{s \in [0,1]} n^{\frac{1}{4}} |\mathcal{V}_s^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ , implies that for every  $f: \mathbb{D}_1 \rightarrow \mathbb{R}$  continuous and bounded it holds that  $|f(\mathcal{M}^n + n^{\frac{1}{4}} \mathcal{V}^n) - f(\mathcal{M}^n)| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$  and so for every bounded continuous function  $f: \mathbb{D}_1 \rightarrow \mathbb{R}$  and bounded measurable random variable  $Y: \Omega \rightarrow \mathbb{R}$  it holds that  $\lim_{n \rightarrow \infty} \mathbb{E} \left[ |f(\mathcal{M}^n + n^{\frac{1}{4}} \mathcal{V}^n) - f(\mathcal{M}^n) Y| \right] = 0$ .

Proposition 4 also ensures that  $\mathcal{M}^n$ ,  $n \in \mathbb{N}$ , satisfies all assumptions, except Item (v), of Proposition 3 (with local martingale  $M = Y$ ,  $\mathcal{Z}^n = \mathcal{M}^n$ ,  $E \equiv 0$  and  $F = K_h L(Y)$  where  $K_h$  is the constant in equation (2.8)). Item (v) of Proposition 3 is trivial due to Lemma 1 in Appendix A.3.

Therefore, applying Proposition 3 as described above completes the proof of Theorem 2.  $\square$

## 4 Applications: two estimators for the local time

Let  $r \in \mathbb{R}$ , and let  $\xi$  be either the OBM with threshold  $r$  solution of (2.4) or a SBM with skewness  $\beta$  in  $r$ . Let  $T \in (0, \infty)$ ,  $N \in \mathbb{N}$ , we observe the process on the discrete time grid  $i \frac{T}{N}$ , for  $i = 0, \dots, N$ . We denote by  $\xi_i = \xi_{i \frac{T}{N}}$ .

Let  $\alpha \in [0, \infty)$  and note that the function  $h_\alpha$  given by  $h_\alpha(x, y) = |y|^\alpha \mathbf{1}_{\{xy < 0\}}$  is in  $\mathbf{I}_\gamma$  for all  $\gamma \in [0, \infty)$ . In fact  $h_\alpha(x, y) \leq c_\alpha e^{-|x|} e^{|y-x|}$  for some constant  $c_\alpha$  depending on  $\alpha$ . We consider two estimators obtained considering the functions proportional to  $h_0$  and  $h_1$ : for  $t \in (0, \infty)$

$$\mathcal{L}_{T,N,t}^r(\xi) = \varepsilon_{\frac{N}{T}, t}^{(r, h_0, \xi)} = \sqrt{\frac{T}{N}} \sum_{i=0}^{\lfloor Nt/T \rfloor - 1} \mathbf{1}_{\{(\xi_i - r)(\xi_{i+1} - r) < 0\}} \quad \text{and} \quad (4.1)$$

$$L_{T,N,t}^r(\xi) = \varepsilon_{\frac{N}{T}, t}^{(r, 2h_1, \xi)} = 2 \sum_{i=0}^{\lfloor Nt/T \rfloor - 1} \mathbf{1}_{\{(\xi_i - r)(\xi_{i+1} - r) < 0\}} |\xi_{i+1} - r|. \quad (4.2)$$

The first counts the number of crossing of the threshold and the second takes into account the distance from the threshold.

As mentioned in the introduction, in the case of Brownian motion, and more general Brownian diffusions, these are consistent estimators of the local time up to a constant (e.g. the already cited [4], [7], [13], and also [23] dealing with fractional Brownian motion).

Throughout this section let the function  $\Phi: \mathbb{R} \rightarrow [0, 1]$  be the cumulative distribution function of a standard Gaussian random variable: for all  $x \in \mathbb{R}$  it holds  $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{y^2}{2}} dy$ .

### 4.1 Estimator counting the number of crossings of the threshold

The next proposition is a consequence of Theorem 2 applied with  $h = h_0$ .

**Proposition 5.** *Let  $Y$  be the OBM solution to (2.4). The estimator  $\mathcal{L}_{T,N}^r(Y)$  in (4.1), counting the number of times the OBM  $Y$  crosses its threshold  $r$ , satisfies that  $\mathcal{L}_{T,N}^r(Y) \xrightarrow[N \rightarrow \infty]{\text{u.c.p.}}$*

*$\frac{2}{\sigma_- + \sigma_+} \sqrt{\frac{2}{\pi}} L^r(Y)$  and  $N^{\frac{1}{4}} \left( \mathcal{L}_{T,N}^r(Y) - \frac{2}{\sigma_- + \sigma_+} \sqrt{\frac{2}{\pi}} L^r(Y) \right) \xrightarrow[N \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_\sigma} B_{L^r(Y)}$  where  $B$  is a Brow-*

nian motion, possibly on an extension of the probability space, independent from  $Y$ ,

$$\begin{aligned} \frac{\sigma_- + \sigma_+}{2} \sqrt{\frac{\pi}{2}} K_\sigma &= 1 + \sqrt{2\pi} \int_{-\infty}^{\infty} \Phi(-|x|) \mathfrak{O}_{\sigma,h}(\sigma(x)x) dx + \frac{4\sigma_- \sigma_+}{(\sigma_- + \sigma_+)^2} \frac{8}{3\pi} \\ &\quad - \frac{4}{\sigma_- + \sigma_+} \sqrt{\frac{2}{\pi}} \int_{-\infty}^{\infty} \sigma(-x) (e^{-\frac{x^2}{2}} - \sqrt{2\pi}|x| \Phi(-|x|)) \mathfrak{O}_{\sigma,h}(\sigma(x)x) dx \\ &\quad - \frac{4\sigma_- \sigma_+}{(\sigma_- + \sigma_+)^2}, \end{aligned} \quad (4.3)$$

and the function  $\mathfrak{O}_{\sigma,h}$  defined in (2.9) has the following expression

$$\mathfrak{O}_{\sigma,h}(\sigma(x)x) = \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} \sigma(y) q_\sigma(j, \sigma(x)x, \sigma(y)y) \mathbf{G}^{(\sigma)}(\sigma(y)y) dy,$$

where  $\mathbf{G}^{(\sigma)}(\sigma(y)y) = \frac{2}{\sigma_+ + \sigma_-} \left( \sigma(y) \Phi(-|y|) - \frac{2}{\pi} \frac{2\sigma_- \sigma_+}{\sigma_+ + \sigma_-} \left( e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y| \Phi(-|y|) \right) \right)$ .

The counterpart for any SBM can be obtained either using the latter proposition and the strong relationship with OBM (see Section 3.1) or applying the specific result for SBM: Theorem 1 with  $h = h_0$ .

**Proposition 6.** *Let  $X$  be the solution to (2.1). Then the estimator  $\mathcal{L}_{T,N}^r(X)$  in (4.1) satisfies that  $\mathcal{L}_{T,N}^r(X) \xrightarrow[N \rightarrow \infty]{\text{u.c.p.}} \sqrt{\frac{2}{\pi}}(1 - \beta^2)L^r(X)$  and*

$$N^{\frac{1}{4}} \left( \mathcal{L}_{T,N}^r(X) - \sqrt{\frac{2}{\pi}}(1 - \beta^2)L^r(X) \right) \xrightarrow[N \rightarrow \infty]{\mathcal{L}\text{-s}} \sqrt{K_\beta} B_{L^r(X)}$$

where  $B$  is a Brownian motion, possibly on an extension of the probability space, independent from  $X$ ,

$$\begin{aligned} \sqrt{\frac{\pi}{2}} \frac{1}{1 - \beta^2} K_\beta &= 1 + \sqrt{2\pi} \int_{-\infty}^{\infty} \Phi(-|x|) \mathcal{P}_\beta(x) dx + \frac{8}{3\pi} (1 - \beta^2) \\ &\quad - 2\sqrt{\frac{2}{\pi}} (1 - \beta^2) \int_{-\infty}^{\infty} (e^{-\frac{x^2}{2}} - \sqrt{2\pi}|x| \Phi(-|x|)) \mathcal{P}_\beta(x) dx - (1 - \beta^2), \end{aligned}$$

and

$$\mathcal{P}_\beta(x) = \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} p_\beta(j, x, y) \left( \Phi(-|y|) - \frac{2}{\pi} (1 + \text{sgn}(y)\beta) \left( e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y| \Phi(-|y|) \right) \right) dy.$$

Clearly the right hand side of (4.3) is equal to the expression for  $\sqrt{\frac{\pi}{2}} \frac{1}{1 - \beta^2} K_\beta$  in the above proposition if  $\beta = \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}$ .

## 4.2 Another estimator

Let us consider the estimator  $L_{T,N}^r$  in (4.2).

In the case of OBM a proof that  $L_{T,N,T}^r(Y) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} L_T^r(Y)$  can be found in [20, Lemma 1].

Applying Proposition 2 we obtain the convergence in a stronger sense:  $L_{T,N}^r(Y) \xrightarrow[N \rightarrow \infty]{\text{u.c.p.}} L^r(Y)$ . And applying Theorem 2 we obtain the convergence rate.

**Proposition 7.** *Let  $Y$  be the OBM solution to (2.4) and let  $L_{T,N}^r(Y)$  be the estimator of the local time  $L^r(Y)$  in (4.2). Then there exists a Brownian motion  $B$  independent of  $Y$  (possibly on an extension of the probability space) such that*

$$N^{1/4}(L_{T,N}^r(Y) - L^r(Y)) \xrightarrow[N \rightarrow \infty]{\mathcal{L}\text{-}s} \sqrt{\frac{16}{3\sqrt{2\pi}} \frac{\sigma_-^2 + \sigma_+^2}{\sigma_- + \sigma_+}} B_{L^r(Y)}.$$

We obtain a simple formula because  $\mathfrak{O}_{\sigma, 2h_1}$  (whose expression is given by (2.9)) is zero, since  $\mathbf{H}_{2h_1} - \mathbf{H}_g = 0$ , and two other terms in equation (2.8) cancel because

$$\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \int_{-\infty}^{\infty} \int_0^1 \int_{-\infty}^{\infty} \frac{|x| e^{-\frac{x^2}{2}} \Phi(-|y|)}{\sqrt{2\pi} \sigma(x) \sigma(y)} \sqrt{\frac{1}{t} - 1} 2h_1(\sigma(x)x\sqrt{t}, \sigma(y)y\sqrt{1-t}) dy dt dx = \frac{4}{3\sqrt{2\pi}}.$$

The following proposition specifies Theorem 1 in case of the estimator (4.2) for SBM.

**Proposition 8.** *Let  $X$  be the solution to (2.1) and  $L_{T,N}^r(X)$  be the estimator in (4.2). There exists a Brownian motion  $B$  independent of  $X$  (possibly on an extension of the probability space) such that  $L_{T,N}^r(X) \xrightarrow[N \rightarrow \infty]{\text{u.c.p.}} (1 - \beta^2)L^r(X)$  and*

$$N^{1/4}(L_{T,N}^r(X) - (1 - \beta^2)L^r(X)) \xrightarrow[N \rightarrow \infty]{\mathcal{L}\text{-}s} \sqrt{K_\beta} B_{L^r(X)}$$

where

$$\frac{1}{1 - \beta^2} K_\beta = \frac{16}{3\sqrt{2\pi}} - 4\beta \int_{-\infty}^{\infty} x \Phi(-|x|) \mathcal{P}_\beta(x) dx + \frac{4}{\sqrt{2\pi}} (1 - \beta)\beta,$$

$\mathcal{P}_\beta(x) = \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} p_\beta(j, x, y) \mathbf{G}^{(\beta)}(y) dy$ , and

$$\mathbf{G}^{(\beta)}(y) = \text{sgn}(y)(1 - \text{sgn}(y)\beta)\beta \frac{1}{\sqrt{2\pi}} \left( e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y|\Phi(-|y|) \right).$$

*Proof.* Theorem 1, applied to  $h = 2h_1$ , yields the result with

$$\begin{aligned} \frac{1}{1 - \beta^2} K_\beta &= \frac{16}{3\sqrt{2\pi}} + 4 \int_{-\infty}^{\infty} |x| \Phi(-|x|) \mathcal{P}_\beta(x) dx \\ &\quad - \frac{4}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (1 + \text{sgn}(x)\beta) (e^{-\frac{x^2}{2}} - \sqrt{2\pi}|x|\Phi(-|x|)) \mathcal{P}_\beta(x) dx. \end{aligned}$$

$K_\beta$  has this expression because two terms cancel. Moreover we now sum up the two integrals to obtain

$$\frac{1}{1 - \beta^2} K_\beta = \frac{16}{3\sqrt{2\pi}} - 4\beta \int_{-\infty}^{\infty} x \Phi(-|x|) \mathcal{P}_\beta(x) dx - 4 \int_{-\infty}^{\infty} p_\beta(1, 0, x) \mathcal{P}_\beta(x) dx$$

and the Chapman-Kolmogorov equation ensures that

$$\int_{-\infty}^{\infty} p_\beta(1, 0, x) \mathcal{P}_\beta(x) dx = \mathcal{P}_\beta(0) - \frac{1}{\sqrt{2\pi}} (1 - \beta)\beta.$$

Note that  $\mathcal{P}_\beta(0)$  can be computed explicitly and it is 0. □

## Appendix A Properties of oscillating Brownian motion

In this section we consider  $Y$  to be an OBM with threshold  $r = 0$ .

## A.1 Scaling property

In this section  $Y^{Y_0}$  denotes the OBM with threshold  $r = 0$  starting from a deterministic point  $Y_0$ , let  $c \in (0, \infty)$ . Let us mention the following well known diffusive scaling properties for OBM:

$$\left( \frac{1}{\sqrt{c}} Y_{ct}^{Y_0} \right)_{t \geq 0} \stackrel{\text{law}}{=} (Y_t^{Y_0/\sqrt{c}})_{t \geq 0}$$

(i.e. “the rescaled OBM is still a OBM with rescaled starting point”) and

$$\left( \frac{1}{\sqrt{c}} Y_{ct}^{\sqrt{c}Y_0}, \frac{1}{\sqrt{c}} L_{ct}(Y^{\sqrt{c}Y_0}) \right)_{t \geq 0} \stackrel{\text{law}}{=} (Y_t^{Y_0}, L_t(Y^{Y_0}))_{t \geq 0}. \quad (\text{A.1})$$

## A.2 The joint density of a standard OBM and its local time

The joint density of a standard OBM and its local time at time  $t$ ,  $\rho_t^\sigma(y, \ell)$  coincides with

$$\rho_t^\sigma(y, \ell) = \frac{1}{(\sigma(y))^2} \rho_t \left( \frac{y}{\sigma(y)}, \frac{\sigma_- + \sigma_+}{2\sigma_- \sigma_+} \ell \right) \quad (\text{A.2})$$

for  $y \neq 0$ , where  $\rho$  is the joint density of the BM and its local time at time  $t$ :

$$\rho_t(y, \ell) = \frac{|y| + \ell}{\sqrt{2\pi t^3}} \exp\left(-\frac{(|y| + \ell)^2}{2t}\right) \mathbf{1}_{(0, \infty)}(\ell). \quad (\text{A.3})$$

In particular  $\rho_t^\sigma(y, \ell) dy d\ell = \rho_t \left( \frac{y}{\sigma(y)}, \frac{\sigma_- + \sigma_+}{2\sigma_- \sigma_+} \ell \right) \lambda_\sigma(dy) d\ell$ .

## A.3 Orthogonal martingales

In this section we show another easy fact that the OBM has in common with the one-dimensional Brownian motion: the only orthogonal square-integrable martingales are the constants.

**Lemma 1.** *Let  $(M_t)_{t \in [0,1]}$  be a square-integrable  $(\mathcal{F}_t)_{t \in [0,1]}$ -martingale such that for all  $t \in [0, 1]$  the cross variation satisfies  $\mathbb{P}(\langle M_t, Y_t \rangle_t = 0) = 1$ . Then  $M$  is constant.*

*Proof.* Without loss of generality we can assume  $M_0 = 0$ , otherwise consider  $M_t - M_0$ . There exists an  $(\mathcal{F}_t)_{t \in [0,1]}$ -progressively measurable process  $\nu$  such that  $\mathbb{E} \left[ \int_0^1 \nu_s^2 ds \right] < \infty$  and  $M_1 = \int_0^1 \nu_s dW_s$  (cf. [15, Problem 4.17 in Chapter 3]). This and the fact that  $M$  is a martingale implies that  $M_t = \mathbb{E}[M_1 | \mathcal{F}_t] = \int_0^t \nu_s dW_s$  for all  $t \in [0, 1]$ . The orthogonality of  $M$  to  $Y$  rewrites as follows for all  $t \in [0, 1]$  it holds  $\mathbb{P}(\int_0^t \nu_s \sigma(Y_s) ds = 0) = 1$ . By continuity we derive that  $\mathbb{P}$ -a.s. for all  $t \in [0, 1]$  it holds that  $\int_0^t \nu_s \sigma(Y_s) ds = 0$ . This together with the fact that  $\sigma$  is positive ensures that  $\mathbb{P}$ -a.s.  $\int_0^1 \mathbf{1}_{\{\nu_s \neq 0\}} ds = 0$ . Hence for all  $t \in [0, 1]$  it holds  $\mathbb{P}$ -a.s. that  $M_t = \int_0^t \nu_s dW_s = 0$ .  $\square$

## A.4 Bounds for the semigroup

For a measurable, bounded function  $f: \mathbb{R} \rightarrow \mathbb{R}$  set

$$Q_t^\sigma f(x) := \int_{-\infty}^{\infty} q_\sigma(t, x, y) f(y) dy \quad \text{and} \quad P_t^{\beta\sigma} f(x) := \int_{-\infty}^{\infty} p_{\beta\sigma}(t, x, y) f(y) dy \quad (\text{A.4})$$

for all  $t \in [0, \infty)$ . They are respectively the semigroup of the standard OBM and of the standard  $\beta_\sigma$ -SBM with skewness parameter  $\beta_\sigma := \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}$  and they satisfy  $Q_t^\sigma f(x) = P_t^{\beta_\sigma} f_\sigma(x/\sigma(x))$ . Note that  $P_t := P_t^0 = Q_t^1$  is the semigroup of the Brownian motion and

$$Q_t^\sigma f(x) = P_t f_\sigma(x/\sigma(x)) + \beta_\sigma P_t(f_\sigma \mathbf{1}_{[0, \infty)})(-|x|/\sigma(x)) - \beta_\sigma P_t(f_\sigma \mathbf{1}_{(-\infty, 0)})(|x|/\sigma(x)) \quad (\text{A.5})$$

where  $f_\sigma(x) = f(\sigma(x)x)$ . From this relationship between the semigroups of OBM and Brownian motion we derive the following properties.

**Lemma 2.** *Let  $f \in \mathbf{L}^{1,b}(\lambda^{(2)})$ , and let us denote by  $p(t, \cdot)$  is the density of a Gaussian random variable with variance  $t$ . Then there exists a positive constant  $K \in (0, \infty)$  such that for all  $x, y \in \mathbb{R}$ ,  $0 \leq s \leq t$  it holds that*

$$i) \quad |Q_t^\sigma f(x)| \leq \frac{1}{\min\{\sigma_-, \sigma_+\}} \left(1 + \frac{|\sigma_- - \sigma_+|}{\sigma_- + \sigma_+}\right) \frac{1}{\sqrt{2\pi}} \frac{\|f\|_1}{\sqrt{t}},$$

$$ii) \quad \left| Q_t^\sigma f(x) - \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \langle \lambda_\sigma, f \rangle p(t, x/\sigma(x)) \right| \leq \frac{K}{\sqrt{t^3}} (\|f\|_{1,2} + \|f\|_{1,1}|x|),$$

iii) *for all  $\zeta \geq 0$  there exists a positive constant  $K_\zeta$  such that*

$$\left| Q_t^\sigma f(x) - \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \langle \lambda_\sigma, f \rangle p(t, x/\sigma(x)) \right| \leq \left(1 + \frac{|\sigma_- - \sigma_+|}{\sigma_- + \sigma_+}\right) \frac{K_\zeta}{t} \left( \frac{\|f\|_{1,1}}{1+(|x|/(\sigma(x)\sqrt{t}))^\zeta} + \frac{\|f\|_{1,1+\zeta}}{1+(|x|/\sigma(x))^\zeta} \right),$$

$$iv) \quad |Q_t^\sigma f(x) - Q_t^\sigma f(y)| \leq \frac{1}{\min\{\sigma_-, \sigma_+\}} \left(1 + \frac{|\sigma_- - \sigma_+|}{\sigma_- + \sigma_+}\right) K \frac{|x-y|}{t} \|f\|_1, \text{ and}$$

$$v) \quad |Q_t^\sigma f(x) - Q_s^\sigma f(x)| \leq \frac{1}{\min\{\sigma_-, \sigma_+\}} \left(1 + \frac{|\sigma_- - \sigma_+|}{\sigma_- + \sigma_+}\right) K \frac{t-s}{\sqrt{s^3}} \|f\|_1.$$

*Proof.* Item (i) is a straightforward consequence of (2.6) and of the fact that

$$p_{\beta_\sigma}(t, x, y) \leq (1 + |\beta_\sigma|) p(t, x - y) \leq \frac{1 + |\beta_\sigma|}{\sqrt{2\pi t}}.$$

To prove the other items we also use the fact for all  $\alpha \geq 0$   $\|f(\sigma(\cdot)\cdot)\|_{1,\alpha} \leq \frac{\|f\|_{1,\alpha}}{(\min\{\sigma_-, \sigma_+\})^{1+\alpha}}$  and  $\|f \mathbf{1}_{[0, \infty)}\|_{1,\alpha} + \|f \mathbf{1}_{(-\infty, 0)}\|_{1,\alpha} = \|f\|_{1,\alpha}$ . Item (ii) follows from (A.5) and [22, Lemma 1] for SBM. Item (iii) follows from (A.5) and the analogous result for Brownian motion: equation (3.2) in [13, Lemma 3.1]. Item (iv) and Item (v) follow from (A.5) and equations (3.4)-(3.5) in [13, Lemma 3.1].  $\square$

The proof of the following lemma follows from Lemma 2 and it is analogous to the one of [13, Lemma 3.3]. It is therefore omitted.

**Lemma 3.** *Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  and*

$$\Gamma_t(n, f) := \sum_{k=1}^{\lfloor nt \rfloor - 1} Q_k^\sigma f(\sqrt{n} Y_0).$$

*Then there exists a positive constant  $K$  (depending on  $\sigma_\pm$ ) such that  $|\Gamma_t(n, f)| \leq K \|f\|_1 \sqrt{nt}$  and if  $\langle \lambda_\sigma, f \rangle = 0$  then*

$$|\Gamma_t(n, f)| \leq K (\|f\|_{1,2} + \|f\|_{1,1} |Y_0| \sqrt{n}) \quad \text{and} \quad |\Gamma_t(n, f)| \leq K \|f\|_{1,1} (1 + \log(nt)).$$

## A.5 Behavior of the local time

In this section we explore some properties of the local time of the OBM  $Y$  and its moments.

**Lemma 4.** *For all  $q \in (2, \infty)$ ,  $\alpha \in (0, \frac{q-2}{2q})$  it holds that (the pathwise continuous version of) the local time  $L(Y)$  is locally  $\alpha$ -Hölder continuous. In particular for all  $\delta \in (-\infty, \frac{1}{2})$ ,  $T \in [0, \infty)$  it holds that*

$$\sup_{t \in [0, T]} n^\delta \left( L_{t+\frac{1}{n}}(Y) - L_t(Y) \right) \xrightarrow[n \rightarrow \infty]{\text{a.s.}} 0.$$

This statement is not surprising since it is well known for the local time of Brownian motion. Although the proof exploits standard techniques, we provide it for the sake of completeness.

*Proof.* Without loss of generality we can assume  $\sigma_-, \sigma_+ \in (0, \infty)$ . In this proof let  $T \in [0, \infty)$  and  $s, t \in [0, T]$  with  $s \leq t$  be fixed. Let us first note that Itô-Tanaka formula implies that  $L_t - L_s = |Y_t| - |Y_s| - \int_s^t \text{sgn}(Y_u) \sigma(Y_u) dW_u$ . The fact that for all  $a, b \in \mathbb{R}$  it holds that  $(a+b)^q \leq 2^{q-1}(a^q + b^q)$  and that  $||a| - |b|| \leq |a - b|$  imply for all  $q \in [1, \infty)$  that

$$|L_t - L_s|^q \leq 2^{q-1} \left| \int_s^t \sigma(Y_u) dW_u \right|^q + 2^{q-1} \left| \int_s^t \text{sgn}(Y_u) \sigma(Y_u) dW_u \right|^q.$$

By Burkholder-Davis-Gundy inequality for all  $q \geq 2$  there exists a constant  $K_q > 0$  such that

$$\mathbb{E}[|L_t - L_s|^q] \leq K_q \mathbb{E} \left[ \left| \int_s^t \sigma^2(Y_u) du \right|^{\frac{q}{2}} \right] \leq \max\{\sigma_-, \sigma_+\}^q K_q (t-s)^{q/2}.$$

Finally Kolmogorov continuity theorem ensures that there exists a continuous version of the local time (that we took already) and it is locally Hölder continuous as required.  $\square$

**Lemma 5.** *Let  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$  be the real function satisfying  $g(x, y) = |y| - |x|$ . Then for all  $t \in [0, \infty)$  it holds that  $\langle \lambda_\sigma, \mathbf{H}_g \rangle = 1$ ,*

$$\frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \mathbf{H}_g(\sqrt{n} Y_{k/n}) = \sum_{k=0}^{\lfloor nt \rfloor - 1} \mathbb{E}[L_{(k+1)/n}(Y) - L_{k/n}(Y) | \mathcal{F}_{k/n}]$$

$$\text{and } \sup_{s \in [0, t]} \left| \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor ns \rfloor - 1} \mathbf{H}_g(\sqrt{n} Y_{k/n}) - L_s(Y) \right| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

*Proof.* Some computations, that we decide to omit, show that  $\langle \lambda_\sigma, \mathbf{H}_g \rangle = 1$ .

Next, let us observe that a simple change of variable (corresponding to the scaling property A.1) and the Markov property yield

$$\begin{aligned} \frac{1}{\sqrt{n}} \mathbf{H}_g(\sqrt{n} Y_{\frac{k}{n}}) &= \frac{1}{\sqrt{n}} \int_{-\infty}^{\infty} (|y| - \sqrt{n} |Y_{\frac{k}{n}}|) q_\sigma(1, \sqrt{n} Y_{\frac{k}{n}}, y) dy \\ &= \int_{-\infty}^{\infty} (|y| - |Y_{\frac{k}{n}}|) q_\sigma\left(\frac{1}{n}, Y_{\frac{k}{n}}, y\right) dy = \mathbb{E}\left[|Y_{\frac{k+1}{n}}| - |Y_{\frac{k}{n}}| \mid |Y_{\frac{k}{n}}|\right] = \mathbb{E}\left[|Y_{\frac{k+1}{n}}| - |Y_{\frac{k}{n}}| \mid \mathcal{F}_{\frac{k}{n}}\right]. \end{aligned}$$

This and Itô-Tanaka formula show  $\frac{1}{\sqrt{n}} \mathbf{H}_g(\sqrt{n} Y_{k/n}) = \mathbb{E}[L_{(k+1)/n}(Y) - L_{k/n}(Y) | \mathcal{F}_{k/n}]$ . Lemma 2.14 in [11] ensures the desired convergence in probability.  $\square$

In the remainder of this section  $Y^x$ ,  $x \in \mathbb{R}$ , denotes the OBM with threshold  $r = 0$  starting from  $Y_0 = x$ . For every  $p \in [0, \infty)$ ,  $x \in \mathbb{R}$ , and function  $f: \mathbb{R} \rightarrow \mathbb{R}$  either non-negative or such that  $(L_1(Y^x))^p f(Y_1^x) \in \mathbf{L}^1(\mathbb{P})$  let

$$\mathbb{L}^{(p)}(f, x) := \mathbb{E}[(L_1(Y^x))^p f(Y_1^x)]. \tag{A.6}$$

In this document we only consider functions  $f: \mathbb{R} \rightarrow \mathbb{R}$  satisfying that there exist  $K, \alpha \in [0, \infty)$  such that  $|f(x)| \leq Ke^{\alpha|x|}$  for all  $x \in \mathbb{R}$ , so  $\mathbb{L}^{(\cdot)}(f, \cdot)$  is well defined. The scaling property (A.1) in Appendix A.1 implies that

$$\mathbb{L}^{(p)}(f, \sqrt{n}Y_{\frac{(k-1)}{n}}^{Y_0}) = n^{\frac{p}{2}} \mathbb{E} \left[ \left( L_{\frac{k}{n}}(Y^{Y_0}) - L_{\frac{k-1}{n}}(Y^{Y_0}) \right)^p f(\sqrt{n}Y_{\frac{k}{n}}^{Y_0}) | \mathcal{F}_{\frac{(k-1)}{n}} \right]. \quad (\text{A.7})$$

In particular note that  $\mathbb{L}^{(1)}(1, \cdot) = \mathbf{H}_g(\cdot)$  with  $g(x, y) = |y| - |x|$ .

**Lemma 6.** *Let  $x \in \mathbb{R}$ , let  $W$  be a Brownian motion with  $W_0 = 0$ , let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function satisfying that there exist  $K, \alpha \in [0, \infty)$  such that  $|f(y)| \leq Ke^{\alpha|y|}$  for all  $y \in \mathbb{R}$ . Then for all  $p \in \mathbb{N}$  it hold that*

$$\begin{aligned} \mathbb{L}^{(p)}(f, x) &= \int_0^1 \frac{|x|}{\sigma(x)} \frac{(1-t)^{\frac{p}{2}}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} \mathbb{E} \left[ (L_1(Y^0))^p f(Y_1^0 \sqrt{1-t}) \right] dt \\ &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^{(p+1)} \frac{|x|}{\sigma(x)} \int_0^1 \frac{(1-t)^{\frac{p}{2}}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} \mathbb{E} \left[ (L_1(W))^p (\sigma(W_1))^{-1} f(\sigma(W_1)W_1 \sqrt{1-t}) \right] dt, \end{aligned}$$

if  $x \neq 0$ , and  $\mathbb{L}^{(p)}(f, 0) = \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^{(p+1)} \mathbb{E} \left[ (L_1(W))^p (\sigma(W_1))^{-1} f(\sigma(W_1)W_1) \right]$ .

(If  $\sigma_{\pm} = \infty$  then replace  $f$  in the right hand side of last two equalities with  $f \mathbf{1}_{\mathbb{R}_{\mp}}$ .)

*Proof.* We reduce to consider the case  $x \neq 0$  because if  $x = 0$  then the statement follows from simple computations using the joint density of the OBM  $Y$  and its local time (A.2).

Let  $X$  be the  $\beta_{\sigma}$ -SBM satisfying (3.1) (recall that  $\beta_{\sigma} = \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+}$ ). In particular  $X_0 = x/\sigma(x)$ . Let  $B$  be a Brownian motion starting at  $x/\sigma(x)$  and let  $X_t^0$  a standard  $\beta_{\sigma}$ -SBM independent of  $B$ . For a process  $\xi$  let us denote by  $T_0(\xi) := \inf(\{\infty\} \cup \{t \geq 0: \xi_t = 0\})$  the first time it hits 0.

One well known property of SBM is that the process behaves as a BM until it reaches the barrier, which is 0. This means that  $T_0(X) \stackrel{\text{law}}{=} T_0(B)$ . After reaching the threshold, by the Markov property, any  $\beta_{\sigma}$ -SBM behaves as a  $\beta_{\sigma}$ -SBM starting at the threshold. This means that conditioned on  $T_0(X)$  it holds that  $X_{t+T_0(X)}$  is distributed as  $X_t^0$ .

This and the relationship between the local times of OBM and SBM (3.2) show that

$$\begin{aligned} \mathbb{L}^{(p)}(f, x) &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^p \mathbb{E} \left[ \mathbf{1}_{\{T_0(X) \leq 1\}} \mathbb{E} \left[ (L_1(X))^p f(\sigma(X_1)X_1) | T_0(X) \right] \right] \\ &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^p \mathbb{E} \left[ \mathbf{1}_{\{T_0(B) \leq 1\}} (L_{1-T_0(B)}(X^0))^p f(\sigma(X_{1-T_0(B)}^0)X_{1-T_0(B)}^0) \right] \end{aligned}$$

Let us recall the well known fact that the random variable  $T_0(B)$  has density w.r.t. the Lebesgue measure given by  $(0, \infty) \ni t \mapsto \frac{|x|}{\sigma(x)} \frac{1}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}}$ . Then the relationship between the local times of OBM and associated SBM (3.2) ensures

$$\begin{aligned} \mathbb{L}^{(p)}(f, x) &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^p \int_0^1 \frac{|x|}{\sigma(x)} \frac{1}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} \mathbb{E} \left[ (L_{1-t}(X^0))^p f(\sigma(X_{1-t}^0)X_{1-t}^0) \right] dt \\ &= \int_0^1 \frac{|x|}{\sigma(x)} \frac{1}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} \mathbb{E} \left[ (L_{1-t}(Y^0))^p f(Y_{1-t}^0) \right] dt. \end{aligned}$$

The scaling property (A.1) and simple changes of variables imply that

$$\mathbb{L}^{(p)}(f, x) = \int_0^1 \frac{|x|}{\sigma(x)} \frac{(1-t)^{\frac{p}{2}}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} \mathbb{E} \left[ (L_1(Y^0))^p f(Y_1^0 \sqrt{1-t}) \right] dt.$$






Figure 2: Map of the proof of Proposition 4.

In the appendix we introduce many auxiliary results. This map show how they intervene in the proof of Proposition 4 and of other results of the appendix.

The relationship between the joint density of the standard OBM and its local time (A.2) and the one for Brownian motion and its local time (A.3) ensures that for all  $t \in [0, 1]$  it holds

$$\mathbb{E}[(L_1(Y^0))^p f(Y_1^0 \sqrt{1-t})] = \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^p \frac{2}{\sigma_- + \sigma_+} \mathbb{E}[(L_1(W))^p \sigma(-W_1) f(\sigma(W_1) W_1 \sqrt{1-t})].$$

which yields the conclusion.  $\square$

## Appendix B Proof of the key Proposition 4

In this section we prove Proposition 4 which was stated in Section 3.3. The section is organized as follows: We first introduce, in Sections B.1-B.2, some auxiliary results and functions. Then we split the proof of Proposition 4 into two parts. The first part, in Section B.3, consists in proving the decomposition (3.3) into a sum of a vanishing term and a martingale part. In the second part, in Section B.4, we demonstrate that the martingale part satisfies Items (i)-(iv) of Proposition 4. Figure 2 show how the results intervenes in the proof.

In this section,  $(Y_t)_{t \in [0,1]}$  is a standard OBM.

### B.1 Auxiliary convergence results

The following is the generalization to the case of OBM of Lemma 4.2 in [13]. The proof is analogous and therefore omitted.

**Lemma 7.** *Let  $(g_n)_{n \in \mathbb{N}}$  be a sequence of real functions satisfying that  $\langle \lambda_\sigma, g_n \rangle = 0$  and for all  $x \in \mathbb{R}$*

$$\lim_{n \rightarrow \infty} \frac{g_n(x\sqrt{n})^2}{n} + \frac{\|g_n^2\|_1}{\sqrt{n}} + \frac{\|g_n\|_{1,1} |g_n(x\sqrt{n})| \log(n)}{n} + \frac{\|g_n\|_{1,1} \|g_n\|_1 \log(n)}{\sqrt{n}} = 0. \quad (\text{B.1})$$

Then  $\lim_{n \rightarrow \infty} \mathbb{E} \left[ \left( \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} g_n(\sqrt{n} Y_{\frac{k}{n}}) \right)^2 \right] = 0$  for all  $t \in [0, 1]$ .

The following propositions corresponds to Theorem 4.1 a) and b) in [13]. The proof of the first is step by step an adaptation to OBM of the proof of Theorem 4.1.a) for Brownian motion and it relies on Lemma 3. The proof of Proposition 9 is therefore omitted.

**Proposition 9.** *Let  $g_n: \mathbb{R} \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}$ , be a sequence of functions satisfying  $\lim_{n \rightarrow \infty} \|g_n\|_1 = 0$  and for all  $x \in \mathbb{R}$  it holds that  $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} g_n(\sqrt{n}x) = 0$ . Then*

$$\lim_{n \rightarrow \infty} \sup_{s \in [0,1]} \mathbb{E} \left[ \left| n^{-\frac{1}{2}} \sum_{k=0}^{\lfloor ns \rfloor - 1} g_n(\sqrt{n} Y_{\frac{k}{n}}) \right| \right] = 0.$$

**Proposition 10.** *Let  $g_n: \mathbb{R} \rightarrow \mathbb{R}$ ,  $n \in \mathbb{N}$ , be a sequence of functions satisfying (B.1) and there exists  $\lambda \in \mathbb{R}$  such that  $\lim_{n \rightarrow \infty} \langle \lambda_\sigma, g_n \rangle = \lambda$ . Then for all  $t \in [0, 1]$  it holds that*

$$\frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} g_n(\sqrt{n} Y_{\frac{k}{n}}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \lambda L_t(Y).$$

If in addition  $\sup_{n \in \mathbb{N}} \|g_n\|_1 < \infty$  then

$$\sup_{s \in [0,1]} \left| n^{-\frac{1}{2}} \sum_{k=0}^{\lfloor ns \rfloor - 1} g_n(\sqrt{n} Y_{\frac{k}{n}}) - \lambda L_s(Y) \right| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

*Proof.* Let us set the sequence  $f_n := g_n - \langle \lambda_\sigma, g_n \rangle \mathbf{H}_g$  with  $g(x, y) := |y| - |x|$ . Note that Lemma 5 ensures that  $\langle \lambda_\sigma, \mathbf{H}_g \rangle = 1$  and that  $\frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \mathbf{H}_g(\sqrt{n} Y_{\frac{k}{n}}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} L_t(Y)$ . Hence  $\langle \lambda_\sigma, f_n \rangle = 0$  and one can easily show that  $f_n$  satisfies (B.1). Lemma 7 yields the result. The additional statemets is the same as [13, Theorem 4.1].  $\square$

*Remark 3* (Proposition 9 and 10 for a constant sequence of functions). Let  $f \in \mathbf{L}^1$  such that for all  $x \in \mathbb{R}$  it holds that  $\lim_{n \rightarrow \infty} \frac{f(\sqrt{n}x)}{\sqrt{n}} = 0$  (e.g.  $f \in \mathbf{L}^{1,b}(\lambda^{(0)})$ ). Then Proposition 9 states that if  $\|f\|_1 = 0$  then

$$\lim_{n \rightarrow \infty} \sup_{s \in [0,1]} \mathbb{E} \left[ \left| n^{-\frac{1}{2}} \sum_{k=0}^{\lfloor ns \rfloor - 1} f(\sqrt{n} Y_{\frac{k}{n}}) \right| \right] = 0.$$

And Proposition 10 states that if  $f^2 \in \mathbf{L}^1$  and  $f \in \mathbf{L}^1(\lambda^{(1)})$  (e.g.  $f \in \mathbf{L}^{1,b}(\lambda^{(1)})$ ) then

$$\sup_{s \in [0,1]} \left| n^{-\frac{1}{2}} \sum_{k=0}^{\lfloor ns \rfloor - 1} f(\sqrt{n} Y_{\frac{k}{n}}) - \langle \lambda_\sigma, f \rangle L_s(Y) \right| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

## B.2 Auxiliary functions

**Lemma 8.** *Let  $\gamma \in (0, \infty)$ ,  $h \in \mathbf{I}_\gamma$ , and let  $\mathbf{G}_h$  be the function*

$$\mathbf{G}_h := \mathbf{H}_h - \langle \lambda_\sigma, \mathbf{H}_h \rangle \mathbf{H}_g \tag{B.2}$$

*with  $g(x, y) = |y| - |x|$ . Then  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = 0$  and  $\mathbf{G}_h \in \mathbf{I}_\gamma$ .*

*Proof.* Throughout this proof let  $K_\sigma = \frac{1}{\min\{\sigma_-^2, \sigma_+^2\}} \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \in (0, \infty)$ . First note that the fact that  $q_\sigma(1, x, y) \leq K_\sigma \frac{1}{\sqrt{2\pi}} e^{-\frac{(x-y)^2}{2}}$  implies that  $|\mathbf{H}_g(x)| \leq K_\sigma \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} |y-x| e^{-\frac{(x-y)^2}{2}} dy \in \mathbf{I}_\alpha$  for all  $\alpha \geq 0$ . And it also implies, together with the fact that  $h \in \mathbf{I}_\gamma$ , that

$$|\mathbf{H}_h(x)| \leq K_\sigma \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} |h(x, y)| e^{-\frac{(x-y)^2}{2}} dy \leq K_\sigma \bar{h}(x) \frac{e^{-\frac{a^2}{2}}}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(|x-y|-a)^2}{2}} dy \leq K_\sigma \bar{h}(x) e^{-\frac{a^2}{2}}$$

with  $\bar{h} \in \mathbf{L}^{1,b}(\lambda^{(\gamma)})$  positive function and  $a$  non negative constant. Hence it holds that  $\mathbf{H}_h \in \mathbf{L}^{1,b}(\lambda^{(\gamma)})$ . In particular it holds  $\langle \lambda_\sigma, \mathbf{H}_h \rangle \leq \|\mathbf{H}_h\|_1 < \infty$ . Therefore  $|\mathbf{G}_h| \leq |\mathbf{H}_h| + \|\mathbf{H}_h\|_1 |\mathbf{H}_g| \in \mathbf{I}_\gamma$  and so  $\mathbf{G}_h \in \mathbf{I}_\gamma$ .

It remains to prove that  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = 0$ . This follows from the fact that  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = \langle \lambda_\sigma, \mathbf{H}_h \rangle (1 - \langle \lambda_\sigma, \mathbf{H}_g \rangle)$  and  $\langle \lambda_\sigma, \mathbf{H}_g \rangle = 1$  by Lemma 5.  $\square$

In the reminder of this section let  $\gamma \in [1, \infty)$ ,  $h \in \mathbf{I}_\gamma$ , let  $\mathbf{G}_h$  be the function in (B.2), and for all  $i, j \in \{0, 1\}$ ,  $\eta \in [0, \infty)$ ,  $n \in [1, \infty)$  let

$$\mathcal{Q}_{n,i,j}^{(\eta)} := \sum_{k=i}^{\lfloor n^\eta \rfloor + j} Q_k^\sigma \mathbf{G}_h \quad \text{and} \quad \mathcal{Q}_{n,i,j} := \mathcal{Q}_{n,i,j}^{(\frac{1}{4})} \quad (\text{B.3})$$

where  $Q^\sigma$  is the semigroup of the OBM given in (A.4). Let  $n \in \mathbb{N}$  be fixed.

The following facts are consequences of Lemma 8, Lemma 2, and the fact that  $\sum_{j=1}^n \frac{1}{j} \leq 2 \log(n)$  for  $n \geq 2$ .

For every  $\zeta \in [0, \gamma - 1]$ , for every  $\eta \in (0, 1)$  the fact that  $\lambda(\mathbf{G}_h) = 0$  and Item (iii) in Lemma 2 imply that for all  $x \in \mathbb{R}$  it holds that

$$\begin{aligned} |Q_{n^\eta+1}^\sigma \mathbf{G}_h(x)| &= |Q_{n^\eta+1}^\sigma \mathbf{G}_h(x) - \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \langle \lambda_\sigma, \mathbf{G}_h \rangle p(n^{\frac{1}{4}} + 1, x/\sigma(x))| \\ &\leq \left(1 + \frac{|\sigma_- \sigma_+|}{\sigma_- + \sigma_+}\right) K_\zeta n^{-\eta} \left( \frac{1}{1+|xn^{-\eta/2}/\sigma(x)|^\zeta} + \frac{1}{1+(|x|/\sigma(x))^\zeta} \right) \end{aligned} \quad (\text{B.4})$$

and

$$|\mathcal{Q}_{n,1,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,1,1}^{(\eta)}(x)| \leq 2K_\zeta \log(n) \left( \frac{1}{1+|xn^{-\frac{1}{2}}/\sigma(x)|^\zeta} + \frac{1}{1+(|x|/\sigma(x))^\zeta} \right) \quad (\text{B.5})$$

for some  $K_\zeta \in (0, \infty)$  depending also on  $\eta$ . Hence (B.5) with  $\zeta = 0$  and Item (ii) in Lemma 2 imply that for all  $x \in \mathbb{R}$  it holds that

$$|\mathcal{Q}_{n,1,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,1,1}^{(\eta)}(x)| \leq 2K \min \{ \log(n), (1 + |x|) \}, \quad (\text{B.6})$$

for some constant  $K \in (0, \infty)$  depending on  $\eta \in (0, 1)$ . This and the fact that  $\mathbf{G}_h$  is bounded ensures that for some positive constant  $K \in (0, \infty)$  it holds that  $|\mathcal{Q}_{n,0,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,0,1}^{(\eta)}(x)| \leq 2|\mathbf{G}_h(x)| + |\mathcal{Q}_{n,1,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,1,1}^{(\eta)}(x)| \leq 2K(\log(n) + 1)$  and

$$|\mathcal{Q}_{n,0,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,0,1}^{(\eta)}(x)| + |\mathcal{Q}_{n,1,0}^{(\eta)}(x)| + |\mathcal{Q}_{n,1,1}^{(\eta)}(x)| \leq 2K(\log(n) + 1). \quad (\text{B.7})$$

**Lemma 9.** *Pointwise  $\lim_{m \rightarrow \infty} \mathcal{Q}_{m,0,0} = \mathfrak{O}_{\sigma,h}$ . (Recall that  $\mathfrak{O}_{\sigma,h}$  is given by (2.9).)*

*Proof.* Straightforward consequence of Item (ii) in Lemma 2 and of the fact that  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = 0$  (see Lemma 8).  $\square$

### B.3 The decomposition as sum martingale and vanishing terms

In this section let  $\gamma$  be an arbitrary non-negative number to be specified in each statement and let  $h \in \mathbf{I}_\gamma$ . We are now determining the terms of the decomposition in equation (3.3).

For every  $n \in \mathbb{N}$  let  $M^{n,1}$ ,  $M^{n,2}$ , and  $N^n$  be the processes satisfying for all  $t \in [0, 1]$  that

$$\begin{aligned} M_t^{n,1} &= \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \left( h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}Y_{\frac{k+1}{n}}) - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) + \langle \lambda_\sigma, \mathbf{H}_h \rangle \mathbf{H}_g(\sqrt{n}Y_{\frac{k}{n}}) \right) - L_{\frac{\lfloor nt \rfloor}{n}}(Y), \\ M_t^{n,2} &= \frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} \left( \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{\frac{k}{n}}) - \mathcal{Q}_{n,1,1}(\sqrt{n}Y_{\frac{k-1}{n}}) \right), \text{ and } N_t^n = \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \mathbf{G}_h(\sqrt{n}Y_{\frac{k}{n}}) \end{aligned} \quad (\text{B.8})$$

(recall the definition for  $\mathbf{G}_h$  in (B.2),  $\mathcal{Q}$  in (B.3), and  $g(x, y) = |y| - |x|$ ). Trivially it holds for all  $n \in \mathbb{N}$ ,  $t \in [0, 1]$  that

$$\begin{aligned} \varepsilon_{n,t}^{(0,h,Y)} - \langle \lambda_\sigma, \mathbf{H}_h \rangle L_t(Y) &= M_t^{n,1} + N_t^n + \langle \lambda_\sigma, \mathbf{H}_h \rangle (L_{\lfloor nt \rfloor / n}(Y) - L_t(Y)) \\ &= M_t^{n,1} + M_t^{n,2} - M_t^{n,2} + N_t^n + \langle \lambda_\sigma, \mathbf{H}_h \rangle (L_{\lfloor nt \rfloor / n}(Y) - L_t(Y)). \end{aligned}$$

The right-hand-side of the latter equality is a sum of an  $(\mathcal{F}_{\lfloor nt \rfloor / n})_{t \geq 0}$ -martingale (denoted by  $n^{-\frac{1}{4}}\mathcal{M}^n$ ) and of a vanishing term with rate of order at least  $1/4$  (denoted by  $\mathcal{V}^n$ ):

$$\begin{aligned} \mathcal{V}_t^n &:= N_t^n - M_t^{n,2} + \langle \lambda_\sigma, \mathbf{H}_h \rangle \left( L_{\frac{\lfloor nt \rfloor}{n}}(Y) - L_t(Y) \right) \quad \text{and} \\ \mathcal{M}_t^n &:= n^{\frac{1}{4}} (M_t^{n,1} + M_t^{n,2}). \end{aligned} \quad (\text{B.9})$$

Lemma 11 below, together with Lemma 4, ensures that  $\sup_{t \in [0,1]} n^{\frac{1}{4}} |\mathcal{V}_t^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ . The two lemmas below prove the martingale property of  $\mathcal{M}^n$ , for  $n \in \mathbb{N}$ .

**Lemma 10.**  *$M^{n,1}$  is a martingale with respect to the filtration  $(\mathcal{F}_{\lfloor nt \rfloor / n})_{t \in [0,1]}$ .*

*Proof.* Throughout this proof let  $A^n$  and  $B^n$  be the processes given by

$$A_t^n = \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \mathbf{H}_g(\sqrt{n}Y_{\frac{k}{n}}) - L_{\frac{\lfloor nt \rfloor}{n}}(Y) \text{ and } B_t^n = M_t^{n,1} - \langle \lambda_\sigma, \mathbf{H}_h \rangle A_t^n.$$

So  $B_t^n = \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor - 1} \left( h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}Y_{\frac{k+1}{n}}) - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) \right)$ . It suffices to show the martingale property for  $A^n$  and  $B^n$ . Let  $t \in [0, 1]$  be fixed. First observe that Lemma 5 ensures that

$$A_t^n = \sum_{k=0}^{\lfloor nt \rfloor - 1} \left( \mathbb{E}[L_{(k+1)/n}(Y) - L_{k/n}(Y) | \mathcal{F}_{k/n}] - (L_{(k+1)/n}(Y) - L_{k/n}(Y)) \right).$$

The martingale property for  $A^n$  is an immediate consequence of the tower property. Let us explicit the case of the process  $B^n$ . For all  $j \in \{0, \dots, \lfloor nt \rfloor - 1\}$  it holds that

$$\mathbb{E} \left[ n^{\frac{1}{2}} B_t^n | \mathcal{F}_{\frac{j}{n}} \right] = \sum_{k=j}^{\lfloor nt \rfloor - 1} \mathbb{E} \left[ h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}Y_{\frac{k+1}{n}}) - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) | \mathcal{F}_{\frac{j}{n}} \right] + B_{\frac{j}{n}}^n = B_{\frac{j}{n}}^n$$

because for all  $k \in \{j, \dots, \lfloor nt \rfloor - 1\}$  it holds that

$$\begin{aligned} \mathbb{E} \left[ h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}Y_{\frac{k+1}{n}}) - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) | \mathcal{F}_{\frac{j}{n}} \right] &= \mathbb{E} \left[ \mathbb{E} \left[ h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}Y_{\frac{k+1}{n}}) - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) | \mathcal{F}_{\frac{k}{n}} \right] | \mathcal{F}_{\frac{j}{n}} \right] \\ &= \mathbb{E} \left[ \int_{-\infty}^{\infty} h(\sqrt{n}Y_{\frac{k}{n}}, \sqrt{n}y) q_\sigma(1/n, \sqrt{n}Y_{\frac{k}{n}}, y) dy - \mathbf{H}_h(\sqrt{n}Y_{\frac{k}{n}}) | \mathcal{F}_{\frac{j}{n}} \right] = \mathbb{E} \left[ 0 | \mathcal{F}_{\frac{j}{n}} \right] = 0. \end{aligned}$$

□

**Lemma 11.** *Let  $h \in \mathbf{I}_3$ . Then  $M^{n,2}$  is a martingale with respect to the filtration  $(\mathcal{F}_{\lfloor nt \rfloor/n})_{t \in [0,1]}$  and it holds that  $\sup_{s \in [0,1]} n^{\frac{1}{4}} |N_s^n - W_s^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ .*

*Proof.* Using (B.3) we rewrite  $M^{n,2}$  in (B.8) as

$$\begin{aligned} M_t^{n,2} &= \frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} \sum_{j=0}^{\lfloor n^{\frac{1}{4}} \rfloor} (\mathbb{E}[\mathbf{G}_h(\sqrt{n}Y_{(j+k)/n}) | \mathcal{F}_{k/n}] - \mathbb{E}[\mathbf{G}_h(\sqrt{n}Y_{(j+k)/n}) | \mathcal{F}_{(k-1)/n}]) \\ &= N_t^n - \frac{1}{\sqrt{n}} (\mathcal{Q}_{n,0,0}(\sqrt{n}Y_0) - \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{\lfloor nt \rfloor/n})) - \frac{1}{\sqrt{n}} \sum_{j=0}^{\lfloor nt \rfloor - 1} Q_{\lfloor n^{\frac{1}{4}} \rfloor + 1}^\sigma \mathbf{G}_h(\sqrt{n}Y_{j/n}). \end{aligned} \quad (\text{B.10})$$

The first equality of the latter equation makes clear that that  $M^{n,2}$  is a martingale with respect to the filtration  $(\mathcal{F}_{\lfloor nt \rfloor/n})_{t \in [0,1]}$ .

Let  $m_t^n := N_t^n - n^{-\frac{1}{2}} (\mathcal{Q}_{n,0,0}^{(3/4)}(\sqrt{n}Y_0) - \mathcal{Q}_{n,0,0}^{(3/4)}(\sqrt{n}Y_{\lfloor nt \rfloor/n})) - n^{-\frac{1}{2}} \sum_{j=0}^{\lfloor nt \rfloor - 1} Q_{\lfloor n^{\frac{3}{4}} \rfloor + 1}^\sigma \mathbf{G}_h(\sqrt{n}Y_{j/n})$ . As

for (B.10) it is clear that  $m^n$  is a martingale with respect to the filtration  $(\mathcal{F}_{\lfloor nt \rfloor/n})_{t \in [0,1]}$ . Therefore  $n^{\frac{1}{4}}(M_t^{n,2} - m_t^n)$  as well. Let us denote by  $D_t^n := n^{\frac{1}{4}}(N_t^n - M_t^{n,2})$  and  $d_t^n := n^{\frac{1}{4}}(N_t^n - m_t^n)$ . Then  $D^n - d^n = n^{\frac{1}{4}}(w^n - M^{n,2})$  is a martingale with respect to the filtration  $(\mathcal{F}_{\lfloor nt \rfloor/n})_{t \in [0,1]}$ . In this notation, the statement we are proving rewrites:  $\sup_{s \in [0,1]} |D_s^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ .

*First step:* For every  $t \in [0,1]$ , let us show that  $\lim_{n \rightarrow \infty} \mathbb{E}[(D_t^n - d_t^n)^2] = 0$  by demonstrating the stronger fact that  $\lim_{n \rightarrow \infty} \mathbb{E}[(D_t^n)^2 + (d_t^n)^2] = 0$ .

Let  $t \in [0,1]$  and  $\eta \in \{\frac{1}{4}, \frac{3}{4}\}$  be fixed. Note that inequality (B.7) implies

$$\sup_{\omega \in \Omega} \sup_{s \in [0,1]} n^{-\frac{1}{4}} \left( \mathcal{Q}_{n,0,0}^{(\eta)}(\sqrt{n}Y_0(\omega)) - \mathcal{Q}_{n,0,0}^{(\eta)}(\sqrt{n}Y_{\lfloor ns \rfloor/n}(\omega)) \right) \leq n^{-\frac{1}{4}} 4K(\log(n) + 1) \xrightarrow[n \rightarrow \infty]{} 0, \quad (\text{B.11})$$

hence it holds also that  $\lim_{n \rightarrow \infty} \mathbb{E} \left[ n^{-\frac{1}{2}} \left( \mathcal{Q}_{n,0,0}^{(\eta)}(\sqrt{n}Y_0) - \mathcal{Q}_{n,0,0}^{(\eta)}(\sqrt{n}Y_{\lfloor nt \rfloor/n}) \right)^2 \right] = 0$ . Next observe that, since  $\lambda_\sigma$  is the stationary measure, the sequences of functions  $g_n^{(\eta)} := n^{\frac{1}{4}} Q_{\lfloor n^\eta \rfloor + 1}^\sigma \mathbf{G}_h$ ,  $\eta \in \{\frac{1}{4}, \frac{3}{4}\}$  satisfy that  $\langle \lambda_\sigma, g_n^{(\eta)} \rangle = n^{\frac{1}{4}} \langle \lambda_\sigma, \mathbf{G}_h \rangle$  which is equal to 0 by Lemma 8. This and inequality (B.4) (with  $\zeta = 2$  since  $h \in \mathbf{I}_{\zeta+1}$ ) ensure that (B.1) holds. Hence, applying Lemma 7 shows that  $\lim_{n \rightarrow \infty} \mathbb{E} \left[ n^{-\frac{1}{2}} \left( \sum_{j=0}^{\lfloor nt \rfloor - 1} Q_{\lfloor n^\eta \rfloor + 1}^\sigma \mathbf{G}_h(\sqrt{n}Y_{j/n}) \right)^2 \right] = 0$ .

*Second step:* It holds that  $\sup_{s \in [0,1]} |D_s^n - d_s^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ .

This follows from [1, Proposition 1.2] as a consequence of the previous step and the martingale property of  $D^n - d^n$ .

*Third step:* It holds that  $\sup_{s \in [0,1]} |d_s^n| \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$ . This follows from (B.11) and from applying Proposition 9 to the sequence  $g_n := n^{\frac{1}{4}} Q_{\lfloor n^{\frac{3}{4}} \rfloor + 1}^\sigma \mathbf{G}_h$ . The assumptions of the latter proposition are satisfied indeed  $\lim_{n \rightarrow \infty} \|g_n\|_1 = 0$  follows from Item (i) in Lemma 2 and  $\lim_{n \rightarrow \infty} \frac{g_n(\sqrt{n}x)}{\sqrt{n}} = 0$  follows from the stronger fact, obtained in the first step, that  $g_n$  satisfies (B.1).

Combining the two last steps yields the conclusion.  $\square$

## B.4 Final steps of the proof

In this section we complete the proof of Proposition 4.

Let  $\gamma > 3$  and  $h \in \mathbf{I}_\gamma$  be fixed. By Definition 1 of  $\mathbf{I}_\gamma$  there exist a non-negative function  $\bar{h} \in \mathbf{L}^{1,b}(\lambda^{(\gamma)})$  and a constant  $a \in [0, \infty)$  such that  $|h(x, y)| \leq \bar{h}(x) e^{a|y-x|}$ . In this section  $\bar{h}$  and

$a$  are fixed.

Let us also recall some notation: let  $\mathbf{H}$  the functional in (2.7),  $Q^\sigma$  the semigroup in (A.4),  $\mathbf{G}_h$  in (B.2),  $\mathcal{Q}$  in (B.3) and its limit  $\mathfrak{Q}_{\sigma,h}$  in (2.9), and  $\mathbb{L}$  in (A.6).

For all  $n \in \mathbb{N}$  the  $(\mathcal{F}_{\lfloor nt/n \rfloor})_{t \in [0,1]}$ -martingale  $\mathcal{M}^n$  in (B.9) rewrites as

$$\mathcal{M}_t^n = \sum_{k=1}^{\lfloor nt \rfloor} \chi_k^n$$

where

$$\begin{aligned} \chi_k^n &:= n^{-\frac{1}{4}} \left( h(\sqrt{n}Y_{(k-1)/n}, \sqrt{n}Y_{k/n}) - \langle \lambda_\sigma, \mathbf{H}_h \rangle \sqrt{n}(L_{k/n}(Y) - L_{(k-1)/n}(Y)) \right) \\ &\quad + n^{-\frac{1}{4}} \left( \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{k/n}) - \mathcal{Q}_{n,0,1}(\sqrt{n}Y_{(k-1)/n}) \right). \end{aligned} \quad (\text{B.12})$$

Now it remains to prove Items (i)-(iv) in Proposition 4.

#### B.4.1 Proof of Item (i) in Proposition 4

For all  $n \in \mathbb{N}$ ,  $k \in \{1, \dots, \lfloor n \rfloor\}$ , the scaling property (A.1), Lemma 5 and (B.3) ensure that

$$\begin{aligned} &\mathbb{E}[\chi_k^n | \mathcal{F}_{(k-1)/n}] \\ &= n^{-\frac{1}{4}} \left( \mathbf{G}_h(\sqrt{n}Y_{(k-1)/n}) + \int_{-\infty}^{\infty} \mathcal{Q}_{n,0,0}(y) q_\sigma(1, \sqrt{n}Y_{(k-1)/n}, y) dy - \mathcal{Q}_{n,0,1}(\sqrt{n}Y_{(k-1)/n}) \right) \\ &= n^{-\frac{1}{4}} \left( Q_1^\sigma \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{(k-1)/n}) - \mathcal{Q}_{n,1,1}(\sqrt{n}Y_{(k-1)/n}) \right) = 0. \end{aligned}$$

#### B.4.2 Proof of Item (ii) in Proposition 4

Let  $t \in [0, 1]$  be fixed.

*First step:* Let us show that

$$\begin{aligned} &\sqrt{n} \mathbb{E}[(\chi_k^n)^2 | \mathcal{F}_{(k-1)/n}] \\ &= f_n(\sqrt{n}Y_{(k-1)/n}) + (\langle \lambda_\sigma, \mathbf{H}_h \rangle)^2 \mathbb{L}^{(2)}(1, \sqrt{n}Y_{(k-1)/n}) - 2\langle \lambda_\sigma, \mathbf{H}_h \rangle h_n(\sqrt{n}Y_{(k-1)/n}) \end{aligned}$$

where  $f_n$  and  $h_n$  are given by

$$\begin{aligned} f_n(x) &:= \mathbf{H}_{h^2}(x) + 2\mathbf{H}_{h, \mathcal{Q}_{n,0,0}}(x) + g_n(x), \quad \text{with } g_n(x) := Q_1^\sigma (\mathcal{Q}_{n,0,0})^2(x) - (\mathcal{Q}_{n,0,1}(x))^2 \quad \text{and} \\ h_n(x) &:= \mathbb{L}^{(1)}(h(x, \cdot) + \mathcal{Q}_{n,0,0}(\cdot), x) = \mathbb{L}^{(1)}(h(x, \cdot), x) + \mathbb{L}^{(1)}(\mathcal{Q}_{n,0,0}, x). \end{aligned}$$

In fact first note that, by (A.7), for all  $k = 1, \dots, \lfloor nt \rfloor$  it holds that

$$h_n(\sqrt{n}Y_{(k-1)/n}) = \sqrt{n} \mathbb{E} \left[ \left( h(\sqrt{n}Y_{\frac{k-1}{n}}, \sqrt{n}Y_{\frac{k}{n}}) + \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{\frac{k}{n}}) \right) (L_{\frac{k}{n}}(Y) - L_{\frac{k-1}{n}}(Y)) \middle| \mathcal{F}_{\frac{k-1}{n}} \right].$$

Let us now consider  $f_n$ . Clearly  $f_h$  have to be the sum of all remaining terms and it has to have the desired form. It does if

$$\begin{aligned} &\mathcal{Q}_{n,0,1}(\sqrt{n}Y_{(k-1)/n}) \\ &= \mathbb{E} \left[ h(\sqrt{n}Y_{(k-1)/n}, \sqrt{n}Y_{k/n}) + \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{k/n}) - \langle \lambda_\sigma, \mathbf{H}_h \rangle \sqrt{n}(L_{k/n}(Y) - L_{(k-1)/n}(Y)) \middle| \mathcal{F}_{(k-1)/n} \right]. \end{aligned}$$

and  $\mathbb{E} \left[ (\mathcal{Q}_{n,0,0}(\sqrt{n}Y_{k/n}))^2 \middle| \mathcal{F}_{(k-1)/n} \right] = Q_1^\sigma ((\mathcal{Q}_{n,0,0}(\sqrt{n}Y_{k/n}))^2)$ . The latter term follows from the scaling property (A.1) and equation (B.3), which also ensures that  $\mathbb{E} \left[ \mathcal{Q}_{n,0,0}(\sqrt{n}Y_{k/n}) \middle| \mathcal{F}_{(k-1)/n} \right] = \mathcal{Q}_{n,1,1}(\sqrt{n}Y_{(k-1)/n})$ . The first equality for  $\mathcal{Q}_{n,0,1}(\sqrt{n}Y_{(k-1)/n})$  follows from this, the definition of

$\mathbf{G}_h$  in (B.2), (A.7), and the fact that  $\mathcal{Q}_{n,1,1}(\sqrt{n}Y_{(k-1)/n}) = \mathcal{Q}_{n,0,1}(\sqrt{n}Y_{(k-1)/n}) - \mathbf{G}_h(\sqrt{n}Y_{(k-1)/n})$ . The proof of the first step is thus completed.

*Second step:* We show that  $\frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} \mathbb{L}^{(2)}(1, \sqrt{n}Y_{(k-1)/n}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \frac{8}{3\sqrt{2\pi}} L_t(Y)$ .

We apply in this step Proposition 10 to the constant sequence of functions  $\mathbb{L}^{(2)}(1, \cdot)$  which shows that  $\frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} \mathbb{L}^{(2)}(1, \sqrt{n}Y_{(k-1)/n}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \langle \lambda_\sigma, \mathbb{L}^{(2)}(1, \cdot) \rangle L_t(Y)$ . To apply it we need to check the assumptions:  $\mathbb{L}^{(2)}(1, \cdot) \in \mathbf{L}^{1,b}(\lambda^{(2)})$  and compute  $\langle \lambda_\sigma, \mathbb{L}^{(2)}(1, \cdot) \rangle$ .

Let us first note that  $\mathbb{E}[(L_1(W))^2] = \int_{-\infty}^{\infty} \int_0^{\infty} \ell^2 \rho_1(y, \ell) d\ell dy = 1$  where  $\rho$  is given by (A.3). This, Lemma 6, and simple computations show that, if  $K = \left(\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+}\right)^3 (\min\{\sigma_-, \sigma_+\})^{-1}$ , then  $\mathbb{L}^{(2)}(1, 0) \leq K$  and

$$\mathbb{L}^{(2)}(1, x) \leq K \frac{|x|}{\sigma(x)} \int_0^1 \frac{1}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} dt = \frac{K\sqrt{2}}{\sqrt{2\pi}} \int_{-\infty}^{-|x|/\sigma(x)} e^{-y^2/2} dy \leq \frac{K}{\sqrt{2}}.$$

We now prove that  $\mathbb{L}^{(2)}(1, \cdot) \in \mathbf{L}^{1,b}(\lambda^{(2)}) \subseteq \mathbf{L}^{1,b}(\lambda^{(1)})$ . The fact that  $\mathbb{E}[(L_1(W))^2] = 1$ , Lemma 6, and simple computations also yield for  $i = 2$

$$\begin{aligned} \|\mathbb{L}^{(2)}(1, \cdot)\|_{1,i} &= \left(\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+}\right)^2 \mathbb{E}[(L_1(W))^2] \int_{-\infty}^{\infty} \int_0^1 \frac{|x|^{i+1} (1-t)}{\sigma(x) \sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} dt dx \\ &= \left(\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+}\right)^2 \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (|x|\sigma(x))^{i+1} e^{-\frac{x^2}{2}} dx \int_0^1 t^{\frac{3}{2}} \frac{(1-t)}{t} dt \end{aligned}$$

(with integral on  $\mathbb{R}_\pm$ , instead of  $\mathbb{R}$  if  $\sigma_\mp = +\infty$ ) which is clearly finite and

$$\langle \lambda_\sigma, \mathbb{L}^{(2)}(1, \cdot) \rangle = \left(\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+}\right)^2 \left(\frac{1}{\sigma_+} + \frac{1}{\sigma_-}\right) \mathbb{E}[(L_1(W))^2] \int_0^1 \frac{1-t}{\sqrt{2\pi t}} dt = \left(\frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+}\right) \frac{8}{3\sqrt{2\pi}}.$$

In the two final steps, we want to apply Proposition 10 to the sequences  $f_n$  and  $h_n$ .

*Third step:* We show that  $\frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} f_n(\sqrt{n}Y_{(k-1)/n}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \langle \lambda_\sigma, \mathbf{H}_{h^2} + 2\mathbf{H}_{h, \mathbf{G}_h} \rangle L_t(Y)$ . applying Proposition 10. To do so we check that the sequence  $f_n$  satisfies (B.1) and that  $\lim_{n \rightarrow \infty} \langle \lambda_\sigma, f_n \rangle = \langle \lambda_\sigma, \mathbf{H}_{h^2} + 2\mathbf{H}_{h, \mathbf{G}_h} \rangle$ .

The fact that  $\lim_{n \rightarrow \infty} \langle \lambda_\sigma, g_n \rangle = 0$  follows from the fact that  $\lambda_\sigma$  is the stationary measure, inequality (B.4) with  $\zeta = \gamma - 1 > 2$ , and inequality (B.5): there exists  $K \in (0, \infty)$  such that

$$\begin{aligned} \langle \lambda_\sigma, g_n \rangle &= \langle \lambda_\sigma, (\mathcal{Q}_{n,0,0})^2 \rangle - \langle \lambda_\sigma, (\mathcal{Q}_{n,0,0} + Q_{\lfloor n^{\frac{1}{4}} \rfloor + 1}^\sigma \mathbf{G}_h)^2 \rangle \\ &= 2\langle \lambda_\sigma, \mathcal{Q}_{n,0,0} Q_{\lfloor n^{\frac{1}{4}} \rfloor + 1}^\sigma \mathbf{G}_h \rangle - \langle \lambda_\sigma, (Q_{\lfloor n^{\frac{1}{4}} \rfloor + 1}^\sigma \mathbf{G}_h)^2 \rangle \\ &\leq K n^{-\frac{1}{4}} (n^{-\frac{1}{4}} + \log(n)) \int_{-\infty}^{\infty} \frac{1}{(\sigma(x))^2} \left(1 + \left(|x| n^{-\frac{1}{8}} / \sigma(x)\right)^{2(\gamma-1)}\right)^{-1} dx \\ &\leq K n^{-\frac{1}{8}} (n^{-\frac{1}{4}} + \log(n)) \int_{-\infty}^{\infty} \frac{1}{(\sigma(x))^2} \left(1 + (|x|/\sigma(x))^{2(\gamma-1)}\right)^{-1} dx \end{aligned}$$

which is finite for  $n$  fixed and vanishes for  $n \rightarrow \infty$ .

Note that  $\mathbf{H}_{h, \mathcal{Q}_{n,0,0}} = \mathbf{H}_{h, \mathbf{G}_h} + \mathbf{H}_{h, \mathcal{Q}_{n,1,0}}$ . Lemma 8 (in particular the fact that  $\mathbf{G}_h$  is bounded) ensures that there exists a constant  $K \in (0, \infty)$  such that

$$|\mathbf{H}_{h, \mathbf{G}_h}(x)| \leq \bar{h}(x) (\sup_{y \in \mathbb{R}} |\mathbf{G}_h(y)|) \int_{-\infty}^{\infty} e^{a|y-x|} q_\sigma(1, x, y) dy \leq K \bar{h}(x).$$

By (B.6) there exists constants  $K_1, K_2 \in (0, \infty)$  (all depending on  $\sigma_{\pm}$  and  $K_2$  depending also on the constant  $a \in [0, \infty)$ ) such that

$$\begin{aligned} |\mathbf{H}_{h, \mathcal{Q}_{n,1,0}}(x)| &\leq K_1 \int_{-\infty}^{\infty} |h(x, y)|(1 + |y|)q_{\sigma}(1, x, y) dy \\ &\leq K_1 \bar{h}(x) \left( \int_{-\infty}^{\infty} (1 + |x| \mathbf{1}_{[-x, x]}(y) + |y| \mathbf{1}_{\mathbb{R} \setminus [-x, x]}(y)) e^{a|y-x|} q_{\sigma}(1, x, y) dy \right) \\ &\leq K_2 \bar{h}(x)(1 + |x|). \end{aligned}$$

The fact that  $h \in \mathbf{L}_{\gamma}$  ensures that  $\bar{h} \in \mathbf{L}^{1,b}(\lambda^{(1)})$  and so  $\mathbf{H}_{h, \mathcal{Q}_{n,0,0}} \in \mathbf{L}^1$ . Hence, dominated convergence and Lemma 9 show that  $\lim_{n \rightarrow \infty} \langle \lambda_{\sigma}, f_n \rangle = \langle \lambda_{\sigma}, \mathbf{H}_{h^2} + 2\mathbf{H}_{h, \mathbf{G}_{\sigma, h}} \rangle$ .

Let us now show that  $f_n$  satisfies equation (B.1). The fact that  $\bar{h} \in \mathbf{L}^{1,b}(\lambda^{(2)})$  ensures that  $\mathbf{H}_{h^2}(x) \leq K\bar{h}(x)$  for some positive constant  $K$  and so  $\mathbf{H}_{h^2} \in \mathbf{L}^{1,b}(\lambda^{(2)})$ .

Let us explore the contribution to (B.1) of the other parts of  $f_n$ . Let us first consider  $\mathbf{H}_{h, \mathcal{Q}_{n,0,0}} = \mathbf{H}_{h, \mathbf{G}_h} + \mathbf{H}_{h, \mathcal{Q}_{n,1,0}}$ . Above we saw that  $|\mathbf{H}_{h, \mathbf{G}_h}| \leq K\bar{h}$  with  $K$  non negative constant. Hölder's inequality implies that  $(\mathbf{H}_{h, \mathcal{Q}_{n,1,0}}(x))^2 \leq \int_{-\infty}^{\infty} h(x, y)^2 q_{\sigma}(1, x, y) dy \int_{-\infty}^{\infty} (\mathcal{Q}_{n,1,0}(y))^2 q_{\sigma}(1, x, y) dy$ . It can be easily shown that

$$\sup_{x \in \mathbb{R}} \int_{-\infty}^{\infty} (h(x, y))^2 q_{\sigma}(1, x, y) dy \leq \sup_{x \in \mathbb{R}} (\bar{h}(x))^2 \int_{-\infty}^{\infty} e^{2a|y-x|} q_{\sigma}(1, x, y) dy < \infty.$$

And by (B.5)(taking  $\zeta = \gamma - 1 > 2$ ) there exist constants  $K_1, K_2 \in (0, \infty)$  such that

$$\begin{aligned} \int_{-\infty}^{\infty} (\mathcal{Q}_{n,1,0}(y))^2 q_{\sigma}(1, x, y) dy &\leq K_1 \int_{-\infty}^{\infty} \frac{(\log n)^2 q_{\sigma}(1, x, y)}{(1 + |yn^{-\frac{1}{8}}/\sigma(y)|^{\gamma-1})^2} dy \\ &\leq K_1 (\log(n))^2 \frac{\left(1 + \left| \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+} \right| \right)}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{e^{-\frac{(y - \frac{x}{\sigma(x)})^2}{2}}}{(1 + |yn^{-\frac{1}{8}}/\sigma(y)|^{\gamma-1})^2} dy \leq \frac{K_2 (\log(n))^2}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{2(\gamma-1)}}. \end{aligned} \quad (\text{B.13})$$

The last two inequalities are consequences of the fact that  $q_{\sigma}(1, x, y) \leq \frac{(1 + \left| \frac{\sigma_- - \sigma_+}{\sigma_- + \sigma_+} \right|)}{\sqrt{2\pi}\sigma(y)} e^{-\frac{1}{2}(\frac{y}{\sigma(y)} - \frac{x}{\sigma(x)})^2}$  and of [13, Lemma 3.2] (or some computations). Therefore  $\mathbf{H}_{h, \mathcal{Q}_{n,1,0}}(x) \leq K \frac{\log(n)}{(1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{2(\gamma-1)})^{\frac{1}{2}}}$

and so  $\mathbf{H}_{h, \mathcal{Q}_{n,0,0}}(x) \leq K \left( \bar{h}(x) + \frac{\log(n)}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{\gamma-1}} \right)$ . Finally we consider the auxiliary function  $g_n$ : note that  $|g_n| \leq Q_1^{\sigma} (2\mathcal{Q}_{n,0,0}^2 + 2\mathbf{G}_h^2) + 2\mathcal{Q}_{n,1,1}^2 + 2\mathbf{G}_h^2$ . Inequality (B.5) and inequality (B.13) imply that  $Q_1^{\sigma} (\mathcal{Q}_{n,0,0}^2 + \mathcal{Q}_{n,1,1}^2) \leq \frac{K(\log(n))^2}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{2(\gamma-1)}}$  for some non negative constant  $K$ . Lemma 8, the fact that  $\gamma \geq 2$  and Item (iii) of Lemma 2 yields  $\mathbf{G}_h^2 \leq \mathbf{G}_h$  and  $Q_1^{\sigma} \mathbf{G}_h^2(x) \leq K(e^{-x^2/2} + \frac{1}{1 + (|x|/\sigma(x))^{\gamma}})$  for some non-negative constant  $K$ .

We conclude that there exists a constant  $K \in (0, \infty)$  such that for all  $x \in \mathbb{R}, n \in \mathbb{N}$

$$f_n(x) \leq K \left( \bar{h}(x) + \frac{1}{1 + |x|^{\gamma}} + \frac{\log(n)}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{\gamma-1}} + \frac{(\log(n))^2}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{2(\gamma-1)}} \right)$$

and this make possible to show that (B.1) is satisfied.

*Forth step:* We show that  $\frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} h_n(\sqrt{n}Y_{(k-1)/n}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} c_h L_t(Y)$  applying Proposition 10, where  $2\langle \lambda_{\sigma}, \mathbf{H}_h \rangle c_h := -K_h + \langle \lambda_{\sigma}, \mathbf{H}_{h^2} + 2\mathbf{H}_{h, \mathbf{G}_{\sigma, h}} \rangle + \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \frac{8}{3\sqrt{2\pi}} (\langle \lambda_{\sigma}, \mathbf{H}_h \rangle)^2$ . To do so we check that the sequence  $h_n$  satisfies (B.1) and that  $\lim_{n \rightarrow \infty} \langle \lambda_{\sigma}, h_n \rangle = c_h$ .



Inequality (B.6) and the fact that  $\mathbf{G}_h$  is bounded (see Lemma 8) imply that there exists a  $K \in (0, \infty)$  such that

$$\begin{aligned} & |\mathbb{L}^{(1)}(\mathcal{Q}_{n,0,0}, x)| \\ & \leq \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \frac{|x|}{\sigma(x)} \int_0^1 \frac{\sqrt{1-t} e^{-\frac{x^2}{2(\sigma(x))^2 t}}}{\sqrt{2\pi t^{\frac{3}{2}}}} \mathbb{E}[L_1(W)(\sigma(W_1))^{-1} |\mathcal{Q}_{n,0,0}(\sigma(W_1)W_1 \sqrt{1-t})|] dt \\ & \leq K \mathbb{E}[L_1(W)(1 + |W_1|)] |x| \int_0^1 \frac{\sqrt{1-t}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} dt. \end{aligned}$$

Similarly there exists a constant  $K \in (0, \infty)$  such that

$$|\mathbb{L}^{(1)}(h(x, \cdot), x)| \leq K \mathbb{E}[L_1(W) e^{a\sigma(W_1)|W_1|} \mathbf{1}_{\{\sigma(W_1) \in \mathbb{R}\}}] \bar{h}(x) |x| \int_0^1 \frac{\sqrt{1-t}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} dt.$$

Observe that  $\mathbb{E}[L_1(W)(1 + |W_1| + e^{a\sigma(W_1)|W_1|} \mathbf{1}_{\{\sigma(W_1) \in \mathbb{R}\}})] < \infty$  (see the joint density of Brownian motion and its local time (A.3)). This, boundedness of  $\bar{h}$ , and the change of variable  $s = \frac{x^2}{(\sigma(x))^2 t}$  show that there exists  $K \in (0, \infty)$  such that

$$\begin{aligned} & |\mathbb{L}^{(1)}(h(x, \cdot), x)| + |\mathbb{L}^{(1)}(\mathcal{Q}_{n,0,0}, x)| \leq K |x| \int_0^1 \frac{\sqrt{1-t}}{\sqrt{2\pi t^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 t}} dt. \\ & = K \frac{x^2}{\sigma(x)} \int_{\frac{x^2}{(\sigma(x))^2}}^\infty \frac{\sqrt{1 - \frac{x^2}{(\sigma(x))^2 s}}}{\sqrt{2\pi s}} e^{-\frac{s}{2}} ds \leq K |x| \int_{\frac{x^2}{(\sigma(x))^2}}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{s}{2}} ds = 2K |x| e^{-\frac{x^2}{(\sigma(x))^2}} \in \mathbf{L}^1. \end{aligned}$$

Hence dominated convergence, and Lemma 9 demonstrates that  $\lim_{n \rightarrow \infty} \langle \lambda_\sigma, \mathbb{L}^{(1)}(\mathcal{Q}_{n,0,0}, \cdot) \rangle = \langle \lambda_\sigma, \mathbb{L}^{(1)}(\mathfrak{G}_{\sigma,h}, \cdot) \rangle$ . Moreover the latter inequalities ensure also that  $h_n$  it clearly satisfy (B.1).

Lemma 6 allows us to rewrite  $c_h := \langle \lambda_\sigma, \mathbb{L}^{(1)}(h(\cdot, \cdot), \cdot) \rangle + \langle \lambda_\sigma, \mathbb{L}^{(1)}(\mathfrak{G}_{\sigma,h}, \cdot) \rangle$  as

$$\begin{aligned} c_h &= \sqrt{\frac{2}{\pi}} \frac{(\sigma_- + \sigma_+)}{2\sigma_- \sigma_+} \int_0^1 \sqrt{\frac{1}{t} - 1} \int_{-\infty}^\infty \int_0^\infty \rho_1^\sigma(y, \ell) \ell \mathfrak{G}_{\sigma,h}(y\sqrt{1-t}) d\ell dy dt \\ & \quad + \int_{-\infty}^\infty \int_0^1 \int_{-\infty}^\infty \int_0^\infty \frac{|x| e^{-\frac{x^2}{2(\sigma(x))^2}}}{\sqrt{2\pi}\sigma(x)} \sqrt{\frac{1}{t} - 1} \rho_1^\sigma(y, \ell) \ell h(x\sqrt{t}, y\sqrt{1-t}) d\ell dy dt \lambda_\sigma(dx) \end{aligned}$$

where  $\rho_1^\sigma$  is the joint density of a standard OBM and its local time at time 1 (given in (A.2)). First note that all integrands are integrable: the fact that  $|\mathfrak{G}_{\sigma,h}(y)| \leq K(1 + |y|)$  for some constant  $K \in (0, \infty)$  follows from Item (ii) in Lemma 2 and of the fact that  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = 0$  (see Lemma 8). Fubini ensures that we can change the order of the integrals. Next observe that for all  $y \in \mathbb{R}$  it holds that  $\int_0^\infty (\sigma(y))^2 \rho_1^\sigma(\sigma(y)y, \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \ell) \ell d\ell = \Phi(-|y|)$  which is the cumulative distribution function of a standard Gaussian random variable. This and simple changes of variable yield

$$\begin{aligned} c_h &= \sqrt{\frac{2}{\pi}} \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \int_0^1 \sqrt{\frac{1}{t} - 1} \int_{-\infty}^\infty \sigma(x) \Phi(-|x|) \mathfrak{G}_{\sigma,h}(\sigma(x)x\sqrt{1-t}) \lambda_\sigma(dx) dt \\ & \quad + \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \int_{-\infty}^\infty \int_0^1 \int_{-\infty}^\infty \frac{|x| e^{-\frac{x^2}{2}} \Phi(-|y|)}{\sqrt{2\pi}\sigma(x)\sigma(y)} \sqrt{\frac{1}{t} - 1} h(\sigma(x)x\sqrt{t}, \sigma(y)y\sqrt{1-t}) dy dt dx. \end{aligned}$$

Note that for all  $t \in [0, 1]$

$$\int_{-\infty}^\infty \Phi(-|y|) \sigma(y) \mathfrak{G}_{\sigma,h}(\sigma(y)y\sqrt{1-t}) \lambda_\sigma(dy) = \int_{-\infty}^\infty \frac{1}{\sqrt{1-t}} \Phi\left(-\frac{|y|}{\sqrt{1-t}}\right) \sigma(y) \mathfrak{G}_{\sigma,h}(\sigma(y)y) \lambda_\sigma(dy).$$

Finally for all  $y \in \mathbb{R}$  the integral  $\int_0^1 \frac{1}{\sqrt{t}} \Phi(-|y|/\sqrt{1-t}) dt$  with the change of variable  $s = \frac{|y|}{\sqrt{1-t}}$  becomes  $\int_{|y|}^{\infty} \Phi(-s) \frac{2y^2}{s^2 \sqrt{s^2 - y^2}} ds = e^{-\frac{y^2}{2}} - \sqrt{2\pi}|y|\Phi(-|y|)$ . We have therefore obtained the desired expression for  $c_h$ .

### B.4.3 Proof of Item (iii) in Proposition 4

Let  $t \in [0, 1]$  be fixed.

*First step:* Let us show that

$$\begin{aligned} & \sum_{k=1}^{\lfloor nt \rfloor} \mathbb{E}[\chi_k^n(Y_{k/n} - Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \\ &= \frac{1}{\sqrt{n}} \sum_{k=1}^{\lfloor nt \rfloor} \left( n^{-\frac{1}{4}} f_1(\sqrt{n}Y_{(k-1)/n}) + g_n(\sqrt{n}Y_{(k-1)/n}) + \langle \lambda_\sigma, \mathbf{H}_h \rangle n^{-\frac{1}{4}} f_2(\sqrt{n}Y_{(k-1)/n}) \right) \end{aligned}$$

where  $f_1, f_2, g_n$  are given by

$$\begin{aligned} f_1(x) &:= \int_{-\infty}^{\infty} (h(x, y) + \mathbf{G}_h(y))(y - x) q_\sigma(1, x, y) dy, \quad f_2(x) := x \mathbb{L}^{(1)}(1, x), \quad \text{and} \\ g_n(x) &:= n^{-\frac{1}{4}} \int_{-\infty}^{\infty} \mathcal{Q}_{n,1,0}(y)(y - x) q_\sigma(1, x, y) dy. \end{aligned}$$

Throughout the proof of this step let  $\mathbb{I}: x \mapsto x$  denote the identity function. It follows from (B.12), the fact that  $Y$  is a martingale, and (A.7) that

$$\begin{aligned} & \mathbb{E}[\chi_k^n(Y_{k/n} - Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \\ &= n^{-\frac{3}{4}} \mathbb{E}[(h(\sqrt{n}Y_{(k-1)/n}, \sqrt{n}Y_{k/n}) + \mathbf{G}_h(\sqrt{n}Y_{k/n}))(\sqrt{n}Y_{k/n} - \sqrt{n}Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \\ &\quad + n^{-\frac{3}{4}} (\sqrt{n}Y_{(k-1)/n} \mathbb{L}^{(1)}(1, \sqrt{n}Y_{(k-1)/n}) - \mathbb{L}^{(1)}(\mathbb{I}, \sqrt{n}Y_{(k-1)/n})) \\ &\quad + n^{-\frac{3}{4}} \mathbb{E}[\mathcal{Q}_{n,1,0}(\sqrt{n}Y_{k/n})(\sqrt{n}Y_{k/n} - \sqrt{n}Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \end{aligned}$$

and the scaling property (A.1) implies that

$$\begin{aligned} & \sqrt{n} \mathbb{E}[\chi_k^n(Y_{k/n} - Y_{(k-1)/n}) | \mathcal{F}_{(k-1)/n}] \\ &= n^{-\frac{1}{4}} (f_1(\sqrt{n}Y_{(k-1)/n}) + f_2(\sqrt{n}Y_{(k-1)/n}) - \mathbb{L}^{(1)}(\mathbb{I}, \sqrt{n}Y_{(k-1)/n})) + g_n(\sqrt{n}Y_{(k-1)/n}). \end{aligned}$$

To conclude the proof of this step we show that  $\mathbb{L}^{(1)}(\mathbb{I}, \cdot) = 0$ . This follows from the fact that for a Brownian motion starting at 0, say  $W$ , it holds that  $\mathbb{E}[L_1(W)W_1] = 0$  and Lemma 6 yields

$$\mathbb{L}^{(1)}(\mathbb{I}, x) = \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \mathbb{E}[L_1(W)W_1] \left( \mathbf{1}_{\{x=0\}} + \mathbf{1}_{\{x \neq 0\}} \frac{|x|}{\sigma(x)} \int_0^1 \frac{1-s}{\sqrt{2\pi s^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 s}} ds \right) = 0.$$

In the next steps we want to check that Proposition 9 can be applied to the sequences  $n^{-\frac{1}{4}} f_1$ ,  $n^{-\frac{1}{4}} f_2$  and  $g_n$ .

*Second step:* We show that  $f_1$  is bounded and integrable:  $f_1 \in \mathbf{L}^{1,b}(\lambda^{(0)})$ .

The fact that  $h \in \mathbf{I}_\gamma \subseteq \mathbf{I}_3$  and Lemma 8 ensure that  $\bar{h}$  and  $\mathbf{G}_h$  are in  $\mathbf{L}^{1,b}(\lambda^{(2)})$ . This, Cauchy-Schwarz, the fact that  $\left( \int_{-\infty}^{\infty} (y-x)^2 q_\sigma(1, x, y) dy \right) \leq \sigma_-^2 \mathbf{1}_{\{\sigma_- \in \mathbb{R}\}} + \sigma_+^2 \mathbf{1}_{\{\sigma_+ \in \mathbb{R}\}}$ , and the fact that  $\int_{-\infty}^{\infty} (\mathbf{G}_h(y))^2 q_\sigma(1, x, y) dy = Q_1^\sigma \mathbf{G}_h^2(x)$  yield

$$|f_1(x)| \leq \bar{h}(x) \int_{-\infty}^{\infty} e^{a|y-x|} |y-x| q_\sigma(1, x, y) dy + \sqrt{\sigma_-^2 \mathbf{1}_{\{\sigma_- \in \mathbb{R}\}} + \sigma_+^2 \mathbf{1}_{\{\sigma_+ \in \mathbb{R}\}}} (|Q_1^\sigma \mathbf{G}_h^2(x)|)^{\frac{1}{2}}.$$

By Lemma 8  $\mathbf{G}_h$  is bounded, i.e. there exists  $K \in (0, \infty)$  such that  $\sup_{x \in \mathbb{R}} |\mathbf{G}_h(x)| \leq K$  and  $\langle \lambda_\sigma, \mathbf{G}_h \rangle = 0$ , and so  $|\langle \lambda_\sigma, \mathbf{G}_h^2 \rangle| \leq K |\langle \lambda_\sigma, \mathbf{G}_h \rangle| = 0$ . And so Item (iii) in Lemma 2 implies, up to increase the constant  $K \in [1, \infty)$ , that

$$|Q_1^\sigma \mathbf{G}_h^2(x)| \leq K \frac{1}{1 + |x/\sigma(x)|^{\gamma-1}} + K \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2(\sigma(x))^2}} \leq 2K^2 \frac{1}{(1 + |x/\sigma(x)|^{\frac{\gamma-1}{2}})^2}.$$

Since  $\gamma > 3$  it holds that  $(|Q_1^\sigma \mathbf{G}_h^2(x)|)^{\frac{1}{2}} \in \mathbf{L}^{1,b}(\lambda^{(0)})$  and so  $|f_1| \in \mathbf{L}^{1,b}(\lambda^{(0)})$ .

*Third step:* We demonstrate that  $f_2$  is bounded and integrable.

Lemma 6 and the change of variable  $r = \frac{x^2}{(\sigma(x))^2 s}$  show that there exists a positive constant  $K$  such that

$$\begin{aligned} f_2(x) &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \frac{x|x|}{\sigma(x)} \mathbb{E} \left[ \frac{L_1(W)}{\sigma(W_1)} \right] \int_0^1 \frac{\sqrt{1-s}}{\sqrt{2\pi s^{\frac{3}{2}}}} e^{-\frac{x^2}{2(\sigma(x))^2 s}} ds \\ &= \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \frac{x|x|}{\sigma(x)} \mathbb{E} \left[ \frac{L_1(W)}{\sigma(W_1)} \right] \int_{\frac{x^2}{(\sigma(x))^2}}^\infty \frac{1}{\sqrt{2\pi}} \sqrt{\frac{(\sigma(x))^2}{x^2} - \frac{1}{r}} e^{-\frac{r}{2}} \frac{1}{\sqrt{r}} dr \mathbf{1}_{\{\sigma(x) \in \mathbb{R}\}} \\ &\leq \left( \frac{2\sigma_- \sigma_+}{\sigma_- + \sigma_+} \right)^2 \frac{|x| \mathbb{E}[L_1(W)]}{\min\{\sigma_-, \sigma_+\}} \int_{\frac{x^2}{(\sigma(x))^2}}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{r}{2}} \frac{1}{\sqrt{r}} dr \mathbf{1}_{\{\sigma(x) \in \mathbb{R}\}} \leq K e^{-\frac{x^2}{4(\sigma(x))^2}} \in \mathbf{L}^{1,b}(\lambda^{(0)}). \end{aligned}$$

In the last inequality we used that  $\mathbb{E}[L_1(W)] \in [0, \infty)$ . This step is thus completed.

*Forth step:* We prove that  $\int_{-\infty}^\infty |g_n(x)| dx$  and  $\frac{1}{\sqrt{n}} g_n(\sqrt{n}x)$  converge to 0.

Cauchy-Schwarz inequality ensures

$$|g_n(x)|^2 \leq n^{-\frac{1}{2}} \left( \int_{-\infty}^\infty (\mathcal{Q}_{n,1,0}(y))^2 q_\sigma(1, x, y) dy \right) \left( \int_{-\infty}^\infty (y-x)^2 q_\sigma(1, x, y) dy \right)$$

Note that  $\left( \int_{-\infty}^\infty (y-x)^2 q_\sigma(1, x, y) dy \right) \leq \sigma_-^2 \mathbf{1}_{\{\sigma_- \in \mathbb{R}\}} + \sigma_+^2 \mathbf{1}_{\{\sigma_+ \in \mathbb{R}\}}$  and by inequality (B.13) there exists a constant  $K \in (0, \infty)$  depending on  $\gamma$  and  $\sigma_\pm$  such that

$$|g_n(x)| \leq \frac{n^{-\frac{1}{4}} K_2 \log(n)}{1 + |xn^{-\frac{1}{8}}/\sigma(x)|^{\gamma-1}}.$$

It is now clear that  $\frac{g_n(\sqrt{n}x)}{\sqrt{n}}$  and the integral of  $g_n$  converge to 0 when  $n \rightarrow \infty$ .

In the last steps we proved that Proposition 9 can be applied and this completes the proof.

#### B.4.4 Proof of Item (iv) in Proposition 4

Let  $\varepsilon \in (0, \infty)$  be fixed. For every  $k$ , Hölder's inequality and Markov's inequality show

$$\begin{aligned} \mathbb{E}[|\chi_k^n|^2 \mathbf{1}_{\{|\chi_k^n| \geq \varepsilon\}} | \mathcal{F}_{(k-1)/n}] &\leq \left( \mathbb{E}[|\chi_k^n|^5 | \mathcal{F}_{(k-1)/n}] \right)^{\frac{2}{5}} \left( \mathbb{E}[\mathbf{1}_{\{|\chi_k^n| \geq \varepsilon\}} | \mathcal{F}_{(k-1)/n}] \right)^{\frac{3}{5}} \\ &\leq \left( \mathbb{E}[|\chi_k^n|^5 | \mathcal{F}_{(k-1)/n}] \right)^{\frac{2}{5}} \left( \mathbb{E}[|\chi_k^n|^5 | \mathcal{F}_{(k-1)/n}] \right)^{\frac{3}{5}} \varepsilon^{-5 \cdot \frac{3}{5}} = \mathbb{E}[|\chi_k^n|^5 | \mathcal{F}_{(k-1)/n}] \varepsilon^{-3}. \end{aligned}$$

The fact that  $h \in \mathbf{I}_\gamma \subseteq \mathbf{I}_0$  ensures that  $\bar{h}$  is bounded and integrable. This combined with Jensen's inequality and (B.7) ensures that  $\sup_{x \in \mathbb{R}} \bar{h}(x) + |\langle \lambda_\sigma, \mathbf{H}_h \rangle|$  is bounded by a constant  $K \in (0, \infty)$  and for all  $n \in \mathbb{N} \setminus \{0, 1, 2, 3\}$  it holds that

$$\begin{aligned} \mathbb{E}[|\chi_k^n|^5 | \mathcal{F}_{(k-1)/n}] &\leq 4^4 n^{-\frac{5}{4}} K^5 \left( \mathbb{E} \left[ e^{5a\sqrt{n}|Y_{k/n} - Y_{(k-1)/n}|} | \mathcal{F}_{(k-1)/n} \right] \right. \\ &\quad \left. + \mathbb{E} \left[ (\sqrt{n}|L_{k/n}(Y) - L_{(k-1)/n}(Y)|)^5 | \mathcal{F}_{(k-1)/n} \right] + (\log n)^5 \right). \end{aligned}$$

The fact that density of the OBM has a Gaussian behavior together with the scaling property (A.1) ensure that  $\mathbb{E}\left[e^{5a\sqrt{n}|Y_{k/n}-Y_{(k-1)/n}|}|\mathcal{F}_{(k-1)/n}\right] = \int_{-\infty}^{\infty} e^{5a|y-x|}q_{\sigma}(1,x,y)dy$  is bounded. By (A.6) and Lemma 6 we can show, similarly to the third step of the proof of Item (iii), that there exist constants  $K_1, K_2 \in (0, \infty)$  such that

$$\begin{aligned} & \mathbb{E}\left[\left(\sqrt{n}|L_{k/n}(Y) - L_{(k-1)/n}(Y)|\right)^5|\mathcal{F}_{(k-1)/n}\right] = \mathbb{L}^{(5)}(1, \sqrt{n}Y_{(k-1)/n}) \\ & \leq K_1 \int_0^1 \frac{(1-t)^{\frac{5}{2}}}{\sqrt{2\pi}t^{\frac{3}{2}}} \frac{\sqrt{n}|Y_{(k-1)/n}|}{\sigma(Y_{(k-1)/n})} e^{-\frac{n(Y_{(k-1)/n})^2}{2(\sigma(Y_{(k-1)/n}))^2 t}} dt \leq K_2. \end{aligned}$$

We conclude that there exists a constant  $K \in (0, \infty)$  such that

$$\sum_{k=1}^n \mathbb{E}[|\chi_k^n|^2 \mathbf{1}_{\{|\chi_k^n| \geq \varepsilon\}}|\mathcal{F}_{(k-1)/n}] \leq \sum_{k=1}^n \mathbb{E}[|\chi_k^n|^5|\mathcal{F}_{(k-1)/n}] \varepsilon^{-3} \leq nKn^{-\frac{5}{4}} \log(n)\varepsilon^{-3} \xrightarrow{n \rightarrow \infty} 0.$$

**Acknowledgments:** The author thanks Sylvie Rœlly and Paolo Pigato for fruitful discussions on the topic of this paper.

## References

- [1] David J. Aldous. Stopping times and tightness. II. *Ann. Probab.*, 17(2):586–595, 1989. ISSN 0091-1798. URL [http://links.jstor.org/sici?sici=0091-1798\(198904\)17:2<586:STATI>2.0.CO;2-5&origin=1](http://links.jstor.org/sici?sici=0091-1798(198904)17:2<586:STATI>2.0.CO;2-5&origin=1)
- [2] David J. Aldous and Geoffrey Kennedy Eagleson. On mixing and stability of limit theorems. *Ann. Probability*, 6(2):325–331, 1978. ISSN 0091-1798. doi: 10.1214/aop/1176995577. URL <https://doi.org/10.1214/aop/1176995577>.
- [3] Jean-Marc Azaïs. Approximation des trajectoires et temps local des diffusions. *Ann. Inst. H. Poincaré Probab. Statist.*, 25(2):175–194, 1989. ISSN 0246-0203. URL [http://www.numdam.org/item?id=AIHPB\\_1989\\_\\_25\\_2\\_175\\_0](http://www.numdam.org/item?id=AIHPB_1989__25_2_175_0).
- [4] Jean-Marc Azaïs. Convergence presque sûre du nombre de franchissements normalisé vers le temps local. *C. R. Acad. Sci. Paris Sér. I Math.*, 312(9):689–694, 1991. ISSN 0764-4442.
- [5] Patrick Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999. doi: 10.1002/9780470316962. A Wiley-Interscience Publication.
- [6] Andrei Borodin and Paavo Salminen. On the local time process of a skew Brownian motion. *Trans. Amer. Math. Soc.*, 372(5):3597–3618, 2019. ISSN 0002-9947. doi: 10.1090/tran/7852. URL <https://doi.org/10.1090/tran/7852>.
- [7] Andrei N. Borodin. On the character of convergence to Brownian local time. I, II. *Probab. Theory Relat. Fields*, 72(2):231–250, 251–277, 1986. ISSN 0178-8051. doi: 10.1007/BF00699105. URL <https://doi.org/10.1007/BF00699105>.
- [8] Andrei N. Borodin. Brownian local time. *Russian Math. Surveys*, 44(2(266)):7–48, 1989. ISSN 0042-1316. doi: 10.1070/RM1989v044n02ABEH002050. URL <https://doi.org/10.1070/RM1989v044n02ABEH002050>.
- [9] Danièle Florens-Zmirou. On estimating the diffusion coefficient from discrete observations. *J. Appl. Probab.*, 30(4):790–804, 1993.

- [10] John Michael Harrison and Lawrence A. Shepp. On skew Brownian motion. *Ann. Probab.*, 9(2):309–313, 1981. ISSN 0091-1798. URL [http://links.jstor.org/sici?sici=0091-1798\(198104\)9:2<309:OSBM>2.0.CO;2-W&origin=MS](http://links.jstor.org/sici?sici=0091-1798(198104)9:2<309:OSBM>2.0.CO;2-W&origin=MS)
- [11] Jean Jacod. Une généralisation des semimartingales: les processus admettant un processus à accroissements indépendants tangent. In *Seminar on probability, XVIII*, volume 1059 of *Lecture Notes in Math.*, pages 91–118. Springer, Berlin, 1984. doi: 10.1007/BFb0100035. URL <https://doi.org/10.1007/BFb0100035>.
- [12] Jean Jacod. On continuous conditional Gaussian martingales and stable convergence in law. In *Séminaire de Probabilités, XXXI*, volume 1655 of *Lecture Notes in Math.*, pages 232–246. Springer, Berlin, 1997. doi: 10.1007/BFb0119308. URL <https://doi.org/10.1007/BFb0119308>.
- [13] Jean Jacod. Rates of convergence to the local time of a diffusion. *Ann. Inst. H. Poincaré Probab. Statist.*, 34(4):505–544, 1998. ISSN 0246-0203. doi: 10.1016/S0246-0203(98)80026-5. URL [https://doi.org/10.1016/S0246-0203\(98\)80026-5](https://doi.org/10.1016/S0246-0203(98)80026-5).
- [14] Jean Jacod and Philip Protter. *Discretization of processes*, volume 67 of *Stochastic Modelling and Applied Probability*. Springer, Heidelberg, 2012. doi: 10.1007/978-3-642-24127-7.
- [15] Ioannis Karatzas and Steven E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991. ISBN 0-387-97655-8. doi: 10.1007/978-1-4612-0949-2. URL <http://dx.doi.org/10.1007/978-1-4612-0949-2>.
- [16] Julian Keilson and Jon A. Wellner. Oscillating Brownian motion. *J. Appl. Probability*, 15(2):300–310, 1978. ISSN 0021-9002. doi: 10.2307/3213403. URL <https://doi.org/10.2307/3213403>.
- [17] Jean-François Le Gall. One-dimensional stochastic differential equations involving the local times of the unknown process. In *Stochastic analysis and applications (Swansea, 1983)*, volume 1095 of *Lecture Notes in Math.*, pages 51–82. Springer, Berlin, 1984. doi: 10.1007/BFb0099122. URL <https://doi.org/10.1007/BFb0099122>.
- [18] Antoine Lejay. On the constructions of the skew Brownian motion. *Probab. Surv.*, 3:413–466, 2006. doi: 10.1214/154957807000000013. URL <https://doi.org/10.1214/154957807000000013>.
- [19] Antoine Lejay and Paolo Pigato. Statistical estimation of the oscillating Brownian motion. *Bernoulli*, 24(4B):3568–3602, 2018. ISSN 1350-7265. doi: 10.3150/17-BEJ969. URL <https://doi.org/10.3150/17-BEJ969>.
- [20] Antoine Lejay and Paolo Pigato. Maximum likelihood drift estimation for a threshold diffusion. *To appear in Scandinavian Journal of Statistics*, <https://doi.org/10.1111/sjos.12417>, 2018.
- [21] Antoine Lejay, Ernesto Mordecki, and Soledad Torres. Is a Brownian motion skew? *Scand. J. Stat.*, 41(2):346–364, 2014. ISSN 0303-6898. doi: 10.1111/sjos.12033. URL <https://doi.org/10.1111/sjos.12033>.
- [22] Antoine Lejay, Ernesto Mordecki, and Soledad Torres. Two consistent estimators for the skew Brownian motion. *ESAIM Probab. Stat.*, 23:567–583, 2019. ISSN 1292-8100. doi: 10.1051/ps/2018018. URL <https://doi.org/10.1051/ps/2018018>.

- [23] Mark Podolskij and Mathieu Rosenbaum. Comment on: Limit of random measures associated with the increments of a brownian semimartingale: Asymptotic behavior of local times related statistics for fractional brownian motion. *Journal of Financial Econometrics*, 16(4):588–598, 2017.
- [24] Alfréd Rényi. On stable sequences of events. *Sankhyā Ser. A*, 25:293–302, 1963. ISSN 0581-572X.
- [25] John B. Walsh. A diffusion with discontinuous local time. In *Temps locaux*, volume 52-53, pages 37–45. Société Mathématique de France, 1978.