

UNCERTAINTY PRINCIPLES ASSOCIATED TO SETS SATISFYING THE GEOMETRIC CONTROL CONDITION

WALTON GREEN, BENJAMIN JAYE, AND MISHKO MITKOVSKI

ABSTRACT. In this paper, we study forms of the uncertainty principle suggested by problems in control theory. First, we prove an analogue of the Paneah-Logvinenko-Sereda Theorem characterizing sets which satisfy the Geometric Control Condition (GCC). This result is applied to get a uniqueness result for functions with spectrum contained in sufficiently flat sets. One corollary is that a function with spectrum in an annulus of a given thickness can be bounded, in L^2 -norm, from above by its restriction to a neighborhood of a GCC set, with constant independent of the radius of the annulus. This result is applied to the energy decay rates for damped fractional wave equations.

1. INTRODUCTION

The goal of this paper is to investigate versions of the uncertainty principle suggested by control theory for PDEs. There is a long history of the relationship between these two fields beginning with Riesz sequence problems for non-harmonic Fourier series and their application to both wave and heat equations by the so-called “moment method” of D. L. Russell [2, 13, 28]. More recently, inequalities of the uncertainty principle type have found application to control theory on unbounded domains [8, 9, 19] and compact Riemannian manifolds of negative curvature, the latter being connected to the fractal uncertainty principle of Bourgain and Dyatlov [5, 7, 10, 15, 16].

Fix $k \in \{1, \dots, d\}$. For $\ell > 0$ and $\gamma > 0$, a set $E \subset \mathbb{R}^d$ satisfies the k -dimensional (ℓ, γ) -*geometric control condition* (GCC) if for any k -dimensional cube $Q \subset \mathbb{R}^d$ of side-length at least ℓ ,

$$\mathcal{H}^k(Q \cap E) \geq \gamma,$$

B.J. supported by NSF through DMS-1847301 and DMS-1800015.

M.M. supported by NSF through DMS-1600874.

where \mathcal{H}^k denotes the k -dimensional Hausdorff measure (which is just the k -dimensional Lebesgue measure on the k -dimensional plane containing Q). We say that E satisfies the k -GCC if it satisfies the k -dimensional (ℓ, γ) -GCC for some $\ell > 0$ and γ .

The GCC arises in the study of control theory for hyperbolic equations in the work of Bardos, Lebeau, Rauch, Taylor and Phillips [3, 27]. Given a Laplacian, an open set ω satisfies the GCC if for some $T > 0$, the Hamiltonian flow always intersects $[0, T] \times \omega$. Choosing the Laplacian to be $-\Delta$ on \mathbb{R}^d , and removing the regularity condition, this simplifies to the above condition with $k = 1$.

On the other hand, when $k = d$, we recover the definition of relatively dense, or thick, sets which are characterized by the Paneah-Logvinenko-Sereda (PLS) theorem [18, 20, 24] as sets E for which

$$(1.1) \quad \|f\|_{L^p(\mathbb{R}^d)} \leq C \|f\|_{L^p(E)}$$

for all f satisfying $\text{spec}(f) \subset Q$. Here $\text{spec}(f)$ denotes the support of the Fourier transform of f and Q is a d -dimensional cube of fixed side length.

Our first main result is a direct analogue of the PLS theorem for any $k \in \{1, \dots, d\}$, characterizing sets which (up to modification by sets of m_d -measure zero) satisfy the k -GCC as those for which (1.1) holds whenever $\text{spec}(f)$ is contained in a $(d-k)$ -dimensional ‘‘strip.’’ To precisely state the result, we need to introduce some additional notation.

For a set $A \subset \mathbb{R}^d$, and $\beta > 0$, $\mathcal{U}_\beta(A)$ denotes the open β neighbourhood of A in \mathbb{R}^d . A $(d-k)$ -plane is a $(d-k)$ -dimensional affine plane (which we interpret as a single point if $k = d$). For a set $S \subset \mathbb{R}^d$, we define¹

$$\beta_{d-k}(S) = \inf_{\substack{L \text{ is a} \\ d-k \text{ plane}}} \sup_{x \in S} \text{dist}(x, L)$$

Therefore, if $\beta_{d-k}(S) < \beta$ then there is a $(d-k)$ -plane L_S such that $S \subset \mathcal{U}_\beta(L_S)$.

Theorem 1. *Fix $k \in \{1, \dots, d\}$, $\ell > 0$ and $\gamma > 0$, $p \in [1, \infty)$. For every $\beta > 0$ there exists $C > 0$ such that if E satisfies the k -dimensional (ℓ, γ) -GCC, and $f \in L^p(\mathbb{R}^d)$ satisfies $\beta_{d-k}(\text{spec}(f)) < \beta$, then*

$$\|f\|_{L^p(\mathbb{R}^d)} \leq C \|f\|_{L^p(E)}.$$

¹The notation comes from Peter Jones’ analysts travelling salesman problem [17].

The constant C in Theorem 1 will take the form

$$C = \left(\frac{C_0 \ell}{\gamma} \right)^{C_0 \beta \ell},$$

where $C_0 = C_0(k) > 0$. The proof of Theorem 1 is a modification of the proof of the aforementioned PLS theorem given by Kovrijkine [18], where it is shown (in the case $k = d$) that the form of constant we obtain is sharp (up to the value of C_0).

Provided that one handles sets of measure zero appropriately, the k -GCC condition is necessary for the conclusion to hold, see Proposition 7 below.

We will use Theorem 1 to derive quantitative uniqueness properties for functions with spectrum contained in more complicated sets. Our general result in this regard is

Theorem 2. *Fix $k \in \{1, \dots, d\}$, $\beta > 0$, $\delta > 0$, $\gamma > 0$ and $\ell > 0$. There exists $R > 0$ and $C > 0$ such that if*

- (1) $\Gamma \subset \mathbb{R}^d$ satisfies that for any ball B of radius R centred on Γ ,

$$\beta_{d-k}(B \cap \Gamma) < \beta, \text{ and}$$

- (2) E satisfies the k -dimensional (ℓ, γ) -GCC,

then for any $f \in L^2(\mathbb{R}^d)$ with $\text{spec}(f) \subset \mathcal{U}_\beta(\Gamma)$,

$$\|f\|_{L^2(\mathbb{R}^d)} \leq C \|f\|_{L^2(\mathcal{U}_\delta(E))}.$$

In contrast with Theorem 1, observe that in the conclusion of Theorem 2, we only control the L^2 -norm of f by its norm on a δ -neighbourhood of a k -GCC set. In the generality that Theorem 2 is stated, one cannot expect R and C to be bounded independently of δ . This can be seen as a direct consequence of the sharpness of the classical Ingham inequality for non-harmonic trigonometric series [12, 14].

Inspecting the proof, we will show that Theorem 2 holds with C and R both taking the form

$$C_1 \left(\frac{\beta}{\delta} \right)^{d+1} \left(\frac{C_0 \ell}{\gamma} \right)^{C_0 \ell \beta}$$

where $C_0 = C_0(k)$ and $C_1 = C_1(k, d)$.

In our application, we will apply Theorem 2 with Γ being dilations of the sphere ($\Gamma = R\mathbb{S}^{d-1}$) in the following form.

Corollary 3. *Let E satisfy the 1-GCC. For any $\beta, \delta > 0$, there exists $C > 0$ such that*

$$(1.2) \quad \|f\|_{L^2(\mathbb{R}^d)} \leq C \|f\|_{L^2(\mathcal{U}_\delta(E))}$$

whenever $f \in L^2(\mathbb{R}^d)$ satisfies $\text{spec } f \subset \{\xi \in \mathbb{R}^d : R - \beta \leq |\xi| \leq R + \beta\}$ for some $R > 0$.

For any particular R , one can derive this inequality with a constant depending on R (either from Theorem 1 or the usual PLS Theorem) by placing the annulus $\{||\xi| - R| \leq \beta\} = \mathcal{U}_\beta(R\mathbb{S}^{d-1})$ inside a ball of radius $R + \beta$, but we emphasize that *the inequality (1.2) holds independent of R* . We pose the question of whether one can remove the δ neighborhood (see Question 5 below). This result on annuli follows from a more general application of Theorem 2, showing that Corollary 3 relies only on the compactness and smoothness of the sphere.

Corollary 4. *Fix $k \in \{1, \dots, d - 1\}$. Suppose that Σ is a compact, C^1 -smooth, $(d - k)$ -dimensional submanifold of \mathbb{R}^d . Suppose that E satisfies the k -GCC. Then for any $\delta > 0$ and $\beta > 0$, there is a constant $C > 0$ such that for every $R > 0$,*

$$\|f\|_{L^2(\mathbb{R}^d)} \leq C \|f\|_{L^2(\mathcal{U}_\delta(E))}$$

whenever $f \in L^2(\mathbb{R}^d)$ satisfies $\text{spec}(f) \subset \mathcal{U}_\beta(R\Sigma)$.

1.1. Application to the decay of damped wave equations. Fix $s > 0$ and a damping function $\gamma : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$.

We consider the fractional damped wave equation recently introduced by Malhi and Stanislavova in [22].

For $(x, t) \in \mathbb{R}^d \times \mathbb{R}_{\geq 0}$, let w satisfy

$$(1.3) \quad w_{tt}(x, t) + \gamma(x)w_t(x, t) + (-\Delta + 1)^{s/2}w(x, t) = 0.$$

The damping force is represented by γw_t and the fractional Laplacian is defined, for $r \in \mathbb{R}$ by

$$(-\Delta + 1)^r f(x) = \int_{\mathbb{R}^d} (|\xi|^2 + 1)^r \widehat{f}(\xi) e^{ix\xi} d\xi.$$

Herein, we study the decay rate of the energy of w , defined by

$$\begin{aligned} E(t) &= \|(w(t), w_t(t))\|_{H^{s/2} \times L^2} \\ &= \left(\int_{\mathbb{R}^d} |(-\Delta + 1)^{s/4} w(x, t)|^2 + |w_t(x, t)|^2 dx \right)^{1/2}. \end{aligned}$$

Standard analysis shows that if $\gamma = 0$, then the energy is conserved, i.e. there is no decay. On the other hand, for constant damping $\gamma = c > 0$,

it can be shown that $E(t)$ decays exponentially.

The classical case of $s = 2$ has been well-studied on bounded domains in the pioneering works of Bardos, Lebeau, Rauch, Taylor, and Phillips [3, 27]. Recently, Burq and Joly have extended these results to \mathbb{R}^d [6], and in particular showed that if γ is uniformly continuous and satisfies the GCC condition (1.4) below, then $E(t)$ decays exponentially in t . The methods in these works are that of microlocal and semiclassical analysis for which we refer to the book of Zworski [30]. These techniques, which allow one to handle very general Laplacians, impose regularity constraints on the damping coefficient γ .

We note two recent works which have, in one dimension, utilized Fourier analysis to prove exponential decay for rough damping [9, 21]. Fourier analytic methods have also proved useful in understanding (polynomial, or logarithmic) decay rates of the semi-group under weaker conditions than the GCC [1, 29].

Building upon the semi-group approach, see e.g. [1, 22, 29], the work of the first author in [9] connects (in one dimension) this problem of exponential decay to the PLS Theorem, specifically a version due to O. Kovrijkine [18], which may be viewed as (a sharper form of) Corollary 4 with Σ being a finite set of points. The proof in [18] uses the Turàn Lemma [23], and, in contrast to Corollary 4, does not require that one take a neighborhood of the GCC set. We wonder if this is the case in higher dimensions, and we specifically pose the following question, regarding the case of a sphere.

Question 5. *Suppose that E satisfies the 1-GCC. Does there exist $C > 0$ such that for every $R > 0$,*

$$\|f\|_{L^2(\mathbb{R}^d)} \leq C\|f\|_{L^2(E)}$$

for every $f \in L^2(\mathbb{R}^d)$ with $\text{spec}(f) \subset \mathcal{U}_1(R\mathbb{S}^{d-1})$?

Using the results of the previous section, particularly Corollary 4, we will prove a resolvent estimate (Proposition 11 below) for the fractional Laplacian, which then, using the strategy in [9], yields a new proof of the Burq-Joly theorem [6], along with its natural extension to the fractional wave equation.

Theorem 6. *Suppose γ is a non-negative, bounded, uniformly continuous function. There exists $L > 0$ and $c > 0$ such that*

$$(1.4) \quad \int_{\ell} \gamma(x) dm_1(x) \geq c > 0$$

for all line segments $\ell \subset \mathbb{R}^d$ of length L if and only if for every $s > 0$ there exists $C, \omega > 0$ such that

$$E(t) \leq \begin{cases} C(1+t)^{\frac{-s}{4-2s}} \|w(0), w_t(0)\|_{H^s \times H^{s/2}} & \text{if } 0 < s < 2 \\ Ce^{-\omega t} E(0) & \text{if } s \geq 2 \end{cases}$$

for all $t > 0$.

We reiterate that, in the case $s = 2$, Theorem 6 was proved by Burq and Joly [6] using semiclassical analysis. Our main goal here was to show how such results follow directly from uncertainty principles. The compactness methods of [6] enable one to prove Corollary 3 for β small enough depending on E and δ , but we do not know how to obtain the full strength of Corollary 3 by these methods. Consequently, the uncertainty principles developed here may have other applications to control theory problems for wave equations. Burq and Joly also pose the question of whether the result can be proved without the assumption of uniform continuity. This would follow from a positive answer to Question 5.

2. PROOF OF THEOREM 1

Throughout the paper, for $k \in \mathbb{N}$, m_k denotes the k -dimensional Lebesgue measure on \mathbb{R}^k .

Without loss of generality, we may assume that a best approximating plane for $\beta_{d-k}(\text{spec}(f))$ is the plane \mathbb{R}^{d-k} . Theorem 1 therefore follows from the following more precise proposition.

Proposition 7. *Fix $E \subset \mathbb{R}^d$. The following two conditions are equivalent:*

- (1) *There exist $\ell > 0$ and $\gamma > 0$ such that for m_{d-k} -almost every $x' \in \mathbb{R}^{d-k}$, whenever $Q \subset \mathbb{R}^k$ is a cube of length at least ℓ , then*

$$m_k(\{t \in Q : (t, x') \in E\}) \geq \gamma.$$

- (2) *For every $\beta > 0$, there exists $C > 0$ such that if $f \in L^p(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ satisfies $\text{supp}(\widehat{f}) \subset [-\beta, \beta]^k \times \mathbb{R}^{d-k}$, then*

$$\int_{\mathbb{R}^d} |f|^p dm_d \leq C \int_E |f|^p dm_d.$$

Moreover, in the direction (1) \implies (2) we will prove that the constant C takes the form

$$C = \left(\frac{C_2 \ell}{\gamma} \right)^{C_2 \beta \ell}$$

for a constant C_2 depending on k .

Proof. We first shall prove **(2)** \implies **(1)**. The proof in this direction follows an idea of Paneah [24] as presented Havin and Jorricke's book [11]. Fix $p \in (1, \infty)$.

Suppose that $f(t, x') = g(t)[h(x')]^{1/p}$, where $g \in L^p(\mathbb{R}^k)$ satisfies $\text{spec}(g) \in [-\beta, \beta]^k$, and $h \in L^1(\mathbb{R}^{d-k})$, $h \geq 0$ and $\|h\|_1 = 1$. Then, by Tonelli's theorem,

$$1 \leq C \int_{\mathbb{R}^{d-k}} h(x') \int_{\{t \in \mathbb{R}^k : (t, x') \in E\}} |g(t)|^p dm_k(t) dm_{d-k}(x').$$

Insofar as the space $L^p(\mathbb{R}^k)$ is separable, we therefore find that, for m_{d-k} -almost every $x' \in \mathbb{R}^{d-k}$,

$$C \int_{\{t \in \mathbb{R}^k : (t, x') \in E\}} |g(t)|^p dm_k(t) \geq 1 \text{ for every } g \in \mathcal{F},$$

where $\mathcal{F} = \{g \in L^p(\mathbb{R}^k) : \|g\|_p = 1 \text{ and } \text{spec}(g) \in [-\beta, \beta]^k\}$.

Fix any $g \in \mathcal{F}$ (it is clearly a non-empty set). Then there exists $M > 0$ such that

$$\int_{\{t \in \mathbb{R}^k : |g(t)| \geq M\}} |g(t)|^p dm_k(t) \leq \frac{1}{4C},$$

along with $\ell > 0$ such that

$$\int_{\mathbb{R}^k \setminus [-\ell, \ell]^k} |g(t)|^p dm_k(t) \leq \frac{1}{4C}.$$

Taking into account the fact that for every $z' \in \mathbb{R}^k$, $g(\cdot - z') \in \mathcal{F}$, we therefore get that, for m_{d-k} -almost every $x' \in \mathbb{R}^{d-k}$,

$$M m_k(\{t \in z' + [-\ell, \ell]^k : (t, x') \in E\}) \geq \frac{1}{2C} \text{ for every } z' \in \mathbb{R}^k.$$

Therefore (1) holds.

(1) \implies **(2)**. We may assume that $\ell = 1$ by replacing β by $\beta\ell$ and γ by $\gamma/\ell \in (0, 1)$. By modifying the set E by a set of Lebesgue measure zero in \mathbb{R}^d (which does not change the integral in (2)), we may assume that the condition in (1) holds for every $x' \in \mathbb{R}^{d-k}$.

Suppose $\|f\|_{L^p(\mathbb{R}^d)} = 1$. We also may assume without loss of generality that $f \in \mathcal{S}(\mathbb{R}^d)$.

Choose $\psi_0 \in \mathcal{S}(\mathbb{R}^k)$ with $\widehat{\psi} \equiv 1$ on $[-1, 1]^k$ and $\text{spec}(\psi_0) \subset [-2, 2]^k$. Put $\psi = \psi_0(\cdot/\beta)$. We first claim that we have $f = f *_k \psi$, where $*_k$ denotes a convolution in the first k variables. To see this, for $\xi \in \mathbb{R}^k$

and $\eta \in \mathbb{R}^{d-k}$, write

$$\begin{aligned} \widehat{f}(\xi, \eta) &= \widehat{f}(\xi, \eta) \widehat{\psi}(\xi) \\ &= \int_{\mathbb{R}^{d-k}} \int_{\mathbb{R}^k} \int_{\mathbb{R}^k} f(t, x') \psi(s) e^{-2\pi i \xi \cdot (t+s)} dm_k(t) dm_k(s) e^{-2\pi i x' \cdot \eta} dm_{d-k}(x') \\ &= \int_{\mathbb{R}^{d-k}} \int_{\mathbb{R}^k} \int_{\mathbb{R}^k} f(t) \psi(\tau - t) dm_k(t) e^{-2\pi i \xi \cdot \tau} dm_k(\tau) e^{-2\pi i x' \cdot \eta} dm_{d-k}(x') \\ &= \widehat{f *_{\mathbb{R}^k} \psi}(\xi, \eta). \end{aligned}$$

Fix $x' \in \mathbb{R}^{d-k}$, and set $F = f(\cdot, x')$ so $F : \mathbb{R}^k \rightarrow \mathbb{R}$. Then for any multi-index $\alpha \in \mathbb{Z}_+^k$, we have $F = \psi *_{\mathbb{R}^k} \psi *_{\mathbb{R}^k} \cdots \psi *_{\mathbb{R}^k} F$, where there are $|\alpha| := \sum_{j=1}^k \alpha_j$ convolutions of ψ . Then, with $C_0 = \|\nabla \psi_0\|_{L^1(\mathbb{R}^k)}$, we have

$$\|D^\alpha F\|_{L^p(\mathbb{R}^k)} \leq \beta^{|\alpha|} \|\nabla \psi_0\|_{L^1(\mathbb{R}^k)}^{|\alpha|} \|F\|_{L^p(\mathbb{R}^k)} = (\beta C_0)^{|\alpha|} \|F\|_{L^p(\mathbb{R}^k)}.$$

Fix $A > C_0$ to be chosen momentarily. We split \mathbb{R}^k into cubes of length 1. We call such a cube I *good* if

$$\int_I |D^\alpha F|^p dm_k \leq (\beta A)^{|\alpha|p} \int_I |F|^p dm_k \text{ for every } \alpha \in \mathbb{Z}_+^k.$$

Observe that, insofar as there are at most $(n+1)^k$ $\alpha \in \mathbb{Z}_+^k$ with $|\alpha| = n$

$$\begin{aligned} \sum_{I \text{ not good}} \int_I |F|^p dm_k &\leq \sum_{n=1}^{\infty} \sum_{\alpha \in \mathbb{Z}_+^k : |\alpha|=n} \frac{1}{(\beta A)^{|\alpha|p}} \int_{\mathbb{R}^k} |D^\alpha F|^p dm_k \\ &\leq \sum_{n=1}^{\infty} (n+1)^k \frac{C_0^{np}}{A^{np}} \int_{\mathbb{R}^k} |F|^p dm_k. \end{aligned}$$

Therefore, if A is large enough in terms of C_0 (which fixes A in terms of k), then

$$\int_{\mathbb{R}^k} |F|^p dm_k \leq 2 \sum_{I \text{ good}} \int_I |F|^p dm_k.$$

Now fix a good cube I , and put $Z = \{t \in I : (t, x') \in E\}$. By assumption, $m_k(Z) \geq \gamma$. Since I is good,

$$\int_I |D^\alpha F|^p dm_k \leq (\beta A)^{p|\alpha|} \int_I |F|^p dm_k \text{ for every } \alpha \in \mathbb{Z}_+^k,$$

and so by appealing to a standard Remez inequality for analytic functions (as in [18], see also [15, Corollary 2.8], [19, Proposition 3.7]),

$$\int_I |F|^p dm_k \leq C(\gamma, \beta A) \int_Z |F|^p dm_k,$$

where $C(\gamma, \beta A) = (C_1/\gamma)^{C_1\beta A}$ for $C_1 = C_1(k)$.

Summing over good intervals, we infer that for every $x' \in \mathbb{R}^{d-k}$,

$$\int_{\mathbb{R}^k} |f(t, x')|^p dm_k(t) \leq C(\gamma, \beta A) \int_{\{t \in \mathbb{R}^k : (t, x') \in E\}} |f(t, x')|^p dm_k(t).$$

Finally, integrating over $x' \in \mathbb{R}^{d-k}$ yields that

$$\int_{\mathbb{R}^d} |f|^p dm_d \leq C(\gamma, \beta A) \int_{\mathbb{R}^{d-k}} \int_{\{t \in \mathbb{R}^k : (t, x') \in E\}} |f(t, x')|^p dm_k(t) dm_{d-k}(x').$$

The right hand side is bounded by $C(\gamma, \beta A) \int_E |f|^p dm_d$. Setting $C_2 = AC_1$ shows that we have found the desired form of constant. \square

Remark 8. It easily follows from the proof of the direction (2) \implies (1) of Proposition 7 that if $E \subset \mathbb{R}^d$ is such that *there exists* $\beta > 0$, $C > 0$ such that

$$\|f\|_{L^p(\mathbb{R}^d)} \leq C \|f\|_{L^p(E)} \text{ whenever } \beta_{d-k}(\text{spec}(f)) \leq \beta,$$

then *any open neighborhood of E satisfies the k-GCC*.

3. PROOF OF THEOREM 2 AND COROLLARY 4

Proof of Theorem 2. Fix $k \in \{1, \dots, d\}$, and $\beta \geq 1$. As before, we may suppose that $\ell = 1$.

Choose ξ_ℓ to be a maximal $\beta/2$ -separated subset of Γ . Then we may cover $\mathcal{U}_\beta(\Gamma)$ by cubes $Q_\ell = Q(\xi_\ell, 2\beta)$ of sidelength 2β centred on $\xi_\ell \in \Gamma$.

Fix $\psi \in C_0^\infty(B(0, 1))$ with $\int_{\mathbb{R}^d} \psi dm_d = 1$ and $|\widehat{\psi}(\xi)| \gtrsim 1$ on $Q(0, 2)$.

Put $\delta_0 = \delta/\beta \leq \delta$. Set $\varphi = \delta_0^{-d} \psi(\delta_0^{-1} \cdot) \in C_0^\infty(B(0, \delta))$ so that $\widehat{\varphi} = \widehat{\psi}(\delta_0 \cdot)$. Then $\int_{\mathbb{R}^d} \varphi dm_d = 1$, $|\widehat{\varphi}(\xi)| \gtrsim 1$ on $Q(0, 2\beta)$, and for any multi-index $m \in \mathbb{N}$,

$$|\widehat{\varphi}(\xi)| \lesssim_m \frac{1}{\delta_0^{d+m} (1 + |\xi|)^m}.$$

Fix ℓ fixed, consider the function

$$g_\ell = f * \mathcal{F}^{-1} \widehat{\varphi}(\cdot - \xi_\ell),$$

so that $\|\widehat{f}\|_{L^2(Q_\ell)} \lesssim \|\widehat{g}_\ell\|_{L^2(\mathbb{R}^d)}$.

Now split $g_\ell = g_\ell^{(1)} + g_\ell^{(2)}$ where $g_\ell^{(1)} = g_\ell \chi_{\{|\xi - \xi_\ell| \leq R\}}$. Observe that, for fixed $m \in \mathbb{N}$,

$$\begin{aligned} \|g_\ell^{(2)}\|_{L^2(\mathbb{R}^d)}^2 &= \int_{|\xi - \xi_\ell| > R} |\widehat{f}(\xi)|^2 |\widehat{\varphi}(\xi - \xi_\ell)|^2 dm_d(\xi) \\ (3.1) \quad &\lesssim_m \delta_0^{-d-m} \int_{|\xi - \xi_\ell| > R} |\widehat{f}(\xi)|^2 \frac{1}{|\xi - \xi_\ell|^m} dm_d(\xi). \end{aligned}$$

By assumption $\text{supp}(\widehat{g_\ell^{(1)}})$ is contained in the β -neighbourhood of a $(d-k)$ -plane, and so Theorem 1, with $A = (\frac{C}{\gamma})^{C\beta}$,

$$\|g_\ell^{(1)}\|_{L^2(\mathbb{R}^d)} \lesssim A \|g_\ell^{(1)}\|_{L^2(E)}.$$

Whence, applying (3.1) twice, we infer that for every $m \in \mathbb{N}$,

$$\begin{aligned} \|\widehat{f}\|_{L^2(Q_\ell)}^2 &\lesssim \|g_\ell\|_{L^2(\mathbb{R}^d)}^2 \\ &\lesssim_m A \|g_\ell^{(1)}\|_{L^2(E)}^2 + \delta_0^{-d-m} \int_{|\xi - \xi_\ell| > R} |\widehat{f}(\xi)|^2 \frac{1}{|\xi - \xi_\ell|^m} dm_d(\xi) \\ &\lesssim_m A \|g_\ell\|_{L^2(E)}^2 + A \delta_0^{-d-m} \int_{|\xi - \xi_\ell| > R} |\widehat{f}(\xi)|^2 \frac{1}{|\xi - \xi_\ell|^m} dm_d(\xi). \end{aligned}$$

We next would like to sum this inequality over ℓ , using the fact that $\mathcal{U}_\beta(\Gamma) \subset \bigcup_\ell Q_\ell$.

Due to the support property of φ , we have that $g_\ell = (f\chi_{\mathcal{U}_\delta(E)}) * \mathcal{F}^{-1}\widehat{\varphi}(\cdot - \xi_\ell)$ on E . Whence,

$$\begin{aligned} \|g_\ell\|_{L^2(E)}^2 &\leq \int_{\mathbb{R}^d} |f\widehat{\chi_{\mathcal{U}_\delta(E)}}(\xi)|^2 |\widehat{\varphi}(\xi - \xi_\ell)|^2 dm_d(\xi) \\ &\lesssim \delta_0^{-d-1} \int_{\mathbb{R}^d} \frac{|f\widehat{\chi_{\mathcal{U}_\delta(E)}}(\xi)|^2}{1 + |\xi - \xi_\ell|^{d+1}} dm_d(\xi), \end{aligned}$$

but,

$$\sum_\ell \int_{\mathbb{R}^d} \frac{|f\widehat{\chi_{\mathcal{U}_\delta(E)}}(\xi)|^2}{1 + |\xi - \xi_\ell|^{d+1}} dm_d(\xi) \lesssim \int_{\mathbb{R}^d} |f\widehat{\chi_{\mathcal{U}_\delta(E)}}|^2 dm_d,$$

and so

$$(3.2) \quad \sum_\ell \|g_\ell\|_{L^2(E)}^2 \lesssim \delta_0^{-d-1} \|f\|_{L^2(\mathcal{U}_\delta(E))}^2.$$

On the other hand, for any $\xi \in \mathbb{R}^d$, and for any $k \geq 1$, there can be at most $C2^{kd}R^d$ of the $\beta/2$ -separated points ξ_ℓ in an annulus $A_k(\xi) := B(\xi, 2^k R) \setminus B(\xi, 2^{k-1} R)$ (crudely using that $\beta \geq 1$). Therefore,

$$(3.3) \quad \begin{aligned} \sum_\ell \frac{\chi_{\{\ell: |\xi_\ell - \xi| > R\}}(\ell)}{|\xi - \xi_\ell|^m} &\leq \sum_{k=1}^{\infty} \sum_{\ell: \xi_\ell \in A_k(\xi)} \frac{1}{|\xi - \xi_\ell|^m} \lesssim \sum_{k=1}^{\infty} \frac{2^{kd} R^d}{2^{(k-1)m} R^m} \\ &\lesssim_m \frac{1}{R^{m-d}} \text{ provided that } m > d. \end{aligned}$$

Consequently, if we set $m = d + 1$, then

$$(3.4) \quad \sum_\ell \int_{\{|\xi - \xi_\ell| > R\}} \frac{|\widehat{f}(\xi)|^2}{|\xi - \xi_\ell|^m} dm_d(\xi) \lesssim \frac{1}{R} \|f\|_{L^2(\mathbb{R}^d)}^2.$$

Combining (3.2) and (3.4) results in

$$\|f\|_{L^2(\mathbb{R}^d)}^2 \lesssim \frac{A}{\delta_0^{d+1}} \|f\|_{L^2(\mathcal{U}_\delta(E))}^2 + \frac{A}{\delta_0^{d+1}R} \|f\|_{L^2(\mathbb{R}^d)}^2.$$

If R is sufficiently large multiple of $A\delta_0^{-d-1}$, then the second term on the right hand side can be hidden in the left hand side, and we get

$$\|f\|_{L^2(\mathbb{R}^d)}^2 \lesssim \frac{A}{\delta_0^{d+1}} \|f\|_{L^2(\mathcal{U}_\delta(E))}^2.$$

The theorem is proved. \square

Proof of Corollary 4. Fix $\delta > 0$. Choose S large enough to be able to apply Proposition 2 with R replaced by S . Since Σ is a compact $(d-k)$ -dimensional C^1 -manifold embedded in \mathbb{R}^d , we can find a function $\sigma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\sigma(r)/r \rightarrow 0$ as $r \rightarrow 0$, such that if $x \in \Sigma$, then $\beta_{d-k}(\Sigma \cap B(x, r)) \leq \sigma(r)$. Fix $R \gg S$. Then for any $x \in R\Sigma$,

$$\beta_{d-k}(R\Sigma \cap B(x, S)) = R\beta_{d-k}(\Sigma \cap B(x, S/R)) \leq R\sigma(S/R) \leq \beta.$$

provided that R is large enough. But then we apply Proposition 2 to conclude that the desired statement holds for sufficiently large R . On the other hand, for small R , we can instead apply Theorem 1 by covering $\mathcal{U}_\beta(R\Sigma)$ by a strip of width $O_{\Sigma, \beta}(R)$. \square

4. DECAY RATES FOR DAMPED WAVE SEMIGROUPS

Throughout this section, for a function f we set $\|f\| := \|f\|_{L^2(\mathbb{R}^d)}$. For an operator A between two normed spaces, $\|A\|$ denotes the operator norm.

We view the equation (1.3) as the following semigroup. Setting $W(t) = (w(t), w_t(t))$,

$$\frac{d}{dt}W(t) = \mathcal{A}_\gamma W(t)$$

where $\mathcal{A}_\gamma : H^s \times H^{s/2} \rightarrow H^{s/2} \times L^2$ is densely defined by $\mathcal{A}_\gamma(u_1, u_2) = (u_2, -(-\Delta + 1)^{s/2}u_1 - \gamma u_2)$. The Sobolev space H^r for $r \in \mathbb{R}$ is defined by the decay of the Fourier transform:

$$H^r := \left\{ u \in L^2 : \|u\|_{H^r}^2 = \int_{\mathbb{R}^d} (|\xi|^2 + 1)^r |\widehat{u}(\xi)|^2 d\xi < \infty \right\}.$$

It is easy to verify that \mathcal{A}_0 is skew adjoint. That $W(t) = e^{t\mathcal{A}_\gamma}$ is a strong semigroup of contractions follows by the standard theory [25] since \mathcal{A}_γ is closed with dense range, and for any $U = (u_1, u_2) \in H^s \times H^{s/2}$,

$$\begin{aligned} \operatorname{Re}\langle \mathcal{A}_\gamma^* U, U \rangle_{H^{s/2} \times L^2} &= \operatorname{Re}\langle \mathcal{A}_\gamma U, U \rangle_{H^{s/2} \times L^2} \\ &= \operatorname{Re}\langle \mathcal{A}_0 U, U \rangle_{H^{s/2} \times L^2} - \langle \gamma u_2, u_2 \rangle_{L^2} = -\langle \gamma u_2, u_2 \rangle_{L^2} \leq 0. \end{aligned}$$

Notice that $E(t) = \|e^{t\mathcal{A}_\gamma}(w(0), w_t(0))\|_{H^{s/2} \times L^2}$, so the energy decay rates in Theorem 6 can be rewritten as

$$\begin{aligned} \|e^{t\mathcal{A}_\gamma} \mathcal{A}_\gamma^{-1}\| &= O(t^{\frac{-s}{4-2s}}) \quad 0 < s < 2, \\ \|e^{t\mathcal{A}_\gamma}\| &\leq C e^{-\omega t} \quad s \geq 2. \end{aligned}$$

Once we can establish

$$(4.1) \quad \|(\mathcal{A}_\gamma - i\lambda)^{-1}\| \leq C \max\{(|\lambda| + 1)^{\frac{4}{s}-2}, 1\},$$

the decay rates will follow from the following two results from semigroup theory.

Theorem 9 (Gearhart-Pruss Test [26]). *Let e^{tA} be a C_0 -semigroup in a Hilbert space and assume there exists $M > 0$ such that $\|e^{tA}\| \leq M$ for all $t \geq 0$. Then, there exists $C, \omega > 0$ such that*

$$\|e^{tA}\| \leq C e^{-\omega t}$$

if and only if $i\mathbb{R} \subset \rho(A)$ and $\sup_{\lambda \in \mathbb{R}} \|(A - i\lambda)^{-1}\| < \infty$.

For the polynomial decay, we use the following result from [4, Theorem 2.4]:

Theorem 10 (Borichev-Tomilov). *Let e^{tA} be a C_0 -semigroup on a Hilbert space. Assume there exists $M > 0$ such that $\|e^{tA}\| \leq M$ for all $t \geq 0$ and $i\mathbb{R} \subset \rho(A)$. Then for a fixed $\alpha > 0$,*

$$\|e^{tA} A^{-1}\| = O(t^{-1/\alpha}) \text{ as } t \rightarrow \infty$$

if and only if $\|(A - i\lambda)^{-1}\| = O(\lambda^\alpha)$ as $\lambda \rightarrow \infty$.

The main step in establishing (4.1) is the following resolvent estimate for the fractional Laplacian from sets which satisfy the Geometric Control Condition.

Proposition 11. *Let $E \subset \mathbb{R}^d$ satisfy the GCC and $\delta > 0$. Then, there exists $C > 0$ such that*

$$\|f\|^2 \leq C(1 + \lambda)^{\frac{2}{s}-2} \|((-\Delta + 1)^{s/2} - \lambda)f\|^2 + \|f\|_{L^2(\mathcal{U}_\delta(E))}^2$$

for all $f \in L^2(\mathbb{R}^d)$, $\lambda \geq 0$.

Proof. First, consider the annulus $A_\mu = \{\xi \in \mathbb{R}^d : ||\xi| - \mu| \leq 1\}$ for $\mu \geq 0$. Appealing to Corollary 3, we find a constant $C > 0$ (independent of μ) such that

$$(4.2) \quad \|f\| \leq C \|f\|_{L^2(\mathcal{U}_\delta(E))} \text{ whenever } \text{spec } f \subset A_\mu.$$

Now, define the Fourier restriction $\widehat{P_\lambda f} := \chi_{\widetilde{A}_\lambda} \widehat{f}$ where

$$\widetilde{A}_\lambda = \{\xi \in \mathbb{R}^d : |(|\xi|^2 + 1)^{1/2} - \lambda^{1/s}| \leq 1/2\}.$$

Then, for $\lambda \geq 3^s$, $\tilde{A}_\lambda \subset A_{\lambda^{1/s}}$ so for any $f \in L^2(\mathbb{R}^d)$,

$$\|f\|^2 \leq \|(I-P_\lambda)f\|^2 + C\|P_\lambda f\|_{L^2(\mathcal{U}_\delta(E))}^2 \leq (2C+1)\|(I-P_\lambda)f\|^2 + 2C\|f\|_{L^2(\mathcal{U}_\delta(E))}^2.$$

Finally, using the fact that $|\tau^s - \lambda| \geq c(1+\lambda)^{1-\frac{1}{s}}$ if $|\tau - \lambda^{1/s}| \geq 1$ (see Lemma 1 in [9]), we have

$$\|(I-P_\lambda)f\|^2 = \int_{\tilde{A}_\lambda^c} |\hat{f}(\xi)|^2 d\xi \leq C(1+\lambda)^{\frac{2}{s}-2} \int_{\mathbb{R}^d} [(|\xi|^2 + 1)^{s/2} - \lambda]^2 |\hat{f}(\xi)|^2 d\xi$$

which completes the proof. \square

To prove (4.1) one can follow the strategy from [9] which we briefly outline. First, the parallelogram identity and the positivity of $(-\Delta + 1)^{s/2}$ yields

$$(4.3) \quad c\|U\|_{H^{s/2} \times L^2}^2 \leq (1+|\lambda|)^{\frac{4}{s}-2} \|(\mathcal{A}_0 - i\lambda)U\|_{H^{s/2} \times L^2}^2 + \|u_2\|_{L^2(\mathcal{U}_\delta(E))}^2$$

from Proposition 11 with s replaced by $s/2$. To get the desired estimate (4.1), notice first that since γ is bounded, for any $\varepsilon > 0$

$$\int_\ell \gamma(x) dm_1(x) \leq \|\gamma\|_\infty m_1(\{\gamma \geq \varepsilon\} \cap \ell) + \varepsilon L$$

all line segments ℓ of length L . Thus, if γ satisfies (1.4), then for ε small enough, $\{\gamma \geq \varepsilon\}$ satisfies the GCC. Moreover, since γ is uniformly continuous, there exists $\delta > 0$ such that $|\gamma(y)| \geq \varepsilon/2$ for y in a δ -neighborhood of $\{\gamma \geq \varepsilon\}$. Thus, taking $E = \{\gamma \geq \varepsilon\}$,

$$\|u_2\|_{L^2(\mathcal{U}_\delta(E))} \leq 2\varepsilon^{-1} \|\gamma u_2\|_{L^2(\mathbb{R}^d)}.$$

Finally, using the triangle inequality and the fact that

$$\begin{aligned} \|\gamma u_2\|^2 &\leq C\langle \gamma u, u \rangle = -C \operatorname{Re} \langle (\mathcal{A}_\gamma - i\lambda)U, U \rangle_{H^{s/2} \times L^2} \\ &\leq \eta^{-1} \|(\mathcal{A}_\gamma - i\lambda)U\|_{H^{s/2} \times L^2}^2 + \eta \|U\|_{H^{s/2} \times L^2}^2 \end{aligned}$$

for any $\eta > 0$, we have

$$\begin{aligned} c\|U\|_{H^{s/2} \times L^2}^2 &\leq (1+|\lambda|)^{\frac{4}{s}-2} \|(\mathcal{A}_\gamma - i\lambda)U\|_{H^{s/2} \times L^2}^2 + ((1+|\lambda|)^{\frac{4}{s}-2} + 4\varepsilon^{-2}) \|\gamma u_2\|_{L^2}^2 \\ &\leq C \max\{(1+|\lambda|)^{\frac{4}{s}-2}, (1+|\lambda|)^{\frac{8}{s}-4}, 1\} \|(\mathcal{A}_\gamma - i\lambda)U\|_{H^{s/2} \times L^2}^2 + c/2 \|U\|_{H^{s/2} \times L^2}^2 \end{aligned}$$

by choosing η appropriately. This gives the desired estimate (4.1).

The converse is a consequence of only the exponential decay case, so we fix $s = 2$. By the Gearhart-Pruss Test, one has the resolvent estimate

$$\|U\|_{H^{s/2} \times L^2} \leq C \|(\mathcal{A}_\gamma - i\lambda)U\|_{H^{s/2} \times L^2}$$

for all $\lambda \in \mathbb{R}$ and all $U = (u_1, u_2) \in H^{s/2} \times L^2$. Taking $U = ((-\Delta + 1)^{-s/4}u, iu)$ for some $u \in L^2(\mathbb{R}^d)$, this implies

$$c\|u\|^2 \leq \|((-\Delta + 1)^{s/4} - \lambda)u\|^2 + \|\gamma u\|^2.$$

Now fix $\varkappa > 0$ to be small, and set $\tilde{A}_\lambda = \{\xi \in \mathbb{R}^d : |(|\xi|^2 + 1) - \lambda| \leq \varkappa\}$, if $\text{supp } \hat{u} \subset \tilde{A}_\lambda$, then

$$\|((-\Delta + 1)^{1/2} - \lambda)u\| \leq \varkappa\|u\|.$$

Consequently, if \varkappa is small enough, then there exists $c > 0$ such that for every $\lambda > 0$, $c\|u\| \leq \|\gamma u\|$ whenever $\text{spec}(u) \subset \tilde{A}_\lambda$. Since this inequality does not see modulation, the conclusion also holds with \tilde{A}_λ replaced by $\tilde{A}_\lambda + \xi$ for any $\xi \in \mathbb{R}^d$.

Next, we notice that inside \tilde{A}_λ , one can fit a rectangle with side lengths $c_0\varkappa \times c_0\lambda^{1/2} \times \dots \times c_0\lambda^{1/2}$ for some constant $c_0 > 0$. Letting $\lambda \rightarrow \infty$, we therefore see that $c\|u\| \leq \|\gamma u\|$ whenever $\beta_{d-1}(\text{spec}(u)) \leq c_0\varkappa$. But now, insofar as γ is bounded, we find $\varepsilon > 0$ such that $c\|u\| \leq \|\chi_{\{|\gamma| > 2\varepsilon\}}u\|$ whenever $\beta_{d-1}(\text{spec}(u)) \leq c_0\varkappa$. Employing Remark 8 (with $\beta = c_0\varkappa$), we see that the set $\{\gamma > \varepsilon\}$ satisfies the 1-GCC, from which we conclude that γ satisfies (1.4).

REFERENCES

- [1] Nalini Anantharaman and Matthieu Léautaud. Sharp polynomial decay rates for the damped wave equation on the torus, *Analysis & PDE* 7 (2014), no. 1, 159–214.
- [2] Sergei A. Avdonin and Sergei A. Ivanov. *Families of exponentials: the method of moments in controllability problems for distributed parameter systems*, volume 48 of *CBMS Regional Conference Series in Mathematics*. 1995.
- [3] Claude Bardos, Gilles Lebeau, and Jeffrey Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM J. Control Optim.*, 30(5):1024–1065, 1992.
- [4] Alexander Borichev and Yuri Tomilov. Optimal polynomial decay of functions and operator semigroups. *Mathematische Annalen*, 347(2):455–478, 2010.
- [5] Jean Bourgain and Semyon Dyatlov. Spectral gaps without the pressure condition. *Annals of Mathematics*, 187(3):825–867, 2018.
- [6] Nicolas Burq and Romain Joly. Exponential decay for the damped wave equation in unbounded domains. *Communications in Contemporary Mathematics*, 18(06):1650012, 2016.
- [7] Semyon Dyatlov and Joshua Zahl. Spectral gaps, additive energy, and a fractal uncertainty principle. *Geometric and Functional Analysis*, 26(4):1011–1094, 2016.
- [8] Michela Egidi and Ivan Veselić. Sharp geometric condition for null-controllability of the heat equation on \mathbb{R}^d and consistent estimates on the control cost. *Archiv der Mathematik*, 111(1):85–99, 2018.

- [9] Walton Green. On the energy decay rate of the fractional wave equation on \mathbb{R} with relatively dense damping. *arXiv preprint arXiv:1904.10946*, 2019.
- [10] Rui Han and Wilhelm Schlag. A higher dimensional Bourgain-Dyatlov fractal uncertainty principle. *arXiv preprint arXiv:1805.04994*, 2018.
- [11] Victor Havin and Burglind Jörnicke. *The uncertainty principle in harmonic analysis*, volume 28. Springer Science & Business Media, 2012.
- [12] Albert Edward Ingham. Some trigonometrical inequalities with applications to the theory of series. *Mathematische Zeitschrift*, 41(1):367–379, 1936.
- [13] Stéphane Jaffard. Contrôle interne exact des vibrations d’une plaque rectangulaire. *Portugaliae mathematica*, 47(4):423–429, 1990.
- [14] Stéphane Jaffard and Sorin Micu. Estimates of the constants in generalized ingham’s inequality and applications to the control of the wave equation. *Asymptotic Analysis*, 28(3, 4):181–214, 2001.
- [15] Benjamin Jaye and Mishko Mitkovski. Quantitative uniqueness properties for L^2 functions with fast decaying, or sparsely supported, Fourier transform. *arXiv preprint arXiv:1808.02149*, 2018.
- [16] Long Jin. Control for Schrödinger equation on hyperbolic surfaces. *Mathematical Research Letters*, 25(6):1865–1877, 2018.
- [17] Peter Jones. Rectifiable sets and the traveling salesman problem. *Invent. Math.*, 102(1):1–15, 1990.
- [18] Oleg Kovrijkine. Some results related to the Logvinenko-Sereda theorem. *Proceedings of the American Mathematical Society*, 129(10):3037–3047, 2001.
- [19] Gilles Lebeau and Iván Moyano. Spectral inequalities for the Schrödinger operator. *arXiv preprint arXiv:1901.03513*, 2019.
- [20] Vladimir Logvinenko and Yu. Sereda. Equivalent norms in spaces of entire functions of exponential type. *Teor. Funkcii Funkcional. Anal. i Priložen. Vyp.*, 20:102–111, 1974.
- [21] Satbir Malhi and Milena Stanislavova. When is the energy of the 1d damped Klein-Gordon equation decaying? *Mathematische Annalen*, 372(3-4):1459–1479, 2018.
- [22] Satbir Malhi and Milena Stanislavova. On the energy decay rates for the 1d damped fractional Klein-Gordon equation. *Mathematische Nachrichten*, 2019.
- [23] Fedor L. Nazarov. Local estimates for exponential polynomials and their applications to inequalities of the uncertainty principle type. *Algebra i analiz*, 5(4):3–66, 1993.
- [24] Boris Petrovich Paneah. Some theorems of Paley–Wiener type. In *Doklady Akademii Nauk*, volume 138, pages 47–50. Russian Academy of Sciences, 1961.
- [25] Amnon Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44. Springer-Verlag, 1983.
- [26] Jan Prüss. On the spectrum of C_0 -semigroups. *Transactions of the American Mathematical Society*, 284(2):847–857, 1984.
- [27] Jeffrey Rauch, Michael Taylor, and Ralph Phillips. Exponential decay of solutions to hyperbolic equations in bounded domains. *Indiana university Mathematics journal*, 24(1):79–86, 1974.
- [28] David L. Russell. Nonharmonic Fourier series in the control theory of distributed parameter systems. *Journal of Mathematical Analysis and Applications*, 18(3):542–560, 1967.

- [29] Jared Wunsch, Periodic damping gives polynomial energy decay *Mathematical Research Letters*, 24(2):571–580, 2017.
- [30] Maciej Zworski. *Semiclassical analysis*, volume 138. American Mathematical Soc., 2012.

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, CLEMSON UNIVERSITY, CLEMSON, SC, USA

E-mail address: awgreen@clemson.edu

E-mail address: bjaye@clemson.edu

E-mail address: mmitkov@clemson.edu