

Semicontinuity of Singularity Invariants in Families of Formal Power Series

Gert-Martin Greuel and Gerhard Pfister

Abstract. The problem we are considering came up in connection with the classification of singularities in positive characteristic. Then it is important that certain invariants like the determinacy can be bounded simultaneously in families of formal power series parametrized by some algebraic variety. In contrast to the case of analytic or algebraic families, where such a bound is well known, the problem is rather subtle, since the modules defining the invariants are quasi-finite but not finite over the base space. In fact, in general the fibre dimension is not semicontinuous and the quasi-finite locus is not open. However, if we pass to the completed fibers in a family of rings or modules we can prove that their fiber dimension is semicontinuous under some mild conditions. We prove this in a rather general framework by introducing and using the completed and the Henselian tensor product, the proof being more involved as one might think. Finally we apply this to the Milnor number and the Tjurina number in families of hypersurfaces and complete intersections and to the determinacy in a family of ideals.

Mathematics Subject Classification (2000). 13B35, 13B40, 14A15, 14B05, 14B07.

Keywords. Formal power series, completed tensor product, Henselian tensor product, semicontinuity, Milnor number, Tjurina number, determinacy.

Table of Contents

Introduction	2
1. Quasi-finite modules and semicontinuity	3
1.1. The completed tensor product	3
1.2. Fiber and completed fiber	7
1.3. Semicontinuity over a 1-dimensional ring	13
1.4. Henselian rings and Henselian tensor product	17
1.5. Semicontinuity for algebraically presented modules	20
2. Singularity invariants	27
2.1. Isolated singularities and flatness	27

2.2. Milnor number and Tjurina number of hypersurface singularities	31
2.3. Determinacy of ideals	32
2.4. Tjurina number of complete intersection singularities	33
References	34

Introduction

In connection with the classification of singularities defined by formal power series over a field a fundamental invariant is the modality of the singularity (with respect to some equivalence relation like right or contact equivalence). To determine the modality one has to investigate adjacent singularities that appear in nearby fibers. This cannot be done by considering families over complete local rings but one has to consider families of power series parametrized by some algebraic variety in the neighbourhood of a given point. To determine potential adjacencies, an important tool is the semicontinuity of certain singularity invariants like, for example, the Milnor or the Tjurina number. Another basic question is if the determinacy of an ideal can be bounded by a semicontinuous invariant. In the complex analytic situation the answer to these questions is well known and positive, but for formal power series the problem is much more subtle as one might think at the first glance. This is mainly due to the fact that ideals or modules that define the invariants are quasi-finite but not finite over the base space.

The modality example shows that the questions treated in this paper are rather natural and appear in important applications. Moreover, the semicontinuity in general is a very basic property with numerous applications in many other contexts. Therefore we decided to choose a rather general framework with families of modules presented by matrices of power series and parametrized by the spectrum of some Noetherian ring. It is not difficult to see that the fiber dimension is in general not semicontinuous and that the quasi-finite locus is in general not open (in contrast the case of ring maps of finite type, where the quasi-finite locus is open by Zariski's Main Theorem, cf. Proposition 47), see Examples 19 and 20. It turns out that the situation is much more satisfactory if we consider not the fibers but the completed fibers and we prove the desired semicontinuity for the completed fiber dimension under some conditions on the family. To guarantee that the completed fibre families behave well under base change we introduce the notion of a (partial) completed tensor product and study its properties in sections 1.1 and 1.2.

Unfortunately, we cannot prove the semicontinuity of the completed fibre dimension in full generality. We prove it if either the base ring has dimension one (in section 1.3), or if the base ring is complete local containing a field, or if the presentation matrix has polynomials or algebraic power series as entries (in section 1.5). Together, these cases cover most applications (see Corollary 45 for a summary). To treat the latter case, we use Henselian rings and the Henselian tensor

product, for which we give a short account in section 1.4. It would be interesting to know, if the result holds for presentation matrices with arbitrary power series as entries or if there are counterexamples. In section 1.5 we consider also the case of families of finite type over the base ring and prove a version of Zariski's main theorem for modules. Moreover, we compare the completed fibre with the usual fibre.

In section 2 we apply our results to singularity invariants. We discuss and compare first the notions of regularity and smoothness (over a field) and show that both notions coincide for the completed fibres (Lemma 60). Under the restrictions mentioned above, we prove the semicontinuity of the Milnor number and Tjurina number for hypersurfaces (section 2.2) and the Tjurina number for complete intersections (section 2.4) as well as an upper bound for the determinacy of an ideal (section 2.3). Since the base ring may be the integers, our results are of some interest for computational purposes. For example, if a power series has integer coefficients then the Milnor number over the rationals is bounded by the Milnor number modulo just one (possibly unlucky) prime number if this is finite (see Corollary 63 and, more generally, Corollary 25 and Remark 26).

We assume all rings to be associative, commutative and with unit. Throughout the paper \mathbb{k} denotes an arbitrary field, A a ring, $R = A[[x]]$, $x = (x_1, \dots, x_n)$, the formal power series ring over A and M an R -module. For our main results we will assume that A is Noetherian and that M is finitely generated as R -module.

1. Quasi-finite modules and semicontinuity

1.1. The completed tensor product

Let A be a ring $R = A[[x]]$ and M an R -module. For any prime ideal \mathfrak{p} of A let $k(\mathfrak{p}) = A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$ be the residue field of \mathfrak{p} . $k(\mathfrak{p}) = \text{Quot}(A/\mathfrak{p})$ is the quotient field of A/\mathfrak{p} and hence $k(\mathfrak{p}) = A/\mathfrak{p}$ if \mathfrak{p} is a maximal ideal. We consider M via the canonical map $A \hookrightarrow R$ as A -module and set

$$M(\mathfrak{p}) := M_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) = (M \otimes_A A_{\mathfrak{p}}) \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) = M \otimes_A k(\mathfrak{p}),$$

which is called the *fiber* of M over \mathfrak{p} . $M(\mathfrak{p})$ is a vector space over $k(\mathfrak{p})$ and its dimension is denoted by

$$d_{\mathfrak{p}}(M) := \dim_{k(\mathfrak{p})} M(\mathfrak{p}).$$

M is called *quasi-finite*¹ over \mathfrak{p} if $d_{\mathfrak{p}}(M) < \infty$. We are interested in the behavior of $d_{\mathfrak{p}}(M)$ if \mathfrak{p} varies in $\text{Spec } A$, in particular in finding conditions under which $d_{\mathfrak{p}}(M)$ is semicontinuous on $\text{Spec } A$.

¹For $M = R/I$, I an ideal, this is the original definition of Grothendieck. Nowadays most authors (e.g. [Sta19]) require in addition that R is of finite type over A .

We say that a function $d : \text{Spec } A \rightarrow \mathbb{R}$, $\mathfrak{p} \mapsto d_{\mathfrak{p}}$, is (*upper*) *semicontinuous* at \mathfrak{p} if \mathfrak{p} has an open neighbourhood $U \subset \text{Spec } A$ such that $d_{\mathfrak{q}} \leq d_{\mathfrak{p}}$ for all $\mathfrak{q} \in U$. d is semicontinuous on $\text{Spec } A$ if it is semicontinuous at every $\mathfrak{p} \in \text{Spec } A$.

For finitely presented A -modules M the semicontinuity of $\mathfrak{p} \rightarrow d_{\mathfrak{p}}(M)$ is true and well known (cf. Lemma 1). However, in many applications M is not finitely generated over A but finite over some A -algebra R . Such a situation appears naturally in algebraic geometry, when one considers families of schemes or of coherent sheaves over $\text{Spec } A$. But then it is usually assumed that the ring R is either of (essentially) finite type over A (in algebraic geometry) or an analytic A -algebra (in complex analytic geometry). When we study families of singularities defined by formal power series (cf. Section 2), we have to consider $R = A[[x]]$, which is not of finite type over A . As far as we know, this situation has not been systematically studied and it leads to some perhaps unexpected results. For example, $d_{\mathfrak{p}}(M)$ is in general not semicontinuous on $\text{Spec } A$ (cf. Examples 19, 20).

It turns out that the situation is much more satisfactory if we pass from the usual fibres to the completed fibres, that is, we consider the *completed fiber dimension*

$$\hat{d}_{\mathfrak{p}}(M) := \dim_{k(\mathfrak{p})} M(\mathfrak{p})^{\wedge},$$

where $M(\mathfrak{p})^{\wedge}$ is the $\langle x \rangle$ -adic completion of the $R(\mathfrak{p})$ -module $M(\mathfrak{p})$. To guarantee that the completed fibres behave well when \mathfrak{p} varies in $\text{Spec } A$, we introduce the notion of a completed tensor product below.

For a finitely presented A -module M the semicontinuity of $\mathfrak{p} \rightarrow d_{\mathfrak{p}}(M)$ is well known:

Lemma 1. *If M is a finitely presented A -module then $d_{\mathfrak{p}}(M)$ is semicontinuous on $\text{Spec } A$. Moreover, if M is A -flat, then $d_{\mathfrak{p}}(M)$ is locally constant on $\text{Spec } A$.*

Proof. Fix $\mathfrak{p} \in \text{Spec } A$ and consider a presentation of M ,

$$A^p \xrightarrow{P} A^q \rightarrow M \rightarrow 0,$$

with matrix $P = (p_{ij})$, $p_{ij} \in A$. Applying $\otimes_A k(\mathfrak{p})$ to this sequence we get the exact sequence of vector spaces

$$k(\mathfrak{p})^p \xrightarrow{\overline{P}_{\mathfrak{p}}} k(\mathfrak{p})^q \rightarrow M(\mathfrak{p}) \rightarrow 0,$$

with entries of $\overline{P}_{\mathfrak{p}}$ being the images of p_{ij} in $k(\mathfrak{p})$. Then $d_{\mathfrak{p}}(M)$ is finite and $d_{\mathfrak{p}}(M) = q - \text{rank}(\overline{P}_{\mathfrak{p}})$. Since $\text{rank}(\overline{P}_{\mathfrak{p}}) \leq \text{rank}(\overline{P}_{\mathfrak{q}})$ for all \mathfrak{q} in some neighbourhood U of \mathfrak{p} , the claim follows.

If M is flat, then $M_{\mathfrak{p}}$ is free over the local ring $A_{\mathfrak{p}}$ for a given $\mathfrak{p} \in \text{Spec } A$. By [Mat86], Theorem 4.10 (ii) (and its proof) there exists an $f \notin \mathfrak{p}$ such that M_f is a free A_f -module of some rank r and hence $d_{\mathfrak{q}}(M) = r$ for \mathfrak{q} in the open neighbourhood $D(f)$ of \mathfrak{p} . \square

We introduce now the completed tensor product. Let us denote by

$$\langle x \rangle := \langle x_1, \dots, x_n \rangle_R$$

the ideal in R generated by x_1, \dots, x_n . More generally, if S is an R -algebra, then $\langle x \rangle_S$ denotes the ideal in S generated by (the images of) x_1, \dots, x_n .

For an R -module N denote by

$$N^\wedge := \varprojlim N/\langle x \rangle^m N$$

the $\langle x \rangle$ -adic completion of N . If N is also an S -module for some R -algebra S , then $\langle x \rangle^m N = (\langle x \rangle_S)^m N$, and hence the $\langle x \rangle$ -adic completion and the $\langle x \rangle_S$ -adic completion of N coincide.

Definition 2. Let A be a ring, $R = A[[x]]$, B an A -algebra and M an R -module. We define the completed tensor product of R and B over A as the ring

$$R \hat{\otimes}_A B := \varprojlim ((R/\langle x \rangle^m) \otimes_A B)$$

and the completed tensor product of M and B over A as the module

$$M \hat{\otimes}_A B := \varprojlim ((M/\langle x \rangle^m M) \otimes_A B).$$

If N is an A -module, we define the R -module

$$M \hat{\otimes}_A N := \varprojlim ((M/\langle x \rangle^m M) \otimes_A N)$$

and call it the completed tensor product of M and N over A .

One reason why we consider the completed tensor product is that it provides the right base change property in the category of rings of the form $A[[x]]$ by the following Proposition 3.1.

Proposition 3. *The completed tensor product has the following properties (assumptions as in Definition 2).*

1. $R \hat{\otimes}_A B = (R \otimes_A B)^\wedge = B[[x]]$.
2. $M \hat{\otimes}_A N = (M \otimes_A N)^\wedge$.
3. If M is finitely presented over R and N finitely presented over B , then

$$M \hat{\otimes}_A N \cong (M \otimes_A N) \otimes_{R \hat{\otimes}_A B} (R \hat{\otimes}_A B).$$

4. The canonical map $M \otimes_A N \rightarrow M \hat{\otimes}_A N$ is injective if A is Noetherian, M finite over R and N finite over A .
5. If $\langle x \rangle^m \subset \text{Ann}_R(M)$ for some m then $M \hat{\otimes}_A N = M \otimes_A N$ for every A -module N .

Proof. 1. We have $\varprojlim (A[[x]]/\langle x \rangle^m \otimes_A B) = \varprojlim (A[x]/\langle x \rangle^m \otimes_A B) = \varprojlim B[x]/\langle x \rangle^m = B[[x]]$, showing the second equality. The first follows since $(R/\langle x \rangle^m) \otimes_A B = (R \otimes_A B)/\langle x \rangle^m (R \otimes_A B)$.

2. Since $(M/\langle x \rangle^m M) \otimes_A N = (M \otimes_A N)/\langle x \rangle^m (M \otimes_A N)$ the equality follows and that $(M \otimes_A N)^\wedge$ is the $\langle x \rangle_R$ as well as the $\langle x \rangle_{R \hat{\otimes}_A B}$ -adic completion of $M \otimes_A N$.

3. If M resp. N are finitely presented over R resp. B , then $M \otimes_A N$ is finitely presented over $R \otimes_A B$. Hence we can apply (the proof of) [AM69, Proposition 10.13] and use 1. to show the isomorphism.

4. If A is Noetherian then R is Noetherian. If M is finitely generated over R and N finitely generated over A then $M \otimes_A N$ is finitely generated over R . The injectivity follows from 2. and [AM69, Theorem 10.17 and Corollary 10.19], since $\langle x \rangle$ is contained in the Jacobson radical of R by Lemma 16.

5. If $\langle x \rangle^m M = 0$ for some m , then $M \hat{\otimes}_A N = M \otimes_A N$ by definition of the completed tensor product. \square

Corollary 4. *The completed tensor product is right-exact on the category of finitely presented modules. That is, let*

$$\begin{aligned} M' \rightarrow M \rightarrow M'' \rightarrow 0, \text{ resp.} \\ N' \rightarrow N \rightarrow N'' \rightarrow 0 \end{aligned}$$

be exact sequences of finitely presented R -modules resp. B -modules. Then the sequences of finitely presented $R \hat{\otimes}_A B$ -modules

$$\begin{aligned} M' \hat{\otimes}_A N \rightarrow M \hat{\otimes}_A N \rightarrow M'' \hat{\otimes}_A N \rightarrow 0, \text{ resp.} \\ M \hat{\otimes}_A N' \rightarrow M \hat{\otimes}_A N \rightarrow M \hat{\otimes}_A N'' \rightarrow 0 \end{aligned}$$

are exact.

Proof. The sequences $M' \otimes_A N \rightarrow M \otimes_A N \rightarrow M'' \otimes_A N \rightarrow 0$ and $M \otimes_A N' \rightarrow M \otimes_A N \rightarrow M \otimes_A N'' \rightarrow 0$ are exact. Now tensor these sequences with $R \hat{\otimes}_A B$ over $R \otimes_A B$ and apply Proposition 3.3. \square

Corollary 5.

- (i) $R \hat{\otimes}_A A = R$.
- (ii) *If S is multiplicatively closed in A then $A[[x]] \hat{\otimes}_A (S^{-1}A) = (S^{-1}A)[[x]]$.*
- (iii) *For any R -module M we have $M \hat{\otimes}_A A = M^\wedge$.*
- (iv) *If M is finitely presented over R then $M = M^\wedge$. If moreover N is finitely presented over A , then $M \hat{\otimes}_A N = M \otimes_A N$.*

Proof. (i) and (ii) follow immediately from Proposition 3.1, (iii) is a special case of Proposition 3.2 and (iv) follows from (i) and Proposition 3.3 with $B = A$. \square

Applying Corollary 4 and Proposition 3.1 we get

Corollary 6. *If $A[[x]]^p \xrightarrow{T} A[[x]]^q \rightarrow M \rightarrow 0$ is an $A[[x]]$ -presentation of M and B an A -algebra, then*

$$M \hat{\otimes}_A B = \text{coker} (B[[x]]^p \xrightarrow{T} B[[x]]^q).$$

Remark 7. Let A be Noetherian and B an A -algebra. If $\langle x \rangle$ is contained in the Jacobson radical of $R \otimes_A B$, then $R \hat{\otimes}_A B$ is faithfully flat over $R \otimes_A B$, by Proposition 3.1 and [Mat86, Theorem 8.14]. Note however that although $\langle x \rangle$ is contained in the Jacobson radical of R by Lemma 16, it need not be in the Jacobson radical of $R \otimes_A B$ (cf. Example 14).

Example 8. Let $\langle f_1, \dots, f_k \rangle \subset R = A[[x]]$ be an ideal and $M = A[[x]]/\langle f_1, \dots, f_k \rangle$. If \mathfrak{p} is a prime ideal in A then $R \hat{\otimes}_A A_{\mathfrak{p}} = A_{\mathfrak{p}}[[x]]$ and from Corollary 6 we get $M \hat{\otimes}_A A_{\mathfrak{p}} = A_{\mathfrak{p}}[[x]]/\langle f_1, \dots, f_k \rangle$. If $k(\mathfrak{p})$ is the residue field of A at \mathfrak{p} then $M \hat{\otimes}_A k(\mathfrak{p}) = k(\mathfrak{p})[[x]]/\langle f_1, \dots, f_k \rangle$, something what one expects as fiber of M over \mathfrak{p} . While $A_{\mathfrak{p}}[[x]]$ and $k(\mathfrak{p})[[x]]$ are nice local rings, the subrings $R \otimes_A A_{\mathfrak{p}} \subset A_{\mathfrak{p}}[[x]]$ and $R \otimes_A k(\mathfrak{p}) \subset k(\mathfrak{p})[[x]]$ are in general not local if \mathfrak{p} is not a maximal ideal (see Example 14).

Remark 9. Proposition 3.1 with $B = A[[y]]$, $y = (y_1, \dots, y_m)$, implies

$$A[[x]] \hat{\otimes}_A A[[y]] = A[[x, y]].$$

Now let A be Noetherian. If I resp. J are ideals in $A[[x]]$ resp. $A[[y]]$, we get from Corollary 4

$$A[[x]]/I \hat{\otimes}_A A[[y]]/J = A[[x, y]]/\langle I, J \rangle_{A[[x, y]]}.$$

We call an A -algebra a *formal A -algebra* if it is isomorphic to an A -algebra $A[[x]]/I$. For two formal A -algebras $B = A[[x]]/I$ and $C = A[[y]]/J$ the completed tensor product can be defined as $B \hat{\otimes}_A C = A[[x, y]]/\langle I, J \rangle_{A[[x, y]]}$. It has the usual universal property of the tensor product in the category of formal A -algebras, analogous to the analytic tensor product for analytic algebras (cf. [GR71, Chapter III.5]). Thus, Definition 2 generalizes the completed tensor product of formal A -algebras.

1.2. Fiber and completed fiber

Let again A be a ring and M an $R = A[[x]]$ -module. We introduce the completed fiber $\hat{M}(\mathfrak{p})$ and the completed fiber dimension $\hat{d}_{\mathfrak{p}}M$ of M for $\mathfrak{p} \in \text{Spec } A$ and compare it with the usual fiber $M(\mathfrak{p})$ and the usual fiber dimension $d_{\mathfrak{p}}M$.

At the end of this section we give examples, showing that semicontinuity of $d_{\mathfrak{p}}(M)$ does not hold in general on $\text{Spec } A$, even if $A = \mathbb{C}[t]$ or $A = \mathbb{Z}$ (Examples 19 and 20). However, we show in the next sections 1.3 and 1.5 that, under some conditions, semicontinuity holds for the completed fiber dimension $\hat{d}_{\mathfrak{p}}(M)$.

Notation 10. We have canonical maps

$$A \xrightarrow{j} R \xrightarrow{\pi} R/\langle x \rangle \xrightarrow{i} A,$$

with $i \circ \pi \circ j = id$ and for an ideal $I \subset R$ we set $\bar{I} := \pi(I)$. On the level of schemes we have the maps $\text{Spec } A \xrightarrow{i^*} V(\langle x \rangle) \xrightarrow{\pi^*} \text{Spec } R \xrightarrow{j^*} \text{Spec } A$, with $i^*(\mathfrak{p}) = \langle \mathfrak{p}, x \rangle$, $j^*(\langle \mathfrak{p}, x \rangle) = \langle \mathfrak{p}, x \rangle \cap A = \mathfrak{p}$ for $\mathfrak{p} \in \text{Spec } A$. We denote by

$$\mathfrak{n}_{\mathfrak{p}} := \langle \mathfrak{p}, x \rangle = \langle \mathfrak{p}, x_1, \dots, x_n \rangle_R$$

the ideal in R generated by $\mathfrak{p} \in \text{Spec } A$ and x_1, \dots, x_n . The family

$$f := j^* : \text{Spec } R \rightarrow \text{Spec } A$$

has the trivial section $\sigma = (i \circ \pi)^* : \text{Spec } A \rightarrow \text{Spec } R$, $\mathfrak{p} \mapsto \mathfrak{n}_{\mathfrak{p}}$, and the composition $h := \pi \circ j : A \cong R/\langle x \rangle$ induces an isomorphism

$$h^* : V(\langle x \rangle) \xrightarrow{\cong} \text{Spec } A,$$

the restriction of f to $V(\langle x \rangle)$. We call $R_{\mathfrak{p}} := R \otimes_A A_{\mathfrak{p}}$ the *stalk* of R over \mathfrak{p} . $R_{\mathfrak{p}}$ is not a local ring, its local ring at $\mathfrak{n}_{\mathfrak{p}}$ is $(R_{\mathfrak{p}})_{\mathfrak{n}_{\mathfrak{p}}} = R_{\mathfrak{n}_{\mathfrak{p}}}$ with residue field $k(\mathfrak{n}_{\mathfrak{p}}) = k(\mathfrak{p})$ (by Lemma 12).

If M is an R -module, we call $M_{\mathfrak{p}} = M \otimes_A A_{\mathfrak{p}}$ the *stalk* of M over \mathfrak{p} and we are interested in the behavior of M along the section σ . However, we are not interested in the $R(\mathfrak{p})$ -modules $M(\mathfrak{p})$ since $R(\mathfrak{p})$ is not a power series ring (and does not behave nicely). We are interested in the completed stalk $\hat{M}_{\mathfrak{p}}$ and in the completed fibers $\hat{M}(\mathfrak{p})$, which we introduce now.

Definition 11. *Let A be a ring, $R = A[[x]]$, M an R -module and $\mathfrak{p} \in \text{Spec } A$.*

1. *We set $\hat{R}_{\mathfrak{p}} := R \hat{\otimes}_A A_{\mathfrak{p}}$, a local ring isomorphic to $A_{\mathfrak{p}}[[x]]$ (Proposition 3.1), and call the $\hat{R}_{\mathfrak{p}}$ -module*

$$\hat{M}_{\mathfrak{p}} := M \hat{\otimes}_A A_{\mathfrak{p}}$$

the completed stalk of M over \mathfrak{p} .

2. *The ring $\hat{R}(\mathfrak{p}) := R \hat{\otimes}_A k(\mathfrak{p})$ is called the completed fiber of R over \mathfrak{p} . It is a local ring isomorphic to $k(\mathfrak{p})[[x]]$ (Proposition 3.1). The $\hat{R}(\mathfrak{p})$ -module*

$$\hat{M}(\mathfrak{p}) := M \hat{\otimes}_A k(\mathfrak{p}) = \hat{M}_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$$

is called the completed fiber of M over \mathfrak{p} .

3. *$\hat{M}(\mathfrak{p})$ is a $k(\mathfrak{p})$ -vector space and we call its dimension*

$$\hat{d}_{\mathfrak{p}}(M) := \dim_{k(\mathfrak{p})} \hat{M}(\mathfrak{p})$$

the completed fiber dimension of M over \mathfrak{p} .

4. *M is called quasi-completed-finite over \mathfrak{p} if $\hat{d}_{\mathfrak{p}}(M) < \infty$.*

The map $A \rightarrow R$ induces a map of local rings $A_{\mathfrak{p}} \rightarrow R_{\mathfrak{n}_{\mathfrak{p}}}$ and for an R -module M we have the fiber $M(\mathfrak{p}) = M \otimes_A k(\mathfrak{p})$ of M w.r.t. $A \rightarrow R$ and the fiber

$$M_{\mathfrak{n}_{\mathfrak{p}}}(\mathfrak{p}) = M_{\mathfrak{n}_{\mathfrak{p}}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) = M_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}M_{\mathfrak{n}_{\mathfrak{p}}}$$

of $M_{\mathfrak{n}_{\mathfrak{p}}}$ w.r.t. $A_{\mathfrak{p}} \rightarrow R_{\mathfrak{n}_{\mathfrak{p}}}$. The fibers are in general different but the completed fibers coincide by Lemma 15 if M is finitely R -presented.

Let us first compare the fiber $M(\mathfrak{p})$ with its completed fiber $\hat{M}(\mathfrak{p})$.

Lemma 12. *For any R -module M the following holds.*

- (i) $\hat{M}_{\mathfrak{p}} = (M_{\mathfrak{p}})^{\wedge}$ and $\hat{M}(\mathfrak{p}) = M(\mathfrak{p})^{\wedge}$.
- (ii) $\mathfrak{n}_{\mathfrak{p}}$ is a prime ideal in R with $\mathfrak{n}_{\mathfrak{p}} \cap A = \mathfrak{p}$ and the residue field of $\mathfrak{n}_{\mathfrak{p}}$ in R satisfies $k(\mathfrak{n}_{\mathfrak{p}}) = k(\mathfrak{p})$.
- (iii) If \mathfrak{n} is any prime ideal in R containing $\langle x \rangle$, then $\mathfrak{n} = \mathfrak{n}_{\mathfrak{p}}$ with $\mathfrak{p} = \mathfrak{n} \cap A \in \text{Spec } A$.

Proof. Statement (i) follows from Proposition 3.1. The first statement of (ii) follows since $R/\mathfrak{n}_{\mathfrak{p}} = A/\mathfrak{p}$ is an integral domain. Since $R/\mathfrak{n}_{\mathfrak{p}} = A/\mathfrak{p}$ we have $k(\mathfrak{n}_{\mathfrak{p}}) = \text{Quot}(R/\mathfrak{n}_{\mathfrak{p}}) = \text{Quot}(A/\mathfrak{p}) = k(\mathfrak{p})$. (iii) is obvious. \square

Remark 13. We have strict flat inclusions $A_{\mathfrak{p}} \subsetneq R_{\mathfrak{p}} \subsetneq R_{\mathfrak{n}_{\mathfrak{p}}} \subsetneq A_{\mathfrak{p}}[[x]]$ of rings that are Noetherian if A is Noetherian.

The strictness is easy to see. E.g. $g_0 + \sum_{|\alpha| \geq 1} (g_{\alpha}/h_{\alpha})x^{\alpha}$, $g_0 \notin \mathfrak{p}$, with arbitrary $h_{\alpha} \in R \setminus \mathfrak{n}_{\mathfrak{p}}$, is a unit in $A_{\mathfrak{p}}[[x]]$ but it is not contained in $R_{\mathfrak{n}_{\mathfrak{p}}}$, where only finitely many different denominators are allowed. We have $R_{\mathfrak{p}} = S^{-1}R$, with S the multiplicative set $A \setminus \mathfrak{p}$ and $R_{\mathfrak{n}_{\mathfrak{p}}} = (R_{\mathfrak{p}})_{\mathfrak{n}_{\mathfrak{p}}}$. Since localization preserves flatness ([AM69, Corollary 3.6]) and the Noether property ([AM69, Proposition 7.3]), the inclusions $A_{\mathfrak{p}} \subset R_{\mathfrak{p}} \subset R_{\mathfrak{n}_{\mathfrak{p}}}$ are flat and the rings are Noetherian if A is Noetherian. The flatness of $A_{\mathfrak{p}}[[x]]$ over $R_{\mathfrak{n}_{\mathfrak{p}}}$ follows, since the first is the $\langle x \rangle$ -adic completion of the second by Lemma 12 (i). Since both rings are local, $R_{\mathfrak{n}_{\mathfrak{p}}} \subset A_{\mathfrak{p}}[[x]]$ is faithfully flat.

The rings $R_{\mathfrak{p}}$ and $R_{\mathfrak{n}_{\mathfrak{p}}}$ are “strange” subrings of $A_{\mathfrak{p}}[[x]]$. The ring $A_{\mathfrak{p}}[[x]]$ is of interest in applications (cf. section 2), while the rings $R_{\mathfrak{p}}$ and $R_{\mathfrak{n}_{\mathfrak{p}}}$ are of minor interest. By the following Lemma 15 we have $(R_{\mathfrak{p}})^{\wedge} = (R_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge} = A_{\mathfrak{p}}[[x]]$.

Example 14. As an example let $A = \mathbb{k}[t]$ and $R = A[[x]]$ with t and x one variable, $\mathfrak{p} = \langle 0 \rangle \in \text{Spec } A$. We have $A_{\mathfrak{p}} = k(\mathfrak{p}) = \mathbb{k}(t)$ and

$$R_{\mathfrak{p}} = \mathbb{k}[t][[x]] \otimes_{\mathbb{k}[t]} \mathbb{k}(t) = \{g/h \mid g \in \mathbb{k}[t][[x]], h \in \mathbb{k}[t] \setminus 0\},$$

$g = g_0 + \sum_{i \geq 1} g_i x^i$, $g_i \in \mathbb{k}[t]$, a subring strictly contained in $R \hat{\otimes}_A A_{\mathfrak{p}} = \mathbb{k}(t)[[x]]$.

- $\langle x \rangle$ is contained in the Jacobson radical of $R \hat{\otimes}_A A_{\mathfrak{p}}$ by Lemma 16.
- $\langle x \rangle$ is not contained in the Jacobson radical of $R_{\mathfrak{p}} = R \otimes A_{\mathfrak{p}}$.

To see this, note that the element $t - x$ is a unit in $R \hat{\otimes}_A A_{\mathfrak{p}}$, since $1/(t - x) = 1/t \sum_{i \geq 0} (x/t)^i$, but $1/(t - x)$ is not an element in $R_{\mathfrak{p}}$. The ideal $\langle t - x \rangle$ is a maximal ideal in $R_{\mathfrak{p}}$, since $R_{\mathfrak{p}}/\langle t - x \rangle \cong \mathbb{k}((t))$ (see Example 19.2), but $x \notin \mathfrak{m}$ since otherwise $t \in \mathfrak{m}$, contradicting the fact that t is a unit in $R_{\mathfrak{p}}$.

- The rings $R_{\mathfrak{p}}$ and $R(\mathfrak{p})$ are in general not local.
Since $R_{\mathfrak{p}}/\langle x \rangle = \mathbb{k}(t)$, the ideals $\langle x \rangle$ and $\langle t - x \rangle$ are two different maximal ideals and $R_{\mathfrak{p}} = R(\mathfrak{p})$ ($\mathfrak{p} = \langle 0 \rangle$) is not local.

Lemma 15. *Let M be a finitely presented R -module and $\mathfrak{p} \in \text{Spec } A$.*

1. *We have isomorphisms*

$$\hat{M}_{\mathfrak{p}} \cong M_{\mathfrak{n}_{\mathfrak{p}}} \hat{\otimes}_{A_{\mathfrak{p}}} A_{\mathfrak{p}} = M_{\mathfrak{n}_{\mathfrak{p}}} \hat{\otimes}_A A = (M_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge}.$$

2. $\hat{M}(\mathfrak{p}) \cong (M_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}M_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge}$.
3. *If $M = \text{coker}(A[[x]]^p \xrightarrow{T} A[[x]]^q)$ then*

$$\begin{aligned} \hat{M}_{\mathfrak{p}} &= M \hat{\otimes}_A A_{\mathfrak{p}} = \text{coker}(A_{\mathfrak{p}}[[x]]^p \xrightarrow{T} A_{\mathfrak{p}}[[x]]^q), \\ \hat{M}(\mathfrak{p}) &= \text{coker}(k(\mathfrak{p})[[x]]^p \xrightarrow{T} k(\mathfrak{p})[[x]]^q). \end{aligned}$$

Note that $\hat{R}_{\mathfrak{p}} = A_{\mathfrak{p}}[[x]] = (R_{\mathfrak{p}})^{\wedge} \cong (R_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge}$ and $\hat{R}(\mathfrak{p}) = k(\mathfrak{p})[[x]] = R(\mathfrak{p})^{\wedge} \cong (R_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}R_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge}$ are local rings but $R_{\mathfrak{p}} \not\cong R_{\mathfrak{n}_{\mathfrak{p}}}$ and $R(\mathfrak{p}) \not\cong R_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}R_{\mathfrak{n}_{\mathfrak{p}}}$, since $R_{\mathfrak{p}}$ and $R(\mathfrak{p})$ are in general not local.

Proof. 1. The natural inclusion $R_{\mathfrak{n}_{\mathfrak{p}}} = A[[x]]_{\mathfrak{n}_{\mathfrak{p}}} \hookrightarrow A_{\mathfrak{p}}[[x]]$ is given as follows. Let $h/g \in R_{\mathfrak{n}_{\mathfrak{p}}}$ with $h, g \in R$, $g \notin \mathfrak{n}_{\mathfrak{p}}$ and write $g = g_0 - g_1$ with $g_0 \in A$ and $g_1 \in \langle x \rangle R$. Then $g \notin \mathfrak{n}_{\mathfrak{p}} = \langle \mathfrak{p}, x \rangle$ iff $g_0 \notin \mathfrak{p}$ and g is a unit in $R_{\mathfrak{n}_{\mathfrak{p}}}$ iff its image in $A_{\mathfrak{p}}[[x]]$ is a unit. We get

$$h/g = \frac{g_0^{-1}h}{(1 - g_1/g_0)} = g_0^{-1}h \sum_{i \geq 0} (g_1/g_0)^i \in A_{\mathfrak{p}}[[x]].$$

Now it is not difficult to see that the induced map $A[[x]]_{\mathfrak{n}_{\mathfrak{p}}}/\langle x \rangle^m \rightarrow A_{\mathfrak{p}}[[x]]/\langle x \rangle^m$ is bijective (a finite sum $\sum_{|\alpha|=0}^{m-1} (a_{\alpha}/b_{\alpha})x^{\alpha}$, $a_{\alpha}, b_{\alpha} \in A$, $b_{\alpha} \notin \mathfrak{p}$ in $A_{\mathfrak{p}}[[x]]$ can be written as $1/b \sum_{|\alpha|=0}^{m-1} (a_{\alpha}b'_{\alpha})x^{\alpha}$ with $b = \prod b_{\alpha} \notin \mathfrak{n}_{\mathfrak{p}}$, $b'_{\alpha} = b/b_{\alpha} \in A$, and hence is in $A[[x]]_{\mathfrak{n}_{\mathfrak{p}}}$). We get

$$R_{\mathfrak{n}_{\mathfrak{p}}} \hat{\otimes}_A A = \varprojlim A[[x]]_{\mathfrak{n}_{\mathfrak{p}}}/\langle x \rangle^m \otimes_A A = \varprojlim A_{\mathfrak{p}}[[x]]/\langle x \rangle^m = A_{\mathfrak{p}}[[x]]$$

and also $R_{\mathfrak{n}_{\mathfrak{p}}} \hat{\otimes}_{A_{\mathfrak{p}}} A_{\mathfrak{p}} = A_{\mathfrak{p}}[[x]] = R_{\mathfrak{n}_{\mathfrak{p}}}^{\wedge}$. Now apply Corollary 4 to the presentation of M and deduce the claim for $M \hat{\otimes}_A A_{\mathfrak{p}}$.

2. $\hat{M}(\mathfrak{p}) = M \hat{\otimes}_A (A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}) = (M \hat{\otimes}_A A_{\mathfrak{p}})/\mathfrak{p}(M \hat{\otimes}_A A_{\mathfrak{p}}) = M_{\mathfrak{n}_{\mathfrak{p}}}^{\wedge}/\mathfrak{p}M_{\mathfrak{n}_{\mathfrak{p}}}^{\wedge}$ by Corollary 5 (iv) and the first statement of this lemma. Since $M_{\mathfrak{n}_{\mathfrak{p}}}$ is finitely presented over $R_{\mathfrak{n}_{\mathfrak{p}}}$ we have $M_{\mathfrak{n}_{\mathfrak{p}}}^{\wedge} = M_{\mathfrak{n}_{\mathfrak{p}}} \otimes_{R_{\mathfrak{n}_{\mathfrak{p}}}} R_{\mathfrak{n}_{\mathfrak{p}}}^{\wedge}$, which implies the result.

3. This follows from Corollary 6. \square

Over maximal ideals the fiber and the completed fiber coincide:

Lemma 16. *Let A be Noetherian and M a finitely generated R -module. For $\mathfrak{a} \subset A$ a maximal ideal the following holds.*

(i)

$$\hat{M}(\mathfrak{a}) = M(\mathfrak{a}), \quad \hat{d}_{\mathfrak{a}}(M) = d_{\mathfrak{a}}(M).$$

(ii) $R/\mathfrak{a}R = R(\mathfrak{a}) = \hat{R}(\mathfrak{a}) = k(\mathfrak{a})[[x]]$ and $\mathfrak{a}R$ is a prime ideal in R .

(iii) $\mathfrak{n}_{\mathfrak{a}}$ is a maximal ideal of R and any maximal ideal of R is of the form $\mathfrak{n}_{\mathfrak{a}}$ for some $\mathfrak{a} \in \text{Max } A$. Hence $\langle x \rangle$ is contained in the Jacobson radical of R .

(iv) $M(\mathfrak{a}) = M/\mathfrak{a}M \cong M_{\mathfrak{n}_{\mathfrak{a}}}/\mathfrak{a}M_{\mathfrak{n}_{\mathfrak{a}}}$.

Proof. (i) Since \mathfrak{a} is maximal, $k(\mathfrak{a}) = A/\mathfrak{a}$ is a finite A -module. Corollary 5 (iv) implies $\hat{M}(\mathfrak{a}) = M \hat{\otimes}_A A/\mathfrak{a} = M \otimes_A A/\mathfrak{a} = M(\mathfrak{a})$.

(ii) This follows from (i) and the fact that $R/\mathfrak{a}R = k(\mathfrak{a})[[x]]$ is integral.

(iii) Cf. [Mat86, §1, Example 1] and [AM69, Chapter 1, Exercise 5].

(iv) $M/\mathfrak{a}M = M \otimes_A A/\mathfrak{a} = M(\mathfrak{a}) = \hat{M}(\mathfrak{a}) = M \hat{\otimes}_A A/\mathfrak{a} = \text{coker}(R_{\mathfrak{n}_{\mathfrak{a}}}^{\mathfrak{p}} \hat{\otimes}_A A/\mathfrak{a} \rightarrow R_{\mathfrak{n}_{\mathfrak{a}}}^{\mathfrak{q}} \hat{\otimes}_A A/\mathfrak{a}) = M_{\mathfrak{n}_{\mathfrak{a}}}/\mathfrak{a}M_{\mathfrak{n}_{\mathfrak{a}}}$, as in the proof of Lemma 15. \square

As a first step to semicontinuity we show below (Lemma 18) that the vanishing locus of $\hat{d}_{\mathfrak{p}}(M)$ is open. For an arbitrary A -module M

$$\text{Supp}_A(M) := \{\mathfrak{p} \in \text{Spec } A \mid M_{\mathfrak{p}} \neq 0\}$$

denotes the support of M and $\text{Ann}_A(M) = \{f \in A \mid fM = 0\}$ the annihilator ideal of M . In general $\text{Supp}_A(M)$ is not closed in $\text{Spec } A$, but if M a finitely generated A -module, then it is well known that $\text{Supp}_A(M) = V(\text{Ann}_A(M))$, which is closed in $\text{Spec } A$. More generally we have:

Remark 17. For any A -module M we have $\text{Supp}_A(M) \subset V(\text{Ann}_A(M))$. If R is an arbitrary A -algebra and M a finite R -module then $\text{Supp}_A(M) = V(\text{Ann}_A(M))$ and hence $\text{Supp}_A(M)$ is closed in $\text{Spec } A$.

To see this, let $\mathfrak{p} \in \text{Spec } A$. Note that $\mathfrak{p} \notin \text{Supp}_A M \Leftrightarrow M_{\mathfrak{p}} = 0 \Leftrightarrow \forall m \in M \exists f \in A, f \notin \mathfrak{p}, fm = 0$ and that $\mathfrak{p} \notin V(\text{Ann}_A(M)) \Leftrightarrow \text{Ann}_A(M) \not\subseteq \mathfrak{p} \Leftrightarrow \exists f \notin \mathfrak{p}, fM = 0$. Hence $\mathfrak{p} \notin V(\text{Ann}_A(M))$ implies $\mathfrak{p} \notin \text{Supp}_A(M)$, i.e. $\text{Supp}_A(M) \subset V(\text{Ann}_A(M))$.

Now let M be generated over R by $m_1, \dots, m_k \in M$. If $M_{\mathfrak{p}} = 0$, choose $f_i \in A, f_i \notin \mathfrak{p}, f_i m_i = 0$. Then $f = f_1 \cdots f_k \notin \mathfrak{p}$ satisfies $fM = 0$ and hence $f \in \text{Ann}_A(M)$. We get $\mathfrak{p} \notin V(\text{Ann}_A(M))$ and hence the other inclusion $\text{Supp}_A(M) \supset V(\text{Ann}_A(M))$.

In our situation for $R = A[[x]]$ and M finitely R -presented we have (possibly strict) inclusions (cf. Lemma 12)

$$\{\mathfrak{p} \in \text{Spec } A \mid \hat{d}_{\mathfrak{p}}(M) \neq 0\} \subset \{\mathfrak{p} \mid d_{\mathfrak{p}}(M) \neq 0\} \subset \text{Supp}_A(M),$$

where the first (Lemma 18) and the last (Remark 17) set are closed, while the middle set may not be closed (Example 19.4).

Lemma 18. *Let M be a finitely presented R -module. Then*

$$V(\langle x \rangle) \cap \text{Supp}_R(M) \xrightarrow{\sim} \{\mathfrak{p} \in \text{Spec } A \mid \hat{d}_{\mathfrak{p}}(M) \neq 0\}$$

and hence the vanishing locus of $\hat{d}_{\mathfrak{p}}(M)$, $\{\mathfrak{p} \in \text{Spec } A \mid \hat{d}_{\mathfrak{p}}(M) = 0\}$, is open in $\text{Spec } A$.

Proof. Let $A[[x]]^p \xrightarrow{T} A[[x]]^q \rightarrow M \rightarrow 0$ be a presentation of M , $T = (t_{ij})$, $t_{ij} \in A[[x]]$, and $\mathfrak{p} \in \text{Spec } A$. Then $k(\mathfrak{p})[[x]]^p \xrightarrow{T'} k(\mathfrak{p})[[x]]^q \rightarrow \hat{M}(\mathfrak{p}) \rightarrow 0$ is a presentation of $\hat{M}(\mathfrak{p})$ with $T' = (t'_{ij})$, $t'_{ij} \in k(\mathfrak{p})[[x]]$, the induced map (Corollary 6).

Now $\hat{M}(\mathfrak{p}) = 0$ iff T' is surjective, i.e., iff the 0-th Fitting ideal (the ideal of q -minors) of T' contains a unit $u' \in k(\mathfrak{p})[[x]]$. Write u' as $u' = u'_0 + u'_1$ with $u'_0 \in k(\mathfrak{p}) \setminus \{0\}$, $u'_1 \in \langle x \rangle k(\mathfrak{p})[[x]]$. Since Fitting ideals are compatible with base change, the 0-th Fitting ideal $F_0 \subset A[[x]]$ of M contains an element $u = u_0 + u_1 \in A[[x]]$ with $u_0 \in A$, $u_1 \in \langle x \rangle A[[x]]$, that maps to u' under $A[[x]] \rightarrow A_{\mathfrak{p}}[[x]] \rightarrow A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}[[x]]$. Hence u' is a unit iff $u_0 \notin \mathfrak{p}$, i.e., iff $\langle x, \mathfrak{p} \rangle \notin V(F_0 + \langle x \rangle)$. The result follows since $\text{Supp}_R(M) = V(F_0)$. \square

At the end of this section we give two examples where $d_{\mathfrak{p}}(M)$ is not semicontinuous on $\text{Spec } A$ while $\hat{d}_{\mathfrak{p}}(M)$ is. The examples show also that $\hat{M}(\mathfrak{p}) = 0$ may happen for $M(\mathfrak{p}) \neq 0$.

Example 19. Let $A = K[t]$, K an algebraically closed field, $R = A[[x]]$, and $M = R/\langle t - x \rangle$, with t and x one variable. The following properties illustrate the difference between the fibers and the completed fibers. Let $\mathfrak{a} = \langle t - c \rangle$, $c \in K$, denote the maximal ideals in A .

1. M is not finitely generated over A , $d_{\mathfrak{a}}(M) = \hat{d}_{\mathfrak{a}}(M) < \infty$ for $\mathfrak{a} \in \text{Max } A$ and $d_{\mathfrak{a}}(M)$ is semicontinuous on $\text{Max } A$:
 $M \cong K[[t]]$ as A -module via $f(x, t) \mapsto f(t, t)$, hence it is not finite over A .
 $M(\mathfrak{a}) \cong K[[t]]/\langle t - c \rangle$ and $d_{\mathfrak{a}}(M) = 1$ if $c = 0$ and 0 if $c \neq 0$.
2. $d_{\mathfrak{p}}(M)$ is not semicontinuous on $\text{Spec } A$:
 The prime ideal $\langle 0 \rangle$ is contained in every neighbourhood of $\mathfrak{a} = \langle t \rangle$ in $\text{Spec } A$. It satisfies $k(\langle 0 \rangle) = K(t)$ and we get $M(\langle 0 \rangle) \cong K[[t]] \otimes_A K(t) = K((t))$, the field of formal Laurent series. Since $\dim_{K(t)} K((t)) = \infty$, $d_{\langle 0 \rangle}(M) = \infty$, while $d_{\mathfrak{a}}(M) \leq 1$ for $\mathfrak{a} \in \text{Spec } A \setminus \langle 0 \rangle$.
3. $\hat{d}_{\mathfrak{p}}(M)$ is semicontinuous on $\text{Spec } A$:
 We have $\hat{M}(\langle 0 \rangle) = K(t)[[x]]/\langle t - x \rangle$ by Corollary 6. Since t is a unit in $K(t)$, $\hat{d}_{\langle 0 \rangle}(M) = 0$.
4. $M(\mathfrak{a}) = M_{\mathfrak{a}}/\mathfrak{a}M_{\mathfrak{a}} = 0$ does not imply $M_{\mathfrak{a}} = 0$:
 In fact, we have $M_{\langle t-c \rangle} \cong K[[t]]_{\langle t-c \rangle}$ (as $K[t]$ -module), which contains $K((t))$ if $c \neq 0$ (since $t \notin \langle t - c \rangle$) while $M(\langle t - c \rangle) = 0$ for $c \neq 0$ by 1. We have $\{\mathfrak{p} \mid \hat{d}_{\mathfrak{p}}(M) \neq 0\} = \{\langle t \rangle\} \subsetneq \{\mathfrak{p} \mid d_{\mathfrak{p}}(M) \neq 0\} = \{\langle t \rangle, \langle 0 \rangle\} \subsetneq \text{Supp}_A(M) = \text{Spec } A$.
5. M is flat over A . By 1. and 3. we cannot expect any continuity of $d_{\mathfrak{p}}(M)$ or $\hat{d}_{\mathfrak{p}}(M)$ on $\text{Max } A$ or on $\text{Spec } A$ for flat A -modules.
6. The quasi-finite locus of $A \rightarrow A[[x]]/\langle t - x \rangle$ is not open:
 The quasi-finite locus $\{\mathfrak{p} \in \text{Spec } A \mid d_{\mathfrak{p}}(M) < \infty\}$ is $\text{Spec } A \setminus \langle 0 \rangle$ by 1. and 2. Recall that if B is a ring of finite type over A , then the quasi-finite locus of $A \rightarrow B$ is open by Zariski's main Theorem (cf. [Sta19, 10.122]).
7. The quasi-completed-finite locus of $A \rightarrow A[[x]]/\langle t - x \rangle$ is open:
 Let us call $\{\mathfrak{p} \in \text{Spec } A \mid \hat{d}_{\mathfrak{p}}(M) < \infty\}$ the *quasi-completed-finite locus*. It is $\text{Spec } A$ in our example. If semicontinuity of $\hat{d}_{\mathfrak{p}}(M)$ holds (Corollary 45) then the quasi-completed-finite locus is open.

Example 20. The following example may be of interest for arithmetic and computational purposes. It goes along similar lines as Example 19.

Let $A = \mathbb{Z}$, $R = \mathbb{Z}[[x]]$, and $M = R/\langle x - p \rangle$, $p \in \mathbb{Z}$ a prime number. Since $R = \varprojlim \mathbb{Z}[x]/\langle x \rangle^n$ we obtain $M = \varprojlim \mathbb{Z}/p^n = \hat{\mathbb{Z}}_{\langle p \rangle}$, the ring of p -adic integers.

Now let $\langle q \rangle \in \text{Max } \mathbb{Z}$.

If $q \neq p$ then q is a unit in $\mathbb{Z}_{\langle p \rangle}$ hence in $\hat{\mathbb{Z}}_{\langle p \rangle}$ and $M \otimes_{\mathbb{Z}} \mathbb{Z}/q = M/\langle q \rangle M = \hat{\mathbb{Z}}_{\langle p \rangle}/q\hat{\mathbb{Z}}_{\langle p \rangle} = 0$.

If $q = p$ then $M \otimes_{\mathbb{Z}} \mathbb{Z}/p = \hat{\mathbb{Z}}_{\langle p \rangle} / p \hat{\mathbb{Z}}_{\langle p \rangle} = \mathbb{Z}/p$.

Hence $d_{\langle q \rangle}(M) = \dim_{\mathbb{Z}/q} M \otimes_{\mathbb{Z}} \mathbb{Z}/q$ is 0 if $q \neq p$ and 1 if $q = p$.

On the other hand, looking at the prime ideal $\langle 0 \rangle$ we get

$$\hat{M}(\langle 0 \rangle) = M \hat{\otimes}_{\mathbb{Z}} \mathbb{Q} = \mathbb{Q}[[x]] / \langle x - p \rangle = 0,$$

while

$$M(\langle 0 \rangle) = M \otimes_{\mathbb{Z}} \mathbb{Q} = \hat{\mathbb{Z}}_{\langle p \rangle} \otimes_{\mathbb{Z}} \mathbb{Q} = \text{Quot}(\hat{\mathbb{Z}}_{\langle p \rangle})$$

has dimension $d_{\langle 0 \rangle}(M) = \dim_{\mathbb{Q}} \text{Quot}(\hat{\mathbb{Z}}_{\langle p \rangle}) = \infty$.

To see the last equality in the formula for $M(\langle 0 \rangle)$ one checks that the following diagram has the universal property of the tensor product:

$$\begin{array}{ccc} \hat{\mathbb{Z}}_{\langle p \rangle} & \xrightarrow{i_1} & \text{Quot}(\hat{\mathbb{Z}}_{\langle p \rangle}) = (\hat{\mathbb{Z}}_{\langle p \rangle})_p \\ \uparrow i_2 & & \uparrow j_1 \\ \mathbb{Z} & \xrightarrow{j_2} & \mathbb{Q} \end{array} .$$

Here i_1, i_2 and j_2 are the canonical inclusions and j_1 is given as follows: if $\alpha, \beta \in \mathbb{Z}$, $p \nmid \beta$, then $j_1(\frac{\alpha}{p^m \beta}) = \frac{1}{p^m} \frac{\alpha}{\beta}$, $\frac{\alpha}{\beta} \in \hat{\mathbb{Z}}_{\langle p \rangle}$ since $p \nmid \beta$. The universality of the diagram is easily seen. If T is a \mathbb{Z} -algebra and $\phi : \hat{\mathbb{Z}}_{\langle p \rangle} \rightarrow T$ and $\psi : \mathbb{Q} \rightarrow T$ are \mathbb{Z} -algebra homomorphisms then the morphism $\sigma : (\hat{\mathbb{Z}}_{\langle p \rangle})_p \rightarrow T$, given as $\sigma(\alpha/p^m) = \phi(\alpha)\psi(1/p^m)$, $p \nmid \alpha$ is the unique one, making the obvious diagram commutative.

1.3. Semicontinuity over a 1-dimensional ring

In this section A will be Noetherian and M a finitely generated R -module (except we say otherwise). Then $R = A[[x]]$ is Noetherian and M is finitely presented as R -module. At the moment we can prove the semicontinuity of $d_{\mathfrak{q}}(M)$ on $\text{Spec } A$ in full generality only under certain assumptions on the irreducible components of $\text{Supp}_R(M)$. This includes the case $\dim A = 1$ where $\dim A$ denotes the Krull dimension of A . The case of arbitrary Noetherian A is treated in the next section under the assumption that the presentation matrix of M is polynomial.

In an important special case semicontinuity holds for arbitrary A :

Proposition 21. *Let A be Noetherian and M a finitely generated R -module. If $\text{Supp}_R(M) \subset V(\langle x \rangle)$ then M is finitely generated over A and $\hat{M}(\mathfrak{q}) = M(\mathfrak{q})$ for all $\mathfrak{q} \in \text{Spec } A$. In particular semicontinuity of $\hat{d}_{\mathfrak{p}}(M) = d_{\mathfrak{p}}(M)$ holds at any $\mathfrak{p} \in \text{Spec } A$.*

Proof. Since $V(\text{Ann}_R(M)) = \text{Supp}_R(M) \subset V(\langle x \rangle)$, we have $\langle x \rangle \subset \sqrt{\text{Ann}_R(M)}$ and there exists an m such that $\langle x \rangle^m \subset \text{Ann}_R(M)$. We get a surjection

$$A[[x]] / \langle x \rangle^m \rightarrow R / \text{Ann}_R(M)$$

and since $A[[x]] / \langle x \rangle^m$ is finitely generated over A this holds for $R / \text{Ann}_R(M)$ too. Since M is finitely generated over $R / \text{Ann}_R(M)$ it is finitely generated over A and

hence finitely presented. By Lemma 1 there there is an open neighborhood U of \mathfrak{p} in $\text{Spec } A$ such that $d_{\mathfrak{q}}(M) \leq d_{\mathfrak{p}}(M)$, $\mathfrak{q} \in U$. By Proposition 3.5, $\hat{M}(\mathfrak{q}) = M(\mathfrak{q})$ for all $\mathfrak{q} \in \text{Spec } A$, showing the claim. \square

Before we formulate the next result, we introduce some notations to be used throughout this section. Consider a minimal primary decomposition of $\text{Ann}_R(M)$,

$$\text{Ann}_R(M) = \bigcap_{i=1}^r Q_i \subset R.$$

Since M is finitely generated over R , $\text{Supp}_R(M) = V(\text{Ann}_R(M)) = \bigcup_{i=1}^r V(Q_i)$ and $\dim M = \dim \text{Supp}_R(M)$.

Let $P_1, \dots, P_s \subset R$ be the minimal associated primes of $\langle x \rangle$. Since they correspond via $h : A \cong R/\langle x \rangle$ to the minimal associated primes $\bar{P}_1, \dots, \bar{P}_s$ of A , we have $\dim V(P_j) \leq \dim A$.

Lemma 22. *For A Noetherian, M finitely generated over R and $\mathfrak{p} \in \text{Spec } A$ the following holds:*

1. *Let A' be the reduction of A , $R' = A'[[x]]$ and M' the R' -module $M \otimes_R R'$. Then $\hat{M}'(\mathfrak{p}) \cong \hat{M}(\mathfrak{p})$ and hence $\hat{d}_{\mathfrak{p}}(M') = \hat{d}_{\mathfrak{p}}(M)$.*
2. *Let $Q \subset R$ be an ideal. Then $\hat{d}_{\mathfrak{p}}(M/QM) \leq \hat{d}_{\mathfrak{p}}(M)$.*
3. *If $Q_i \not\subset \mathfrak{n}_{\mathfrak{p}}$ for some $1 \leq i \leq r$, then $\hat{d}_{\mathfrak{p}}(M) = \hat{d}_{\mathfrak{p}}(M/QM)$, with $Q = \bigcap_{j \neq i} Q_j$.*
4. *If $Q_i \subset \mathfrak{n}_{\mathfrak{p}}$ and $\dim V(Q_i) > \dim A_{\mathfrak{p}}$ for some $1 \leq i \leq r$, then $\hat{d}_{\mathfrak{p}}(M) = \infty$.*
5. *Let $U = \text{Spec } B \subset \text{Spec } A$ be an affine open neighbourhood of \mathfrak{p} and $M_B = M \otimes_A B$ the restriction of M to U . Then $\hat{M}_B(\mathfrak{q}) = \hat{M}(\mathfrak{q})$ for all $\mathfrak{q} \in U$.*

Proof. 1. Since $\mathfrak{p} \in \text{Spec } A$ contains the nilpotent elements, $A'/\mathfrak{p}' = A/\mathfrak{p}$, where \mathfrak{p}' is the image of \mathfrak{p} in A' , and hence the residue field does not change if we pass from A to A' . By Proposition 3.1 we have $\hat{R}'(\mathfrak{p}') = R' \hat{\otimes}_{A'} k(\mathfrak{p}) = k(\mathfrak{p})[[x]] = \hat{R}(\mathfrak{p})$. Consider a presentation $R^p \xrightarrow{T} R^q \rightarrow M \rightarrow 0$ of M . Applying $\otimes_R R'$, we get a presentation of M' , $R'^p \xrightarrow{T} R'^q \rightarrow M' \rightarrow 0$. Apply $\hat{\otimes}_A k(\mathfrak{p})$ to the first resp. $\hat{\otimes}_{A'} k(\mathfrak{p})$ to the second exact sequence above. The sequences stay exact by Corollary 4. Since $(\hat{R}(\mathfrak{p}))^k = (\hat{R}'(\mathfrak{p}'))^k$ it follows that the canonical morphism $M \rightarrow M'$ induces an isomorphism $\hat{M}(\mathfrak{p}) \cong \hat{M}'(\mathfrak{p})$.

2. Since $(M/QM) \hat{\otimes}_A k(\mathfrak{p}) = M \hat{\otimes}_A k(\mathfrak{p}) / Q(M \hat{\otimes}_A k(\mathfrak{p}))$ by Corollary 4, the result follows.

3. If $Q_i \not\subset \mathfrak{n}_{\mathfrak{p}}$, then $M_{\mathfrak{n}_{\mathfrak{p}}} = (M/Q)_{\mathfrak{n}_{\mathfrak{p}}}$ and the result follows from Lemma 15.

4. Set $\bar{R} := R/Q_i$ and $\bar{M} := M/Q_i M$. Then $\dim \bar{R}_{\mathfrak{n}_{\mathfrak{p}}} = \dim V(Q_i) > \dim A_{\mathfrak{p}}$ by assumption. Considering \bar{M} as R -module, we have $\bar{M} \hat{\otimes}_A k(\mathfrak{p}) = \bar{M}(\mathfrak{p})^{\wedge} = (\bar{M}_{\mathfrak{n}_{\mathfrak{p}}} / \mathfrak{p} \bar{M}_{\mathfrak{n}_{\mathfrak{p}}})^{\wedge}$ by Lemma 15. Since the $\bar{R}_{\mathfrak{n}_{\mathfrak{p}}}$ -modules $\bar{M}(\mathfrak{p})$ and its $\langle x \rangle$ -adic completion $\bar{M}(\mathfrak{p})^{\wedge}$ have the same Hilbert-Samuel function w.r.t. $\mathfrak{n}_{\mathfrak{p}}$, their dimension coincides (c.f. [GP08, Corollary 5.6.6]). Moreover, $\mathfrak{p} \bar{R}_{\mathfrak{n}_{\mathfrak{p}}}$ is the annihilator of $\bar{M}_{\mathfrak{n}_{\mathfrak{p}}} / \mathfrak{p} \bar{M}_{\mathfrak{n}_{\mathfrak{p}}}$ and therefore $\dim \bar{M}(\mathfrak{p})^{\wedge} = \dim \bar{R}_{\mathfrak{n}_{\mathfrak{p}}} / \mathfrak{p} \bar{R}_{\mathfrak{n}_{\mathfrak{p}}}$.

We apply now [Mat86, Theorem 15.1] to the map of local rings $A_{\mathfrak{p}} \rightarrow \bar{R}_{\mathfrak{p}}$ and get that $\dim \bar{R}_{\mathfrak{p}}/\mathfrak{p}\bar{R}_{\mathfrak{p}} \geq \dim \bar{R}_{\mathfrak{p}} - \dim A_{\mathfrak{p}} > 0$ and hence $\dim_{k(\mathfrak{p})} \bar{M}(\mathfrak{p})^\wedge = \infty$. Then $\hat{d}_{\mathfrak{p}}(M) = \dim_{k(\mathfrak{p})} M(\mathfrak{p})^\wedge = \infty$ by 2. of this lemma.

5. We may assume that $B = A_f$ for some $f \notin \mathfrak{p}$. Since $A_{\mathfrak{q}} = (A_f)_{\mathfrak{q}}$ for $\mathfrak{q} \in U = D(f)$, we have $k(\mathfrak{q}) = A_f \otimes_A k(\mathfrak{q})$. Now Proposition 3.1 implies $\hat{M}_{A_f}(\mathfrak{q}) = (M \otimes_A A_f) \hat{\otimes}_A k(\mathfrak{q}) = (M \otimes_A A_f \otimes_A k(\mathfrak{q}))^\wedge = M(\mathfrak{q})^\wedge = \hat{M}(\mathfrak{q})$. \square

Proposition 23. *Let A be Noetherian, M a finitely generated R -module and fix $\mathfrak{p} \in \text{Spec } A$. Let Q_1, \dots, Q_r be the primary components of $\text{Ann}_R(M)$ and P_1, \dots, P_s the minimal primes of $V(\langle x \rangle)$. After reenumeration we have two disjoint sets of primary components of $\text{Ann}_R(M)$ (any one may be empty):*

- I. Q_1, \dots, Q_k such that for each $1 \leq i \leq k$ there exists a j with $P_j \subset \sqrt{Q_i}$.
- II. Q_{k+1}, \dots, Q_r such that for each $k+1 \leq i \leq r$ we have $P_j \not\subset \sqrt{Q_i}$ for all j .

We set

$$\begin{aligned} Q_I &:= \bigcap_{i=1}^k Q_i, & Q_{II} &:= \bigcap_{i=k+1}^r Q_i, \\ M_I &:= M/Q_I M, & M_{II} &:= M/Q_{II} M. \end{aligned}$$

1. If case II does not occur (i.e., $k = r$), then there is an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$ such that $\hat{d}_{\mathfrak{q}}(M) \leq \hat{d}_{\mathfrak{p}}(M)$ for all prime ideals $\mathfrak{q} \in U$.
2. If $\dim V(\langle x \rangle) \cap V(Q_I) \cap V(Q_{II}) \leq 0$ then there is an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$ such that $\hat{d}_{\mathfrak{q}}(M) \leq \hat{d}_{\mathfrak{p}}(M)$ for all prime ideals $\mathfrak{q} \in U$.

Proof. Note that $V(Q_I) \subset V(\langle x \rangle)$ and $V(Q_i) \not\subset V(\langle x \rangle)$ for $k+1 \leq i \leq r$.

1. follows from Proposition 21, since the assumption says that $\text{Supp}(M) = \bigcup_{i=1}^r V(Q_i) \subset \bigcup_{i=1}^s V(P_i) = V(\langle x \rangle)$.

2. We renumber the P_j such that P_1, \dots, P_t satisfy $P_j \subset \sqrt{Q_i}$ for some $i = 1, \dots, k$ and P_{t+1}, \dots, P_s satisfy $P_j \not\subset \sqrt{Q_i}$ for any $i = 1, \dots, r$. We set

$$P_I := \bigcap_{j=1}^t P_j, \text{ satisfying } P_I \subset \sqrt{Q_I}.$$

We have $\text{Supp}_R(M_I) = V(Q_I) \subset V(P_I) \subset V(\langle x \rangle)$. By Proposition 21 there is an open neighborhood U_1 of \mathfrak{p} in $\text{Spec } A$ such that

$$\hat{d}_{\mathfrak{q}}(M_I) \leq \hat{d}_{\mathfrak{p}}(M_I), \quad \mathfrak{q} \in U_1. \quad (1)$$

We set

$$d := \dim R/(\text{Ann}_R(M) + \langle x \rangle)$$

and make induction on d . If $d = 0$ then $\dim \text{Supp}(M) \cap V(\langle x \rangle) = 0$ and \mathfrak{p} is an isolated point of $\{\mathfrak{q} \in \text{Spec } A \mid \hat{d}_{\mathfrak{q}}(M) \neq 0\}$ by Lemma 18 and the theorem is trivially true in this case. Hence we may assume that $d \geq 1$. By Lemma 22 we may assume that

- (α) A is reduced (Lemma 22 1.),
- (β) $Q_i \subset \mathfrak{n}_{\mathfrak{p}}$ for all $1 \leq i \leq r$ (Lemma 22 3.),

(γ) $\dim V(Q_i) \leq d$ for all $1 \leq i \leq r$ (Lemma 22 4., assuming (β)).

We have $\dim V(Q_{II}) \leq d$ by (γ). Since $P_j \not\subset \sqrt{Q_i}$ and since $V(Q_i)$ and $V(P_j)$ are irreducible we have for each $k+1 \leq i \leq r$ and for $t+1 \leq j \leq s$

$$\dim V(Q_i) \cap V(P_j) < \min\{\dim V(Q_i), \dim V(P_j)\}.$$

In particular $\dim V(\langle x \rangle) \cap V(Q_{II}) < d$ and $\dim R/\text{Ann}_R(M_{II}) + \langle x \rangle < d$ since $Q_{II} \subset \text{Ann}_R(M_{II})$.

By the induction hypothesis there exists an open neighbourhood $U_2 \subset \text{Spec } A$ of \mathfrak{p} such that

$$\hat{d}_{\mathfrak{q}}(M_{II}) \leq \hat{d}_{\mathfrak{p}}(M_{II}), \quad \mathfrak{q} \in U_2. \quad (2)$$

By Lemma 22 2. we have

$$\hat{d}_{\mathfrak{q}}(M_I), \hat{d}_{\mathfrak{q}}(M_{II}) \leq \hat{d}_{\mathfrak{q}}(M), \quad \mathfrak{q} \in \text{Spec } A. \quad (3)$$

Note that

$$\begin{aligned} M_{I,\mathfrak{q}} &= M_{\mathfrak{q}}, & \mathfrak{n}_{\mathfrak{q}} &\in V(Q_I) \setminus V(Q_I) \cap V(Q_{II}), \\ M_{II,\mathfrak{q}} &= M_{\mathfrak{q}}, & \mathfrak{n}_{\mathfrak{q}} &\in V(Q_{II}) \setminus V(Q_I) \cap V(Q_{II}). \end{aligned} \quad (4)$$

Using Notation 10, we set $V_1 = U_1 \cap V(\bar{Q}_I) \setminus (V(\bar{Q}_I) \cap V(\bar{Q}_{II}))$ and $V_2 = U_2 \cap V(\bar{Q}_{II}) \setminus (V(\bar{Q}_I) \cap V(\bar{Q}_{II}))$. From (1), (2), (3) and (4), we deduce

$$\begin{aligned} \hat{d}_{\mathfrak{q}}(M) &= \hat{d}_{\mathfrak{q}}(M_I) \leq \hat{d}_{\mathfrak{p}}(M), & \mathfrak{q} \in V_1, \\ \hat{d}_{\mathfrak{q}}(M) &= \hat{d}_{\mathfrak{q}}(M_{II}) \leq \hat{d}_{\mathfrak{p}}(M), & \mathfrak{q} \in V_2. \end{aligned} \quad (5)$$

The set $U := V_1 \cup V_2 \cup (V(\bar{Q}_I) \cap V(\bar{Q}_{II}) \cap (V_1 \cup V_2))$ is an open neighbourhood of \mathfrak{p} . In general we cannot compare $\hat{d}_{\mathfrak{p}}(M)$ with $\hat{d}_{\mathfrak{q}}(M)$ for $\mathfrak{q} \in V(\bar{Q}_I) \cap V(\bar{Q}_{II})$, except if this intersection consists just of \mathfrak{p} , which is the case if $V(\bar{Q}_I) \cap V(\bar{Q}_{II}) \cong V(\langle x \rangle) \cap V(Q_I) \cap V(Q_{II})$ has dimension 0. Then the result follows from (5). \square

As a corollary we get the following theorem, which was already proved for maximal ideals in $A = \mathbb{k}[t]$ in [GPh19].

Theorem 24. *Let A be Noetherian, M a finitely generated R -module and $\mathfrak{p} \in \text{Spec } A$. If $\dim A_{\mathfrak{p}} \leq 1$ then there is an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$ such that*

$$\hat{d}_{\mathfrak{q}}(M) \leq \hat{d}_{\mathfrak{p}}(M) \text{ for all } \mathfrak{q} \in U.$$

Proof. We may assume that $\hat{d}_{\mathfrak{p}}(M) < \infty$ and, as shown in the proof of Proposition 23, that $\mathfrak{n}_{\mathfrak{p}} \supset \text{Ann}_R(M) = Q_I \cap Q_{II}$. By Proposition 21, the result holds if case II. does not occur. If it occurs, then $\dim A_{\mathfrak{p}} \leq 1$ implies $\dim R/(\text{Ann}_R(M) + \langle x \rangle) \leq 1$ and $\dim V(\langle x \rangle) \cap V(Q_I) \cap V(Q_{II}) = 0$ as shown in the proof of Proposition 23 and the result follows. \square

Corollary 25. *Let $A = \mathbb{Z}$ and M be a finitely generated $\mathbb{Z}[[x]]$ -module, $x = (x_1 \cdots x_n)$, given by a presentation*

$$\mathbb{Z}[[x]]^r \rightarrow \mathbb{Z}[[x]]^s \rightarrow M \rightarrow 0.$$

Denote by

$$M_p := \hat{M}(\langle p \rangle) = \text{coker}(\mathbb{F}_p[[x]]^r \xrightarrow{\hat{T}} \mathbb{F}_p[[x]]^s)$$

if $p \in \mathbb{Z}$ is a prime number and by

$$M_0 := \hat{M}(\langle 0 \rangle) = \text{coker}(\mathbb{Q}[[x]]^r \xrightarrow{\hat{T}} \mathbb{Q}[[x]]^s)$$

the induced modules.

1. If $\dim_{\mathbb{F}_p} M_p < \infty$ then $\dim_{\mathbb{F}_p} M_p \geq \dim_{\mathbb{Q}} M_0$. Moreover, for all except finitely many prime numbers $q \in \mathbb{Z}$, $\dim_{\mathbb{F}_p} M_p \geq \dim_{\mathbb{F}_q} M_q$.
2. If $\dim_{\mathbb{Q}} M_0 < \infty$ then $\dim_{\mathbb{Q}} M_0 \geq \dim_{\mathbb{F}_q} M_q$ for all except finitely many prime numbers $q \in \mathbb{Z}$, and hence “ \geq ” for all except finitely many prime numbers $q \in \mathbb{Z}$.

The first part of statement 1. follows, since $\langle 0 \rangle$ is in every neighbourhood of p . In particular $\dim_{\mathbb{Q}} M_0$ is finite if $\dim_{\mathbb{F}_p} M_p$ is finite for some prime number p .

Remark 26. The corollary is important for practical computations in computer algebra systems. For simplicity let I be an ideal in $\mathbb{Z}[[x]]$ generated by polynomials, $M = \mathbb{Z}[[x]]/I$, and I_p the image of I in $\mathbb{F}_p[[x]]$. The dimension of $\mathbb{Q}[[x]]/I$ resp. of $\mathbb{F}_p[[x]]/I_p$, if finite, is equal to the dimension of $\mathbb{Q}[x]_{\langle x \rangle}/I$ resp. of $\mathbb{F}_p[x]_{\langle x \rangle}/I_p$. These dimensions can be computed in the localizations $\mathbb{Q}[x]_{\langle x \rangle}$ resp. $\mathbb{F}_p[x]_{\langle x \rangle}$ by computing a Groebner or standard basis of I resp I_p w.r.t. a local monomial ordering (cf. [GP08]). Such algorithms are implemented e.g. in SINGULAR [DGPS]. Usually the computations over \mathbb{Q} are very time consuming or do not finish, due to extreme coefficient growths, and therefore often modular methods are used. The above corollary says that for all except finally many prime numbers p we have equality $\dim_{\mathbb{Q}} \mathbb{Q}[x]_{\langle x \rangle}/I = \dim_{\mathbb{F}_p} \mathbb{F}_p[x]_{\langle x \rangle}/I_p$, and if this holds p is sometimes called a “lucky” prime. This fact can also be proved by Groebner basis methods. More interesting is however that $\dim_{\mathbb{Q}} \mathbb{Q}[x]_{\langle x \rangle}/I < \infty$ if there exists just one p (lucky or not) such that $\dim_{\mathbb{F}_p} \mathbb{F}_p[x]_{\langle x \rangle}/I_p < \infty$ and that the first dimension is bounded by the latter. This was stated in [Pf07] without proof.

1.4. Henselian rings and Henselian tensor product

In this section we recall some basic facts about Henselian rings and introduce similarly to the complete tensor product a Henselian tensor product. For details about Henselian rings see [Sta19] or [KPR75]. The Henselian tensor product is needed in Section 1.5 for algebraically presented modules. We start with some basic facts about étale ring maps.

- Definition 27.**
1. A ring map $\phi : A \rightarrow B$ is called *étale* if it is flat, unramified and of finite presentation.²
 2. ϕ is called *standard étale* if $B = (A[T]/F)_G$, $F, G \in A[T]$, the univariate polynomial ring, F monic and F' a unit in B .
 3. ϕ is called *étale at* $\mathfrak{q} \in \text{Spec}(B)$ if there exist $g \in B \setminus \mathfrak{q}$ such that $A \rightarrow B_g$ is étale.

The following proposition lists some basic properties of étale maps. The results can be found in section 10.142 of [Sta19].

- Proposition 28.**
1. The map $A \rightarrow A_f$ is étale.
 2. A standard étale map is an étale map.
 3. The composition of étale maps is étale.
 4. A base change of étale maps is étale.
 5. An étale map is open.
 6. An étale map is quasi-finite.
 7. Given $\phi : A \rightarrow B$ and $g_1, \dots, g_m \in B$ generating the unit ideal³ such that $A \rightarrow B_{g_i}$ is étale for all i then $\phi : A \rightarrow B$ is étale.
 8. Let $\phi : A \rightarrow B$ be étale then there exist $g_1, \dots, g_m \in B$ generating the unit ideal such that $A \rightarrow B_{g_i}$ is standard étale for all i .
 9. Let $S \subset A$ be a multiplicatively closed subset and assume that $\phi' : S^{-1}A \rightarrow B'$ is étale. Then there exist an étale map $\phi : A \rightarrow B$ such that $B' = S^{-1}B$ and $\phi' = S^{-1}\phi$.
 10. Let $\phi' : A/I \rightarrow B'$ be étale for some ideal $I \subset A$ then there exist an étale map $\phi : A \rightarrow B$ such that $B' = B/IB$ and the obvious diagram commutes.

Definition 29. Let A be a ring and $I \subset A$ an ideal. A is called *Henselian* with respect to I if the following holds⁴ (Univariate Implicit Function Theorem): Let $F \in A[T]$, the univariate polynomial ring, such that $F(0) \in I$ and $F'(0)$ is a unit modulo I then there exists $a \in I$ such⁵ that $F(a) = 0$.

Next we associate to any pair (A, I) , $I \subset A$ an ideal, the Henselization A_I^h , i.e. the "smallest" Henselian ring with respect to I , such that $A_I^h \subset \hat{A}_I = \varprojlim (A/I^n)$ the I -adic completion.

Definition 30.

1. Let A be a ring and $I \subset A$ an ideal. The ring

$$A_I^h = \varinjlim (B \mid A \rightarrow B \text{ an étale ring map inducing } A/I \cong B/IB)$$

is called the Henselization of A with respect to I .

² ϕ is unramified if it is of finite type and the module of Kähler differentials $\Omega_{B/A}$ vanishes. ϕ is of finite presentation if $B \cong A[x_1, \dots, x_n]/\langle f_1, \dots, f_m \rangle$ as A -algebras.

³ i.e., $\text{Spec}(A) = \cup D(g_i)$.

⁴Note, that (similarly to the I -adic completion) the condition implies that I is contained in the Jacobson radical of A . If we start with an ideal contained in the Jacobson radical then it is enough to consider monic polynomials F in the definition.

⁵Note that a is uniquely determined by the condition $a \in I$, [KPR75].

2. The Henselization of $A[x]$, A any ring, $x = (x_1, \dots, x_n)$, with respect to $I = \langle x \rangle$ is denoted by $A\langle x \rangle$. We call $A\langle x \rangle$ the ring of algebraic power series over A .

The Henselization has the following properties (cf. section 15.11 and 15.12 of [Sta19]):

Proposition 31. *Let A be ring and $I \subset A$ an ideal.*

1. A_I^h is Henselian with respect to $I^h = IA_I^h$ and $A/I^m = A_I^h/(I^h)^m$ for all m .
2. A is Henselian with respect to I if and only if $A = A_I^h$.
3. If A is Noetherian then the canonical map $A \rightarrow A_I^h$ is flat.
4. If A is Noetherian then the canonical map $A_I^h \rightarrow \hat{A}_I$ is faithfully flat and \hat{A}_I is the I^h -adic completion of A_I^h .

Remark 32. The definition of the Henselization implies that A_I^h is contained in the algebraic closure of A in \hat{A}_I . If A is excellent⁶ then A_I^h is the algebraic closure of A in \hat{A}_I . This is even true under milder conditions, see [KPR75]. In this situation $C\langle x \rangle$ is called the *ring of algebraic power series* of $C[[x]]$.

Next we prove a lemma which we need later in the applications.

Lemma 33. *Let A be a ring and $\mathfrak{p} \in \text{Spec}(A)$ a prime ideal. Let $C = A_{\mathfrak{p}}$, $I = \mathfrak{p}C$ and $f_1, \dots, f_m \in C_I^h$. Then there exist an étale map $A \rightarrow B$ such that*

1. $f_1, \dots, f_m \in B$,
2. there exists a prime ideal $\mathfrak{q} \in \text{Spec}(B)$ such that $\mathfrak{q} \cap A = \mathfrak{p}$.

Proof. By definition we have

$$C_I^h = \varinjlim (D | C \rightarrow D \text{ étale inducing } C/I = D/ID).$$

We choose D from the inductive system above such that $f_1, \dots, f_m \in D$. Since (C, I) is a local ring and $C/I = D/ID$, the ideal ID is a maximal ideal in D and we have $ID \cap C = I$. Using Proposition 28 (8) for the multiplicatively closed system $S = A \setminus \mathfrak{p}$ we find an étale map $A \rightarrow B'$ such that $D = S^{-1}B'$. This implies that $f_1, \dots, f_m \in B'_g$ for a suitable $g \in S$. Let $B = B'_g$ and $\mathfrak{q} = ID \cap B$ then $A \rightarrow B$ is étale having the properties 1. and 2. \square

Next we define the Henselization of an A -module M with respect to an ideal $I \subset A$ similarly to the definition of the Henselization of A with respect to I .

Definition 34. *Let A be a ring, $I \subset A$ an ideal and M an A -module. The module*

$$M_I^h = \varinjlim (M \otimes_A B | A \rightarrow B \text{ étale inducing } A/I = B/IB)$$

is called the Henselization of M with respect to I .

Lemma 35. $M_I^h = M \otimes_A A_I^h$.

⁶For the definition of excellence see 15.51 [Sta19].

Proof. The lemma follows since the direct limit commutes with the tensor product (cf. 10.75.2 [Sta19]). \square

Definition 36. Let A be a ring, $R = A\langle x \rangle$, B an A -algebra and M an R -module. We define the henselian tensor product of R and B over A as the ring

$$R \otimes_A^h B := \varinjlim (C \mid B[x] \longrightarrow C \text{ étale inducing } B = C/\langle x \rangle C = B\langle x \rangle).$$

$$M \otimes_A^h B := \varinjlim (M \otimes_A C \mid B[x] \longrightarrow C \text{ étale inducing } B = C/\langle x \rangle C = M \otimes_A B\langle x \rangle).$$

The Henselian tensor product has similar properties as the complete tensor product. Especially we obtain the following lemma.

Lemma 37. If $A\langle x \rangle^p \xrightarrow{T} A\langle x \rangle^q \rightarrow M \rightarrow 0$ is an $A\langle x \rangle$ -presentation of M then

$$M \otimes_A^h B = \text{coker} (B\langle x \rangle^p \xrightarrow{T} B\langle x \rangle^q).$$

In particular $R \otimes_A^h k(\mathfrak{p}) = k(\mathfrak{p})\langle x \rangle$ for $\mathfrak{p} \in \text{Spec } A$.

Definition 38. Let A be a ring, $R = A\langle x \rangle$ and M an R -module. We define for $\mathfrak{p} \in \text{Spec } A$ the $R \otimes_A^h k(\mathfrak{p}) = k(\mathfrak{p})\langle x \rangle$ -module

$$M^h(\mathfrak{p}) := M \otimes_A^h k(\mathfrak{p})$$

and call it the Henselian fiber of M over \mathfrak{p} . Moreover, we set

$$d_{\mathfrak{p}}^h(M) := \dim_{k(\mathfrak{p})} M^h(\mathfrak{p}).$$

1.5. Semicontinuity for algebraically presented modules

Let A be Noetherian and M finitely generated as $R = A[[x]]$ -module. Then M is finitely R -presented and in this section we assume that M has an algebraic presentation matrix. That is, there exists a presentation

$$R^p \xrightarrow{T} R^q \rightarrow M \rightarrow 0$$

with $T = (t_{ij})$ a $q \times p$ matrix such that $t_{ij} \in A\langle x \rangle$, $x = (x_1, \dots, x_n)$, the ring of algebraic power series over A (cf. Definition 30), e.g. $t_{ij} \in A[x]$. Under this assumption we shall prove the semicontinuity of $\hat{d}_{\mathfrak{b}}(M)$ for $\mathfrak{b} \in \text{Spec } A$.

We set $R_0 = A\langle x \rangle$ and $M_0 = \text{coker}(R_0^p \xrightarrow{T} R_0^q)$. Then using the $\langle x \rangle$ -adic completion we obtain $R_0^\wedge = R$ and $M_0^\wedge = M$.

Lemma 39. Let $B \supset A$ be an A -algebra, $\mathfrak{b} \in \text{Spec } B$ and $\mathfrak{a} = \mathfrak{b} \cap A$. Then

$$\hat{d}_{\mathfrak{a}}(M) < \infty \Leftrightarrow \hat{d}_{\mathfrak{b}}(M \hat{\otimes}_A B) < \infty \Leftrightarrow d_{\mathfrak{b}}^h(M_0 \otimes_A^h B) < \infty$$

and

$$\hat{d}_{\mathfrak{a}}(M) = \hat{d}_{\mathfrak{b}}(M \hat{\otimes}_A B) = d_{\mathfrak{b}}^h(M_0 \otimes_A^h B).$$

Proof. $M \hat{\otimes}_A B$ is considered as $R \hat{\otimes}_A B = B[[x]]$ -module and $M_0 \otimes_A^h B$ as $R_0 \otimes_A^h B = B\langle x \rangle$ -module. Therefore we have

$$\begin{aligned} \hat{d}_{\mathfrak{a}}(M) &= \dim_{k(\mathfrak{a})}(M \hat{\otimes}_A k(\mathfrak{a})) \\ \hat{d}_{\mathfrak{b}}(M \hat{\otimes}_A B) &= \dim_{k(\mathfrak{b})}(M \hat{\otimes}_A B \hat{\otimes}_B k(\mathfrak{b})) \\ d_{\mathfrak{b}}^h(M_0 \otimes_A^h B) &= \dim_{k(\mathfrak{b})}(M_0 \otimes_A^h B \otimes_B^h k(\mathfrak{b})) \end{aligned}$$

and

$$\begin{aligned} M \hat{\otimes}_A k(\mathfrak{a}) &= \operatorname{coker}(k(\mathfrak{a})[[x]]^p \xrightarrow{\bar{T}} k(\mathfrak{a})[[x]]^q) \\ M \hat{\otimes}_A B \hat{\otimes}_B k(\mathfrak{b}) &= \operatorname{coker}(k(\mathfrak{b})[[x]]^p \xrightarrow{\bar{T}} k(\mathfrak{b})[[x]]^q) \\ M_0 \otimes_A^h B \otimes_B^h k(\mathfrak{b}) &= \operatorname{coker}(k(\mathfrak{b})\langle x \rangle^p \xrightarrow{\bar{T}} k(\mathfrak{b})\langle x \rangle^q) \end{aligned}$$

with $\bar{T} = (\bar{t}_{ij})$ and \bar{t}_{ij} the induced elements in $k(\mathfrak{a})[x]$ resp. $k(\mathfrak{b})[x]$.

If $\hat{d}_{\mathfrak{b}}(M \hat{\otimes}_A B) < \infty$ there exists an N_0 such that $\langle x \rangle^N M \hat{\otimes}_A B \hat{\otimes}_B k(\mathfrak{b}) = 0$ for $N \geq N_0$ and hence

$$\begin{aligned} M \hat{\otimes}_A B \hat{\otimes}_B k(\mathfrak{b}) &= \operatorname{coker}(k(\mathfrak{b})[[x]]^p / \langle x \rangle^N \xrightarrow{\bar{T}} (k(\mathfrak{b})[[x]]^q / \langle x \rangle^N) \\ &= (\operatorname{coker}(k(\mathfrak{a})[[x]]^p / \langle x \rangle^N) \xrightarrow{\bar{T}} (k(\mathfrak{a})[[x]]^q / \langle x \rangle^N)) \otimes_{k(\mathfrak{a})} k(\mathfrak{b}). \end{aligned}$$

Since this hold for every $N \geq N_0$, we obtain $\hat{d}_{\mathfrak{a}}(M) < \infty$. Similarly we can see that $\hat{d}_{\mathfrak{a}}(M) < \infty$ implies $\hat{d}_{\mathfrak{b}}(M \otimes_A B) < \infty$ and in both cases we obtain $\hat{d}_{\mathfrak{a}}(M) = \hat{d}_{\mathfrak{b}}(M \otimes_A B)$. This gives the first equality in the Lemma. Since $B\langle x \rangle / \langle x \rangle^N = B[[x]] / \langle x \rangle^N$ we get the remaining claims. \square

Lemma 40. *Let $(A, \mathfrak{m}, \mathbb{k})$ be a local Noetherian Henselian ring and R a local quasi-finite (i.e. $\dim_{\mathbb{k}} R/\mathfrak{m}R < \infty$) and finite type A -algebra in the Henselian sense⁷. Then R is a finite A -algebra, i.e., finitely generated as A -module.*

Proof. This is an immediate consequence of Proposition 1.5 of [KPP78]. \square

Corollary 41. *Let $(A, \mathfrak{m}, \mathbb{k})$ be a local Noetherian Henselian ring and R a local and finite type A -algebra in the Henselian sense. If M is a finitely generated and quasi-finite (i.e. $\dim_{\mathbb{k}} M/\mathfrak{m}M < \infty$) R -module, then M is a finitely generated A -module.*

Proof. Passing from R to $R/\operatorname{Ann}_R(M)$ we may assume that $\operatorname{Ann}_R(M) = 0$. In this case $\dim_{\mathbb{k}} M/\mathfrak{m}M < \infty$ implies $\dim_{\mathbb{k}} R/\mathfrak{m}R < \infty$. Lemma 40 implies that R is a finitely generated A -module. Since M is finitely generated over R it follows that M is a finitely generated A -module. \square

Theorem 42. *Let A be a Noetherian ring, $R = A[[x]]$, $x = (x_1, \dots, x_n)$, and M a finitely generated R -module admitting a presentation*

$$R^p \xrightarrow{T} R^q \rightarrow M \rightarrow 0$$

⁷ R is an A -algebra of finite type in the Henselian sense if $R = A\langle t_1, \dots, t_s \rangle$ for suitable $t_1, \dots, t_s \in R$

with algebraic presentation matrix $T = (t_{ij})$, $t_{ij} \in A[x]$ or, more generally, $\in A\langle x \rangle$. Fix $\mathfrak{p} \in \text{Spec } A$ with $\hat{d}_{\mathfrak{p}}(M) < \infty$. Then there is an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$ such that

$$\hat{d}_{\mathfrak{q}}(M) \leq \hat{d}_{\mathfrak{p}}(M) \text{ for all } \mathfrak{q} \in U.$$

Proof. Recall that $R_0 = A\langle x \rangle$ is the Henselization of $A[x]$ with respect to $\langle x \rangle$ and $M_0 = \text{coker}(R_0^p \xrightarrow{T} R_0^p)$. Denote by A^h the henselization of the local ring $A_{\mathfrak{p}}$ with respect to its maximal ideal. We set $R^h := A^h\langle x \rangle$ and $M^h := \text{coker}((R^h)^p \xrightarrow{T} (R^h)^p) = M_0 \otimes_A^h R^h$. Then Lemma 39 implies $\hat{d}_{\mathfrak{p}}(M) = d_{\mathfrak{p}}^h(M^h)$ and Corollary 41 that M^h is a finitely generated A^h -module (R^h is a finite type A^h -algebra in the Henselian sense). Lemma 33 implies that there is an étale neighbourhood $\pi : \text{Spec } B \rightarrow \text{Spec } A$ of \mathfrak{p} such that $M_0 \otimes_A^h B = \text{coker}((R_0 \otimes_A^h B)^p \xrightarrow{T} (R_0 \otimes_A^h B)^p)$ is a finitely generated B -module and $M_0 \otimes_A^h B \otimes_B^h A^h = M^h$. Choose $\mathfrak{b} \in \text{Spec } B$ such that $\mathfrak{b} \cap A = \mathfrak{p}$. This is possible because of Lemma 33. Corollary 41 and Lemma 1 imply that there is an open neighbourhood $\tilde{U} \subset \text{Spec } B$ of \mathfrak{b} such that for $\mathfrak{c} \in \tilde{U}$ we have $d_{\mathfrak{c}}(M_0 \otimes_A^h B) \leq d_{\mathfrak{b}}(M_0 \otimes_A^h B)$. Since π is étale it is open (Proposition 28), $U := \pi(\tilde{U})$ is an open neighbourhood of \mathfrak{p} in $\text{Spec } A$ and for any $\mathfrak{q} \in U \cap \text{Spec } A$ there exists a $\mathfrak{c} \in \tilde{U} \cap \text{Spec } B$ with $\mathfrak{c} \cap A = \mathfrak{q}$. From Lemma 39 we obtain $\hat{d}_{\mathfrak{q}}(M) = d_{\mathfrak{c}}(M_0 \otimes_A^h B) \leq d_{\mathfrak{b}}(M_0 \otimes_A^h B) = \hat{d}_{\mathfrak{p}}(M)$. \square

The important property of Henselian local rings is that quasi-finite implies finite (in the sense of Corollary 41). Examples of Henselian local rings are quotient rings of the algebraic power series rings $A = \mathbb{k}\langle y \rangle / I$ over some field \mathbb{k} , and analytic \mathbb{k} -algebras.⁹

If A is a complete local ring containing a field, then any finitely generated R -module M can be polynomially presented and semicontinuity of $\hat{d}_{\mathfrak{p}}(M)$ holds, as we show now. We start with the following proposition, based on the Weierstrass division theorem.

Proposition 43. *Let $(A, \mathfrak{m}, \mathbb{k})$ be a Noetherian complete local ring containing \mathbb{k} , $R = A[[x]]$, $x = (x_1, \dots, x_n)$, and M a finitely generated R -module such that $\dim_{\mathbb{k}} M/\mathfrak{m}M < \infty$. Let $J = \text{Ann}_R(M)$. Then there exists f_1, \dots, f_s , $s \geq n$, with the following properties:*

1. $f_i \in A[x]$ for all i .
2. $f_{n-i+1} \in A[x_1, \dots, x_i]$ is a Weierstrass polynomial with respect to x_i for $i = 1, \dots, n$.
3. $J = \langle f_1, \dots, f_s \rangle A[[x]]$.

⁸Note that R^h is the Henselization of $A_{\mathfrak{p}}[[x]]$ with respect to the maximal ideal $\langle \mathfrak{p}, x \rangle$.

⁹An analytic \mathbb{k} -algebra is the quotient $\mathbb{k}\{y\}/I$, $y = (y_1, \dots, y_s)$, of a convergent power series ring over a complete real-valued field \mathbb{k} (cf. [GLS07]). E.g., if \mathbb{k} is any field with the trivial valuation, then $\mathbb{k}\{y\} = \mathbb{k}[[y]]$ is the formal power series ring; if $\mathbb{k} \in \{\mathbb{R}, \mathbb{C}\}$, then $\mathbb{k}\{y\}$ is the usual convergent power series ring.

Proof. To prove the statements we use induction on n , the number of the variables x . The assumption implies that $\dim_{\mathbb{k}}(R/J + \mathfrak{m}R) < \infty$, i.e. the ideal $J + \mathfrak{m}R$ is primary to the maximal ideal $\langle x \rangle + \mathfrak{m}R$ of R . This implies that $x_n^b \in J + \mathfrak{m}R$ for some b . Therefore there exists $g \in J$, $g = x_n^b + f$ with $f \in \mathfrak{m}R$. We know by Cohen's structure theorem that $A = \mathbb{k}[[y]]/I$ for suitable variables y and an ideal $I \subset \mathbb{k}[[y]]$. We can apply in the following the Weierstrass preparation and division theorem to representatives in $\mathbb{k}[[y, x]]$ and then take residue classes mod I . Obviously g is x_n -general. The Weierstrass preparation theorem implies $g = uh$, u a unit in R , and $h \in A[[x_1, \dots, x_{n-1}]][[x_n]]$ a Weierstrass polynomial with respect to x_n . To simplify notations we assume that g is already a Weierstrass polynomial with respect to x_n . Setting $R_0 = A[[x_1, \dots, x_{n-1}]]$, the Weierstrass division theorem (cf. [GLS07, Theorem I.1.8]) says that for any f in R there exist unique $h \in R$ and $r \in R_0[x_n]$ such that $f = hg + r$, $\deg_{x_n}(r) \leq b - 1$. In other words, as R_0 modules we have

$$R = R \cdot g \oplus R_0 \cdot x_n^{b-1} \oplus R_0 \cdot x_n^{b-2} \oplus \dots \oplus R_0. \quad (*)$$

We may thus assume that $J = \langle g_1, \dots, g_r \rangle$ with $g_1 = g$ and $g_i \in R_0[x_n]$ with $\deg_{x_n}(g_i) \leq b - 1$.

If $n = 1$ then $R_0 = A$ and the claim follows from (*). If $n \geq 2$ then M is a finitely generated R_0 -module since

- $R/\langle g \rangle$ is finite over R_0 and
- $g \in \text{Ann}_R(M)$, i.e. M is a finitely generated $R/\langle g \rangle$ -module.

Now let $J_0 = \text{Ann}_{R_0}(M)$. By induction hypothesis there are f_2, \dots, f_l , $l \geq n$, such that

1. $f_i \in A[x_1, \dots, x_{n-1}]$ for all i .
2. $f_{n-i+1} \in A[x_1, \dots, x_i]$ is a Weierstrass polynomial with respect to x_i for $i = 1, \dots, n - 1$.
3. $J_0 = \langle f_2, \dots, f_l \rangle R_0$.

Now denote by f_1 be the remainder of the division of g successively by f_2, \dots, f_n and by f_{l+i} the remainder of g_i by f_2, \dots, f_n for $i > 1$. These are polynomials in x_1, \dots, x_n . Then f_1, \dots, f_s satisfy the conditions 1. to 3. of the proposition. \square

Corollary 44. *Let $(A, \mathfrak{m}, \mathbb{k})$ be a Noetherian complete local ring containing \mathbb{k} , $R = A[[x]]$ and M a finitely generated R -module such that $\dim_{\mathbb{k}} M/\mathfrak{m}M < \infty$. Then M is polynomially presented.*

Proof. Assume M has a presentation matrix $T = (g_{ij})$, $g_{ij} \in A[[x]]$. Let $J = \text{Ann}_R(M)$. The assumption implies that $\dim_{\mathbb{k}} R/(J + \mathfrak{m}R) < \infty$. Using Proposition 43 we obtain that R/J is a A -finite and $J = \langle f_1, \dots, f_s \rangle$ with $f_{n-i+1} \in A[x_1, \dots, x_i]$ a Weierstrass polynomial with respect to x_i for $i = 1, \dots, n$, $n \leq s$. This implies that M has a presentation as R/J -module with presentation matrix T having entries in R/J . Now we can divide representatives in R of the entries of T successively by the Weierstrass polynomials f_{n-i+1} , $i = 1, \dots, n$. The remainders are polynomials in $A[x]$ representing the entries of T , which proves the claim. \square

Let us collect the cases for which we proved that semicontinuity of $\hat{d}_{\mathfrak{p}}(M)$ holds.

Corollary 45. *Let A be Noetherian and M a finitely generated R -module. Let $\mathfrak{p} \in \text{Spec } A$ and assume that one of the following conditions is satisfied:*

1. $\text{Supp}_R(M) \subset V(\langle x \rangle)$, or
2. $\dim A = 1$, or
3. M is algebraically presented, or
4. $(A, \mathfrak{m}, \mathbb{k})$ is a complete local ring containing a field¹⁰.

Then there is an open neighbourhood $U \subset \text{Spec } A$ of \mathfrak{p} such that $\hat{d}_{\mathfrak{q}}(M) \leq \hat{d}_{\mathfrak{p}}(M)$ for all $\mathfrak{q} \in U$. In particular, the quasi-completed-finite locus $\{\mathfrak{p} \in \text{Spec } A \mid \hat{d}_{\mathfrak{p}}(M) < \infty\}$ is open.

Proof. Statement 1. follows from Proposition 21, statement 2. from Theorem 24 and 3. from Theorem 42. Statement 4. follows from 3. and Corollary 44. \square

We do not know if semicontinuity of $\hat{d}_{\mathfrak{p}}(M)$ holds in general for A Noetherian of any dimension and M finitely but not necessarily algebraically presented.

Remark 46. For completeness we recall cases where semicontinuity of the usual fibre dimension $d_{\mathfrak{p}}(M)$ on $\text{Spec } A$ holds if M is an arbitrary finitely presented R -module, for different (local) rings A and R .

- $(A, \mathfrak{m}, \mathbb{k})$ local Noetherian Henselian, R a finite type A -algebra in the Henselian sense (by Corollary 41 and Lemma 1).
- $A = \mathbb{k}\{y\}/I$ an analytic \mathbb{k} -algebra and $R = \mathbb{k}\{y, x\}/J$ with $I\mathbb{k}\{y, x\} \subset J$ (by [GLS07, Theorem I.1.10]).
- A a Noetherian complete local ring containing a field, $R = A[[x]]$. This is a special case of the previous item. We mention it, since R is of the form considered in this paper.
- In the complex analytic situation with $A = \mathbb{C}\{y\}/I$, $y = (y_1, \dots, y_s)$, and $R = \mathbb{C}\{y, x\}/J$, $I\mathbb{C}\{y, x\} \subset J$, $x = (x_1, \dots, x_n)$, we have the following stronger statement: $A \rightarrow R$ induces a morphism of complex germs $f : (X, 0) \rightarrow (Y, 0)$, $(X, 0) = V(J) \subset (\mathbb{C}^n \times \mathbb{C}^s, 0)$, $(Y, 0) = V(I) \subset (\mathbb{C}^s, 0)$ and f the projection. For a sufficiently small representative $f : X \rightarrow Y$, M induces a coherent \mathcal{O}_X -module \mathcal{F} on X and $d_{\mathfrak{m}}(M) < \infty$, \mathfrak{m} the maximal ideal of $A = \mathcal{O}_{Y,0}$, means that the fibre dimension over $0 \in Y$ is finite, i.e. $d_0(\mathcal{F}) := \dim_{\mathbb{C}} \mathcal{F}_0/\mathfrak{m}\mathcal{F}_0 < \infty$. Then, for sufficiently small suitable X and Y , is $f|_{\text{Supp } \mathcal{F}}$ a finite morphism and $f_*\mathcal{F}$ is a coherent \mathcal{O}_Y -module (cf. [GLS07, Theorem I.1.67]). It follows that

$$d_y(\mathcal{F}) := \dim_{\mathbb{C}} f_*\mathcal{F} \otimes_{\mathcal{O}_{Y,y}} \mathbb{C} = \sum_{z \in f^{-1}(y)} \dim_{\mathbb{C}} \mathcal{F}_z/\mathfrak{m}_y\mathcal{F}_z$$

is upper semicontinuous at $0 \in Y$, i.e. 0 has an open neighbourhood $U \subset Y$ such that $d_y(\mathcal{F}) \leq d_0(\mathcal{F})$ for all $y \in U$.

¹⁰By Cohen's structure theorem this is equivalent to $A \cong \mathbb{k}[[y]]/I$, \mathbb{k} a field.

In the above cases M is finite over A if it is quasi-finite over A and hence semicontinuity of $d_{\mathfrak{p}}(M)$ holds by Lemma 1. Example 19 shows that for A an affine ring and $R = A[[x]]$ semicontinuity of $d_{\mathfrak{p}}(M)$ does in general not even hold for polynomially presented modules.

Instead of families of power series let us now consider families of algebras of finite type, a situation which is quite common in algebraic geometry. We treat the more general case of families of modules.

Let A be a ring, $R = A[x]/I$ of finite type over A and M a finitely presented R -module. M is called *quasi-finite at $\mathfrak{n} \in \text{Spec } R$ over A* if $\dim_{k(\mathfrak{p})} M_{\mathfrak{n}}/\mathfrak{p}M_{\mathfrak{n}} < \infty$ with $\mathfrak{p} \in \text{Spec } A$ lying under \mathfrak{n} . M is called *quasi-finite over $\mathfrak{p} \in \text{Spec } A$* if it is quasi-finite at all primes $\mathfrak{n} \in \text{Spec } R$ lying over \mathfrak{p} , and M is *quasi-finite over A* if it is quasi-finite at all primes $\mathfrak{n} \in \text{Spec } R$. The following proposition is a generalization of results from [Sta19], where the case of ring maps is treated.

Proposition 47. *Let A be a ring, R an A -algebra of finite type over A , M a finitely presented R -module and $f : \text{Spec } R \rightarrow \text{Spec } A$ the induced map of schemes.*

1. *The following are equivalent:*
 - (a) *M is quasi-finite over A ,*
 - (b) *$d_{\mathfrak{p}}(M) = \dim_{k(\mathfrak{p})} M(\mathfrak{p}) = \sum_{\mathfrak{n} \in f^{-1}(\mathfrak{p})} \dim_{k(\mathfrak{p})} M_{\mathfrak{n}}/\mathfrak{p}M_{\mathfrak{n}} < \infty \forall \mathfrak{p} \in \text{Spec } A$,*
 - (c) *The induced map $A \rightarrow S := R/\text{Ann}_R(M)$ is quasi-finite.*
2. *(Zariski's main theorem for modules). The quasi-finite locus of M*

$$\{\mathfrak{n} \in \text{Spec } R \mid M \text{ is quasi-finite at } \mathfrak{n}\}$$

is open in $\text{Spec } R$.

Proof. 1. (a) \Rightarrow (b): We have to show that the support of $M(\mathfrak{p})$ is finite. By [Sta19, Lemma 29.19.10], if $R = A[x]/I$ is a ring of finite type and quasi-finite over A , the induced map $f : \text{Spec } R \rightarrow \text{Spec } A$ has finite fibres $R(\mathfrak{p}) = R \otimes_A k(\mathfrak{p}) = k(\mathfrak{p})[x]/I(\mathfrak{p})$. It follows that 2. holds if M is a ring of finite type over A .

In the general case let $I = \text{Ann}_R(M)$. Then $S = R/I$ is of finite type over A , M is finitely presented over S and hence $\text{Supp}(S) = \text{Supp}(M)$. Moreover, let $J(\mathfrak{p})$ be the annihilator of the finitely generated $R(\mathfrak{p})$ -module $M(\mathfrak{p})$ satisfying $V(J(\mathfrak{p})) = \text{Supp}(M(\mathfrak{p}))$. Since $R(\mathfrak{p})$ is Noetherian and $\dim_{k(\mathfrak{p})} M_{\mathfrak{n}}(\mathfrak{p}) < \infty$ by assumption, we have $\mathfrak{n}^N M_{\mathfrak{n}}(\mathfrak{p}) = 0$ for some N by Nakayama's lemma. Hence $\mathfrak{n}^N \subset J(\mathfrak{p})R_{\mathfrak{n}}(\mathfrak{p})$ and $\dim_{k(\mathfrak{p})} R_{\mathfrak{n}}(\mathfrak{p})/J(\mathfrak{p})R_{\mathfrak{n}}(\mathfrak{p}) < \infty$. In general, the annihilator is not compatible with base change, hence $I(\mathfrak{p})$ is in general different from $J(\mathfrak{p})$. But for a finitely presented module the annihilator coincides up to radical with a Fitting ideal, and Fitting ideals are compatible with base change. It follows that $\sqrt{J(\mathfrak{p})R_{\mathfrak{n}}(\mathfrak{p})} = \sqrt{I(\mathfrak{p})R_{\mathfrak{n}}(\mathfrak{p})}$ and therefore $\dim_{k(\mathfrak{p})} R_{\mathfrak{n}}(\mathfrak{p})/I(\mathfrak{p})R_{\mathfrak{n}}(\mathfrak{p}) = \dim_{k(\mathfrak{p})} S_{\mathfrak{n}}(\mathfrak{p}) < \infty$, which means that $A \rightarrow S$ is quasi-finite at \mathfrak{n} ([Sta19, Definition 10.121.3]). Since this holds for each $\mathfrak{n} \in \text{Spec } R$, the map $A \rightarrow S$ is quasi-finite and by [Sta19, Lemma 29.19.10] the set $\text{Supp}(M(\mathfrak{p})) = \text{Supp}(S(\mathfrak{p}))$ is finite.

(b) \Rightarrow (c): If $d_{\mathfrak{p}}(M) < \infty$ for all $\mathfrak{p} \in \text{Spec } A$, then $\dim_{k(\mathfrak{p})} M_{\mathfrak{n}}/\mathfrak{p}M_{\mathfrak{n}} < \infty$ for all \mathfrak{n} and \mathfrak{p} under \mathfrak{n} . Then $A \rightarrow S$ is quasi-finite by the previous step.

(c) \Rightarrow (a): If $A \rightarrow S$ is quasi-finite then $\dim_{k(\mathfrak{p})} S_{\mathfrak{n}}(\mathfrak{p}) < \infty$ for all \mathfrak{p} and \mathfrak{n} over \mathfrak{p} . Since $M_{\mathfrak{n}}(\mathfrak{p})$ is finitely presented as $S_{\mathfrak{n}}(\mathfrak{p})$ -module, $\dim_{k(\mathfrak{p})} M_{\mathfrak{n}}(\mathfrak{p}) < \infty$ for all \mathfrak{p} and \mathfrak{n} over \mathfrak{p} and M is quasifinite over A .

2. In the first and third step of 1. we proved $\dim_{k(\mathfrak{p})} S_{\mathfrak{n}}(\mathfrak{p}) < \infty$ if and only if $\dim_{k(\mathfrak{p})} M_{\mathfrak{n}}(\mathfrak{p}) < \infty$, and hence M is quasi-finite at \mathfrak{n} iff $A \rightarrow S$ is quasi-finite at \mathfrak{n} . It follows from a version of Zarisk's main theorem as proved in [Sta19, Lemma 10.122.13] that the set $\{\mathfrak{n} \in \text{Spec } S \mid A \rightarrow S \text{ is quasi-finite at } \mathfrak{n}\}$ is open in $\text{Spec } S$ and thus of the form $U \cap S$ with U open in $\text{Spec } R$. If $\mathfrak{n} \in V = \text{Spec } R \setminus \text{Spec } S$ then $M_{\mathfrak{n}}(\mathfrak{p}) = 0$, hence M is quasi-finite at $\mathfrak{n} \in V$. Thus, the quasi-finite locus of M is the open set $U \cup V$. \square

Example 48. In the situation of Proposition 47, although the quasi-finite locus of M is open in $\text{Spec } R$, we cannot expect semicontinuity of $d_{\mathfrak{p}}(M)$ on $\text{Spec } A$. The following example shows that the vanishing locus of $d_{\mathfrak{p}}(M)$ is not open in $\text{Spec } A$. Let K be an algebraically closed field, $A = K[y]$, $R = A[x]$ and $M = R/\langle xy - 1 \rangle$. Then M is quasi-finite over A but $d_{\mathfrak{p}}(M)$ is not semicontinuous since $d_{\mathfrak{p}}(M) = 0$ if $\mathfrak{p} = \langle y \rangle$ and $d_{\mathfrak{p}}(M) = 1$ otherwise.

Problem 49. Let M be a finitely presented $R = A[[x]]$ -module and assume, moreover, that M is also finitely presented as A -module, in particular $d_{\mathfrak{p}}(M) < \infty$ for $\mathfrak{p} \in \text{Spec } A$. By Lemma 1 we have that $d_{\mathfrak{p}}(M)$ is semicontinuous on $\text{Spec } A$. The answer to the following questions is in general not known to us.

1. Is $\hat{d}_{\mathfrak{p}}(M)$ always semicontinuous on $\text{Spec } A$ (we know this only for $d_{\mathfrak{p}}(M)$)? The answer is positive under the assumptions of Corollary 45.
2. What is the relation between $\hat{d}_{\mathfrak{p}}(M)$ and $d_{\mathfrak{p}}(M)$? Is there always an inequality (c.f. Example 50)? The question is only interesting if \mathfrak{p} is not a maximal ideal, since we have $\hat{d}_{\mathfrak{p}}(M) = d_{\mathfrak{p}}(M)$ if $\mathfrak{p} \in \text{Max } A$ by Lemma 16.
3. Let (A, \mathfrak{m}) be a local ring and M a finitely presented A -module with $\text{Supp}(M)$ not contained in $V(\langle x \rangle)$ and $\text{Ann}_A(M) = 0$. Does this imply that A is \mathfrak{m} -adically complete? This is suggested by examples.

Example 50. A modification of Example 19 shows that $\hat{d}_{\mathfrak{p}}(M) \neq d_{\mathfrak{p}}(M)$ may happen for A a power series ring. Let $A = K[[t]]$, K an algebraically closed field, $R = A[[x]]$, and $M = R/\langle t - x \rangle \cong K[[t]]$. For the two prime ideals $\langle 0 \rangle$ and $\langle t \rangle$ of A we get:

$k(\langle 0 \rangle) = K((t))$, $k(\langle t \rangle) = K$, $M(\langle 0 \rangle) \cong K[[t]] \otimes_{K[[t]]} K((t)) = K((t))$, $M(\langle t \rangle) \cong K$ and $d_{\langle 0 \rangle}(M) = d_{\langle t \rangle}(M) = 1$. Hence $d_{\mathfrak{p}}(M)$ is semicontinuous (even continuous) on $\text{Spec } A$ as predicted in Remark 46, third item.

$\hat{M}(\langle 0 \rangle) \cong K((t))[[x]]/\langle t - x \rangle = 0$, $\hat{M}(\langle t \rangle) \cong K$ and $\hat{d}_{\langle 0 \rangle}(M) = 0$, $\hat{d}_{\langle t \rangle}(M) = 1$. Hence $d_{\mathfrak{p}}(M)$ is semicontinuous on $\text{Spec } A$ as predicted by Corollary 45. Note that M is finitely presented as A -module and we have $\hat{d}_{\langle 0 \rangle}(M) < d_{\langle 0 \rangle}(M)$.

2. Singularity invariants

2.1. Isolated singularities and flatness

Recall that a local Noetherian ring (A, \mathfrak{m}) is said to be *regular* if \mathfrak{m} can be generated by $\dim A$ elements. A Noetherian ring A is said to be regular if the local ring $A_{\mathfrak{p}}$ is regular for all $\mathfrak{p} \in \operatorname{Spec} A$. For arbitrary Noetherian rings the *regular locus* $\operatorname{Reg} A := \{\mathfrak{p} \in \operatorname{Spec} A \mid A_{\mathfrak{p}} \text{ is regular}\}$ need not be open in $\operatorname{Spec} A$. However, $\operatorname{Reg} A$ is open if A is a complete Noetherian local ring ([Mat86, Corollary of Theorem 30.10]) and the *non-regular locus* $\{\mathfrak{p} \in \operatorname{Spec} A \mid A_{\mathfrak{p}} \text{ is not regular}\}$ is closed.

However, in our situation of families of power series, the notion of formal smoothness is more appropriate than that of regularity. Formal smoothness is a relative notion and refers to a morphism, while regularity is an absolute property of the ring. The notions are related as follows. Let (A, \mathfrak{m}) be a local ring containing a field \mathbb{k} . If A is formally smooth over \mathbb{k} (w.r.t. the \mathfrak{m} -adic topology) then A is regular and the converse holds if the residue field A/\mathfrak{m} is separable over \mathbb{k} (see Remark 52). Hence formal smoothness of A over \mathbb{k} coincides with regularity if \mathbb{k} is a perfect field. The notions do also coincide for arbitrary \mathbb{k} if A is the quotient ring of a formal power series ring over \mathbb{k} by an ideal (cf. Lemma 57).

We recall now basic facts about formal smoothness. For details and proofs see [Mat86] and [Maj10].

Definition 51. *Let A be a ring, B an A -algebra defined by $\phi : A \rightarrow B$ and I an ideal in B . The A -algebra B is called formally smooth with respect to the I -adic topology (for short B is I -smooth over A) if for any A -algebra C and any continuous¹¹ A -algebra homomorphism $u : B \rightarrow C/N$, N an ideal in C with $N^2 = 0$, there exist $\sigma : B \rightarrow C$ such that $\pi\sigma = u$.*

$$\begin{array}{ccc}
 B & \xrightarrow{u} & C/N \\
 \uparrow \phi & \searrow \sigma & \uparrow \pi \\
 A & \xrightarrow{v} & C
 \end{array}$$

If $I = 0$ then B is called a formally smooth A -algebra.

Remark 52. 1. A formally smooth map of finite presentation is smooth ([Sta19] Proposition 10.137.13).

2. $A[x]$, $x = (x_1, \dots, x_n)$, is smooth over A ([Sta19] Lemma 10.137.4).

3. $A[[x]]$ is $\langle x \rangle$ -smooth over A ([Mat86] page 215).

4. Let (A, \mathfrak{m}) be a local ring containing a field \mathbb{k} .

(a) A is \mathfrak{m} -smooth over \mathbb{k} iff A is geometrically regular, i.e. $A \otimes_{\mathbb{k}} \mathbb{k}'$ is a regular ring for every finite extension field \mathbb{k}' of \mathbb{k} ([Mat86, Theorem 28.7]).

¹¹ Here we consider B with the I -adic topology and C/N with the discrete topology; u is continuous if $u(I^m) = 0$ for some m .

- (b) Assume that A/\mathfrak{m} is separable over \mathbb{k} . Then A is \mathfrak{m} -smooth over \mathbb{k} iff A is regular ([Mat86] Lemma 1, page 216).

We now generalize example 1 on page 215 of [Mat86].

Lemma 53. *Let A be a ring, B a A -algebra, I an ideal in B and \widehat{B} the I -adic completion of B . $\phi : A \rightarrow B$ is I -smooth iff $\widehat{\phi} : A \rightarrow \widehat{B}$ is $I\widehat{B}$ -smooth.*

Proof. Assume that B is I -smooth over A and consider the following commutative diagram:

$$\begin{array}{ccc} \widehat{B} & \xrightarrow{\hat{u}} & C/N \\ \hat{\phi} \uparrow & \searrow \hat{\sigma} & \uparrow \pi \\ A & \xrightarrow{v} & C \end{array}$$

with $N^2 = 0$. We have to prove that there exists $\hat{\sigma}$ such that $\pi\hat{\sigma} = \hat{u}$. Since \hat{u} is continuous there exist m such that $\hat{u}(I^m\widehat{B}) = 0$. Let $i : B \rightarrow \widehat{B}$ be the cononical map such that $\widehat{\phi} = i\phi$. The I -smoothness of B implies that there exists $\sigma : B \rightarrow C$ such that $\sigma\pi = \hat{u}i$. $\hat{u}(I^m\widehat{B}) = 0$ implies $\sigma(I^m) \subset N$. Since $N^2 = 0$ we obtain $\sigma(I^{2m}) = 0$. We obtain the following commutative diagram:

$$\begin{array}{ccccc} & & \widehat{B} & & \\ & & \downarrow \hat{i}_{2m} & & \\ B & \xrightarrow{i_{2m}} & B/I^{2m} & \xrightarrow{u_{2m}} & C/N \\ & \searrow \phi & & \searrow \sigma_{2m} & \uparrow \pi \\ & & A & \xrightarrow{v} & C \end{array}$$

Now we define $\hat{\sigma} = \sigma_{2m}\hat{i}_{2m}$. This proves that $\widehat{\phi} : A \rightarrow \widehat{B}$ is $I\widehat{B}$ -smooth.

Now assume that $\widehat{\phi} : A \rightarrow \widehat{B}$ is $I\widehat{B}$ -smooth. Consider the following commutative diagram:

$$\begin{array}{ccc} B & \xrightarrow{u} & C/N \\ \phi \uparrow & \searrow \sigma & \uparrow \pi \\ A & \xrightarrow{v} & C \end{array}$$

with $N^2 = 0$. We have to prove that there exists σ such that $\pi\sigma = u$. Since $\widehat{\phi} : A \rightarrow \widehat{B}$ is $I\widehat{B}$ -smooth there exists $\hat{\sigma} : \widehat{B} \rightarrow C$ with $\pi\hat{\sigma} = \hat{u}$. Now we define $\sigma = \hat{\sigma}i$ and obtain $\pi\sigma = u$. \square

The following important theorem is due to Grothendieck ([Mat86] Theorem 28.9).

Theorem 54. *Let (A, \mathfrak{m}) be a local ring and (B, \mathfrak{n}) a local A -algebra. Let $\bar{B} = B/\mathfrak{m}B$ and $\bar{\mathfrak{n}} = \mathfrak{n}/\mathfrak{m}B$. Then B is \mathfrak{n} -smooth over A iff \bar{B} is $\bar{\mathfrak{n}}$ -smooth over A/\mathfrak{m} and B is flat over A .*

Definition 55. *Let A be a ring and B an A -algebra defined by $\phi : A \rightarrow B$. We define the smooth locus of ϕ by*

$$\text{Sm}(\phi) := \{P \in \text{Spec}(B) \mid A_{\phi^{-1}(P)} \rightarrow B_P \text{ is } P\text{-smooth}\}.$$

and the singular locus of ϕ by

$$\text{Sing}(\phi) := \text{Spec}(B) \setminus \text{Sm}(\phi).$$

Remark 56. Let A be a ring and B a A -algebra defined by $\phi : A \rightarrow B$. The Theorem 54 of Grothendieck implies that

$$\begin{aligned} \text{Sm}(\phi) = \{ & P \in \text{Spec}(B) \mid A_Q \rightarrow B_P, Q = \phi^{-1}(P), \text{ is flat} \\ & \text{and } B_P/QB_P \text{ is } PB_P/QB_P\text{-smooth over } k(Q)\}. \end{aligned}$$

Now let \mathbb{k} be a field, $\mathbb{k}[[x]]$, $x = (x_1, \dots, x_n)$, the formal power series ring over \mathbb{k} and I an ideal in $\langle x \rangle \mathbb{k}[[x]]$. If I is generated by f_1, \dots, f_m we denote by $Jac(I)$ the Jacobian matrix $(\partial f_j / \partial x_i)$ and by $I_k(Jac(I))$ the ideal generated by the $k \times k$ -minors of $Jac(I)$ (which is independent of the chosen generators f_j). The following lemma gives equivalent conditions for the maximal ideal $\langle x \rangle \in B = \mathbb{k}[[x]]/I$ to be contained in the smooth locus $\text{Sm}(\phi)$ of the map $\phi : \mathbb{k} \rightarrow B$ (Remark 56).

Lemma 57. *If $\dim \mathbb{k}[[x]]/I = d$ the following are equivalent.*

1. $\mathbb{k}[[x]]/I$ is $\langle x \rangle$ -smooth over \mathbb{k} .
2. $\mathbb{k}[[x]]/I$ is regular.
3. $I_d(Jac(I)) = \mathbb{k}[[x]]$ (Jacobian criterion).
4. $\mathbb{k}[[x]]/I \cong \mathbb{k}[[y_1, \dots, y_d]]$.

Proof. The equivalence of 1. and 2. follows from [Mat86, Lemma 1, p. 216], the equivalence of 3. and 4. is the inverse mapping theorem for formal power series¹². Obviously 4. implies 2. From [Mat86, Theorem 29.7, p. 228 in] we deduce that 2. implies 4. \square

Remark 58. Part of the lemma can be generalized by extending the proof of Theorem 30.3 in [Mat86] as follows:

Let P be a prime ideal in $\mathbb{k}[[x]]$ containing $I = \langle f_1, \dots, f_m \rangle$ and \mathfrak{m} the maximal ideal of $A = \mathbb{k}[[x]]_P/I\mathbb{k}[[x]]_P$. Then $I_d(Jac(I)) = \mathbb{k}[[x]]_P$ implies that A is \mathfrak{m} -smooth over \mathbb{k} (or geometric regular by Remark 52).

We use the Jacobian criterion to define the singular locus of ideals in power series rings over a field.

¹²Given $f_1, \dots, f_n \in \mathbb{k}[[x_1, \dots, x_n]]$ then $\det(\frac{\partial f_i}{\partial x_j})$ is a unit iff $\mathbb{k}[[x_1, \dots, x_n]] = \mathbb{k}[[f_1, \dots, f_n]]$ ([GLS07, Theorem I.1.18]).

Definition 59. 1. If $B = \mathbb{k}[[x]]/I$ is pure d -dimensional (i.e. $\dim B/P = d$ for all minimal primes $P \in \text{Spec } B$) we define the singular locus of B (or of I) as

$$\text{Sing}(B) = V(I + I_d(\text{Jac}(I))).$$

2. If B is not pure dimensional we consider the minimal primes P_1, \dots, P_r of B . Then B/P_i is pure dimensional and we define the singular locus of B as

$$\text{Sing}(B) = \bigcup_{i=1}^r \text{Sing}(B/P_i) \cup \bigcup_{i \neq j} V(P_i) \cap V(P_j),$$

which is a closed subvariety of $\text{Spec } B$. The points in $\text{Spec } B \setminus \text{Sing}(B)$ are called non-singular points of B .

3. We say that $\mathbb{k}[[x]]/I$ (or I) has an isolated singularity (at 0) if the maximal ideal $\langle x \rangle$ is an isolated point of $\text{Sing}(\mathbb{k}[[x]]/I)$ or if $\langle x \rangle$ is a non-singular point.

Note that $\text{Sing}(B)$ carries a natural scheme structure given by the Fitting ideal $I + I_d(\text{Jac}(I)) \subset \mathbb{k}[[x]]$ if B is pure d -dimensional. In general we endow $\text{Sing}(B)$ with its reduced structure.

Now let us consider families. Let A be a Noetherian ring, $F_1, \dots, F_m \in \langle x \rangle A[[x]]$, $I \subset A[[x]]$ the ideal generated by F_1, \dots, F_m and set $B := A[[x]]/I$. We describe now the smooth locus of the map $\phi : A \rightarrow B$ along the section $\sigma : \text{Spec } A \rightarrow \text{Spec } B$, $\mathfrak{p} \mapsto \mathfrak{n}_{\mathfrak{p}} = \langle x, \mathfrak{p} \rangle$, of $\text{Spec } \phi$.

For $\mathfrak{p} \in \text{Spec } A$ denote by $F_i(\mathfrak{p})$ the image of F_i in $k(\mathfrak{p})[[x]]$. Note that $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$ generate the ideal $\hat{I}(\mathfrak{p}) \subset k(\mathfrak{p})[[x]]$, and that we have (by Lemma 15.3) for the completed fiber of ϕ over \mathfrak{p}

$$\hat{B}(\mathfrak{p}) = (B_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}B_{\mathfrak{n}_{\mathfrak{p}}})^\wedge = k(\mathfrak{p})[[x]]/\hat{I}(\mathfrak{p}).$$

The maximal ideals of the local rings of the fiber $B_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}B_{\mathfrak{n}_{\mathfrak{p}}}$ and the completed fiber $\hat{B}(\mathfrak{p})$ are generated by $\mathfrak{n}_{\mathfrak{p}}/\mathfrak{p} = \langle x \rangle$. Assume that $\phi : A \rightarrow B$ is flat. Then the theorem of Grothendieck says

$$\begin{aligned} \mathfrak{n}_{\mathfrak{p}} \in \text{Sm}(\phi) &\Leftrightarrow B_{\mathfrak{n}_{\mathfrak{p}}} \text{ is } \mathfrak{n}_{\mathfrak{p}}\text{-smooth over } A_{\mathfrak{p}} \\ &\Leftrightarrow B_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}B_{\mathfrak{n}_{\mathfrak{p}}} \text{ is } \langle x \rangle\text{-smooth over } k(\mathfrak{p}). \end{aligned}$$

Lemma 60. With the above notations assume that $\phi : A \rightarrow B$ is flat. Denote by

$$\text{Sing}_\sigma(\phi) := \{\mathfrak{n}_{\mathfrak{p}} \in \text{Spec } B \mid B_{\mathfrak{n}_{\mathfrak{p}}} \text{ is not } \mathfrak{n}_{\mathfrak{p}}\text{-smooth over } A_{\mathfrak{p}}\}$$

the singular locus of ϕ along the section σ . Then

$$\text{Sing}_\sigma(\phi) = \{\mathfrak{n}_{\mathfrak{p}} \in \text{Spec } B \mid \hat{B}(\mathfrak{p}) \text{ is not regular}\}.$$

Proof. $B_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}B_{\mathfrak{n}_{\mathfrak{p}}}$ is $\langle x \rangle$ -smooth over $k(\mathfrak{p})$ iff $(B_{\mathfrak{n}_{\mathfrak{p}}}/\mathfrak{p}B_{\mathfrak{n}_{\mathfrak{p}}})^\wedge = k(\mathfrak{p})[[x]]/\hat{I}(\mathfrak{p})$ is $\langle x \rangle$ -smooth over $k(\mathfrak{p})$ by Lemma 53. The claim follows from Lemma 57. \square

Since we assumed B to be flat over A , we have $\dim \hat{B}(\mathfrak{p}) = \dim B_{\mathfrak{n}_{\mathfrak{p}}} - \dim A_{\mathfrak{p}}$ (by [Mat86, Theorem 15.1]). If ϕ is of pure relative dimension d (i.e. $\hat{B}(\mathfrak{p})$ is pure d -dimensional for all \mathfrak{p}) then Lemma 57 implies

$$\text{Sing}_\sigma(\phi) = \{\mathfrak{n}_{\mathfrak{p}} \in \text{Spec } B \mid \hat{I}_d(\text{Jac}(I))(\mathfrak{p}) \text{ is a proper ideal of } k(\mathfrak{p})[[x]]\},$$

where $Jac(I)$ is the Jacobian matrix $(\partial F_j / \partial x_i)$ and $I_d(Jac(I)) \subset A[[x]]$ the ideal defined by the $d \times d$ -minors.

2.2. Milnor number and Tjurina number of hypersurface singularities

Let \mathbb{k} be a field and $f \in \mathbb{k}[[x]]$, $x = (x_1, \dots, x_n)$ a formal power series. The most important invariants are the *Milnor number* $\mu(f)$ and the *Tjurina number* $\tau(f)$, defined as

$$\begin{aligned}\mu(f) &= \dim_{\mathbb{k}} \mathbb{k}[[x]]/j(f), \\ \tau(f) &= \dim_{\mathbb{k}} \mathbb{k}[[x]]/\langle f, j(f) \rangle,\end{aligned}$$

where $j(f) = \langle \partial f / \partial x_1, \dots, \partial f / \partial x_n \rangle$ is the *Jacobian ideal* of f . We say that f has an *isolated critical point* (at 0) resp. an *isolated singularity* (at 0) if $\mu(f) < \infty$ resp. $\tau(f) < \infty$. Note that $\tau(f) < \infty$ iff $\mathbb{k}[[x]]/\langle f \rangle$ has an isolated singularity in the sense of Definition 59.

Remark 61. Let $\text{char}(\mathbb{k}) = 0$. It is proved in [BGM12, Theorem 2] that for $f \in \langle x \rangle$, $\mu(f) < \infty \Leftrightarrow \tau(f) < \infty$ but it is easy to see that this is not true in positive characteristic. We have always $\tau(f) \leq \mu(f)$ and $\tau(f) = \mu(f) \Leftrightarrow f \in j(f)$. If $\mathbb{k} = \mathbb{C}$ and if $f \in \langle x \rangle^2$ has an isolated singularity, this is equivalent to f being quasi homogeneous by a theorem of K. Saito (see [Sa71]). His proof generalises to any algebraically closed field of characteristic zero (cf. [BGM11, Theorem 2.1]).

We consider now families of singularities. Let A be a Noetherian ring and $F \in R = A[[x]]$. Set

$$j(F) := \langle \partial F / \partial x_1, \dots, \partial F / \partial x_n \rangle$$

and for $\mathfrak{p} \in \text{Spec } A$ denote by $F(\mathfrak{p})$ the image of F in $k(\mathfrak{p})[[x]]$. Then the Milnor number

$$\mu(F(\mathfrak{p})) = \dim_{k(\mathfrak{p})} k(\mathfrak{p})[[x]]/j(F(\mathfrak{p}))$$

and the Tjurina number

$$\tau(F(\mathfrak{p})) = \dim_{k(\mathfrak{p})} k(\mathfrak{p})[[x]]/\langle F(\mathfrak{p}), j(F(\mathfrak{p})) \rangle$$

are defined, and we deduce now the semicontinuity of $\mu(F(\mathfrak{p}))$ and $\tau(F(\mathfrak{p}))$.

Proposition 62. *Let A be Noetherian, $F \in R = A[[x]]$ and $\mathfrak{p} \in \text{Spec } A$. Assume that $V(j(F)) \subset V(\langle x \rangle)$ resp. $V(\langle F, j(F) \rangle) \subset V(\langle x \rangle)$ (as sets), or $\dim A = 1$, or $F \in A[x]$, or A is a complete local ring containing a field. Then $\mu(F(\mathfrak{p}))$ and $\tau(F(\mathfrak{p}))$ are semicontinuous at $\mathfrak{p} \in \text{Spec } A$.*

Proof. Set $M = R/j(F)$ resp. $M = R/\langle F, j(F) \rangle$, then $\text{Supp}_R(M) = V(j(F))$ resp. $\text{Supp}_R(M) = V(\langle F, j(F) \rangle)$. Using Lemma 15 we get $\hat{d}_{\mathfrak{q}}(M) = \mu(F(\mathfrak{q}))$ resp. $\hat{d}_{\mathfrak{q}}(M) = \tau(F(\mathfrak{q}))$ for $\mathfrak{q} \in \text{Spec } A$. The result follows from Corollary 45. \square

Corollary 63. *Let $F \in \mathbb{Z}[x]$, $p \in \mathbb{Z}$ a prime number and denote by F_p the image of F in $\mathbb{F}_p[[x]]$ and by F_0 the image of F in $\mathbb{Q}[[x]]$.*

If $\mu(F_p)$ is finite, then $\mu(F_p) \geq \mu(F_0)$ and $\mu(F_p) \geq \mu(F_q)$ for all except finitely many prime numbers $q \in \mathbb{Z}$. In particular, if $\mu(F_p)$ is finite for some p then $\mu(F_0)$ is finite.

If $\mu(F_0)$ is finite, then $\mu(F_0) \geq \mu(F_q)$ (and hence “=”) for all except finitely many prime numbers $q \in \mathbb{Z}$.

The same holds for the Tjurina number.

Example 64. We illustrate the corollary by a simple example. Let $F = F_0 = x^p + x^{(p+1)} + y^q$ with p, q prime numbers. Then $\mu(F_0) = (p-1)(q-1)$, $\mu(F_p) = p(q-1) \geq \mu(F_0)$ while $\mu(F_q) = \infty$. Moreover, for any prime number $r \neq p, q$ we have $\mu(F_r) = \mu(F_0)$.

2.3. Determinacy of ideals

Let I be a proper ideal of $\mathbb{k}[[x]]$ and f_1, \dots, f_m a minimal set of generators of I . I is called *contact k -determined* if for every ideal J of $\mathbb{k}[[x]]$ that can be generated by m elements g_1, \dots, g_m with $g_i - f_i \in \langle x \rangle^{k+1}$ for $i = 1, \dots, m$, the local \mathbb{k} -algebras $\mathbb{k}[[x]]/I$ and $\mathbb{k}[[x]]/J$ are isomorphic. I is called *finitely contact determined* if I is contact k -determined for some k . It is easy to see (cf. [GPh19, Proposition 4.3]) that these notions depend only on the ideal and not on the set of generators.

The ideal I or the ring $\mathbb{k}[[x]]/I$ is called a *complete intersection* if $\dim \mathbb{k}[[x]]/I = n - m$ and an *isolated complete intersection singularity (ICIS)* if it has moreover an isolated singularity.

Set $f = (f_1, \dots, f_m) \in \mathbb{k}[[x]]^m$ and denote by $\langle \partial f / \partial x_1, \dots, \partial f / \partial x_n \rangle$ the submodule of $\mathbb{k}[[x]]^m$, generated by the m -tuples $\partial f / \partial x_i = (\partial f_1 / \partial x_i, \dots, \partial f_m / \partial x_i)$, $i = 1, \dots, n$. We define

$$T_I := \mathbb{k}[[x]]^m / I\mathbb{k}[[x]]^m + \langle \partial f / \partial x_1, \dots, \partial f / \partial x_n \rangle.$$

If I is a complete intersection, then $\tau(I) := \dim_{\mathbb{k}} T_I$ is called the *Tjurina number of I* . For a complete intersection T_I is concentrated on the singular locus of $\mathbb{k}[[x]]/I$ (Definiton 59) and $\tau(I)$ is finite iff I has an isolated singularity. This follows from [GPh19, Lemma 3.1], where it is shown that the ideals $I + I_{n-m}(\text{Jac}(I))$ and $\text{Ann}_{\mathbb{k}[[x]]}(T_I)$ have the same radical.

The module T_I is used in the following theorem.

Theorem 65 ([GPh19], **Theorem 4.6**). *Let $I \subset \mathbb{k}[[x]]$ be a proper ideal and \mathbb{k} infinite. Then the following are equivalent:*

- (i) I is finitely contact determined.
- (ii) $\dim_{\mathbb{k}} T_I < \infty$.
- (iii) R/I is an isolated complete intersection singularity.

If one of these condition is satisfied then I is contact $(2 \dim_{\mathbb{k}} T_I - \text{ord}(I) + 2)$ -determined, where $\text{ord}(I) = \max\{k \mid I \subset \langle x \rangle^k\}$. The implications (iii) \Leftrightarrow (ii) \Rightarrow (i) hold for any field \mathbb{k} , as well as (i) \Rightarrow (ii) for hypersurfaces.

Proposition 66. *Let A be Noetherian, $F_1, \dots, F_m \in \langle x \rangle A[[x]]$. Let $I \subset A[[x]]$ be the ideal generated by F_1, \dots, F_m and $\hat{I}(\mathfrak{p}) \subset k(\mathfrak{p})[[x]]$, $\mathfrak{p} \in \text{Spec } A$, the ideal generated by $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p}) \in k(\mathfrak{p})[[x]]$.*

Assume that $V(I + I_{n-m}(\text{Jac}(I))) \subset V(\langle x \rangle)$ (as set), or $\dim A = 1$, or $F_i \in \langle x \rangle A[x]$ for $i = 1, \dots, m$, or A is a complete local ring containing a field.

Then any $\mathfrak{p} \in \text{Spec } A$ has an open neighbourhood $U \subset \text{Spec } A$ such that for all $\mathfrak{q} \in U$ $\dim_{k(\mathfrak{p})} T_{\hat{I}(\mathfrak{p})} \geq \dim_{k(\mathfrak{q})} T_{\hat{I}(\mathfrak{q})}$.

Proof. By [GPh19, Lemma 3.1] $\text{Supp}(T_I) = V(I + I_{n-m}(\text{Jac}(I)))$. The claim follows from Corollary 45. \square

2.4. Tjurina number of complete intersection singularities

We show first that being a regular sequence in a flat family of power series in $R = A[[x]]$ is an open property.

Proposition 67. *Let A be a Noetherian ring, $F_i \in \langle x \rangle R$, $i = 1, \dots, m$ and M a finitely generated R -module. For $\mathfrak{p} \in \text{Spec } A$ we denote by $F_i(\mathfrak{p})$ the image of F_i in $\hat{R}(\mathfrak{p}) = k(\mathfrak{p})[[x]]$ and by $F_{i, \mathfrak{p}}$ the image of F_i in $R_{\mathfrak{p}}(\mathfrak{p})$ (cf. Definition 11).*

- (i) *If $\mathfrak{p} \in \text{Spec } A$ then $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$ is an $\hat{M}(\mathfrak{p})$ -sequence iff $F_{1, \mathfrak{p}}, \dots, F_{m, \mathfrak{p}}$ is an $M_{\mathfrak{p}}(\mathfrak{p})$ -sequence.*
- (ii) *Let F_1, \dots, F_m be an M -sequence and let $M/\langle F_1, \dots, F_m \rangle M$ be A -flat. Then $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$ is an $\hat{M}(\mathfrak{p})$ -sequence for all $\mathfrak{p} \in \text{Spec } A$.*
- (iii) *Let $\mathfrak{p} \in \text{Spec } A$ and $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$ an $\hat{M}(\mathfrak{p})$ -sequence. If $M/\langle F_1, \dots, F_m \rangle M$ is flat over A , then there exists an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$ such that $F_1(\mathfrak{q}), \dots, F_m(\mathfrak{q})$ is a $\hat{M}(\mathfrak{q})$ -sequence for all \mathfrak{q} in U .*

Proof. Set $M_0 = M, M_i = M/\langle F_1, \dots, F_i \rangle M$ and consider for $i = 1, \dots, m$ the exact sequence

$$0 \rightarrow K_{i-1} \rightarrow M_{i-1} \xrightarrow{F_i} M_{i-1} \rightarrow M_i \rightarrow 0, \quad (*)$$

with K_{i-1} the kernel of F_i .

(i) By Lemma 15.2 $\hat{R}(\mathfrak{p}) = R_{\mathfrak{p}}(\mathfrak{p})^\wedge$ and $\hat{M}_i(\mathfrak{p}) = M_{i, \mathfrak{p}}(\mathfrak{p})^\wedge$ for all i and hence

$$F_i(\mathfrak{p}) = F_{i, \mathfrak{p}}^\wedge : (M_{i-1, \mathfrak{p}}(\mathfrak{p}))^\wedge \rightarrow (M_{i-1, \mathfrak{p}}(\mathfrak{p}))^\wedge.$$

Since $M_{i, \mathfrak{p}}(\mathfrak{p})$ is a finite $R_{\mathfrak{p}}(\mathfrak{p})$ -module we have (by [AM69, Proposition 10.13]) $\hat{M}_i(\mathfrak{p}) = M_{i, \mathfrak{p}}(\mathfrak{p}) \otimes_{R_{\mathfrak{p}}(\mathfrak{p})} R_{\mathfrak{p}}(\mathfrak{p})^\wedge$. Moreover $R_{\mathfrak{p}}(\mathfrak{p})^\wedge$ is faithfully flat over the local ring $R_{\mathfrak{p}}(\mathfrak{p})$ ([Mat86, Theorem 8.14]). Hence $F_{i, \mathfrak{p}} : M_{i-1, \mathfrak{p}}(\mathfrak{p}) \rightarrow M_{i-1, \mathfrak{p}}(\mathfrak{p})$ is injective iff $F_i(\mathfrak{p}) : \hat{M}_{i-1}(\mathfrak{p}) \rightarrow \hat{M}_{i-1}(\mathfrak{p})$ is injective.

(ii) By assumption $K_{i-1} = 0$ for $i = 1, \dots, m$ and $M/\langle F_1, \dots, F_m \rangle M$ is A -flat. By Lemma 16 the Jacobson radical of R contains $\langle x \rangle$ and F_1, \dots, F_m is a regular sequence contained in the Jacobson radical. Hence M_i is A -flat and $M_{i, \mathfrak{p}}$ is $A_{\mathfrak{p}}$ -flat for all i (repeated application of [Mat86, Theorem 22.2]). Tensoring $0 \rightarrow M_{i-1, \mathfrak{p}} \rightarrow M_{i-1, \mathfrak{p}} \rightarrow M_{i, \mathfrak{p}} \rightarrow 0$ with $\otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$ we get an exact sequence $0 \rightarrow M_{i-1, \mathfrak{p}}(\mathfrak{p}) \rightarrow M_{i-1, \mathfrak{p}}(\mathfrak{p}) \rightarrow M_{i, \mathfrak{p}}(\mathfrak{p}) \rightarrow 0$ for $i = 1, \dots, m$ by [Mat86, Theorem 22.3]. Now apply (i).

(iii) Localizing the exact sequence (*) at $\mathfrak{n}_{\mathfrak{p}}$ we get an exact sequence of finite $R_{\mathfrak{n}_{\mathfrak{p}}}$ -modules. Taking the $\langle x \rangle$ -adic completion, the sequence stays exact and we see that $(K_{i-1, \mathfrak{n}_{\mathfrak{p}}})^\wedge = \ker(F_{i, \mathfrak{n}_{\mathfrak{p}}}^\wedge : (M_{i-1, \mathfrak{n}_{\mathfrak{p}}})^\wedge \rightarrow (M_{i-1, \mathfrak{n}_{\mathfrak{p}}})^\wedge)$. By Lemma 15 $\hat{M}_{i-1}(\mathfrak{p}) = (M_{i-1, \mathfrak{n}_{\mathfrak{p}}})^\wedge \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$ and $F_i(\mathfrak{p}) = F_{i, \mathfrak{n}_{\mathfrak{p}}}^\wedge \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$, and by assumption $F_i(\mathfrak{p})$ is injective. We apply now repeatedly [Mat86, Theorem 22.5] to $A_{\mathfrak{p}} \rightarrow \hat{R}_{\mathfrak{n}_{\mathfrak{p}}} = A_{\mathfrak{p}}[[x]]$ and to $F_{i, \mathfrak{n}_{\mathfrak{p}}}^\wedge$ to get that $(K_{i-1, \mathfrak{n}_{\mathfrak{p}}})^\wedge = K_{i-1, \mathfrak{n}_{\mathfrak{p}}} \otimes_{R_{\mathfrak{n}_{\mathfrak{p}}}} R_{\mathfrak{n}_{\mathfrak{p}}}^\wedge = 0$ and that $(M_{i, \mathfrak{n}_{\mathfrak{p}}})^\wedge = M_{i, \mathfrak{n}_{\mathfrak{p}}} \otimes_{R_{\mathfrak{n}_{\mathfrak{p}}}} R_{\mathfrak{n}_{\mathfrak{p}}}^\wedge$ is flat over $A_{\mathfrak{p}}$ for all i . Since $R_{\mathfrak{n}_{\mathfrak{p}}}^\wedge$ is faithfully flat over $R_{\mathfrak{n}_{\mathfrak{p}}}$ this implies $K_{i-1, \mathfrak{n}_{\mathfrak{p}}} = 0$ and that $M_{i, \mathfrak{n}_{\mathfrak{p}}}$ is flat over $A_{\mathfrak{p}}$.

The support of the R -module K_{i-1} is closed and hence $(K_{i-1})_{\mathfrak{n}_{\mathfrak{q}}}^\wedge = 0$ for \mathfrak{q} in an open neighbourhood U of \mathfrak{p} in $\text{Spec } A$. Moreover the flatness of $M/\langle F_1, \dots, F_m \rangle M$ implies that $M_{\mathfrak{n}_{\mathfrak{q}}}^\wedge / \langle F_1, \dots, F_m \rangle M_{\mathfrak{n}_{\mathfrak{q}}}^\wedge$ is $A_{\mathfrak{q}}$ -flat. Applying [Mat86, Theorem 22.5] now to $F_{i, \mathfrak{n}_{\mathfrak{q}}}^\wedge : (M_{i-1})_{\mathfrak{n}_{\mathfrak{q}}}^\wedge \rightarrow (M_{i-1})_{\mathfrak{n}_{\mathfrak{q}}}^\wedge$ we get that $\hat{M}_{i-1}(\mathfrak{q}) \rightarrow \hat{M}_{i-1}(\mathfrak{q})$ is injective and that $F_1(\mathfrak{q}), \dots, F_m(\mathfrak{q})$ is an $\hat{M}(\mathfrak{q})$ -sequence. \square

Proposition 68. *Let A be a Noetherian ring and $I \subset \langle x \rangle A[[x]]$ an ideal generated by F_1, \dots, F_m , such that $A[[x]]/IA[[x]]$ is A -flat. For $\mathfrak{p} \in \text{Spec } A$ denote by $\hat{I}(\mathfrak{p}) \subset k(\mathfrak{p})[[x]]$ the ideal generated by $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$.*

1. *If $\hat{I}(\mathfrak{p})$ is a complete intersection, then $\hat{I}(\mathfrak{q})$ is a complete intersection for \mathfrak{q} in an open neighbourhood of \mathfrak{p} in $\text{Spec } A$.*
2. *Assume that $\hat{I}(\mathfrak{p})$ is an ICIS and that the hypotheses of Proposition 66 are satisfied. Then $\hat{I}(\mathfrak{q})$ is an ICIS with $\tau(I(\mathfrak{p})) \geq \tau(I(\mathfrak{q}))$ for \mathfrak{q} in an open neighbourhood of \mathfrak{p} in $\text{Spec } A$.*

Proof. 1. We may assume that $F_1(\mathfrak{p}), \dots, F_m(\mathfrak{p})$ is a $k(\mathfrak{p})[[x]]$ -sequence. By Proposition 67 $F_1(\mathfrak{q}), \dots, F_m(\mathfrak{q})$ is a $k(\mathfrak{q})[[x]]$ -sequence, hence $\hat{I}(\mathfrak{q})$ is a complete intersection, for \mathfrak{q} in an open neighbourhood of \mathfrak{p} in $\text{Spec } A$.

2. follows from Proposition 66 since for $\hat{I}(\mathfrak{q}) \subset k(\mathfrak{q})[[x]]$ a complete intersection $\dim_{k(\mathfrak{q})} T_{\hat{I}(\mathfrak{q})}$ is the Tjurina number of $\hat{I}(\mathfrak{q})$. \square

References

- [AM69] Atiyah, M.F., Macdonald I.G.: *Introduction to commutative algebra*. Addison-Wesley, 1969.
- [BGM11] Boubakri, Y.; Greuel, G.-M.; Markwig, T. *Normal Forms of Hypersurface Singularities in Positive Characteristic*. Mosc. Math. J. 11, No. 4, 657-683 (2011).
- [BGM12] Boubakri, Y.; Greuel, G.-M.; Markwig, T. *Invariants of hypersurface singularities in positive characteristic*. Rev. Mat. Complut. 25, No. 1, 61-85 (2012).
- [DGPS] Decker, W.; Greuel, G.-M.; Pfister, G.; Schönemann, H.: SINGULAR 4-1-2 – A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2019).
- [Eis95] Eisenbud, D.; *Commutative algebra. With a view toward algebraic geometry*. Graduate Texts in Mathematics 150. Springer (1995).

- [GR71] Grauert, H.; Remmert, R.: *Analytische Stellenalgebren*, Grundlehren der mathematischen Wissenschaften, Springer (1971).
- [GLS07] Greuel, G.-M.; Lossen, C.; Shustin, E.: *Introduction to Singularities and Deformations*. Springer (2007).
- [GP08] Greuel, G.-M., Pfister, G.: *A **Singular** Introduction to Commutative Algebra*, 2nd Edition, Springer, Berlin (2008). With contributions by O. Bachmann, C. Lossen and H. Schönemann.
- [GPh19] Greuel, G.-M.; Pham, T.H.: *Finite determinacy of matrices and ideals in arbitrary characteristics*, J. Algebra 530 (2019).
- [KPR75] Kurke, H.; Pfister, G.; Roczen, M.: *Henselsche Ringe und algebraische Geometrie*. Mathematische Monographien. Band 11. Berlin: VEB Deutscher Verlag der Wissenschaften. 223 S. M 57.00 (1975).
- [KPP78] Kurke, H.; Pfister, G.; Popescu, D.; Roczen, M.; Mostowski T.: *Die Approximationseigenschaft lokaler Ringe*. Lecture Notes in Mathematics 634, Springer-Verlag Berlin Heidelberg New York (1978).
- [La67] Lafou, J. P. : *Anneaux henséliens et théorèmes de préparation*. C. R. Acad. Sci., Paris, A1161 -A1162 (1967).
- [Maj10] Majadas, J., Rodicio, A.G.: : *Smoothness, Regularity and Complete Intersection*. LN London Math. Soc. 373 (2010).
- [Mat86] Matsumura, H.: *Commutative ring theory*. Cambridge Univ. Press (1986).
- [Pf07] Pfister, G. *On modular computation of standard basis*. An. Stiint. Univ. Ovidius Constana, Ser. Mat. 15, No. 1, 129-138 (2007).
- [Sa71] Saito, K. *Quasihomogene isolierte Singularitäten von Hyperflächen*. Invent. Math. 14, 123-142 (1971).
- [Sta19] The Stacks project authors. *The Stacks project* (2019)
<https://stacks.math.columbia.edu>

Gert-Martin Greuel and Gerhard Pfister
University of Kaiserslautern
Department of Mathematics
Erwin-Schroedinger Str.
67663 Kaiserslautern
Germany
e-mail: greuel@mathematik.uni-kl.de
e-mail: pfister@mathematik.uni-kl.de