# Long-time Asymptotic Behavior of the Fifth-order Modified KdV Equation in Low Regularity Spaces

Nan Liu<sup> $\dagger$ </sup>, Mingjuan Chen<sup> $\dagger$ ,\*</sup>, Boling Guo<sup> $\dagger$ </sup>

<sup>†</sup> Institute of Applied Physics and Computational Mathematics, Beijing 100088, China

**Abstract.** Based on the nonlinear steepest descent method of Deift and Zhou for oscillatory Riemann–Hilbert problems and the Dbar approach, the long-time asymptotic behavior of solutions to the fifth-order modified Korteweg-de Vries equation on the line is studied in the case of initial conditions that belong to some weighted Sobolev spaces. Using techniques in Fourier analysis and the idea of *I*-method, we give its global well-posedness in lower regularity Sobolev spaces, and then obtain the asymptotic behavior in these spaces with weights.

**Keywords:** Fifth-order modified Korteweg-de Vries equation; Nonlinear steepest descent method; Dbar method; Long-time asymptotics; Fourier analysis, Low regularity.

# 1 Introduction

The modified KdV (Korteweg-de Vries) equation is a fundamental completely integrable model in solitary waves theory, and usually is written as

$$\partial_t u + 6u^2 \partial_x u + \partial_x^3 u = 0, \tag{1.1}$$

where u = u(x,t) is a real function with evolution variable t and transverse variable x. This equation is well-known as a canonical model for the description of nonlinear long waves in many branches of physics when there is polarity symmetry. For instance, applications in the context of electrodynamics are described in [46], in the context of stratified fluids in reference [23]. Recently, a fifth-order modified KdV taking the form

$$\partial_t u + \partial_x^5 u + 30u^4 \partial_x u - 10u^2 \partial_x^3 u - 10(\partial_x u)^3 - 40u \partial_x u \partial_x^2 u = 0,$$
(1.2)

was reported in [49, 50], which also describes nonlinear wave propagation in physical systems with polarity symmetry. Meanwhile, Equation (1.2) also has certain application for the description of nonlinear internal waves in a fluid stratified by both density and current [22, 45].

Equation (1.2) is endowed with two important features. One is the complete integrability: in the sense that there are Lax pair formulations, so (1.2) can be solved by means of the inverse scattering transformation formalism both in the case of vanishing as well as nonvanishing boundary conditions. Being integrable, Equation (1.2) admits an infinite number of conservation laws and its

<sup>\*</sup>Corresponding author.

E-mail addresses: ln10475@163.com (N. Liu), mjchenhappy@pku.edu.cn (M. Chen), gbl@iapcm.ac.cn (B. Guo).

solution exists globally in time for any Schwartz initial data. The other one is the scaling symmetry: for any solution u(x,t) of (1.2) with initial data  $u_0(x)$ , the scaling functions  $u_{\lambda}(x,t) := \lambda u(\lambda x, \lambda^5 t)$ also solve (1.2) with initial data  $u_{0,\lambda} := \lambda u_0(\lambda x)$ . The Sobolev space  $\dot{H}^{s_c}$  is called the critical space if it satisfies

$$||u_{0,\lambda}||_{\dot{H}^{s_c}} \sim ||u_0||_{\dot{H}^{s_c}}.$$

By simple calculation, one can know that  $s_c = -1/2$ . This scale invariance suggests ill-posedness for  $H^s$ , s < -1/2.

Using the Fourier analysis techniques and the theory of dispersive equations, one can study the low regularity theory for the fifth-order modified KdV equation. It seems natural that the regularity index of the well-posedness theory can be decreased to -1/2. However, it is extremely difficult to achieve this goal by the known methods in Sobolev spaces. The best local well-posedness theory so far was given by Kwon [38]. He expressed that Equation (1.2) is locally well-posed in Sobolev space  $H^s(\mathbb{R})$  for  $s \geq 3/4$  via the  $X^{s,b}$  method which was posed by Bourgain, and showed that it is ill-posed when s < 3/4 in the sense that the data-to-solution map fails to be uniformly continuous.

In addition, the solutions to (1.2) enjoy mass conservation and energy conservation:

$$M(u) = \int_{\mathbb{R}} u^2 \mathrm{d}x; \qquad E(u) = \int_{\mathbb{R}} (\partial_x u)^2 + u^4 \mathrm{d}x.$$
(1.3)

Energy conservation and the local well-posedness immediately yield the global well-posedness of (1.2) from initial data in  $H^1(\mathbb{R})$ . The biggest obstacle in getting global solutions in  $H^s$  with 0 < s < 1 is the lack of any conservation law. In this paper, by utilizing the idea of *I*-method, we modify the  $H^1$  energy to obtain an "almost conservation law". Therefore, we can get the global well-posedness in  $H^s$ , s > 19/22.

There are many researchers studying the local and global well-posteness theory of fifth-order KdV-type equations. For the fifth-order KdV equation, whose nonlinearities are quadratic, Chen and Guo [10] gave the local and global well-posedness in  $H^s$  with  $s \ge -7/4$  for the equation with the first-order derivative nonlinearities. The fifth-order KdV equation with the third-order derivative nonlinearities was considered by Kwon [37], and he showed that it is locally well-posedness in  $H^s$ . s > 5/2. In [24], the authors prove a priori bound of solutions to this equation in  $H^s$ ,  $s \ge 5/4$ , and showed that the equation is globally well-posed in  $H^2$  due to the second energy conservation law. For the fifth-order modified KdV equation, whose nonlinearities are cubic, Linares [40] gave the local well-posedness in  $H^2$  by using the dispersive smoothing effect, and then the global well-posedness is immediately obtained via the conservation law. Kwon [38] improved the local well-posedness theory to the Sobolev spaces  $H^s$  with  $s \geq 3/4$ . In addition, by using *I*-method, Gao [20] prove the global well-posedness in  $H^s$ , s > -3/22 for the fifth-order modified KdV equation with a first-order derivative nonlinearity. Via the energy method and the short-time Fourier restriction norm method, Kwak [36] considered the Equation (1.2) under the periodic boundary condition and proved the local well-posedness in  $H^{s}(\mathbb{T})$ , s > 2. For more results, we lead the readers to [7,30,31,47] and the references therein.

From the integrable point of view, in the context of inverse scattering, the first work to provide explicit formulas depending only on initial conditions for long-time asymptotics of solutions is due to Zakharov and Manakov [52], where the model considered was the nonlinear Schrödinger (NLS) equation. In this setting, the inverse scattering map and the reconstruction of the potential is formulated through an oscillatory Riemann–Hilbert (RH) problem. Then the now well-known nonlinear steepest descent method for oscillatory RH problems introduced by Deift and Zhou in [16] provides a rigorous and transparent proof to analyze the long-time asymptotics of initial value problems for a large range of nonlinear integrable evolution equations with initial conditions lying in Schwartz space  $\mathscr{S}(\mathbb{R})$ . Numerous new significant results for the initial value or initial-boundary value problems of a lot of integrable systems were obtained under the assumptions that the initial or initial boundary data belong to the Schwartz space [2, 11, 16, 17, 42, 51].

To consider the asymptotic behavior of the solution in lower regularity space, we have to mention another meaningful result developed by Zhou in [54], that is, the  $L^2$ -Sobolev space bijectivity of the direct and inverse scattering of the 2 × 2 AKNS system for the initial data  $u_0(x)$  belonging to the weighted Solobev space  $H^{i,j}(\mathbb{R})$ , where

$$H^{i,j}(\mathbb{R}) = \left\{ f(x) \in L^2(\mathbb{R}) : x^j f(x), f^{(i)}(x) \in L^2(\mathbb{R}) \right\},$$

which is endowed with the following norm

$$\|f(x)\|_{H^{i,j}(\mathbb{R})} = \left(\|f(x)\|_{L^2(\mathbb{R})}^2 + \|x^j f(x)\|_{L^2(\mathbb{R})}^2 + \|f^{(i)}(x)\|_{L^2(\mathbb{R})}^2\right)^{\frac{1}{2}}.$$

And then Deift and Zhou in [18] obtained the long-time asymptotics for solutions of the defocusing NLS equation with initial data in a weighted Sobolev space  $H^{1,1}$  based on this work. Moreover, inspired by the work [43] and combined Zhou's result [54], the Dbar generalization of the nonlinear steepest descent method proposed by Dieng and McLaughlin [19] was implemented in the analysis of the long-time asymptotic behavior of the solutions, where they derived the asymptotics for the defocusing NLS equation with the initial conditions  $u_0(x)$  belonging to the weighted Sobolev space  $H^{1,1}(\mathbb{R})$ . This approach replaces the rational approximation of the reflection coefficient in Deift-Zhou method [16] by some non-analytic extension of the jump matrices off the real axis, leading to a  $\bar{\partial}$  problem to be solved in some sectors of the complex plane. The new  $\bar{\partial}$  problem can be recast into an integral equation and solved by Neumann series, which contributed to the error estimate. In the context of NLS equatin with soliton solutions, they were successfully applied to prove asymptotic stability of N-soliton solutions to defocusing NLS equation [14] and address the soliton resolution problem for focusing NLS equation in [5]. Whereafter, the long-time asymptotic behavior of derivative NLS equation for generic initial data in  $H^{2,2}(\mathbb{R})$  that do not support solitons and can support solitons (but exclude spectral singularities) were analyzed in [41] and [29], respectively. Recently, long-time behavior of the defocusing modified KdV equation and the soliton resolution of the focusing modified KdV equation in weighted Sobolev spaces were reported in [8,9].

It is worth noting that some researchers also study the long-time asymptotic behavior of the dispersive equations by PDE techniques, which do not need the complete integrability. For instance, one can see the references [27], [21] and [26] about the methods of factorization of operators, the space-time resonance, and wave packets analysis, respectively. However, they all need some smallness assumptions on the initial data. Specially, employing the method of testing by wave packets, Okamoto [44] has proved the small-data global existence and derived the asymptotic behavior of solutions in both the self-similar and oscillatory regions to a fifth-order modified KdV type equation in spaces  $H^{2,1}(\mathbb{R})$ , whereas the complex-valued function W arising in the leading-order term can not be written exactly in oscillatory region.

In this paper, relying on complete integrability, we drop the smallness conditions and consider the long-time asymptotic behavior of the fifth-order modified KdV equation (1.2) by using the  $\bar{\partial}$  generalization of the nonlinear steepest descent method with the initial conditions  $u_0(x)$  belonging to the weighted Sobolev space  $H^{4,1}(\mathbb{R})$ .

More precisely, in the physically interesting Region I:  $k_0 \leq M, \tau \to \infty$ , where M is a positive constant, the phase function  $\Phi(k)$  defined in (1.20) has two different real critical points located at  $\pm k_0$ , where  $k_0 = \sqrt[4]{x/(80t)}$ . To implement the  $\bar{\partial}$  approach, our first step is to introduce a scale function  $\delta(k)$  conjugated to the solution M(x,t;k) of the original RH problem 1.1. This operation is aimed to factorize the jump matrix (1.22) into a product of a upper/lower triangular and lower/upper triangular matrix, which is necessary in the following contour deformation. The second step is to deform the contour from  $\mathbb{R}$  to a new contour depicted in Fig. 3 so that the jump matrix involves the exponential factor  $e^{-t\Phi}$  on the parts of the contour where  $Re\Phi$  is positive and the factor  $e^{t\Phi}$  on the parts where  $Re\Phi$  is negative. To achieve this goal, the idea is to construct  $\bar{\partial}$ extensions  $R_i(k)$  in  $\Omega_i$  to have the prescribed boundary values and  $\bar{\partial}R_i(k)$  small in the sector (see Lemma 2.1). This will allow us to reformulate RH problem 2.1 as a mixed  $\bar{\partial}$ -RH problem 2.2 for a new matrix-valued function  $M^{(2)}(x,t;k)$ . The next step is to extract from  $M^{(2)}$  a contribution that is a pure RH problem. More exactly, we factorize  $M^{(2)}(k) = M^{(3)}(k)M^{mod}(k)$ , where  $M^{mod}(k)$  is a solution of the RH problem 2.3 below with the jump matrix  $J^{mod} = J^{(2)}$ , and  $M^{(3)}(k)$  has no jump across  $\Gamma$  which will prove to be satisfied a pure  $\bar{\partial}$  problem 2.6. Since the reflection coefficient r(k) in the jump matrix  $J^{mod}$  is only fixed at  $\pm k_0$  along the deformed contours, we then can solve this RH problem in terms of parabolic cylinder functions and the large-k expansion can be exactly written with the estimate of decay rate as  $\tau \to \infty$  (see Subsection 2.3). The remaining  $\bar{\partial}$  problem may be written as an integral equation (refer to equation (2.71)) whose integral operator has small norm at large times (see equation (2.74)) and the large-time contribution to the asymptotics of u(x,t) is negligible. The final step is to regroup all the transformations to find the behavior of the solution of fifth-order modified KdV equation as  $\tau \to \infty$  using the large-k behavior of the RH problem solutions.

However, our main RH problem has only jump across the real axis  $\mathbb{R}$ , the main contributions to the asymptotic formula of the solution u(x,t) come from the local RH problems near the two real critical points  $\pm k_0$  even though there are two real and two pure imaginary critical points of the phase function  $\Phi(k)$  located at the points  $\pm k_0$  and  $\pm ik_0$ . Due to symmetries of reflection coefficient r(k), the leading-order aysmptotics of u(x,t) exhibits decaying, of order  $O(t^{-1/2})$ , modulated oscillations in Region I.

Another interesting region is the Region III:  $\tau \leq M'$ . Especially, as  $t \to \infty$ , the critical points  $\pm k_0$  approach 0. The above steps are much easier to operate in this case. We can show that the asymptotics of the solution u(x,t) in this region is expressed in terms of the solution of a fourth-order Painlevé II equation (see [35])

$$u_p^{\prime\prime\prime\prime}(y) - 40u_p^2(y)u_p^{\prime\prime}(y) - 40u_p(y)u_p^{\prime 2}(y) + 96u_p^5(y) - 4yu_p(y) = 0.$$
(1.4)

It is a beautiful example that the asymptotic behavior of the solution of an integrable equation is expressed by the solution of the high order Painlevé II equation in the asymptotic analysis. Nevertheless, the asymptotic behaviors of solutions to the standard defocusing mKdV in sector  $0 < x < Mt^{\frac{1}{3}}$  correspond to the solution of the standard Painlevé II equation  $u''_p(y) - 2u^3_p(y) - yu_p(y) = 0$ , both on the line [16] and half-line problem [39]. We also note that for the Camassa–Holm and Sasa– Satsuma equation, there also exists Painlevé II-type asymptotics in some certain regions [6, 28]. Interestingly, the Painlevé III hierarchy has appeared in recent study of the fundamental rogue wave solutions of the focusing nonlinear Schrödinger equation in the limit of large order [4]. However, for the fifth-order modified KdV equation (1.2), we find another interesting new asymptotic result which related to the solution of a fourth-order Painlevé II equation (1.4). The study of asymptotic behaviors of the solution u(x,t) in remaining regions is derived in Section 4, which follows the same strategy.

All in all, our asymptotic results can be divided into three categories: in Region I (which is a oscillatory region), the leading-order asymptotics of u(x,t) is described by the form of cosine function; in the self-similar Regions II-IV, the leading-order asymptotic behaviors can be expressed in terms of the solution of a fourth-order Painlevé II equation; in the Region V, the solution u(x,t)tends to 0 with fast decay.

To extend these asymptotic behavior results to lower regularity spaces, we have to obtain the global well-posedness theory in some low regularity spaces by PDE techniques first. As mentioned before, the global solution in space  $H^1(\mathbb{R})$  is easily to obtain owing to the energy conservation. The biggest obstacle in getting global solutions in  $H^s$  with 0 < s < 1 is the lack of any conservation law. In this paper, by utilizing the idea of *I*-method, we modify the  $H^1$  energy to obtain an "almost conservation law", whose increment is very small. Therefore, we can get the global well-posedness in low regularity Sobolev spaces and then extend the long-time asymptotic formulae to the rough data  $u_0(x) \in H^{s,1}(\mathbb{R})$ , s > 19/22.

Broadly speaking, the asymptotic analysis of (1.2) presents following innovation points: (i) the phase function  $\Phi(k)$  in jump matrix raises to the fifth power of k, this will lead to the computations in the scaling transform and related estimates about the  $\bar{\partial}$  problem more involved; (ii) we should establish a new suitable model RH problem which arise in the study of long-time asymptotics in the Region III; (iii) because of the lack of any conservation law in  $H^s$ , s < 1, we have to get an "almost conservation law" by utilizing *I*-method. As far as the authors can see, it is the first time that using the idea of *I*-method to hand with the third-order derivative nonlinearities. Compared with the first-order derivative nonlinearities, the absence of the symmetries and the vanished property in the case of third-order derivatives makes the problem fairly tricky. Thanks to an important observation, substantial technical difficulties cased by the third-order derivative nonlinearities could be solved.

### 1.1 Formulation of the Riemann–Hilbert problem

Equation (1.2) is the compatibility condition for the simultaneous linear equations of a Lax pair

$$\Psi_x = X\Psi, \quad X = ik\sigma_3 + U, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad U = \begin{pmatrix} 0 & u \\ u & 0 \end{pmatrix}, \tag{1.5}$$

and

$$\Psi_t = T\Psi, \quad T = -16ik^5\sigma_3 + V, \tag{1.6}$$
  

$$V = -8ik^3U^2\sigma_3 + 2ik(2UU_{xx} - U_x^2 - 3U^4)\sigma_3 - 16k^4U + 8ik^3\sigma_3U_x + k^2(4U_{xx} - 8U^3)$$
  

$$+ik\sigma_3(12U^2U_x - 2U_{xxx}) - 6U^5 + 10U^2U_{xx} + 10UU_x^2 - U_{xxxx}.$$

governing a  $2 \times 2$  matrix-valued function  $\Psi(x,t;k)$  and the spectral parameter  $k \in \mathbb{C}$ . For the given initial data  $u_0(x) \in L^1(\mathbb{R})$ , we define the Jost eigenfunctions  $\Psi_{\pm}(x;k) = \psi_{\pm}(x;k)e^{ikx\sigma_3}$  by

$$\psi_{\pm}(x;k) = I + \int_{\pm\infty}^{x} e^{ik(x-x')\sigma_3} [U(x',0)\psi_{\pm}(x';k)] e^{-ik(x-x')\sigma_3} dx', \qquad (1.7)$$

which satisfy

$$\psi_{\pm x}(x;k) - ik[\sigma_3,\psi_{\pm}(x;k)] = U(x,0)\psi_{\pm}(x;k), \qquad (1.8)$$

and the normalization conditions

$$\lim_{x \to \pm \infty} \psi_{\pm}(x;k) = I.$$

If we denote  $\psi = (\psi^1, \psi^2)$  for a 2 × 2 matrix  $\psi$ , it follows from (1.7) that for all (x, t):

- (i)  $\det[\psi_{\pm}] = 1.$
- (i)  $\psi_+^1$  and  $\psi_-^2$  are analytic and bounded in  $\{k \in \mathbb{C} | \text{Im}k > 0\}$ , and  $(\psi_+^1, \psi_-^2) \to I$  as  $k \to \infty$ . (iii)  $\psi_-^1$  and  $\psi_+^2$  are analytic and bounded in  $\{k \in \mathbb{C} | \text{Im}k < 0\}$ , and  $(\psi_-^1, \psi_+^2) \to I$  as  $k \to \infty$ .
- (iv)  $\psi_{\pm}$  are continuous up to the real axis.
- (v) Symmetry:

$$\psi_{\pm}(x;\bar{k}) = \psi_{\pm}(x;-k) = \sigma_1 \psi_{\pm}(x;k)\sigma_1, \qquad (1.9)$$

where  $\sigma_1$  is the first Pauli matrix. The symmetry relation (1.9) can be proved easily due to the symmetries of the matrix X:

$$X(x,t;\bar{k}) = X(x,t;-k) = \sigma_1 X(x,t;k)\sigma_1.$$

Both  $\psi_+$  and  $\psi_-$  define a fundamental solution matrix for (1.8) and so there exists a continuous matrix s(k) independent of x, satisfying

$$\Psi_{+}(x;k) = \Psi_{-}(x;k)s(k), \quad k \in \mathbb{R}.$$
(1.10)

Evaluation at  $x \to -\infty$  gives

$$s(k) = \lim_{x \to -\infty} e^{-ikx\sigma_3} \psi_+(x;k) e^{ikx\sigma_3},$$
  
$$= I - \int_{-\infty}^{+\infty} e^{-ikx\sigma_3} [U(x,0)\psi_+(x;k)] e^{ikx\sigma_3} dx.$$
 (1.11)

It follows from det[ $\psi_{\pm}(x;k)$ ] = 1 and the symmetries (1.9) that the matrix-valued function s(k)can be expressed as

$$s(k) = \begin{pmatrix} a(k) & \overline{b(k)} \\ b(k) & \overline{a(k)} \end{pmatrix}, \quad \det[s(k)] = 1, \quad k \in \mathbb{R}.$$
(1.12)

Moreover, by (1.11) and (1.12), we find that a(k) is analytic in  $\mathbb{C}_+$ , and

$$a(k) = \overline{a(-\overline{k})}, \quad b(k) = \overline{b(-k)}.$$
 (1.13)

However, it follows from det[s(k)] = 1, i.e.,

$$|a(k)|^2 - |b(k)|^2 = 1$$
(1.14)

that a(k) is zero-free.

At each time t, we define the spectral matrix s(k, t) according to (1.11), i.e.

$$s(k,t) = \lim_{x \to -\infty} e^{-ikx\sigma_3} \psi_+(x,t;k) e^{ikx\sigma_3}.$$
(1.15)

Then from the t-part Lax pair of (1.2), the evolution of the spectral matrix is given by

$$s(k,t) = e^{-16ik^5 t\sigma_3} s(k) e^{16ik^5 t\sigma_3}.$$
 (1.16)

More precisely, we have

$$a(k,t) = a(k), \ b(k,t) = b(k)e^{32ik^5t}.$$
 (1.17)

Now we define the Beals–Coifman solution

$$M(x,t;k) = \begin{cases} \left(\frac{\psi_{+}^{1}(x,t;k)}{a(k)}, \psi_{-}^{2}(x,t;k)\right), & k \in \mathbb{C}_{+}, \\ \left(\psi_{-}^{1}(x,t;k), \frac{\psi_{+}^{2}(x,t;k)}{\overline{a(\bar{k})}}\right), & k \in \mathbb{C}_{-}. \end{cases}$$
(1.18)

Then, the boundary values  $M_{\pm}(x,t;k)$  of M as k approaches  $\mathbb{R}$  from the sides  $\pm \text{Im}k > 0$  are related as follows:

$$M_{+}(x,t;k) = M_{-}(x,t;k)J(x,t;k), \quad k \in \mathbb{R},$$
(1.19)

with

$$J(x,t;k) = \begin{pmatrix} 1 - |r(k)|^2 & -\overline{r(k)}e^{-t\Phi(k)} \\ r(k)e^{t\Phi(k)} & 1 \end{pmatrix},$$
  
$$r(k) = \frac{b(k)}{a(k)}, \quad \Phi(k) = 32ik^5 - 2i\frac{x}{t}k.$$
 (1.20)

It obviously follows from Zhou's results [54] that the following map holds and is Lipschitz:

$$\mathcal{D}: H^{4,1}(\mathbb{R}) \ni \{u_0(x)\} \mapsto \{r(k)\} \in H^{1,4}(\mathbb{R}).$$

Then, we arrive at our main RH problem which is formulated as follows.

**Riemann–Hilbert problem 1.1.** Given  $r(k) \in H^{1,4}(\mathbb{R})$ . Find an analytic 2 × 2 matrix-valued function M(x,t;k) on  $\mathbb{C} \setminus \mathbb{R}$  with the following properties:

- 1. M(x,t;k) is analytic for  $k \in \mathbb{C} \setminus \mathbb{R}$  and is continuous for  $k \in \mathbb{R}$ .
- 2. The boundary values  $M_{\pm}(x,t;k)$  satisfy the jump condition

$$M_{+}(x,t;k) = M_{-}(x,t;k)J(x,t;k), \qquad (1.21)$$

where

$$J(x,t;k) = \begin{pmatrix} 1 - |r(k)|^2 & -\overline{r(k)}e^{-t\Phi(k)} \\ r(k)e^{t\Phi(k)} & 1 \end{pmatrix}, \quad r(-k) = \overline{r(k)}, \quad k \in \mathbb{R}.$$
 (1.22)

3. M(x,t;k) has the asymptotics:

$$M(x,t;k) = I + O\left(\frac{1}{k}\right), \ k \to \infty.$$

The jump matrix admits the following factorization:

$$J(x,t;k) = J_{-}^{-1}(x,t;k)J_{+}(x,t;k) = (I - w_{-}(x,t;k))^{-1}(I + w_{+}(x,t;k)),$$
(1.23)

where

$$w_{-} = \begin{pmatrix} 0 & -\overline{r(k)} e^{-t\Phi(k)} \\ 0 & 0 \end{pmatrix}, \quad w_{+} = \begin{pmatrix} 0 & 0 \\ r(k) e^{t\Phi(k)} & 0 \end{pmatrix}, \quad k \in \mathbb{R}$$

By standard RH theory, the existence and uniqueness of the solution to Riemann–Hilbert problem 1.1 is determined by the existence and uniqueness of the following singular integral equation

$$\mu(x,t;k) = I + \mathcal{C}_w \mu(x,t;k) = I + \mathcal{C}_+(\mu w_-)(x,t;k) + \mathcal{C}_-(\mu w_+)(x,t;k),$$
(1.24)

where

$$\mu(x,t;k) = M_{+}(x,t;k)(I + w_{+}(x,t;k))^{-1} = M_{-}(x,t;k)(I - w_{-}(x,t;k))^{-1}, \qquad (1.25)$$

and  $\mathcal{C}_{\pm}$  denote the Cauchy operators:

$$(\mathcal{C}_{\pm}f)(k) = \lim_{\epsilon \to 0+} \int_{\mathbb{R}} \frac{f(\zeta)}{\zeta - (k \pm i\epsilon)} \frac{\mathrm{d}\zeta}{2\pi \mathrm{i}}.$$

It then follows from [53] that  $I - C_w$  is invertible, as a result that RH problem 1.1 has a unique solution. Then the solution M(x, t; k) is given by

$$M(x,t;k) = I + \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{[\mu(w_+ + w_-)](x,t;s)}{s-k} ds.$$
(1.26)

The solution u(x,t) of (1.2) in terms of M(x,t;k) is given by

$$u(x,t) = -2i \lim_{k \to \infty} (kM(x,t;k))_{12} = \frac{1}{\pi} \left( \int_{\mathbb{R}} [\mu(w_+ + w_-)](x,t;s) \mathrm{d}s \right)_{12}.$$
 (1.27)

### 1.2 Main results

$$\begin{array}{c|c} x < 0 & x > 0 \\ \hline & & \\ IV : \underline{k_0 \le M, \tau \ge \tilde{M}} & II : \underline{o(t^{\frac{2}{7}}) = \tau} \ge \tilde{M} \\ \hline & & \\ \hline & & \\ V : k_0 \ge M, x \to -\infty & \overline{III : \tau \le M'} & \overline{I : k_0 \le M, \tau} \to \infty \end{array}$$

Fig. 1: The five regions.

Our main results in this paper are summarized by the following theorems. The first one is about

the asymptotic behavior of the Beals–Coifman solution to Equation (1.2).

**Theorem 1.1** Let  $u_0(x) \in H^{4,1}(\mathbb{R})$ , M, M' and  $\tilde{M}$  be fixed constants larger than 1. Select a suitable constant  $\rho > 0$ . Then the solution u(x,t) of fifth-order modified KdV equation (1.2) with the initial data  $u_0(x)$  admits the following uniform asymptotics as  $t \to \infty$  in the five regions depicted in Fig. 1. More precisely, set

$$k_0 = \sqrt[4]{\frac{x}{80t}}, \quad \tau = tk_0^5, \tag{1.28}$$

and let  $u_p(y)$  be the solution of the fourth order Painlevé II equation (A.5). Then, we have In Region I,

$$u(x,t) = u_{as}(x,t) + O\left(\tau^{-1} + (k_0^3 t)^{-\frac{3}{4}}\right),$$

where

$$\begin{aligned} u_{as}(x,t) &= \frac{\sqrt{\nu}}{2k_0\sqrt{10k_0t}} \cos\left(128tk_0^5 + \nu\ln(2560tk_0^5) + \varphi(k_0)\right), \\ \varphi(k_0) &= -\frac{3\pi}{4} - \arg r(k_0) + \arg \Gamma(i\nu) - \frac{1}{\pi} \int_{-k_0}^{k_0} \ln\left(\frac{1 - |r(s)|^2}{1 - |r(k_0)|^2}\right) \frac{ds}{s - k_0} \\ \nu &= -\frac{1}{2\pi} \ln(1 - |r(k_0)|^2) > 0, \end{aligned}$$

and  $\Gamma(\cdot)$  denotes the standard Gamma function. In Region II,

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O\left(t^{-\frac{3}{10}} + \left(\frac{\tau}{t}\right)^{\frac{2}{5}}\right).$$

In Region III,

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O(t^{-\frac{3}{10}}).$$

In Region IV,

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O(t^{-\frac{3}{10}} e^{-8(20\tau)^{\frac{4}{5}}\rho})$$

In Region V,

$$u(x,t) = O\left(t^{-\frac{1}{5}}e^{-c\tau} + t^{-\frac{3}{10}}e^{-8(20\tau)\frac{4}{5}\rho}\right).$$

**Remark 1.1** The significance of the restriction  $\tau = o(t^{\frac{2}{7}})$  in Region II is to ensure that the asymptotic formulas of the solution match up in the overlap section of Region I and Region II. For more details, one can see Remark 4.2.

We next consider Equation (1.2) in low regularity spaces. From the PDE point of view, one

always consider the strong solution given by an integral form. For convenience, we denote

$$F(u) = 30u^4 \partial_x u - 10u^2 \partial_x^3 u - 10(\partial_x u)^3 - 40u \partial_x u \partial_x^2 u,$$

then by Duhamel's formula, (1.2) is equivalent to the following integral equation:

$$u(x,t) = e^{-t\partial_x^5} u_0 - \int_0^t e^{-(t-t')\partial_x^5} F(u)(t') \, \mathrm{d}t', \quad \text{where} \quad e^{-t\partial_x^5} = \mathscr{F}^{-1} e^{-\mathrm{i}t\xi^5} \mathscr{F}.$$
(1.29)

Using the idea of I-method [12, 13], we obtain the global well-posedness theory.

**Theorem 1.2** Let  $19/22 < s \leq 1$ , the initial value problem of fifth-order modified equation (1.2) is globally well-posed from initial data  $u_0 \in H^s(\mathbb{R})$ .

In the end, cooperating the integrable structure with PDE techniques, we can give the long-time asymptotic behavior in lower regularity spaces.

**Theorem 1.3** Let  $u_0 \in H^{s,1}(\mathbb{R})$ ,  $19/22 < s \leq 1$ , the strong solution given by the integral form (1.29) has the same asymptotic behavior as in Theorem 1.1.

**Notations.** C denotes a universal positive constant which can always be different at different lines.  $\mathscr{F}(\mathscr{F}^{-1})$  denotes the (inverse) Fourier transform on  $\mathbb{R}$ . We also denote  $\widehat{\phi}$  the Fourier transform of a distribution  $\phi$ . We write  $a \leq b$  if  $a \leq Cb$ , and analogous for  $a \geq b$ . We use the notation  $a \sim b$  if  $a \leq b \leq a$ . We denote by  $a + = a + \varepsilon$  for some  $0 < \varepsilon \ll 1$ . We denote by  $\overline{a}$  the complex conjugate of a complex number a. The three Pauli matrices are defined by

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The organization of this paper is as follows. In Section 2, we derive the long-time asymptotics of the solution for Equation (1.2) in Region I. Section 3 is aim to consider the asymptotics of u(x, t)in the Region III. The asymptotic behavior of the solution in remaining regions is derived in Section 4. We then extend the long-time asymptotic behavior to the low regularity spaces in Section 5. A few facts related to the RH problem associated with the fourth order Painlevé II equation are collected in the Appendix A and B, for the detailed proof one can see [42].

### 2 Asymptotics in Region I

The jump matrix J defined in (1.22) involves the exponentials  $e^{\pm t\Phi}$ , therefore, the sign structure of the quantity  $\operatorname{Re}\Phi(k)$  plays an important role as  $t \to \infty$ . In particular, in region I, it follows that there are two different real stationary points located at the points where  $\frac{\partial\Phi}{\partial k} = 0$ , namely, at

$$\pm k_0 = \pm \sqrt[4]{\frac{x}{80t}}.$$
 (2.1)

The signature table for  $\operatorname{Re}\Phi(k)$  is shown in Fig. 2.



Fig. 2: The signature table for  $\operatorname{Re}\Phi(k)$  in the complex k-plane.

### 2.1 Conjugation

In order to apply the method of steepest descent, one goes from the original RH problem 1.1 for M to the equivalent RH problem for the new unknown function  $M^{(1)}$  defined by

$$M^{(1)}(x,t;k) = M(x,t;k)\delta^{-\sigma_3}(k), \qquad (2.2)$$

where the complex-valued function  $\delta(k)$  is given by

$$\delta(k) = \exp\left\{\frac{1}{2\pi i} \int_{-k_0}^{k_0} \frac{\ln(1-|r(s)|^2)}{s-k} ds\right\}, \ k \in \mathbb{C} \setminus [-k_0, k_0].$$
(2.3)

**Proposition 2.1** The function  $\delta(k)$  has the following properties:

(i)  $\delta(k)$  is bounded and analytic function of  $k \in \mathbb{C} \setminus [-k_0, k_0]$  with continuous boundary values on  $(-k_0, k_0)$ , and satisfies the jump condition across the real axis oriented from  $-\infty$  to  $\infty$ :

$$\delta_{+}(k) = \begin{cases} \delta_{-}(k)(1 - |r(k)|^{2}), & k \in (-k_{0}, k_{0}), \\ \delta_{-}(k), & k \in \mathbb{R} \setminus (-k_{0}, k_{0}). \end{cases}$$

(ii) As  $k \to \infty$ ,  $\delta(k)$  satisfies the asymptotic formula

$$\delta(k) = 1 + O(k^{-1}), \quad k \to \infty.$$
(2.4)

(iii)  $\delta(k)$  obeys the symmetry

$$\delta(k) = \overline{\delta(\bar{k})}^{-1} = \overline{\delta(-\bar{k})}, \quad k \in \mathbb{C} \setminus [-k_0, k_0].$$

(iv)

$$\delta(k) = \left(\frac{k - k_0}{k + k_0}\right)^{i\nu} e^{\chi(k)},\tag{2.5}$$

where

$$\nu = -\frac{1}{2\pi} \ln(1 - |r(k_0)|^2) > 0, \qquad (2.6)$$

$$\chi(z) = \frac{1}{2\pi i} \int_{-k_0}^{k_0} \ln\left(\frac{1 - |r(s)|^2}{1 - |r(k_0)|^2}\right) \frac{ds}{s - k}.$$
(2.7)

Moreover, along any ray of the form  $\pm k_0 + e^{i\phi}\mathbb{R}_+$  with  $0 < \phi < \pi$  or  $\pi < \phi < 2\pi$ ,

$$\left|\delta(k) - \left(\frac{k - k_0}{k + k_0}\right)^{i\nu} e^{\chi(\pm k_0)}\right| \le C |k \mp k_0|^{\frac{1}{2}}.$$
(2.8)

**Proof.** See [8] and references therein.

Then  $M^{(1)}(x,t;k)$  satisfies the following RH problem:

**Riemann–Hilbert problem 2.1.** Given  $r(k) \in H^1(\mathbb{R})$ . Find an analytic 2 × 2 matrix-valued function  $M^{(1)}(x,t;k)$  on  $\mathbb{C} \setminus \mathbb{R}$  with the following properties:

1.  $M^{(1)}(x,t;k)$  is analytic for  $k \in \mathbb{C} \setminus \mathbb{R}$  and is continuous for  $k \in \mathbb{R}$ .

2. The boundary values  $M^{(1)}_{\pm}(x,t;k)$  satisfy the jump condition

$$M_{+}^{(1)}(x,t;k) = M_{-}^{(1)}(x,t;k)J^{(1)}(x,t;k), \quad k \in \mathbb{R},$$
(2.9)

where

$$J^{(1)}(x,t;k) = \begin{cases} \begin{pmatrix} 1 & -\overline{r(k)}\delta^2(k)\mathrm{e}^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(k)\delta^{-2}(k)\mathrm{e}^{t\Phi(k)} & 1 \end{pmatrix}, & |k| > k_0, \\ \begin{pmatrix} 1 & 0 \\ \frac{r(k)}{1-|r(k)|^2}\delta^{-2}_{-}(k)\mathrm{e}^{t\Phi(k)} & 1 \end{pmatrix} \begin{pmatrix} 1 & -\frac{\overline{r(k)}}{1-|r(k)|^2}\delta^2_{+}(k)\mathrm{e}^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & |k| < k_0. \end{cases}$$
(2.10)

3.  $M^{(1)}(x,t;k) = I + O(\frac{1}{k})$ , as  $k \to \infty$ .

## 2.2 Contour deformation and introducing $\bar{\partial}$ extensions

The next step is to introduce factorizations of the jump matrix whose factors admit continuous but not necessarily analytic extensions off the real axis by exploiting the methods in [5, 8, 19, 29, 41]. More precisely, define the contours

$$L := \left\{ k \in \mathbb{C} | k = k_0 + k_0 \alpha e^{\frac{5\pi i}{6}}, -\infty < \alpha \le \frac{2\sqrt{3}}{3} \right\}$$
$$\cup \left\{ k \in \mathbb{C} | k = -k_0 + k_0 \alpha e^{\frac{\pi i}{6}}, -\infty < \alpha \le \frac{2\sqrt{3}}{3} \right\}.$$
(2.11)

Then, L and  $\overline{L}$  split the complex plane  $\mathbb{C}$  into eight regions  $\{\Omega_j\}_1^8$ , for convenience, we also write  $L \cup \overline{L} = \bigcup_{i=1}^{8} \Gamma_i$ , see Fig. 3. Now, we define the  $\overline{\partial}$  extensions in the following lemma.

Lemma 2.1 Let

$$T(k;k_0) = \left(\frac{k - k_0}{k + k_0}\right)^{i\nu}.$$
(2.12)



Fig. 3: The open sets  $\{\Omega_j\}_1^8$  and the contours  $\{\Gamma_j\}_1^8$  in the complex k-plane.

It is possible to define functions  $R_j: \overline{\Omega}_j \mapsto \mathbb{C}, \ j = 1, 3, 4, 5, 6, 8$  with boundary values satisfying

$$R_{1}(k) = \begin{cases} -r(k)\delta^{-2}(k), & k \in (k_{0}, \infty), \\ -r(k_{0})T^{-2}(k; k_{0})e^{-2\chi(k_{0})}, & k \in \Gamma_{1}, \end{cases}$$

$$(2.13)$$

$$R_{3}^{+}(k) = \begin{cases} \frac{T(k)}{1 - |r(k)|^{2}} \delta_{+}^{2}(k), & k \in (0, k_{0}), \\ \frac{1}{1 - |r(k_{0})|^{2}} T^{2}(k; k_{0}) e^{2\chi(k_{0})}, & k \in \Gamma_{2}, \end{cases}$$
(2.14)

$$R_{3}^{-}(k) = \begin{cases} \frac{r(k)}{1 - |r(k)|^{2}} \delta_{+}^{2}(k), & k \in (-k_{0}, 0), \\ \frac{r(-k_{0})}{r(-k_{0})} & T^{2}(k; k) e^{2\chi(-k_{0})} & k \in \Gamma \end{cases}$$

$$(2.15)$$

$$\begin{cases}
\frac{1}{1-|r(-k_0)|^2}I^{-}(k;k_0)e^{-k(1-\delta)}, & k \in I_3, \\
R_4(k) = \begin{cases}
-r(k)\delta^{-2}(k), & k \in (-\infty, -k_0), \\
0 & 0 & 0 & 0
\end{cases}$$
(2.16)

$$\begin{cases} -r(-k_0)T^{-2}(k;k_0)e^{-2\chi(-k_0)}, & k \in \Gamma_4, \\ \left( -\overline{r(k)}\delta^2(k), & k \in (-\infty, -k_0), \end{cases} \right) \end{cases}$$

$$(2.10)$$

$$R_{5}(k) = \begin{cases} -\frac{r(k)\sigma(k)}{r(-k_{0})}T^{2}(k;k_{0})e^{2\chi(-k_{0})}, & k \in \Gamma_{5}, \end{cases}$$
(2.17)

$$R_{6}^{-}(k) = \begin{cases} \frac{r(n)}{1 - |r(k)|^{2}} \delta_{-}^{-2}(k), & k \in (-k_{0}, 0), \\ \frac{r(-k_{0})}{1 - |r(-k_{0})|^{2}} T^{-2}(k; k_{0}) e^{-2\chi(-k_{0})}, & k \in \Gamma_{6}, \end{cases}$$
(2.18)

$$R_{6}^{+}(k) = \begin{cases} \frac{r(k)}{1 - |r(k)|^{2}} \delta_{-}^{-2}(k), & k \in (0, k_{0}), \\ r(k_{0}) & \pi^{-2}(k, k_{0}) = -2\chi(k_{0}) & k \in \mathbb{N} \end{cases}$$
(2.19)

$$R_{8}(k) = \begin{cases} -\overline{r(k_{0})}^{2} T^{-2}(k;k_{0}) e^{-2\chi(k_{0})}, & k \in \Gamma_{7}, \\ -\overline{r(k_{0})} \delta^{2}(k), & k \in (k_{0},\infty), \\ -\overline{r(k_{0})} T^{2}(k;k_{0}) e^{2\chi(k_{0})}, & k \in \Gamma_{8}, \end{cases}$$
(2.20)

such that

$$|\bar{\partial}R_j(k)| \le c_1 |r'(Rek)| + c_2 |k \mp k_0|^{-\frac{1}{2}}, \qquad (2.21)$$

for two positive constants  $c_1, c_2$  depended on  $||r||_{H^1(\mathbb{R})}$ .

**Proof.** Define the functions

$$f_1(k) = -r(k_0)T^{-2}(k;k_0)e^{-2\chi(k_0)}\delta^2(k), \qquad k \in \bar{\Omega}_1,$$
  
$$f_3^+(k) = \frac{\overline{r(k_0)}}{1 - |r(k_0)|^2}T^2(k;k_0)e^{2\chi(k_0)}\delta^{-2}(k), \quad k \in \bar{\Omega}_3^+.$$

Then, we can define the extensions for  $k \in \overline{\Omega}_1$ ,

$$R_{1}(k) = \left[f_{1}(k) + \left(-r(\operatorname{Re}k) - f_{1}(k)\right)\mathcal{X}(\phi)\right]\delta^{-2}(k), \qquad (2.22)$$

where  $\phi = \arg(k - k_0)$  and  $\mathcal{X}(\phi)$  is a smooth cut-off function with

$$\mathcal{X}(\phi) = \begin{cases} 1, & \phi \in \left[0, \frac{\pi}{12}\right], \\ 0, & \phi \in \left[\frac{\pi}{9}, \frac{\pi}{6}\right]. \end{cases}$$
(2.23)

It is easy to check that  $R_1(k)$  as constructed has the boundary values (2.13). Let  $k - k_0 = s e^{i\phi}$ . It follows from

$$\bar{\partial} = \frac{1}{2} \left( \frac{\partial}{\partial k_1} + i \frac{\partial}{\partial k_2} \right) = \frac{1}{2} e^{i\phi} \left( \frac{\partial}{\partial s} + \frac{i}{s} \frac{\partial}{\partial \phi} \right)$$

and Proposition 2.1 (iv) that

$$\begin{aligned} |\bar{\partial}R_1(k)| &= -\frac{1}{2}r'(\mathrm{Re}k)\mathcal{X}(\phi)\delta^{-2}(k) + \frac{1}{2}\mathrm{i}\mathrm{e}^{\mathrm{i}\phi}\frac{-r(\mathrm{Re}k) - f_1(k)}{|k - k_0|}\mathcal{X}'(\phi)\delta^{-2}(k) \\ &\leq c_1|r'(\mathrm{Re}k)| + c_2|k - k_0|^{-\frac{1}{2}}. \end{aligned}$$
(2.24)

For  $k \in \overline{\Omega}_3^+$ , let

$$R_{3}^{+}(k) = \left[f_{3}^{+}(k) + \left(\frac{\overline{r(\text{Re}k)}}{1 - |r(\text{Re}k)|^{2}} - f_{3}^{+}(k)\right)\tilde{\mathcal{X}}(\phi)\right]\delta^{2}(k),$$
(2.25)

where  $\tilde{\mathcal{X}}(\phi)$  is a smooth cut-off function with

$$\tilde{\mathcal{X}}(\phi) = \begin{cases} 1, & \phi \in \left[\frac{11\pi}{12}, \pi\right], \\ 0, & \phi \in \left[\frac{3\pi}{4}, \frac{5\pi}{6}\right]. \end{cases}$$
(2.26)

A direct computation implies that  $R_3^+(k)$  satisfies the boundary values (2.14) and the relation (2.21). In other sectors, the extensions can find similarly.  $\Box$ We now use the extensions in Lemma 2.1 to define a new unknown function  $M^{(2)}$  by

 $(0) \qquad \qquad (1) \qquad \qquad (2) \qquad \qquad (2)$ 

$$M^{(2)}(x,t;k) = M^{(1)}(x,t;k)\mathcal{R}^{(2)}(k), \qquad (2.27)$$

where

$$\mathcal{R}^{(2)}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1(k)e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Omega_1, & \begin{pmatrix} 1 & R_3^{\pm}(k)e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Omega_3^{\pm}, \\ \begin{pmatrix} 1 & 0 \\ R_4(k)e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Omega_4, & \begin{pmatrix} 1 & R_5(k)e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Omega_5, \\ \begin{pmatrix} 1 & 0 \\ R_6^{\pm}(k)e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Omega_6^{\pm}, & \begin{pmatrix} 1 & R_8^{\pm}(k)e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Omega_8, \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & k \in \Omega_2 \cup \Omega_7. \end{cases}$$
(2.28)

Let  $\Gamma = {\Gamma_j}_1^8 \cup \frac{ik_0}{\sqrt{3}}(-1, 1)$ . It is an immediate consequence of Lemma 2.1 and RH problem 2.1 that  $M^{(2)}$  satisfies a mixed  $\bar{\partial}$ -Riemann–Hilbert problem.

 $\bar{\partial}$ -Riemann–Hilbert problem 2.2. Find a function  $M^{(2)}(x,t;k)$  with the following properties: 1.  $M^{(2)}(x,t;k)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \Gamma$ .

2. Across  $\Gamma$ , the boundary values satisfy the jump relation

$$M_{+}^{(2)}(x,t;k) = M_{-}^{(2)}(x,t;k)J^{(2)}(x,t;k), \quad k \in \Gamma,$$
(2.29)

where the jump matrix  $J^{(2)}(x,t;k)$  is given by

$$J^{(2)}(x,t;k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ -r(k_0)T^{-2}(k;k_0)e^{-2\chi(k_0)}e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Gamma_1, \\ \begin{pmatrix} 1 & -\frac{\overline{r(k_0)}}{1-|r(k_0)|^2}T^2(k;k_0)e^{2\chi(k_0)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Gamma_2, \\ \begin{pmatrix} 1 & -\frac{\overline{r(-k_0)}}{1-|r(-k_0)|^2}T^2(k;k_0)e^{2\chi(-k_0)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Gamma_3, \\ \begin{pmatrix} 1 & -r(-k_0)T^{-2}(k;k_0)e^{-2\chi(-k_0)}e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Gamma_4, \\ \begin{pmatrix} 1 & \overline{r(-k_0)}T^2(k;k_0)e^{-2\chi(-k_0)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Gamma_5, \end{cases} \\ \begin{pmatrix} \frac{r(-k_0)}{1-|r(-k_0)|^2}T^{-2}(k;k_0)e^{-2\chi(-k_0)}e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Gamma_6, \\ \begin{pmatrix} \frac{r(k_0)}{1-|r(-k_0)|^2}T^{-2}(k;k_0)e^{-2\chi(-k_0)}e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in \Gamma_7, \\ \begin{pmatrix} 1 & \overline{r(k_0)}T^2(k;k_0)e^{2\chi(k_0)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \Gamma_8, \\ \begin{pmatrix} 1 & \overline{r(k_0)}T^2(k;k_0)e^{2\chi(k_0)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in \kappa_8, \\ \begin{pmatrix} 1 & [R_3^-(k) - R_3^+(k)]e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix}, & k \in ik_0 \left(\tan\frac{\pi}{12}, \frac{\sqrt{3}}{3}\right), \\ \begin{pmatrix} R_6^-(k) - R_6^+(k)]e^{t\Phi(k)} & 1 \end{pmatrix}, & k \in ik_0 \left(-\frac{\sqrt{3}}{3}, -\tan\frac{\pi}{12}\right), \\ I, & k \in ik_0 \tan\frac{\pi}{12}(-1, 1). \end{cases}$$

3. For  $k \in \mathbb{C} \setminus \Gamma$ , we have

$$\bar{\partial}M^{(2)}(k) = M^{(2)}(k)\bar{\partial}\mathcal{R}^{(2)}(k).$$
 (2.31)

4.  $M^{(2)}(x,t;k)$  enjoys asymptotics:  $M^{(2)}(x,t;k) = I + O(\frac{1}{k}), \ k \to \infty.$ 

In the next subsection, we aim to extract from  $M^{(2)}$  a contribution that is a pure RH problem. More precisely, we write

$$M^{(2)}(k) = M^{(3)}(k)M^{mod}(k)$$
(2.32)

and we request that  $M^{(3)}(k)$  has no jump across  $\Gamma$ . Thus we look for  $M^{mod}(k)$  solution of the RH problem 2.3 below with the jump matrix  $J^{mod} = J^{(2)}$ .

### 2.3 Analysis of the Riemann–Hilbert problem 2.3

We now focus on  $M^{mod}(k)$ , which satisfies the following RH problem.

**Riemann–Hilbert problem 2.3.** Find a 2 × 2 matrix-valued function  $M^{mod}(x,t;k)$ , analytic on  $\mathbb{C} \setminus \Gamma$  with the following properties:

1.  $M^{mod}(x,t;k)$  is analytic for  $k \in \mathbb{C} \setminus \Gamma$  and is continuous for  $k \in \Gamma$ .

2. The boundary values  $M^{mod}_{\pm}(x,t;k)$  satisfy the jump condition

$$M^{mod}_{+}(x,t;k) = M^{mod}_{-}(x,t;k)J^{mod}(x,t;k), \quad k \in \Gamma,$$
(2.33)

where  $J^{mod}(x,t;k) = J^{(2)}(x,t;k)$ .

3.  $M^{mod}(x,t;k) = I + O(\frac{1}{k})$ , as  $k \to \infty$ .

By using the method of Beals and Coifman in [3], we set

$$J^{mod} = (I - w_{-}^{mod})^{-1} (I + w_{+}^{mod}), \quad w^{mod} = w_{+}^{mod} + w_{-}^{mod},$$

and let

$$(C_{\pm}f)(k) = \int_{\Gamma} \frac{f(s)}{s - k_{\pm}} \frac{\mathrm{d}s}{2\pi \mathrm{i}}, \quad k \in \Gamma, \ f \in L^2(\Gamma),$$

denote the Cauchy operator on  $\Gamma$ . Define the operator  $C_{w^{mod}}: L^2(\Gamma) + L^{\infty}(\Gamma) \to L^2(\Gamma)$  by

$$C_{w^{mod}}f = C_{+}(fw_{-}^{mod}) + C_{-}(fw_{+}^{mod})$$
(2.34)

for any  $2 \times 2$  matrix-valued function f. Let  $\mu^{mod}(x,t;k) \in L^2(\Gamma) + L^{\infty}(\Gamma)$  be the solution of the basic inverse equation

$$\mu^{mod} = I + C_{w^{mod}} \mu^{mod}. \tag{2.35}$$

Then

$$M^{mod}(x,t;k) = I + \int_{\Gamma} \frac{(\mu^{mod}w^{mod})(x,t;s)}{s-k} \frac{\mathrm{d}s}{2\pi \mathrm{i}}, \quad k \in \mathbb{C} \setminus \Gamma$$
(2.36)

is the unique solution of RH problem 2.3.

For a small enough constant  $\varepsilon > 0$ , we define

$$L_{\varepsilon} := \left\{ k \in \mathbb{C} | k = k_0 + k_0 \alpha e^{\frac{5\pi i}{6}}, \varepsilon < \alpha \le \frac{2\sqrt{3}}{3} \right\}$$
$$\cup \left\{ k \in \mathbb{C} | k = -k_0 + k_0 \alpha e^{\frac{\pi i}{6}}, \varepsilon < \alpha \le \frac{2\sqrt{3}}{3} \right\}.$$
(2.37)

Denote  $\Gamma_9 = \frac{ik_0}{\sqrt{3}}(-1,1)$ . Set  $\Gamma' = \Gamma \setminus (\Gamma_9 \cup L_{\varepsilon} \cup \bar{L}_{\varepsilon})$  with the orientation as in Fig. 4. In the following, we shall reduce the RH problem 2.3 to a RH problem on the truncated contour  $\Gamma'$  with controlled error terms. Let  $w^{mod} = w^e + w'$ , where  $w' = w^{mod} \mid_{\Gamma'}$  and  $w^e = w^{mod} \mid_{(\Gamma_9 \cup L_{\varepsilon} \cup \bar{L}_{\varepsilon})}$ .

**Lemma 2.2** For sufficiently small  $\varepsilon > 0$ , we have

$$\|w^e\|_{L^n(\Gamma_9 \cup L_\varepsilon \cup \bar{L}_\varepsilon)} \le c e^{-32\varepsilon^2 \tau}, \quad 1 \le n \le \infty,$$
(2.38)

$$\|w'\|_{L^{\infty}(\Gamma')} \le c e^{-32\varepsilon^2 \tau}.$$
(2.39)

Furthermore,

$$\|w'\|_{L^1(\Gamma')} \le c\tau^{-\frac{1}{2}}, \ \|w'\|_{L^2(\Gamma')} \le c\tau^{-\frac{1}{4}}.$$
 (2.40)



Fig. 4: The oriented contour  $\Gamma'$ .

**Proof.** For  $k \in L_{\varepsilon}$ , we have

$$\operatorname{Re}\Phi(k) = 32\alpha^2 k_0^5 \left( 5\sqrt{3} - 10u + \frac{5\sqrt{3}}{2}u^2 - \frac{1}{2}u^3 \right)$$
  

$$\geq 32\alpha^2 k_0^5 \left( 5\sqrt{3} + (5\sqrt{3} - 10)u - \frac{5\sqrt{3}}{2} - \frac{1}{2}u^3 \right) \geq 32\alpha^2 k_0^5.$$
(2.41)

Thus, we find

$$|R_3^{\pm} e^{-t\Phi}| \le c e^{-32\alpha^2 t k_0^5} \le c e^{-32\varepsilon^2 \tau}.$$
(2.42)

This yields (2.38). On the other hand, a simple calculation implies

$$\|e^{-32\alpha^2 tk_0^5}\|_{L^1(\Gamma')} \le c\tau^{-\frac{1}{2}}, \quad \|e^{-32\alpha^2 tk_0^5}\|_{L^2(\Gamma')} \le c\tau^{-\frac{1}{4}}.$$
(2.43)

The estimate (2.40) immediately follows.

**Lemma 2.3** In the case  $0 < k_0 \leq M$ , as  $\tau \to \infty$ , that is, in Region I,  $(1 - C_{w'})^{-1} : L^2(\Gamma) \to L^2(\Gamma)$  exists and is uniformly bounded:

$$||(1 - C_{w'})^{-1}||_{L^2(\Gamma)} \le c.$$

Moreover,

$$\|(1 - C_{w^{mod}})^{-1}\|_{L^2(\Gamma)} \le c.$$

**Proof.** See [16] and references therein. A simple computation implies that

$$((1 - C_{w^{mod}})^{-1}I)w^{mod} = ((1 - C_{w'})^{-1}I)w' + w^{e} + ((1 - C_{w'})^{-1}(C_{w^{e}}I))w^{mod} + ((1 - C_{w'})^{-1}(C_{w'}I))w^{e} + ((1 - C_{w'})^{-1}C_{w^{e}}(1 - C_{w^{mod}})^{-1})(C_{w^{mod}}I)w^{mod}.$$

However, it follows from Lemma 2.3 that

$$\begin{split} \|w^e\|_{L^1(\Gamma)} &\leq c e^{-32\varepsilon^2 \tau}, \\ \|((1-C_{w'})^{-1}(C_{w^e}I))w^{mod}\|_{L^1(\Gamma)} &\leq \|(1-C_{w'})^{-1}\|_{L^2(\Gamma)} \|C_{w^e}I\|_{L^2(\Sigma)} \|w^{mod}\|_{L^2(\Gamma)} \\ &\leq c \|w^e\|_{L^2(\Sigma)} \|w^{mod}\|_{L^2(\Gamma)} \leq c e^{-32\varepsilon^2 \tau} \tau^{-\frac{1}{4}}, \\ \|(1-C_{w'})^{-1}(C_{w'}I))w^e\|_{L^1(\Gamma)} &\leq \|((1-C_{w'})^{-1}\|_{L^2(\Sigma)} \|C_{w'}I\|_{L^2(\Gamma)} \|w^e\|_{L^2(\Gamma)} \\ &\leq c \|w'\|_{L^2(\Gamma)} \|w^e\|_{L^2(\Gamma)} \leq c e^{-32\varepsilon^2 \tau} \tau^{-\frac{1}{4}}, \\ -C_{w'})^{-1} C_{w^e} (1-C_{w^{mod}})^{-1}) (C_{w^{mod}}I) w^{mod}\|_{L^1(\Gamma)} \leq c \|w^e\|_{L^\infty(\Gamma)} \|w^{mod}\|_{L^2(\Gamma)}^2 \leq c e^{-32\varepsilon^2 \tau} \tau^{-\frac{1}{2}}. \end{split}$$

Thus, as  $\tau \to \infty$ , we have the following result:

 $\|((1$ 

$$\int_{\Gamma} \left( ((1 - C_{w^{mod}})^{-1}I)w^{mod} \right)(x, t; s) \mathrm{d}s = \int_{\Gamma'} \left( ((1 - C_{w'})^{-1}I)w' \right)(x, t; s) \mathrm{d}s + O(e^{-c\tau}).$$
(2.44)

On  $\Gamma'$ , set  $\mu' = (1 - C_{w'})^{-1}I$ . Then it follows that

$$M'(x,t;k) = I + \int_{\Gamma'} \frac{(\mu'w')(x,t;s)}{s-k} \frac{\mathrm{d}s}{2\pi \mathrm{i}}$$
(2.45)

satisfies the following RH problem:

**Riemann–Hilbert problem 2.4.** Find a  $2 \times 2$  matrix-valued function M'(x, t; k) satisfying:

- 1. M'(x,t;k) is analytic for  $k \in \mathbb{C} \setminus \Gamma'$  and is continuous for  $k \in \Gamma'$ .
- 2. The boundary values  $M'_{\pm}(x,t;k)$  satisfy the jump condition

$$M'_{+}(x,t;k) = M'_{-}(x,t;k)J'(x,t;k), \quad k \in \Gamma',$$
(2.46)

where  $J'(x,t;k) = J^{mod}(x,t;k) \mid_{\Gamma'}$ .

3.  $M'(x,t;k) = I + O\left(\frac{1}{k}\right)$ , as  $k \to \infty$ .

Split  $\Gamma'$  into a union of two disjoint crosses,  $\Gamma' = \Gamma'_{-k_0} \cup \Gamma'_{k_0}$ , where  $\Gamma'_{\pm k_0}$  denote the cross of  $\Gamma'$  centered at  $\pm k_0$ . Write  $w' = w'_{-k_0} + w'_{k_0}$ , where  $w'_{-k_0} = 0$  for  $k \in \Gamma'_{k_0}$  and  $w'_{k_0} = 0$  for  $k \in \Gamma'_{-k_0}$ . Define the operators  $C_{w'_{-k_0}}$  and  $C_{w'_{k_0}}$  as in definition (2.34). Then, analogous to Lemma 3.5 in [16], we obtain

$$\begin{aligned} \|C_{w'_{-k_0}}C_{w'_{k_0}}\|_{L^2(\Gamma')} &= \|C_{w'_{k_0}}C_{w'_{-k_0}}\|_{L^2(\Gamma')} \le c\tau^{-\frac{1}{2}}, \\ \|C_{w'_{k_0}}C_{w'_{-k_0}}\|_{L^{\infty}(\Gamma')\to L^2(\Gamma')} \le c\tau^{-\frac{3}{4}}, \\ \|C_{w'_{-k_0}}C_{w'_{k_0}}\|_{L^{\infty}(\Gamma')\to L^2(\Gamma')} \le c\tau^{-\frac{3}{4}}. \end{aligned}$$

$$(2.47)$$

From the identity

$$(1 - C_{w'_{-k_0}} - C_{w'_{k_0}}) \left( 1 + C_{w'_{-k_0}} (1 - C_{w'_{-k_0}})^{-1} + C_{w'_{k_0}} (1 - C_{w'_{k_0}})^{-1} \right)$$
  
=  $1 - C_{w'_{k_0}} C_{w'_{-k_0}} (1 - C_{w'_{-k_0}})^{-1} - C_{w'_{-k_0}} C_{w'_{k_0}} (1 - C_{w'_{k_0}})^{-1},$ 

it follows that

$$(1 - C_{w'})^{-1} = 1 + C_{w'_{-k_0}}(1 - C_{w'_{-k_0}})^{-1} + C_{w'_{k_0}}(1 - C_{w'_{k_0}})^{-1}$$

$$+ \left[1 + C_{w'_{-k_0}}(1 - C_{w'_{-k_0}})^{-1} + C_{w'_{k_0}}(1 - C_{w'_{k_0}})^{-1}\right] \left[1 - C_{w'_{k_0}}C_{w'_{-k_0}}(1 - C_{w'_{-k_0}})^{-1} - C_{w'_{-k_0}}C_{w'_{k_0}}(1 - C_{w'_{k_0}})^{-1}\right]^{-1} \left[C_{w'_{k_0}}C_{w'_{-k_0}}(1 - C_{w'_{-k_0}})^{-1} + C_{w'_{-k_0}}C_{w'_{k_0}}(1 - C_{w'_{k_0}})^{-1}\right]^{-1}$$

According to Lemma 2.2, 2.3 and estimate (2.47), proceeding the estimates as in [16], we obtain an important result stated as follows:

Lemma 2.4 As  $\tau \to \infty$ ,

$$\int_{\Gamma'} \left( ((1 - C_{w'})^{-1}I)w' \right) (x, t; s) ds = \int_{\Gamma'_{-k_0}} \left( ((1 - C_{w'_{-k_0}})^{-1}I)w'_{-k_0} \right) (x, t; s) ds + \int_{\Gamma'_{k_0}} \left( ((1 - C_{w'_{k_0}})^{-1}I)w'_{k_0} \right) (x, t; s) ds + O(\tau^{-1}).$$

$$(2.48)$$

In the following, we aim to perform a scaling transform and then formulate the model RH problem. Extend the contours  $\Gamma'_{-k_0}$  and  $\Gamma'_{k_0}$  to the contours

$$\hat{\Gamma}'_{-k_0} = \{k = -k_0 + k_0 \alpha e^{\pm \frac{i\pi}{6}} : -\infty < \alpha < \infty\},\$$
$$\hat{\Gamma}'_{k_0} = \{k = k_0 + k_0 \alpha e^{\pm \frac{5i\pi}{6}} : -\infty < \alpha < \infty\},\$$

respectively, and define  $\hat{w}'_{-k_0}$ ,  $\hat{w}'_{k_0}$  on  $\hat{\Gamma}'_{-k_0}$ ,  $\hat{\Gamma}'_{k_0}$ , respectively, through

$$\hat{w}_{-k_0}' = \begin{cases} w_{-k_0}', & k \in \Gamma_{-k_0}' \subset \hat{\Gamma}_{-k_0}', \\ 0, & k \in \hat{\Gamma}_{-k_0}' \setminus \Gamma_{-k_0}', \end{cases} \quad \hat{w}_{k_0}' = \begin{cases} w_{k_0}', & k \in \Gamma_{k_0}' \subset \hat{\Gamma}_{k_0}', \\ 0, & k \in \hat{\Gamma}_{k_0}' \setminus \Gamma_{k_0}'. \end{cases}$$

Let  $\Gamma_{-k_0}$  and  $\Gamma_{k_0}$  denote the contours  $\{k = \alpha e^{\pm \frac{i\pi}{6}} : -\infty < \alpha < \infty\}$  centered at original point and oriented as shown in Fig. 5.



Fig. 5: The contours  $\Gamma_{-k_0}$  and  $\Gamma_{k_0}$ .

We next introduce the following scaling operators

$$S_{-k_0}: k \mapsto \frac{z}{8k_0\sqrt{10k_0t}} - k_0,$$
 (2.49)

$$S_{k_0}: k \mapsto \frac{z}{8k_0\sqrt{10k_0t}} + k_0.$$
 (2.50)

Set  $w_{-k_0} = S_{-k_0} \hat{w}'_{-k_0}$ ,  $w_{k_0} = S_{k_0} \hat{w}'_{k_0}$ . A simple change of variables argument shows that

$$C_{\hat{w}'_{-k_0}} = S_{-k_0}^{-1} C_{w_{-k_0}} S_{-k_0}, \quad C_{\hat{w}'_{k_0}} = S_{k_0}^{-1} C_{w_{k_0}} S_{k_0},$$

where  $C_{w_{\pm k_0}}$  is a bounded map from  $\Gamma_{\pm k_0}$  to  $\Gamma_{\pm k_0}$ . On the part

$$L_{k_0} : \{ z = 8k_0^2 \sqrt{10k_0 t} \alpha e^{-\frac{i\pi}{6}} : -\varepsilon < \alpha < \infty \},\$$

we have

$$w_{k_0} = w_{k_0+} = \begin{cases} \begin{pmatrix} 0 & -\frac{\overline{r(k_0)}}{1 - |r(k_0)|^2} (\delta_{k_0}^0)^{-1} (\delta_{k_0}^1)^{-1} \\ 0 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & \overline{r(k_0)} (\delta_{k_0}^0)^{-1} (\delta_{k_0}^1)^{-1} \\ 0 & 0 \end{pmatrix}, \end{cases}$$

and on  $\bar{L}_{k_0}$ ,

$$w_{k_0} = w_{k_0-} = \begin{cases} \begin{pmatrix} 0 & 0 \\ \frac{r(k_0)}{1 - |r(k_0)|^2} \delta_{k_0}^0 \delta_{k_0}^1 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & 0 \\ -r(k_0) \delta_{k_0}^0 \delta_{k_0}^1 & 0 \end{pmatrix}, \end{cases}$$

where

$$\delta_{k_0}^0 = e^{-2\chi(k_0) - 128ik_0^5 t} (2560k_0^5 t)^{i\nu}, \qquad (2.51)$$
  

$$\delta_{k_0}^1 = z^{-2i\nu} \exp\left\{\frac{iz^2}{2} \left(1 + \frac{z}{8k_0^2\sqrt{10k_0t}} + \frac{z^2}{1280k_0^5 t} + \frac{z^3}{51200k_0^7 t\sqrt{10\epsilon k_0t}}\right)\right\} \\ \times \left(\frac{2k_0}{z/8k_0\sqrt{10k_0t} + 2k_0}\right)^{-2i\nu}. \qquad (2.52)$$

If we set  $J^{k_0}(z; r(k_0)) = (I - w_-^{k_0})^{-1}(I + w_+^{k_0})$ , where

$$w^{k_0} = w_+^{k_0} = \begin{cases} \begin{pmatrix} 0 & (\delta_{k_0}^0)^{-1} z^{2i\nu} e^{-\frac{iz^2}{2}} \overline{r(k_0)} \\ 0 & 0 \end{pmatrix}, & z \in \Gamma_{k_0}^1, \\ \begin{pmatrix} 0 & -(\delta_{k_0}^0)^{-1} z^{2i\nu} e^{-\frac{iz^2}{2}} \frac{\overline{r(k_0)}}{1 - |r(k_0)|^2} \\ 0 & 0 \end{pmatrix}, & z \in \Gamma_{k_0}^3, \end{cases}$$
(2.53)

$$w^{k_0} = w_{-}^{k_0} = \begin{cases} \begin{pmatrix} 0 & 0 \\ -\delta_{k_0}^0 z^{-2i\nu} e^{\frac{iz^2}{2}} r(k_0) & 1 \end{pmatrix}, & z \in \Gamma_{k_0}^2, \\ \begin{pmatrix} 0 & 0 \\ \delta_{k_0}^0 z^{-2i\nu} e^{\frac{iz^2}{2}} \frac{r(k_0)}{1+|r(k_0)|^2} & 0 \end{pmatrix}, & z \in \Gamma_{k_0}^4. \end{cases}$$
(2.54)

**Lemma 2.5** Let  $0 < \kappa < 1/2$  be a fixed small number, then we have as  $\tau \to \infty$ 

$$\|\delta_{k_0}^1 - z^{-2i\nu} e^{\frac{iz^2}{2}}\|_{L^{\infty}(\bar{L}_{k_0})} \le c |e^{\frac{i\kappa z^2}{2}} |\tau^{-\frac{1}{2}}.$$
(2.55)

As a result, we have

$$\|\delta_{k_0}^1 - z^{-2i\nu} e^{\frac{iz^2}{2}}\|_{(L^{\infty} \cap L^1 \cap L^2)(\bar{L}_{k_0})} \le c\tau^{-\frac{1}{2}}.$$
(2.56)

 $\mathbf{Proof.}\ \mathrm{Let}$ 

$$\triangle = \frac{z}{8k_0^2\sqrt{10k_0t}(1-2\kappa)} + \frac{z^2}{1280k_0^5t(1-2\kappa)} + \frac{z^3}{51200k_0^7t\sqrt{10k_0t}(1-2\kappa)}.$$

A direct calculation yieds

$$\begin{split} |\delta_{k_0}^1 - z^{-2i\nu} e^{\frac{iz^2}{2}}| &\leq c |e^{\frac{i\kappa z^2}{2}} |\left\{ \left| e^{\frac{i\kappa z^2}{2}} \left[ \left( \frac{2k_0}{z/8k_0\sqrt{10k_0t} + 2k_0} \right)^{-2i\nu} - 1 \right] \right| \\ &+ \left| \exp\left( \frac{i(1-2\kappa)z^2}{2}(1+\Delta) \right) - e^{\frac{i(1-2\kappa)z^2}{2}} \right| \right\} \\ &\leq c |e^{\frac{i\kappa z^2}{2}} |\tau^{-\frac{1}{2}}. \end{split}$$

Therefore, we have

$$\begin{split} &\int_{\Gamma'_{k_0}} \left( \left( (1 - C_{w'_{k_0}})^{-1} I \right) w'_{k_0} \right) (s) \mathrm{d}s = \int_{\hat{\Gamma}'_{k_0}} \left( ((1 - C_{\hat{w}'_{k_0}})^{-1} I) \hat{w}'_{k_0} \right) (s) \mathrm{d}s \\ &= \int_{\hat{\Gamma}'_{k_0}} (S_{k_0}^{-1} (1 - C_{w_{k_0}})^{-1} S_{k_0} I) (s) \hat{w}'_{k_0} (s) \mathrm{d}s \\ &= \int_{\hat{\Gamma}'_{k_0}} \left( (1 - C_{w_{k_0}})^{-1} I) \left( (s - k_0) 8k_0 \sqrt{10k_0 t} \right) (S_{k_0} \hat{w}'_{k_0}) \left( (s - k_0) 8k_0 \sqrt{10k_0 t} \right) \mathrm{d}s \\ &= \frac{1}{8k_0 \sqrt{10k_0 t}} \int_{\Gamma_{k_0}} \left( ((1 - C_{w_{k_0}})^{-1} I) w_{k_0} \right) (s) \mathrm{d}s \\ &= \frac{1}{8k_0 \sqrt{10k_0 t}} \int_{\Gamma_{k_0}} \left( ((1 - C_{w^{k_0}})^{-1} I) w^{k_0} \right) (s) \mathrm{d}s + O(\tau^{-1}). \end{split}$$

Together with a similar computation for  $C_{w'_{-k_0}}$  and (2.44), (2.48), we have obtained as  $\tau \to \infty$ 

$$\int_{\Gamma} \left( ((1 - C_{w^{mod}})^{-1}I)w^{mod} \right)(s) \mathrm{d}s = \frac{1}{8k_0\sqrt{10k_0t}} \left( \int_{\Gamma_{-k_0}} \left( ((1 - C_{w^{-k_0}})^{-1}I)w^{-k_0} \right)(s) \mathrm{d}s + \int_{\Gamma_{k_0}} \left( ((1 - C_{w^{k_0}})^{-1}I)w^{k_0} \right)(s) \mathrm{d}s \right) + O(\tau^{-1}). \quad (2.57)$$

Е		1
L		I
L		I

For  $z \in \mathbb{C} \setminus \Gamma_{k_0}$ , let

$$M^{k_0}(z) = I + \int_{\Gamma_{k_0}} \frac{\left( ((1 - C_{w^{k_0}})^{-1} I) w^{k_0} \right)(s)}{s - z} \frac{\mathrm{d}s}{2\pi \mathrm{i}},$$
(2.58)

then  $M^{k_0}(z)$  satisfies the following RH problem:

**Riemann–Hilbert problem 2.5.** Find a  $2 \times 2$  matrix-valued function  $M^{k_0}(z)$  with the following properties:

- 1.  $M^{k_0}(z)$  is analytic for  $z \in \mathbb{C} \setminus \Gamma_{k_0}$  and is continuous for  $z \in \Gamma_{k_0}$ .
- 2. The boundary values  $M_{\pm}^{k_0}(z)$  satisfy the jump condition

$$M_{+}^{k_{0}}(z) = M_{-}^{k_{0}}(z)J^{k_{0}}(z;r(k_{0})), \quad z \in \Gamma_{k_{0}}.$$
(2.59)

3.  $M^{k_0}(z) = I + O(\frac{1}{z})$ , as  $z \to \infty$ . In particular, if we assume

 $M^{k_0}(z) = I + \frac{M_1^{k_0}}{z} + O(z^{-2}), \quad z \to \infty,$ (2.60)

then

$$M_1^{k_0} = -\int_{\Gamma_{k_0}} \left( ((1 - C_{w^{k_0}})^{-1} I) w^{k_0} \right) (s) \frac{\mathrm{d}s}{2\pi \mathrm{i}}.$$
 (2.61)

There is a analogous RH problem on  $\Gamma_{-k_0}$ , which satisfies

$$\begin{cases} M_{+}^{-k_{0}}(z) = M_{-}^{-k_{0}}(z)J^{-k_{0}}(z;r(-k_{0})), & z \in \Gamma_{-k_{0}}, \\ M^{-k_{0}}(z) \to I, & z \to \infty. \end{cases}$$
(2.62)

According (2.30),  $r(-k) = \overline{r(k)}$  and the analogously computation for  $w^{-k_0}$ , one can find that

$$J^{-k_0}(z; r(-k_0)) = \overline{J^{k_0}(-\bar{z}; r(k_0))}, \qquad (2.63)$$

which in turn implies, by uniqueness, that

$$M^{-k_0}(z) = \overline{M^{k_0}(-\bar{z})}, \quad M_1^{-k_0} = -\overline{M_1^{k_0}}.$$
(2.64)

On the other hand, it follows from [16, 17] that the solution  $M^{k_0}(z)$  of RH problem 2.5 can be explicitly solved in terms of parabolic cylinder functions

$$M^{k_0}(z) = I + \frac{\mathrm{i}}{z} \begin{pmatrix} 0 & -\beta(r(k_0))(\delta_{k_0}^1)^{-1} \\ \overline{\beta(r(k_0))}\delta_{k_0}^1 & 0 \end{pmatrix} + O\left(\frac{1}{z^2}\right),$$
(2.65)

where the function  $\beta(r(k_0))$  is defined by

$$\beta(r(k_0)) = \sqrt{\nu} e^{i(\frac{\pi}{4} - \arg r(k_0) + \arg \Gamma(i\nu))}, \qquad (2.66)$$

and  $\Gamma(\cdot)$  denotes the standard Gamma function. Thus, we have find that

$$M_1^{k_0} = i \begin{pmatrix} 0 & -\beta(r(k_0))(\delta_{k_0}^0)^{-1} \\ \overline{\beta(r(k_0))}\delta_{k_0}^0 & 0 \end{pmatrix}.$$
 (2.67)

Taking into account that (2.36), (2.57), (2.60), (2.64) and (2.65), we get

$$M^{mod}(x,t;k) = I + \frac{M_1^{mod}}{k} + O\left(\frac{1}{k^2}\right), \quad k \to \infty,$$
(2.68)

where

$$M_1^{mod} = \frac{1}{8k_0\sqrt{10k_0t}} (2i\text{Im}M_1^{k_0}) + O(\tau^{-1}), \text{ as } \tau \to \infty.$$
(2.69)

### 2.4 Analysis of the remaining $\bar{\partial}$ problem

Recalling the transform (2.32), it is easy to verify that  $M^{(3)}(k)$  is a continuously differentiable function satisfying the following pure  $\bar{\partial}$  problem.

 $\bar{\partial}$  problem 2.6. Find a function  $M^{(3)}(k)$  with the following properties:

1.  $M^{(3)}(k)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \Gamma$ .

2. For  $k \in \mathbb{C} \setminus \Gamma$ , we have

$$\bar{\partial}M^{(3)}(k) = M^{(3)}(k)W^{(3)}(k),$$

$$W^{(3)}(k) = M^{mod}(k)\bar{\partial}\mathcal{R}^{(2)}(k)[M^{mod}(k)]^{-1}.$$
(2.70)

3.  $M^{(3)}(k)$  admits asymptotics:  $M^{(3)}(k) = I + O(\frac{1}{k}), \ k \to \infty.$ 

It follows from [1] that the  $\bar{\partial}$  problem 2.6 is equivalent to the integral equation

$$M^{(3)}(k) = I - \frac{1}{\pi} \iint_{\mathbb{C}} \frac{M^{(3)}(s)W^{(3)}(s)}{s-k} dA(s),$$
(2.71)

where dA(s) is Lebesgue measure on the plane. Equation (2.71) can be rewritten as the following operator form

$$(1 - K_W)[M^{(3)}(k)] = I, (2.72)$$

where the integral operator  $K_W$  is defined by

$$(K_W f)(k) = -\frac{1}{\pi} \iint_{\mathbb{C}} \frac{f(s)W^{(3)}(s)}{s-k} dA(s).$$
(2.73)

**Proposition 2.2** There exists a constant C > 0 such that for large t > 0, the operator (2.73) obeys the estimate

$$\|K_W\|_{L^{\infty} \to L^{\infty}} \le C(k_0^3 t)^{-\frac{1}{4}}.$$
(2.74)

**Proof.** We only discuss the case in  $\Omega_1$ . We write  $s = k_0 + u + iv$ , then  $u \ge \sqrt{3}v \ge 0$ . Note that

$$\operatorname{Re}\Phi = 32v \left( 10u^2v^2 + 20uv^2k_0 + 10k_0^2v^2 - 5u^4 - 20u^3k_0 - 20uk_0^3 - 30u^2k_0^2 - v^4 \right) \le -640k_0^3uv.$$

Set  $k = \alpha + i\beta$ . Let  $f \in L^{\infty}(\Omega_1)$ , we have

$$|(K_W f)(k)| \le ||f||_{L^{\infty}(\Omega_1)} \iint_{\Omega_1} \frac{|W^{(3)}(s)|}{|s-k|} dA(s) \le C ||f||_{L^{\infty}(\Omega_1)} \iint_{\Omega_1} \frac{|\bar{\partial}R_1(s)|e^{-640k_0^3 tuv}}{|s-k|} dA(s).$$
(2.75)

Thus, we find

$$\|K_W\|_{L^{\infty} \to L^{\infty}} \le C(I_1 + I_2), \tag{2.76}$$

where

$$I_{1} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{|s-k|} |r'(u+k_{0})| e^{-640k_{0}^{3}tuv} dudv,$$
  
$$I_{2} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{|u+iv|^{-\frac{1}{2}}}{|s-k|} e^{-640k_{0}^{3}tuv} dudv.$$

Recalling the estimates from [19], we can similarly get

$$|I_1|, |I_2| \le C(k_0^3 t)^{-\frac{1}{4}}.$$
(2.77)

To recover the long-time asymptotic behavior of u(x,t) using (1.27), it is necessary to determine the asymptotic behavior of the coefficient of the  $k^{-1}$  term in the Laurent expansion of  $M^{(3)}$  at infinity. In fact, we have

$$M^{(3)}(k) = I - \frac{1}{\pi} \iint_{\mathbb{C}} \frac{M^{(3)}(s)W^{(3)}(s)}{s-k} dA(s) = I + \frac{M_1^{(3)}}{k} + \frac{1}{\pi} \iint_{\mathbb{C}} \frac{sM^{(3)}(s)W^{(3)}(s)}{k(s-k)} dA(s), \quad (2.78)$$

where

$$M_1^{(3)} = \frac{1}{\pi} \iint_{\mathbb{C}} M^{(3)}(s) W^{(3)}(s) dA(s).$$
(2.79)

**Proposition 2.3** For all large t > 0, there exists a constant C > 0 such that

$$|M_1^{(3)}| \le C(k_0^3 t)^{-\frac{3}{4}}.$$
(2.80)

**Proof.** We estimate the integral (2.80) as follows:

$$\begin{split} |M_1^{(3)}| &\leq C \iint_{\Omega_1} |\bar{\partial}R_1(s)| e^{-640k_0^3 tuv} \mathrm{d}A(s) \leq C \bigg( \int_0^\infty \int_{\sqrt{3}v}^\infty \left| r'(u+k_0) \right| e^{-640k_0^3 tuv} \mathrm{d}u \mathrm{d}v \\ &+ \int_0^\infty \int_{\sqrt{3}v}^\infty \frac{1}{|u+\mathrm{i}v|^{\frac{1}{2}}} e^{-640k_0^3 tuv} \mathrm{d}u \mathrm{d}v \bigg) \leq C(I_3+I_4). \end{split}$$

It follows from the Proposition D.2 in [5] that  $I_3$  and  $I_4$  satisfy

$$|I_3|, |I_4| \le C(k_0^3 t)^{-\frac{3}{4}}.$$
(2.81)

### **2.5** Asymptotics for u(x,t)

We are now ready to find the asymptotic behavior of the solution u(x,t) to fifth-order modified KdV equation (1.2) in Region I. Working through the transformations, we have

$$M(x,t;k) = M^{(3)}(k)M^{mod}(k)[\mathcal{R}^{(2)}(k)]^{-1}\delta^{\sigma_3}(k).$$
(2.82)

Using the reconstruction formula (1.27) and (2.67)-(2.69), (2.78) and (2.80), we immediately find

$$u(x,t) = \frac{\sqrt{\nu}}{2k_0\sqrt{10k_0t}} \cos\left(128tk_0^5 + \nu\ln(2560tk_0^5) + \varphi(k_0)\right) + O\left(\tau^{-1} + (k_0^3t)^{-\frac{3}{4}}\right), \quad (2.83)$$

where

$$\begin{split} \varphi(k_0) &= -\frac{3\pi}{4} - \arg r(k_0) + \arg \Gamma(i\nu) - \frac{1}{\pi} \int_{-k_0}^{k_0} \ln \left( \frac{1 - |r(s)|^2}{1 - |r(k_0)|^2} \right) \frac{\mathrm{d}s}{s - k_0},\\ \nu &= -\frac{1}{2\pi} \ln(1 - |r(k_0)|^2) > 0. \end{split}$$

## 3 Asymptotics in Region III

We now consider the asymptotics of u(x,t) in the Region III:  $\tau \leq M'$ , that is,  $|x| \leq 80M'^{\frac{4}{5}}t^{\frac{1}{5}}$ . First, for x > 0, the function  $\Phi(k)$  has two real stationary phase points

$$\pm k_0 = \pm \sqrt[4]{\frac{x}{80t}},$$
(3.1)

however, as  $t \to \infty$ , the critical points  $\pm k_0$  approach 0 at least as fast as  $t^{-\frac{1}{5}}$ , i.e.,  $k_0 \leq M'^{\frac{1}{5}}t^{-\frac{1}{5}}$ . Note that the jump matrix J(x,t;k) enjoys the factorization

$$J(x,t;k) = \begin{pmatrix} 1 & -\overline{r(k)}e^{-t\Phi(k)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(k)e^{t\Phi(k)} & 1 \end{pmatrix}.$$
 (3.2)

Introducing the following scaling transform:

$$k \to (20t)^{-\frac{1}{5}}z,$$
 (3.3)

and letting

$$y = \frac{x}{(20t)^{\frac{1}{5}}} \tag{3.4}$$

we then have

$$J(x,t;z) = \begin{pmatrix} 1 & -\overline{r((20t)^{-\frac{1}{5}}z)}e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r((20t)^{-\frac{1}{5}}z)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}.$$
 (3.5)

The first step also is to introduce the continuous but not necessarily analytic extensions. Let

$$z_0 = (20t)^{\frac{1}{5}} k_0. \tag{3.6}$$

Define the new contour  $\Sigma = \Sigma_1 \cup \Sigma_2 \cup \Sigma_3$ , where the line segments

$$\Sigma_{1} = \{z_{0} + le^{\frac{\pi i}{6}} | l \ge 0\} \cup \{-z_{0} + le^{\frac{5\pi i}{6}} | l \ge 0\},$$
  

$$\Sigma_{2} = \{z_{0} + le^{-\frac{\pi i}{6}} | l \ge 0\} \cup \{-z_{0} + le^{-\frac{5\pi i}{6}} | l \ge 0\},$$
  

$$\Sigma_{3} = \{l| - z_{0} \le l \le z_{0}\}$$
(3.7)

oriented with increasing real part and denote the four open sectors  $\{D_j\}_1^4$  in  $\mathbb{C}$ , see Fig. 6.



Fig. 6: The contour  $\Sigma$  and the open sets  $\{D_j\}_1^4$  in the complex z-plane.

**Lemma 3.1** There exist functions  $R_j(z)$  on  $\overline{D}_j$  for j = 1, 4 with boundary values satisfying

$$R_1(z) = \begin{cases} -r((20t)^{-\frac{1}{5}}z), & z \in (-\infty, -z_0) \cup (z_0, \infty), \\ -r(k_0), & z \in \Sigma_1, \end{cases}$$
(3.8)

$$R_4(z) = \begin{cases} -\overline{r((20t)^{-\frac{1}{5}}z)}, & z \in (-\infty, -z_0) \cup (z_0, \infty), \\ -\overline{r(k_0)}, & z \in \Sigma_2, \end{cases}$$
(3.9)

such that

$$|\bar{\partial}R_j(z)| \le c_1 t^{-\frac{1}{5}} |r'((20t)^{-\frac{1}{5}} Rez)| + c_2 t^{-\frac{1}{5}} |(20t)^{-\frac{1}{5}} z - k_0|^{-\frac{1}{2}}$$
(3.10)

for two positive constants  $c_1, c_2$  depended on  $||r||_{H^1(\mathbb{R})}$ .

**Proof.** We only consider  $z \in \overline{\Omega}_1 \cap \{ \operatorname{Re} z > z_0 \}$ . Define the extension

$$R_1(z) = -r(k_0) + \left(r(k_0) - r((20t)^{-\frac{1}{5}} \text{Re}z)\right) \cos(3\phi).$$
(3.11)

Let  $z - z_0 = s e^{i\phi}$ . It follows from

$$\bar{\partial} = \frac{1}{2} \left( \frac{\partial}{\partial z_1} + i \frac{\partial}{\partial z_2} \right) = \frac{1}{2} e^{i\phi} \left( \frac{\partial}{\partial s} + \frac{i}{s} \frac{\partial}{\partial \phi} \right)$$

that

$$\begin{aligned} |\bar{\partial}R_1(z)| &= \left| -\frac{1}{2} (20t)^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Re} z) \cos(3\phi) + \frac{3i}{2} e^{\mathrm{i}\phi} \frac{r((20t)^{-\frac{1}{5}} \operatorname{Re} z) - r(k_0)}{|z - z_0|} \sin(3\phi) \right| \\ &\leq c_1 t^{-\frac{1}{5}} |r'((20t)^{-\frac{1}{5}} \operatorname{Re} z)| + c_2 t^{-\frac{1}{5}} |(20t)^{-\frac{1}{5}} z - k_0|^{-\frac{1}{2}}. \end{aligned}$$
(3.12)

Next, we use the extensions in Lemma 3.1 to define

$$M^{(1)}(y,t;z) = M(x,t;k)\mathcal{R}^{(1)}(z), \qquad (3.13)$$

where

$$\mathcal{R}^{(1)}(z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1(z)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in D_1, \\ \begin{pmatrix} 1 & R_4(z)e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in D_4. \end{cases}$$
(3.14)

Then it is an immediate consequence of Lemma 3.1 and RH problem 1.1 that  $M^{(1)}(y,t;z)$  satisfies the following  $\bar{\partial}$ -Riemann–Hilbert problem.

 $\bar{\partial}$ -Riemann-Hilbert problem 3.1. Find a function  $M^{(1)}(y,t;z)$  with the following properties: 1.  $M^{(1)}(y,t;z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \Sigma$ .

2. Across  $\Sigma$ , the boundary values satisfy the jump relation

$$M_{+}^{(1)}(y,t;z) = M_{-}^{(1)}(y,t;z)J^{(1)}(y,t;z), \quad z \in \Sigma,$$
(3.15)

where the jump matrix  $J^{(1)}(y,t;z)$  is given by

$$J^{(1)}(y,t;z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ r(k_0)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in \Sigma_1, \\ \begin{pmatrix} 1 & -\overline{r(k_0)}e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_2, \\ \begin{pmatrix} 1 & -\overline{r((20t)^{-\frac{1}{5}}z)}e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r((20t)^{-\frac{1}{5}}z)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in \Sigma_3. \end{cases}$$
(3.16)

3. For  $z \in \mathbb{C} \setminus \Sigma$ , we have

$$\bar{\partial}M^{(1)}(z) = M^{(1)}(z)\bar{\partial}\mathcal{R}^{(1)}(z),$$

$$\bar{\partial}R^{(1)}(z) = \begin{cases} \begin{pmatrix} 1 & 0\\ \bar{\partial}R_1(z)\mathrm{e}^{2\mathrm{i}(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in D_1, \\ \begin{pmatrix} 1 & \bar{\partial}R_4(z)\mathrm{e}^{-2\mathrm{i}(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in D_4, \\ 0, & z \in D_2 \cup D_3. \end{cases}$$
(3.17)

4.  $M^{(1)}(y,t;z)$  enjoys asymptotics:  $M^{(1)}(y,t;z) = I + O(\frac{1}{z}), \ z \to \infty.$ 

Let  $M^{\text{RHP}}(y,t;z)$  be the solution of the RH problem resulting from setting  $\bar{\partial}\mathcal{R}^{(1)} \equiv 0$  in  $\bar{\partial}$ -RH problem 3.1. Then the vanishing lemma of Zhou [53] implies that  $M^{\text{RHP}}$  exists and is unique. Moreover, we have the following lemma.

**Lemma 3.2** As  $t \to \infty$ , we have

$$M^{RHP}(y,t;z) = \left(I + O(t^{-\frac{1}{5}})\right) M^{Z}(y;z,z_{0}),$$
(3.18)

where matrix-valued function  $M^{Z}(y; z, z_{0})$  is the solution of model RH problem B.1 with s = r(0)and satisfies

$$M^{Z}(y;z,z_{0}) = I + \frac{M_{1}^{Z}(y)}{z} + O\left(\frac{1}{z^{2}}\right), \quad z \to \infty, \quad (M_{1}^{Z}(y))_{12} = -iu_{p}(y), \tag{3.19}$$

and  $u_p(y)$  denotes the solution of the fourth order Painlevé II equation (A.5).

**Proof.** For  $z = z_0 + le^{\frac{\pi i}{6}}$  and  $z = -z_0 + le^{\frac{5\pi i}{6}}$  with  $l \ge 0, z_0 \ge 0$  and  $y = 4z_0^4$ , we find

$$\operatorname{Re}\left(2\mathrm{i}\left(\frac{4}{5}z^{5} - yz\right)\right) = l^{2}\left(-\frac{4}{5}l^{3} - 4\sqrt{3}z_{0}l^{2} - 16z_{0}^{2}l - 8\sqrt{3}z_{0}^{3}\right).$$

For  $z \in \Sigma_3$ ,  $|e^{\pm 2i(\frac{4}{5}z^5 - yz)}| = 1$ . Then, it follows that

$$\|r(k_0)e^{2i(\frac{4}{5}z^5-yz)} - r(0)e^{2i(\frac{4}{5}z^5-yz)}\|_{L^p(\Sigma)} \le ct^{-\frac{1}{5}}, \quad p \in [1,\infty].$$
(3.20)

A simple calculation shows that the jumps  $J^e$  of the quantity  $e(z) = M^{\text{RHP}}(z)[M^Z(z)]^{-1}$ , satisfy

$$\|J^e - I\|_{L^p(\Sigma)} \le ct^{-\frac{1}{5}}.$$
(3.21)

Using the theory of small-norm RH problems, one shows that e(z) exists and that  $e(z) = I + O(t^{-\frac{1}{5}})$ .

Next we define the ratio

$$M^{(2)}(z) = M^{(1)}(z)[M^{RHP}(z)]^{-1}, (3.22)$$

then it is easy to verify that  $M^{(2)}(z)$  is a continuously differentiable function satisfying the following pure  $\bar{\partial}$  problem.

 $\bar{\partial}$  problem 3.2. Find a function  $M^{(2)}(z)$  with the following properties:

- 1.  $M^{(2)}(z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \Sigma$ .
- 2. For  $z \in \mathbb{C} \setminus \Sigma$ , we have

$$\bar{\partial}M^{(2)}(z) = M^{(2)}(z)W^{(2)}(z),$$

$$W^{(2)}(z) = M^{RHP}(z)\bar{\partial}\mathcal{R}^{(1)}(z)[M^{RHP}(z)]^{-1},$$
(3.23)

where  $\bar{\partial}\mathcal{R}^{(1)}(z)$  is defined by (3.17).

3.  $M^{(2)}(z)$  admits asymptotics:  $M^{(2)}(z) = I + O(\frac{1}{z}), \ z \to \infty.$ 

Proceeding as in the previous section, we find that the  $\bar{\partial}$  problem 3.2 is equivalent to the integral equation

$$M^{(2)}(z) = I - \frac{1}{\pi} \iint_{\mathbb{C}} \frac{M^{(2)}(s)W^{(2)}(s)}{s-z} dA(s).$$
(3.24)

**Proposition 3.4** For sufficiently large t > 0, the operator  $K_W$  defined in (2.73) satisfies

$$||K_W||_{L^{\infty} \to L^{\infty}} \le Ct^{-\frac{1}{10}}.$$
 (3.25)

**Proof.** In  $D_1^1 = D_1 \cap \{ \text{Re} z > z_0 \}$ , we write  $s = z_0 + u + iv$ , then  $u \ge \sqrt{3}v \ge 0$ . We first note that

$$\operatorname{Re}\left(2\mathrm{i}\left(\frac{4}{5}s^{5}-ys\right)\right) = \frac{8}{5}v\left(10u^{2}v^{2}+20uv^{2}z_{0}+10v^{2}z_{0}^{2}-5u^{4}-20u^{3}z_{0}-20uz_{0}^{3}-30u^{2}z_{0}^{2}-v^{4}\right)$$
$$\leq \frac{8}{5}v\left(10u^{2}\cdot\frac{u^{2}}{3}-5u^{4}\right) \leq -2u^{4}v.$$

Set  $z = \alpha + i\beta$ . Let  $f \in L^{\infty}(D_1^1)$ , we have

$$|(K_W f)(z)| \le ||f||_{L^{\infty}(D_1^1)} \iint_{D_1^1} \frac{|W^{(2)}(s)|}{|s-z|} dA(s) \le C ||f||_{L^{\infty}(D_1^1)} \iint_{D_1^1} \frac{|\bar{\partial}R_1(s)|e^{-2u^4v}}{|s-z|} dA(s).$$
(3.26)

Thus, we find

$$\|K_W\|_{L^{\infty} \to L^{\infty}} \le C(I_1 + I_2), \tag{3.27}$$

where

$$I_{1} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{|s-z|} \left| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \text{Res}) \right| e^{-2u^{4}v} du dv,$$
  
$$I_{2} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{|s-z|} \frac{1}{t^{\frac{1}{10}} |u+iv|^{\frac{1}{2}}} e^{-2u^{4}v} du dv.$$

We first note that

$$\left(\int_{\mathbb{R}} \left| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Res}) \right|^2 \mathrm{d}u \right)^{\frac{1}{2}} \le C t^{-\frac{1}{10}}.$$

Recall the bound

$$\left\|\frac{1}{|s-z|}\right\|_{L^2(\sqrt{3}v,\infty)} \le \left(\frac{\pi}{|v-\beta|}\right)^{\frac{1}{2}}.$$

Using these results and Schwarz's inequality on the u-integration we may bound  $I_1$  by

$$|I_1| \le Ct^{-\frac{1}{10}} \int_0^\infty \frac{\mathrm{e}^{-18v^5}}{|v-\beta|^{\frac{1}{2}}} \mathrm{d}v \le Ct^{-\frac{1}{10}}.$$
(3.28)

For p > 2,  $\frac{1}{p} + \frac{1}{q} = 1$ , we recall the estimate from Appendix D of [5],

$$\left\|\frac{1}{|u+\mathrm{i}v|^{\frac{1}{2}}}\right\|_{L^{p}(\sqrt{3}v,\infty)} \leq Cv^{\frac{1}{p}-\frac{1}{2}}, \quad \left\|\frac{1}{|s-z|}\right\|_{L^{q}(\sqrt{3}v,\infty)} \leq C|v-\beta|^{\frac{1}{q}-1}.$$

Thus, we get

$$|I_2| \le Ct^{-\frac{1}{10}} \int_0^\infty v^{\frac{1}{p} - \frac{1}{2}} |v - \beta|^{\frac{1}{q} - 1} e^{-18v^5} dv \le Ct^{-\frac{1}{10}}.$$
(3.29)

Finally, we consider the Laurent expansion of  $M^{(2)}$  as  $z \to \infty$ . In fact, we have

$$M^{(2)}(z) = I + \frac{M_1^{(2)}}{z} + \frac{1}{\pi} \iint_{\mathbb{C}} \frac{sM^{(2)}(s)W^{(2)}(s)}{z(s-z)} dA(s),$$
(3.30)

where

$$M_1^{(2)} = \frac{1}{\pi} \iint_{\mathbb{C}} M^{(2)}(s) W^{(2)}(s) dA(s).$$
(3.31)

**Proposition 3.5** For all large t > 0, the following estimate holds:

$$|M_1^{(2)}| \le Ct^{-\frac{1}{10}}.\tag{3.32}$$

**Proof.** We estimate the integral (3.31) as follows:

$$\begin{split} |M_{1}^{(2)}| &\leq C \iint_{D_{1}^{1}} |\bar{\partial}R_{1}(s)|e^{-2u^{4}v} \mathrm{d}A(s) \leq C \bigg( \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \bigg| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \mathrm{Re}s) \bigg| e^{-2u^{4}v} \mathrm{d}u \mathrm{d}v \\ &+ \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{t^{\frac{1}{10}} |u + \mathrm{i}v|^{\frac{1}{2}}} e^{-2u^{4}v} \mathrm{d}u \mathrm{d}v \bigg) \leq C(I_{3} + I_{4}). \end{split}$$

We bound  ${\cal I}_3$  by applying the Cauchy–Schwarz inequality:

$$|I_{3}| \leq Ct^{-\frac{1}{10}} \int_{0}^{\infty} e^{-9v^{5}} \left( \int_{\sqrt{3}v}^{\infty} e^{-2u^{4}v} du \right)^{\frac{1}{2}} dv$$
  
$$\leq Ct^{-\frac{1}{10}} \sqrt{\Gamma(1/4)} \int_{0}^{\infty} \frac{e^{-9v^{5}}}{\sqrt[8]{2v}} dv \leq Ct^{-\frac{1}{10}}.$$
(3.33)

For  $I_4$ , applying Hölder's inequality, we find

$$|I_4| \le Ct^{-\frac{1}{10}} \int_0^\infty v^{\frac{1}{p} - \frac{1}{2}} e^{-9v^5} \left( \int_{\sqrt{3}v}^\infty e^{-qu^4v} du \right)^{\frac{1}{q}} dv$$
  
$$\le Ct^{-\frac{1}{10}} \int_0^\infty v^{\frac{5}{4p} - \frac{3}{4}} e^{-9v^5} dv \le Ct^{-\frac{1}{10}}.$$
(3.34)

Recalling the transformations (3.13) and (3.22), we have

$$M(x,t;k) = M^{(2)}(z)M^{RHP}(z)[\mathcal{R}^{(1)}(z)]^{-1}.$$
(3.35)

Using the reconstruction formula (1.27) and (3.18)-(3.19), (3.30)-(3.32), we immediately find the asymptotics of the solution u(x,t) in Region III when x > 0:

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O(t^{-\frac{3}{10}}).$$
(3.36)

For x < 0, the two stationary points become

$$\pm k_0 = \pm i \sqrt[4]{\frac{-x}{80t}}.$$
(3.37)

We again perform the scaling  $k \to (20t)^{-\frac{1}{5}}z$  and the contour deformation as shown in Fig. 7. Now



Fig. 7: The contour  $\Sigma$  and the open sets  $\{D_j\}_1^4$  in the complex z-plane.

the  $\bar{\partial}$  extension in  $\bar{D}_1$  turns into

$$R_1(z) = \begin{cases} -r((20t)^{-\frac{1}{5}}z), & z \in (-\infty, \infty), \\ -r(0), & z \in \Sigma_1, \end{cases}$$
(3.38)

and the interpolation is given by

$$R_1(z) = -r(0) + \left( -r((20t)^{-\frac{1}{5}}z) + r(0) \right) \cos(3\phi).$$
(3.39)

Then we have

$$\begin{aligned} |\bar{\partial}R_1(z)| &= \left| -\frac{1}{2} (20t)^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Re} z) \cos(3\phi) + \frac{3\mathrm{i}}{2} \mathrm{e}^{\mathrm{i}\phi} \frac{r((20t)^{-\frac{1}{5}} \operatorname{Re} z) - r(0)}{|z|} \sin(3\phi) \right| \\ &\leq c_1 t^{-\frac{1}{5}} |r'((20t)^{-\frac{1}{5}} \operatorname{Re} z)| + c_2 t^{-\frac{1}{10}} |z|^{-\frac{1}{2}}. \end{aligned}$$
(3.40)

And if we write s = u + iv in  $D_1^1 = D_1 \cap \{ \text{Re}z > 0 \}$ , then  $u \ge \sqrt{3}v \ge 0$ . Note that y < 0, and then

$$\operatorname{Re}\left(2\mathrm{i}\left(\frac{4}{5}s^{5} - ys\right)\right) = \frac{8}{5}v(10u^{2}v^{2} - 5u^{4} - v^{4}) + 2yv \le -2u^{4}v.$$
(3.41)

Thus we can repeat the analysis as the case above for x > 0 and obtain the same long-time asymptotics as (3.36) for the solution u(x, t).

# 4 Asymptotics in Regions II, IV and V

We first derive the asymptotics for the fifth-order modified KdV equation (1.2) in region IV, then V and finally II.

# 4.1 Region IV: $k_0 \leq M, \tau \geq \tilde{M}$

Again, we perform the scaling transformation

$$k \to (20t)^{-\frac{1}{5}}z.$$
 (4.1)

Letting s = u + iv, we find  $\operatorname{Re}\left(2i\left(\frac{4}{5}s^5 - ys\right)\right) = \frac{8}{5}v(10u^2v^2 - 5u^4 - v^4) + 2yv$ . As  $y = -4(20\tau)^{\frac{4}{5}} < 0$ , we can select  $\rho > 0$  which is sufficiently small and independent of y and construct a new contour  $\gamma = \gamma_1 \cup \gamma_2$  given in Fig. 8, such that in sector  $E_1^1$ ,

$$\operatorname{Re}\left(2\mathrm{i}\left(\frac{4}{5}s^{5} - ys\right)\right) \le \frac{8}{5}vu^{2}(10\rho^{2} - 5u^{2}) - 8(20\tau)^{\frac{4}{5}}v \le -u^{2}v < 0.$$

$$(4.2)$$

We define the extension in  $\bar{E}_1^1$  by  $R_1^1(z) = -r((20t)^{-\frac{1}{5}}\text{Re}z)$ , in  $\bar{E}_3^1$  by  $R_3^1(z) = -\overline{r((20t)^{-\frac{1}{5}}\text{Re}z)}$ .



Fig. 8: The contour  $\gamma = \gamma_1 \cup \gamma_2$  and the open sets  $\{E_j\}_1^4$  in the complex z-plane.

Then, we set

$$M^{(1)}(y,t;z) = M(x,t;k)\mathcal{R}^{(1)}(z),$$
(4.3)

where

$$\mathcal{R}^{(1)}(z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1^1(z) e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in E_1^1, \\ \begin{pmatrix} 1 & R_3^1(z) e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in E_3^1 \\ I, & \text{elsewhere.} \end{cases}$$
(4.4)

Thus,  $M^{(1)}(y,t;z)$  satisfies the following  $\bar{\partial}$ -Riemann–Hilbert problem.

 $\bar{\partial}$ -Riemann-Hilbert problem 4.1. Find a function  $M^{(1)}(y,t;z)$  with the following properties: 1.  $M^{(1)}(y,t;z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \{z | \text{Im} z = \pm \rho\}$ .

2. Across  $\text{Im}z = \pm \rho$ , the boundary values satisfy the jump relation

$$M_{+}^{(1)}(y,t;z) = M_{-}^{(1)}(y,t;z)J^{(1)}(y,t;z),$$
(4.5)

where

$$J^{(1)}(y,t;z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ r((20t)^{-\frac{1}{5}} \operatorname{Re} z) e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in \{z | \operatorname{Im} z = \rho\}, \\ \begin{pmatrix} 1 & -\overline{r((20t)^{-\frac{1}{5}} \operatorname{Re} z)} e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in \{z | \operatorname{Im} z = -\rho\}. \end{cases}$$
(4.6)

3. For  $z \in \mathbb{C} \setminus \{z | \text{Im} z = \pm \rho\}$ , we have

$$\bar{\partial}M^{(1)}(z) = M^{(1)}(z)\bar{\partial}\mathcal{R}^{(1)}(z),$$

$$\bar{\partial}R^{(1)}(z) = \begin{cases} \begin{pmatrix} 1 & 0\\ \bar{\partial}R_1^1(z)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in E_1^1, \\ \begin{pmatrix} 1 & \bar{\partial}R_3^1(z)e^{-2i(\frac{4}{5}z^5 - yz)}\\ 0 & 1 \end{pmatrix}, & z \in E_3^1, \\ 0, & \text{elsewhere.} \end{cases}$$

$$(4.7)$$

4.  $M^{(1)}(y,t;z)$  enjoys asymptotics:  $M^{(1)}(y,t;z) = I + O(\frac{1}{z}), z \to \infty$ . Let  $M^{(2)}(y,t;z)$  be the solution of the RH problem resulting from setting  $\bar{\partial}\mathcal{R}^{(1)} \equiv 0$  in  $\bar{\partial}$ -RH problem 4.1. We now define the ratio

$$M^{(3)}(z) = M^{(1)}(z)[M^{(2)}(z)]^{-1},$$
(4.8)

then it is easy to verify that  $M^{(3)}(z)$  satisfies the following pure  $\bar{\partial}$  problem.

 $\bar{\partial}$  problem 4.2. Find a function  $M^{(3)}(z)$  satisfying the following properties:

1.  $M^{(3)}(z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \{z | \text{Im} z = \pm \rho\}$ . 2. For  $z \in \mathbb{C} \setminus \{z | \text{Im} z = \pm \rho\}$ , we have

$$\bar{\partial}M^{(3)}(z) = M^{(3)}(z)W^{(3)}(z), 
W^{(3)}(z) = M^{(2)}(z)\bar{\partial}\mathcal{R}^{(1)}(z)[M^{(2)}(z)]^{-1},$$
(4.9)

where  $\bar{\partial}\mathcal{R}^{(1)}(z)$  is defined by (4.7).

3.  $M^{(3)}(z)$  admits asymptotics:  $M^{(3)}(z) = I + O(\frac{1}{z}), \ z \to \infty$ . Consider the integral

$$\iint_{E_{1}^{1}} \frac{|W^{(3)}(s)|}{|s-z|} dA(s) \leq \iint_{E_{1}^{1}} \frac{|\bar{\partial}R_{1}(s)|e^{-u^{2}v}}{|s-z|} dA(s) \\
\leq \int_{0}^{\rho} \int_{-\infty}^{\infty} \frac{1}{|s-z|} \left| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Res}) \right| e^{-u^{2}v} du dv \\
\leq Ct^{-\frac{1}{10}} \int_{0}^{\rho} \frac{1}{|v-\beta|^{\frac{1}{2}}} dv \leq Ct^{-\frac{1}{10}}.$$
(4.10)

On the other hand, we have

$$\iint_{E_{1}^{1}} |W^{(3)}(s)| dA(s) \leq \iint_{E_{1}^{1}} |\bar{\partial}R_{1}(s)| e^{-u^{2}v} dA(s) \\
\leq \int_{0}^{\rho} \int_{-\infty}^{\infty} \left| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Res}) \right| e^{-u^{2}v} du dv \\
\leq Ct^{-\frac{1}{10}} \int_{0}^{\rho} \frac{1}{\sqrt[4]{v}} dv \leq Ct^{-\frac{1}{10}}.$$
(4.11)

Thus,  $M^{(3)}(z)$  exists and if we expand  $M^{(3)}(z)$  as

$$M^{(3)}(z) = I + \frac{M_1^{(3)}}{z} + O\left(\frac{1}{z^2}\right), \quad z \to \infty,$$
(4.12)

we have

$$|M_1^{(3)}| \le Ct^{-\frac{1}{10}}.\tag{4.13}$$

Next, we analyze  $M^{(2)}(y,t;z)$  which satisfies the conditions 1, 2, 4 of  $\bar{\partial}$ -RH problem 4.1. We aim to deform the contour  $\{z | \text{Im} z = \pm \rho\}$  to  $\gamma = \gamma_1 \cup \gamma_2$ . For this purpose, we introduce a new unknown  $M^{(4)}$  obtained from  $M^{(2)}$  as

$$M^{(4)}(z) = M^{(2)}(z)\mathcal{R}^{(2)}(z).$$
(4.14)

We choose  $\mathcal{R}^{(2)}(z)$  to remove the jump on the contour  $\{z | \text{Im} z = \pm \rho\}$ . More precisely, we define functions  $R_1^2$  and  $R_3^2$  satisfying

$$R_1^2(z) = \begin{cases} -r((20t)^{-\frac{1}{5}}z), & \text{Im}z = \rho, \\ -r(0), & z \in \gamma_1, \end{cases}$$
(4.15)

$$R_3^2(z) = \begin{cases} -\overline{r((20t)^{-\frac{1}{5}}z)}, & \text{Im}z = -\rho, \\ -\overline{r(0)}, & z \in \gamma_2, \end{cases}$$
(4.16)

and then we can select  $\mathcal{R}^{(2)}(z)$  as follows

$$\mathcal{R}^{(2)}(z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1^2(z) e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in E_1^2 \cup E_1^3, \\ \begin{pmatrix} 1 & R_3^2(z) e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in E_3^2 \cup E_3^3, \\ I, & \text{elsewhere.} \end{cases}$$
(4.17)

**Lemma 4.1** There exists functions  $R_1^2(z)$  in  $E_1^2 \cup E_1^3$  and  $R_3^2$  in  $E_3^2 \cup E_3^3$  satisfying the boundary value conditions (4.15)-(4.16), such that

$$|\bar{\partial}R_j^2| \le c_1 t^{-\frac{1}{5}} |r'((20t)^{-\frac{1}{5}}Rez)| + c_2 t^{-\frac{1}{10}} |z|^{-\frac{1}{2}}, \quad j = 1, 3.$$
(4.18)

Thus,  $M^{(4)}(z)$  satisfies the following  $\bar{\partial}$ -Riemann–Hilbert problem.

 $\bar{\partial}$ -Riemann–Hilbert problem 4.3. Find a function  $M^{(4)}(z)$  with the following properties:

1.  $M^{(4)}(z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \gamma$ .

2. Across  $\gamma$ , the boundary values  $M^{(4)}_+(z)$  satisfy the jump relation

$$M_{+}^{(4)}(z) = M_{-}^{(4)}(z)J^{(4)}(y,t;z), \qquad (4.19)$$

where

$$J^{(4)}(y,t;z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ r(0)e^{2i(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in \gamma_1, \\ \begin{pmatrix} 1 & -\overline{r(0)}e^{-2i(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in \gamma_2. \end{cases}$$
(4.20)

3. For  $z \in \mathbb{C} \setminus \gamma$ , we have

$$\bar{\partial}M^{(4)}(z) = M^{(4)}(z)\bar{\partial}\mathcal{R}^{(2)}(z),$$

$$\bar{\partial}R^{(2)}(z) = \begin{cases} \begin{pmatrix} 1 & 0\\ \bar{\partial}R_1^2(z)\mathrm{e}^{2\mathrm{i}(\frac{4}{5}z^5 - yz)} & 1 \end{pmatrix}, & z \in E_1^2 \cup E_1^3, \\ \begin{pmatrix} 1 & \bar{\partial}R_3^2(z)\mathrm{e}^{-2\mathrm{i}(\frac{4}{5}z^5 - yz)} \\ 0 & 1 \end{pmatrix}, & z \in E_3^2 \cup E_3^3, \\ 0, & \text{elsewhere.} \end{cases}$$

$$(4.21)$$

4.  $M^{(4)}(z) = I + O(\frac{1}{z}), \ z \to \infty.$ 

Letting  $\bar{\partial}\mathcal{R}^{(2)} = 0$  in  $\bar{\partial}$ -RH problem 4.3, it easy to see that the remaining pure RH problem is equivalent to the fourth order Painlevé II RH problem A.1 by setting s = r(0) up to a trivial contour deformation. Thus, the ratio

$$M^{(5)}(z) = M^{(4)}(z)[M^P(z)]^{-1}$$
(4.22)

is a continuously differentiable function satisfying the following  $\bar{\partial}$  problem.

 $\bar{\partial}$  problem 4.4. Find a function  $M^{(5)}(z)$  with the following properties:

1.  $M^{(5)}(z)$  is continuous with sectionally continuous first partial derivatives in  $\mathbb{C} \setminus \gamma$ .

2. For  $z \in \mathbb{C} \setminus \gamma$ , we have

$$\bar{\partial}M^{(5)}(z) = M^{(5)}(z)W^{(5)}(z),$$

$$W^{(5)}(z) = M^{P}(z)\bar{\partial}\mathcal{R}^{(2)}(z)[M^{P}(z)]^{-1},$$
(4.23)

where  $\bar{\partial}\mathcal{R}^{(2)}(z)$  is defined by (4.21).

3.  $M^{(5)}(z)$  admits asymptotics:  $M^{(5)}(z) = I + O(\frac{1}{z}), \ z \to \infty.$ 

In order to show the existence of  $M^{(5)}(z)$ , as the discussion in Section 2, we need to check the boundedness of the operator  $K_W$  defined by

$$(K_W f)(z) = -\frac{1}{\pi} \iint_{\mathbb{C}} \frac{f(s) W^{(5)}(s)}{s-z} dA(s).$$
(4.24)

Indeed, we have the following proposition.

**Proposition 4.1** For large time, the integral operator  $K_W$  given by (4.24) obeys the estimate

$$\|K_W\|_{L^{\infty} \to L^{\infty}} \le Ct^{-\frac{1}{10}} e^{-8(20\tau)^{\frac{4}{5}}\rho}.$$
(4.25)

**Proof.** In  $E_1^2$ , we write  $s = u + i(v + \rho)$ , where  $u \ge \sqrt{3}v \ge 0$ . Then we have

$$\operatorname{Re}\left(2\mathrm{i}\left(\frac{4}{5}s^{5}-ys\right)\right) = \frac{8}{5}(v+\rho)\left(10u^{2}(v+\rho)^{2}-5u^{4}-(v+\rho)^{4}\right)+2y(v+\rho)$$
  
$$\leq -u^{4}v-8(20\tau)^{\frac{4}{5}}\rho.$$
(4.26)

Thus, it follows from (4.18) that

$$||K_W||_{L^{\infty} \to L^{\infty}} \le C(I_1 + I_2),$$
(4.27)

where

$$I_{1} = \int_{0}^{\infty} \int_{\sqrt{3}(v+\rho)}^{\infty} \frac{1}{|s-z|} \left| t^{-\frac{1}{5}} r'((20t)^{-\frac{1}{5}} \operatorname{Res}) \right| e^{-u^{4}v - 8(20\tau)^{\frac{4}{5}\rho}} du dv,$$
  

$$I_{2} = \int_{0}^{\infty} \int_{\sqrt{3}(v+\rho)}^{\infty} \frac{1}{|s-z|} \frac{1}{t^{\frac{1}{10}} |u+i(v+\rho)|^{\frac{1}{2}}} e^{-u^{4}v - 8(20\tau)^{\frac{4}{5}\rho}} du dv.$$

Proceeding the same procedure, we can get following estimates

$$|I_1|, |I_2| \le Ct^{-\frac{1}{10}} e^{-8(20\tau)^{\frac{4}{5}}\rho}.$$
(4.28)

Moreover, if we assume that

$$M^{(5)}(z) = I + \frac{M_1^{(5)}}{z} + O\left(\frac{1}{z^2}\right), \ z \to \infty,$$
(4.29)

we then have

$$|M_1^{(5)}| \le Ct^{-\frac{1}{10}} \mathrm{e}^{-8(20\tau)^{\frac{4}{5}}\rho}.$$
(4.30)

Combining all the transforms (4.3), (4.8), (4.14) and (4.22), we find

$$M(x,t;k) = M^{(3)}(z)M^{(5)}(z)M^{P}(z)[\mathcal{R}^{(2)}(z)]^{-1}[\mathcal{R}^{(1)}(z)]^{-1}.$$
(4.31)

We now can use (4.1), (4.12)-(4.13), (4.29)-(4.30) and (A.3)-(A.4) to obtain the long-time asymptotics of u(x,t) in Region IV,

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O(t^{-\frac{3}{10}} e^{-8(20\tau)^{\frac{4}{5}}\rho}).$$
(4.32)

# 4.2 Region V: $k_0 \ge M, x \to -\infty$

Observe that on  $\gamma_1$ ,

$$\|r(0)e^{2i(\frac{4}{5}z^5 - yz)}\|_{L^1 \cap L^2 \cap L^\infty} \le Ce^{-8(20\tau)^{\frac{4}{5}}\rho}.$$
(4.33)

However, we may choose  $\rho \ge (20\tau)^{\frac{1}{5}}$ . It then follows that in Region V

$$u(x,t) = O\left(t^{-\frac{1}{5}}e^{-c\tau} + t^{-\frac{3}{10}}e^{-8(20\tau)^{\frac{4}{5}}\rho}\right).$$
(4.34)

# 4.3 Region II: $o(t^{\frac{2}{7}}) = \tau \ge \tilde{M}$

We now scale:

$$k \to k_0 z, \tag{4.35}$$

and then have

$$J(x,t;z) = \begin{pmatrix} 1 & -\overline{r(k_0 z)} e^{-32i\tau(z^5 - 5z)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(k_0 z) e^{32i\tau(z^5 - 5z)} & 1 \end{pmatrix}.$$
 (4.36)

Construct the contour  $\Upsilon$  as shown in Fig. 9, and define the function  $R_1(z)$  in  $F_1$  with the boundary values

$$R_1(z) = \begin{cases} -r(k_0 z), & z \in (-\infty, -1) \cup (1, \infty), \\ -r(k_0), & z \in \Upsilon_1. \end{cases}$$
(4.37)

In fact, we can choose

$$R_1(z) = -r(k_0) + \left(r(k_0) - r(k_0 \operatorname{Re} z)\right) \cos(3\phi).$$
(4.38)

A simple computation shows that

$$|\bar{\partial}R_1(z)| \le c_1 k_0 r'(k_0 \operatorname{Re} z) + c_2 k_0^{\frac{1}{2}} |z - 1|^{-\frac{1}{2}}.$$
(4.39)

However, if we write z = u + 1 + iv in  $F_1 \cap \{\text{Re}z > 1\}$ , then one can get

$$\operatorname{Re}\left(32\mathrm{i}\tau(z^{5}-5z)\right) = 32\tau v(10u^{2}v^{2}+20uv^{2}+10v^{2}-5u^{4}-20u^{3}-30u^{2}-20u-v^{4})$$
  
$$\leq -2\tau u^{4}v.$$
(4.40)

On the other hand, on  $\Upsilon_1$ , for  $\tau \geq \tilde{M}$ , we have

$$\|r(k_0)e^{32i\tau(z^5-5z)} - r(0)e^{32i\tau(z^5-5z)}\|_{L^p} \le ck_0, \quad p \in [1,\infty].$$
(4.41)

Following the notation and analysis of Section 3, we then only need to give the following estimates. **Proposition 4.2** The integral operator  $K_W$  given by (2.73) obeys the estimate

$$\|K_W\|_{L^{\infty} \to L^{\infty}} \le Ck_0^{\frac{1}{2}} \tau^{-\frac{1}{10}}.$$
(4.42)



Fig. 9: The contour  $\Upsilon$  and the open sets  $\{F_j\}_1^4$  in the complex z-plane.

**Proof.** In  $F_1^1 := F_1 \cap \{ \text{Re}z > 1 \}$ , it follows from (4.39)-(4.40) that

$$||K_W||_{L^{\infty} \to L^{\infty}} \le C(I_1 + I_2),$$
(4.43)

where

$$I_{1} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{|s-z|} \left| k_{0}r'(k_{0} \operatorname{Re} s) \right| e^{-2\tau u^{4}v} \mathrm{d}u \mathrm{d}v,$$
  
$$I_{2} = \int_{0}^{\infty} \int_{\sqrt{3}v}^{\infty} \frac{1}{|s-z|} \frac{k_{0}^{\frac{1}{2}}}{|u+\mathrm{i}v|^{\frac{1}{2}}} e^{-2\tau u^{4}v} \mathrm{d}u \mathrm{d}v.$$

For  $I_1$ , we can get following estimates

$$|I_1| \le Ck_0^{\frac{1}{2}} \int_0^\infty \frac{\mathrm{e}^{-18\tau v^5}}{|v-\beta|^{\frac{1}{2}}} \mathrm{d}v \le Ck_0^{\frac{1}{2}} \tau^{-\frac{1}{10}} \int_0^\infty \frac{\mathrm{e}^{-18(w+\tau^{\frac{1}{5}}\beta)^5}}{|w|^{\frac{1}{2}}} \mathrm{d}w \le Ck_0^{\frac{1}{2}} \tau^{-\frac{1}{10}}.$$
 (4.44)

For  $I_2$ , we have

$$|I_2| \le k_0^{\frac{1}{2}} \int_0^\infty e^{-18\tau v^5} v^{\frac{1}{p} - \frac{1}{2}} |v - \beta|^{\frac{1}{q} - 1} dv.$$
(4.45)

Observe that  $e^{-m} \le m^{-\frac{1}{10}}$ , thus, one can get

$$\int_{0}^{\beta} e^{-18\tau v^{5}} v^{\frac{1}{p}-\frac{1}{2}} |v-\beta|^{\frac{1}{q}-1} dv = \int_{0}^{1} \beta^{\frac{1}{2}} e^{-18\tau \beta^{5} w^{5}} w^{\frac{1}{p}-\frac{1}{2}} (1-w)^{\frac{1}{q}-1} dw$$
$$\leq \tau^{-\frac{1}{10}} \int_{0}^{1} w^{\frac{1}{p}-1} (1-w)^{\frac{1}{q}-1} dw \leq \tau^{-\frac{1}{10}}$$

Finally, we have

$$\int_{\beta}^{\infty} e^{-18\tau v^5} v^{\frac{1}{p} - \frac{1}{2}} (v - \beta)^{\frac{1}{q} - 1} dv \le \int_{0}^{\infty} e^{-18\tau w^5} w^{-\frac{1}{2}} dw \le \tau^{-\frac{1}{10}}.$$

Next, we estimate the integral defined in (3.31).

**Proposition 4.3** For all large t > 0, we find

$$|M_1^{(2)}| \le Ck_0^{\frac{1}{2}}\tau^{-\frac{3}{10}}.$$
(4.46)

**Proof.** We estimate the integral (3.31) as follows:

$$\begin{split} |M_1^{(2)}| &\leq C \iint_{F_1^1} |\bar{\partial}R_1(s)| e^{-2\tau u^4 v} \mathrm{d}A(s) \leq C \bigg( \int_0^\infty \int_{\sqrt{3}v}^\infty \bigg| k_0 r'(k_0 \mathrm{Re}s) \bigg| e^{-2\tau u^4 v} \mathrm{d}u \mathrm{d}v \\ &+ \int_0^\infty \int_{\sqrt{3}v}^\infty \frac{k_0^{\frac{1}{2}}}{|u+\mathrm{i}v|^{\frac{1}{2}}} e^{-2\tau u^4 v} \mathrm{d}u \mathrm{d}v \bigg) \leq C(I_3 + I_4). \end{split}$$

We bound  $I_3$  as follows:

$$|I_{3}| \leq Ck_{0}^{\frac{1}{2}} \int_{0}^{\infty} e^{-9\tau v^{5}} \left( \int_{\sqrt{3}v}^{\infty} e^{-2\tau u^{4}v} du \right)^{\frac{1}{2}} dv$$
  
$$\leq Ck_{0}^{\frac{1}{2}} \sqrt{\Gamma(1/4)} \int_{0}^{\infty} \frac{e^{-9\tau v^{5}}}{\sqrt[8]{2}\tau v} dv \leq Ck_{0}^{\frac{1}{2}} \tau^{-\frac{3}{10}}.$$
 (4.47)

For  $I_4$ , applying Hölder's inequality, we find

$$|I_4| \le Ck_0^{\frac{1}{2}} \int_0^\infty v^{\frac{1}{p} - \frac{1}{2}} e^{-9\tau v^5} \left( \int_{\sqrt{3}v}^\infty e^{-q\tau u^4 v} du \right)^{\frac{1}{q}} dv$$
  
$$\le Ck_0^{\frac{1}{2}} \tau^{-\frac{1}{4q}} \int_0^\infty v^{\frac{5}{4p} - \frac{3}{4}} e^{-9\tau v^5} dv \le Ck_0^{\frac{1}{2}} \tau^{-\frac{3}{10}}.$$
(4.48)

Using these estimates and the scaling transform (4.35), we immediately find the asymptotics of the solution u(x, t) in Region II:

$$u(x,t) = \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O(k_0^{\frac{3}{2}}\tau^{-\frac{3}{10}} + k_0^2)$$
$$= \left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) + O\left(t^{-\frac{3}{10}} + \left(\frac{\tau}{t}\right)^{\frac{2}{5}}\right).$$
(4.49)

**Remark 4.2** We show how to match the asymptotic formulas of solution u(x,t) in the overlaps of Regions I and II. It is easy to see that the final model RH problem on  $\Upsilon$  corresponding to a special case of in Region I in which one replace

$$t \to \tau$$
,  $k_0 \to 1$ ,  $r(z) \to r(0)$ .

Then, we find as  $\tau \to \infty$ ,

$$\left(\frac{8}{5t}\right)^{\frac{1}{5}} u_p\left(\frac{x}{(20t)^{\frac{1}{5}}}\right) = k_0 \frac{\sqrt{\nu(r(0))}}{2\sqrt{10\tau}} \cos\left(128\tau + \nu(r(0))\ln(2560\tau)\right)$$

$$-\frac{3\pi}{4} - \arg r(0) + \arg \Gamma(i\nu(r(0))) + k_0 O\left(\tau^{-\frac{3}{4}}\right), \qquad (4.50)$$

and thus in Region II

$$u(x,t) = \frac{\sqrt{\nu(r(0))}}{2k_0\sqrt{10tk_0}} \cos\left(128\tau + \nu(r(0))\ln(2560\tau) - \frac{3\pi}{4} - \arg r(0) + \arg \Gamma(i\nu(r(0)))\right) + k_0 O\left(\tau^{-\frac{3}{4}}\right) + O\left(\left(\frac{\tau}{t}\right)^{\frac{2}{5}}\right).$$
(4.51)

On the other hand, in Region I, it follows from the similar analysis in [16] that

$$u(x,t) = \frac{\sqrt{\nu(r(0))}}{2k_0\sqrt{10k_0t}} \cos\left(128\tau + \nu(r(0))\ln(2560\tau) - \frac{3\pi}{4} - \arg r(0) + \arg \Gamma(i\nu(r(0)))\right) + (tk_0^3)^{-\frac{1}{2}}O\left(\tau^{-1}(tk_0^3)^{\frac{1}{2}} + (k_0^3t)^{-\frac{1}{4}} + k_0\ln\tau\right).$$
(4.52)

However, for  $\tau = o(t^{\frac{2}{7}})$ , we find

$$k_0 \tau^{-\frac{3}{4}} = (tk_0^3)^{-\frac{1}{2}} \tau^{-\frac{1}{4}}, \quad k_0 \ln \tau = o\left(\frac{\ln t}{t^{\frac{1}{7}}}\right), \quad \left(\frac{\tau}{t}\right)^{\frac{2}{5}} = (tk_0^3)^{-\frac{1}{2}} \left(\frac{\tau^{\frac{7}{2}}}{t}\right)^{\frac{1}{5}}.$$
 (4.53)

Therefore, we have

$$|u_I(x,t) - u_{II}(x,t)| = (tk_0^3)^{-\frac{1}{2}}O\left(\tau^{-\frac{1}{4}} + \left(\frac{\tau^{\frac{7}{2}}}{t}\right)^{\frac{1}{5}} + \frac{\ln t}{t^{\frac{1}{7}}} + \tau^{-1}(tk_0^3)^{\frac{1}{2}} + (k_0^3t)^{-\frac{1}{4}}\right)$$
(4.54)

which is of order  $o(\sup |u_{as}(x,t)|)$  as  $\tau \to \infty$  with  $\tau = o(t^{\frac{2}{7}})$ .

# 5 Asymptotic behavior in low regularity spaces

In this section, we first obtain the global well-posedness for the initial value problem of Equation (1.2), and then extend the long-time asymptotic behavior of the solution to the low regularity spaces  $H^{s,1}$ ,  $19/22 < s \leq 1$ .

 $X^{s,b}$  method is extremely useful for studying the Cauchy problem of rough initial data. The motivation behind  $X^{s,b}$  method is based on the dispersion relation. We consider the corresponding linear equation of fifth-order KdV equation:

$$\partial_t u + \partial_x^5 u = 0.$$

Taking Fourier transform with respect to both space and time variable, we can get

$$(\tau - \xi^5)\widehat{u}(\xi, \tau) = 0.$$

It is easy to see that  $\hat{u}(\xi, \tau)$  is supported on the surface  $\{(\tau, \xi) : \tau = \xi^5\}$ .  $\tau = \xi^5$  and  $|\tau - \xi^5|$  are called the dispersion relation and dispersion modulation, respectively. Let  $s, b \in \mathbb{R}$ , the Bourgain space  $X^{s,b}_{\tau=\xi^5}(\mathbb{R} \times \mathbb{R})$ , or simply denoted as  $X^{s,b}$ , is defined to be the closure of the Schwartz functions

 $\mathscr{S}(\mathbb{R} \times \mathbb{R})$  under the norm

$$\|u\|_{X^{s,b}_{\tau=\xi^5}} = \left\| \langle \xi \rangle^s \langle \tau - \xi^5 \rangle^b \widehat{u}(\xi,\tau) \right\|_{L^2_{\tau,\xi}}.$$

We will also need the truncated version of the above norm:

$$\|u\|_{X^{s,b}_{\delta}} = \inf_{\tilde{u}=u \text{ on } t \in [0,\delta]} \|\tilde{u}\|_{X^{s,b}}.$$
(5.1)

### 5.1 Estimates in Bourgain spaces

Let  $\psi : \mathbb{R} \to [0,1]$  denote an even smooth function supported in [-2,2] and equal to 1 in [-1,1]. Denote  $\psi_{\mu}(\cdot) = \psi(\cdot/\mu)$  for any  $\mu > 0$ .  $X^{s,b}$  method reduces to the following crucial estimates.

**Lemma 5.1** For  $s \in \mathbb{R}$  and  $1/2 < b < b' < 1/2 + \epsilon$ , then

- (1) (Embedding) For any  $u \in X^{s,b}$ , we have  $||u||_{L^{\infty}(\mathbb{R};H^{s}(\mathbb{R}))} \lesssim ||u||_{X^{s,b}}$ ;
- (2) (Homogeneous estimate)  $\|\psi(t)e^{-t\partial_x^5}u_0\|_{X^{s,b}} \lesssim \|u_0\|_{H^s};$
- (3) (Inhomogeneous estimate)  $\|\psi(t)\int_0^t e^{-(t-t')\partial_x^5}F(u)(t') dt'\|_{X^{s,b}} \lesssim \|F(u)\|_{X^{s,b-1}};$
- (4) (Property) Let  $0 < \mu < 1$ , then  $\|\psi_{\mu}(t)F\|_{X^{s,b-1}} \lesssim \mu^{b'-b} \|F\|_{X^{s,b'-1}}$ ;
- (5) (Nonlinear estimates) Let  $s \ge 3/4$ , we have

$$\begin{aligned} \|u_1 u_2 \partial_x^3 u_3\|_{X^{s,b'-1}} + \|\partial_x u_1 \partial_x u_2 \partial_x u_3\|_{X^{s,b'-1}} + \|u_1 \partial_x u_2 \partial_x^2 u_3\|_{X^{s,b'-1}} \lesssim \prod_{j=1}^3 \|u_j\|_{X^{s,b}}; \\ \|\partial_x (u_1 u_2 u_3 u_4 u_5)\|_{X^{s,b'-1}} \lesssim \prod_{j=1}^5 \|u_j\|_{X^{s,b}}. \end{aligned}$$

**Proof.** The proofs of (1-4) can be found in many literatures, we refer to [34,38,48]. The nonlinear estimate (5) needs condition  $s \ge 3/4$ , and its proof was followed by [38].

We recall some important estimates for the fifth-order KdV equations. These estimates are due to semi-group estimates and the Extension Lemma of [48], their proofs can be obtained by the methods in [15, 25, 32, 33].

**Lemma 5.2** Let  $u \in \mathscr{S}(\mathbb{R} \times \mathbb{R})$ , then we have

(1) (Strichartz estimates) Let  $2 \le q, r \le \infty$  and 2/q + 1/r = 1/2, then

$$\|D_x^{3/q}u\|_{L^q_t L^r_x} \lesssim \|u\|_{X^{0,\frac{1}{2}+}}.$$
(5.2)

(2) (Local smoothing effect estimates)

$$\|D_x^2 u\|_{L^{\infty}_x L^2_t} \lesssim \|u\|_{X^{0,\frac{1}{2}+}}.$$
(5.3)

(3) (Maximal function estimates)

$$\|u\|_{L^4_x L^\infty_t} \lesssim \|u\|_{X^{\frac{1}{4}, \frac{1}{2}+}}.$$
(5.4)

**Lemma 5.3** (Bilinear estimates) Suppose  $u, v \in X^{0,\frac{1}{2}+}$  be supported on spatial frequencies on  $|\xi| \sim N_1, N_2$ , respectively. Let  $N_1 \gg N_2$ , then we have

$$\|uv\|_{L^2_{x,t}} \lesssim N_1^{-2} \|u\|_{X^{0,\frac{1}{2}+}} \|v\|_{X^{0,\frac{1}{2}+}}.$$
(5.5)

### **5.2** Local well-posedness in $H^s, s \ge 3/4$

In this subsection, we use the  $X^{s,b}$  method and the contraction principle to obtain the local existence of the solution in Sobolev space  $H^s, s \ge 3/4$ . This result was mainly stated by S. Kwon [38]. For the sake of completeness, we give the outline of its proof.

**Theorem 5.1** Let  $s \ge 3/4$  and  $u_0 \in H^s(\mathbb{R})$ . Then there exists a time  $T = T(||u_0||_{H^s(\mathbb{R})}) > 0$  such that (1.2) has a unique solution in  $C([0,T]; H^s(\mathbb{R}))$ . Moreover, the solution map from data to the solutions is real-analytic.

**Proof.** We construct the mapping:

$$\mathfrak{M}: u(x,t) \longrightarrow \psi(t) \mathrm{e}^{-t\partial_x^5} u_0 - \psi(t) \int_0^t \mathrm{e}^{-(t-t')\partial_x^5} \psi_T(t') F(u)(t') \,\mathrm{d}t', \tag{5.6}$$

and show that it is a contraction if T < 1 is sufficiently small. Assume  $u_0 \in H^s(s \ge 3/4)$ , we define the metric space:

$$\mathfrak{D} = \left\{ u : \|u\|_{X^{s,b}} \le 2C \|u_0\|_{H^s} \right\}; \qquad d(u,v) = \|u-v\|_{X^{s,b}}$$

Therefore, we have from the estimates in Lemma 5.1 that

$$\begin{aligned} \|\mathfrak{M}u\|_{X^{s,b}} &\leq C \|u_0\|_{H^s} + CT^{b'-b} \|u\|_{X^{s,b}}^3 + CT^{b'-b} \|u\|_{X^{s,b}}^5 \\ &\leq C \|u_0\|_{H^s} + CT^{b'-b} \big( (2C)^3 \|u_0\|_{H^s}^2 + (2C)^5 \|u_0\|_{H^s}^4 \big) \|u_0\|_{H^s}. \end{aligned}$$
(5.7)

Thus we choose T small enough such that

$$T^{b'-b} ((2C)^3 \|u_0\|_{H^s}^2 + (2C)^5 \|u_0\|_{H^s}^4) < 1/2,$$

then we know that  $\mathfrak{M}u \in \mathfrak{D}$ . Similarly, assume  $(u_1, u_2) \in \mathfrak{D}$ , we have

$$\|\mathfrak{M}u_1 - \mathfrak{M}u_2\|_{X^{s,b}} \le 1/2 \|u_1 - u_2\|_{X^{s,b}}.$$

This implies the local well-posedness for large data. By similar argument, we can get the solution map is even real-analytic.  $\hfill \Box$ 

### 5.3 Global well-posedness in $H^s$ , $19/22 < s \le 1$

As mentioned in the introduction, the global well-posedness of (1.2) in  $H^1(\mathbb{R})$  can be immediately obtained by energy conservation. One can conjecture that (1.2) is in fact globally well-posed in time from all initial data contained in the local theory. The biggest obstacle in getting global solutions in  $H^s$  with 0 < s < 1 is the lack of any conservation law. Therefore, in this subsection we introduce an "almost conservation law" by utilizing *I*-method, then we can get the global well-posedness. First of all, we introduce the definition of *I*-operator. Given s < 1, assume  $\varphi(\xi) : \mathbb{R} \to \mathbb{R}$  is a smooth, even, real-valued function, which is equal to 1 in [-1, 1], and defined by  $|\xi|^{s-1}$  in the set  $\{\xi : |\xi| \ge 2\}$ . For a parameter  $N \gg 1$ , define

$$m_N(\xi) := \varphi(\xi/N) = \begin{cases} 1 & |\xi| \le N; \\ \left(\frac{N}{|\xi|}\right)^{1-s} & |\xi| \ge 2N. \end{cases}$$
(5.8)

and the *I*-operator as follows:

$$I_N u = \mathscr{F}_{\xi}^{-1} m_N(\xi) \mathscr{F} u \tag{5.9}$$

For convenient, we will omit the parameter N, writing  $m(\xi)$  and Iu in (5.8) and (5.9).

Recall the energy E(u) in (1.3), the quantity E(Iu)(t) can be compared with  $||u(\cdot, t)||_{H^s}$ , indeed, we have

$$E(Iu)(t) \lesssim N^{2(1-s)} \|u(\cdot,t)\|_{\dot{H}^s}^2 + \|u(\cdot,t)\|_{L^4}^4;$$
(5.10)

$$\|u(\cdot,t)\|_{H^s}^2 \lesssim E(Iu)(t) + \|u_0\|_{L^2}^2.$$
(5.11)

Moreover, we will show that E(Iu)(t) is an "almost conservation law" in the  $H^s$  level, and its increment can be described as the following proposition, whose proof shall be given later.

**Proposition 5.1** Let 19/22 < s < 1,  $N \gg 1$ . Given initial data  $u_0 \in C_0^{\infty}(\mathbb{R})$  with  $E(Iu_0) \leq 1$ , then there exists a  $\delta = \delta(||u_0||_{L^2}) > 0$  so that the solution

$$u(x,t) \in C([0,\delta], H^s(\mathbb{R}))$$
(5.12)

of (1.2) satisfies

$$E(Iu)(t) = E(Iu)(0) + O(N^{-1/2}) \quad for \ all \ t \in [0, \delta].$$
(5.13)

**Proof of Theorem 1.2.** We only need to consider 19/22 < s < 1. We may assume  $u_0 \in C_0^{\infty}(\mathbb{R})$ , and then show that the solution in the  $H^s$  norm grows at most polynomially, that is to say there exist  $C_1$ ,  $C_2$  and M, which depend only on  $||u_0||_{H^s}$  such that

$$\|u(\cdot,t)\|_{H^s} \le C_1 t^M + C_2. \tag{5.14}$$

Then the global result is immediately obtained by (5.14), the local well-posedness and a standard density argument. From (5.11), we only need to show that

$$E(Iu)(t) \lesssim (1+t)^M.$$
 (5.15)

Recall the scaling symmetry, let  $0 < \lambda < 1$ , then (5.10) and the Sobolev embedding inequality imply that

$$E(Iu_{0,\lambda}) \lesssim N^{2(1-s)} \lambda^{2s+1} \|u_0\|_{\dot{H}^s}^2 + \lambda^3 \|u_0\|_{L^4}^4 \lesssim N^{2(1-s)} \lambda^{2s+1} (1 + \|u_0\|_{H^s})^4.$$
(5.16)

Assuming  $N \gg 1$  is given, we choose  $\lambda = \lambda(N, ||u_0||_{H^s})$ 

$$\lambda = N^{-\frac{2(1-s)}{2s+1}} (2C_0)^{-\frac{1}{2s+1}} \left(1 + \|u_0\|_{H^s}\right)^{-\frac{4}{2s+1}}$$
(5.17)

such that  $E(Iu_{0,\lambda}) \leq 1/2$ , where  $C_0$  is the implicit constant in (5.16). We apply Proposition 5.1 at least  $CN^{\frac{1}{2}}$  times to get

$$E(Iu_{\lambda})(CN^{\frac{1}{2}}\delta) \sim 1.$$
(5.18)

For any  $T_0 \gg 1$ , we choose  $N \gg 1$  so that

$$T_0 \sim \lambda^5 N^{\frac{1}{2}} \delta \sim N^{\frac{22s-19}{2(2s+1)}},\tag{5.19}$$

where we used (5.17). In order to keep the exponent of N is positive, we need the condition s > 19/22. From the scaling symmetry, we have that

$$E(Iu)(t) \le \lambda^{-3} E(Iu_{\lambda})(\lambda^{-5}t).$$
(5.20)

Therefore, from (5.17), (5.19) and (5.20), for any  $T_0 \gg 1$ , we have

$$E(Iu)(T_0) \le CT_0^{\frac{12(1-s)}{22s-19}},$$
(5.21)

where N is determined by (5.19) and  $C = C(||u_0||_{H^s}, \delta)$ . We now get the conclusion (5.15), and complete the proof.

To prove Proposition 5.1, we need the following variant local well-posedness result:

**Proposition 5.2** Let  $3/4 \le s < 1$ . Assume  $u_0$  satisfies  $E(Iu_0) \le 1$ , then there exists a constant  $\delta = \delta(||u_0||_{L^2})$  and a unique solution u to (1.2) such that

$$\|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}} \lesssim 1, \tag{5.22}$$

where the implicit constant is independent of N.

**Proof.** We have the following estimates, which are analogous to Lemma 5.1 (2-3),

$$\left\| e^{-t\partial_x^5} u_0 \right\|_{X^{1,\frac{1}{2}+}_{\delta}} \lesssim \|u_0\|_{H^1};$$
(5.23)

$$\left\| \int_{0}^{t} e^{-(t-t')\partial_{x}^{5}} F(u)(t') dt' \right\|_{X_{\delta}^{1,\frac{1}{2}+}} \lesssim \|F(u)\|_{X_{\delta}^{1,-\frac{1}{2}+}}.$$
(5.24)

Iu satisfies the integral equation on  $t \in [0, \delta]$ :

$$Iu(x,t) = e^{-t\partial_x^5} Iu_0 - \int_0^t e^{-(t-t')\partial_x^5} \psi_{\delta}(t') IF(u)(t') dt'.$$

By the definition of the restricted norm (5.1), we can choose  $\tilde{u} \in X^{1,\frac{1}{2}+}$ ,  $\tilde{u}|_{[0,\delta]} = u$  such that

$$\|I\tilde{u}\|_{X^{1,\frac{1}{2}+}} \sim \|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}}.$$
(5.25)

Duhamel's principle, (5.23), (5.24) and (4) in Lemma 5.1 give us

$$\begin{aligned} \|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}} &\lesssim \|Iu_0\|_{H^1} + \left\|\psi_{\delta}(t)IF(\tilde{u})(t)\right\|_{X^{1,-\frac{1}{2}+}} \\ &\lesssim \|Iu_0\|_{H^1} + \delta^{\varepsilon} \|IF(\tilde{u})(t)\|_{X^{1,-\frac{1}{2}++}}. \end{aligned}$$
(5.26)

We divide each  $\tilde{u}$  into a part supported on frequencies on  $|\xi| \leq N$  and a part supported on frequencies  $|\xi| \gg N$ , then from nonlinear estimates in Lemma 5.1 we immediately obtain that

$$\left\| IF(\tilde{u})(t) \right\|_{X^{1,-\frac{1}{2}++}} \lesssim \| I\tilde{u} \|_{X^{1,\frac{1}{2}+}}^3 + \| I\tilde{u} \|_{X^{1,\frac{1}{2}+}}^5.$$
(5.27)

Thus (5.25) and (5.26) yield that

$$\|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}} \lesssim \|Iu_0\|_{H^1} + \delta^{\varepsilon} \Big( \|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}}^3 + \|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}}^5 \Big).$$
(5.28)

It is easy to see that

$$\|Iu_0\|_{H^1} \lesssim \left(E(Iu_0)\right)^{1/2} + \|u_0\|_{L^2} \le 1 + \|u_0\|_{L^2}.$$
(5.29)

Since  $Q(\delta) := \|Iu\|_{X^{1,\frac{1}{2}+}_{\delta}}$  is continuous in the variable  $\delta$ , a continuous argument implies the conclusion from (5.28) and (5.29).

In the end, we turn to prove Proposition 5.1.

**Proof of Proposition 5.1.** By simple calculation, we can change all nonlinear terms in (1.2) into the divergence form, that is to say (1.2) becomes the following form:

$$\partial_t u + \partial_x^5 u + 6\partial_x (u^5) - 5\partial_x \left( u \partial_x^2 (u^2) \right) = 0.$$
(5.30)

The energy in (1.3) is known to be conserved by differentiating in time, using the equation (5.30) and integrating by parts, we have that

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t}E(u) &= \int_{\mathbb{R}} (-2\partial_x^2 u + 4u^3) \cdot \partial_t u \, \mathrm{d}x \\ &= \int_{\mathbb{R}} (-2\partial_x^2 u + 4u^3) \Big( -\partial_x^5 u - 6\partial_x (u^5) + 5\partial_x \Big( u\partial_x^2 (u^2) \Big) \Big) \, \mathrm{d}x \\ &= 2\int_{\mathbb{R}} \partial_x^2 u \partial_x^5 u \, \mathrm{d}x - 24\int_{\mathbb{R}} u^3 \partial_x (u^5) \, \mathrm{d}x \\ &- 2\int_{\mathbb{R}} 2u^3 \partial_x^5 u + 5\partial_x^2 u \partial_x \Big( u\partial_x^2 (u^2) \Big) \, \mathrm{d}x + 4\int_{\mathbb{R}} 3\partial_x^2 u \partial_x (u^5) + 5u^3 \partial_x \Big( u\partial_x^2 (u^2) \Big) \, \mathrm{d}x \\ &:= T_1 + T_2 + T_3 + T_4 = 0, \end{aligned}$$
(5.31)

In fact, we can get  $T_1 = T_2 = T_3 = T_4 = 0$  only by integrating by parts, which will be used in the following discussion. We apply I to (5.30), then

$$\partial_t I u = -\partial_x^5 I u - 6\partial_x I(u^5) + 5\partial_x I\left(u\partial_x^2(u^2)\right).$$
(5.32)

Following the same strategy, we estimate the growth of E(Iu)(t). By the equation (5.32), we know

that

$$\frac{d}{dt}E(Iu)(t) = \int_{\mathbb{R}} (-2\partial_x^2 Iu + 4(Iu)^3) \cdot \partial_t Iu \, dx$$

$$= \int_{\mathbb{R}} (-2\partial_x^2 Iu + 4(Iu)^3) \left( -\partial_x^5 Iu - 6\partial_x I(u^5) + 5\partial_x I(u\partial_x^2(u^2)) \right) \, dx$$

$$= 2\int_{\mathbb{R}} \partial_x^2 Iu \cdot \partial_x^5 Iu \, dx + 72\int_{\mathbb{R}} \partial_x Iu \cdot (Iu)^2 \cdot I(u^5) \, dx$$

$$+ 4\int_{\mathbb{R}} 3\partial_x^2 Iu \cdot \partial_x I(u^5) + 5(Iu)^3 \cdot \partial_x I(u\partial_x^2(u^2)) \, dx$$

$$- 2\int_{\mathbb{R}} 2(Iu)^3 \cdot \partial_x^5 Iu + 5\partial_x^2 Iu \cdot \partial_x I(u\partial_x^2(u^2)) \, dx$$

$$:= T_{I1} + T_{I2} + T_{I3} + T_{I4}.$$
(5.33)

It is easy to see that

$$T_{I1} = -2 \int_{\mathbb{R}} \partial_x^3 I u \cdot \partial_x^4 I u \, dx = -\int_{\mathbb{R}} \partial_x \left( (\partial_x^3 I u)^2 \right) \, dx = 0.$$

We recall the k-Parseval formula:

$$\int_{\mathbb{R}} f_1(x) f_2(x) \cdots f_k(x) \, \mathrm{d}x = \int_{\xi_1 + \xi_2 + \dots + \xi_k = 0} \widehat{f}_1(\xi_1) \widehat{f}_2(\xi_2) \cdots \widehat{f}_k(\xi_k), \tag{5.34}$$

For simplicity, denote  $\Gamma_k = \{(\xi_1, \xi_2, \dots, \xi_k) \in \mathbb{R}^k, \xi_1 + \xi_2 + \dots + \xi_k = 0\}$ . Therefore, k-Parseval formula combined with integrating by parts, yields that

$$T_{I2} = 72i \int_{\Gamma_8} \xi_1 \cdot \frac{m(\xi_4 + \xi_5 + \xi_6 + \xi_7 + \xi_8)}{m(\xi_4)m(\xi_5)m(\xi_6)m(\xi_7)m(\xi_8)} \widehat{Iu}(\xi_1)\widehat{Iu}(\xi_2) \cdots \widehat{Iu}(\xi_8);$$
(5.35)

$$T_{I3} = 12i \int_{\Gamma_6} \left( \xi_1^3 \frac{m(\xi_2 + \xi_3 + \dots + \xi_6)}{m(\xi_2)m(\xi_3)\cdots m(\xi_6)} + 5\xi_1(\xi_2 + \xi_3)^2 \cdot \frac{m(\xi_2 + \xi_3 + \xi_4)}{m(\xi_2)m(\xi_3)m(\xi_4)} \right) \\ \widehat{Iu}(\xi_1)\widehat{Iu}(\xi_2)\cdots \widehat{Iu}(\xi_6);$$
(5.36)

$$T_{I4} = -2i \int_{\Gamma_4} \left( 2\xi_1^5 - 5\xi_1^3 (\xi_2 + \xi_3)^2 \cdot \frac{m(\xi_2 + \xi_3 + \xi_4)}{m(\xi_2)m(\xi_3)m(\xi_4)} \right) \widehat{Iu}(\xi_1) \widehat{Iu}(\xi_2) \cdots \widehat{Iu}(\xi_4).$$
(5.37)

Integrating in time for (5.33), it suffices to control

$$E(Iu(\delta)) - E(Iu(0)) = \int_0^{\delta} T_{I2} + T_{I3} + T_{I4}.$$
(5.38)

The term  $T_{I2}$  is the easiest one to estimate, since there are fewer derivatives and more *u*'s. On the contrary, the term  $T_{I4}$  is the worst one. We will consider every term in the following three propositions. Have these propositions in hand, we can complete the proof by utilizing Proposition 5.2.

It remains to control every term in (5.38). Before that, we give some observations. The same argument as  $T_2 = T_3 = T_4 = 0$ , using k-Parseval formula and integrating by parts, show that

$$\int_{\Gamma_8} \xi_1 \widehat{Iu}(\xi_1) \widehat{Iu}(\xi_2) \cdots \widehat{Iu}(\xi_8) = 0; \qquad (5.39)$$

$$\int_{\Gamma_6} \left( \xi_1^3 + 5\xi_1 (\xi_2 + \xi_3)^2 \right) \widehat{Iu}(\xi_1) \widehat{Iu}(\xi_2) \cdots \widehat{Iu}(\xi_6) = 0;$$
(5.40)

$$\int_{\Gamma_4} \left( 2\xi_1^5 - 5\xi_1^3 (\xi_2 + \xi_3)^2 \right) \widehat{Iu}(\xi_1) \widehat{Iu}(\xi_2) \cdots \widehat{Iu}(\xi_4) = 0.$$
(5.41)

**Proposition 5.3** Let  $3/4 \leq s < 1$  and  $T_{I2}$  be as in (5.35), then

$$\left| \int_{0}^{\delta} T_{I2} \right| \lesssim N^{-5} \| Iu \|_{X_{\delta}^{1, \frac{1}{2} +}}^{8}.$$
(5.42)

**Proof.** We divide Iu into a sum of dyadic parts  $P_kIu$ , whose frequency is supported on  $\{\xi : |\xi| \sim 2^k\}$ ,  $k = 0, 1, 2, \cdots$ . It suffices to show that

$$\sum_{N_1,\dots,N_8} \left| \int_0^\delta \int_{\Gamma_8} \xi_1 \cdot \frac{m(\xi_4 + \dots + \xi_8)}{m(\xi_4) \cdots m(\xi_8)} \widehat{u_1}(\xi_1) \cdots \widehat{u_8}(\xi_8) \right| \lesssim N^{-5} \prod_{j=1}^8 \|u_j\|_{X_{\delta}^{1,\frac{1}{2}+}}, \tag{5.43}$$

for any function  $u_j$ ,  $j = 1, 2, \dots, 8$  with the frequency supported on  $|\xi_j| \sim 2^{k_j} \equiv N_j$ ,  $k_j \in \{0, 1, \dots\}$ . We may assume all  $\hat{u_j}$ 's are non-negative. In the following discussion, we will pull the symbol out of the integral, then reverse the k-Parseval formula and use Hölder's inequality to estimate the remaining integrals.

Note that the derivative is located in  $u_1$ , and by the symmetry of the above multiplier in  $\xi_4, \xi_5, \dots, \xi_8$ , we may assume that

$$N_1 \ge N_2 \ge N_3, \qquad N_4 \ge N_5 \ge \dots \ge N_8.$$
 (5.44)

Let  $N_{max}$  and  $N_{sec}$  denote the maximum and the second maximum of the numbers  $N_1, N_2, \dots, N_8$ . From the constraint condition  $\Gamma_8 = \left\{ \sum_{j=1}^8 \xi_j = 0 \right\}$  in the above integration, we know that

$$N_{max} \sim N_{sec}$$
.

Since  $m(\xi) = 1$  for  $\xi \leq N$ , (5.35) and (5.39) shows that  $T_{I2}$  vanishes when  $|\xi_4| \leq N/5$ . Thus, we may assume that

$$N_4 \gtrsim N.$$

According to  $N_{max}$  and  $N_{sec}$ , we split the proof into three cases:  $\{N_{max}, N_{sec}\} = \{N_1, N_4\}, \{N_{max}, N_{sec}\} = \{N_4, N_5\}, \{N_{max}, N_{sec}\} = \{N_1, N_2\}$ . For the sake of brevity, we only consider the first case since the other two cases can be obtained by the same techniques.

If  $\{N_{max}, N_{sec}\} = \{N_1, N_4\}$ , we have here  $N_1 \sim N_4 \gtrsim N$ . Using local smoothing effect estimates (5.3) to  $u_1, u_4$ , maximal function estimates (5.4) to  $u_5$ - $u_8$ , it follows that

LHS of 
$$(5.43)$$

$$\lesssim \sum_{N_1 \sim N_4 \gtrsim N \atop N_1 \cdots N_8} \frac{N_1 N_4^{1-s}}{N^{1-s} m(N_5) \cdots m(N_8)} \|u_1\|_{L_x^{\infty} L_t^2} \|u_4\|_{L_x^{\infty} L_t^2} \prod_{i=2}^3 \|u_i\|_{L_x^{\infty} L_t^{\infty}} \prod_{j=5}^8 \|u_j\|_{L_x^4 L_t^{\infty}} \\ \lesssim \sum_{N_1 \sim N_4 \gtrsim N \atop N_1 \cdots N_8} \frac{N_1 N_4^{1-s}}{N^{1-s} m(N_5) \cdots m(N_8)} \|u_1\|_{X_{\delta}^{-2, \frac{1}{2}+}} \|u_4\|_{X_{\delta}^{-2, \frac{1}{2}+}} \prod_{i=2}^3 \|u_i\|_{X_{\delta}^{\frac{1}{2}+, \frac{1}{2}+}} \prod_{j=5}^8 \|u_j\|_{X_{\delta}^{\frac{1}{4}, \frac{1}{2}+}}$$

$$\lesssim N^{s-1} \sum_{N_1 \sim N_4 \gtrsim N \atop N_1, \dots, N_8} N_1^{-2} N_4^{-2-s} (N_2 N_3)^{-\frac{1}{2}+} (N_5 N_6 N_7 N_8)^{-\frac{1}{2}} \prod_{j=1}^8 \|u_j\|_{X_{\delta}^{1,\frac{1}{2}+}} \lesssim N^{-5} \prod_{j=1}^8 \|u_j\|_{X_{\delta}^{1,\frac{1}{2}+}},$$

where we used Sobolev embedding and Strichart estimate to get  $\|u_i\|_{L_t^{\infty}L_x^{\infty}} \lesssim \|\langle \nabla \rangle^{\frac{1}{2}+} u_i\|_{L_t^{\infty}L_x^2} \lesssim \|u_i\|_{X_{\delta}^{\frac{1}{2}+,\frac{1}{2}+}}$ . Since  $3/4 \leq s < 1$ , we know that  $m(N_j)N_j^{1/4} \geq 1$  for  $j = 5, \cdots, 8$ .

**Proposition 5.4** Let  $3/4 \le s < 1$  and  $T_{I3}$  be as in (5.36), then

$$\left| \int_{0}^{\delta} T_{I3} \right| \lesssim N^{-3} \| Iu \|_{X_{\delta}^{1, \frac{1}{2} +}}^{6}.$$
(5.45)

**Proof.** At first, for getting some vanishing properties, we insert the zero term (5.40) into  $T_{I3}$ . Then as we discussed in the above proposition, it suffices to control

$$I_1 := \sum_{N_1, \cdots, N_6} \left| \int_0^\delta \int_{\Gamma_6} \xi_1^3 \left( \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \cdots m(\xi_6)} - 1 \right) \widehat{u_1}(\xi_1) \cdots \widehat{u_6}(\xi_6) \right|$$

and

$$I_2 := \sum_{N_1, \cdots, N_6} \left| \int_0^\delta \int_{\Gamma_6} 5\xi_1 (\xi_2 + \xi_3)^2 \cdot \left( \frac{m(\xi_2 + \xi_3 + \xi_4)}{m(\xi_2)m(\xi_3)m(\xi_4)} - 1 \right) \widehat{u_1}(\xi_1) \cdots \widehat{u_6}(\xi_6) \right|,$$

for any function  $u_j$ ,  $j = 1, 2, \dots, 6$  with the frequency supported on  $|\xi_j| \sim N_j$ .

For term  $I_1$ , by the symmetry of the multiplier in  $\xi_2, \xi_3, \dots, \xi_6$ , we can assume that

$$N_2 \ge N_3 \ge \dots \ge N_6, \qquad N_2 \gtrsim N, \tag{5.46}$$

where we use that if  $N_2 \leq N/10$ ,  $I_1$  vanishes. Because of  $N_{max} \sim N_{sec}$ , we divide the proof into two cases:  $N_1 \sim N_2 \gtrsim N$  and  $N_2 \sim N_3 \gtrsim N$ ,  $N_3 \geq N_1$ .

If  $N_1 \sim N_2 \gtrsim N$ , we take local smoothing effect estimates (5.3) to  $u_1$ ,  $u_2$ , maximal function estimates (5.4) to  $u_3$ - $u_6$ , and then get that

$$I_{1} \lesssim \sum_{\substack{N_{1} \sim N_{2} \gtrsim N \\ N_{1}, \cdots, N_{6} \\ N_{1}, \cdots, N_{6} \\ N_{1}, \cdots, N_{6} \\ N_{1}, \cdots, N_{6} \\ N_{1} \sim N_{2} \gtrsim N \\ N_{1} \sim N_{2} \gtrsim N \\ \sum_{\substack{N_{1} \sim N_{2} \gtrsim N \\ N_{1}, \cdots, N_{6} \\ N_{1}, \cdots, N_{6} \\ N_{2} \sim N_{2} \\ N_{1} \sim N_{2} \gtrsim N \\ N_{2}^{-3} (N_{3}N_{4}N_{5}N_{6})^{-\frac{1}{2}} \prod_{j=1}^{6} \|u_{j}\|_{X_{\delta}^{-\frac{1}{2}+}} \|u_{2}\|_{X_{\delta}^{-2,\frac{1}{2}+}} \prod_{j=3}^{6} \|u_{j}\|_{X_{\delta}^{\frac{1}{4},\frac{1}{2}+}}$$

$$(5.47)$$

If  $N_2 \sim N_3 \gtrsim N$ ,  $N_3 \geq N_1$ , we can get similarly that

$$I_1 \lesssim \sum_{N_2 \sim N_3 \gtrsim N \atop N_1, \dots, N_6} \frac{N_1^3 m(N_1)}{m(N_2) \cdots m(N_6)} \|u_2\|_{L_x^{\infty} L_t^2} \|u_3\|_{L_x^{\infty} L_t^2} \|u_1\|_{L_x^4 L_t^{\infty}} \prod_{j=4}^6 \|u_j\|_{L_x^4 L_t^{\infty}}$$

$$\lesssim N^{1-s} \sum_{\substack{N_2 \sim N_3 \gtrsim N \\ N_1, \cdots, N_6}} \frac{N_1^{s+2}}{m(N_2) \cdots m(N_6)} \|u_1\|_{X_{\delta}^{\frac{1}{4}, \frac{1}{2}+}} \|u_2\|_{X_{\delta}^{-2, \frac{1}{2}+}} \|u_3\|_{X_{\delta}^{-2, \frac{1}{2}+}} \prod_{j=4}^6 \|u_j\|_{X_{\delta}^{\frac{1}{4}, \frac{1}{2}+}} \\ \lesssim N^{1-s} \sum_{\substack{N_2 \sim N_3 \gtrsim N \\ N_1, \cdots, N_6}} N_1^{s+\frac{5}{4}} N_2^{-\frac{11}{4}} N_3^{-\frac{11}{4}} (N_4 N_5 N_6)^{-\frac{1}{2}} \prod_{j=1}^6 \|u_j\|_{X_{\delta}^{1, \frac{1}{2}+}} \\ \lesssim N^{-3} \prod_{j=1}^6 \|u_j\|_{X_{\delta}^{1, \frac{1}{2}+}}.$$

$$(5.48)$$

For term  $I_2$ , the same argument as before, we may assume that

$$N_1 \ge N_5 \ge N_6, \qquad N_2 \ge N_3 \ge N_4, \qquad N_2 \gtrsim N.$$
 (5.49)

If  $N_1 \sim N_2 \gtrsim N$ , the symbol is controlled by  $N_1^3 m(N_1)/m(N_2)m(N_3)m(N_4)$ , which is smaller than  $N_1^3 m(N_1)/m(N_2)\cdots m(N_6)$ , therefore, we can get the same bound as (5.47). If  $N_2 \sim N_3 \gtrsim N$ , using the same techniques as (5.48) yields the estimates. If  $N_1 \sim N_5 \gtrsim N$ , this case is better, thus we can obtain the conclusion as before.

**Proposition 5.5** Let  $3/4 \le s < 1$  and  $T_{I4}$  be as in (5.51), then

$$\left| \int_{0}^{\delta} T_{I4} \right| \lesssim N^{-\frac{1}{2}} \| Iu \|_{X_{\delta}^{1,\frac{1}{2}+}}^{4}.$$
(5.50)

**Proof.** As the above proposition, inserting the zero term (5.41) into  $T_{I4}$ , it suffices to control

$$\sum_{N_1,\dots,N_4} \left| \int_0^\delta \int_{\Gamma_4} \xi_1^3 (\xi_2 + \xi_3)^2 \cdot \left( \frac{m(\xi_2 + \xi_3 + \xi_4)}{m(\xi_2)m(\xi_3)m(\xi_4)} - 1 \right) \widehat{u_1}(\xi_1) \cdots \widehat{u_4}(\xi_4) \right|, \tag{5.51}$$

for any function  $u_j$ , j = 1, 2, 3, 4 with the frequency supported on  $|\xi_j| \sim N_j$ . A glance at the symbol, we may assume that

$$N_2 \ge N_3 \ge N_4, \qquad N_2 \gtrsim N, \tag{5.52}$$

since that if  $N_2 \leq N/8$ , the symbol vanishes. The other orders of  $N_2$ ,  $N_3$  and  $N_4$  are similar and much easier. Then we control the above summation by considering the following three cases:

- 1. the non-resonant case, where  $N_2 \gg N_3 \ge N_4$ ;
- 2. the semi-resonant case, where  $N_2 \sim N_3 \gg N_4$ ;
- 3. the resonant case, where  $N_2 \sim N_3 \sim N_4$ .

1. In the non-resonant case, from the constraint condition  $\Gamma_4 = \left\{ \sum_{j=1}^4 \xi_j = 0 \right\}$ , it follows immediately that  $N_1 \sim N_2 \gg N_3 \geq N_4$ . Utilizing bilinear estimates (5.5) can efficiently control (5.51) by

$$\sum_{\substack{N_1 \sim N_2 \gtrsim N \\ N_1 \sim N_2 \gg N_3 \geq N_4}} \frac{N_1^3 N_2^2 m(N_1)}{m(N_2) m(N_3) m(N_4)} \|u_1 u_3\|_{L^2_{x,t}} \|u_2 u_4\|_{L^2_{x,t}}$$

$$\lesssim \sum_{\substack{N_1 \sim N_2 \gtrsim N \\ N_1 \sim N_2 \gg N_3 \geq N_4}} \frac{N_1}{m(N_3)m(N_4)} \prod_{j=1}^4 \|u_j\|_{X_{\delta}^{0,\frac{1}{2}+}}$$
  
$$\lesssim \sum_{\substack{N_1 \sim N_2 \gtrsim N \\ N_1 \sim N_2 \gg N_3 \geq N_4}} N_2^{-1} (N_3 N_4)^{-\frac{3}{4}} \prod_{j=1}^4 \|u_j\|_{X_{\delta}^{1,\frac{1}{2}+}} \lesssim N^{-1} \prod_{j=1}^4 \|u_j\|_{X_{\delta}^{1,\frac{1}{2}+}}.$$
(5.53)

2. In the semi-resonant case, it is easy to see that  $N_2 \sim N_3 \gg N_4 \sim N_1$ . Bilinear estimates (5.5) imply that (5.51) is controlled by

$$\sum_{\substack{N_{2}\sim N_{3}\gtrsim N\\N_{2}\sim N_{3}\gg N_{4}\sim N_{1}}} \frac{N_{1}^{3}N_{2}^{2}m(N_{1})}{m(N_{2})m(N_{3})m(N_{4})} \|u_{1}u_{3}\|_{L_{x,t}^{2}} \|u_{2}u_{4}\|_{L_{x,t}^{2}}$$

$$\lesssim \sum_{\substack{N_{2}\sim N_{3}\gtrsim N\\N_{2}\sim N_{3}\gg N_{4}\sim N_{1}}} \frac{N_{1}^{3}N_{3}^{-2}}{m(N_{2})m(N_{3})} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{0,\frac{1}{2}+}}$$

$$\lesssim \sum_{\substack{N_{2}\sim N_{3}\gtrsim N\\N_{2}\sim N_{3}\gg N_{4}\sim N_{1}}} N_{1}N_{2}^{-\frac{3}{4}}N_{3}^{-\frac{11}{4}} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{1,\frac{1}{2}+}} \lesssim N^{-\frac{5}{2}} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{1,\frac{1}{2}+}}.$$
(5.54)

3. In the resonant case, we have  $N_1 \sim N_2 \sim N_3 \sim N_4$ . Strichartz estimates (5.2) yield that

$$(5.51) \lesssim \delta^{\frac{1}{2}} \sum_{N_{1} \sim N_{2} \sim N_{3} \sim N_{4} \gtrsim N} \frac{N_{1}^{3} N_{2}^{2} m(N_{1})}{m(N_{2}) m(N_{3}) m(N_{4})} \prod_{j=1}^{4} \|u_{j}\|_{L_{t}^{8} L_{x}^{4}}$$

$$\lesssim \delta^{\frac{1}{2}} \sum_{N_{1} \sim N_{2} \sim N_{3} \sim N_{4} \gtrsim N} \frac{N_{1}^{3} N_{2}^{2}}{m(N_{3}) m(N_{4})} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{-\frac{3}{8}, \frac{1}{2}+}}$$

$$\lesssim \delta^{\frac{1}{2}} N^{2s-2} \sum_{N_{1} \gtrsim N} N_{1}^{\frac{3}{2}-2s} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{\frac{1}{2}+}} \lesssim \delta^{\frac{1}{2}} N^{-\frac{1}{2}} \prod_{j=1}^{4} \|u_{j}\|_{X_{\delta}^{1, \frac{1}{2}+}}.$$
(5.55)

We now complete the proof.

### 5.4 Global approximation in low regularity spaces

In this subsection, we shall extend the long-time asymptotic behavior to the rough data. At the beginning, we show that the Beals-Coifman solution is equal to the strong solution given by (1.29).

**Lemma 5.4** Let  $u_0 \in H^{4,1}(\mathbb{R})$ , the Beals-Coifman solution and the strong solution are the same (up to a measure zero set)

$$u = \frac{1}{\pi} \left( \int \mu (w^+ + w^-) \right)_{12} = e^{-t\partial_x^5} u_0 - \int_0^t e^{-(t-t')\partial_x^5} F(u)(t') \, dt'$$

in [-T, T] where T is given as in Theorem 5.1.

**Proof.** For  $u_0 \in H^{4,1}(\mathbb{R})$ , there exists a Cauchy sequence  $u_{0,k} \in \mathscr{S}(\mathbb{R})$ , such that

$$\lim_{k \to \infty} \|u_{0,k} - u_0\|_{H^{4,1}} = 0, \quad \text{and} \quad \|u_{0,k}\|_{H^4} \le \|u_{0,k}\|_{H^{4,1}} \le C.$$

On the one hand, from Theorem 5.1, there exist a strong solution  $u_k$  with initial data  $u_{0,k}$  such that for b > 1/2,

$$||u_k - u_l||_{X^{4,b}} \lesssim ||u_{0,k} - u_{0,l}||_{H^4} \to 0 \text{ as } k, l \to \infty,$$

Therefore, there exists  $u_{\infty}$  such that

$$\sup_{t \in [-T,T]} \|u_k - u_\infty\|_{L^{\infty}} \lesssim \|u_k - u_\infty\|_{X^{4,b}} \to 0, \quad \text{as} \quad k \to \infty,$$
(5.56)

On the other hand, from the inverse scattering transform, we have the Beals-Coifman solutions  $\tilde{u}_k$  with initial data  $u_{0,k}$ . By utilizing the bijectivity of the transformation and bi-Lipschitz continuity, we know reflection coefficients satisfy

$$\|r_k - r_l\|_{H^{1,4}} \lesssim \|u_{0,k} - u_{0,l}\|_{H^{4,1}} \to 0 \quad as \quad k, l \to \infty.$$

By the resolvent estimates,

$$\|\tilde{u}_k - \tilde{u}_l\|_{L^{\infty}} \lesssim \|r_k - r_l\|_{H^1} \to 0 \quad as \quad k, l \to \infty.$$

Therefore, there exists  $\tilde{u}_{\infty}$  such that

$$\tilde{u}_{\infty} = \lim_{k \to \infty} \tilde{u}_k. \tag{5.57}$$

Since  $u_{0,k} \in \mathscr{S}(\mathbb{R})$ ,  $u_k$  and  $\tilde{u}_k$  are also Schwartz functions, then  $u_k = \tilde{u}_k$ . Therefore, we can get the conclusion from (5.56) and (5.57).

In the end, we divide the proof of Theorem 1.3 into the following two theorems. We first consider the long-time behavior in the Sobolev space  $H^{1,1}(\mathbb{R})$ . It is well known that  $u(\cdot, t)$  is uniformly bounded in  $H^1(\mathbb{R})$ . Indeed, from the mass conservation and energy conservation, we know that

$$||u||_{H^1(\mathbb{R})} = ||u||_{L^2} + ||u||_{\dot{H}^1} \le ||u_0||_{L^2} + ||\partial_x u_0||_{L^2} + ||u_0||_{L^4}^2.$$

From Gagliardo-Nirenberg's inequality and Young's inequality,

$$\|u_0\|_{L^4}^2 \lesssim \|\partial_x u_0\|_{L^2}^{1/2} \|u_0\|_{L^2}^{3/2} \lesssim \frac{1}{\epsilon} \|\partial_x u_0\|_{L^2} + \epsilon \|u_0\|_{L^2}^3.$$

We may choose  $\epsilon$  satisfying  $\epsilon ||u_0||_{L^2}^2 \leq 1$ , and get that

$$\|u(\cdot,t)\|_{H^1} \lesssim \|u_0\|_{H^1}. \tag{5.58}$$

**Theorem 5.2** Let  $u_0 \in H^{1,1}(\mathbb{R})$ , the strong solution given by the integral form (1.29) has the same asymptotic behavior as in Theorem 1.1.

**Proof.** For  $u_0 \in H^{1,1}(\mathbb{R})$ , there exists a sequence  $u_{0,k} \in H^{4,1}(\mathbb{R})$ , such that

$$\lim_{k \to \infty} \|u_{0,k} - u_0\|_{H^{1,1}} = 0, \quad and \quad \|u_{0,k}\|_{H^1} \le \|u_{0,k}\|_{H^{1,1}} \le C.$$

Following the same strategy as in the proof of Lemma 5.4, we know that there exist a pointwise limit  $u_{\infty}$  of the strong solutions  $u_k$ , and a pointwise limit  $\tilde{u}_{\infty}$  of Beals-Coifman solutions  $\tilde{u}_k$ , with initial data  $u_{0,k}$ . Moreover, one has  $u_{\infty} = \tilde{u}_{\infty}$  pointwise for  $t \in [-T, T]$ . Since  $u(\cdot, t)$  is uniformly bounded in  $H^1(\mathbb{R})$  as (5.58), one can repeat the above argument many times to extend the time interval to whole line  $\mathbb{R}$ . Because  $\tilde{u}_{\infty}$  has asymptotic behavior in Theorem 1.1,  $u_{\infty}$  also does.  $\Box$ 

**Theorem 5.3** Let  $u_0 \in H^{s,1}(\mathbb{R})$ , 19/22 < s < 1, the strong solution given by the integral form (1.29) has the same asymptotic behavior as in Theorem 1.1.

**Proof.** For  $u_0 \in H^{s,1}(\mathbb{R})$ , 19/22 < s < 1, there exists a sequence  $u_{0,k} \in H^{4,1}(\mathbb{R})$ , such that

$$\lim_{k \to \infty} \|u_{0,k} - u_0\|_{H^{s,1}} = 0, \quad and \quad \|u_{0,k}\|_{H^s} \le \|u_{0,k}\|_{H^{s,1}} \le C$$

Denote  $u_{\infty}$  and  $\tilde{u}_{\infty}$  the pointwise limit of the strong solutions  $u_k$  and Beals-Coifman solutions  $\tilde{u}_k$  with initial data  $u_{0,k}$ , respectively. As Lemma 5.4, we know that  $u_{\infty} = \tilde{u}_{\infty}$  pointwise for  $t \in [-T, T]$ .

From the global well-posedness theory Theorem 1.2,  $u_{\infty}$  exists in  $H^s$  globally. One can also construct  $\tilde{u}_{\infty}$  globally. Suppose that  $u_{\infty} = \tilde{u}_{\infty}$  does not hold for all t. We may assume

$$T_* = \inf \{ t \ge 0; u_{\infty}(t) \neq \tilde{u}_{\infty}(t) \}.$$
(5.59)

From (5.14), we know that

$$\|u(\cdot, T_*)\|_{H^s} \le C_1 T_*^M + C_2, \tag{5.60}$$

then as in the proof of Theorem 5.1, there exists  $t_0 > 0$  such that

$$\sup_{t \in [0,T_*+t_0]} \|u_k - u_\infty\|_{L^\infty} \lesssim \|u_k - u_\infty\|_{X^{s,b}} \to 0, \quad as \quad k \to \infty,$$

Combining

$$\sup_{t \in [0,T_*+t_0]} \|\tilde{u}_k - \tilde{u}_\infty\|_{L^{\infty}} \to 0, \quad as \quad k \to \infty, \quad and \quad u_k = \tilde{u}_k,$$

we can conclude that  $u_{\infty}(t) = \tilde{u}_{\infty}(t)$  pointwise for  $t \in [0, T_* + t_0]$ , this is a contradiction with (5.59). Hence we can obtain that  $u_{\infty}(t) = \tilde{u}_{\infty}(t)$  pointwise for all  $t \ge 0$ . Since  $\tilde{u}_{\infty}$  has asymptotic behavior in Theorem 1.1,  $u_{\infty}$  also does.

#### Acknowledgments.

N. Liu is supported by the China Postdoctoral Science Foundation under Grant Nos. 2019TQ0041, 2019M660553; M. Chen is supported in part by the China Postdoctoral Science Foundation under Grant No. 2019M650019.

### A Fourth order Painlevé II RH problem

Let P denote the contour  $P = P_1 \cup P_2$  oriented as in Fig. 10, where

$$P_1 = \{ le^{\frac{\pi i}{6}} | l \ge 0 \} \cup \{ le^{\frac{5\pi i}{6}} | l \ge 0 \}, \quad P_2 = \{ le^{-\frac{\pi i}{6}} | l \ge 0 \} \cup \{ le^{-\frac{5\pi i}{6}} | l \ge 0 \}.$$



Fig. 10: The oriented contour P.

Fourth order Painlevé II RH problem A.1. Let  $s \in \mathbb{R}$  be a real number. Find an analytic function  $M^P(y; z)$  in  $\mathbb{C} \setminus P$  parametrized by  $y \in \mathbb{R}, s \in \mathbb{R}$  such that:

1. For  $z \in P$ , the continuous boundary values  $M^P_{\pm}(y; z)$  satisfy

$$M^{P}_{+}(y;z) = M^{P}_{-}(y;z)v^{P}(s,y;z), \quad z \in P,$$
(A.1)

where the jump matrix  $v^P(s, y; z)$  is defined by

$$v^{P}(s,y;z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ se^{2i(\frac{4}{5}z^{5}-yz)} & 1 \end{pmatrix}, & z \in P_{1}, \\ \begin{pmatrix} 1 & -se^{-2i(\frac{4}{5}z^{5}-yz)} \\ 0 & 1 \end{pmatrix}, & z \in P_{2}. \end{cases}$$
(A.2)

2. 
$$M^P(y;z) = I + O(\frac{1}{z})$$
, as  $z \to \infty$ .

**Lemma A.1** The RH problem A.1 has a unique solution  $M^P(y; z)$  for each  $y \in \mathbb{R}$ . Moreover, there exists smooth functions  $\{M_j^P(y)\}^4$  of  $y \in \mathbb{R}$  with decay as  $y \to -\infty$  such that

$$M^{P}(y;z) = I + \sum_{j=1}^{4} \frac{M_{j}^{P}(y)}{z^{j}} + O(z^{-5}), \quad z \to \infty,$$
(A.3)

uniformly for y in compact subsets of  $\mathbb{R}$  and for  $\arg z \in [0, 2\pi]$ . The leading coefficient  $M_1^Y$  is given by

$$M_1^P(y) = i \begin{pmatrix} -2 \int_{-\infty}^y u_p^2(y') \, dy' & u_p(y) \\ -u_p(y) & 2 \int_{-\infty}^y u_p^2(y') \, dy' \end{pmatrix}, \tag{A.4}$$

where the real-valued function  $u_p(y)$  satisfies the following fourth order Painlevé II equation (see [35])

$$u_p^{\prime\prime\prime\prime}(y) - 40u_p^2(y)u_p^{\prime\prime}(y) - 40u_p(y)u_p^{\prime 2}(y) + 96u_p^5(y) - 4yu_p(y) = 0.$$
(A.5)



Fig. 11: The oriented contour Z.

# B Model RH problem in Region III

Given a number  $z_0 \ge 0$ , let Z denote the contour  $Z = Z_1 \cup Z_1 \cup Z_3$ , where the line segments

$$Z_{1} = \{z_{0} + le^{\frac{\pi i}{6}} | l \ge 0\} \cup \{-z_{0} + le^{\frac{5\pi i}{6}} | l \ge 0\},$$

$$Z_{2} = \{z_{0} + le^{-\frac{\pi i}{6}} | l \ge 0\} \cup \{-z_{0} + le^{-\frac{5\pi i}{6}} | l \ge 0\},$$

$$Z_{3} = \{l | -z_{0} \le l \le z_{0}\}$$
(B.1)

are oriented as in Fig. 11.

Model RH problem B.1. Find an analytic function  $M^Z(y; z, z_0)$  in  $\mathbb{C} \setminus Z$  parametrized by  $y > 0, s \in \mathbb{R}, z_0 \ge 0$  such that:

1. The continuous boundary values  $M^Z_{\pm}(y;z,z_0)$  satisfy the jump condition

$$M_{+}^{Z}(y;z,z_{0}) = M_{-}^{Z}(y;z,z_{0})v^{Z}(s,y;z,z_{0}), \quad z \in \mathbb{Z},$$
(B.2)

where the jump matrix  $v^{Z}(s, y, z; z_{0})$  is defined by

$$v^{Z}(s,y;z,z_{0}) = \begin{cases} \begin{pmatrix} 1 & 0 \\ se^{2i(\frac{4}{5}z^{5}-yz)} & 1 \end{pmatrix}, & z \in Z_{1}, \\ \begin{pmatrix} 1 & -se^{-2i(\frac{4}{5}z^{5}-yz)} \\ 0 & 1 \end{pmatrix}, & z \in Z_{2}, \\ \begin{pmatrix} 1 & -se^{-2i(\frac{4}{5}z^{5}-yz)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ se^{2i(\frac{4}{5}z^{5}-yz)} & 1 \end{pmatrix}, & z \in Z_{3}. \end{cases}$$
(B.3)

2. As  $z \to \infty$ ,  $M^Z(y; z, z_0) = I + O(\frac{1}{z})$ .

Lemma B.2 Define the parameter subset

$$\mathbb{P} = \{ (y, t, z_0) \in \mathbb{R}^3 | 0 < y < C_1, t \ge 1, \sqrt[4]{y} / \sqrt{2} \le z_0 \le C_2 \},$$
(B.4)

where  $C_1, C_2 > 0$  are constants. Then for  $(y, t, z_0) \in \mathbb{P}$ , the RH problem B.1 has a unique solution  $M^Z(s, y; z, z_0)$  which satisfies

$$M^{Z}(y;z,z_{0}) = I + \frac{i}{z} \begin{pmatrix} -2\int_{\infty}^{y} u_{p}^{2}(y') dy' & -u_{p}(y) \\ u_{p}(y) & 2\int_{\infty}^{y} u_{p}^{2}(y') dy' \end{pmatrix} + O\left(\frac{1}{z^{2}}\right), \quad z \to \infty,$$
(B.5)

where  $u_p(y)$  denotes the solution of the fourth order Painlevé II equation (A.5) and  $M^Z(y, z, z_0)$  is uniformly bounded for  $z \in \mathbb{C} \setminus Z$ . Furthermore,  $M^Z$  obeys the symmetries

$$M^{Z}(y; -z, z_{0}) = \overline{M^{Z}(y; \bar{z}, z_{0})} = \sigma_{1} M^{Z}(y; z, z_{0}) \sigma_{1}.$$
 (B.6)

# References

- M.J. Ablowitz and A.S. Fokas, Complex Variables: Introduction and Applications, 2nd edn. Cambridge University Press, Cambridge 2003.
- [2] L.K. Arruda and J. Lenells, Long-time asymptotics for the derivative nonlinear Schrödinger equation on the half-line, Nonlinearity, 30 (2017), 4141–4172.
- [3] R. Beals and R. Coifman, Scattering and inverse scattering for first order systems, Commun. Pure Appl. Math., 37 (1984), 39–90.
- [4] D. Bilman, L. Ling and P. Miller, Extreme superposition: rogue waves of infinite order and the Painlevé-III hierarchy, arXiv:1806.00545, 2018. To appear in Duke Math. J.
- [5] M. Borghese, R. Jenkins and K.D.T.-R. McLaughlin, Long time asymptotic behavior of the focusing nonlinear Schrödinger equation, Ann. I. H. Poincaré-AN, 35 (2018), 887–920.
- [6] A. Boutet de Monvel, A. Its and D. Shepelsky, Painlevé-type asymptotics for the Camassa-Holm equation, SIAM J. Math. Anal., 42 (2010), 1854–1873.
- [7] B. Bringmann, R. Killip and M. Visan, Global well-posedness for the fifth-order KdV equation in  $H^{-1}(\mathbb{R})$ , arXiv:1912.01536v1, 2019.
- [8] G. Chen and J. Liu, Long-time asymptotics of the modified KdV equation in weighted Sobolov spaces, arXiv:1903.03855, 2019.
- [9] G. Chen and J. Liu, Soliton resolution for the modified KdV equation, arXiv:1907.07115, 2019.
- W. Chen and Z. Guo, Global well-posedness and I method for the fifth order Korteweg-de Vries equation, J. Anal. Math., 114 (2011), 121–156.
- [11] Po-Jen Cheng, S. Venakides and X. Zhou, Long-time asymptotics for the pure radiation solution of the sine-Gordon equation, Commun. Partial Differential Equations, 24 (1999), 1195–1262.
- [12] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Almost conservation laws and global rough solutions to a Nonlinear Schrödinger equation, Math. Res. Lett., 9 (2002), 659–682.
- [13] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Sharp Global well-posedness for KdV and modified KdV on ℝ and T, J. Amer. Math. Soc., 16 (2003), 705–749.
- [14] S. Cuccagna and R. Jenkins, On the asymptotic stability of N-soliton solutions of the defocusing nonlinear Schrödinger equation, Commun. Math. Phys., 343 (2016), 921–969.
- S. Cui and S. Tao, Strichartz estimates for dispersive equations and solvability of the Kawahara equation, J. Math. Anal. Appl., 304 (2005), 683–702.
- [16] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Ann. Math., 137 (1993), 295–368.
- [17] P. Deift and X. Zhou, Long-time behavior of the non-focusing nonlinear Schrödinger equation-a case study, Lectures in Mathematical Sciences, University of Tokyo, Tokyo, 1995.
- [18] P. Deift and X. Zhou, Long-time asymptotics for solutions of the NLS equation with initial data in a weighted Sobolev space, Commun. Pure Appl. Math., 56 (2003), 1029–1077.

- [19] M. Dieng, K.D.T.-R. McLaughlin, Long-time asymptotics for the NLS equation via dbar methods, Preprint, arXiv:0805.2807, 2008.
- [20] X. Gao, Global well-posedness for the fifth-order mKdV equation, Acta Math. Sin. (Engl. Ser.), 34 (2018), 1015–1027.
- [21] P. Germain, F. Pusateri and F. Rousset, Asymptotic stability of solitons for mKdV, Adv. Math., 299 (2016), 272–330.
- [22] R. Grimshaw, E. Pelinovsky and O. Poloukhina, Higher-order Korteweg-de Vries models for internal solitary waves in a stratified shear flow with a free surface, Nonlinear Processes in Geophysics, 9 (2002), 221–235.
- [23] R. Grimshaw, E. Pelinovsky and T. Talipova, The modified Korteweg-de Vries equation in the theory of large-amplitude internal waves, Nonlinear Processes in Geophysics, 4 (1997), 237–250.
- [24] Z. Guo, C. Kwak and S. Kwon, Rough solutions of the fifth-order KdV equations, J. Funct. Anal., 265 (2013), 2791–2829.
- [25] Z. Guo, L. Peng and B. Wang, Decay estimates for a class of wave equations, J. Funct. Anal., 254 (2008), 1642–1660.
- [26] B. Harrop-Griffiths, Long time behavior of solutions to the mKdV, Commun. Partial Differential Equations, 41 (2016), 282–317.
- [27] N. Hayashi and P. Naumkin, Large time behavior of solutions for the modified Korteweg-de Vries equation, Int. Math. Res. Not. IMRN, 1999 (1999), 395–418.
- [28] L. Huang and J. Lenells, Asymptotics for the Sasa–Satsuma equation in terms of a modified Painlevé II transcendent, J. Differential Equations, https://doi.org/10.1016/j.jde.2019.11.062, (2019).
- [29] R. Jenkins, J. Liu, P. Perry and C. Sulem, Soliton resolution for the derivative nonlinear Schrödinger equation, Commun. Math. Phys., 363 (2018), 1003–1049.
- [30] Y. Jia and Z. Huo, Well-posedness for the fifth-order shallow water equations, J. Differential Equations, 246 (2009), 2448–2467.
- [31] C. Kenig, G. Ponce and L. Vega, On the hierarchy of the generalized KdV equations. In: N.M. Ercolani, I.R. Gabitov, C.D. Levermore, D. Serre (eds) Singular Limits of Dispersive Waves. NATO ASI Series (Series B: Physics), vol 320. Springer, Boston, MA, 1994.
- [32] C. Kenig, G. Ponce and L. Vega, Well-posedness of the initial value problem for the Korteweg-de Vries equation, J. Amer. Math. Soc., 4 (1991), 323–347.
- [33] C. Kenig, G. Ponce and L. Vega, Oscillatory integrals and regularity of dispersive equations, Indiana Univ. Math. J., 40 (1991), 33–69
- [34] C. Kenig, G. Ponce and L. Vega, A bilinear estimate with applications to the KdV equation, J. Amer. Math. Soc., 9 (1996), 573–603.
- [35] N.A. Kudryashov and M.B. Soukharev, Uniformization and transcendence of solutions for the first and second Painlevé hierarchies, Phys. Lett. A, 237 (1998), 206–216.
- [36] C. Kwak, Low regularity Cauchy problem for the fifth-order modified KdV equations on T, J. Hyperbolic Differential Equations, 15 (2018), 463–557.
- [37] S. Kwon, On the fifth-order KdV equation: Local well-posedness and lack of uniform continuity of the solution map, J. Differential Equations, 245 (2008), 2627–2659.
- [38] S. Kwon, Well-posedness and ill-posedness of the fifth-order modified KdV equation, Electronic J. Differential Equations, 2008 (2008), 1–15.

- [39] J. Lenells, The nonlinear steepest descent method: asymptotics for initial-boundary value problems, SIAM J. Math. Anal., 48 (2016), 2076–2118.
- [40] F. Linares, A higher order modified Korteweg-de Vries equation, Comp. Appl. Math., 14 (1995), 253– 267.
- [41] J. Liu, P. Perry and C. Sulem, Long-time behavior of solutions to the derivative nonlinear Schrödinger equation for soliton-free initial data, Ann. I. H. Poincaré-AN, 35 (2018), 217–265.
- [42] N. Liu, B. Guo, D. Wang and Y. Wang, Long-time asymptotic behavior for an extended modified Korteweg-de Vries equation, arXiv:1907.03067v2, 2019. To appear in Commun. Math. Sci.
- [43] K.D.T.-R. McLaughlin and P. Miller, The  $\bar{\partial}$  steepest descent method for orthogonal polynomials on the real line with varying weights, Int. Math. Res. Not. IMRN, 2008 (2008), rnn075.
- [44] M. Okamoto, Long-time behavior of solutions to the fifth-order modified KdV-type equation, Adv. Differential Equations, 23 (2018), 751–792.
- [45] E. Pelinovskii, O. Polukhina and K. Lamb, Nonlinear internal waves in the ocean stratified in density and current, Oceanology, 40 (2000), 757–765.
- [46] T. Perelman, A. Fridman and M. El'yashevich, A modified Korteweg-de Vries equation in electrodynamics, Sov. Phys. JETP, 39 (1974), 643–646.
- [47] G. Ponce, Lax pairs and higher order models for water waves, J. Differential Equations, 102 (1993), 360–381.
- [48] B. Wang, Z. Huo, C. Hao and Z. Guo, Harmonic Analysis Method for Nonlinear Evolution Equations. I. World Scientific Publishing Co., Pte. Ltd., Hackensack, NJ 2011.
- [49] X. Wang, J. Zhang and L. Wang, Conservation laws, periodic and rational solutions for an extended modified Korteweg-de Vries equation, Nonlinear Dyn., 92 (2018), 1507–1516.
- [50] A.M. Wazwaz, Two new integrable modified KdV equations, of third-and fifth-order, with variable coefficients: multiple real and multiple complex soliton solutions, Waves Random Complex Media, https://doi.org/10.1080/17455030.2019.1631504, 2019.
- [51] J. Xu and E. Fan, Long-time asymptotics for the Fokas-Lenells equation with decaying initial value problem: Without solitons, J. Differential Equations, 259 (2015), 1098–1148.
- [52] V.E. Zakharov and S.V. Manakov, Asymptotic behavior of non-linear wave systems integrated by the inverse scattering method, 30 Years of the Landau Institute–Selected Papers (1996), 358–364.
- [53] X. Zhou, The Riemann–Hilbert problem and inverse scattering, SIAM J. Math. Anal., 20 (1989), 966– 986.
- [54] X. Zhou, L<sup>2</sup>-Sobolev space bijectivity of the scattering and inverse scattering transforms, Commun. Pure Appl. Math., 51 (1998), 697–731.