

AN INTRINSICALLY HYDRODYNAMIC APPROACH TO MULTIDIMENSIONAL QHD SYSTEMS

PAOLO ANTONELLI, PIERANGELO MARCATI, AND HAO ZHENG

ABSTRACT. In this paper we consider the multi-dimensional Quantum Hydrodynamics (QHD) system, by adopting an intrinsically hydrodynamic approach. The present work continues the analysis initiated in [6] where the one dimensional case was studied. Here we extend the analysis to the multi-dimensional problem, in particular by considering two physically relevant classes of solutions. First of all we consider two-dimensional initial data endowed with point vortices; by assuming the continuity of the mass density and a quantization rule for the vorticity we are able to study the Cauchy problem and provide global finite energy weak solutions. The same result can be obtained also by considering spherically symmetric initial data in the multi-dimensional setting. For rough solutions with finite energy, we are able to provide suitable dispersive estimates, which also apply to a more general class of Euler-Korteweg equations.

Moreover we are also able to show the sequential stability of weak solutions with positive density. Analogously to the one dimensional case this is achieved through the a priori bounds given by a new functional first introduced in [6].

1. INTRODUCTION

The quantum hydrodynamic (QHD) system is used as a model in various contexts in physics and engineering, such as superfluidity [50], Bose-Einstein condensation [24], quantum plasmas [37], or semiconductor devices [30]. In general such system describes phenomena whose macroscopic characterization still exhibit quantum features. From the mathematical point of view it is an Euler type system for compressible fluids in which in the current density balance equation a third order tensor appears provided by quantum potentials encoding fluid macroscopic properties due to quantum interactions. The system reads

$$(1.1) \quad \begin{cases} \partial_t \rho + \operatorname{div} J = 0 \\ \partial_t J + \operatorname{div} \left(\frac{J \otimes J}{\rho} \right) + \nabla p(\rho) = \frac{1}{2} \rho \nabla \left(\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}} \right), \end{cases}$$

with $(t, x) \in \mathbf{R} \times \mathbf{R}^d$, where the unknowns ρ and J describe the fluid particle and momentum densities, respectively. The term $p(\rho)$ is the pressure, which for convenience will be assumed to satisfy a power law, namely $p(\rho) = \frac{\gamma-1}{\gamma} \rho^\gamma$ with $1 < \gamma < \infty$ for $d = 2$ and $1 < \gamma < 3$ for $d = 3$. More general pressure laws can be taken into considerations, for instance it is possible to treat non-monotone cases, see Remark 18 for more details.

Date: March 3, 2022.

1991 Mathematics Subject Classification. Primary: 35Q40; Secondary: 76Y05, 35Q35, 82D50.

Key words and phrases. Quantum Hydrodynamics, global existence, stability, finite energy, chemical potential.

The energy of the system, formally conserved along the flow of solutions, is given by

$$(1.2) \quad E(t) = \int_{\mathbf{R}^d} \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} \frac{|J|^2}{\rho} + f(\rho) \, dx,$$

where the internal energy $f(\rho)$ is related to the pressure term by the usual relation $f(\rho) = \rho \int_0^\rho \frac{p(s)}{s^2} \, ds$.

The natural bounds given by (1.2) imply that the only available control for the velocity field v , defined by $J = \rho v$, is in L^2 with respect to the measure $\rho \, dx$. In particular, there is no control of the velocity field in the vacuum region. For this reason in order to deal with finite energy weak solutions to (1.1), it is more convenient to consider the unknowns $(\sqrt{\rho}, \Lambda)$, which define the hydrodynamic quantities by $\rho = (\sqrt{\rho})^2$, $J = \sqrt{\rho} \Lambda$, see Definition 15 below for more details.

The lack of suitable a priori bounds prevents the study of solutions to (1.1) by using compactness arguments like it is done for viscous systems, see for example [8, 7, 51, 58] where a viscous counterpart of system (1.1) is considered, see also [9, 10] where a similar system is studied by using a suitable truncation argument. On the contrary, for the QHD system most of the existing results in the literature are perturbative [47, 53, 41, 42].

On the other hand, the QHD system (1.1) enjoys an useful analogy with nonlinear Schrödinger equations, established through the Madelung transformations [59]. More specifically, given a wave function it is possible to define suitable hydrodynamical quantities, in such a way that a solution to the NLS equation determines a solution to the QHD system by means of the Madelung transform. This analogy was rigorously set up and exploited in [2, 3] in order to prove the global existence of finite energy weak solutions to (1.1) without any smallness or regularity assumptions. In particular, in [2, 3] the authors develop a polar factorization technique which overcome the difficulty of defining the velocity field in the vacuum region and allows to define the quantities $(\sqrt{\rho}, \Lambda)$ almost everywhere. The main drawback of this approach is that the initial data for (1.1) need to be consistent with some given wave function, in order to define an initial datum for the wave function dynamics. In [6] we overcome this problem in the one dimensional case by developing a wave function lifting, namely given a set of hydrodynamical quantities $(\sqrt{\rho}, \Lambda)$ it is possible to determine an associated wave function. By using the wave function lifting it is then possible to show the existence of global in time finite energy weak solutions to the one dimensional QHD system without the assumption that the initial data are generated by a wave function. Moreover, by introducing a new functional which, at least formally, is bounded over compact time intervals, it is also possible to determine a class of solutions enjoying a stability property. More precisely, by assuming the uniform bounds yielded by the total energy and the new functional, it is possible to prove that a bounded sequence of solutions in this class has a converging subsequence and that the limit is a finite energy weak solution to the one dimensional QHD system. The stability result given in [6] holds for general solutions satisfying some uniform bounds, not necessarily those originated by solutions to NLS equations.

The present paper continues the analysis initiated in [6], in order to develop an intrinsically hydrodynamic theory to QHD systems, by extending it to the multidimensional case. It is straightforward to see that considering the problem for

$d \geq 2$ entails more mathematical difficulties. If we want to follow a similar strategy as in [6], one of the main obstacles is the characterization of the vacuum region $\{\rho = 0\}$. Indeed, while in one dimension the Sobolev embedding yields the mass density to be continuous, which consequently implies that $\{\rho = 0\}$ is a closed set, in the multidimensional case we lose this information and in general it is not possible to characterize the structure of the vacuum region. Moreover, for $d \geq 2$ an extra structural property of solutions needs to be taken into account, namely the generalized irrotationality condition [2, 3], see Definition 15 and the following Remark 16. Heuristically, the generalized irrotationality condition imposes to consider solutions such that the velocity field is irrotational in the region $\{\rho > 0\}$, whereas in the vacuum region $\{\rho = 0\}$ vortices may appear. This condition is dictated by physical observations and numerical simulations for Bose-Einstein condensates and superfluid Helium II. Moreover, theoretical predictions [63, 29] say that the vorticity of a quantum fluid must be quantized, namely the circulation of the velocity field on a closed curve around a vortex must be an integer multiple of the vorticity quantum. This is consistent with the formal analogy given by the Madelung transform. Indeed given a wave function ψ , by expressing it in terms of its amplitude and phase $\psi = \sqrt{\rho}e^{iS}$, then we see that the velocity field v is given by the gradient of the phase, $v = \nabla S$. In this way the velocity field is straightforwardly irrotational, however the formal description given by the Madelung transform fails in the vacuum region, where the phase can not be defined. In [2, 3] the authors overcome the formal analogy given by the Madelung transformation by means of the polar decomposition approach. In this way it is possible to consider the hydrodynamical quantities $(\sqrt{\rho}, \Lambda)$ which are defined almost everywhere and satisfy the generalized irrotationality condition. It is straightforward to see that this condition plays also a role for the wave function lifting in the multidimensional setting.

In the present paper we are able to consider a class of initial data such that there is a countable number of non-accumulating quantized vortices and the mass density is continuous. In this framework we set up the wave function lifting, which then allows to prove the existence of global in time finite energy weak solutions to (1.1). This result allows to deal in a straightforward way also with a countable number of vortex lines on a three dimensional framework. Moreover we are able also to study another physically relevant case, namely by assuming radial symmetry of the initial data. Also in this case it is possible to establish a wave function lifting result for hydrodynamical quantities and hence to infer global existence of finite energy weak solutions to (1.1).

Moreover, in the present paper we also provide a class of purely hydrodynamical dispersive estimates for the QHD system; in fact some of them can be also stated for general Euler-Korteweg systems, see Proposition 29.

Analogously to the one dimensional case studied in [6], in our paper we also show a stability result for weak solutions to (1.1). In contrast to [6], here we need to assume positivity for the sequence of mass densities and only one of the cases proven in [6] can be extended to the multi-dimensional case.

As we already mentioned before, one of the main novelties of our analysis relies on the introduction of a new functional which provides suitable a priori bounds for solutions to (1.1). Formally, this higher order functional gives a control in $L^2(\rho dx)$

of the chemical potential associated to system (1.1), which is defined by

$$(1.3) \quad \mu = \frac{\delta E}{\delta \rho} = -\frac{1}{2} \frac{\Delta \sqrt{\rho}}{\sqrt{\rho}} + \frac{1}{2} |v|^2 + f'(\rho),$$

see also [32]. Unfortunately the chemical potential μ cannot be used to carry out a satisfactory mathematical analysis in the framework of weak solutions to (1.1). For this reason it will be more convenient to consider

$$(1.4) \quad \xi := \rho \mu = -\frac{1}{4} \Delta \rho + e + p(\rho),$$

where

$$(1.5) \quad e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho)$$

is the energy density. In this way we define

$$(1.6) \quad I(t) = \int \frac{\xi^2}{\rho} + (\partial_t \sqrt{\rho})^2 dx.$$

More precisely, by denoting

$$\lambda = \frac{\xi}{\sqrt{\rho}} \chi_{\{\rho > 0\}}$$

we can write $I(t) = \int \lambda^2 + (\partial_t \sqrt{\rho})^2 dx$. A more detailed discussion about λ and $I(t)$ will be given later, see also Section 5 in [6].

We can now present the main results of our paper.

1.1. Global existence of weak solutions for initial data with point vortices.

The first result we show deals with the existence of global in time finite energy weak solutions. As mentioned before we are going to exploit a wave function lifting argument in the two dimensional case by considering initial data with point vortices. More precisely we consider hydrodynamic data $(\rho_0(x), J_0(x))$ with $x \in \mathbf{R}^2$ such that ρ_0 is continuous, and the vacuum set $V = \{x; \rho_0(x) = 0\}$ consists of isolated points,

$$(1.7) \quad V = \{x_{(1)}, x_{(2)}, \dots, x_{(j)}, \dots\} \subset \mathbf{R}^2,$$

where in the case that V is an infinite set, we further require $\inf |x_{(j)} - x_{(k)}| = \alpha > 0$ for $j \neq k$. Let $v_0 = J_0/\rho_0$ be the velocity field, which is well-defined almost everywhere, then we say (ρ_0, J_0) satisfies the quantised vorticity condition if

$$(1.8) \quad \begin{cases} v_0 \in D'(\mathbf{R}^2) \\ \nabla \wedge v_0 = 2\pi \sum_j k_j \delta_{x_{(j)}}, \quad k_j \in \mathbf{Z}, \end{cases}$$

where $D'(\mathbf{R}^2)$ is the space of distribution, and $\delta_{x_{(j)}}$ is the Dirac-delta function supported at $x_{(j)}$ for $j = 1, 2, \dots, m$.

The condition (1.8) is dictated by physical motivations. Indeed as we discussed before one of the main features of quantum fluids is the appearance of quantized vortices [29, 61, 75]. More precisely, quantum fluids are characterized by the fact that the flow is irrotational in the region $\{\rho > 0\}$ and on the vacuum region quantized vortices may appear. This means that if $x_0 \in \mathbf{R}^2$ is a vacuum point, then the circulation of the velocity field along a closed curve around x_0 must be quantized,

$$(1.9) \quad \oint_{\gamma} v(x) \cdot d\vec{l}(x) = 2\pi k$$

for some $k \in \mathbb{Z}$. By continuity, the integer k is uniquely determined by the homotopy class of curve γ . Formally by Stokes' theorem, the quantised vorticity condition (1.8) is the distributional formulation of identity (1.9).

Theorem 1 (Global Existence of finite energy weak solutions). *Let us consider a pair $(\sqrt{\rho_0}, \Lambda_0)$ satisfying the bound*

$$(1.10) \quad \|\sqrt{\rho_0}\|_{L^\infty(0,T;H^1(\mathbf{R}^2))} + \|\Lambda_0\|_{L^\infty(0,T;L^2(\mathbf{R}^2))} \leq M_1.$$

and let us further assume that ρ_0 is continuous with isolated point vacuum

$$V = \{\rho_0 = 0\} = \{x_{(1)}, x_{(2)}, \dots, x_{(j)}, \dots\} \subset \mathbf{R}^2, \quad \inf_{j \neq k} |x_{(j)} - x_{(k)}| \geq \alpha > 0.$$

and $v_0 = J_0/\rho_0$ has quantised vorticity

$$\begin{cases} v_0 \in D'(\mathbf{R}^2) \\ \nabla \wedge v_0 = 2\pi \sum_j k_j \delta_{x_{(j)}}, \quad k_j \in \mathbf{Z}. \end{cases}$$

Then there exists a global in time finite energy weak solution to the Cauchy problem (1.1) which conserves the total energy for all time, $E(t) = E(0)$. In particular we have

$$(1.11) \quad \|\sqrt{\rho}\|_{L^\infty(0,T;H^1(\mathbf{R}^2))} + \|\Lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^2))} \leq M_1.$$

Furthermore, we are able to construct a family of more regular weak solutions. As we will see, the result we establish is not given in terms of the Sobolev regularity for $(\sqrt{\rho}, \Lambda)$ but rather in terms of the quantities involved in the analysis of the functional (1.6).

Theorem 2 (Global existence of regular weak solutions). *Consider a pair of finite energy initial data $(\sqrt{\rho_0}, \Lambda_0)$ satisfying the bound (1.10) and the quantised vorticity rules (1.8). Let us further assume that $\triangle \rho, \nabla J \in L^1_{loc}(\mathbf{R}^2)$ and $(\sqrt{\rho_0}, \Lambda_0)$ satisfies*

$$(1.12) \quad \left\| \frac{\Lambda_{0,j}^2}{\sqrt{\rho_0}} - \partial_{x_j}^2 \sqrt{\rho_0} \right\|_{L^2(\mathbf{R}^2)} + \left\| \frac{\partial_{x_j} J_{0,j}}{\sqrt{\rho_0}} \right\|_{L^2(\mathbf{R}^2)} \leq M_2, \quad j = 1, 2.$$

Then the solution to (1.1) constructed in Theorem 1 enjoys the following properties:

- *there exists $\lambda \in L^\infty(0, T; L^2(\mathbf{R}^2))$ such that*

$$(1.13) \quad \sqrt{\rho} \lambda = -\frac{1}{4} \triangle \rho + e + \rho f'(\rho)$$

where e is the total energy density

$$e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho);$$

- *for any $0 < T < \infty$ we have*

$$(1.14) \quad \|\partial_t \sqrt{\rho}\|_{L^\infty(0,T;L^2(\mathbf{R}^2))}^2 + \|\lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^2))} \leq C(T, M_1, M_2).$$

Furthermore, for such solutions the following bounds hold true

$$(1.15) \quad \|\rho\|_{L^\infty(0,T;H^2(\mathbf{R}^2))} + \|J\|_{L^\infty(0,T;H^1(\mathbf{R}^2))} + \|\sqrt{e}\|_{L^\infty(0,T;H^1(\mathbf{R}^2))} \leq C(T, M_1, M_2).$$

Remark 3. Since in Theorem 2 we assume the initial data $(\sqrt{\rho_0}, \Lambda_0)$ has points vacuum and $\triangle \rho, \nabla J \in L^1_{loc}(\mathbf{R}^2)$, the functions in condition (1.12) are well-defined almost everywhere on \mathbf{R}^2 for $j = 1, 2$. The condition (1.12) plays an essential role in characterising the higher order regularity of the lifted wave function in Proposition

21. We also remark that the bound (1.12) is preserved by the solutions we obtained in Theorem 2.

The argument of Theorem 1 and Theorem 2 also applies to typical 3D initial data. For example we can consider the 3D hydrodynamic data $(\rho_0(x), J_0(x))$, $x = (x_1, x_2, x_3) \in \mathbf{R}^3$ and $J \in \mathbf{R}^3$, with planar symmetry given in the following form:

$$(1.16) \quad \rho_0(x) = \rho_1(x_1, x_2)\rho_2(x_3) \quad \text{and} \quad J_0(x) = (J_1(x_1, x_2)\rho_2(x_3), 0) = \sqrt{\rho_0}\Lambda_0,$$

where $\rho_j \in \mathbf{R}$ and $J_1 \in \mathbf{R}^2$. We assume $(\rho_1, J_1 = \sqrt{\rho_1}\Lambda_1)$ has point vortices, namely ρ_1 is continuous with vacuum structure (1.7), and the velocity field $v_1 = J_1/\rho_1$ satisfies the quantised vorticity condition (1.8). In this case the vortices of $(\rho_0(x), J_0(x))$ appear as parallel straight lines in x_3 direction, which is a widely studied vortex structure of quantum flows in 3D space as vortex lines [13]. For such initial data, we can still prove the global existence of finite energy weak solutions and weak solutions in the regular class of the functional $I(t)$.

Theorem 4. *Let (ρ_0, J_0) be a pair of hydrodynamic data as in (1.16), with (ρ_1, J_1) satisfying the condition of point vortices (1.7) and (1.8). We assume that (ρ_1, J_1) satisfies the bound*

$$\|\sqrt{\rho_1}\|_{L^\infty(0,T;H^1(\mathbf{R}^2))} + \|\Lambda_1\|_{L^\infty(0,T;L^2(\mathbf{R}^2))} \leq M_1.$$

and $\sqrt{\rho_2} \in H^1(\mathbf{R})$. Then there exists a global in time finite energy weak solution to the Cauchy problem (1.1) which conserves the total energy for all times, $E(t) = E(0)$. In particular we have

$$(1.17) \quad \|\sqrt{\rho}\|_{L^\infty(0,T;H^1(\mathbf{R}^3))} + \|\Lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^3))} \leq M_1.$$

Theorem 5. *If we furthermore assume the controls*

$$\left\| \frac{\Lambda_{1,j}^2}{\sqrt{\rho_1}} - \partial_{x_j}^2 \sqrt{\rho_1} \right\|_{L^2(\mathbf{R}^2)} + \left\| \frac{\partial_{x_j} J_{1,j}}{\sqrt{\rho_1}} \right\|_{L^2(\mathbf{R}^2)} \leq M_2, \quad j = 1, 2.$$

on (ρ_1, J_1) and $\sqrt{\rho_2} \in H^2(\mathbf{R})$, then the functional $I(t)$ stays finite along the solution to (1.1) constructed in Theorem 4, namely exists $\lambda \in L^\infty(0, T; L^2(\mathbf{R}^3))$ such that

$$\sqrt{\rho}\lambda = -\frac{1}{4}\triangle\rho + e + \rho f'(\rho)$$

and for any $0 < T < \infty$ we have

$$\|\partial_t \sqrt{\rho}\|_{L^\infty(0,T;L^2(\mathbf{R}^3))}^2 + \|\lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^3))} \leq C(T, M_1, M_2).$$

Furthermore, for such solutions the following bounds hold true

$$\|\rho\|_{L^\infty(0,T;H^2(\mathbf{R}^3))} + \|J\|_{L^\infty(0,T;H^1(\mathbf{R}^3))} + \|\sqrt{e}\|_{L^\infty(0,T;H^1(\mathbf{R}^3))} \leq C(T, M_1, M_2).$$

1.2. Global existence of spherically symmetric weak solutions. Now we consider the spherically symmetric hydrodynamic data in the sense that

$$\sqrt{\rho_0(x)} = \sqrt{\rho_0(r)} \quad \text{and} \quad \Lambda_0(x) = \Lambda_0(r) \frac{x}{|x|},$$

where $x \in \mathbf{R}^d$, $r = |x|$ and $\Lambda_0(r) \in \mathbf{R}$. With suitable regularity assumption on $(\sqrt{\rho_0}, \Lambda_0)$ we can prove the global existence of spherically symmetric weak solutions to the QHD system in the finite energy space and the regular class of the functional (1.6).

Theorem 6 (Global Existence of finite energy weak solutions). *Let $(\sqrt{\rho_0}, \Lambda_0)$ be a pair of spherically symmetric hydrodynamic data*

$$\sqrt{\rho_0(x)} = \sqrt{\rho_0(r)} \quad \text{and} \quad \Lambda_0(x) = \Lambda_0(r) \frac{x}{|x|}$$

with $\Lambda_0 = 0$ a.e. on the set $\{\rho_0 = 0\}$. Let us further assume the bounds

$$(1.18) \quad \|\sqrt{\rho_0}\|_{H^1(\mathbf{R}^d)} + \|\Lambda_0\|_{L^2(\mathbf{R}^d)} \leq M_1,$$

Then there exists a spherically symmetric global in time finite energy weak solution to the Cauchy problem (1.1) which conserves the total energy for all times, i.e. $E(t) = E(0)$. In particular we have

$$\|\sqrt{\rho}\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} + \|\Lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq M_1.$$

Analogously to the case with point vortices, also here we provide an existence result for regular weak solutions.

Theorem 7 (Global existence of regular weak solutions). *Let $(\sqrt{\rho_0}, \Lambda_0)$ be a pair of spherically symmetric finite energy initial satisfying (1.18). We further assume that the initial data satisfy the higher order bounds*

$$(1.19) \quad \left\| \frac{\Lambda_0^2}{\sqrt{\rho_0}} \mathbf{1}_{\{\rho_0 > 0\}} \right\|_{L^2(\mathbf{R}^d)} + \|\Delta \sqrt{\rho_0}\|_{L^2(\mathbf{R}^d)} + \left\| \frac{\operatorname{div} J_0}{\sqrt{\rho_0}} \mathbf{1}_{\{\rho_0 > 0\}} \right\|_{L^2(\mathbf{R}^d)} \leq M_2,$$

then there exists $\lambda \in L^\infty(0, T; L^2(\mathbf{R}^d))$ such that $\lambda(x) = \lambda(|x|)$ and

$$(1.20) \quad \sqrt{\rho} \lambda = -\frac{1}{4} \Delta \rho + e + p(\rho),$$

where e is the total energy density

$$e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho),$$

and for any $0 < T < \infty$ we have

$$(1.21) \quad \|\partial_t \sqrt{\rho}\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} + \|\lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C(T, M_1, M_2).$$

Furthermore, for such solutions the following bounds hold true

$$\|\rho\|_{L^\infty(0,T;H^2(\mathbf{R}^d))} + \|J\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} + \|\sqrt{e}\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} \leq C(T, M_1, M_2),$$

Remark 8. Let us notice that assumption (1.19) is stronger than its analogue (1.12) in Theorem 2. This is somehow reminiscent of the one dimensional case, where in Theorem 2 in [6] a similar assumption is considered. Contrarily to (1.12), condition (1.19) in general is not preserved along the dynamics of (1.1). It is interesting to weaken this assumption in such a way that it can be propagated along the flow of solutions to (1.1); this will be the subject of future investigations.

1.3. Dispersive estimates. The Theorems above show the existence of global in time weak solutions to (1.1), moreover they also show that for such solutions the functional $I(t)$ defined in (1.6) remains bounded over compact time intervals. However no uniqueness is provided here and in general such result does not hold true, see [26, 60]. On the other hand, we are able to prove some dispersive estimates in genuine hydrodynamic approaches, which hold for general finite energy weak solutions.

Theorem 9 (Dispersive estimates). *Let (ρ, J) be a finite energy weak solution to (1.1) satisfying the bound*

$$\|\sqrt{\rho}\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} + \|\Lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq M_1.$$

(i) *We have*

$$(1.22) \quad \||\nabla|^{\frac{3-d}{2}}\rho\|_{L^2(\mathbf{R}_t \times \mathbf{R}_x^d)}^2 + \||\nabla|^{\frac{1-d}{2}}\sqrt{\rho p(\rho)}\|_{L^2(\mathbf{R}_t \times \mathbf{R}_x^d)}^2 \lesssim M_1^4.$$

(ii) *If we further assume that $\int_{\mathbf{R}^d} |x|^2 \rho_0(x) dx < \infty$, then*

$$(1.23) \quad \|\nabla\sqrt{\rho}(t)\|_{L^2} + \|\Lambda(t) - \frac{x}{t}\sqrt{\rho}(t)\|_{L^2} + \|\rho^{\frac{\gamma}{2}}(t)\|_{L^2} \lesssim t^{-\sigma},$$

where $\sigma = \min\{1, \frac{d}{2}(\gamma - 1)\}$.

Remark 10. The dispersive estimates (1.22) and (1.23) hold for general weak solutions to the QHD system with finite energy. In fact such estimates apply to a more general class of Euler-Korteweg equations, which is stated in Proposition 29.

1.4. Stability. Next we prove a compactness property that the class of weak solutions with positive mass density and satisfying the uniform bounds given by the energy and the functional $I(t)$. Let us remark that in contrast to existing literature on QHD system, we do not need a uniform low bound on the mass density, and also our result below hold for a general class of solutions, not necessarily those one constructed in the existence Theorems given above.

Theorem 11 (Stability). *Let us assume $\{(\sqrt{\rho_n}, \Lambda_n)\}_{n \geq 1}$ is a sequence of weak solutions to (1.1) in the compactness class, which satisfies the uniform bounds*

$$(1.24) \quad \begin{aligned} &\|\sqrt{\rho_n}\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} + \|\sqrt{\rho_n}\|_{W^{1,\infty}(0,T;L^2(\mathbf{R}^d))} \leq C \\ &\|\Lambda_n\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C, \quad \|\lambda_n\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C, \end{aligned}$$

with λ_n satisfying (1.13). *Let us also assume that for all $n \in \mathbf{N}$ and almost every $t \in [0, T]$, $\rho_n(t, \cdot)$ is continuous with $\rho_n(t, \cdot) > 0$. Then up to subsequences we have that for any $0 < T < \infty$*

$$\begin{aligned} \sqrt{\rho_n} &\rightarrow \sqrt{\rho} \quad \text{in } L^p(0, T; H_{loc}^1(\mathbf{R}^d)), \\ \Lambda_n &\rightarrow \Lambda \quad \text{in } L^p(0, T; L_{loc}^2(\mathbf{R}^d)), \end{aligned}$$

where $(\sqrt{\rho}, \Lambda)$ is a finite energy weak solution to (1.1) and $2 \leq p < \infty$.

We conclude this introduction by discussing some results already existing in the mathematical literature. The QHD system, together with similar evolutionary PDE models, is widely studied. Previous results can be found in [47, 53, 31, 45, 41, 42]. System (1.1) is also intimately related to the class of Euler-Korteweg fluids [15], encoding capillary effects in the description; local and global analysis of small, regular perturbations of constant states are discussed in [17, 11]. The method of convex integration can be applied also to Euler-Korteweg system in order to show the existence of infinitely many solutions emanating from the same initial data [26], see also [60] where a non-uniqueness result for system (1.1) is given by adopting a different strategy, related to the underlying wave function dynamics. Recently also a class of viscous quantum fluid dynamical systems was considered. Such models can be derived from the Wigner-Fokker-Planck equation [46], see also the review [43]. The analysis of finite energy weak solutions for the quantum Navier-Stokes

system was done in [44, 8, 7, 51], see also [52, 54] where similar arguments are used to study the compressible Navier-Stokes system with degenerate viscosity.

We structure the contents of this chapter as follows. Some notations and preliminary results that will be used through this paper are given in Section 2. The wave function lifting method, together with the existence Theorem 1, for hydrodynamic data with pointwise vacuum and quantised vorticity is proved in Section 3. On the other hand in Section 4 we introduce the spherically symmetric QHD system and prove the wave function lifting method for spherically symmetric data, and as a consequence the global existence Theorem 6 for spherically symmetric initial data is also obtained in Section 4. In section 5 we provide some a priori estimates on solutions to system (1.1), and finally we show the stability of solutions in Section 6.

2. NOTATIONS AND PRELIMINARIES

In this Section we fix the notation that will be used through this paper.

We use the standard notation for Lebesgue and Sobolev norms

$$\|f\|_{L_x^p} := \left(\int_{\mathbf{R}^d} |f(x)|^p dx \right)^{\frac{1}{p}},$$

$$\|f\|_{W_x^{k,p}} := \sum_{j=0}^k \|\nabla^j f\|_{L_x^p},$$

and we denote $H_x^k := H^k(\mathbf{R}^d) = W^{k,2}(\mathbf{R}^d)$. The mixed Lebesgue norm of functions $f : I \rightarrow L^r(\mathbf{R}^d)$ is defined as

$$\|f\|_{L_t^q L_x^r} := \left(\int_I \|f(t)\|_{L_x^r}^q dt \right)^{\frac{1}{q}} = \left(\int_I \left(\int_{\mathbf{R}^d} |f(x)|^r dx \right)^{\frac{q}{r}} dt \right)^{\frac{1}{q}},$$

where $I \subset [0, \infty)$ is a time interval. Similarly the mixed Sobolev norm $L_t^q W_x^{k,r}$ is defined.

We recall here some basic properties of the following one-dimensional nonlinear Schrödinger equation

$$(2.1) \quad \begin{cases} i\partial_t \psi = -\frac{1}{2}\Delta \psi + |\psi|^{2(\gamma-1)}\psi \\ \psi(0) = \psi_0, \end{cases}$$

and we mainly consider $\gamma \in (1, \infty)$ for $x \in \mathbf{R}^2$ and $1 < \gamma < 3$ for $x \in \mathbf{R}^3$.

The reader can find more details and the proofs of the statements of the following Theorem in the comprehensive monographs [19, 56, 74].

Theorem 12. *Let $\psi_0 \in H^1(\mathbf{R}^d)$, $\gamma > 1$ for $d = 2$ and $1 < \gamma < 3$ for $d = 3$, then there exists a unique global solution $\psi \in \mathcal{C}(\mathbf{R}; H^1(\mathbf{R}^d))$ to (2.1) such that the total mass and energy are conserved at all times, i.e.*

$$(2.2) \quad \begin{aligned} M(t) &= \int_{\mathbf{R}^d} |\psi(t, x)|^2 dx = M(0) \\ E(t) &= \int_{\mathbf{R}^d} \frac{1}{2} |\nabla \psi|^2 + \frac{1}{\gamma} |\psi|^{2\gamma} dx = E(0). \end{aligned}$$

If moreover $\psi_0 \in H^2(\mathbf{R}^d)$, then we also have $\psi \in \mathcal{C}(\mathbf{R}; H^2(\mathbf{R}^d)) \cap \mathcal{C}^1(\mathbf{R}; L^2(\mathbf{R}^d))$ and for any $0 < T < \infty$

$$(2.3) \quad \|\psi\|_{L^\infty(0,T;H^2(\mathbf{R}^d))} + \|\partial_t \psi\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C(T, \|\psi_0\|_{H^2(\mathbf{R}^d)}).$$

Next we are going to recall the polar factorization technique developed in [2, 3], see also the reviews [4, 5]. This method allows to define the hydrodynamic quantities $(\sqrt{\rho}, \Lambda)$ and sets up a correspondence between the wave function dynamics and the hydrodynamical system. The main advantage of this approach with respect to the usual WKB method for instance is that vacuum regions are allowed in the theory. Here we only give a brief introduction and state the results we will exploit later, for a more detailed presentation and for a proof of the statements in Lemma 13 below we address to Section 3 in [1].

Given any function $\psi \in H^1(\mathbf{R}^d)$ we can define the set of polar factors as

$$(2.4) \quad P(\psi) := \{\phi \in L^\infty(\mathbf{R}^d) \mid \|\phi\|_{L^\infty} \leq 1, \sqrt{\rho}\phi = \psi \text{ a.e.}\},$$

where $\sqrt{\rho} := |\psi|$. In general this set can contain more than one element, due to the possible appearance of vacuum regions. Nevertheless, as it will be shown in the next Lemma, the hydrodynamical quantities $(\sqrt{\rho}, \Lambda)$ are well defined and they furthermore enjoy suitable stability properties.

Lemma 13 (Polar factorization [2, 3]). *Let $\psi \in H^1(\mathbf{R}^d)$ and $\sqrt{\rho} := |\psi|$, and let $\phi \in P(\psi)$. Then we have $\nabla \sqrt{\rho} = \text{Re}(\bar{\phi} \nabla \psi) \in L^2(\mathbf{R}^d)$ and if we set $\Lambda := \text{Im}(\bar{\phi} \nabla \psi)$, we have*

$$(2.5) \quad \text{Re}(\nabla \bar{\psi} \otimes \nabla \psi) = \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho} + \Lambda \otimes \Lambda, \quad \text{a.e. in } \mathbf{R}^d.$$

Moreover if $\{\psi_n\} \subset H^1$ satisfies $\|\psi_n - \psi\|_{H^1} \rightarrow 0$ as $n \rightarrow \infty$, then we have

$$\nabla \sqrt{\rho_n} \rightarrow \nabla \sqrt{\rho}, \quad \Lambda_n \rightarrow \Lambda, \quad \text{in } L^2,$$

with $\sqrt{\rho_n} := |\psi_n|$, $\Lambda_n := \text{Im}(\bar{\phi}_n \nabla \psi_n)$, ϕ_n being a unitary factor for ψ_n .

We stress here the fact that $\Lambda = \text{Im}(\bar{\phi} \nabla \psi)$ is well-defined even if ϕ is not uniquely determined, this is a consequence of Theorem 6.19 in [55], which we will state below as it will be thoroughly used in our exposition.

Lemma 14. *Let $g : \Omega \rightarrow \mathbb{R}$ be in $H^1(\Omega)$, and*

$$B = g^{-1}(\{0\}) = \{x \in \Omega : g(x) = 0\}.$$

Then $\nabla g(x) = 0$ for almost every $x \in B$.

By combining the well-posedness result for the NLS equation (2.1) stated in Theorem 12 and the polar factorization method recalled in Lemma 13 we can prove an existence result for finite energy weak solutions to (1.1), see Theorem 17 below. This result is proved in [2, 3] for the two and three dimensional cases, see also the review [1].

Before stating the results let us first recall a useful identity regarding the non-linear dispersive term on the right hand side of the equation for the momentum density in (1.1), namely it can also be written as

$$(2.6) \quad \frac{1}{2} \rho \nabla \left(\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}} \right) = \frac{1}{4} \text{div}(\rho \nabla^2 \log \rho) = \frac{1}{4} \nabla \Delta \rho - \text{div}(\nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho}).$$

By using identity (2.6), system (1.1) then becomes

$$(2.7) \quad \begin{cases} \partial_t \rho + \operatorname{div} J = 0 \\ \partial_t J + \operatorname{div} (\Lambda \otimes \Lambda + \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho}) + \nabla p(\rho) = \frac{1}{4} \nabla \Delta \rho \end{cases}$$

We can now give the definition of finite energy weak solutions.

Definition 15 (Finite energy weak solutions). Let $\rho_0, J_0 \in L^1_{loc}(\mathbf{R}^d)$, we say the pair (ρ, J) is a finite energy weak solution to the Cauchy problem for (1.1) with initial data $\rho(0) = \rho_0, J(0) = J_0$, in the space-time slab $[0, T) \times \mathbf{R}^d$ if there exist $\sqrt{\rho} \in L^2_{loc}(0, T; H^1_{loc}(\mathbf{R}^d)), \Lambda \in L^2_{loc}(0, T; L^2_{loc}(\mathbf{R}^d))$ such that

- (i) $\rho := (\sqrt{\rho})^2, J := \sqrt{\rho} \Lambda;$
- (ii) $\forall \eta \in \mathcal{C}_0^\infty([0, T) \times \mathbf{R}^d),$

$$\int_0^T \int_{\mathbf{R}^d} \rho \partial_t \eta + J \cdot \nabla \eta \, dx dt + \int_{\mathbf{R}^d} \rho_0(x) \eta(0, x) \, dx = 0;$$

- (iii) $\forall \zeta \in \mathcal{C}_0^\infty([0, T) \times \mathbf{R}^d; \mathbf{R}^d),$

$$\begin{aligned} & \int_0^T \int_{\mathbf{R}^d} J \cdot \partial_t \zeta + \Lambda \otimes \Lambda : \nabla \zeta + p(\rho) \operatorname{div} \zeta + \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho} : \nabla \zeta \\ & + \frac{1}{4} \rho \Delta \operatorname{div} \zeta \, dx dt + \int_{\mathbf{R}^d} J_0(x) \cdot \zeta(0, x) \, dx = 0; \end{aligned}$$

- (iv) (finite energy) the total energy defined by (1.2) is finite for almost every $t \in [0, T)$;
- (v) (generalized irrotationality condition) for almost every $t \in (0, T)$

$$\nabla \wedge J = 2 \nabla \sqrt{\rho} \wedge \Lambda,$$

holds in the sense of distributions.

We say (ρ, J) is a global in time finite energy weak solution if we can take $T = \infty$ in the above definition.

Remark 16. In the case of a smooth solution (ρ, J) , for which we can write $J = \rho v$, for some smooth velocity field v , the generalized irrotationality condition defined above is equivalent to $\rho \nabla \wedge v = 0$, i.e. the velocity field v is irrotational $\rho \, dx$ almost everywhere. It shows that the previous definition is the right weak formulation of the classical irrotationality condition $\nabla \wedge v = 0$ valid away from vacuum in the WKB approach. The generalized irrotationality condition is also motivated by the quantised vortices of quantum flows, namely the flow is irrotational outside the set $\{\rho = 0\}$ and in the vacuum the vorticity becomes singular. In this respect the solutions introduced in Definition 15 are more general than those obtained by using the WKB ansatz, since in the latter case the velocity field $v = \nabla S$ is always irrotational and there is no vacuum. Our study is motivated also by physical considerations, since quantized vortices have very rich structures which lead to interesting phenomena in quantum turbulence [75].

For initial data generated by wave functions, the existence of finite energy weak solutions was proved in [2, 3]. We address the interested reader to [2, 3], see also Theorem 4.2. in [1].

Theorem 17. *Let $\psi_0 \in H^1(\mathbf{R}^d)$ and define the initial data for the QHD system (1.1) as $\rho_0 := |\psi_0|^2$, $J_0 := \text{Im}(\bar{\psi}_0 \nabla \psi_0)$. Then there exists a global in time finite energy weak solution such that $\sqrt{\rho} \in L^\infty(\mathbf{R}; H^1(\mathbf{R}^d))$, $\Lambda \in L^\infty(\mathbf{R}; L^2(\mathbf{R}^d))$, which conserves the energy at all times.*

Remark 18. Let us remark that the proof of Theorem 17 relies on two facts, the polar factorization technique of Lemma 13 and the well-posedness result in the energy space for the associated NLS equation. This means that in general the result in Theorem 17 remains true as long as the associated NLS equation is globally well-posed and enjoys suitable stability properties. In particular this implies that it is also possible to study QHD systems with non-monotone pressure terms.

We conclude this Section by giving a discussion about the higher order functional introduced in (1.6) and a definition for the quantity λ involved, formally related to the chemical potential μ by $\lambda = \sqrt{\rho}\mu$.

As mentioned in the Introduction, to obtain the stability property of weak solutions, we will exploit the a priori bounds inferred from the functional

$$I(t) = \int_{\mathbf{R}^d} \lambda^2 + (\partial_t \sqrt{\rho})^2 dx,$$

where $\lambda \in L^\infty(0, T; L^2(\mathbf{R}^d))$ is implicitly given by

$$(2.8) \quad \xi = \sqrt{\rho}\lambda = -\frac{1}{4}\Delta\rho + \frac{1}{2}|\nabla\sqrt{\rho}|^2 + \frac{1}{2}|\Lambda|^2 + f'(\rho)\rho.$$

The functional $I(t)$ has a direct interpretation for Schrödinger-generated solutions. Indeed let us assume for the moment that we have $\rho = |\psi|^2$ and $J = \text{Im}(\bar{\psi}\nabla\psi)$, for some solutions ψ to (2.1). Then we can see that the functional (1.6) actually equals

$$I(t) = \int |\partial_t \psi|^2 dx.$$

Indeed by using the polar factorization we have

$$\partial_t \sqrt{\rho} = \text{Re}(\bar{\phi} \partial_t \psi), \quad \lambda = -\text{Im}(\bar{\phi} \partial_t \psi),$$

where $\phi \in P(\psi)$ is a polar factor. On the other hand, by invoking Lemma 14, we see that for Schrödinger-generated functions, λ vanishes almost everywhere on the vacuum region $\{\rho = 0\}$. Let us remark that in this case identity (2.8) is recovered from $\xi = \sqrt{\rho}\lambda = -\text{Im}(\bar{\psi}\partial_t \psi)$ and by exploiting the fact that ψ is a solution to (2.1). Following this intuition, we give a rigorous definition for arbitrary, not necessarily Schrödinger-generated, solutions.

Definition 19. Let (ρ, J) be a finite energy weak solution to (1.1) such that ρ is continuous, $\Delta\rho \in L^1_{loc}(\mathbf{R}^d)$ and $\Lambda = 0$ a.e. on $\{\rho = 0\}$. Then we define λ to be measurable function given by

$$(2.9) \quad \lambda = \begin{cases} -\frac{1}{2}\Delta\sqrt{\rho} + \frac{1}{2}\frac{|\Lambda|^2}{\sqrt{\rho}} + f'(\rho)\sqrt{\rho} & \text{in } \{\rho > 0\} \\ 0 & \text{elsewhere} \end{cases}$$

Note however that in order to prove our stability result given in Theorem 11 we will only need λ to satisfy $\lambda \in L^\infty_t L^2_x$ and identity (2.8). In fact we will identify the class of weak solutions for which we can prove our stability result exactly by those two properties satisfied by λ .

Definition 20. Let (ρ, J) be a finite energy weak solution to (1.1) on $[0, T]$ as in Definition 15. We say (ρ, J) belongs to compactness class if there exists a $\lambda \in L^\infty(0, T; L^2(\mathbf{R}^d))$ such that

$$\sqrt{\rho}\lambda = -\frac{1}{4}\Delta\rho + e + p(\rho),$$

where e is the energy density

$$e = \frac{1}{2}|\nabla\sqrt{\rho}|^2 + \frac{1}{2}|\Lambda|^2 + f(\rho)$$

and for a.e. $t \in [0, T]$ the functional

$$I(t) = \int_{\mathbf{R}^d} \lambda^2 + (\partial_t \sqrt{\rho})^2 dx \leq C.$$

Resuming, a finite energy weak solution $(\sqrt{\rho}, \Lambda)$ belonging to the compactness class specified in the Definition above satisfies the following bounds

$$\begin{aligned} \|\sqrt{\rho}\|_{L^\infty H^1} + \|\Lambda\|_{L^\infty L^2} &\leq M_1, \\ \|\partial_t \sqrt{\rho}\|_{L^\infty L^2} + \|\lambda\|_{L^\infty L^2} &\leq M_2, \end{aligned}$$

with λ satisfying identity (2.8).

3. WAVE FUNCTION LIFTING: HYDRODYNAMIC DATA WITH POINTS VACUUM

The main tool we use to prove the global existence results is the wave function lifting method. For given hydrodynamic data $(\sqrt{\rho}, \lambda)$ with suitable assumptions, we show that it is possible to define a wave function ψ associated to $(\sqrt{\rho}, \lambda)$. The global existence of weak solutions to (1.1) is obtained by solving the NLS equation with initial data ψ and using the polar factorization and Theorem 17.

This section mostly focuses on the two-dimensional case. We will assume that the mass density $\rho = \rho(x)$, $x \in \mathbf{R}^2$, is continuous and that the vacuum set $V = \{x; \rho(x) = 0\}$ consists of isolated points,

$$(3.1) \quad V = \{x_{(1)}, x_{(2)}, \dots, x_{(j)}, \dots\} \subset \mathbf{R}^2, \quad \inf_{j \neq k} |x_{(j)} - x_{(k)}| \geq \alpha > 0.$$

Moreover we also consider a quantization condition for the vorticity. More precisely, let $v = J/\rho$ be the velocity field, which is well-defined almost everywhere, then the quantised vorticity condition is stated as following:

$$(3.2) \quad \begin{cases} v \in D'(\mathbf{R}^2) \\ \nabla \wedge v = 2\pi \sum_j k_j \delta_{x_{(j)}}, \quad k_j \in \mathbf{Z}, \end{cases}$$

where $D'(\mathbf{R}^2)$ is the space of distribution, and $\delta_{x_{(j)}}$ is the Dirac-delta function supported at $x_{(j)}$ for $j = 1, 2, \dots, m$. As discussed in Remark 16, the quantised vorticity condition is a special case of the generalised irrotationality condition

$$\nabla \wedge J = 2\nabla\sqrt{\rho} \wedge \Lambda.$$

We always assume $(\rho, J = \sqrt{\rho}\Lambda)$ has finite mass $M(\rho)$ and finite energy $E(\rho, J)$, which is equivalent to

$$\|\sqrt{\rho}\|_{H^1(\mathbf{R}^2)} + \|\Lambda\|_{L^2(\mathbf{R}^2)} \leq M_1.$$

On the other hand, in the case of pointwise vacuum to characterise the higher order regularity of the data, we assume the condition that $\Delta\rho, \nabla J \in L^1_{loc}(\mathbf{R}^2)$ and

$$\left\| \frac{\Lambda_j^2}{\sqrt{\rho}} - \partial_{x_j}^2 \sqrt{\rho} \right\|_{L^2(\mathbf{R}^2)} + \left\| \frac{\partial_{x_j} J_j}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}^2)} \leq M_2, \quad j = 1, 2,$$

where the functions in the right hand side of the inequality are well-defined a.e. on \mathbf{R}^2 .

As a technical tool, we refer the theory of difference quotient, which will be used to prove the regularity of the wave function constructed from given hydrodynamic data. The difference quotient of a function $f \in L^1_{loc}(\mathbf{R}^2)$ in direction $h \in \mathbf{R}^2$, $|h| > 0$ is defined by

$$D^h f(x) = \frac{f(x+h) - f(x)}{|h|},$$

and there exists a equivalence between the L^p norms of $D^h f$ uniform in h and the $W^{1,p}$ norms of f for $1 < p < \infty$. For readers interested in the details, we refer the Theorem 3 of Section 5.8.2 in [27].

Now we are at the point to prove the wave function lifting proposition. We remark that the continuity of density ρ , which can not be inferred by the finite energy assumption and Sobolev embedding in multi-dimension, is essential to give a proper definition to the point vacuum, and it is also needed to provide a local lower bound on the density when strictly away from the vacuum. Nevertheless, in the physical and mathematical literature the most widely studied vortex structures fall in this framework, see for example [18, 64, 70].

Proposition 21. *Let $(\sqrt{\rho}, \Lambda)$ be a pair of finite energy hydrodynamic data satisfying the bounds*

$$(3.3) \quad \|\sqrt{\rho}\|_{H^1(\mathbf{R}^2)} + \|\Lambda\|_{L^2(\mathbf{R}^2)} \leq M_1.$$

Let us further assume that $\sqrt{\rho}$ is continuous with isolated point vacuum

$$(3.4) \quad V = \{\rho = 0\} = \{x_{(1)}, x_{(2)}, \dots, x_{(j)}, \dots\} \subset \mathbf{R}^2, \quad \inf_{j \neq k} |x_{(j)} - x_{(k)}| = \alpha > 0.$$

and quantised vorticity

$$(3.5) \quad \begin{cases} v \in D'(\mathbf{R}^2) \\ \nabla \wedge v = 2\pi \sum_j k_j \delta_{x_{(j)}}, \quad k_j \in \mathbf{Z}. \end{cases}$$

Then there exists a wave function $\psi \in H^1(\mathbf{R}^2)$ such that

$$\sqrt{\rho} = |\psi|, \quad \Lambda = \text{Im}(\bar{\phi} \nabla \psi),$$

where $\phi \in P(\psi)$ is a polar factor defined in (2.4).

If we furthermore assume that $\Delta\rho \in L^1_{loc}(\mathbf{R}^2)$ and $(\sqrt{\rho}, \Lambda)$ satisfy also the bounds

$$(3.6) \quad \left\| \frac{\Lambda_j^2}{\sqrt{\rho}} - \partial_{x_j}^2 \sqrt{\rho} \right\|_{L^2(\mathbf{R}^2)} + \left\| \frac{\partial_{x_j} J_j}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}^2)} \leq M_2, \quad j = 1, 2,$$

then $\psi \in H^2(\mathbf{R}^2)$ and we have

$$\|\psi\|_{H^2(\mathbf{R}^2)} \leq C(M_1, M_2).$$

Proof. We start with the simplest case, the vacuum is a single point, and without loss of generality we assume $V = \{0\}$. In this case the quantised vorticity condition becomes

$$\begin{cases} v \in D'(\mathbf{R}^2) \\ \nabla \wedge v = 2k\pi\delta_0, \quad k \in \mathbf{Z}. \end{cases}$$

For isolated vacuum points, the proof follows essentially same idea.

To construct a wave function we take a sequence of smooth mollifier $\{\chi_\epsilon\}$ such that $\text{supp}(\chi_\epsilon) \subset B_\epsilon(0)$, where $B_r(x)$ is the ball centred at x with radius r . Since $v \in D'(\mathbf{R}^2)$ we can mollify

$$v^\epsilon(x) = v * \chi_\epsilon(x) = \langle v(\cdot), \chi_\epsilon(x - \cdot) \rangle,$$

where $\langle \cdot \rangle$ is the pairing of distributions and test functions. By the theory of distribution v^ϵ is a smooth function, and

$$\begin{aligned} \nabla \wedge v^\epsilon &= (\nabla \wedge v) * \chi_\epsilon \\ &= 2k\pi\delta_0 * \chi_\epsilon = 2k\pi\chi_\epsilon. \end{aligned}$$

We fix a base point $x_0 \notin B_\epsilon(0)$, then for any $x \notin B_\epsilon(0)$ we can take a piecewise smooth curve $\gamma(x)$ that connects x_0 to x and is strictly away from $B_\epsilon(0)$. The approximating polar factor ϕ^ϵ is defined as

$$\phi^\epsilon(x) = \exp(i \int_{\gamma(x)} v^\epsilon(y) \cdot d\vec{l}(y))$$

for $x \notin B_\epsilon(0)$, and for $x \in B_\epsilon(0)$ we set $\phi^\epsilon(x) = \phi^\epsilon(\epsilon x/|x|)$.

The definition of ϕ^ϵ is independent of the choice of curve γ : for any two curves γ_1 and γ_2 connecting x_0 and x , we can attach them into a piecewise smooth closed curve $\tilde{\gamma}$. Denote Ω the domain inside $\tilde{\gamma}$, then we have two cases: $B_\epsilon(0) \subset \Omega$ or $B_\epsilon(0) \cap \Omega = \emptyset$. In either case by the Stokes' theorem

$$\begin{aligned} \int_{\gamma_1} v^\epsilon(y) \cdot d\vec{l}(y) - \int_{\gamma_2} v^\epsilon(y) \cdot d\vec{l}(y) &= \int_{\tilde{\gamma}} v^\epsilon(y) \cdot d\vec{l}(y) \\ &= \int_{\Omega} \nabla \wedge v^\epsilon(y) dy \\ &= 2k\pi \text{ or } 0, \end{aligned}$$

thus we have

$$\exp(i \int_{\gamma_1} v^\epsilon(y) \cdot d\vec{l}(y)) = \exp(i \int_{\gamma_2} v^\epsilon(y) \cdot d\vec{l}(y)).$$

The polar factors $\{\phi^\epsilon\}$ have uniform L^∞ bound 1, therefore we can take a subsequence which has a weak* limit ϕ with $\|\phi\|_{L^\infty} \leq 1$. Without changing the notation we have $\phi^\epsilon \xrightarrow{*} \phi$ in $L^\infty(\mathbf{R}^2)$. The target wave function is defined by $\psi = \sqrt{\rho}\phi$. Obviously $\psi \in L^2(U)$, then next step is to prove $\psi \in H^1(\mathbf{R}^2)$.

To prove $\psi \in H^1(\mathbf{R}^2)$ we only need to show

$$\|D^h\psi\|_{L^2(\mathbf{R}^2)} \leq C$$

for some constant C and all $h \in \mathbf{R}^2$ with $|h| \neq 0$ small. Given a fixed $h \in \mathbf{R}^2$, we divide \mathbf{R}^2 into $B_{|h|}(0) \cup B_{|h|}(0)^c$, and write

$$\|D^h\psi\|_{L^2(\mathbf{R}^2)} = \|D^h\psi\|_{L^2(B_{|h|}(0))} + \|D^h\psi\|_{L^2(B_{|h|}(0)^c)}.$$

For the part near the vacuum we simply bound it by

$$\begin{aligned} \|D^h \psi\|_{L^2(B_{|h|}(0))} &= \frac{1}{|h|} \|\psi(\cdot + h) - \psi(\cdot)\|_{L^2(B_{|h|}(0))} \\ &\leq \frac{2}{|h|} \|\sqrt{\rho}\|_{L^\infty(B_1(0))} |B_{|h|}(0)|^{\frac{1}{2}} = 2\sqrt{\pi} \|\sqrt{\rho}\|_{L^\infty(B_1(0))}, \end{aligned}$$

which is uniformly bounded since $\sqrt{\rho}$ is continuous.

Then we consider the term away from the vacuum. For any open set $U \subset B_{|h|}(0)^c$ such that \overline{U} is compact, and for any $\epsilon < h$ we have explicit formula

$$\begin{aligned} \phi^\epsilon &= \exp(i \int_\gamma v^\epsilon(y) \cdot d\vec{l}(y)), \\ \nabla \phi^\epsilon &= i v^\epsilon \phi^\epsilon. \end{aligned}$$

Note that $\sqrt{\rho}$ is continuous and U is strictly away from the vacuum, therefore $\inf_{x \in U} \{\sqrt{\rho}(x)\} = \alpha > 0$ and the velocity field $v = \Lambda/\sqrt{\rho} \in L^2(U)$. By the theory of mollifier we have $v^\epsilon \rightarrow v$ in $L^2(U)$, and combining it with the weak* limit of ϕ^ϵ we obtain

$$(3.7) \quad \nabla \phi^\epsilon = i v^\epsilon \phi^\epsilon \rightharpoonup i v \phi = \nabla \phi \quad \text{in } L^2(U).$$

Thus we obtain

$$\begin{aligned} \nabla \psi &= (\nabla \sqrt{\rho} + i \sqrt{\rho} v) \phi \\ &= (\nabla \sqrt{\rho} + i \Lambda) \phi \end{aligned}$$

with uniform L^2 bound

$$\|\nabla \psi\|_{L^2(U)} = \|\nabla \sqrt{\rho}\|_{L^2(U)} + \|\Lambda\|_{L^2(U)} \leq M_1.$$

Extending the open set U to $B_{|h|}(0)^c$, we get

$$\|\nabla \psi\|_{L^2(B_h(0)^c)} \leq M_1,$$

and we conclude $\psi \in H^1(\mathbf{R}^2)$. Furthermore, the above argument implies for a.e. $x \in \mathbf{R}^2$

$$(3.8) \quad \nabla \psi = (\nabla \sqrt{\rho} + i \Lambda) \phi.$$

If further assume condition (3.6), we are going to prove $\psi \in H^2(\mathbf{R}^2)$. The proof follows similar idea as before using the theorem of difference quotient, but here we need a small technique since we don't have a L^∞ bound on $\nabla \psi$.

First we also take arbitrary open set U such that \overline{U} is compact and strictly away from the vacuum. By (3.7) and (3.8) we can directly compute that for $j = 1, 2$,

$$\begin{aligned} \partial_{x_j}^2 \psi &= (\partial_{x_j}^2 \sqrt{\rho} + i \partial_{x_j} \Lambda_j + i \partial_{x_j} \sqrt{\rho} u_j - \Lambda_j u_j) \phi \\ &= [(\partial_{x_j}^2 \sqrt{\rho} - \frac{\Lambda_j^2}{\sqrt{\rho}}) + i \frac{\partial_{x_j} J_j}{\sqrt{\rho}}] \phi. \end{aligned}$$

Therefore condition 3.6 implies $\partial_{x_j}^2 \psi \in L^2(U)$ with uniform bound

$$\|\partial_{x_j}^2 \psi\|_{L^2(U)} \leq M_2.$$

On the other hand, for $\partial_{x_j}\psi$, $j = 1, 2$ we can decompose it via polar factor into the form $\partial_{x_j}\psi = |\partial_{x_j}\psi|\phi_j$, where the polar factor $\phi_j = \partial_{x_j}\psi/|\partial_{x_j}\psi|$ when $|\partial_{x_j}\psi| \neq 0$. In this case direct computation shows

$$(3.9) \quad \partial_{x_j}\phi_j = i \frac{\operatorname{Im}(\partial_{x_j}\bar{\psi}\partial_{x_j}^2\psi)}{|\partial_{x_j}\psi|^2}\phi_j,$$

and by the same argument as polar factorization we have

$$(3.10) \quad |\partial_{x_j}^2\psi|^2 = (\partial_{x_j}|\partial_{x_j}\psi|)^2 + \left(\frac{\operatorname{Im}(\partial_{x_j}\bar{\psi}\partial_{x_j}^2\psi)}{|\partial_{x_j}\psi|} \right)^2.$$

Then instead of $\partial_{x_j}\psi$, we consider a sequence of functions $\{f_{j,n}\}$ such that

$$f_{j,n} = \min\{|\partial_{x_j}\psi|, n\}\phi_j, \quad j = 1, 2.$$

By definition $f_{j,n}$ is a L^∞ cut-off of $\partial_{x_j}\psi$ with L^∞ bound n , and this cut-off process is legal since it occurs only when $|\partial_{x_j}\psi| \geq n$. Now we consider the difference quotient of $f_{j,n}$ in the direction e_j , where

$$e_1 = (1, 0) \quad \text{and} \quad e_2 = (0, 1),$$

and we let $h_j = |h|e_j$ for $|h| \neq 0$. We have

$$\|D^{h_j}f_n\|_{L^2(\mathbf{R}^2)} = \|D^{h_j}f_{j,n}\|_{L^2(B_{|h|}(0))} + \|D^{h_j}f_{j,n}\|_{L^2(B_{|h|}(0)^c)}$$

for $h \neq 0$ small, and the part near the vacuum can be controlled by

$$\|D^{h_j}f_{j,n}\|_{L^2(B_{|h|}(0))} \leq 2\sqrt{\pi}\|f_{j,n}\|_{L^\infty} \leq 2\sqrt{\pi}n.$$

To deal with the part away from the vacuum, we again approximate it by open sets $U \subset B_{|h|}(0)^c$ with \bar{U} compact. By definition $|f_{j,n}| \leq |\partial_{x_j}\psi|$, and since $\partial_{x_j}^2\psi \in L^2(U)$, the x_j -derivative of the truncated function $\partial_{x_j}f_{j,n}$ is also in $L^2(U)$. Moreover by using (3.9) and (3.10) we compute (for $|\partial_j\psi| \geq n$)

$$\begin{aligned} |\partial_{x_j}f_{j,n}|^2 &= (\partial_{x_j}|f_{j,n}|)^2 + \left(f_{j,n} \frac{\operatorname{Im}(\partial_{x_j}\bar{\psi}\partial_{x_j}^2\psi)}{|\partial_{x_j}\psi|^2} \right)^2 \\ &\leq (\partial_{x_j}|\partial_j\psi|)^2 + \left(\frac{\operatorname{Im}(\partial_{x_j}\bar{\psi}\partial_{x_j}^2\psi)}{|\partial_{x_j}\psi|} \right)^2 = |\partial_{x_j}^2\psi|^2 \end{aligned}$$

Therefore we have uniform bound

$$\|\partial_{x_j}f_{j,n}\|_{L^2(U)} \leq \|\partial_{x_j}^2\psi\|_{L^2(U)} \leq M_2,$$

for any $U \subset B_{|h|}(0)^c$, hence it also works on $B_{|h|}(0)^c$, therefore we conclude $\partial_{x_j}f_{j,n} \in L^2(\mathbf{R}^2)$.

Note that even though the control of $\|D^{h_j}f_n\|_{L^2}$ obtained above depends on n , as long as we proved $\partial_{x_j}f_{j,n} \in L^2(\mathbf{R}^2)$, a zero measure set has no effect on the L^2 norm of $\partial_{x_j}f_{j,n}$. Thus by extending the open set U to $\mathbf{R}^2 - \{0\}$ we obtain that

$$\|\partial_{x_j}f_{j,n}\|_{L^2(\mathbf{R}^2)} \leq M_2, \quad j = 1, 2,$$

which shows $\{\partial_{x_j}f_{j,n}\}$ is a uniformly bounded sequence in $L^2(\mathbf{R}^2)$. Passing to the limit, $\partial_{x_j}f_{j,n}$ converges (upto a subsequence) to a limit in $L^2(\mathbf{R})$ weakly. On the

other hand, it is straightforward to see that $f_{j,n}$ converges to $\partial_{x_j}\psi$ strongly by construction. Therefore we conclude $\partial_{x_j}^2\psi \in L^2(\mathbf{R}^2)$ for $j = 1, 2$, with bound

$$\|\partial_{x_j}^2\psi\|_{L^2(\mathbf{R}^2)} \leq M_2.$$

It implies $\Delta\psi \in L^2(\mathbf{R}^2)$ with control

$$\|\Delta\psi\|_{L^2(\mathbf{R}^2)} \leq 2M_2,$$

then by the standard argument we obtain $\psi \in H^2(\mathbf{R}^2)$ with bound

$$\|\psi\|_{H^2(\mathbf{R}^2)} \leq C(M_1, M_2).$$

□

The next Proposition is a simplified analogue of Proposition 21 which will be used in later contents. We consider hydrodynamic data (ρ, J) such that ρ is continuous with $\rho > 0$, and let us assume the data is irrotational in the sense of Definition 15 i.e.

$$(3.11) \quad \nabla \wedge J = 2\nabla\sqrt{\rho} \wedge \Lambda.$$

By Remark 16 we see that when the density ρ is positive, identity (3.11) is equivalent to the classical irrotationality condition $\nabla \wedge v = 0$. For such hydrodynamic data with regularity assumption (3.3) and (3.6), we can lift it to a wave function $\psi \in H^2(\mathbf{R}^d)$ following the same argument as Proposition 21. Moreover the Proposition also applies to general multi-dimensional data.

Proposition 22. *Let $(\sqrt{\rho}, \Lambda)$ be a pair of finite energy hydrodynamic data on \mathbf{R}^d satisfying bounds*

$$(3.12) \quad \|\sqrt{\rho}\|_{H^1(\mathbf{R}^d)} + \|\Lambda\|_{L^2(\mathbf{R}^d)} \leq M_1.$$

and

$$(3.13) \quad \left\| \frac{\Lambda^2}{\sqrt{\rho}} - \Delta\sqrt{\rho} \right\|_{L^2(\mathbf{R}^d)} + \left\| \frac{\operatorname{div} J}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}^d)} \leq M_2,$$

Let us further assume that $\sqrt{\rho}$ is continuous with $\rho > 0$ and the (ρ, J) satisfies the generalised irrotationality condition (3.11). Then there exists a wave function $\psi \in H^2(\mathbf{R}^d)$ such that

$$\sqrt{\rho} = |\psi|, \quad \Lambda = \operatorname{Im}(\bar{\phi}\nabla\psi),$$

where $\phi \in P(\psi)$ is a polar factor, and we have controls

$$\|\psi\|_{H^2(\mathbf{R}^d)} \leq M_2 + C(M_1).$$

Proof. The proof is an analogue of the proof to Proposition 21, but the argument becomes simpler due to the absence of vacuum.

Since $\sqrt{\rho}$ is continuous in space and positive, it has a positive lower bound on any compact set in \mathbf{R}^d . Therefore the velocity field $v = \Lambda/\sqrt{\rho}$ belongs to $L_{loc}^2(\mathbf{R}^d)$. Let $\{\chi_\epsilon\}$, $\epsilon > 0$ be a sequence of smooth mollifier. We define approximating velocity

$$v^\epsilon = v * \chi_\epsilon,$$

and define the approximating polar factor

$$\phi^\epsilon(x) = \exp(i \int_{\gamma(x)} v^\epsilon(y) \cdot d\vec{l}(y)),$$

where $\gamma(x)$ is the straight line in \mathbf{R}^d connecting 0 and x . By Remark 16, the identity (3.11) is equivalent to the classical irrotationality condition

$$\nabla \wedge v = 0$$

when then density ρ is always positive. Hence the approximating velocity field v^ϵ is also irrotational, and the polar factor ϕ^ϵ is well-defined. Moreover it is straightforward to compute that

$$\nabla \phi^\epsilon = i v^\epsilon \phi^\epsilon.$$

The polar factors $\{\phi^\epsilon\}$ have uniform L^∞ bound 1, therefore we can take a subsequence which has a weak* limit ϕ with $\|\phi\|_L^\infty \leq 1$. Without changing the notation we assume $\phi^\epsilon \xrightarrow{*} \phi$ in $L^\infty(\mathbf{R}^d)$. On the other hand, by the basic property of mollifier, $v^\epsilon \rightarrow v$ in $L_{loc}^2(\mathbf{R}^d)$ as $\epsilon \rightarrow 0$. Thus we get

$$\nabla \phi^\epsilon = i v^\epsilon \phi^\epsilon \rightharpoonup i v \phi \quad \text{in } L_{loc}^2(\mathbf{R}^d),$$

and by the weak derivative formulation we conclude

$$(3.14) \quad \nabla \phi = i v \phi \quad a.e.$$

We define $\psi = \sqrt{\rho} \phi$, then we can directly check that ψ is the wave function associated to $(\sqrt{\rho}, \Lambda)$ and $\psi \in H^2(\mathbf{R}^2)$. First we have $|\psi| = \sqrt{\rho}$. By identity (3.14) we can compute

$$(3.15) \quad \nabla \psi = (\sqrt{\rho} + i \sqrt{\rho} v) \phi = (\sqrt{\rho} + i \Lambda) \phi,$$

hence $\Lambda = \text{Im}(\bar{\phi} \nabla \psi)$ and

$$\|\psi\|_{H^1(\mathbf{R}^2)} \leq \|\sqrt{\rho}\|_{H^1(\mathbf{R}^2)} + \|\Lambda\|_{L^2(\mathbf{R}^2)} \leq M_1.$$

Moreover by (3.14) and (3.15) we can directly compute

$$\begin{aligned} \Delta \psi &= \text{div } \nabla \psi \\ &= (\Delta \sqrt{\rho} + i \text{div } \Lambda + i \nabla \sqrt{\rho} \cdot u - \Lambda \cdot u) \phi \\ &= \left[(\Delta \sqrt{\rho} - \frac{|\Lambda|^2}{\sqrt{\rho}}) + i \frac{\text{div } J}{\sqrt{\rho}} \right] \phi. \end{aligned}$$

Using (3.12) we obtain that

$$\|\Delta \psi\|_{L^2(\mathbf{R}^2)} \leq \left\| \frac{\Lambda^2}{\sqrt{\rho}} - \Delta \sqrt{\rho} \right\|_{L^2(\mathbf{R}^2)} + \left\| \frac{\text{div } J}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}^2)} \leq M_2,$$

which conclude the proof. \square

To conclude this section we prove Theorem 1 and Theorem 2, i.e. the global well-posedness Theorem for the Cauchy of 2D QHD system (1.1) with initial data satisfying the point vacuum and quantised vorticity condition, by combining the Proposition 25, the global well-posedness of NLS equations (Theorem 12) and the polar factorization method (Theorem 17).

Proof of Theorem 1 and Theorem 2. For given initial data $(\sqrt{\rho_0}, \Lambda_0)$ satisfying (1.10) and the condition (1.7) and (1.8), by Proposition 21 we know there exists a associated wave function $\psi_0 \in H^1$ associated to $(\sqrt{\rho_0}, \Lambda_0)$. Then using Theorem 12 and Theorem 17 we obtain a global in time finite energy weak solution $(\sqrt{\rho}, \Lambda)$ to (1.1) which conserves the energy.

If $(\sqrt{\rho_0}, \Lambda_0)$ satisfy both (1.10) and (1.12) then Proposition 21 shows the wave function ψ_0 is actually $\psi_0 \in H^2(\mathbf{R}^2)$. By the Theorem 12 for the NLS, we have $\psi \in \mathcal{C}([0, T]; H^2(\mathbf{R}^2)) \cap \mathcal{C}^1([0, T]; L^2(\mathbf{R}^2))$ with

$$(3.16) \quad \|\psi\|_{L^\infty(0, T; H^2(\mathbf{R}^2))} + \|\partial_t \psi\|_{L^\infty(0, T; L^2(\mathbf{R}^2))} \leq C(T, M_1, M_2).$$

holds for any $0 < T < \infty$. Now, let us recall that $\partial_t \sqrt{\rho} = \text{Re}(\bar{\phi} \partial_t \psi)$ and $\lambda = -\text{Im}(\bar{\phi} \partial_t \psi)$, with $\phi \in P(\psi)$, then by the polar factorization and (3.16) we also infer

$$\|\partial_t \sqrt{\rho}\|_{L_t^\infty L_x^2} + \|\lambda\|_{L_t^\infty L_x^2} \leq C(T, M_1, M_2).$$

Moreover using the NLS equation (2.1) we can directly compute that

$$\sqrt{\rho} \lambda = -\text{Im}(\bar{\psi} \partial_t \psi) = -\frac{1}{4} \Delta \rho + e + p(\rho)$$

so that (1.20) holds. To conclude the proof of Theorem 2 it only remains to prove the higher bounds. We already know that, by the conservation of energy, we have

$$\|\sqrt{\rho}\|_{L^\infty(\mathbf{R}; H^1(\mathbf{R}^2))} + \|\Lambda\|_{L^\infty(\mathbf{R}; L^2(\mathbf{R}^2))} \leq M_1.$$

Let $0 < T < \infty$ be fixed, by using $\Delta \rho = 2 \text{Re}(\bar{\psi} \Delta \psi) + 2|\nabla \psi|^2$, Hölder's inequality and Sobolev embedding we have

$$\|\Delta \rho\|_{L^\infty(0, T; L^2(\mathbf{R}^2))} \lesssim \|\psi\|_{L^\infty(0, T; H^2(\mathbf{R}^2))}^2 \leq C(T, M_1, M_2).$$

On the other hand, by the Madelung transformation

$$\|\nabla J\|_{L_t^\infty L_x^2} = \|\nabla \text{Im}(\bar{\psi} \nabla \psi)\|_{L_t^\infty L_x^2} \leq \|\psi\|_{L_t^\infty H_x^2}^2 \leq C(T, M_1, M_2).$$

Finally, let us recall that by the polar factorization we have

$$e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho) = \frac{1}{2} |\nabla \psi|^2 + f(|\psi|^2).$$

It is straightforward to see that $\nabla |\nabla \psi| \leq |\nabla^2 \psi|$ a.e. $x \in \mathbf{R}^2$, so that

$$\|\nabla \sqrt{e}\|_{L_t^\infty L_x^2} \leq \|\psi\|_{L_t^\infty H_x^2} \leq C(T, M_1, M_2).$$

□

3.1. Example of 3D hydrodynamic data: straight vortex lines. Here we extend wave function lifting argument to a simple example of 3D hydrodynamic data, where we assume planar symmetry to the data and the vortices appear as finite number of parallel straight lines, which is a typical example of vortex structure of quantum flows in 3D space. More precisely we consider hydrodynamic data $(\rho(x), J(x))$, $x = (x_1, x_2, x_3) \in \mathbf{R}^3$ and $J \in \mathbf{R}^3$, which has the following form:

$$(3.17) \quad \rho(x) = \rho_1(x_1, x_2) \rho_2(x_3) \quad \text{and} \quad J(x) = (J_1(x_1, x_2) \rho_2(x_3), 0) = \sqrt{\rho} \Lambda,$$

where $\rho, \rho_j \in \mathbf{R}$ and $J_1 \in \mathbf{R}^2$. The data (ρ, J) is essentially a product of a 2D data (ρ_1, J_1) and a 1D data $(\rho_2, 0)$. We assume $(\rho_1, J_1 = \sqrt{\rho_1} \Lambda_1)$ has pointwise vacuum and quantised vorticity, i.e. we assume ρ_1 to be continuous with vacuum structure (3.4), and assume the velocity field $v_1 = J_1/\rho_1$ to satisfies the quantised vorticity condition (3.5). The regularity of (ρ_1, J_1) is characterised by the bounds (3.3) and (3.6), and let us assume $\sqrt{\rho_2} \in H^s(\mathbf{R})$ with $s = 1$ or 2 .

With the assumption above, we can state the following Proposition as an extension of the wave function lifting argument to 3D data. Moreover the the global existence Theorem 4 and Theorem 5 for initial data of the form (3.17) can be proved in the same strategy as before by using Proposition 23.

Proposition 23. *Let (ρ, J) be a pair of hydrodynamic data as in (3.17), with (ρ_1, J_1) satisfying the condition (3.4) and (3.5). Furthermore we assume that (ρ_1, J_1) satisfy the mass and energy bounds (3.3) and $\sqrt{\rho_2} \in H^1(\mathbf{R})$. Then there exists a wave function $\psi \in H^1(\mathbf{R}^3)$ such that*

$$\sqrt{\rho} = |\psi|, \quad \Lambda = \text{Im}(\bar{\phi} \nabla \psi),$$

where $\phi \in P(\psi)$ is a polar factor defined in (2.4).

If we furthermore assume the controls (3.6) on (ρ_1, J_1) and $\sqrt{\rho_2} \in H^2(\mathbf{R})$, then $\psi \in H^2(\mathbf{R}^3)$ and we have

$$\|\psi\|_{H^2(\mathbf{R}^3)} \leq C(M_1, M_2).$$

Proof. By the assumption we can apply Proposition 21 to (ρ_1, J_1) to lift it to a wave function $\psi_1 \in H^1(\mathbf{R}^2)$ such that

$$\sqrt{\rho_1} = |\psi_1|, \quad \Lambda_1 = \text{Im}(\bar{\phi}_1 \nabla \psi_1),$$

where ϕ_1 is a polar factor of ψ_1 . Then we claim

$$\psi(x) = \psi_1(x_1, x_2) \sqrt{\rho_2(x_3)}$$

is the wave function we need. Since $\psi_1 \in H^1(\mathbf{R}^2)$ and $\sqrt{\rho_2} \in H^1(\mathbf{R})$ we have $\psi \in H^1(\mathbf{R}^3)$. Furthermore

$$|\psi| = |\psi_1| \sqrt{\rho_2} = \sqrt{\rho_1 \rho_2} = \sqrt{\rho}.$$

Obviously $\phi(x) = \phi_1(x_1, x_2)$ is a polar factor of ψ , then for $j = 1, 2$

$$\text{Im}(\bar{\phi} \partial_{x_j} \psi) = \text{Im}(\bar{\phi}_1 \partial_{x_j} \psi_1) \sqrt{\rho_2} = \Lambda_{1,j} \sqrt{\rho_2} = \Lambda_j,$$

and

$$\text{Im}(\bar{\phi} \partial_{x_3} \psi) = \text{Im}(\bar{\phi}_1 \psi_1) \partial_{x_3} \sqrt{\rho_2} = 0.$$

Hence we conclude

$$\text{Im}(\bar{\phi} \nabla \psi) = \Lambda,$$

which finish the proof of the first part.

If we further assume (ρ_1, J_1) to satisfy (3.6), which implies $\psi_1 \in H^2(\mathbf{R}^2)$ by Proposition 21, and assume $\sqrt{\rho_2} \in H^2(\mathbf{R})$, then it is direct to obtain $\psi = \psi_1 \sqrt{\rho_2} \in H^2(\mathbf{R}^3)$ with

$$\|\psi\|_{H^2(\mathbf{R}^3)} \leq \|\psi_1\|_{H^2(\mathbf{R}^2)} \|\sqrt{\rho_2}\|_{H^2(\mathbf{R})} \leq C(M_1, M_2).$$

□

4. WAVE FUNCTION LIFTING: SPHERICAL SYMMETRIC DATA

This Section concerns the wave function lifting method for spherically symmetric data, which is essentially an analogue of the method we applied in [6] to one-dimensional data. Then we prove the global in time existence of spherically symmetric solutions to system (1.1).

4.1. Spherically symmetric QHD system. We consider spherically symmetric solutions to the QHD system (1.1) of the following form:

$$(4.1) \quad \sqrt{\rho}(t, x) = \sqrt{\rho}(t, r) \text{ and } \Lambda(t, x) = \Lambda(t, r) \frac{x}{|x|},$$

for $x \in \mathbf{R}^d$ with $r = |x|$, and $\Lambda(t, r) \in \mathbf{R}$. In our works we mainly focus on the case $d = 2$ or 3 , but here we give an argument that applies to general dimension d . Now we need to transform the equations (1.1) into scalar equations for $(\sqrt{\rho}(t, r), \Lambda(t, r))$.

With a slight abuse of notations, we denote $f(x) = f(r)$ for scalar functions and $F(x) = F(r) \frac{x}{|x|}$ for vector fields. In this way, the action of gradient, divergence and Laplace operator on f and F are the following

$$(4.2) \quad \nabla_x f = \partial_r f \frac{x}{|x|}, \quad \operatorname{div}_x(F) = \partial_r F + \frac{d-1}{r} F, \quad \Delta_x f = \partial_r^2 f + \frac{d-1}{r} \partial_r f.$$

By using (4.2) and coordinate change, it is straightforward to obtain the formulation of QHD system (1.1) in the spherically symmetric case:

Proposition 24. *For spherically symmetric solutions of the form*

$$\sqrt{\rho}(t, x) = \sqrt{\rho}(t, r) \text{ and } \Lambda(t, x) = \Lambda(t, r) \frac{x}{|x|},$$

the QHD system (1.1) is equivalent to the system of scalars $(\sqrt{\rho}, \Lambda)(t, r)$ given by

$$(4.3) \quad \begin{cases} \partial_t \rho + (\partial_r + \frac{d-1}{r})(\sqrt{\rho} \Lambda) = 0 \\ \partial_t(\sqrt{\rho} \Lambda) + (\partial_r + \frac{d-1}{r})(\Lambda^2 + (\partial_r \sqrt{\rho})^2) + \partial_r p(\rho) = \frac{1}{4} \partial_r (\partial_r + \frac{d-1}{r}) \partial_r \rho. \end{cases}$$

for $(t, r) \in [0, \infty)^2$. The total mass and energy of (4.3) are defined as

$$(4.4) \quad M(t) = C(d) \int_0^\infty r^{d-1} \rho \, dr,$$

and

$$(4.5) \quad E(t) = C(d) \int_0^t r^{d-1} \left[\frac{1}{2} (|\partial_r \sqrt{\rho}|^2 + |\Lambda|^2) + f(\rho) \right] dr,$$

where $C(d)$ is the area of $(d-1)$ -dimensional unit sphere, and we denote the total energy density by

$$(4.6) \quad e = \frac{1}{2} (\partial_r \sqrt{\rho})^2 + \frac{1}{2} \Lambda^2 + f(\rho).$$

Furthermore for spherically symmetric solutions, the higher order functional $I(t)$, introduced by (1.6), has the form

$$(4.7) \quad I(t) = C(d) \int_0^\infty r^{d-1} [\lambda^2 + (\partial_t \sqrt{\rho})^2] \, dr,$$

where

$$(4.8) \quad \sqrt{\rho} \lambda = -\frac{1}{4} (\partial_r^2 \rho + \frac{d-1}{r} \partial_r \rho) + e + p(\rho).$$

Last we define the Lebesgue space and Sobolev space of functions $f(r)$, $r \in \mathbf{R}^+$, to be

$$\|f\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} = \left(\int_0^\infty f(r)^2 r^{d-1} dr \right)^{\frac{1}{2}},$$

and

$$\|f\|_{H^1(\mathbf{R}_+, r^{d-1}dr)} = \|f\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} + \|\partial_r f\|_{L^2(\mathbf{R}_+, r^{d-1}dr)}.$$

4.2. Wave function lifting for spherically symmetric data. In this section we will establish the wave function lifting method for multi-dimensional spherically symmetric hydrodynamic data. More precisely, given hydrodynamic data $(\sqrt{\rho}(r), \Lambda(r))$ on $\mathbf{R}_+ = [0, \infty)$, with finite mass $M(\rho)$, energy $E(\rho, J)$ and $\Lambda = 0$ a.e. on the vacuum region $\{\rho = 0\}$, we will construct a spherically symmetric wave function $\psi \in H^1(\mathbf{R}^d)$ associated to $(\sqrt{\rho}, \Lambda)$. If we further assume suitable higher order regularity on $(\sqrt{\rho}, \Lambda)$, then the wave function ψ can be chosen in $H^2(\mathbf{R}^d)$.

Proposition 25. *Let $(\sqrt{\rho}, \Lambda)$ be a spherically symmetric hydrodynamic data on \mathbf{R}^d , i.e.*

$$\sqrt{\rho(x)} = \sqrt{\rho(r)} \text{ and } \Lambda(x) = \Lambda(r) \frac{x}{|x|},$$

where $r = |x|$ and $\Lambda(r) \in \mathbf{R}$ such that $\Lambda = 0$ a.e. on the set $\{\rho = 0\}$. Let us further assume the bounds

$$(4.9) \quad \|\sqrt{\rho}\|_{H^1(\mathbf{R}_+, r^{d-1}dr)} + \|\Lambda\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} \leq M_1,$$

Then there exists a spherically symmetric wave function $\psi \in H^1(\mathbf{R}^d)$ such that

$$\sqrt{\rho} = |\psi|, \quad \Lambda = \text{Im}(\bar{\phi} \partial_r \psi),$$

where $\phi \in P(\psi)$ is a polar factor of ψ .

If we further assume that $(\sqrt{\rho}, \Lambda)$ satisfy also the bounds

$$(4.10) \quad \begin{aligned} \|\partial_r^2 \sqrt{\rho} + (d-1) \frac{\partial_r \sqrt{\rho}}{r}\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} + \left\| \frac{\Lambda^2}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} \\ + \left\| \frac{\partial_r J + (d-1)J/r}{\sqrt{\rho}} \right\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} \leq M_2. \end{aligned}$$

then $\psi \in H^2(\mathbf{R}^d)$ and we have

$$(4.11) \quad \|\psi\|_{H^2(\mathbf{R}^d)} \leq C(M_1, M_2).$$

Proof. Let us consider a sequence $\{\delta_n\}$ of Schwartz functions such that $\delta_n(r) > 0$ for all $r \in \mathbf{R}^+$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$. For instance we may consider $\delta_n(r) = \frac{1}{n} e^{-r^2/2}$. We define the following hydrodynamical quantities

$$\sqrt{\rho_n} = \sqrt{\rho} + \delta_n, \quad \Lambda_n = \frac{J}{\sqrt{\rho_n}} = \frac{\sqrt{\rho}}{\sqrt{\rho_n}} \Lambda,$$

then we can check that also $(\sqrt{\rho_n}, \Lambda_n)$ satisfy (4.9) uniformly in $n \in \mathbf{N}$. Indeed,

$$\|\sqrt{\rho_n}\|_{H^1(\mathbf{R}_+, r^{d-1}dr)} \leq \|\sqrt{\rho}\|_{H^1(\mathbf{R}_+, r^{d-1}dr)} + \|\delta_n\|_{H^1(\mathbf{R}_+, r^{d-1}dr)} \leq C$$

and since $\sqrt{\rho_n}(r) \geq \sqrt{\rho}(r)$ for every $r \in \mathbf{R}^+$, we have $|\Lambda_n(r)| \leq |\Lambda(r)|$ a.e. and consequently

$$\|\Lambda_n\|_{L^2(\mathbf{R}_+, r^{d-1}dr)} \leq \|\Lambda\|_{L^2(\mathbf{R}_+, r^{d-1}dr)}.$$

By construction we straightforwardly have that $\sqrt{\rho_n} \rightarrow \sqrt{\rho}$ in $H^1(\mathbf{R}^+, r^{d-1}dr)$. Moreover, by assumption we have $\Lambda(r) = 0$ for a.e. $r \in \{\rho = 0\}$ and by construction the same holds also for Λ_n . Hence the pointwise and monotonic convergence $\sqrt{\rho_n}(r) \rightarrow \sqrt{\rho}(r)$ also implies

$$\Lambda_n(r) \rightarrow \Lambda(r), \quad \text{a.e. } r \in \mathbf{R}^+,$$

and $|\Lambda_n|$ converges to $|\Lambda|$ monotonically. Then dominated convergence theorem yields $\Lambda_n \rightarrow \Lambda$ in $L^2(\mathbf{R}^+, r^{d-1}dr)$. Furthermore, since $\sqrt{\rho_n}(r) > 0$ it is possible to define the velocity field

$$v_n = \frac{\Lambda_n}{\sqrt{\rho_n}}.$$

Notice that since $\sqrt{\rho_n}$ is uniformly bounded away from zero on compact intervals we have $v_n \in L^1_{loc}(\mathbf{R}^+)$, hence it makes sense to define the phase function for $r > 0$ by

$$S_n(r) = \int_{r_0}^r v_n(s) ds,$$

where $r_0 > 0$ is a given point, and consequently the radially symmetric wave function

$$\psi_n(r) = \sqrt{\rho_n}(r) e^{iS_n(r)}.$$

We can now show that the sequence ψ_n has a limit $\psi \in H^1(\mathbf{R}^+, r^{d-1}dr)$ which satisfies $|\psi| = \sqrt{\rho}$, $\text{Im}(\bar{\phi} \partial_r \psi) = \Lambda$. Indeed, since $\partial_r \psi_n = e^{iS_n}(\partial_r \sqrt{\rho_n} + i\Lambda_n)$, we have

$$\|\psi_n\|_{H^1(\mathbf{R}^+, r^{d-1}dr)}^2 = \|\sqrt{\rho_n}\|_{H^1(\mathbf{R}^+, r^{d-1}dr)}^2 + \|\Lambda_n\|_{L^2(\mathbf{R}^+, r^{d-1}dr)}^2 \leq C,$$

thus, up to subsequences, $\psi_n \rightharpoonup \psi$ in $H^1(\mathbf{R}^+, r^{d-1}dr)$. On the other hand, we also have $\sqrt{\rho_n} \rightarrow \sqrt{\rho}$ in $H^1(\mathbf{R}^+, r^{d-1}dr)$, $\Lambda_n \rightarrow \Lambda$ in $L^2(\mathbf{R}^+, r^{d-1}dr)$ and moreover $e^{iS_n} \rightharpoonup \phi$ weakly* in $L^\infty(\mathbf{R}^+)$, for some $\phi \in L^\infty(\mathbf{R}^+)$. It is straightforward to check that ϕ is a polar factor for ψ , since $\psi_n = \sqrt{\rho_n} e^{iS_n} \rightharpoonup \sqrt{\rho} \phi$. Furthermore

$$\partial_r \psi_n = e^{iS_n}(\partial_r \sqrt{\rho_n} + i\Lambda_n) \rightharpoonup \phi(\partial_r \sqrt{\rho} + i\Lambda),$$

so that $\partial_r \psi = \phi(\partial_r \sqrt{\rho} + i\Lambda)$ and hence $(\sqrt{\rho}, \Lambda)$ are the hydrodynamical quantities associated to ψ .

We will now prove the second part of the Proposition, so let us assume that $(\sqrt{\rho}, \Lambda)$ further satisfy the bounds (4.10). We are going to prove that the sequence ψ_n constructed above satisfies that $(\partial_r + \frac{d-1}{r})\partial_r \psi_n$ is uniformly bounded in $L^2(\mathbf{R}^+, r^{d-1}dr)$, namely

$$\|(\partial_r + \frac{d-1}{r})\partial_r \psi_n\|_{L^2(\mathbf{R}^+, r^{d-1}dr)} \leq C(M_2), \quad \forall n \in \mathbf{N},$$

which by (4.2) is equivalent to $\|\Delta \psi\|_{L^2(\mathbf{R}^d)} \leq C(M_2)$. Using the formula of $\partial_r \psi_n$ we compute

$$\begin{aligned} (\partial_r + \frac{d-1}{r})\partial_r \psi_n &= (\partial_r + \frac{d-1}{r})[(\partial_r \sqrt{\rho_n} + i\Lambda_n)e^{iS_n}] \\ &= (\partial_r^2 \sqrt{\rho_n} + i\partial_r \sqrt{\rho_n} v_n + i\partial_r \Lambda_n \\ &\quad - v_n \Lambda_n + \frac{d-1}{r}\partial_r \sqrt{\rho_n} + \frac{d-1}{r}\Lambda_n) e^{iS_n} \\ &= \left[(\partial_r^2 \sqrt{\rho_n} + (d-1)\frac{\partial_r \sqrt{\rho_n}}{r}) + i\frac{\partial_r J + (d-1)J/r}{\sqrt{\rho_n}} - \frac{\Lambda_n^2}{\sqrt{\rho_n}} \right] e^{iS_n}. \end{aligned}$$

Because of the definition of $\sqrt{\rho_n}$, we also have

$$\frac{|\partial_r J + (d-1)J/r|}{\sqrt{\rho_n}} \leq \frac{|\partial_r J + (d-1)J/r|}{\sqrt{\rho}}, \quad \frac{\Lambda_n^2}{\sqrt{\rho_n}} \leq \frac{\Lambda^2}{\sqrt{\rho}}, \quad a.e.$$

so that

$$\|(\partial_r + \frac{d-1}{r})\partial_r \psi_n\|_{L^2(\mathbf{R}^+, r^{d-1}dr)} \leq C(M_2).$$

Since the bound is uniform in $n \in \mathbf{N}$, by lower semicontinuity it also holds for $(\partial_r + \frac{d-1}{r})\partial_r \psi \in L^2(\mathbf{R}^+, r^{d-1}dr)$. \square

As a conclusion of this section, we can prove the global existence of spherically symmetric weak solutions to (1.1). As Theorem 1 and Theorem 2, the well-posedness result is a direct consequence of the wave function lifting Proposition 21, Theorem 12 and polar factorization method 17.

Proof of Theorem 6 and Theorem 7. For a given spherically symmetric initial data $(\sqrt{\rho_0}, \Lambda_0)$ satisfying (1.18) and such that $\Lambda_0 = 0$ a.e. on $\{\sqrt{\rho_0} = 0\}$, by Proposition 25 we know there exists a spherically symmetric $\psi_0 \in H^1$ associated to $(\sqrt{\rho_0}, \Lambda_0)$. Then using Theorem 12 and Theorem 17 we obtain a global in time finite energy weak solution $(\sqrt{\rho}, \Lambda)$ to (1.1) which conserves the energy. Furthermore since the solution is obtained by solving the associated NLS equation with initial data ψ_0 and spherical symmetry is preserved by NLS equation, we know that the solution $(\sqrt{\rho}, \Lambda)$ is also spherical symmetric

If we further assume $(\sqrt{\rho_0}, \Lambda_0)$ satisfy the bounds (1.19) as in Theorem 7, which is equivalent to (4.10) by the discussion in the last Section, then Proposition 25 shows the wave function ψ_0 is actually $\psi_0 \in H^2(\mathbf{R}^d)$. By the persistence of regularity for the NLS, we have $\psi \in \mathcal{C}([0, T]; H^2(\mathbf{R}^d)) \cap \mathcal{C}^1([0, T]; L^2(\mathbf{R}^d))$ with

$$(4.12) \quad \|\psi\|_{L^\infty(0, T; H^2(\mathbf{R}^d))} + \|\partial_t \psi\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} \leq C(T, M_1, M_2).$$

holds for any $0 < T < \infty$. Now, let us recall that $\partial_t \sqrt{\rho} = \text{Re}(\bar{\phi} \partial_t \psi)$ and $\lambda = -\text{Im}(\bar{\phi} \partial_t \psi)$, with $\phi \in P(\psi)$, then by the polar factorization and (4.12) we also infer

$$\|\partial_t \sqrt{\rho}\|_{L_t^\infty L_x^2} + \|\lambda\|_{L_t^\infty L_x^2} \leq C(T, M_1, M_2).$$

Moreover using the NLS equation (2.1) we can directly compute that

$$\sqrt{\rho} \lambda = -\frac{1}{4} \Delta \rho + e + p(\rho),$$

so that (1.20) holds. To conclude the proof of Theorem 6 it only remains to prove the higher bounds. We already know that, by the conservation of energy, we have

$$\|\sqrt{\rho}\|_{L^\infty(\mathbf{R}; H^1(\mathbf{R}^d))} + \|\Lambda\|_{L^\infty(\mathbf{R}; L^2(\mathbf{R}^d))} \leq M_1.$$

Let $0 < T < \infty$ be fixed, by using $\Delta \rho = 2 \text{Re}(\bar{\psi} \Delta \psi) + 2|\nabla \psi|^2$, Hölder's inequality and Sobolev embedding we have

$$\|\Delta \rho\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} \lesssim \|\psi\|_{L^\infty(0, T; H^2(\mathbf{R}^d))}^2 \leq C(T, M_1, M_2),$$

where in the last inequality we used (4.12). On the other hand, by the Madelung transformation

$$\|\nabla J\|_{L_t^\infty L_x^2} = \|\nabla \text{Im}(\bar{\psi} \nabla \psi)\|_{L_t^\infty L_x^2} \leq C \|\psi\|_{L_t^\infty H_x^2}^2 \leq C(T, M_1, M_2).$$

Finally, let us recall that by the polar factorization we have

$$e = \frac{1}{2}|\nabla\sqrt{\rho}|^2 + \frac{1}{2}|\Lambda|^2 + f(\rho) = \frac{1}{2}|\nabla\psi|^2 + f(|\psi|^2).$$

It is straightforward to see that $\nabla|\nabla\psi| \leq |\nabla^2\psi|$ a.e. x , so that

$$\|\nabla\sqrt{e}\|_{L_t^\infty L_x^2} \leq \|\psi\|_{L_t^\infty H_x^2} \leq C(T, M_1, M_2).$$

□

5. DISPERSIVE ESTIMATES AND BOUNDS ON $I(t)$

In this section we will collect some a priori estimates satisfied by finite energy weak solutions to (1.1). If we restricted our analysis to Schrödinger-generated solutions - like the ones constructed in the proof of Theorem 1 and Theorem 6 - then the dispersive estimates inherited by the NLS dynamics would yield a wide range of information. For general solutions that is not the case; the quasi-linear nature of system (1.1) prevents to successfully exploit semigroup techniques to infer suitable smoothing estimates. However we are still able to prove some a priori estimates for general solutions to QHD system in genuine hydrodynamic approaches.

We first define the functional

$$(5.1) \quad V(t) = \int_{\mathbf{R}^d} \frac{|x|^2}{2} \rho(t, x) dx - t \int_{\mathbf{R}^d} x \cdot J(t, x) dx + t^2 E(t),$$

where the total energy $E(t)$ is given by (1.2). The functional $V(t)$ is the hydrodynamic analogue of the pseudo-conformal energy for NLS solutions [34], see also [38, 39].

Proposition 26. *Let (ρ, J) be a finite energy weak solution to (1.1) as in Definition 15 such that*

$$(5.2) \quad \int_{\mathbf{R}^d} |x|^2 \rho_0(x) dx < \infty.$$

Then we have

$$V(t) + d \int_0^t s \int_{\mathbf{R}^d} \rho \left[f'(\rho) - \left(\frac{2}{d} + 1 \right) f(\rho) \right] dx ds = \int_{\mathbf{R}^d} \frac{|x|^2}{2} \rho_0(x) dx.$$

Proof. We first prove that the variance is finite at any time if we assume (5.2). Take $\Theta \in C_c^\infty(\mathbf{R})$ to be a cut-off function such that $0 \leq \Theta \leq 1$, $\Theta(|x|) \equiv 1$ for $|x| \leq 1$ and $\Theta(|x|) = 0$ for $|x| \geq 2$. Let $\Theta_r(x) = \Theta(|x|/r)$ for $r > 0$, then $\int_{\mathbf{R}^d} |x|^2 \Theta_r(x)^2 \rho(t, x) dx < \infty$ is well-defined. Using the continuity equation in (1.1) we can compute

$$\begin{aligned} \frac{d}{dt} \int_{\mathbf{R}^d} |x|^2 \Theta_r^2 \rho(t, x) dx &= - \int_{\mathbf{R}^d} |x|^2 \Theta_r^2 \operatorname{div} J dx \\ &= 2 \int_{\mathbf{R}^d} \Theta_r^2 x \cdot J dx + 2 \int_{\mathbf{R}^d} |x|^2 \Theta_r \nabla \Theta_r \cdot J dx \\ &\leq 2 \|\Theta_r + |x| |\nabla \Theta_r|\|_{L^\infty} \|\Lambda(t)\|_{L^2} \left(\int_{\mathbf{R}^d} |x|^2 \Theta_r^2 \rho(t, x) dx \right)^{\frac{1}{2}} \\ &\leq 4E(t) \left(\int_{\mathbf{R}^d} |x|^2 \Theta_r^2 \rho(t, x) dx \right)^{\frac{1}{2}}, \end{aligned}$$

therefore by Gronwall and the finiteness of energy we obtain

$$\int_{\mathbf{R}^d} \frac{|x|^2}{2} \Theta_r^2 \rho(t, x) dx \lesssim t^2 + \int_{\mathbf{R}^d} \frac{|x|^2}{2} \Theta_r^2 \rho_0(x) dx.$$

Passing to the limit as $r \rightarrow \infty$ we get

$$\int_{\mathbf{R}^d} \frac{|x|^2}{2} \rho(t, x) dx \lesssim t^2 + \int_{\mathbf{R}^d} \frac{|x|^2}{2} \rho_0(x) dx.$$

Now let us consider $V(t)$ and differentiate it with respect to time. By using the equations (2.7) and the conservation of energy we find

$$\begin{aligned} \frac{d}{dt} V(t) &= \int_{\mathbf{R}^d} \frac{|x|^2}{2} \partial_t \rho dx - \int_{\mathbf{R}^d} x \cdot J(t, x) dx - t \int_{\mathbf{R}^d} x \cdot \partial_t J dx + 2tE(t) \\ &= - \int_{\mathbf{R}^d} \frac{|x|^2}{2} \operatorname{div} J dx - \int_{\mathbf{R}^d} x \cdot J(t, x) dx \\ &\quad + t \int_{\mathbf{R}^d} x \cdot \operatorname{div} (\Lambda \otimes \Lambda + \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho}) dx \\ &\quad + t \int_{\mathbf{R}^d} x \cdot \nabla \left(p(\rho) - \frac{1}{4} \nabla \Delta \rho \right) dx + 2tE(t) \\ &= 2tE(t) - t \int_{\mathbf{R}^d} |\Lambda|^2 + |\nabla \sqrt{\rho}|^2 + dp(\rho) dx \\ &= t \int_{\mathbf{R}} 2f(\rho) - dp(\rho) dx = t \int_{\mathbf{R}} (2 + d)f(\rho) - df'(\rho) \rho dx. \end{aligned}$$

By integrating in time we then prove the Proposition. \square

In the case under our consideration we have $f(\rho) = \frac{1}{\gamma} \rho^\gamma$ so that

$$(5.3) \quad V(t) + d \left(1 - \frac{1 + 2/d}{\gamma} \right) \int_0^t s \int \rho^\gamma dx = \int \frac{|x|^2}{2} \rho_0(x) dx.$$

Furthermore, the functional in (5.1) can also be expressed as

$$(5.4) \quad V(t) = \int_{\mathbf{R}^d} \frac{t^2}{2} |\nabla \sqrt{\rho}|^2 + \frac{t^2}{2} |\Lambda - \frac{x}{t} \sqrt{\rho}|^2 + t^2 f(\rho) dx.$$

We exploit (5.3), (5.4) in order to infer a dispersive type estimate for solutions to (1.1).

Proposition 27. *Let (ρ, J) be a finite energy weak solution to system (1.1) and let us further assume that $\int_{\mathbf{R}^d} |x|^2 \rho_0(x) dx < \infty$. Then we have*

$$(5.5) \quad V(t) \lesssim t^{2(1-\sigma)} + \int_{\mathbf{R}^d} \frac{|x|^2}{2} \rho_0(x) dx,$$

where $\sigma = \min\{1, \frac{d}{2}(\gamma - 1)\}$. In particular we have

$$(5.6) \quad \|\nabla \sqrt{\rho}(t)\|_{L^2} \lesssim t^{-\sigma},$$

$$(5.7) \quad \int |\Lambda(t, x) - \frac{x}{t} \sqrt{\rho}(t, x)|^2 dx \lesssim t^{-2\sigma}.$$

and

$$(5.8) \quad \int \rho^\gamma dx \lesssim t^{-2\sigma}.$$

Proof. Let us consider the identity (5.3), hence for $\gamma > 1 + \frac{2}{d}$ we have

$$V(t) \leq \int \frac{|x|^2}{2} \rho_0(x) dx$$

and (5.5) holds for $\sigma = 1$. Let us now consider the case $1 < \gamma \leq 1 + \frac{2}{d}$, if we define

$$F(t) = \frac{t^2}{\gamma} \int \rho^\gamma(t, x) dx$$

then by (5.3) and (5.4) we have

$$F(t) \leq V(t) = [2 + d(1 - \gamma)] \int_0^t \frac{1}{s} F(s) ds + \int \frac{|x|^2}{2} \rho_0(x) dx.$$

By Gronwall we then have

$$F(t) \lesssim t^{2+d(1-\gamma)} F(1) + \int \frac{|x|^2}{2} \rho_0(x) dx,$$

which also implies

$$\int \rho^\gamma(t, x) dx \lesssim t^{d(1-\gamma)}.$$

We can now plug the above estimate in (5.4) in order to obtain

$$V(t) \lesssim \int_0^t s^{1+d(1-\gamma)} ds + \int \frac{x^2}{2} \rho_0(x) dx,$$

which then implies (5.5). \square

Remark 28. Proposition 27 shows that the gradient of mass density converges to zero and formally the velocity field asymptotically approaches a rarefaction wave, namely $v(t, x) \sim x/t$ as $t \rightarrow \infty$. From the conservation of energy we also infer that

$$\lim_{t \rightarrow \infty} \frac{1}{2} \|\Lambda(t)\|_{L^2}^2 = E(0),$$

i.e. for large times all the energy is transferred to the kinetic part. Moreover by Gagliardo-Nirenberg the decay of $\|\nabla \sqrt{\rho}(t)\|_{L^2}$ implies the following dispersive estimate

$$\|\sqrt{\rho}(t)\|_{L^p} \lesssim \|\sqrt{\rho}(t)\|_{L^2}^{1-\alpha} \|\nabla \sqrt{\rho}(t)\|_{L^2}^\alpha \lesssim t^{-\alpha\sigma},$$

where $0 \leq \alpha \leq 1$ such that

$$\frac{1}{p} = \frac{1}{2} - \frac{\alpha}{d},$$

and $2 \leq p < \infty$ for $d = 2$, $2 \leq p \leq 6$ for $d = 3$. Notice that in the case $\gamma \geq 1 + 2/d$ this is still consistent with the dispersive estimate for the free Schrödinger equation.

In general similar estimates appear in many contexts for the study of evolutionary PDEs, see [73, 36, 72, 62] and many others. In particular estimates like (5.5) have been considered in classical compressible fluid dynamics [21, 71, 72], or in [14, 66, 40] for the analogue (5.7) in kinetic theory. Those are also somehow reminiscent of the vector field method [49] used to study nonlinear wave equations, see also [28] for recent applications. In the wave function dynamics context, the fact that solutions to the nonlinear problem disperse as much as the linear solutions gives some information about their asymptotic behavior, see [34]. Recently an alternative proof, based on interaction Morawetz estimates [62, 57], was developed in order to show asymptotic completeness for mass-supercritical, energy-subcritical NLS

equations [23, 22, 35, 69]. It turns out that such estimates hold also for arbitrary weak solutions to (1.1).

In this section we also present the Morawetz-type estimates, in a general form for the Euler-Korteweg system

$$(5.9) \quad \begin{cases} \partial_t \rho + \operatorname{div} J = 0 \\ \partial_t J + \operatorname{div} \left(\frac{J \otimes J}{\rho} \right) + \nabla p(\rho) = \rho \nabla \left(\operatorname{div}(\kappa(\rho) \nabla \rho) - \frac{1}{2} \kappa'(\rho) |\nabla \rho|^2 \right), \end{cases}$$

where $\kappa : (0, \infty) \rightarrow (0, \infty)$ is a smooth function, and by choosing $\kappa(\rho) = \frac{1}{4\rho}$ we recover the QHD system. Moreover, while for the dispersive estimates in Proposition 27 we need to assume the initial mass density to have finite variance (5.2), in the next Proposition we only need the weak solutions to be finite energy.

Proposition 29. *Let (ρ, J) be a weak solution to the system (5.9) such that*

$$\|\sqrt{\rho}\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} + \|\kappa(\rho)^{\frac{1}{2}} \nabla \rho\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} + \|\Lambda\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq M_1.$$

We define $K(\rho) = \int_0^\rho s \kappa(s) ds$ and further assume $K(\rho)$ is a convex function in ρ , namely $\kappa(\rho) + \rho \kappa'(\rho) \geq 0$ for $\rho \geq 0$. Then we have

$$\| |\nabla|^{\frac{3-d}{2}} \sqrt{\rho K(\rho)} \|_{L^2(\mathbf{R}_t \times \mathbf{R}_x^d)}^2 + \| |\nabla|^{\frac{1-d}{2}} \sqrt{\rho p(\rho)} \|_{L^2(\mathbf{R}_t \times \mathbf{R}_x^d)}^2 \lesssim M_1^4.$$

Proof. Through a direct computation, the right hand side of the equation for momentum density J in (5.9) can be written as

$$\nabla \operatorname{div}(\rho \kappa(\rho) \nabla \rho) - \operatorname{div}(\kappa(\rho) \nabla \rho \otimes \nabla \rho) - \frac{1}{2} \nabla [(\kappa(\rho) + \rho \kappa'(\rho)) |\nabla \rho|^2].$$

By the definition of $K(\rho)$ we have

$$\nabla \operatorname{div}(\rho \kappa(\rho) \nabla \rho) = \nabla \Delta K(\rho),$$

then we can rewrite the system (5.9) as

$$(5.10) \quad \begin{cases} \partial_t \rho + \operatorname{div} J = 0 \\ \partial_t J + \operatorname{div} \left(\frac{J \otimes J}{\rho} + \kappa(\rho) \nabla \rho \otimes \nabla \rho \right) + \nabla p(\rho) = \nabla \Delta K(\rho) - \frac{1}{2} \nabla [(\kappa(\rho) + \rho \kappa'(\rho)) |\nabla \rho|^2] \end{cases}$$

Now we consider the interaction functional

$$H(t) = \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{x-y}{|x-y|} \cdot J(t, x) dx dy.$$

The functional $H(t)$ is well-defined on $[0, T]$ with bound

$$|H(t)| \leq \|\rho\|_{L_t^\infty L_x^1} \|J\|_{L_t^\infty L_x^1} \leq M_1^4.$$

By differentiating $H(t)$ in time using the system (5.10) and integrating by parts we obtain

$$\begin{aligned}
 (5.11) \quad \frac{d}{dt}H(t) = & - \int_{\mathbf{R}^{2d}} J(t, y) \cdot B(x, y) \cdot J(t, x) dx dy + \int_{\mathbf{R}^{2d}} \rho(t, y) \Lambda(t, x) \cdot B(x, y) \cdot \Lambda(t, x) dx dy \\
 & + \int_{\mathbf{R}^{2d}} \rho(t, y) \kappa(\rho)(t, x) \nabla \rho(t, x) \cdot B(x, y) \cdot \nabla \rho(t, x) dx dy \\
 & + \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} \left[p(\rho) + \frac{1}{2} (\kappa(\rho) + \rho \kappa'(\rho)) |\nabla \rho|^2 \right] (t, x) dx dy \\
 & - \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} \triangle K(\rho)(t, x) dx dy,
 \end{aligned}$$

where

$$B(x, y) = \nabla_x \left(\frac{x-y}{|x-y|} \right) = \frac{1}{|x-y|} \left[Id - \frac{(x-y) \otimes (x-y)}{|x-y|^2} \right].$$

Since $B(x, y)$ is a semi-positive definite matrix and symmetric in x and y , we have

$$\begin{aligned}
 \int_{\mathbf{R}^{2d}} J(t, y) \cdot B(x, y) \cdot J(t, x) dx dy &= \int_{\mathbf{R}^{2d}} (\sqrt{\rho} \Lambda)(t, y) \cdot B(x, y) \cdot (\sqrt{\rho} \Lambda)(t, x) dx dy \\
 &\leq \frac{1}{2} \int_{\mathbf{R}^{2d}} \rho(t, y) \Lambda(t, x) \cdot B(x, y) \cdot \Lambda(t, x) dx dy \\
 &\quad + \frac{1}{2} \int_{\mathbf{R}^{2d}} \rho(t, x) \Lambda(t, y) \cdot B(x, y) \cdot \Lambda(t, y) dx dy \\
 &= \int_{\mathbf{R}^{2d}} \rho(t, y) \Lambda(t, x) \cdot B(x, y) \cdot \Lambda(t, x) dx dy.
 \end{aligned}$$

Therefore the first and second lines in the right hand side of (5.11) are non-negative. Also by our assumption

$$\int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} [(\kappa(\rho) + \rho \kappa'(\rho)) |\nabla \rho|^2] (t, x) dx dy \geq 0.$$

To estimate the remaining integrals in the right hand side of (5.11), we first notice that by the symmetry in x and y , we have

$$\begin{aligned}
 \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} p(\rho)(t, x) dx dy &= \frac{1}{2} \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} p(\rho)(t, x) dx dy \\
 &\quad + \frac{1}{2} \int_{\mathbf{R}^{2d}} \rho(t, x) \frac{d-1}{|x-y|} p(\rho)(t, y) dx dy \\
 &\geq \int_{\mathbf{R}^{2d}} \sqrt{\rho p(\rho)}(t, y) \frac{d-1}{|x-y|} \sqrt{\rho p(\rho)}(t, x) dx dy
 \end{aligned}$$

The convolution of the kernel $|x-y|^{-1}$ gives

$$\int_{\mathbf{R}^d} \frac{1}{|x-y|} \sqrt{\rho p(\rho)}(t, x) dx = C |\nabla|^{1-d} \sqrt{\rho p(\rho)}(t, y),$$

then we obtain

$$\begin{aligned}
 \int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} p(\rho)(t, x) dx dy &\geq C \int_{\mathbf{R}^d} \sqrt{\rho p(\rho)}(t, y) |\nabla|^{1-d} \sqrt{\rho p(\rho)}(t, y) dy \\
 &= C \| |\nabla|^{\frac{1-d}{2}} \sqrt{\rho p(\rho)} \|_{L_x^2}^2(t).
 \end{aligned}$$

Similarly the last line of (5.11) can be written as

$$-\int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} \Delta K(\rho)(t, x) dx dy = C \int_{\mathbf{R}^d} \rho(t, y) |\nabla|^{3-d} K(\rho)(t, y) dy,$$

and again by the symmetry we obtain

$$-\int_{\mathbf{R}^{2d}} \rho(t, y) \frac{d-1}{|x-y|} \Delta K(\rho)(t, x) dx dy \geq C \|\nabla\|^{\frac{3-d}{2}} \sqrt{\rho K(\rho)}\|_{L_x^2}^2(t).$$

By summarising the argument above, we get

$$(5.12) \quad \frac{d}{dt} H(t) \geq C \|\nabla\|^{\frac{1-d}{2}} \sqrt{\rho p(\rho)}\|_{L_x^2}^2(t) + C \|\nabla\|^{\frac{3-d}{2}} \sqrt{\rho K(\rho)}\|_{L_x^2}^2(t),$$

then integrating (5.12) in time shows

$$\|\nabla\|^{\frac{1-d}{2}} \sqrt{\rho p(\rho)}\|_{L_t^2 L_x^2}^2 + C \|\nabla\|^{\frac{3-d}{2}} \sqrt{\rho K(\rho)}\|_{L_t^2 L_x^2}^2 \lesssim H(T) - H(0) \lesssim M_1^4.$$

□

In the remaining part of this Section we are going to discuss the 2nd order energy functional introduced in (1.6) in multi-dimension. As already discussed in Section 2, if we consider Schrödinger-generated solutions to the QHD system (1.1), say $\rho = |\psi|^2$, $J = \text{Im}(\bar{\psi} \nabla \psi)$, for some H^2 solutions ψ to the nonlinear Schrödinger equation

$$\begin{cases} i\partial_t \psi = -\frac{1}{2} \Delta \psi + f'(|\psi|^2) \psi \\ \psi(0) = \psi_0 \end{cases}$$

then the functional consists in the square L^2 norm of $\partial_t \psi$,

$$I(t) = \int_{\mathbf{R}^d} |\partial_t \psi|^2 dx.$$

By using the polar factorization we see that it is possible to write

$$I(t) = \int_{\mathbf{R}^d} (\partial_t \sqrt{\rho})^2 + \lambda^2 dx,$$

where we define $\lambda := \text{Im}(\bar{\psi} \partial_t \psi)$ and direct computation shows

$$\xi = \sqrt{\rho} \lambda = \text{Im}(\bar{\psi} \partial_t \psi) = -\frac{1}{4} \Delta \rho + \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + \rho f'(\rho)$$

Moreover by using the regularity of ψ we also have that $\lambda = 0$ a.e. in the vacuum set $\{\rho = 0\}$. Therefore it is reasonable to define

$$\lambda = \begin{cases} -\frac{1}{2} \Delta \sqrt{\rho} + \frac{1}{2} \frac{|\Lambda|^2}{\sqrt{\rho}} + f'(\rho) \sqrt{\rho} & \text{in } \{\rho > 0\} \\ 0 & \text{elsewhere} \end{cases}$$

For Schrödinger-generated solutions, $I(t)$ is uniformly bounded on compact time intervals due to the well-posedness results of NLS in $H^2(\mathbf{R}^d)$. Alternatively, here we give a direct calculation of the time derivative to $I(t)$ for smooth, non-vanishing solutions $(\sqrt{\rho}, \Lambda)$ to (1.1). To present this computation, we first state the following Lemma, which shows the conservation law for the total energy density e of smooth, non-vanishing solutions.

Lemma 30. *Let $(\sqrt{\rho}, \Lambda)$ be a smooth solution to (1.1) such that $\rho > 0$. Then the energy density e satisfies the following conservation law*

$$(5.13) \quad \partial_t e + \text{div}(\Lambda \lambda - \partial_t \sqrt{\rho} \nabla \sqrt{\rho}) = 0.$$

Proof. Since we are dealing with smooth solutions and we have $\rho > 0$, then we can write system (1.1) as

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho v) = 0 \\ \rho \partial_t v + \rho v \cdot \nabla v + \rho \nabla f'(\rho) = \frac{1}{2} \rho \nabla \left(\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}} \right) \end{cases}$$

and we notice that the equation for the momentum density can be equivalently written as

$$\rho \partial_t v + \rho \nabla \mu = 0,$$

where μ is the chemical potential defined in (1.3). Analogously, the energy density reads

$$e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} \rho |v|^2 + f(\rho).$$

By using this expression we can differentiate it with respect to time and find

$$\begin{aligned} \partial_t e &= \nabla \sqrt{\rho} \cdot \partial_t \nabla \sqrt{\rho} + \left(\frac{1}{2} |v|^2 + f'(\rho) \right) \partial_t \rho + \rho v \partial_t v \\ &= \operatorname{div}(\nabla \sqrt{\rho} \partial_t \sqrt{\rho}) + \left(-\frac{1}{2} \frac{\Delta \sqrt{\rho}}{\sqrt{\rho}} + \frac{1}{2} |v|^2 + f'(\rho) \right) \partial_t \rho + \rho v \partial_t v, \end{aligned}$$

where in the last identity we used the Leibniz formula for the first term. By using the evolution equations above and definition (1.3) we then have

$$\begin{aligned} \partial_t e &= \operatorname{div}(\nabla \sqrt{\rho} \partial_t \sqrt{\rho}) - \mu \operatorname{div}(\rho v) - \rho v \cdot \nabla \mu \\ &= \partial_x (\partial_x \sqrt{\rho} \partial_t \sqrt{\rho} - \rho v \mu). \end{aligned}$$

□

Remark 31. The calculation in Lemma above requires smoothness and positivity of solutions in order to be rigorously justified. On the other hand it is an interesting question to see whether this conservation law, or its weaker version with the inequality, hold for a larger class of solutions. For this problem we refer the paper [25] where the authors determine some conditions on the velocity field and the mass density which allow to show the conservation of energy.

Proposition 32. *Let (ρ, J) be a smooth solution to (1.1) such that $\rho > 0$. Then we have*

$$(5.14) \quad \frac{d}{dt} I(t) = 2 \int_{\mathbf{R}^d} \lambda \partial_t \sqrt{\rho} p'(\rho) dx.$$

Proof. Since we consider smooth, positive solutions to (1.1), then we write our functional $I(t)$ as

$$I(t) = \frac{1}{2} \int_{\mathbf{R}^d} \rho (\mu^2 + \sigma^2) dx,$$

where chemical potential

$$\mu = \lambda / \sqrt{\rho} = -\frac{\Delta \sqrt{\rho}}{2\sqrt{\rho}} + \frac{1}{2} |v|^2 + f'(\rho),$$

and we define $\sigma = \partial_t \log \sqrt{\rho}$. In this framework it is possible to write down the evolution equations for μ, σ .

By using the formula

$$\rho \mu = -\frac{1}{4} \Delta \rho + e + p(\rho)$$

and the conservation law (5.13) of e , we have

$$\begin{aligned}\partial_t(\rho\mu) &= \partial_t \left(e - \frac{1}{4}\Delta\rho + p(\rho) \right) \\ &= \operatorname{div}(\nabla\sqrt{\rho}\partial_t\sqrt{\rho} - \rho v\mu) - \frac{1}{4}\partial_t\Delta\rho + \partial_t p(\rho).\end{aligned}$$

Again by using the continuity equation we may write

$$(5.15) \quad \rho\partial_t\mu + \rho v \cdot \nabla\mu = \operatorname{div}(\nabla\sqrt{\rho}\partial_t\sqrt{\rho}) - \frac{1}{4}\partial_t\Delta\rho + \partial_t p(\rho).$$

Now, to write the equation for σ we may proceed in the following way. By writing the continuity equation as below

$$\partial_t\rho + v \cdot \nabla\rho + \rho \operatorname{div} v = 0,$$

we find the equation for $\log\sqrt{\rho}$, namely

$$\partial_t \log\sqrt{\rho} + v \cdot \nabla \log\sqrt{\rho} + \frac{1}{2} \operatorname{div} v = 0.$$

By differentiating the last equation with respect to time and by using $\partial_t v = -\nabla\mu$ we then obtain

$$\partial_t\sigma + v \cdot \nabla\sigma - \nabla\mu \cdot \nabla \log\sqrt{\rho} - \frac{1}{2}\Delta\mu = 0.$$

By multiplying this by ρ we get

$$(5.16) \quad \rho\partial_t\sigma + \rho v \cdot \nabla\sigma = \frac{1}{2} \operatorname{div}(\rho\nabla\mu).$$

Now we can use the equations (5.15) and (5.16) to compute the time derivative of the functional $I(t)$. By differentiating the functional $I(t)$ in time using the continuity equation of ρ , the evolution (5.15) and (5.16) of μ and σ , we obtain

$$\begin{aligned}\frac{d}{dt}I(t) &= \int \frac{1}{2}(\mu^2 + \sigma^2)\partial_t\rho + \mu\rho\partial_t\mu + \sigma\rho\partial_t\sigma \, dx \\ &= \int \mu \left(\operatorname{div}(\nabla\sqrt{\rho}\partial_t\sqrt{\rho}) - \frac{1}{4}\Delta\partial_t\rho + \partial_t p(\rho) \right) \\ &\quad + \frac{\sigma}{2} \operatorname{div}(\rho\nabla\mu) \, dx.\end{aligned}$$

By using that $\sqrt{\rho}\sigma = \partial_t\sqrt{\rho}$, $\rho\sigma = \frac{1}{2}\partial_t\rho$ and integration by parts, we get

$$\frac{d}{dt}I(t) = \int \mu\partial_t p(\rho) \, dx = 2 \int \lambda\partial_t\rho p'(\rho) \, dx,$$

which finishes the proof. \square

Recall that $p(\rho) = \frac{\gamma-1}{\gamma}\rho^\gamma$, then as a direct consequence of identity (5.14) we have

$$(5.17) \quad \frac{d}{dt}I(t) = 2(\gamma-1) \int_{\mathbf{R}^d} \rho^{\gamma-1} \lambda \partial_t \sqrt{\rho} \, dx \lesssim \|\rho(t)\|_{L_x^\infty}^{\gamma-1} I(t).$$

To apply the Gronwall's argument to functional $I(t)$, we need further control on $\|\rho(t)\|_{L_x^\infty}$, which can not be directly obtained from the energy bounds in multi-dimension. Here we recall the following 2D log-type Sobolev inequality [16].

Lemma 33. *For any $f \in H^2(\mathbf{R}^2)$ we have*

$$(5.18) \quad \|f\|_{L^\infty(\mathbf{R}^2)} \leq C(1 + \sqrt{\log(1 + \|f\|_{H^2(\mathbf{R}^2))}).$$

By using Proposition 32 and the log-type Sobolev inequality in 2D, we obtain the uniform bound of functional $I(t)$ via standard Gronwall's argument.

Corollary 1. *Let (ρ, J) be a smooth solution to (1.1) on $[0, T] \times \mathbf{R}^2$ such that $\rho > 0$, and we further assume the pressure $p(\rho) = (1 - \frac{1}{\gamma})\rho^\gamma$ with $1 < \gamma \leq 2$. Then for all $t \in [0, T]$ we have*

$$I(t) \leq C(T)I(0).$$

Proof. First we notice that for all $t \in [0, T]$, the hydrodynamic data $(\rho(t, \cdot), J(t, \cdot))$ satisfies the assumption of Proposition 22, therefore we can lift $(\rho, J)(t)$ to a wave function $\psi(t) \in H_x^2(\mathbf{R}^2)$. Furthermore following the proof of Proposition 22 we have

$$\|\psi(t)\|_{H_x^1}^2 \leq M(t) + E(t),$$

where $M(t)$ and $E(t)$ are the conserved total mass and energy of (ρ, J) , and

$$\|\Delta\psi(t)\|_{L_x^2}^2 \leq C I(t) + C(M(t), E(t)).$$

We remark that the wave function ψ is constructed at each isolated time $t \in [0, T]$, hence it is not a solution to any equation. By using the log-type Sobolev inequality (5.18), we have

$$\begin{aligned} \|\rho(t)\|_{L_x^\infty} &= \|\psi(t)\|_{L_x^\infty}^2 \\ &\leq C(1 + \log(1 + \|\psi(t)\|_{H_x^2})). \end{aligned}$$

Then the H_x^2 bound of $\psi(t)$ implies

$$(5.19) \quad \|\rho(t)\|_{L_x^\infty} \leq C \log I(t) + C(M(t), E(t)).$$

Now by Proposition 32 and (5.19) we have

$$\frac{d}{dt} I(t) \leq (\gamma - 1) \|\rho(t)\|_{L_x^\infty}^{\gamma-1} I(t) \leq C(\log I(t))^{\gamma-1} I(t) + C(M(t), E(t))I(t),$$

which implies

$$\frac{d}{dt} \log I(t) \leq C[(\log I(t))^{\gamma-1} + 1].$$

Hence by standard Gronwall's argument we conclude the proof. \square

Remark 34. The Gronwall inequality used in the proof above clearly restricts the boundedness result of $I(t)$ to $\gamma \leq 2$. The case $\gamma > 2$ in 2D and the higher dimensional cases would require a finer analysis, for instance by using some dispersive type estimates. This is consistent for example with what is discussed in [68].

6. STABILITY OF WEAK SOLUTIONS WITH POSITIVE DENSITY

In the last Section of this paper on multi-dimensional QHD system, we will consider the compactness of weak solutions with non-vanishing density, in the framework of finite energy and functional $I(t)$. Let us now consider a sequence of weak solutions $\{(\sqrt{\rho_n}, \Lambda_n)\}$ to (1.1) in the compactness class (Definition 20), which satisfies the following uniform bounds,

$$\begin{aligned} \|\sqrt{\rho_n}\|_{L^\infty(0, T; H^1(\mathbf{R}^d))} + \|\sqrt{\rho_n}\|_{W^{1, \infty}(0, T; L^2(\mathbf{R}^d))} &\leq C \\ \|\Lambda_n\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} &\leq C, \quad \|\lambda_n\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} \leq C, \end{aligned}$$

where λ_n is such that

$$\sqrt{\rho_n} \lambda_n = -\frac{1}{4} \Delta \rho_n + e_n + p(\rho)$$

and

$$e_n = \frac{1}{2} |\nabla \sqrt{\rho_n}|^2 + \frac{1}{2} |\Lambda_n|^2 + f(\rho_n).$$

We further assume for all $n \in \mathbf{N}$ a.e. $t \in [0, T]$, $\rho_n(t, \cdot)$ is continuous and positive, i.e. $\rho_n(t, \cdot) > 0$. Since the density is always positive, the quantity λ_n can be written as

$$(6.1) \quad \lambda_n = \frac{1}{2} \left(-\Delta \sqrt{\rho_n} + \frac{|\Lambda_n|^2}{\sqrt{\rho_n}} \right) + f'(\rho_n) \sqrt{\rho_n}.$$

Proposition 35. *Let $(\sqrt{\rho_n}, \Lambda_n)$ be a finite energy weak solution to the QHD system (1.1) satisfying the bounds*

$$(6.2) \quad \begin{aligned} & \|\sqrt{\rho_n}\|_{L^\infty(0,T;H^1(\mathbf{R}^d))} + \|\sqrt{\rho_n}\|_{W^{1,\infty}(0,T;L^2(\mathbf{R}^d))} \leq C \\ & \|\Lambda_n\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C, \quad \|\lambda_n\|_{L^\infty(0,T;L^2(\mathbf{R}^d))} \leq C. \end{aligned}$$

Moreover we assume for a.e. $t \in [0, T]$, $\rho_n(t, \cdot)$ is continuous and $\rho_n(t, \cdot) > 0$. Then we have

$$(6.3) \quad \|\Delta \rho_n\|_{L_t^\infty L_x^2} + \|\nabla J_n\|_{L_t^\infty L_x^2} + \|\nabla \sqrt{e_n}\|_{L_t^\infty L_x^2} \leq C.$$

Proof. Since we only consider a fix n in Proposition 35, we will drop out the subscript n in this proof without causing any confusing, and we use C to denote the uniform constant independent of the sequence, which may change from line to line.

To obtain the estimates (6.3), we notice that for a.e. $t \in [0, T]$, the hydrodynamic data $(\sqrt{\rho}, \Lambda)(t, \cdot)$ satisfies the assumption of Proposition 22, hence it can be lifted to a wave function $\psi(t, \cdot) \in H_x^2(\mathbf{R}^d)$ with uniform bounds

$$\|\psi\|_{L_t^\infty H_x^2} \leq C.$$

The estimates (6.3) is a direct consequence of the $L_t^\infty H_x^2$ bound of ψ . By using $\Delta \rho = 2 \operatorname{Re}(\bar{\psi} \Delta \psi) + 2 |\nabla \psi|^2$, Hölder's inequality and Sobolev embedding we have

$$\|\Delta \rho\|_{L_t^\infty L_x^2} \lesssim \|\psi\|_{L_t^\infty H_x^2}^2 \leq C,$$

On the other hand, by the Madelung transformation

$$\|\nabla J\|_{L_t^\infty L_x^2} = \|\nabla \operatorname{Im}(\bar{\psi} \nabla \psi)\|_{L_t^\infty L_x^2} \leq \|\psi\|_{L_t^\infty L_x^2}^2 \leq C.$$

Finally, let us recall that by the polar factorization we have

$$e = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho) = \frac{1}{2} |\nabla \psi|^2 + f(|\psi|^2).$$

It is straightforward to see that $\nabla |\nabla \psi| \leq |\nabla^2 \psi|$ a.e. x , so that

$$\|\nabla \sqrt{e}\|_{L_t^\infty L_x^2} \leq \|\psi\|_{L_t^\infty H_x^2} \leq C.$$

□

Remark 36. As discussed in the last Section the wave function ψ constructed in the proof is not a solution to the NLS equations. On the other hand the argument of Proposition 35 can be extended to weak solutions $(\sqrt{\rho}, \Lambda)$ such that, for a.e. $t \in [0, T]$, $(\sqrt{\rho}, \Lambda)(t, \cdot)$ has suitable vacuum and vortex structure which is compatible with a wave function lift method, for instance the pointwise vacuum and quantised vorticity structure in 2D.

The bounds in (6.2) imply that, up to passing to subsequences, we have

$$(6.4) \quad \sqrt{\rho_n} \rightharpoonup \sqrt{\rho}, \quad L^\infty(0, T; H^1(\mathbf{R}^d)) \cap W^{1, \infty}(0, T; L^2(\mathbf{R}^d)),$$

$$(6.5) \quad \Lambda_n \rightharpoonup \Lambda, \quad L^\infty(0, T; L^2(\mathbf{R}^d)),$$

and the uniform bound of e_n in (6.3) implies local strong convergence

$$(6.6) \quad \sqrt{e_n} \rightarrow \omega, \quad L^\infty(0, T; L^2_{loc}(\mathbf{R}^d)).$$

it is not straightforward to show the identity for the limit ω and $(\sqrt{\rho}, \Lambda)$, because of the lack of compactness for $(\nabla \sqrt{\rho_n}, \Lambda_n)$. However the next proposition shows $(\sqrt{\rho_n}, \Lambda_n)$ indeed converges strongly. Furthermore, we prove that ω and Λ vanish almost everywhere in the vacuum $\{\rho = 0\}$, which matches the physical interpretation of the energy density.

Proposition 37. *Let $(\sqrt{\rho_n}, \Lambda_n)$ satisfy the bounds*

$$(6.7) \quad \begin{aligned} & \|\sqrt{\rho_n}\|_{L^\infty(0, T; H^1(\mathbf{R}^d))} + \|\sqrt{\rho_n}\|_{W^{1, \infty}(0, T; L^2(\mathbf{R}^d))} \leq C \\ & \|\Lambda_n\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} \leq C, \quad \|\lambda_n\|_{L^\infty(0, T; L^2(\mathbf{R}^d))} \leq C. \end{aligned}$$

Moreover we assume for all $n \in \mathbf{N}$ and a.e. $t \in [0, T]$, $\rho_n(t, \cdot)$ is continuous and $\rho_n(t, \cdot) > 0$. Then we have

$$(6.8) \quad \omega^2 = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho)$$

is satisfied for a.e. $(t, x) \in [0, T] \times \mathbf{R}^d$, and $\omega = 0$ a.e. $(t, x) \in \{\rho = 0\}$. Furthermore we have the following local strong convergence

$$(6.9) \quad \begin{aligned} & \nabla \sqrt{\rho_n} \rightarrow \nabla \sqrt{\rho}, \quad L^p(0, T; L^2_{loc}(\mathbf{R}^d)), \\ & \Lambda_n \rightarrow \Lambda, \quad L^p(0, T; L^2_{loc}(\mathbf{R}^d)), \end{aligned}$$

for $2 \leq p < \infty$.

Proof. Multiplying the energy density by ρ_n , we have that

$$(6.10) \quad \rho_n e_n = \frac{1}{8} |\nabla \rho_n|^2 + \frac{1}{2} |J_n|^2 + f(\rho_n) \rho_n.$$

Now, from the uniform bounds (6.3) we know that, up to subsequences,

$$\begin{aligned} \rho_n &\rightarrow \rho, & L^2(0, T; H^1_{loc}(\mathbf{R}^d)) \\ J_n &\rightarrow J, & L^2(0, T; L^2_{loc}(\mathbf{R}^d)), \end{aligned}$$

hence, by passing to the limit in (6.10) we obtain

$$(6.11) \quad \rho \omega^2 = \frac{1}{8} |\nabla \rho|^2 + \frac{1}{2} |J|^2 + f(\rho), \quad \text{a.e.}$$

Recall that

$$J_n = \sqrt{\rho_n} \Lambda_n,$$

then again by passing to the limit we have

$$J = \sqrt{\rho} \Lambda, \quad \text{a.e.}$$

which implies that we can write (6.11) as

$$(6.12) \quad \rho \left(\omega^2 - \frac{1}{2} |\nabla \sqrt{\rho}|^2 - \frac{1}{2} |\Lambda|^2 - f(\rho) \right) = 0, \quad \text{a.e.}$$

We now claim that $\omega^2 = 0$ a.e. on $\{\rho = 0\}$. From (6.1) we know that

$$e_n = \sqrt{\rho_n} \lambda_n + \frac{1}{4} \Delta \rho_n - p(\rho_n)$$

and by passing to the limit we obtain

$$\omega^2 = \sqrt{\rho} \lambda + \frac{1}{4} \Delta \rho - p(\rho).$$

By the vanishing derivative Lemma 14 we have $\Delta \rho = 0$ a.e. on $\{\rho = 0\}$, consequently we have that $\omega = 0$ a.e. on $\{\rho = 0\}$. On the other hand, for any $0 < R < \infty$ let us define $V_R = \{\rho = 0\} \cap B_R(0)$, then by using (6.3) and (6.6) we have

$$\begin{aligned} \int_{V_R} \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 dx &\leq \liminf_{n \rightarrow \infty} \int_{V_R} \frac{1}{2} |\nabla \sqrt{\rho_n}|^2 + \frac{1}{2} |\Lambda_n|^2 dx \\ &= \lim_{n \rightarrow \infty} \int_{V_R} e_n - f(\rho_n) dx \\ &= \int_{V_R} \omega^2 - f(\rho) dx = 0, \end{aligned}$$

where in the last equality we use the local strong convergence of $\sqrt{e_n}$ and ρ_n . This implies that $\Lambda = 0$ a.e. on $\{\rho = 0\}$ and consequently from (6.11) we also have

$$\omega^2 = \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 + f(\rho).$$

Last, to prove local strong convergence we notice that for any $R > 0$, similar as before we have at a.e $t \in [0, T]$

$$\lim_{n \rightarrow \infty} \int_{B_R(0)} e_n - f(\rho_n) dx = \int_{B_R(0)} \omega^2 - f(\rho) dx dt.$$

Therefore by same argument as above

$$\begin{aligned} \int_{B_R(0)} \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 dx &\leq \liminf_{n \rightarrow \infty} \int_{B_R(0)} \frac{1}{2} |\nabla \sqrt{\rho_n}|^2 + \frac{1}{2} |\Lambda_n|^2 dx \\ &= \lim_{n \rightarrow \infty} \int_{B_R(0)} e_n - f(\rho_n) dx \\ &= \int_{B_R(0)} \omega^2 - f(\rho) dx \\ &= \int_{B_R(0)} \frac{1}{2} |\nabla \sqrt{\rho}|^2 + \frac{1}{2} |\Lambda|^2 dx, \end{aligned}$$

and we conclude for $2 \leq p < \infty$.

$$\|\nabla \sqrt{\rho}\|_{L_t^p L_{x,loc}^2} = \lim_{n \rightarrow \infty} \|\nabla \sqrt{\rho_n}\|_{L_t^p L_{x,loc}^2} \text{ and } \|\Lambda\|_{L_t^p L_{x,loc}^2} = \lim_{n \rightarrow \infty} \|\Lambda_n\|_{L_t^p L_{x,loc}^2}.$$

Then by standard argument we can improve the weak convergence to a strong one. \square

To conclude this Section, we prove the stability Theorem 11 of weak solutions $(\sqrt{\rho_n}, \Lambda_n)$ with positive density, which is a direct consequence of the compactness Proposition 37. As we remarked in the introduction that we do not require any uniform lower bound on the density, and the limiting weak solution may contain vacuum.

Proof of Theorem 11. By the definition of weak solutions, sequence $\{(\rho_n, J_n)\}$ satisfy, for any $n \geq 1$, test function $\eta \in \mathcal{C}_c^\infty([0, T] \times \mathbf{R}^d)$ and vector test function $\xi \in \mathcal{C}_c^\infty([0, T] \times \mathbf{R}^d; \mathbf{R}^d)$

$$\begin{aligned} & \int_0^T \int_{\mathbf{R}^d} \rho_n \partial_t \eta + J_n \cdot \nabla \eta dx dt + \int_{\mathbf{R}^d} \rho_{n,0} \eta(0) dx = 0, \\ & \int_0^T \int_{\mathbf{R}^d} [J_n \cdot \partial_t \xi + (\Lambda_n \otimes \Lambda_n + \nabla \sqrt{\rho_n} \otimes \nabla \sqrt{\rho_n}) : \nabla \xi \\ & \quad p(\rho_n) \operatorname{div} \xi - \frac{1}{4} \rho_n \Delta \operatorname{div} \xi] dx dt + \int_{\mathbf{R}^d} J_{n,0} \cdot \xi(0) dx = 0. \end{aligned}$$

Then by Proposition 37, the strong convergence of ρ_n and J_n in $L_t^p L_{x,loc}^2$, $2 \leq p < \infty$, implies

$$\begin{aligned} & \int_0^T \int_{\mathbf{R}^d} \rho_n \partial_t \eta dx dt \rightarrow \int_0^T \int_{\mathbf{R}^d} \rho \partial_t \eta dx dt, \\ & \int_0^T \int_{\mathbf{R}^d} J_n \cdot \nabla \xi dx dt \rightarrow \int_0^T \int_{\mathbf{R}^d} J \cdot \nabla \xi dx dt, \\ & \int_0^T \int_{\mathbf{R}^d} J_n \cdot \nabla \eta dx dt \rightarrow \int_0^T \int_{\mathbf{R}^d} J \cdot \nabla \eta dx dt, \\ & \int_0^T \int_{\mathbf{R}^d} p(\rho_n) \operatorname{div} \xi dx dt \rightarrow \int_0^T \int_{\mathbf{R}^d} p(\rho) \operatorname{div} \xi dx dt, \\ & - \int_0^T \int_{\mathbf{R}^d} \frac{1}{4} \rho_n \Delta \operatorname{div} \xi dx dt \rightarrow - \int_0^T \int_{\mathbf{R}^d} \frac{1}{4} \rho \Delta \operatorname{div} \xi dx dt, \end{aligned}$$

and also we have

$$\begin{aligned} & \int_0^T \int_{\mathbf{R}^d} (\Lambda_n \otimes \Lambda_n + \nabla \sqrt{\rho_n} \otimes \nabla \sqrt{\rho_n}) : \nabla \xi dx dt \\ & \rightarrow \int_0^T \int_{\mathbf{R}^d} (\Lambda \otimes \Lambda + \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho}) : \nabla \xi dx dt. \end{aligned}$$

Similarly we can prove the convergence of the integrals of initial data.

Thus the limit functions (ρ, J) satisfy

$$\begin{aligned} & \int_0^T \int_{\mathbf{R}^d} \rho \partial_t \eta + J \cdot \nabla \eta dx dt + \int_{\mathbf{R}^d} \rho_0 \eta(0) dx = 0, \\ & \int_0^T \int_{\mathbf{R}^d} [J \cdot \partial_t \xi + (\Lambda \otimes \Lambda + \nabla \sqrt{\rho} \otimes \nabla \sqrt{\rho}) : \nabla \xi \\ & \quad p(\rho) \operatorname{div} \xi - \frac{1}{4} \rho \Delta \operatorname{div} \xi] dx dt + \int_{\mathbf{R}^d} J_{n,0} \cdot \xi(0) dx = 0. \end{aligned}$$

Last we need to check the generalised irrotationality condition is preserved. For weak solutions (ρ_n, J_n) , by Definition 15 we have

$$\nabla \wedge J_n = 2 \nabla \sqrt{\rho_n} \wedge \Lambda_n.$$

Using the compactness of J_n , $\nabla \sqrt{\rho_n}$ and Λ_n and passing to the limit, we conclude

$$\nabla \wedge J = 2 \nabla \sqrt{\rho} \wedge \Lambda, \quad a.e. (t, x) \in [0, T] \times \mathbf{R}^d.$$

Therefore by definition (ρ, J) is a weak solution to the Cauchy problem of the QHD system. \square

REFERENCES

- [1] P. Antonelli, L.E. Hientzsch, P. Marcati, H. Zheng *On some results for quantum hydrodynamical models*, Mathematical Analysis in Fluid and Gas Dynamics, Proceeding RIMS Kôkyûroku **2070** (2018), 107–129.
- [2] P. Antonelli, P. Marcati, *On the finite energy weak solutions to a system in Quantum Fluid Dynamics*, Comm. Math. Phys. **287** (2009), no 2, 657–686.
- [3] P. Antonelli, P. Marcati, *The Quantum Hydrodynamics system in two space dimensions*, Arch. Rat. Mech. Anal. **203** (2012), 499–527.
- [4] P. Antonelli, P. Marcati, *Quantum hydrodynamics with nonlinear interactions*, Disc. Cont. Dyn. Sys. Ser. S **9**, no. 1 (2016), 1–13.
- [5] P. Antonelli, P. Marcati, *Some results on systems for quantum fluids*, (2016), Recent Advances in Partial Differential Equations and Application, Cont. Math. **666**, 41–54.
- [6] P. Antonelli, P. Marcati, H. Zheng, *Genuine Hydrodynamic Analysis to the 1-D QHD system: Existence, Dispersion and Stability*, preprint <https://arxiv.org/abs/1910.08104>
- [7] P. Antonelli, S. Spirito, *On the compactness of finite energy weak solutions to the quantum Navier-Stokes equations*, J. Hyperbolic Differ. Equ., **15** (2018), 133–147.
- [8] P. Antonelli, S. Spirito, *Global existence of finite energy weak solutions of quantum Navier-Stokes equations*, Arch. Ration. Mech. Anal., **3** (2017), 1161–1199.
- [9] P. Antonelli, S. Spirito, *On the compactness of weak solutions to the Navier-Stokes-Korteweg equations for capillary fluids*, Nonlin. Anal. **187** (2019), 110–124.
- [10] P. Antonelli, S. Spirito, *Global existence of weak solutions to the Navier-Stokes-Korteweg equations*, preprint <https://arxiv.org/abs/1903.02441>.
- [11] C. Audiard, B. Haspot, *Global well-posedness of the Euler-Korteweg system for small irrotational data*, Comm. Math. Phys. **351**, no. 1 (2017), 2017–247.
- [12] J. Barab, *Nonexistence of asymptotically free solutions for a nonlinear Schrödinger equation*, J. Math. Phys. **25** (1984), 3270.
- [13] I. Bialynicki-Birula, Z. Bialynicka-Birula, C. Sliwa, *Motion of vortex lines in quantum mechanics*, Physical Review A **61**, (1999), 032110.
- [14] C. Bardos, P. Degond, *Global existence for the Vlasov-Poisson equation in 3 space variables with small initial data*, Ann. I.H.P. sect. C **2**, no. 2 (1985), 101–118.
- [15] S. Benzoni-Gavage, *Propagating phase boundaries and capillary fluids*, available online at <http://math.univ-lyon1.fr/~benzoni/Levico.pdf>.
- [16] H. Brezis, T. Gallouet, *Nonlinear Schrödinger evolution equations*, Nonlinear Analysis: Theory, Methods and Applications **4** (1980), 677–681.
- [17] S. Benzoni-Gavage, R. Danchin, S. Descombes, *On the well-posedness for the Euler-Korteweg model in several space dimensions*, Indiana Univ. Math. J. **56** (2007), 1499–1579.
- [18] F. Bethuel, R. Jerrard, D. Smets, *On the NLS dynamics for infinite energy vortex configurations on the plane*, Rev. Mat. Iberoamericana **24**, no. 2 (2008), 671–702.
- [19] T. Cazenave, *Semilinear Schrödinger Equations*. Courant Lecture Notes in Mathematics vol. 10, New York University, Courant Institute of Mathematical Sciences, AMS, 2003.
- [20] R. Carles, R. Danchin, J.-C. Saut, *Madelung, Gross-Pitaevskii and Korteweg*, Nonlinearity **25** (2012), 2843–2873.
- [21] J.Y. Chemin, *Dynamique des gaz à masse totale finie*, Asympt. Anal. **3** (1990), 215–220.
- [22] J. Colliander, M. Grillakis, N. Tzirakis, *Tensor products and correlation estimates with applications to nonlinear Schrödinger equations*, Comm. Pure Appl. Math. **62**, no. 7 (2009), 920–968.
- [23] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on \mathbf{R}^3* , Comm. Pure Appl. Math. **57**, no. 8 (2004), 987–1014.
- [24] F. Dalfovo, S. Giorgini, L. Pitaevskii, S. Stringari, *Theory of Bose-Einstein condensation in trapped gases*, Rev. Mod. Phys., **71** (1999), 463–512.
- [25] T. Debiec, P. Gwiazda, A. Swierczewska-Gwiazda, A. Tzavaras, *Conservation of energy for the Euler-Korteweg equations*, Calc. Var. and PDEs **57** (2018), no.6.

- [26] D. Donatelli, E. Feireisl, P. Marcati, *Well/ill posedness for the Euler- Korteweg-Poisson system and related problems*, Communications in Partial Differential Equations, **40** (2015), 1314–1335.
- [27] L. C. Evans, *Partial differential equations - second edition*, American Mathematical Society, Providence, R.I., (2010)
- [28] D. Fajman, J. Joudioux, J. Smulevici, *A vector field method for relativistic transport equations with applications*, Anal. PDE **10**, no. 7 (2017), 1539–1612.
- [29] R.P. Feynman, *Application of quantum mechanics to liquid helium*, Progr. Low Temp. Phys. **1** (1955), 17–53.
- [30] C. Gardner, *The quantum hydrodynamic model for semiconductor devices*, SIAM J. Appl. Math. **54** (1994), 409–427.
- [31] I. Gasser, P. Markowich, *Quantum hydrodynamics, Wigner transforms and the classical limit*, Asympt. Anal. **14** (1997), 97–116.
- [32] J. Giesselmann, C. Lattanzio, A. Tzavaras, *Relative energy for the Korteweg theory and related Hamiltonian flows in gas dynamics*, Arch. Ration. Mech. Anal. **223** (2017), no. 3, 1427–1484.
- [33] J. Giesselmann, A. Tzavaras, *Stability properties of the Euler-Korteweg system with non-monotone pressures*, Appl. Anal. **96** (2017), no. 9, 1528–1546.
- [34] J. Ginibre, G. Velo, *On a class of nonlinear Schrödinger equations. II. Scattering theory, general case*, J. Funct. Anal. **32** (1979), 33–71.
- [35] J. Ginibre, G. Velo, *Quadratic Morawetz inequalities and asymptotic completeness in the energy space for nonlinear Schrödinger and Hartree equations*, Quart. Appl. Math. **68**, no. 1 (2010), 113–134.
- [36] R. Glassey, *On the blowing up of solutions to the Cauchy problem for nonlinear Schrödinger equations*, J. Math. Phys. **18**, no. 9 (1977), 1794–1799.
- [37] F. Haas, *Quantum plasmas: An hydrodynamic approach*, New York: Springer.
- [38] N. Hayashi, K. Nakamitsu, M. Tsutsumi, *On solutions of the initial value problem for the nonlinear Schrödinger equations in one space dimension*, Math. Z. **192** (1986), 637–650.
- [39] N. Hayashi, K. Nakamitsu, M. Tsutsumi, *On solutions of the initial value problem for the nonlinear Schrödinger equations*, J. Funct. Anal. **71** (1987), 218–245.
- [40] R. Illner, G. Rein, *Time decay of the solutions of the Vlasov-Poisson system in the plasma physical case*, Math. Methods Appl. Sci. **6** (1984), 1409–1413.
- [41] F. Huang, H.-L. Li, A. Matsumura, *Existence and stability of steady-state of one-dimensional quantum hydrodynamic system for semiconductors*, J. Diff. Equ. **225** (2006), 1–25.
- [42] F. Huang, H.-L. Li, A. Matsumura, S. Odanaka, *Well-posedness and stability of quantum hydrodynamics of semiconductors in \mathbb{R}^3* , in “Some problems on nonlinear hyperbolic equations and applications”, Ser. Contemp. Appl. Math. CAM, vol. 15, p. 131–160. Higher Ed. Press, Beijing, 2010.
- [43] A. Jüngel, *Dissipative quantum fluid models*, Riv. Mat. Univ. Parma **3** (2012), 217–290.
- [44] A. Jüngel, *Global weak solutions to compressible Navier-Stokes equations for quantum fluids*, SIAM J. Math. Anal. **42**, no. 3 (2010), 1025–1045.
- [45] A. Jüngel, H.-L. Li, A. Matsumura, *The relaxation-time limit in the quantum hydrodynamic equations for semiconductors*, J. Diff. Equ. **225**, no. 2 (2006), 440–464.
- [46] A. Jüngel, J.L. López, J. Montejó-Gámez, *A new derivation of the quantum Navier-Stokes equations in the Wigner-Fokker-Planck approach*, J. Stat. Phys. **145**, no. 6 (2011), 1661–1673.
- [47] A. Jüngel, M.C. Mariani, D. Rial, *Local existence of solutions to the transient quantum hydrodynamic equations*, Math. Mod. Meth. Appl. Sci. **12** (2002), 485.
- [48] I. Khalatnikov, *An Introduction to the Theory of Superfluidity*, 2000.
- [49] S. Klainerman, *Uniform decay estimates and the Lorentz invariance of the classical wave equation*, Comm. Pure Appl. Math. **38**, no. 3 (1985), 321–332.
- [50] L. Landau, *Theory of the Superfluidity of Helium II*, Phys. Rev. **60**, 356 (1941).
- [51] I. Lacroix-Violet, A. Vasseur, *Global weak solutions to the compressible quantum Navier-Stokes and its semi-classical limit*, preprint arxiv.org/abs/1607.06646.
- [52] H.-L. Li, J. Li, Z. Xin, *Vanishing of vacuum states and blow-up phenomena of the compressible Navier-Stokes equations*, Commun. Math. Phys. **281** (2008) 401–444.
- [53] H.-L. Li, P. Marcati, *Existence and asymptotic behavior of multi-dimensional quantum hydrodynamic model for semiconductors*, Comm. Math. Phys. **245** (2004), 215–247.

- [54] J. Li, Z. Xin, *Global existence of weak solutions to the barotropic compressible Navier-Stokes flows with degenerate viscosities*, archived as arxiv.org/abs/1504.06826
- [55] E. Lieb, M. Loss, *Analysis*, Graduate Studies in Mathematics, vol. 14, AMS, 2001.
- [56] F. Linares and G. Ponce, *Introduction to nonlinear dispersive equations*. Springer-Verlag, New York, 2009.
- [57] J-E. Lin and W. A. Strauss, *Decay and scattering of solutions of a nonlinear Schrödinger equation*. Journal of Functional Analysis **30**, Issue 2, (1978), 245-263.
- [58] B. Lü, R. Zhang, X. Zhong, *Global existence of weak solutions to the compressible quantum Navier-Stokes equations with degenerate viscosity*, J. Math. Phys. **60**, no. 12 (2019), 121502.
- [59] E. Madelung, *Quantentheorie in hydrodynamischer form*, Z. Physik **40** (1927), 322.
- [60] P. Markowich, J. Sierra, *Non-uniqueness of weak solutions of the Quantum-Hydrodynamic system*, Kin. Rel. Models **12**, no. 2 (2019), 347-356.
- [61] M.R. Matthews, B.P. Anderson, P.C. Haljain, D.S. Hall, C.E. Wieman, E.A. Cornell, *Vortices in a Bose-Einstein condensate*, Phys. Rev. Lett. **83** (1999), 2498.
- [62] C. Morawetz, *Time decay for the nonlinear Klein-Gordon equations*, Proc. Roy. Soc. Ser. A **306** (1968), 291-296.
- [63] L. Onsager, *Statistical hydrodynamics*, Nuovo Cimento Suppl. **6** (1949), 249-250.
- [64] Y. Ovchinnikov, I. M. Sigal *Ginzburg-Landau Equation I. Static Vortices**, CRM Proc. Lecture Notes. **12** (1998).
- [65] L. Pitaevskii *Vortex lines in an imperfect Bose gas*, Sov. Phys. JETP. **13** (1961), 451-454.
- [66] B. Perthame, *Time decay, propagation of low moments and dispersive effects for kinetic equations*, Comm. PDEs **21** (1996), 659-686.
- [67] L. Pitaevskii, S. Stringari, *Bose-Einstein condensation and superfluidity*, Clarendon Press, Oxford, 2016.
- [68] F. Planchon, N. Tzvetkov, N. Visciglia, *On the growth of Sobolev norms for NLS on 2-and 3-dimensional manifolds*, Anal. PDEs **10**, no. 5 (2017), 1123-1147.
- [69] F. Planchon, L. Vega, *Bilinear virial identities and applications*, Ann. Sci. E.N.S., Série 4 **42**, no. 2 (2009), 261-290.
- [70] P. H. Roberts, J. Grant, *Motions in a Bose condensate I. The structure of the large circular vortex*, Journal of Physics A: General Physics **4** (1971), 55-72.
- [71] D. Serre, *Solutions classiques globales des équations d'Euler pour un fluide parfait compressible*, Ann. Inst. Four. **47**, no. 1 (1997), 139-153.
- [72] T. Sideris, *Formation of singularities in three-dimensional compressible fluids*, Comm. Math. Phys. **101**, no. 4 (1985), 475-485.
- [73] W. Strauss, *Nonlinear invariant wave equations*, Invariant wave equations (Proc. "Ettore Majorana" Internat. School of Math. Phys., Erice, 1977), pp. 197-249, Lecture Notes in Phys., 73, Springer, Berlin-New York, 1978.
- [74] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*. CBMS Regional Conference Series in Mathematics, AMS 2006.
- [75] M. Tsubota, *Quantized Vortices in Superfluid Helium and Bose-Einstein Condensates*, J. Phys.: Conf. Ser. **31** (2006) 88-94.

GRAN SASSO SCIENCE INSTITUTE, VIALE FRANCESCO CRISPI, 7, 67100 L'AQUILA
E-mail address: paolo.antonelli@gssi.it

GRAN SASSO SCIENCE INSTITUTE, VIALE FRANCESCO CRISPI, 7, 67100 L'AQUILA
E-mail address: pierangelo.marcatti@gssi.it

GRAN SASSO SCIENCE INSTITUTE, VIALE FRANCESCO CRISPI, 7, 67100 L'AQUILA
E-mail address: hao.zheng@gssi.it