

ON THE STABILITY OF TANGENT BUNDLES ON CYCLIC COVERINGS

YONGMING ZHANG

ABSTRACT. Let Y be a smooth projective surface defined over an algebraically closed field k with char $k \nmid n$, and let $\pi : X \rightarrow Y$ be a n -cyclic covering branched along a smooth divisor B . We show that under some conditions \mathcal{T}_X is semi-stable with respect to $\pi^*\mathcal{H}$ if the tangent bundle \mathcal{T}_Y is semi-stable with respect to some ample line bundle \mathcal{H} on Y .

1. INTRODUCTION

In this paper we correct some results in [9] and improve them to cyclic coverings. Let $\pi : X \rightarrow Y$ be a finite covering defined over k . We know that the pull back of semi-stable sheaves under π^* are semi-stable in characteristic zero (see [3, lemma 1.1]), but it does not generally hold that the pull back of stable sheaves are stable. In some special cases, the pull back of stable sheaves are stable:

Theorem 1.1. (Theorem 3.2) *Let Y be a smooth projective variety over k with char $k \nmid n$ and let $\pi : X \rightarrow Y$ be a n -cyclic covering branched along a smooth effective divisor $B \in \Gamma(Y, \mathcal{L}^{\otimes n})$ for some line bundle $\mathcal{L} \in \text{Pic}(Y)$. If \mathcal{F} is locally free sheaf on Y and (semi-)stable with respect to an ample line bundle \mathcal{H} on Y then $\pi^*\mathcal{F}$ is (semi-)stable with respect to the ample line bundle $\pi^*\mathcal{H}$.*

Then we study the stability of tangent bundle on a surface which is a n -cyclic covering of another surface and obtain the following results.

Theorem 1.2. (Theorem 3.4) *Let $\pi : X \rightarrow Y$ be a n -cyclic covering determined by a line bundle $L \in \text{Pic}(Y)$ and a smooth divisor $B \in \Gamma(Y, L^{\otimes n})$, where Y is a smooth projective surface over an algebraically closed field k with char $k \nmid n$. And let $\mathcal{H} \in \text{Pic}(Y)$ be an ample line bundles satisfying $\mathcal{H} \equiv lB$ for some $0 < l \in \mathbb{Q}$. Assume that the inequality holds: $n \deg \Omega_Y + (n+1) \deg B \geq 0$ and the cotangent bundle Ω_Y is semi-stable with respect to \mathcal{H} . Then Ω_X is semi-stable with respect to $\pi^*\mathcal{H} \in \text{Pic}(X)$.*

Theorem 1.3. (Theorem 3.5) *With the same conditions in Theorem 1.2, if the strict inequality holds: $n \deg \Omega_Y + (n+1) \deg B > 0$ and the cotangent bundle Ω_Y is semi-stable with respect to \mathcal{H} , then Ω_X is stable with respect to $\pi^*\mathcal{H} \in \text{Pic}(X)$.*

Using this result we find a class of surfaces whose tangent bundle are stable:

Corollary 1.4. (Corollary 3.7) *Let k be an algebraically closed field with char $k \nmid n$ and $\pi : X \rightarrow \mathbb{P}_k^2$ be a n -cyclic covering branched along a smooth curve B in \mathbb{P}_k^2 . Then Ω_X is semi-stable with respect to $\pi^*\mathcal{O}_{\mathbb{P}^N}(1)$, and it is stable with respect to $\pi^*\mathcal{O}_{\mathbb{P}^N}(1)$ when $n = 2$ and $\deg B \geq 4$ or $n > 2$.*

In addition, we already know that the tangent bundle of a K3 surface of characteristic zero is stable (see Chapter 9 of [5]). By the above corollary we obtain a partial result about the stability of the tangent bundle of a K3 surface in positive characteristic: when char $k \neq 2$, the tangent bundle of a K3 surface, which can be realized as a double covering of \mathbb{P}^2 branched along a smooth curve of degree 6 or a 4-cyclic covering of \mathbb{P}^2 branched along a smooth curve of degree 1, is stable.

2. PRELIMINARIES

2.1. cyclic coverings. Let k denote an algebraically closed field with char $k \nmid n$. First we recall the definition of n -cyclic coverings. Let Y be a smooth variety over k and let B be an effective divisor

on Y . Suppose that $\mathcal{O}_Y(B)$ is divisible by n in $\text{Pic}(Y)$, i.e. there is a line bundle $\mathcal{L} \in \text{Pic}(Y)$ such that $\mathcal{L}^{\otimes n} = \mathcal{O}_Y(B)$. Note that the divisor B gives a morphism $\mathcal{L}^{\otimes -n} \rightarrow \mathcal{O}_Y$, and this morphism gives $\mathcal{E} = \mathcal{O}_Y \oplus \mathcal{L}^{-1} \oplus \dots \oplus \mathcal{L}^{-n+1}$ a ring structure. Let $X = \text{Spec}(\mathcal{E})$, then with the canonical projection we get a n -cyclic covering over Y :

$$\pi : X \longrightarrow Y.$$

Let B_1 denote the reduced divisor of the inverse image $\pi^{-1}(B)$. Then we have the following lemma:

Lemma 2.1. ([1, 1.17]) *Keep the above notation. Then we have:*

- (1) $\pi^*\mathcal{L} = \mathcal{O}_X(B_1)$;
- (2) $K_X = \pi^*(K_Y \otimes \mathcal{L}^{\otimes(n-1)})$;
- (3) $\pi_*\mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{L}^{-1} \oplus \dots \oplus \mathcal{L}^{\otimes(1-n)}$.

2.2. Stability. We collect here some well known facts about stability of coherent sheaves. Let X be a smooth projective variety of dimension n over an algebraically closed field k and \mathcal{H} an ample line bundle on X . Let E be a torsion free sheaf on X . The rank of E , denoted by $\text{rank } E$, is the rank of E at the generic point of X . The first Chern class of E is denoted by $c_1(E)$, then the degree of E , denoted by $\text{deg } E$, is the intersection number $c_1(E) \cdot \mathcal{H}^{n-1}$. We define:

$$\mu(E) := \frac{\text{deg } E}{\text{rank } E}$$

which depends on the choice of the ample line bundle \mathcal{H} on X .

Definition 2.2. E is stable (resp. semi-stable) if for any sub-sheaf F with $0 < \text{rank } F < \text{rank } E$, we have:

$$\mu(F) < \mu(E) \quad (\text{resp. } \mu(F) \leq \mu(E)).$$

Actually, in the above definition we only need to consider the saturated sub-sheaves of E . If E is a torsion free sheaf on X , then there is a unique filtration (the Harder-Narasimhan filtration)

$$0 = E_0 \subsetneq E_1 \subsetneq \dots \subsetneq E_l = E$$

such that the factors $gr_i^{HN} := E_i/E_{i-1}$ for $i = 1, \dots, l$ are semi-stable sheaves and

$$\mu_{max}(E) := \mu(gr_1^{HN}) > \dots > \mu(gr_l^{HN}) =: \mu_{min}(E).$$

The instability of E is defined as $I(E) := \mu_{max}(E) - \mu_{min}(E)$, and E is semi-stable if and only if $I(E) = 0$.

If E is a semi-stable torsion free sheaf on X , then there is a filtration (the Jordan-Hölder filtration)

$$0 = E_0 \subsetneq E_1 \subsetneq \dots \subsetneq E_l = E$$

such that the factors $gr_i = E_i/E_{i-1}$ for $i = 1, \dots, l$ are stable sheaves and

$$\mu(gr_1) = \dots = \mu(gr_l) = \mu(E).$$

Note that the Jordan-Hölder filtration is not unique but the factors $gr_i := E_i/E_{i-1}$ for $i = 1, \dots, l$ are unique determined by E .

3. THE MAIN RESULTS AND PROOF

3.1. stability of the pull back of stable sheaves. For a Galois covering $\pi : X \rightarrow Y$ with Galois group G and coherent sheaf \mathcal{E} on Y , then there is an isomorphism $\sigma^*\pi^*\mathcal{E} \simeq \pi^*\mathcal{E}$. For any sub-sheaf $\mathcal{F} \subset \pi^*\mathcal{E}$ we will denote the image of sub-sheaf $\sigma^*\mathcal{F} \subset \sigma^*\pi^*\mathcal{E}$ under this isomorphism by \mathcal{F}^σ for $\sigma \in G$.

Next we relate the following fact in Galois decent theory:

Theorem 3.1. ([2, Theorem 28, Chapte 9]) *Let $p : X \rightarrow Y$ be a finite flat Galois morphism with Galois group G , and suppose that X and Y are both integral. Let \mathcal{F} be a locally free sheaf on Y and suppose that \mathcal{G}' is a coherent sub-sheaf of $p^*\mathcal{F}$ such that:*

- *For all $\sigma \in G$, $(\mathcal{G}')^\sigma = \mathcal{G}'$ as a sub-sheaf of $p^*\mathcal{F}$;*
- *$p^*\mathcal{F}/\mathcal{G}'$ is torsion free.*

Then there exists a coherent sub-sheaf \mathcal{G} of \mathcal{F} such that $\mathcal{G}' = p^\mathcal{G}$ as a sub-sheaf of $p^*\mathcal{F}$*

Immediately, we have the following observation:

Theorem 3.2. *Let Y be a smooth projective variety over k with $\text{char } k \nmid n$ and let $\pi : X \rightarrow Y$ be a n -cyclic covering branched along a smooth effective divisor $B \in \Gamma(Y, \mathcal{L}^{\otimes n})$ for some line bundle $\mathcal{L} \in \text{Pic}(Y)$. If \mathcal{F} is locally free sheaf on Y and (semi-)stable with respect to an ample line bundle \mathcal{H} on Y then $\pi^*\mathcal{F}$ is (semi-)stable with respect to the ample line bundle $\pi^*\mathcal{H}$.*

Proof. Let $G = \{1, \sigma, \dots, \sigma^{n-1}\}$ be the Galois group acting on X . We first prove $\pi^*\mathcal{F}$ is semi-stable. Suppose $\pi^*\mathcal{F}$ is not semi-stable. then there is a unique maximal destabilizing sub-sheaf

$$\mathcal{M} \subset \pi^*\mathcal{F} \quad \text{with} \quad \mu(\mathcal{M}) > \mu(\pi^*\mathcal{F}).$$

Note that \mathcal{M} is a saturated sub-sheaf and $\mathcal{M}^\sigma = \mathcal{M}$. By Theorem 3.1, $\mathcal{M} = \pi^*\mathcal{G}$ for some sub-sheaf $\mathcal{G} \subset \mathcal{F}$. Since $\mu(\pi^*\mathcal{G}) = \deg(\pi) \mu(\mathcal{G})$ and $\mu(\pi^*\mathcal{F}) = \deg(\pi) \mu(\mathcal{F})$ we get $\mu(\mathcal{G}) > \mu(\mathcal{F})$ which is a contradiction.

Second, we suppose $\pi^*\mathcal{F}$ is not stable. Then there is a filtration (the Jordan-Hölder filtration)

$$0 = \mathcal{F}_0 \subsetneq \mathcal{F}_1 \subsetneq \dots \subsetneq \mathcal{F}_l = \pi^*\mathcal{F}$$

such that the factors $gr_i = \mathcal{F}_i/\mathcal{F}_{i-1}$ for $i = 1, \dots, l$ ($l \geq 2$) are stable sheaves and

$$\mu(gr_1) = \dots = \mu(gr_l) = \mu(\pi^*\mathcal{F}).$$

There is no saturated sub-sheaf $\mathcal{E} \subset \pi^*\mathcal{F}$ with $\mu(\mathcal{E}) \geq \mu(\pi^*\mathcal{F})$ and $\mathcal{E}^\sigma = \mathcal{E}$ (if not then $\mathcal{E} = \pi^*\mathcal{E}'$ for some sub-sheaf $\mathcal{E}' \subset \mathcal{F}$ and by the argument as before we will get a contradiction), so $\mathcal{F}_i^\sigma \neq \mathcal{F}_i$ for $i < l$. Using the fact that the only nontrivial map between two stable sheaves with the same slope is an isomorphism, we have $l \mid n$,

$$\pi^*\mathcal{F} = gr_1 \oplus \dots \oplus gr_l \quad \text{and} \quad gr_1 \simeq \dots \simeq gr_l.$$

More over G acts on those factors transversally. In fact, after the action of σ we get another Jordan-Hölder filtration:

$$0 = \mathcal{F}_0^\sigma \subsetneq \mathcal{F}_1^\sigma \subsetneq \dots \subsetneq \mathcal{F}_l^\sigma = \pi^*\mathcal{F}$$

such that the factors $gr'_i = \mathcal{F}_i^\sigma/\mathcal{F}_{i-1}^\sigma$ for $i = 1, \dots, l$ are stable sheaves and

$$\mu(gr'_1) = \dots = \mu(gr'_l) = \mu(\pi^*\mathcal{F}).$$

Let l_0 be the smallest number satisfies $\mathcal{F}_1^\sigma \subset \mathcal{F}_{l_0}$, then we get a nontrivial morphism between stable sheaves with the same slope: $\mathcal{F}_1^\sigma \rightarrow gr_{l_0} = \mathcal{F}_{l_0}/\mathcal{F}_{l_0-1}$, so it is an isomorphism, and $\mathcal{F}_1^\sigma \cap \mathcal{F}_1 = 0$. It follows that: $\mathcal{F}_1^\sigma \oplus \mathcal{F}_1 \subseteq \pi^*\mathcal{F}$. If we compare the original filtration with the filtration obtained by the action of σ^2 it we will get $\mathcal{F}_1^{\sigma^2} \oplus \mathcal{F}_1^\sigma \oplus \mathcal{F}_1 \subseteq \pi^*\mathcal{F}$. Let us do this process and we will get the result finally.

On the other hand, after pushing forward we obtain:

$$\pi_*\pi^*(\mathcal{F}) = \mathcal{F} \oplus \mathcal{F}(\mathcal{L}^{-1}) \oplus \dots \oplus \mathcal{F}(\mathcal{L}^{-n+1}) = \pi_*gr_1 \oplus \pi_*gr_2 \oplus \dots \oplus \pi_*gr_l.$$

Note that σ acts on $\mathcal{F}(\mathcal{L}^{-i})$ by multiplying ξ^i where ξ is the n -th root of unity, and on π_*gr_1 , $i = 1, \dots, l$ transversally. Consider the map $\pi_*gr_1 \rightarrow \mathcal{F}(\mathcal{L}^{-i})$, $a \mapsto \sum_{j=0}^{l-1} (\sigma^j)^* \xi^{ij} a$ at the generic point we will find that $\text{rank } \pi_*gr_1 = \text{rank } \mathcal{F}$ and $\pi_*gr_1 \cap \mathcal{F} \oplus \mathcal{F}(\mathcal{L}^{-1}) \oplus \dots \oplus \mathcal{F}(\mathcal{L}^{-n+2}) = 0$, hence we can get an injection:

$$\pi_*gr_1 \hookrightarrow \mathcal{F}(\mathcal{L}^{-n+1}),$$

whose quotient is a torsion sheaf. So we will get

$$\mu(\pi_*gr_1) \leq \mu(\mathcal{F}(\mathcal{L}^{-n+1})) < \mu(\mathcal{F} \oplus \mathcal{F}(\mathcal{L}^{-1}) \oplus \cdots \oplus \mathcal{F}(\mathcal{L}^{-n+1})) = \mu(\pi_*\pi^*(\mathcal{F})),$$

which contradicts to $\mu(\pi_*\pi^*(\mathcal{F})) = \mu(\pi_*gr_1)$. So $\pi^*\mathcal{F}$ is stable. \square

3.2. sub-sheaves in cotangent bundle. Let $\pi : X \rightarrow Y$ be a n -cyclic covering determined by a line bundle $L \in \text{Pic}(Y)$ and a smooth divisor $B \in \Gamma(Y, L^{\otimes n})$, where Y is a smooth projective variety over an algebraically closed field k with $\text{char } k \nmid n$. Let B_1 denote the reduced divisor of the inverse image $\pi^{-1}(B)$.

Now we consider a saturated sub-sheaf $\mathcal{M}_1 \subset \Omega_X$ which is stable under action of σ .

Consider the following diagram

$$(3.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{N}_2 & \longrightarrow & \mathcal{M}_2 & \longrightarrow & \mathcal{Q}_2 \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & \pi^*\Omega_Y & \longrightarrow & \Omega_X & \longrightarrow & \Omega_{X/Y} \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & \mathcal{N}'_1 & \longrightarrow & \mathcal{M}_1 & \xrightarrow{\varphi} & \mathcal{Q}_1 \longrightarrow 0 \end{array}$$

where $\mathcal{N}'_1 = \pi^*\Omega_Y \cap \mathcal{M}_1 \subset \pi^*\Omega_Y$ is also saturated and stable under the action of σ . By lemma 3.1 there is a sub-sheaf $N'_1 \subset \Omega_Y$ such that

$$\pi^*N'_1 = \mathcal{N}'_1.$$

Let (x_1, x_2, \dots, x_m) be local coordinates on Y , centred at a point $x \in Y$, such that B is defined by $x_1 = 0$. Then $(t, \pi^*x_2, \dots, \pi^*x_m)$ is local coordinates on X , centred at $\pi^{-1}(x) \in X$, where t is the vertical coordinate in L satisfying $\pi^*x_1 - t^n = 0$. Locally any element of \mathcal{M}_1 should look like

$$t^{n-1} \sum_{i \geq 2} \pi^*b_{n-1,i} d\pi^*x_i + \cdots + t \sum_{i \geq 2} \pi^*b_{1,i} d\pi^*x_i + \pi^*cdt, \text{ where } b_{j,i}, c \in \mathcal{O}_Y,$$

where σ acts on it by multiplying n -th unit root ξ on t and dt .

Since \mathcal{M}_1 is stable under action of σ , locally it can always be generated by 1-forms of the two kinds

$$\sum_{i \geq 2} \pi^*a_i d\pi^*x_i, \text{ where } a_i \in \mathcal{O}_Y$$

and

$$t \sum_{i \geq 2} \pi^*b_i d\pi^*x_i + \pi^*cdt, \text{ where } b_i, 0 \neq c \in \mathcal{O}_Y.$$

We will call them the first and second kind generators respectively. It is easy to find that $\mathcal{N}'_1 = \mathcal{M}_1$ if and only if \mathcal{M}_1 can be generated by the first kind generators only.

If $\mathcal{M}_1 \neq \mathcal{N}'_1$ then its generators should contain the second kind. And by some local calculation we find that \mathcal{Q}_1 is a sub-sheaf of $\Omega_{X/Y}$ of rank $n-1$ as \mathcal{O}_{B_1} -module. In this case we consider the following diagram

$$(3.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & I_B/I_B^2 & \longrightarrow & \Omega_Y|_B & \longrightarrow & \Omega_B \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & I_B/I_B^2 \cap \mathcal{N}'_1|_B & \longrightarrow & \mathcal{N}'_1|_B & \longrightarrow & \mathcal{G} \longrightarrow 0. \end{array}$$

First, Multiplying a second kind generators of \mathcal{M}_1 by t^{n-1} ,

$$t^{n-1} [t \sum_{i \geq 2} \pi^*b_i d\pi^*x_i + \pi^*cdt] = \pi^*x_1 \sum_{i \geq 2} \pi^*b_i d\pi^*x_i + \frac{1}{n} \pi^*cd\pi^*x_1,$$

we will get a element in $I_B/I_B^2 \cap \mathcal{N}'_1|_B$ under the restriction $\Omega_Y \rightarrow \Omega_Y|_B$. Conversely, for any element in $I_B/I_B^2 \cap \mathcal{N}'_1|_B$, we could get an element in \mathcal{N}_1 which can factorize as a second kind generator and t^{n-1} . Since \mathcal{M}_1 is saturated, \mathcal{M}_1 would have the second kind generators. So we get the following observation.

Lemma 3.3. \mathcal{M}_1 is not contained in $\pi^*\Omega_Y$ if and only if

$$\mathcal{N}'_1|_B \cap I_B/I_B^2 \neq 0.$$

And in this case, $\text{rank } Q_1 = n - 1$ as \mathcal{O}_{B_1} -module.

3.3. stability of tangent bundle on cyclic coverings.

Theorem 3.4. Let $\pi : X \rightarrow Y$ be a n -cyclic covering determined by a line bundle $L \in \text{Pic}(Y)$ and a smooth divisor $B \in \Gamma(Y, L^{\otimes n})$, where Y is a smooth projective surface over an algebraically closed field k with $\text{char } k \nmid n$. And let $\mathcal{H} \in \text{Pic}(Y)$ be an ample line bundles satisfying $\mathcal{H} \equiv lB$ for some $0 < l \in \mathbb{Q}$. Assume that the inequality holds: $n \deg \Omega_Y + (n+1) \deg B \geq 0$ and the cotangent bundle Ω_Y is semi-stable with respect to \mathcal{H} . Then Ω_X is semi-stable with respect to $\pi^*\mathcal{H} \in \text{Pic}(X)$.

Proof. Suppose that Ω_X is not semi-stable, then there is a unique maximally destabilizing sub-line bundle $\mathcal{M}_1 \subset \Omega_X$ which is saturated and stable under the action of σ .

If $\mathcal{M}_1 \subset \pi^*\Omega_Y \subset \Omega_X$, then by lemma 3.2 $\pi^*\Omega_Y$ is semi-stable and $\mu(\mathcal{M}_1) \leq \mu(\pi^*\Omega_Y) < \mu(\Omega_X)$ which contradicts the property of the maximally destabilizing sub-sheaves $\mathcal{M}_1 \subset \Omega_X$.

Now we have $\mathcal{M}_1 \not\subset \pi^*\Omega_Y$. By lemma 3.3 $\text{rank } I_B/I_B^2 \cap \mathcal{N}'_1|_B = \text{rank } \mathcal{N}'_1 = 1$ as \mathcal{O}_B -mod. So $\mathcal{G} \subset \Omega_B$ is a torsion sheaf as \mathcal{O}_B -mod and it is zero. Therefore we have

$$\mathcal{N}'_1|_B \subseteq I_B/I_B^2$$

and

$$\deg(\mathcal{N}'_1|_B) \leq \deg(I_B/I_B^2) = \frac{-\deg B}{l}$$

and it is easy to check the following (in)equalities

$$\begin{aligned} \mu(\mathcal{M}_1) &= \deg \mathcal{N}_1 + (n-1) \deg B_1 && \text{(by the last line in diagram 3.2 and lemma 3.3)} \\ &= n \deg \mathcal{N}'_1 + (n-1) \deg B = nl \deg \mathcal{N}'_1|_B + (n-1) \deg B \\ &\leq -\deg B && \text{(by the above inequality)} \\ &\leq \frac{n \deg \Omega_Y + (n-1) \deg B}{2} && \text{(by the condition in the theorem)} \\ &= \mu(\Omega_X) && \text{(by the second line in diagram 3.2 and } \deg B = \deg B_1 \text{).} \end{aligned}$$

We also get a contraction to the assumption. So Ω_X is semi-stable with respect to $\pi^*\mathcal{H}$. \square

Theorem 3.5. With the same conditions in Theorem 3.4, if the strict inequality holds: $n \deg \Omega_Y + (n+1) \deg B > 0$ and the cotangent bundle Ω_Y is semi-stable with respect to \mathcal{H} , then Ω_X is stable with respect to $\pi^*\mathcal{H} \in \text{Pic}(X)$.

Proof. We know that Ω_X is semi-stable by Theorem 3.4. We suppose that Ω_X is not stable with respect to $\pi^*\mathcal{H}$, then there is a Jordan-Hölder filtration:

$$0 \subset \mathcal{M}_1 \subset \Omega_X \quad \text{with} \quad \mu(\mathcal{M}_1) = \mu(\Omega_X).$$

Now we get a saturated sub-line bundle \mathcal{M}_1 . By the same argument as Theorem 3.4 we know that $\mathcal{M}_1^\sigma \neq \mathcal{M}_1$, then we get another saturated stable sub-line bundle \mathcal{M}_1^σ with $\mu(\mathcal{M}_1^\sigma) = \mu(\Omega_X)$. Since the graded factors of Jordan-Hölder filtration are uniquely determined, the filtration splits:

$$\Omega_X = \mathcal{M}_1 \oplus \mathcal{M}_1^\sigma.$$

Now we want to prove that there is no such decomposition. Since π_* is exact we have the following diagram by pushing forward diagram 3.2:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \pi_*\mathcal{N}_2 & \longrightarrow & \pi_*\mathcal{M}_1^\sigma & \longrightarrow & \pi_*Q_2 & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & \pi_*\pi^*\Omega_Y & \longrightarrow & \pi_*\Omega_X & \longrightarrow & \pi_*\Omega_{X/Y} & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & \pi_*\mathcal{N}_1 & \longrightarrow & \pi_*\mathcal{M}_1 & \longrightarrow & \pi_*Q_1 & \longrightarrow & 0.
\end{array}$$

where $\mathcal{N}_1 = \pi^*\Omega_Y \cap \mathcal{M}_1$ and $\mathcal{N}_2 = \pi^*\Omega_Y/\mathcal{N}_1$. Note that

$$\pi_*\mathcal{N}_1 \oplus \pi_*\mathcal{N}_1^\sigma \subseteq \pi_*\pi^*(\Omega_Y) = \Omega_Y \oplus \Omega_Y(\mathcal{L}^{-1}) \cdots \oplus \Omega_Y(\mathcal{L}^{\otimes(-n+1)}).$$

where $\mathcal{N}_1^\sigma = \pi^*\Omega_Y \cap \mathcal{M}_1^\sigma$ and σ acts on $\Omega_Y(\mathcal{L}^{-i})$ by multiplying ξ^i , where ξ is the n -th root of unity, and acts $\pi_*\mathcal{N}_1$ onto $\pi_*\mathcal{N}_1^\sigma$. Consider the sub-sheaf generated by $a + \sigma^*(a)$, $a \in \pi_*\mathcal{N}_1$ whose rank is equal to $\text{rank } \pi_*\mathcal{N}_1$, we have $\text{rank } \Omega_Y = \text{rank } \pi_*\mathcal{N}_1$ and $n = 2$. So we only need to prove that such decomposition does not occur when $n = 2$.

Note that $\pi_*\mathcal{N}_1 \cap \Omega_Y = 0$, and we get an injection:

$$\pi_*\mathcal{N}_1 \hookrightarrow \Omega_Y(\mathcal{L}^{-1})$$

with a nonzero torsion sheaf as its quotient when $n = 2$. So we have

$$\mu(\pi_*\mathcal{N}_1) < \mu(\Omega_Y(\mathcal{L}^{-1})) = \mu(\Omega_Y) - \text{deg}(\mathcal{L}).$$

Now we relate some facts which will be used later. For any coherent sheaf \mathcal{E} on X we have

$$\text{deg}(\pi_*\mathcal{E}) = \frac{\text{rank}(\pi_*\mathcal{E})}{2} \text{deg}(\Omega_Y) - \frac{\text{rank}(\mathcal{E})}{2} \text{deg}(\Omega_X) + \text{deg}(\mathcal{E}),$$

which can be calculated using Grothendieck-Riemann-Roch theorem. Note that $\text{rank}(\pi_*\mathcal{E}) = n \text{rank}(\mathcal{E})$, we have

$$\mu(\pi_*\mathcal{E}) = \mu(\Omega_Y) - \frac{1}{n}\mu(\Omega_X) + \frac{1}{n}\mu(\mathcal{E})$$

if $\text{rank}(\mathcal{E}) > 0$. By Lemma 2.1 ii) we can get $\mu(\Omega_X) = \mu(\pi^*\Omega_Y) + \text{deg}(\pi^*\mathcal{L}^{\otimes(n-1)}) = n\mu(\Omega_Y) + n(n-1)\text{deg}(\mathcal{L})$, so we have

$$\mu(\pi_*\mathcal{E}) = \frac{1}{n}\mu(\mathcal{E}) - (n-1)\text{deg}(\mathcal{L})$$

if $\text{rank}(\mathcal{E}) > 0$.

Next we will get a contradiction by calculating the slope of sheaves.

When $n = 2$, by the above facts we have

$$\mu(\pi_*\mathcal{N}_2) = \frac{1}{2}\mu(\mathcal{N}_2) - \text{deg}(\mathcal{L})$$

and since the quotient Q_2 is torsion, $\mu(\mathcal{N}_2) < \mu(\mathcal{M}_1^\sigma) = \mu(\Omega_X)$. So we have

$$\mu(\pi_*\mathcal{N}_2) < \frac{1}{2}\mu(\Omega_X) - \text{deg}(\mathcal{L}) = \mu(\Omega_Y).$$

finally we conclude that

$$\mu(\pi_*\pi^*(\Omega_Y)) = \frac{1}{2}(\mu(\pi_*\mathcal{N}_1) + \mu(\pi_*\mathcal{N}_2)) < \frac{1}{2}(\mu(\Omega_Y) - \text{deg}(\mathcal{L}) + \mu(\Omega_Y)) = \mu(\Omega_Y) - \frac{1}{2}\text{deg}(\mathcal{L}),$$

but $\mu(\pi_*\pi^*(\Omega_Y)) = \mu(\Omega_Y \oplus \Omega_Y(\mathcal{L}^{-1})) = \mu(\Omega_Y) - \frac{1}{2}\text{deg}(\mathcal{L})$. So there is no such decomposition and Ω_X is stable with respect to $\pi^*\mathcal{H}$. \square

Next, we relate the following well known result.

Theorem 3.6. ([5, 1.4]) *Let k be an algebraically closed field, and $\Omega_{\mathbb{P}^N}^1$ is the cotangent bundle on \mathbb{P}_k^N . Then $\Omega_{\mathbb{P}^N}^1$ is stable with respect to $\mathcal{O}_{\mathbb{P}^N}(1)$.*

Then we get the following result using Theorem 3.6.

Corollary 3.7. *Let k be an algebraically closed field with $\text{char } k \nmid n$ and $\pi : X \rightarrow \mathbb{P}_k^2$ be a n -cyclic covering branched along a smooth curve B in \mathbb{P}_k^2 . Then Ω_X is semi-stable with respect to $\pi^*\mathcal{O}_{\mathbb{P}^N}(1)$, and it is stable with respect to $\pi^*\mathcal{O}_{\mathbb{P}^N}(1)$ when $n = 2$ and $\deg B \geq 4$ or $n > 2$.*

Remark 3.8. In particular, when $n = 2$ and $\deg B = 6$ or $n = 4$ and $\deg B = 4$, X is a K3 surface of degree 2 or 4 respectively. And the cotangent bundle Ω_X is stable with respect to $\pi^*\mathcal{O}_{\mathbb{P}^N}(1)$.

4. APPLICATION

Let X be a smooth projective variety over an algebraically closed field k with $\text{char } k = p > 0$. The absolute Frobenius $F_X : X \rightarrow X$ is defined by the map: $f \mapsto f^p$ in \mathcal{O}_X . Let $F : X \rightarrow X_1 := X \times_k k$ denote the relative Frobenius morphism over k . We know that F_* preserves semi-stability of vector bundles on curves of genus $g \geq 1$. When W is a vector bundle on a higher dimensional variety, the instability of F_*W is bounded by instability of $W \otimes T^l(\Omega_X^1)$ ($0 \leq l \leq n(p-1)$) which is proved by X. Sun in [8] and we will use this result to get our corollary.

Definition 4.1. ([6, 5.1]) Let $V_0 := V = F^*(F_*W)$, $V_1 = \ker(F^*(F_*W) \rightarrow W)$, and $V_{l+1} := \ker(V_l \xrightarrow{\nabla} V \otimes_{\mathcal{O}_X} \Omega_X^1 \rightarrow (V/V_l) \otimes_{\mathcal{O}_X} \Omega_X^1)$ where $\nabla : V \rightarrow V \otimes_{\mathcal{O}_X} \Omega_X^1$ is the canonical connection with zero p -curvature (see [7, Theorem 5.1]). Then we have the filtration

$$0 = V_{n(p-1)+1} \subset V_{n(p-1)} \subset \cdots \subset V_1 \subset V_0 = V = F^*(F_*W).$$

Next, we recall the definition of $T^l(V)$. Let V be a n -dimensional vector space over k with standard representation of $\text{GL}(n)$, S_l be the symmetric group with natural action on $V^{\otimes l}$ by $(v_1 \otimes \cdots \otimes v_l) \cdot \sigma = v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(l)}$ for $v_i \in V$ and $\sigma \in S_l$. Let e_1, \dots, e_n be a basis of V , for $k_i \geq 0$ with $k_1 + \cdots + k_n = l$ then define

$$v(k_1, \dots, k_n) = \sum_{\sigma \in S_l} (e_1^{\otimes k_1} \otimes \cdots \otimes e_n^{\otimes k_n}) \cdot \sigma.$$

Let $T^l(V) \subset V^{\otimes l}$ be the linear subspace generated by all vectors $v(k_1, \dots, k_n)$ for all $k_i \geq 0$ satisfying $k_1 + \cdots + k_n = l$. Then we have the following property.

Theorem 4.2. ([8, Theorem 3.7]) *The filtration defined in Definition 4.1 has the following properties*

- (i) $\nabla(V_{l+1}) \subset V_l \otimes \Omega_X^1$ for $l \geq 1$, and $V_0/V_1 \cong W$.
- (ii) $V_l/V_{l+1} \xrightarrow{\nabla} (V_{l-1}/V_l) \otimes \Omega_X^1$ are injective morphisms of vector bundles for $1 \leq l \leq n(p-1)$, which induced isomorphisms $\nabla^l : V_l/V_{l+1} \cong W \otimes_{\mathcal{O}_X} T^l(\Omega_X^1)$, $0 \leq l \leq n(p-1)$.

Let \mathcal{H} be an ample line bundle on X . For any vector bundle W on X , let

$$I(W, X) := \max\{I(W \otimes T^l(\Omega_X^1)) \mid 0 \leq l \leq n(p-1)\}$$

be the maximal value of instabilities $I(W \otimes T^l(\Omega_X^1))$.

Theorem 4.3. ([8, Theorem 4.8]) *When $K_X \cdot \mathcal{H}^{n-1} \geq 0$, we have, for any $\mathcal{E} \subset F_*W$,*

$$\mu(F_*W) - \mu(\mathcal{E}) \geq -\frac{I(W, X)}{p}.$$

*In particular, if $W \otimes T^l(\Omega_X^1)$, $0 \leq l \leq n(p-1)$, are semi-stable, then F_*W is semi-stable. Moreover, if $K_X \cdot \mathcal{H}^{n-1} > 0$, the stability of the bundles $W \otimes T^l(\Omega_X^1)$, $0 \leq l \leq n(p-1)$, implies the stability of F_*W .*

When X is a surface we have a better description of $T^l(\Omega_X^1)$.

Theorem 4.4. ([8, Theorem 3.8]) *When $\dim(X) = 2$, we have*

$$V_l/V_{l+1} = \begin{cases} W \otimes \text{Sym}^l(\Omega_X^1) & \text{when } l < p \\ W \otimes \text{Sym}^{2(p-1)-l}(\Omega_X^1) \otimes \omega_X^{l-(p-1)} & \text{when } l \geq p \end{cases}$$

Now using the above results we get the following corollary.

Corollary 4.5. *Let $\pi : X \rightarrow \mathbb{P}^2$ be a n -cyclic covering over an algebraically closed field k with $\text{char } k \nmid n$ branched along a smooth curve B of degree d and $\mathcal{H} = \pi^*\mathcal{O}(1)$. Then for any torsion free sheaf \mathcal{L} of rank 1, $F_*\mathcal{L}$ is semi-stable if $d \geq 5$ or $n = 4$ and $d \geq 4$. Moreover, F_*W is stable if $n = 2$ and $d \geq 6$ or $n \geq 3$ and $d \geq 5$.*

Proof. First, by Lemma 2.1 we know that $K_X \cdot \mathcal{H} \geq 0$ if $d \geq 5$ or $n = 4$ and $d \geq 4$ and $K_X \cdot \mathcal{H} > 0$ if $n = 2$ and $d \geq 6$ or $n \geq 3$ and $d \geq 5$. And we have proved that Ω_X^1 is stable in Corollary 3.7, so $\text{Sym}^l(\Omega_X^1)$ is semi-stable when $l < p$. Then $T^l(\Omega_X^1)$ is semi-stable by Theorem 4.4 and $I(\mathcal{L} \otimes T^l(\Omega_X^1)) = I(T^l(\Omega_X^1)) = 0$ since \mathcal{L} is of rank 1. At last by using Theorem 4.3 we obtain our result. \square

REFERENCES

- [1] W. P. Barth, K. Hulek, C. A. M. Peters, and A. Van de Ven, *Compact complex surfaces*, 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 4, Springer-Verlag, Berlin, 2004. [MR2030225](#) $\uparrow 2$
- [2] R. Friedman, *Algebraic surfaces and holomorphic vector bundles*, Universitext, Springer-Verlag, New York, 1998. [MR1600388](#) $\uparrow 3$
- [3] D. Gieseker, *On a theorem of Bogomolov on Chern classes of stable bundles*, Amer. J. Math. **101** (1979), no. 1, 77–85. [MR527826](#) $\uparrow 1$
- [4] D. Huybrechts, *Lectures on K3 surfaces*, Cambridge Studies in Advanced Mathematics, vol. 158, Cambridge University Press, Cambridge, 2016. [MR3586372](#) \uparrow
- [5] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, 2nd ed., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2010. [MR2665168](#) $\uparrow 1, 7$
- [6] K. Joshi, S. Ramanan, E. Z. Xia, and J.-K. Yu, *On vector bundles destabilized by Frobenius pull-back*, Compos. Math. **142** (2006), no. 3, 616–630. [MR2231194](#) $\uparrow 7$
- [7] N. M. Katz, *Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin*, Inst. Hautes Études Sci. Publ. Math. **39** (1970), 175–232. [MR0291177](#) $\uparrow 7$
- [8] X. Sun, *Direct images of bundles under Frobenius morphism*, Invent. Math. **173** (2008), no. 2, 427–447. [MR2415312](#) $\uparrow 7, 8$
- [9] Y. Zhang, *On the stability of tangent bundle on double coverings*, Acta Math. Sin. (Engl. Ser.) **33** (2017), no. 8, 1039–1047. [MR3668639](#) $\uparrow 1$

E-mail address: zhangyongming@amss.ac.cn

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI, 200433, CHINA