

# Comment on "Direct Observation of Proton Emission in $^{11}\text{Be}$ "

H.O.U. Fynbo,<sup>1</sup> Z. Janas,<sup>2</sup> C. Mazzocchi,<sup>2</sup> M. Pfützner,<sup>2,\*</sup> J. Refsgaard,<sup>3,4</sup> K. Riisager,<sup>1</sup> and N. Sokołowska<sup>2</sup>

<sup>1</sup>*Institut for Fysik og Astronomi, Aarhus Universitet, DK-8000 Aarhus, Denmark*

<sup>2</sup>*Faculty of Physics, University of Warsaw, 02-093 Warszawa, Poland*

<sup>3</sup>*Department of Astronomy and Physics, Saint Mary's University, Halifax, Nova Scotia, B3H 3C3 Canada*

<sup>4</sup>*TRIUMF, 4004 Wesbrook Mall, Vancouver BC, V6T 2A3 Canada*

We argue that conclusions of [PRL 123, 082501 (2019)] are incorrect. The authors present the direct observation of beta-delayed proton emission in the beta decay of  $^{11}\text{Be}$ . From the determined branching ratio for this process and from the energy spectrum of emitted protons the existence of a so far unobserved narrow resonance in  $^{11}\text{B}$  was deduced. The given beta strength for the transition to this state is however wrong. In addition, we show that the combination of peak position and branching ratio is in strong disagreement with models considered by the authors. Furthermore, we identify several deficiencies in the analysis, and we provide possible sources of background, that could explain the error.

In their recent Letter Ayyad et al. reported the first direct observation of delayed protons emitted in the  $\beta$  decay of  $^{11}\text{Be}$  [1]. The authors claim that the decay proceeds through a narrow resonance in  $^{11}\text{B}$  at an excitation energy of 11.425(2) MeV with a branching ratio of  $1.3(3) \times 10^{-5}$ . From the position of the resonance and the measured branching ratio Ayyad et al. calculate the  $\log(ft)$  value to be 4.8(4). This value is wrong. A simple estimate based on the resonance position and experimental branching ratio gives a  $\log(ft)$  value of 2.9 for a sharp resonance. Inserting a realistic width will not change this value much, which means that the beta strength  $\langle GT \rangle^2$  would be around 4.8, in significant disagreement with the model presented in Ref. [1]. Such a large value cannot be explained in any single-nucleon model where  $\langle GT \rangle^2$  at most can be 3. The combination of the resonance position and branching ratio reported by Ayyad et al. is in contradiction with any model presented so far.

The direct observation of protons emitted in the  $\beta$  decay of  $^{11}\text{Be}$  is extremely challenging because of the small energy window available of about 280 keV and the very low branching ratio [2]. Riisager et al. [3] reported the indirect observation of this decay channel with a branching ratio of  $(8.3 \pm 0.9) \times 10^{-6}$  through identification with the AMS technique of atoms of  $^{10}\text{Be}$  in collected samples of  $^{11}\text{Be}$ . Importantly,  $^{11}\text{Be}$  may also decay by  $\beta$ -delayed  $\alpha$  emission ( $\beta\alpha$ ) [4] with the much larger total branching of 3.30(10)%. In the experiment of Ayyad et al. the selected ions of  $^{11}\text{Be}$  were stopped inside a gaseous time projection chamber. About 90% of them drifted to the cathode before decaying - in such case only one heavy decay product could enter the active volume and be recorded.

The most likely cause of error is a contribution to the spectrum from particles other than protons. A key ingredient of the analysis was the selection of candidate proton events from a large background coming from the  $\beta\alpha$  channel. We show in Fig. 1 the expected spectra of recoiling  $^7\text{Li}$  ions and  $\alpha$  particles at low energy based on fits obtained in a recent experiment [4]. In their analysis Ayyad et al. entirely neglected low-energy  $\alpha$  particles as a possible source of background. However, the extrapolation shown in Fig. 1 indicates that about 150  $\alpha$  particles could be expected below 250 keV, which




FIG. 1. The  $\beta$ -delayed proton spectrum from  $^{11}\text{Be}$  reported in Ref. [1] is shown along with extrapolated intensities of  $\alpha$  particles and  $^7\text{Li}$  recoils from the  $\beta\alpha$  branch as determined in Ref. [4].

amounts to 40% of the reported branching ratio. In Ref. [1] the  $^7\text{Li}$  ions were taken into account as a source of background, but the quality of the proton- $^7\text{Li}$  discrimination was not documented; Fig. 2 of Ref. [1] shows only two example events of unknown energy. Using the absolute values of  $\chi^2$ , as in their Fig. 3, is not appropriate as they depend on the number of data points (samples) fitted, and thus on the particle energy and the emission angle. For event classification the normalized  $\chi^2$  per number of degrees of freedom should have been used instead. Hence, a fraction of  $^7\text{Li}$  ions could also be a source of background leading to the unphysical peak position and branching ratio reported by Ayyad et al.

An additional issue with the analysis of Ayyad et al. concerns the extraction of the width of the resonance. The width of the proton peak shown in Fig. 3 in Ref. [1] is 42 keV (FWHM), which is inconsistent with the quoted energy resolution of 15 keV and the width of the fitted Breit-Wigner distribution ( $\Gamma = 15$  keV).

\* pfutzner@fuw.edu.pl

[1] Y. Ayyad et al., Phys. Rev. Lett **123**, 082501 (2019).

[2] M.J. Borge et al., J. Phys. G: Nucl. Part. Phys. **40**, 035109 (2013).

[3] K. Riisager et al., Phys. Lett. B **732**, 305 (2014).

[4] J. Refsgaard et al., Phys. Rev. C **99**, 044316 (2019).