

The UV/optical peak and X-ray brightening in TDE candidate AT2019azh: A case of stream-stream collision and delayed accretion

XIAO-LONG LIU,¹ LI-MING DOU,^{2,3} RONG-FENG SHEN,^{1,*} AND JIN-HONG CHEN¹

¹*School of Physics and Astronomy, Sun Yat-Sen University, Zhuhai, 519082, China*

²*Center for Astrophysics, Guangzhou University, Guangzhou 510006, China*

³*Astronomy Science and Technology Research Laboratory of Department of Education of Guangdong Province, Guangzhou 510006, China*

ABSTRACT

We present and analyze the optical/UV and X-ray observations of a nearby tidal disruption event (TDE) candidate AT2019azh, spanning from 30 d before to ~ 250 d after its early optical peak. The X-rays show a late brightening by a factor of ~ 30 -100 around 250 days after discovery, while the UV/opticals continuously decayed. The early X-rays show two flaring episodes of variation, temporally uncorrelated with the early UV/opticals. We found a clear sign of X-ray hardness evolution, i.e., the source is harder at early times, and becomes softer as it brightens later. The drastically different temporal behaviors in X-rays and UV/opticals suggest that the two bands are physically distinct emission components, and probably arise from different locations. These properties argue against the reprocessing of X-rays by any outflow as the origin of the UV/optical peak. The full data are best explained by a two-process scenario, in which the UV/optical peak is produced by the debris stream-stream collisions during the circularization phase; some low angular momentum, shocked gas forms an early, low-mass accretion disk which emits the early X-rays. The major body of the disk is formed after the circularization finishes, whose enhanced accretion rate produces the late X-ray brightening. AT2019azh is a strong case of TDE whose emission signatures of stream-stream collision and delayed accretion are both identified.

Keywords: supermassive black holes - tidal disruption - galaxy accretion disks - transient source

1. INTRODUCTION

In a tidal disruption event (TDE), a star wanders toward the center of a galactic nucleus and is tidally disrupted by a supermassive black hole (SMBH). A bright, multi-wavelength flare is produced when the stellar debris falls back and accretes toward the black hole. Observing such incidents is an important way to probe the otherwise dormant SMBHs in the center of many non-active galaxies and to study the feeding process of SMBHs.

An unsettled issue still remains regarding the physical origin of the observed emission (i.e., where the emission is produced). Initially it was expected that the emission is produced in the accretion disk, which should peak in soft X-rays (Rees 1988). However, most of the observed TDEs are X-ray weak¹, only a small fraction of TDE sample have been detected with X-ray

emission in follow-up observations: GALEX D1-9 and D3-13 (Gezari et al. 2008), ASASSN-14li (Holoien et al. 2016b), AT2018fyk (Wevers et al. 2019) and ASASSN-15oi (Holoien et al. 2016a), with upper limits in a few cases: PS1-10jh (Gezari et al. 2012), iPTF16fnl (Blagorodnova et al. 2017; Brown et al. 2018), and iPTF16axa (Hung et al. 2017), while the majority were discovered in optical or UV (e.g., Gezari et al. 2012; Chornock et al. 2014; Holoien et al. 2014; van Velzen & Farrar 2014; Arcavi et al. 2014).

Two categories of models are proposed to explain the optical emissions. One involves ejected mass which reprocesses the high energy emission from the center to lower energies (Ulmer et al. 1997; Strubbe & Quataert 2009). The other focuses on the stream-stream interaction at the apocenter as the major energy dissipation site (Piran et al. 2015; Jiang et al. 2016). Recently Dai et al. (2018) explain the X-ray/optical dichotomy by considering the inclination effect and the angular distribution of the mass outflow's property such as density, speed and temperature.

Some TDEs have been discovered to have surprisingly low blackbody temperatures of $1 \sim 3 \times 10^4$ K through wide-field UV and optical surveys (Gezari et al. 2009, 2012; van Velzen et al. 2011; Arcavi et al. 2014; Holoien et al.

* Corresponding author: R.-F. Shen (shenrf3@mail.sysu.edu.cn)

¹ Note that a separate class of TDEs are non-thermal X-ray dominated, and they are thought to originate from a relativistic jet and require special orientation in order to be seen (Giannios & Metzger 2011; Burrows et al. 2011; Bloom et al. 2011; Levan et al. 2011; Zauderer et al. 2011; Cenko et al. 2012; Brown et al. 2015).

2014, 2016a,b; Blagorodnova et al. 2017; Hung et al. 2017), which can not be explained through radiation from traditional accretion process. These are attributed to larger radii associated with a reprocessing layer (Loeb & Ulmer 1997; Guillochon et al. 2014; Roth et al. 2016), which form from a radiatively driven wind (Miller et al. 2015; Strubbe & Murray 2015; Metzger & Stone 2016) or the radiation from stream-stream collisions during the circularization to form the disk (Lodato 2012; Piran et al. 2015; Shiokawa et al. 2015; Jiang et al. 2016; Krolik et al. 2016; Bonnerot et al. 2017; Wevers et al. 2017). On the other hand, one should expect the X-rays to show up eventually, once the wind subsides or the circularization finishes.

In this paper, we analyse the observations of a recent, nearby TDE candidate AT2019azh which shows a long (by ~ 200 days) delayed X-ray brightening with respect to its UV/optical peak, a pattern similar to that seen in the past TDE candidate ASASSN-15oi (Gezari et al. 2017). In §2 we describes the observations and data reduction in optical, UV and X-ray bands. Then we analyze its multi-wavelength light curves in §3, present the spectral energy distribution (SED) fit and show the source’s bolometric behavior (evolution of temperature and photospheric radius) in §4. We discuss three physical scenarios in TDEs in order to explain its multi-waveband behavior in §5. We summarize the results and conclude in §6.

2. OBSERVATIONS

2.1. Optical discovery

The bright nuclear transient AT2019azh was discovered by All Sky Automated Survey for SuperNovae (ASSA-SN) on UT 2019 Feb 22.02 (Brimacombe et al. 2019) and by Zwicky Transient Facility (ZTF) on 2019 Feb 12.40 (Van Velzen et al. 2019) in the center of an E+A galaxy at $z = 0.022$ (luminosity distance $D = 96$ Mpc). Follow-up spectroscopic observations by NUTS (Heikkilä et al. 2019) and ePESSTO (Barbarino et al. 2019) show a featureless blue spectrum.

Van Velzen et al. (2019) reported the ZTF and *Swift* UVOT photometry which show a ~ 15 -day long slowly-rising or plateau phase starting from 2 days after the ASSA-SN trigger. However, later monitoring showed that the flux continued to rise until it peaked at $g = 14.4$, about 31 days after the trigger (see Figures 1). The ZTF and host-subtracted UVOT photometry indicates a temperature of $\log(T) = 4.5 \pm 0.1$, and the ZTF photometry confirms that the transient is consistent with originating from the center of its host galaxy, with a mean offset of 0.07 ± 0.31 arcsec (Van Velzen et al. 2019). *Swift* XRT observations on 2019 Mar 11.45 detected 5 soft photons corresponding to a luminosity of $L_X = 2.5 \times 10^{41}$ erg s $^{-1}$ (Van Velzen et al. 2019).




Figure 1. Observed light curves of AT2019azh in optical, *Swift* UV and X-ray bands. The host contribution has already been subtracted in the optical and UV bands (*top*). The *bottom* panel shows the XRT 0.3-2 keV count rate light curve. The two data points with downward arrows are upper limits.

Based on its unclear position, persistent blue color, high blackbody temperature and lack of spectroscopic features associated with a supernova or AGN, Van Velzen et al. (2019) identified AT2019azh as a TDE, which we will follow hereafter.

We collect the public available ASSA-SN² g and V band, the ZTF³ g and r band, *Swift* UVOT, Gaia G band photometry data. We plot the host-subtracted source light curves in Figure 1.

2.2. *Swift* UV and X-rays

AT2019azh has been observed for 36 times with the *Swift* Observatory since March 2, 2019 (update to November 11, 2019). We download and reduce the *Swift* Observations with the software HEASoft V.6.26 and the latest updated calibration files of *Swift*. The UVOT Telescope observed the source with multi-wavelength filters (V, B, U, UVM2, UVW1, UVW2). We extract the source photometry from the source region of radius of 3", and the background from a source-free region with radius of 20" near the source position, using the task of 'uvotsource'. The UVOT photometry results are presented in Figure 1.

² https://asas-sn.osu.edu/light_curves/07988c67-2399-46f1-a9dc-3608c7e8141c

³ <https://lasair.roe.ac.uk/object/ZTF17aaazdba/>




Figure 2. Stacking *Swift* XRT images in the 2-10 keV (upper panels) and 0.3-2 keV (lower panels) bands.




Figure 3. The evolution of X-ray hardness of AT2019azh. Asterisks and crosses are the stacking count rates in 0.3-2 keV and 2-10 keV, respectively. The hardness ratio is defined as the count rate ratio between the two bands: $HR = (2-10 \text{ keV}) / (0.3-2 \text{ keV})$.

For *XRT*, we reprocess the event files with the task ‘xrt-pipeline’, and select the event files which operated in Photon Counting mode. The source is detected in almost all observations, with a count rate in the range of 0.001 - 0.1 cts s^{-1} in 0.3 - 2 keV. The source file is extracted using a source region of radius of 30''. The background is estimated in an annulus region centered on the source position, with an inner radius of 60'' and outer radius of 90''. Due to the low counts, we only extract the 0.3-2 keV band count rate for single observations. They are listed in Table 1 and plotted in Figure 1.

We also stack the *XRT* observations into five groups and list them in Table 1, which are marked as O01_09, O10_18, O20_31, O01_31, and O32_40. The 0.3-2 and 2-10 keV band stacking images for each group are presented in Figure 2. We find that the most of X-ray photons are detected in the soft band (0.3-2 keV). Interestingly, about two dozen of X-ray

photons are also detected in 2-10 keV band during the observations before June 4, 2019 (O01_31), while half of the hard band X-ray photons are detected during the 2019 April observations (O10_18). We extract the source and background files from the stacking events files. The stacking count rates in 0.3-2 keV and 2-10 keV, respectively, are listed in Table 1, and are plotted in Figure 3 as well.

Interestingly, the soft band X-ray count rate shows a factor of ~ 25 increase from O01_31 to O32_40, while the hard band X-ray count rate dims to be undetectable, as was shown in Figure 3. We will come back to this strong hardness evolution feature in §4.2.

2.3. Radio

Perez-Torres et al. (2019) reported radio detections of the source at 90 - 110 days after the ASAS-SN detection, from which we take the data and plot it in Figure 4. For comparison, we also plot the data of another radio bright TDE ASASSN-14li.

The host galaxy of AT2019azh is a non-active galaxy based on the lack of related spectroscopic feature (Brimacombe et al. 2019). Therefore, its radio emission ($\sim 10^{37} \text{ erg s}^{-1}$) is unlikely due to AGN activity. It is commonly believed that radio emission from TDEs are produced from the interaction of a relativistic jet or non-relativistic outflow with ambient medium. We discuss this later in §5.

3. TEMPORAL BEHAVIOR

The ASAS-SN and ZTF data make AT2019azh one of the few TDE candidates with a well-sampled rising light curve. It rises in the ASAS-SN *g*, *V* and ZTF *g*, *r* bands to the peak within 35 days, then decays gradually in all UV / optical filters. In addition, there is a minor re-brightening in the ASASSN *g* and *V* bands at $t \approx 90$ d.

Two features of the UV / optical light curves shown in Figure 1 are notable. First, the data in the several filters indicate very little color evolution in a time span of ~ 100 days, ex-




Figure 4. Radio light curve of AT2019azh. Data is from Perez-Torres et al. (2019). Its optical discovery date is set to be MJD58536. Another TDE ASASSN-14li (data from Alexander et al. 2016, discovery date MJD56983) is plotted for comparison.

cept for the earliest 10 days of the rising. This is supported by the inferred photospheric temperature evolution to be shown later in Figure 6. It is also in line with earlier discovery report by Van Velzen et al. (2019) and is a characteristic of most previously found TDEs (e.g., Holoien et al. 2019). Second, the post-peak magnitudes decay linearly with time, which means the flux decays in an exponential rather than a power law with time.

The X-ray count rate light curve in Figure 1 shows a very different behavior. Apparently it has two slow flares during the optical bright phase, each varying by a factor of ~ 10 with a duration of ~ 50 days. Most importantly, the X-ray brightens by a factor of ~ 30 at around 250 days after discovery. This late X-ray brightening is unusual for TDEs, and the only past similar case is ASASSN-15oi (Gezari et al. 2017).

The behaviors in X-rays and in UV/optical are quite different. The early ($t < 120$ d) X-rays show significant variations while UV/optical are in smooth rise and fall. Later, UV/optical decay monotonically for the rest of the time, but X-rays rise to its peak in about 250 days. These strongly suggest that the two emission components are produced probably at different locations and by different dissipation processes. This is helpful in discerning the most appropriate physical scenario in §5.

4. SPECTRAL ANALYSIS

4.1. UV/optical

To better quantify the physical parameters of the system, we modeled the UV and Optical SED of AT2019azh for epochs, which is shown in Figure 5, the blackbody fits provide good fits to the data. The resulting temperature evolution in days from detection is shown in Figure 6. Additionally, the evolution of the effective radius for the UV/optical and X-ray emission is shown in Figure 7. The bolometric luminosity is shown in Figure 9.




Figure 5. Blackbody fits to the multi-epoch SEDs composed of UV and optical photometry data. The numbers mark the time in days since the optical detection.




Figure 6. Temperature evolution of AT2019azh from blackbody fits to the UV/Optical SED.




Figure 7. Evolution of the photospheric radius derived from the blackbody SED fits to the UV/Optical.

The blackbody fits indicate that after a rise between 10 and 15 days after detection, the temperature of AT2019azh holds relatively constant around $T \simeq 26,000$ K for about 90 days. This temperature is similar to those of other TDEs.

4.2. X-rays

For Swift X-ray data, we fit the two stacking spectra which are grouped in before and after June 4, 2019 (O01_31 and O32_40). We group the data to have at least 4 counts in each bin, and adopt mainly the C-statistic for the Swift spectral fittings which are performed using XSPEC (v.12.9; Arnaud 1996). We try to fit the spectra with five different models, which are a single power-law, a double power-laws, a single blackbody, a double blackbodies, and a single power-law plus single blackbody, respectively. For Galactic absorption, we adopt a column density of $N_H = 4.15 \times 10^{20} \text{ cm}^{-2}$ (HI4PI Collaboration et al. 2016) in the direction of AT2019azh.

The fitting results are listed in the Table 2. We find that the best fit and accepted model for O01_31 is a single power-law, with photon index $\Gamma = 1.9 \pm 0.6$, plus a single blackbody of temperature $kT = 56 \pm 9$ eV. The best fit and accepted model for O32_40 is a double blackbodies, with $kT_1 = 51 \pm 4$ eV and $kT_2 = 120 \pm 28$ eV. They are shown in Figure 8.

Based on the best fit results, we estimate the X-ray fluxes of the two stacking spectra. The 0.3-2, and 2-10 keV band unabsorbed fluxes in O01_31 are $2.34_{-0.37}^{+0.10} \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$, and $2.5_{-0.8}^{+1.0} \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$, respectively. The 0.3-2 keV band unabsorbed flux in O32_40 is $7.10_{-0.50}^{+0.08} \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$, while the 0.3-2 keV band flux of the second blackbody component is only $\sim 7\%$ of the flux of first one. Using the stacking spectral fitting results, we convert the count rate of each XRT exposure to the 0.3 - 10 keV flux. Replacing the blackbody component in the best fit models with a multi-blackbody from accretion disk model ('diskbb'), we also derive the X-ray blackbody radii, which are $\sim 7 \times 10^{10}$, 6×10^{11} and $1 \times 10^{10} \text{ cm}$, respectively, for the blackbody component in O01_31, the first, and the second one in O32_40. These radii are 0.07-4 times of R_g ($R_g = GM/c^2$), if considering an SMBH mass of $10^6 M_\odot$.

As was shown in Figure 3, the X-ray hardness ratio drops substantially toward later times and becomes softer when the source is brighter, which is reminiscent of the state transitions behavior typically seen in BH X-ray binaries (e.g., Remillard & McClintock 2006). Thus, the X-rays here can be considered as the signature of the BH accretion. Therefore, the early X-rays probably correspond to a low accretion state which is harder, whereas the late X-ray brightening at $t = 250$ d corresponds to a high accretion rate where the hard photons disappear.

4.3. Comparison between UV/optical and X-rays

Figure 9 shows the UV/optical and X-ray luminosity evolution of AT2019azh. In the figure we also show $t^{-5/3}$ power-law and $e^{-t/\tau}$ exponential fits to the data. However, it is hard to compare the two fitting methods because of the large error, although in Figure 1 the exponential fits well to data.

In addition, we plot the evolution of the X-ray-to-UV/optical luminosity ratio L_X/L_{opt} in Figure 10. It shows that AT2019azh and ASASSN-15oi are generally very similar, in that this ratio rises from 0.001-0.01 during the early optically bright phase to ~ 1 later. They differ from ASASSN-14li which shows almost constant $L_X/L_{\text{opt}} \sim 1$.

5. PHYSICAL INTERPRETATION

In order to compare the evolution of AT2019azh to the relevant timescales for a TDE, we adopt an radius relation $r_* \approx m_*^{0.89}$ for low-mass main sequence stars (Torres et al. 2010), where $R_* = r_* \times R_\odot$ and $M_* = m_* \times M_\odot$ is the star's radius and mass. The characteristic timescale for a TDE is set by the orbital period of the most tightly bound debris, known as the fallback time

$$t_{\text{fb}} = 41 M_6^{1/2} m_*^{0.34} d \quad (1)$$

where, $M = M_6 \times 10^6 M_\odot$ is the mass of the black hole. The rise time of a TDE light curve could be a rough estimate of t_{fb} ; in the case of AT2019azh, from Figure 1 we estimate $t_{\text{fb}} \approx 30$ d. After disruption, the less bound debris follows the most bound debris in returning, in a rate that drops with time as (Rees 1988; Phinney 1989; Lodato et al. 2009; Ramirez-Ruiz & Rosswog 2009; Guillochon & Ramirez-Ruiz 2013)

$$\dot{M}_{\text{fb}} \simeq \dot{M}_{\text{peak}} \left(\frac{t}{t_{\text{fb}}} \right)^{-5/3}. \quad (2)$$

In the following we discuss three possible scenarios for interpreting the UV/optical and X-ray behavior of AT2019azh.

5.1. Super-Eddington accretion for the UV/optical peak

The optical bright stage could be super-Eddington which peaks in the UV/optical bands, during which various energy dissipation processes will produce winds or outflows (Strubbe & Quataert 2009; Lodato & Rossi 2011; Metzger & Stone 2016) which will regulate the luminosity (Krolik & Piran 2012), and block the X-rays or reprocess them into UV/opticals (e.g., Dai et al. 2018). After the accretion rate drops below the Eddington rate, the bolometric luminosity falls and X-rays from the inner accretion disk start to be seen (Chen & Shen 2018). This transition time can be estimated from Eq. (2) as

$$t_{\text{Edd}} \simeq 2.2 \eta_{0.1}^{3/5} M_6^{2/5} r_*^{3/5} m_*^{1/5} \text{ yr}, \quad (3)$$




Figure 8. Stacking *Swift* XRT spectra and best fit models. The data is marked as black color, the best fit models are marked as blue color, the blackbody model component is marked as red dot line (the second blackbody in O32_40 is marked as pink dot line) and the power-law model component is marked as green dot line.




Figure 9. UV/optical luminosity evolution of AT2019azh from the blackbody SED fits (red) and X-ray luminosity evolution from *Swift* (black).

where $\eta = 0.1 \times \eta_{0.1}$ is the efficiency of converting accretion power to luminosity. AT2019azh is visible in X-rays during the early, optical bright phase. This suggests that, in Dai et al. (2018)’s model, the line of sight is somewhat close to the pole direction.

However, there are two evidences against this early super-Eddington accretion scenario. First, the X-rays and UV/optical light curves of AT2019azh behave very differently during the early time ($t < 100$ d), as was shown in Figure 1 and mentioned in §3. They show no sign of temporal correlation between the two bands that one should expect to see in this scenario, since there the two bands are produced by the same accretion process⁴.

⁴ We notice that the early UV/optical and X-rays in AT2018fyk do show a weak temporal correlation (Wevers et al. 2019). Therefore, AT2018fyk may belong to the early super-Eddington accretion scenario.




Figure 10. Evolution of the luminosity ratio of X-rays over UV/optical. The data source for the other three TDEs: ASASSN-14li and ASASSN-15oi are from Gezari et al. (2017), and AT2018fyk is from Wevers et al. (2019).

Second, as was shown in Figure 3 and mentioned in §4.2, hard X-ray photons appeared during early time but disappeared later. The resemblance of this particular behavior to the state transition pattern of BH X-ray binaries suggests that those X-rays, both the early and the late, are signatures of accretion. The early X-rays probably correspond to a low accretion state which is harder, whereas the late X-ray brightening at $t \simeq 250$ d corresponds to a high accretion rate so the hard photons disappear. This is in sharp contradiction with the early UV/optical peak being a super Eddington accretion phase.

5.2. Stream-stream collision followed by delayed accretion

Next we consider a two-process scenario, in which the late X-ray brightening comes from the delayed accretion

through a recently formed accretion disk, and the early UV/optical emission is from the stream-stream collisions (e.g., Piran et al. 2015; Jiang et al. 2016) before the major body of the accretion disk is formed.

Bonnerot et al. (2016) estimated the circularization timescale, which is driven by relativistic apsidal precession of the debris streams, to be

$$t_{\text{cir}} = 8.3 t_{fb} M_6^{-5/3} \beta^{-3}, \quad (4)$$

where $\beta = R_T/R_P$, $R_T = R_*(M/M_*)^{1/3}$ (Rees 1988; Phinney 1989). Once a disk is formed with radius R , the viscous inflow timescale for a standard α -disk model (Shakura & Sunyaev 1973) is $t_{\text{vis}} = [\alpha \Omega_K(R)]^{-1} (h/r)^{-2}$, where α is the standard viscous parameter, $\Omega_K(R)$ is the Keplerian angular speed, h/r is the disk's scale-height to radius ratio. Since the disk forms at radius of about $2R_p$, the ratio of t_{vis} over t_{fb} is

$$\frac{t_{\text{vis}}}{t_{fb}} = 0.1 \beta^{-3/2} \left(\frac{0.01}{\alpha} \right) \left(\frac{m_*}{M_6} \right)^{1/2} \left(\frac{h}{r} \right)^{-2}. \quad (5)$$

In the following we attempt to model how the energy release rate from the stream-stream collisions evolves. During the fallback process, the orbit of the most bound debris has a semi-major axis of $a_{\text{mb}} \simeq R_T^2/(2R_*)$, an orbital period of t_{fb} and an specific energy of $E_0 = -GM/2a_{\text{mb}}$, upon its first pericenter passage. During subsequent passages its orbit shrinks (circularizes) due to self-crossing and collision, with its semi-major axis $a(t)$ being ever decreasing. We suppose the specific energy dissipation rate is

$$-\dot{q} = \frac{dE}{dt} = \delta(a) \frac{E(a)}{t(a)} \quad (6)$$

where the efficiency factor $\delta \ll 1$ represents the fraction of orbital energy that is dissipated per orbit, and the orbital period $t(a) = t_0(a/a_0)^{3/2}$. We assume $\delta(a) = \delta_0(a/a_0)^\gamma$, where $\gamma > 0$, i.e., the dissipation efficiency drops progressively as the stream is being circularized. With algebra, the equation becomes

$$\frac{d}{dt} \left(\frac{a}{a_0} \right) = -\frac{\delta_0}{t_0} \left(\frac{a}{a_0} \right)^{\gamma-1/2} \quad (7)$$

The solution is

$$a(t) = a_0 \left[1 + (\gamma - \frac{3}{2}) \frac{\delta_0}{t_0} (t - t_0) \right]^{-\frac{1}{\gamma-3/2}} \quad (8)$$

Therefore, we get the specific energy $E(t) = -\frac{GM}{2a}$, and the dissipation rate

$$\dot{q}(t) = \frac{GM}{2a_0} \frac{\delta_0}{t_0} \left[1 + (\gamma - \frac{3}{2}) \frac{\delta_0}{t_0} (t - t_0) \right]^{\frac{1}{\gamma-3/2}-1}. \quad (9)$$

If $\gamma > 5/2$, then $\dot{q}(t)$ decreases with time as $t^{-\alpha}$, with $\alpha = (\gamma - 5/2)/(\gamma - 3/2)$. We assume the relationship

between release energy and luminosity is $L(t) = \lambda \dot{q}(t) c^2$, where λ is the energy conversion efficiency c is light speed. We take $\gamma = 10$ and plot $L(t)$ in Figure 9, which roughly shows the feasibility of matching the radiation from stream-stream collisions to the UV/optical data.

What produced the early ($t < 100$ d) low-level X-ray activity? During the stream-stream collisions, the hydrodynamic numerical simulation by Shiokawa et al. (2015) shows that some minor amount of debris lost a significant fraction of their angular momentum, such that their pericenter radius could shrink significantly and they could form an early, low mass accretion disk. If this is possible, then one naturally expects that the early accretion rate is low (i.e., sub-Eddington). The hard X-ray photons appearing during this 'low hard' state, similar to what happens in BH X-ray binaries.

The above scenario is naturally consistent with the absence of temporal correlation between early X-rays and UV/opticals, since they are produced by different dissipation processes and at very different locations.

Once the stream-stream collision process was over, the major body of the disk should have formed disk formed around $t \sim 250$ d. The accretion rate has risen to the peak, so does the soft X-ray flux, while hard X-ray photons disappear, probably with the hot corona.

5.3. CIO's reprocessing of X-rays

Lu & Bonnerot (2019) argue that during the stream-stream collisions, a considerable number of the shocked gas will become unbound and ejected as the so-called collision-induced outflow (CIO), which could reprocess early X-rays (presumably coming from an inner disk which is formed in the same way as was described in §5.2) into optical bands.

This scenario is disfavored for the UV/optical peak in AT2019azh due to the following reasons. If the content of CIO is large and massive so that the CIO could cover the whole sphere around the source, then the early X-rays should not be visible, which is clearly not the case. If the CIO coverage is partial and it does not fully block the line of sight, then one should expect: 1) the observed X-ray flux shall exceed or at least be comparable to the UV/optical flux, not the opposite, because only a portion of L_X gets reprocessed and becomes L_{opt} ; 2) the UV/optical should show a temporal correlation with the early X-rays, which is in contradiction with what the early light curves show in Figures 1 and 9.

Note that AT2019azh is detected in radio at $t \sim 100$ d (see §2.3 and Figure 4). This suggests that CIO may actually exist and its interaction with ambient medium produced the radio emission (Lu & Bonnerot 2019). However, the CIO's reprocessing of X-rays can not be the origin of the UV/optical peak.

6. SUMMARY AND CONCLUSION

We present and analyze a large data set by ASAS-SN, ZTF, *Swift* and Gaia of the light curves of TDE candidate AT2019azh in optical/UV and X-ray bands. We highlight a rare case in which the late X-rays brightened by a factor of ~ 30 -100 around 250 days after discovery, while the UV/opticals continuously decayed. The early X-rays show two flaring episodes of variation, temporally uncorrelated with the early UV/opticals. In addition, we present the evolution of temperature and photospheric radius from the fitting of SED. We found a clear sign of evolution of the X-ray hardness ratio which drops substantially toward later times and becomes softer when the source is brighter.

The drastically different temporal behaviors in X-rays and UV/opticals suggest that the two bands are physically distinct emission components, and probably arise from different locations. The hard X-ray (2 -10 keV) photons found during $t < 100$ d suggest that the early X-rays must be of accretion origin as well.

Putting all pieces together, we conclude that the full data are best explained by a two-process scenario, in which the UV/Optical peak is produced by the stream-stream collisions during the circularization phase; the same process causes some low angular momentum, shocked gas to form an early, low-mass accretion disk which emits the early X-rays. The major body of the disk is formed after the circularization finishes, at $t \sim 250$ d. Its enhanced rate of accretion toward the black hole produces the late X-ray brightening.

AT2019azh is the second case, after ASASSN-15oi (Gezari et al. 2017), of TDEs that shows a clear sign of delayed accretion. However, the early detection and full multi-waveband coverage make AT2019azh the first strong case that the emission signature of stream-stream collision is identified and early steps of disk formation can be inferred. At the time of the paper is written, AT2019azh is still detectable in X-rays, so deeper and broader understanding of this event is reachable.

REFERENCES

- Arcavi, I., et al. 2014, *ApJ*, 793, 38
- Alexander, K., D., Berger, E., Guillochon, J., et al. 2016, *ApJ*, 819, 25
- Barbarino, C.; Carracedo, A. S.; Tartaglia, L., et al., 2019, *ATel* #12530
- Blagorodnova, N., Gezari, S., Hung, T., et al. 2017, *ApJ*, 844, 46
- Bloom J. S. et al., 2011, *Science*, 333, 203
- Bonnerot, C., Rossi, E. M., Lodato, G., & Price, D. J. 2016, *MNRAS*, 455, 2253
- Bonnerot, C., Rossi, E. M., & Lodato, G. 2017, *MNRAS*, 464, 2816
- Brimacombe, J.; Kendurkar, M.R.; Masi, G., et al., 2019, *ATel* #12526
- Brown, G. C.; Levan, A. J.; Stanway, E. R., et al., 2015, *MNRAS* 452, 4297
- Brown, J. S., Kochanek, C. S., Holoien, T. W. S., et al. 2018, *MNRAS*, 473, 1130
- Burrows D. N., et al., 2011, *Nature*, 476, 421
- Cenko, S. B.; Krimm, H. A.; Horesh, A., et al., 2012, *ApJ*, 753, 77
- Chen, J.-H., & Shen, R.-F., 2018, *APJ*, 867, 20
- Chornock, R., et al. 2014, *ApJ*, 780, 44
- Dai, L.-X.; McKinney, J. C.; Roth, N., 2018, *ApJ*, 859L, 20
- Gezari, S., Basa, S., Martin, D. C., et al. 2008, *ApJ*, 676, 944
- Gezari, S., Heckman, T., Cenko, S. B., et al. 2009, *ApJ*, 698, 1367
- Gezari, S., Chornock, R., Rest, A., et al., 2012, *Nature*, 485, 217
- Gezari, S.; Cenko, S. B.; Arcavi, I., 2017, *ApJL*, 851, L47
- Giannios, D., & Metzger, B. D. 2011, *MNRAS*, 416, 2102
- Guillochon, J., Manukian, H., & Ramirez-Ruiz, E. 2014, *ApJ*, 783, 23
- Guillochon, J., & Ramirez-Ruiz, E. 2013, *ApJ*, 767, 25
- Holoien, T. W.-S., Prieto, J. L., Bersier, D., et al., 2014, *MNRAS*, 445, 3263
- Holoien, T. W.-S., Kochanek, C. S., Prieto, J. L., et al. 2016a, *MNRAS*, 463, 3813
- Holoien, T. W.-S., Kochanek, C. S., Prieto, J. L., et al. 2016b, *MNRAS*, 455, 2918
- Holoien, T. W.-S., Vallely, P. J.; Auchettl, K., et al., 2019, *ApJ*, 883, 111
- HI4PI Collaboration, N. Ben Bekhti, L. Floer, et al., 2016, *A&A*, 594, A116
- Hung, T., Gezari, S., Blagorodnova, N., et al., 2017, *ApJ*, 842, 29
- Heikkilä, T.; Reynolds, T.; Kankare, E., et al., 2019, *ATel* #1252
- Jiang, Y.-F., Guillochon, J., & Loeb, A., 2016, *ApJ*, 830, 125
- Krolik, J. H., & Piran, T., 2011, *ApJ*, 743, 134
- Krolik, J., Piran, T., Svirski, G., & Cheng, R. M., 2016, *ApJ*, 827, 127
- Lodato, G., King, A. R., & Pringle, J. E., 2009, *MNRAS*, 392, 332
- Lodato, G., & Rossi, E. M., 2011, *MNRAS*, 410, 359
- Lodato, G., 2012, *EPJWC*, 39, 01001
- Loeb, A., & Ulmer, A., 1997, *ApJ*, 489, 573
- Lacy, J. H., Townes, C. H., & Hollenbach, D. J. 1982, *ApJ*, 262, 120
- Levan A. J., et al., 2011, *Science*, 333, 199
- Li, L., Narayan, R., & Menou, K., 2002, *ApJ*, 576, 753
- Lu, W., & Bonnerot, C., 2019, *arXiv:1904.12018*
- Miller, J. M., Kaastra, J. S., Miller, M. C., et al. 2015, *Natur*, 526, 542
- Metzger, B. D., & Stone, N. C., 2016, *MNRAS*, 461, 948
- Piran, T., Svirski, G., Krolik, J., 2015, *ApJ*, 806, 164
- Perez-Torres, M., Moldon, J., Mattila, S., et al., 2019, *ATel* #12870

- Phinney, E. S., 1989, IAUS, 136, 543
- Ramirez-Ruiz, E., & Rosswog, S., 2009, ApJL, 697, L77
- Rees, M. J., 1988, Nature, 333, 523
- Remillard, R. A., & McClintock, J. E., 2006, ARA&A, 44, 49
- Roth, N., Kasen, D., Guillochon, J., & Ramirez-Ruiz, E. 2016, ApJ, 827, 3
- Strubbe, L., E., Quataert, E., 2009, MNRAS, 400, 2070
- Strubbe, L. E., & Murray, N., 2015, MNRAS, 454, 2321
- Shiokawa, H., Krolik, J. H., Cheng, R. M., Piran, T., & Noble, S. C., 2015, ApJ, 804, 85
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337
- Torres, G., Andersen, J. & Gimenez, A., 2010, A&ARv, 18, 67
- Ulmer, A., Paczyński, B., Goodman, J., 1997, A&A, 333, 379
- van Velzen, S., Farrar, G., Gezari, S., et al., 2011, ApJ, 741, 73
- van Velzen, S., & Farrar, G. R., 2014, ArXiv e-prints
- van Velzen, S., Gezari, S., Hung, T., Gatkine, T., Cenko, S. B., Ho, A., Kulkarni, S. R., Mahabal, A., 2019, ATel #12568
- Wevers, T., van Velzen, S., Jonker, P. G., et al., 2017, MNRAS, 471, 1694
- Wevers, T., Pasham, D., R., van Velzen, S., et al., 2019, MNRAS, 488, 4816
- Zauderer B. A., et al., 2011, Nature, 476, 425

Table 1. The log of *Swift*-XRT observation.

ObsID (11186-)	ObsDate	Exposure	Count rate 0.3-2 keV (10^{-3} cts/s)	Flux 0.3-2 keV (10^{-13} erg/s/cm ²)		
		(ks)				
001	2019-03-02T18:00:36	2.198	<0.45	<0.4		
003	2019-03-11T10:42:36	0.529	7.55±3.78	6.2±3.16		
004	2019-03-14T02:26:34	0.484	6.19±3.57	5.1±2.97		
005	2019-03-20T22:38:35	1.066	5.43±2.30	4.4±1.94		
006	2019-03-23T22:17:34	0.996	6.02±2.46	4.9±2.07		
007	2019-03-26T06:32:34	1.166	5.14±2.10	4.2±1.77		
008	2019-03-26T01:17:35	0.302	6.62±4.7	5.4±3.87		
009	2019-03-26T01:22:35	2.040	3.92±1.38	3.2±1.18		
010	2019-04-01T18:40:36	1.366	2.04±1.28	1.7±1.06		
011	2019-04-04T02:13:36	3.264	2.15±0.81	1.8±0.69		
012	2019-04-07T14:39:34	2.879	1.79±0.86	1.5±0.72		
013	2019-04-10T06:20:35	2.977	1.68±7.51	1.4±0.63		
014	2019-04-16T12:23:04	2.814	1.20±0.72	1.0±0.60		
015	2019-04-19T00:53:36	2.974	2.02±0.82	1.7±0.69		
016	2019-04-22T08:25:34	3.179	3.08±1.00	2.5±0.85		
017	2019-04-25T06:38:34	2.972	4.38±1.21	3.6±1.06		
018	2019-04-27T07:56:35	0.092	≤10.82	≤8.9		
020	2019-05-03T02:39:35	3.014	6.16±1.45	5.1±1.29		
021	2019-05-06T03:58:34	2.829	3.89±1.17	3.2±1.01		
022	2019-05-08T10:00:35	2.707	7.31±1.65	6.0±1.48		
024	2019-05-15T07:44:34	2.505	5.02±1.44	4.1±1.25		
025	2019-05-18T07:27:36	2.807	2.42±0.94	2.0±0.8		
026	2019-05-21T10:21:36	2.667	2.98±1.06	2.4±0.9		
027	2019-05-24T00:32:36	2.829	4.17±1.23	3.4±1.06		
029	2019-05-27T08:14:36	1.493	2.01±1.16	1.6±0.96		
030	2019-05-31T04:42:36	1.536	2.60±1.30	2.1±1.09		
031	2019-06-04T06:21:36	2.322	2.09±0.96	1.7±0.81		
032	2019-10-11T22:12:36	1.618	74.74±0.68	61.2±8.27		
033	2019-10-17T20:27:35	1.094	96.33±0.94	78.9±11.01		
034	2019-10-23T21:11:35	1.656	95.87±0.76	78.6±10.02		
035	2019-10-26T16:02:34	2.008	163.8±0.9	134.2±15.29		
036	2019-10-31T04:45:36	1.930	66.36±0.59	54.4±7.26		
037	2019-11-03T14:01:14	0.499	130.9±0.2	107.3±17.08		
038	2019-11-07T15:08:35	1.878	32.03±0.42	26.2±4.30		
039	2019-11-09T14:42:36	2.030	48.43±0.49	39.7±5.65		
040	2019-11-12T16:25:36	1.905	95.07±0.71	77.9±9.68		
ObsID 11186- stacking	ObsDate	Exposure total (ks)	Count rate 0.3-2 keV (10^{-3} cts/s)	Count rate 2-10 keV (10^{-3} cts/s)	Flux 0.3-2 keV (10^{-13} erg/s/ cm ²)	Flux 2-10 keV (10^{-13} erg/s/cm ²)
001-009	March	8.783	3.94±0.67	0.24±0.20	2.71 ^{-0.66} _{+0.52}	0.19 ^{-0.18} _{+0.19}
010-018	April	22.52	2.41±0.34	0.39±0.15	1.66 ^{-0.37} _{+0.27}	0.32 ^{-0.18} _{+0.20}
020-031	May-Jun	24.71	4.13±0.42	0.19±0.11	2.84 ^{-0.57} _{+0.38}	0.15 ^{-0.11} _{+0.12}
001-031	Mar-Jun	56.01	3.40±0.25	0.309±0.09	2.34 ^{-0.37} _{+0.10}	0.25 ^{-0.08} _{+0.10}
032-040	Oct-Nov	14.62	85.29±0.24	<0.1041	58.70 ^{-10.24} _{+4.99}	<0.084

Table 2. Stacking X-ray spectral fitting results. The uncertainties are given at 90% confidence level.

NOTE: The Galactic absorbed models "phabs * (1. or 2. or 3. or 4. or 5.)" are used for fitting (1. powerlaw, 2. powerlaw + powerlaw, 3. zbody, 4. zbody+powerlaw, 5. zbody+zbody).

The best-fit model is 4 for the stacking O01_31, and 5 for the stacking O32_40.

Stacking O01_31				
	Model	Parameter	Value	C-Statistic/d.o.f
1	powerlaw	Γ	$4.95^{+0.44}_{-0.40}$	64.4/36
2	powerlaw1	Γ_1	$6.10^{+0.90}_{-0.73}$	33.4/34
	powerlaw2	Γ_2	$1.34^{+0.77}_{-0.89}$	
3	zbbody	kT (keV)	$0.077^{+0.008}_{-0.008}$	104.72/36
4	zbbody	kT (keV)	$0.056^{+0.009}_{-0.009}$	33.6/34
	powerlaw	PhoIndex	$1.95^{+0.64}_{-0.59}$	
Stacking O32_40				
	Model	Parameter	Value	C-Statistic/d.o.f
1	powerlaw	Γ	$6.75^{+0.20}_{-0.19}$	63.6/46
3	zbbody	kT (keV)	$0.060^{+0.002}_{-0.002}$	98.2/46
4	zbbody	kT (keV)	$0.053^{+0.006}_{-0.004}$	46.4/44
	powerlaw	PhoIndex	$6.00^{+0.70}_{-0.96}$	
5	zbbody1	kT_1 (keV)	$0.051^{+0.003}_{-0.004}$	43.9/44
	zbbody2	kT_2 (keV)	$0.120^{+0.028}_{-0.020}$	