

The Geometry of H_4 Polytopes

Tomme Denney, Da'Shay Hooker, De'Janeke Johnson,
Tianna Robinson, Majid Butler and Sandernisha Claiborne*

We describe the geometry of an arrangement of 24-cells inscribed in the 600-cell. In §7 we apply our results to the even unimodular lattice E_8 and show how the 600-cell transforms $E_8/2E_8$, an 8-space over the field \mathbf{F}_2 , into a 4-space over \mathbf{F}_4 whose points, lines and planes are labeled by the geometric objects of the 600-cell.

May 25, 2019

1 Introduction

The 600-cell occupies a difficult niche in Euclidean geometry. Because it exists in four dimensions the Greeks never considered it, and it will never achieve the fame of the five Platonic solids. Because it's the last of its kind (dimensions higher than four admit only simplexes, orthoplexes and n-cubes ([14], Chapter X)), no higher-dimensional polytope analogues of the 600-cell exist, so it can never be part of a systematic treatment of polytopes and must always be treated as an outlier. Even its most natural coordinates in 4-space are a disadvantage: they lie in $\mathbb{Z}[\frac{1+\sqrt{5}}{2}]$ and not in \mathbb{Z} , so the regular H_4 polytopes (the 600-cell and its dual, the 120-cell) seem less natural than polytopes whose coordinates are all integers.

And yet these drawbacks account for the 600-cell's fascination. The 600-cell is scarcely known because it exists in four dimensions and not three; but a well-known theorem of Hurwitz states that composition algebras exist only in dimensions 1, 2, 4 and 8—these are the real numbers, the complex numbers,

*The authors were supported by the Student Research Fellowship program at McDonogh 35 High School in New Orleans. The authors thank Professor Cathy Kriloff of Idaho State University for suggesting they use the techniques they developed in [1] to study H_3 (and H_4) symmetry, they thank their advisor Rich Margolin for several helpful conversations, and they thank the referee for a number of suggestions that improved the text and corrected the terminology of this paper.

the quaternions and the octonions ([9], §7.6)—which raises the possibility that the 600-cell has some connection to the quaternions. (In fact, it does: see Fact 8 and §2 below). And because the four coordinates for H_4 polytopes lie in $\mathbb{Q}(\sqrt{5})$, the variety of “norm reduction” maps from this extension field to \mathbb{Q} results in numerous different ways to view these polytopes (§6, Examples 1-3). Still, despite these varied approaches to H_4 objects, the geometry of the 600-cell (and other H_4 polytopes) has remained all too mysterious.

Even the experts have sometimes been confused about the structure of H_4 polytopes. In 1907 the Dutch mathematician P.H. Schoute published, without proof, the following claim: “Die 120 Eckpunkte eines $Z_{600}^{(1)}$ bilden auf zehn verschiedene Arten die Eckpunkte von fünf $Z_{24}^{\frac{1}{2}(1+\sqrt{5})}$.” ([13], p. 231) (“The 120 vertices of a 600-cell form in ten different ways the vertices of five 24-cells.”) Evidently this claim was viewed skeptically: in 1933 H.S.M. Coxeter published his opinion that “But surely *zehn* should be *fünf* in the phrase *auf zehn verschiedene Arten*,” ([4], p. 337) i.e., that surely there were only five and not ten such partitions.

But in fact there are ten, and not five, such partitions of the vertices of the 600-cell. Coxeter realized his mistake and corrected it: “Thus Schoute was right when he said the 120 vertices of $\{3, 3, 5\}$ belong to five $\{3, 4, 3\}$ ’s in ten different ways. The disparaging remark in the second footnote to Coxeter [4], p. 337, should be deleted.” ([5], p. 270) And although it was later fixed, Coxeter’s mistake highlights a difficulty that has plagued H_4 polytopes (and their symmetry group) since their study began roughly 150 years ago: H_4 remains more complicated than it should.

So the goal of our paper is to make the geometry of the 600-cell a little more accessible. We’ve tried to achieve this in four ways: (i) by collecting a number of results (some old and some new) about the 600-cell and listing them as Facts later in our Introduction; (ii) by proving Schoute’s century-old statement that there are exactly ten ways to partition the 120 vertices of a 600-cell into five disjoint 24-cells¹ ([13], p. 231 and §3); (iii) by using a refinement of these ten partitions to label the vertices of the 600-cell and the 120-cell so that H_4 symmetry becomes more transparent (§2); and (iv) by providing new ways to look at the 600-cell, in the hope of finding even more avenues towards understanding H_4 geometry (§§6 and 7 and also [10].)

We begin with our ten Facts:

Fact 1: The 600-cell is a regular H_4 polytope that consists of 120 vertices, 720 edges, 1200 triangular faces and 600 tetrahedral cells. ([12], p. 386; [5], p. 153)

Fact 2: The 600-cell contains exactly 25 24-cells, 75 16-cells and 75 8-cells, with each 16-cell and each 8-cell lying in just one 24-cell. (§3)

¹Throughout this paper we identify a polytope with its vertex set.

Fact 3: The rotation group of the 600-cell is the central product $2A_5 \circ 2A_5 \cong 2(A_5 \times A_5)$ ([11], p. 17) so the full symmetry group (including reflections) of a 600-cell H is $\text{Aut}(H) \cong 2(A_5 \times A_5)2$.

Fact 4: The subgroup of $\text{Aut}(H)$ that fixes a vertex is isomorphic to $2 \times A_5$ and the subgroup that stabilizes a 24-cell inscribed into a 600-cell is isomorphic to $2(A_4 \times A_4)2$. (§7)

Fact 5: The 25 24-cells can be placed in a 5×5 array, so that each row and each column of the array partition the 120 vertices of the 600-cell into five disjoint 24-cells. The rows and columns of the array are the only ten such partitions of the 600-cell. ([13], p. 231, stated without proof, and §§2, 3)

Fact 6: The quotient group $\text{Aut}(H)/\{\pm 1\} \cong (A_5 \times A_5)2$ is isomorphic to a subgroup of the symmetric group S_{10} , where the first A_5 permutes five symbols 1, 2, 3, 4 and 5 and fixes five symbols 6, 7, 8, 9 and X; the second A_5 permutes 6, 7, 8, 9 and X and fixes 1, 2, 3, 4 and 5; and elements outside the subquotient $A_5 \times A_5$ interchange the pentads $\{1, 2, 3, 4, 5\}$ and $\{6, 7, 8, 9, X\}$. (§§2, 3)

Fact 7: The 25 24-cells of a 600-cell may be labeled by duads $(i j)$, where $1 \leq i \leq 5$ and $6 \leq j \leq X$, and the ± 60 vertices² of the 600-cell may be labeled by five duads $(1 j_1)(2 j_2)(3 j_3)(4 j_4)(5 j_5)$ where $(j_1, j_2, j_3, j_4, j_5)$ is an even permutation of $(6, 7, 8, 9, X)$. The symmetry group $\text{Aut}(H)$ permutes the 24-cells and the vertices of the 600-cell as $(A_5 \times A_5)2 \cong \text{Aut}(H)/\{\pm 1\}$ permutes their labels. (§2)

Fact 8: The 120 vertices of a 600-cell may be viewed as the 120 *icosians*, the units of a quaternion algebra whose coefficients lie in $\mathbb{Q}(\sqrt{5})$. ([6], p. 74)

Fact 9: A 600-cell H can be "embedded" in the E_8 lattice by means of a "norm reduction" map, so that the 120 vertices of H become 120 of the 240 root vectors of E_8 . The remaining 120 root vectors are the vertices of the embedded 600-cell φH (a scale copy of H), where φ is the golden ratio $\frac{1}{2}(1+\sqrt{5})$. ([15], p. 59)

Fact 10: The "embeddings" of the 600-cells H and φH in E_8 turn $E_8/2E_8$ (an 8-space over \mathbf{F}_2) into a 4-space over \mathbf{F}_4 whose 85 points are labeled by the ± 60 vertices and the 25 24-cells of H . (§7)

In addition, a few other aspects of H_4 geometry (of lesser importance) are italicized in §§2-4.

2 The Labeling of H_4

In this section (much of which follows §4.27 of [6]) we exploit the algebraic properties of the 600-cell that are evident when we view its 120 vertices as the 120 icosians—that is, when we consider the vertices

²By " ± 60 vertices" we mean the 60 pairs $\{v, -v\}$ of opposite vertices.

$$\begin{aligned}
& (\pm 1, 0, 0, 0)^S, & (4 \text{ pairs of vertices}) \\
& \frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1), & (8) \\
& \frac{1}{2}(0, \pm 1, \pm \varphi, \pm \varphi^{-1})^A & (48)
\end{aligned}$$

as a set of 120 quaternions closed under multiplication, where the four coordinates are scalars for $(1, i, j, k)$ and where $\varphi = \frac{1}{2}(1+\sqrt{5})$, $\varphi^{-1} = \frac{1}{2}(-1+\sqrt{5})$, $S \equiv$ all permutations of the coordinates and $A \equiv$ all even permutations are permitted. ([6], p. 74; [11], p. 16)

We will let $2A_5$ denote both the set of 120 vertices and also the symmetry subgroup these vertices generate, and we will sometimes write $2A_{5L}$ or $2A_{5R}$ to emphasize that $2A_5$ is acting by left or right icosian multiplication, respectively. Similarly $2A_4$ will denote both the sub-polytope of 24 vertices (a 24-cell with 8 vertices $(\pm 1, 0, 0, 0)^S$ and 16 vertices $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$) and also the subgroup these vertices generate, with $2A_{4L}$ or $2A_{4R}$ specifically denoting left or right multiplication. It will be clear from the context which meaning is intended.

We begin with the 24-cell $2A_4$. If g is an icosian of order 5 then $\{g^i\}$ ($0 \leq i \leq 4$) is a left transversal of $2A_4$ in $2A_5$ and $\{g^{-j}\}$ ($0 \leq j \leq 4$) is a right transversal. Since $2A_4$ is stable under both $2A_{4L}$ and $2A_{4R}$ and since $2A_4$ is self-normalizing in $2A_5$, we obtain 25 distinct 24-cells $\{g^i 2A_4 g^{-j}\}_{0 \leq i, j \leq 4}$ under left and right multiplication by $2A_5$; that is, the rotation subgroup $2(A_5 \times A_5)$ of $\text{Aut}(H)$ yields 25 different 24-cells. In the next section we prove that these 25 are the only 24-cells within a 600-cell.

These 25 24-cells may be arranged in a 5×5 array (**Fact 5**) by letting the $(i, j)^{th}$ entry be the 24-cell $g^i 2A_4 g^{-j}$, so that A_{5L} permutes the rows of the array and A_{5R} permutes the columns. Since the cosets of $2A_4$ in $2A_5$ contain each element once, the five 24-cells $\{g^i 2A_4\}$ (the first column of the array) form one of Schoute's partitions of the 600-cell; since right multiplication by any right coset representative g^{-j} yields a symmetry of the 600-cell, each column of the array is also a partition. But the same arguments apply to the top row $\{2A_4 g^{-j}\}$ of the array and to each of the five rows in general. Thus we obtain ten partitions of the 600-cell into disjoint 24-cells—the five rows and five columns of the array—and in the next section we prove that these ten are the only ones possible.

In the meantime we'll use the array and its ten partitions to label the 60 vertex pairs and the 25 24-cells of the 600-cell, creating a simple way to calculate in the quotient group $\text{Aut}(H)/\{\pm 1\} \cong (A_5 \times A_5)2$. First, we label the five rows of the array by the symbols 1, 2, 3, 4 and 5 and the five columns by 6, 7, 8, 9 and X, so that each of Schoute's partitions corresponds to one of the ten symbols. Next, we label each of the 25 24-cells in the array by a duad $(i j)$, where $1 \leq i \leq 5$ and $6 \leq j \leq X$, according to its position (**Fact 7**).

Furthermore, since a 24-cell contains 12 vertex pairs $\{\pm v\}$, each of the 60 vertex pairs of the 600-cell must lie in five of the 25 24-cells, so we can name a vertex (pair) by the five different 24-cells that contain it. Since all ± 60 vertices of the 600-cell appear exactly once in each row and once in each column, each vertex (pair) of the 600-cell is labeled by five duads $(1 j_1)(2 j_2)(3 j_3)(4 j_4)(5 j_5)$ where $(j_1, j_2, j_3, j_4, j_5)$ is some permutation of $(6, 7, 8, 9, X)$. The unit icosian $1_I = (1, 0, 0, 0)$ is labeled $(16)(27)(38)(49)(5X)$, as the 24-cells along the main diagonal of the array are the five 24-cells $g^i 2A_4 g^{-i}$ that contain $g^i 1_I g^{-i} = 1_I$.

If a vertex v has label $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$, then v_R (right multiplication by the icosian v) must send partition 6 to partition j_1 , and sends 7 to j_2 , etc., since v_R sends the 24-cell (16) that contains the unit icosian 1_I to the 24-cell in row 1 that contains v , namely, $(1\ j_1)$. Thus v_R induces an even permutation of $\{6, 7, 8, 9, X\}$. Alternatively, the 24-cells that contain $\{\pm v\}$ can instead demonstrate how the rows are permuted by v_L : if the same five duads $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$ are now written (in a different order, perhaps) as $(i_1\ 6)(i_2\ 7)(i_3\ 8)(i_4\ 9)(i_5\ X)$, then evidently v_L sends partition 1 to partition i_1 , sends 2 to i_2 , etc. Thus v_L induces an even permutation of $\{1, 2, 3, 4, 5\}$, and so the permutations induced by v_R and v_L are indicated by the label of v .

The 60 reflections that generate $\text{Aut}(H)$ are also easy to describe using the ten symbols. In the next section we will show that elements in the odd half of $\text{Aut}(H)$ —the elements outside the rotation subgroup $2(A_5 \times A_5)$ of index 2—interchange the rows and columns of our 5×5 array. In other words, each of these elements (including the 60 reflections) interchanges the symbols $\{1, 2, 3, 4, 5\}$ with $\{6, 7, 8, 9, X\}$. But for any vertex v of the 600-cell, the reflection r_v (that uses v^\perp as its hyperplane of reflection) stabilizes each 24-cell that contains v , since $2A_4$ is itself a closed reflection group; and because the five 24-cells $(1\ j_1)$, $(2\ j_2)$, $(3\ j_3)$, $(4\ j_4)$, $(5\ j_5)$ maintain their positions in the 5×5 array, the reflection r_v permutes Schoute's ten partitions as $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$. Hence the label of a vertex pair $\{\pm v\}$ displays the permutation induced by r_v , and the entire quotient group $\text{Aut}(H)/\{\pm 1\} \cong (A_5 \times A_5)2$ obtains a straightforward description as a transitive but imprimitive subgroup of S_{10} that stabilizes the two pentads $\{1, 2, 3, 4, 5\}$ and $\{6, 7, 8, 9, X\}$ (**Fact 6**).

A similar process labels the 300 pairs of vertices of the 120-cell. We start with the tetrahedral cell of the 600-cell that has vertices

$$\begin{aligned} &\frac{1}{2}(\varphi, 1, \varphi^{-1}, 0), \\ &\frac{1}{2}(\varphi, 1, -\varphi^{-1}, 0), \\ &\frac{1}{2}(1, \varphi, 0, \varphi^{-1}) \text{ and} \\ &\frac{1}{2}(1, \varphi, 0, -\varphi^{-1}). \end{aligned}$$

The center $\frac{1}{4}(\varphi+1, \varphi+1, 0, 0)$ of this tetrahedron is then a vertex of the dual of the 600-cell—that is, a vertex of the 120-cell. If we rescale we obtain the standard coordinates $(2, 2, 0, 0)$ of a vertex v_1 of the 120-cell, and if we rotate v_1 by elements of $2A_4$ (acting on the left and/or the right) we obtain a 24-cell $C = 2A_4 v_1 2A_4 = \{(\pm 2, \pm 2, 0, 0)^S\}$ as a sub-polytope of the 120-cell.

As before, we obtain 25 distinct 24-cells $\{g^i C g^{-j}\}$ permuted by $\text{Aut}(H)$ in a 5×5 array; since $2(A_5 \times A_5)$ is transitive on the 600 vertices of the 120-cell these 25 24-cells are mutually disjoint. Hence each 24-cell (and each of its 12 pairs of vertices) can initially be labeled by a duad $(i\ j)$, where $1 \leq i \leq 5$ and $6 \leq j \leq X$. If we further label each vertex (pair) by appending the label of each of its four neighbors—the four other vertices with which it shares an edge—then each of the ± 300 vertices of the 120-cell is uniquely labeled by $1+4$ duads. For example, $(38)|(16)(27)(4X)(59)$ denotes the vertex in the 24-cell (38) whose four neighbors lie in the 24-cells $\{(16), (27), (4X), (59)\}$. Naturally the labels for

the vertices of the 120-cell are "duals" of the labels for the vertices of the 600-cell, in the sense that our example's duads $\{(38), (16), (27), (4X), (59)\}$ are transformed to $\{(16), (27), (38), (49), (5X)\}$ (the duads for the icosian unit 1_I of the 600-cell) by an odd permutation of $\{6, \dots, X\}$, whereas labels for vertices of the 600-cell are conjugate under even permutations.

Finally, since the first column of the 120-cell's array is the orbit (under icosian multiplication $2A_{5L}$) of a single vertex of the 120-cell, the 120 vertices in the first column obtain the geometry of the icosians, i.e., they form a 600-cell. The same is true for each of the four other columns, which shows that *the 600 vertices of the 120-cell may be partitioned into five disjoint 600-cells*. Likewise, by using $2A_{5R}$ we find that the five rows partition the 120-cell into disjoint 600-cells as well.

3 An Overdue Proof of Schoute's Result

In this section we prove Schoute's stated result that there are exactly ten ways to partition the 120 vertices of a 600-cell into five disjoint 24-cells ([13], p. 231). To keep things simple, we'll use the standard coordinates for the 600-cell (i.e., twice the icosian coordinates of §2) where the inner products of vertices are the natural inner product divided by 2.

We begin by proving **Fact 2**.

Theorem: The 600-cell contains exactly 25 24-cells, 75 16-cells and 75 8-cells, with each 16-cell and each 8-cell lying in just one 24-cell.

Proof: To begin with, $2(A_5 \times A_5)2$ is transitive on the 120 vertices of the 600-cell, since given any two vertices v and w , icosian multiplication by $(v^{-1}w)_R \in 2A_{5R}$ sends v to w . Furthermore, given a vertex v its stabilizer $\text{Stab}(v)$ is transitive on the set of all vertices that have a given inner product with v ; for example, if we let $v = 1_I = (2, 0, 0, 0)$ then the monomial subgroup $2^3:A_3 \leq \text{Stab}(v)$ (which leaves the first coordinate untouched) is itself transitive on the set \mathcal{S}_φ of 12 vertices that have inner product φ with v , and $2^3:A_3$ is likewise transitive on each set $\mathcal{S}_{-\varphi}$, $\mathcal{S}_{\varphi^{-1}}$ and $\mathcal{S}_{-\varphi^{-1}}$, which are the sets of vertices whose inner product with v is $-\varphi$, φ^{-1} or $-\varphi^{-1}$, respectively.

This monomial subgroup has two orbits on the set of 30 vertices orthogonal to v ; these are the six vertices of shape $(0, 2, 0, 0)$ and the 24 of shape $(0, 1, \varphi, \varphi^{-1})$. But if w is the vertex $(0, -1, \varphi, \varphi^{-1})$ then the reflection r_w (that uses w^\perp as its hyperplane of reflection) fixes v and fuses the two orbits, so that $\text{Stab}(v)$ is transitive on vertices orthogonal to v . Similarly, the monomial subgroup $2^3:A_3$ has two orbits on the set of 8+12 vertices that have inner product 1 with v (these orbits consist of vertices with shapes $(1, 1, 1, 1)$ and $(1, 0, \varphi^{-1}, \varphi)$) and once again r_w fuses the orbits. The same is clearly true for the remaining orbit under $\text{Stab}(v)$, which is the set of 20 vertices that have inner product -1 with v .

Now, since each of the ± 60 vertices of the 600-cell is orthogonal to 15 others, the 600-cell contains $60 \cdot 15 / 2 = 450$ pairs of orthogonal vertices, and these are

permuted transitively by $2(A_5 \times A_5)2$ since $\text{Stab}(v)$ is transitive on the ± 15 vertices orthogonal to v . If we choose one of these orthogonal pairs—say, the vertices $(\pm 2, 0, 0, 0)$ and $(0, \pm 2, 0, 0)$ —then only two of the remaining 58 vertices are orthogonal to both (these are $(0, 0, \pm 2, 0)$ and $(0, 0, 0, \pm 2)$, of course) so each of the 450 orthogonal pairs lies in a unique tetrad of mutually orthogonal vertices, i.e., each pair lies in a unique 16-cell. Since a 16-cell contains $\binom{4}{2} = 6$ such pairs, the 600-cell contains exactly $450/6 = 75$ 16-cells.

But each vertex (pair) in a 24-cell is orthogonal to three others and has inner product ± 1 with the remaining eight, so each vertex in a 24-cell lies in exactly one of the three 16-cells in the 24-cell. This inner product structure also means that we can complete a 16-cell to a 24-cell only by adding vertices that have inner products ± 1 with all four vertices of our given 16-cell. For example, if we start with the 16-cell $\{(\pm 2, 0, 0, 0)^S\}$ then the ± 8 other vertices that comprise a 24-cell must have coordinates that are all ± 1 , and so we can only add the ± 8 vertices $(\pm 1, \pm 1, \pm 1, \pm 1)$ from the 600-cell to obtain a 24-cell. Thus each 16-cell lies in a unique 24-cell, and a 600-cell therefore contains exactly $75/3 = 25$ 24-cells.

Moreover, since an 8-cell is comprised of two 16-cells, where a vertex in either 16-cell has inner product ± 1 with each vertex in the other 16-cell, an 8-cell must consist of two of the three 16-cells within the same 24-cell. Hence there are $\binom{3}{2} \cdot 25 = 75$ 8-cells in a 600-cell. And because a 16-cell lies in a unique 24-cell, each 8-cell (being the union of two 16-cells) must lie in a unique 24-cell as well. QED

Beyond that, the structure of the 5×5 array described earlier makes it easy to show that *each of the 25 24-cells is disjoint from 8 24-cells and intersects the other 16*. Certainly each 24-cell is disjoint from the $4 + 4$ others that lie in the same row or the same column of the array. But given a 24-cell $(i\ j)$, its stabilizer $S_{ij} \cong 2(A_4 \times A_4)$ is transitive on the four rows of the array other than i and the four columns other than j , so S_{ij} is transitive on the 16 24-cells whose labels contain neither i nor j . Thus each of these 16 24-cells has the same cardinality of intersection with (ij) , and this cardinality cannot be 0—otherwise each 24-cell would be disjoint from all the others—so that a 24-cell intersects each of the 16 24-cells outside its row and column of the array. (In the next section we'll see that each intersection is a regular hexagon.) But since two 24-cells are disjoint if and only if they lie in the same row or the same column, three or more mutually disjoint 24-cells must all lie in the same exact row or column, so the only partitions of a 600-cell into five disjoint 24-cells are the five rows and the five columns of the array. Thus Schoute was right all along: there are precisely ten partitions of a 600-cell into five disjoint 24-cells (**Fact 5**).

And now it's easy to see that elements in the odd half of $\text{Aut}(H)$ interchange the rows and columns of the array (**Fact 6**). For, each row and each column have a single 24-cell in common, whereas any two rows are disjoint (i.e., their $5 + 5$ 24-cells are distinct) and any two columns are disjoint as well. Hence the rows and columns form a bipartite graph, so $\text{Aut}(H)$ permutes Schoute's ten partitions as two pentads: the rows $\{1, 2, 3, 4, 5\}$ and the columns $\{6, 7, 8, 9,$

$X\}$. If v is any vertex of the 600-cell (so that v is labeled $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$ by the five 24-cells that contain it) then, as in §2, the reflection r_v fixes each of these five duads, so the reflection either fixes or interchanges the two symbols in each duad. If r_v mapped rows to rows and columns to columns, then r_v would fix each of the ten symbols (and thus fix each of the ± 60 vertices of the 600-cell), so we would have $r_v = \pm 1$. But this can't be true, and the contradiction shows that r_v and all the other elements of $\text{Aut}(H)$ outside $2(A_5 \times A_5)$ interchange the rows and columns of the array.

4 The Planes of the 600-cell

With §7 in mind we describe how pairs of vertices of the 600-cell can interact.

To begin with, the unit icosian 1_I (§2) is the only vertex that lies in all three 24-cells (16), (27) and (38): the 120 vertices of the 600-cell are conjugate by $2A_{5R}$ (right multiplication by icosians, which fix $\{1, 2, 3, 4, 5\}$ pointwise) and so a vertex labeled $(16)(27)(38)(4X)(59)$ cannot exist, since it would be conjugate to 1_I by only the odd permutation $(9X)$ of $\{6, 7, 8, 9, X\}$; and in general, naming any three of a vertex's five duads is sufficient to identify the vertex uniquely. Thus the labels of distinct vertices $\{\pm v\}$ and $\{\pm w\}$ cannot share three duads, i.e., pairs of vertices of a 600-cell can lie together in at most two of the 25 24-cells.

If the labels of $\{\pm v\}$ and $\{\pm w\}$ share at least one duad—if v and w are in at least one 24-cell together—then the inner product of v and w is either 0 or ± 1 . If the vertices are orthogonal then they lie in a 16-cell, and therefore in a unique 24-cell, so their labels share only one duad. It was shown in §3 that there are $60 \cdot 15 / 2 = 450$ pairs of orthogonal vertices in a 600-cell, with $\binom{4}{2}$ pairs in each of the 75 16-cells. The significance of these 450 pairs can be seen in [10].

If, on the other hand, the labels of $\{\pm v\}$ and $\{\pm w\}$ share two duads (i.e., if they lie in the same two 24-cells) then a third vertex must also lie in both 24-cells in question, since $60 / (5 \cdot 4) = 3$ vertex labels contain any two given (disjoint) duads. For example, the 24-cells labeled (16) and (27) contain the trio of vertices $(16)(27)(38)(49)(5X)$, $(16)(27)(39)(4X)(58)$ and $(16)(27)(3X)(48)(59)$. These ± 3 vertices all have inner product ± 1 with each other—if any pair were orthogonal, they'd lie in a 16-cell and therefore in a unique 24-cell—so the vertices form a (planar) regular hexagon.

A 24-cell contains $12 \cdot 8 / (3 \cdot 2) = 16$ hexagons, so *a 24-cell is disjoint from eight 24-cells and intersects each of the other 16 in six vertices that form a hexagon*. Since no two of the ± 3 vertices in a hexagon have inner product 0, each of the three 16-cells of a 24-cell contains one pair of vertices of a hexagon. A 600-cell contains $25 \cdot 16 / 2 = 200$ such hexagons, whose significance will be seen in §7.

The final possibility for two vertices v_1 and v_2 is that their labels have no duads in common—that is, if v_1 is labeled $(1\ i_1)(2\ i_2)(3\ i_3)(4\ i_4)(5\ i_5)$ and v_2 is labeled $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$ then each of the i 's is a different element of $\{6, 7, 8, 9, X\}$ than its j counterpart. Since $2A_{5R}$ is transitive on the ± 60

vertices of the 600-cell, we can find an icosian w such that w_R fixes $\{1, 2, 3, 4, 5\}$ while sending i_1 to j_1 , i_2 to j_2 , etc., so that w_R has order 5 and sends v_1 to v_2 , where we can replace v_2 by its negative if necessary. The orbit of v_1 under w_R is therefore five vertices $\{v_1, v_2, v_3, v_4, v_5\}$ that are easily shown to form a (planar) regular pentagon. Moreover, since w_R fixes $\{1, \dots, 5\}$ and is transitive on $\{6, \dots, X\}$, all 25 duads $(i j)$ appear among the labels of these five vertices—in other words, *each of the 25 24-cells of the 600-cell contains exactly one vertex of a regular pentagon.*

Naturally the five antipodal vertices $\{-v_1, -v_2, -v_3, -v_4, -v_5\}$ also form a pentagon, and the two pentagons create a regular decagon. Since each of the ± 60 vertices of a 600-cell has inner product φ with 12 others and inner product φ^{-1} with 12, a 600-cell contains $(60 \cdot 24)/(5 \cdot 4) = 72$ decagons whose significance will be seen in §7. Since each of the 720 edges of the 600-cell is also an edge of a decagon—the two vertices in question have inner product φ in either case—*each of the 720 edges of the 600-cell lies on just one of the 72 decagons.*

When we generalize the construction of §2 (which created a 5×5 array from the vertices of the 600-cell), we find that *the vertices of the 600-cell create an $n \times n$ array for each of the 600-cell's basic planar shapes*—the square (vertices v and w with $v \cdot w = 0$; $n = 5$), the triangle ($v \cdot w = \pm 1$; $n = 10$) and the pentagon ($v \cdot w = \pm \varphi$; $\pm \varphi^{-1}$, $n = 6$.)

We begin with any subgroup $K \leq A_5$ of the icosians. (We work projectively for the rest of this section and write A_5 instead of $2A_5$, etc.) Let $\{g_i\}$ be a left transversal of K in A_5 , so that $\{g_i^{-1}\}$ is a right transversal. Since our goal is to form an array of sets $\{g^i K g^{-j}\}$ that are distinct for all choices of i, j we require K to be self-normalizing.

The only self-normalizing subgroups of A_5 are the normalizers of the Sylow- p subgroups ($p = 2, 3$ or 5), so we consider the 600-cell from the perspective of these primes. To each of the three finite fields \mathbf{F}_p we adjoin the roots of the polynomial $f(x) = x^2 - x - 1$ in order to incorporate the 600-cell's scalar φ , which is a root of $f(x)$. Since $f(x)$ is irreducible mod 2 and mod 3 but splits mod 5, we obtain fields \mathbf{F}_q , where $q = 4, 9$ or 5 , depending on p . For the primes 2, 3 and 5 the group A_5 contains $q + 1 = 5, 10$ and 6 Sylow- p subgroups, respectively.

Let N denote both the subgroup $N(P)$ (the normalizer of a Sylow- p subgroup P of A_5 , where p is 2, 3 or 5) and also the set of vertices we obtain from $N(P)$ when we consider the ± 60 vertices of the 600-cell as the icosians. Let $\{g_i\}$ be a left transversal of N in A_5 , so that $\{g_i^{-1}\}$ is a right transversal. As in §2, we obtain from N a $(q + 1) \times (q + 1)$ array (q is 4, 9 or 5, depending on p) where the $(i, j)^{th}$ entry is the set $g^i N g^{-j}$ of vertices of the 600-cell. Since N is self-normalizing, these sets are all distinct. Furthermore, since $\{g_i\}$ is a transversal of N in A_5 , the first column $\{g_i N\}$ of the array partitions the 120 vertices of the 600-cell into $q + 1$ disjoint sets. But right multiplication by the icosian g^{-j} is a symmetry of the 600-cell, so each column of the array is a partition; of course the same is also true for each row.

The case $p = 2$ was already considered in §3, where $N \cong A_4$ creates a 5×5

array that displays Schoute's ten partitions. When $p = 3$, the normalizer N of a Sylow- p subgroup of A_5 has order 6; for example, we can take this subgroup (of A_{5R}) to be $\langle (89X), (67)(89) \rangle$. In this case the ± 6 vertices N of the 600-cell are the two hexagons (of ± 3 vertices)

$$h = \{(16)(27)(38)(49)(5X), (16)(27)(39)(4X)(58), (16)(27)(3X)(48)(59)\}$$

$$\text{and } h' = \{(17)(26)(38)(4X)(59), (17)(26)(39)(48)(5X), (17)(26)(3X)(49)(58)\}.$$

Since each vertex of h shares a single duad with each vertex of h' , the two hexagons are orthogonal, so *the 200 hexagons in a 600-cell form 100 orthogonal pairs that create a 10×10 array.*

When $p = 5$, the normalizer N of a Sylow- p subgroup of A_5 has order 10; for example, we can take this subgroup to be $\langle (6789X), (7X)(89) \rangle$. In this case, the ± 10 vertices N are the decagons

$$d = \{(16)(27)(38)(49)(5X), (17)(28)(39)(4X)(56), (18)(29)(3X)(46)(57),$$

$$(19)(2X)(36)(47)(58), (1X)(26)(37)(48)(59)\}$$

$$\text{and } d' = \{(16)(2X)(39)(48)(57), (17)(26)(3X)(49)(58), (18)(27)(36)(4X)(59),$$

$$(19)(28)(37)(46)(5X), (1X)(29)(38)(47)(56)\}.$$

Since each vertex of d shares a single duad with each vertex of d' , the two decagons are orthogonal, so *the 72 decagons in a 600-cell form 36 orthogonal pairs that create a 6×6 array.*

Finally, since $L_2(q) \times L_2(q) \cong O_4^+(q)$ ([2], p. xii) and since $L_2(4) \cong L_2(5) \cong A_5$ while $L_2(9) \cong A_6$, each group $O_4^+(q).2$ ($q = 4, 9$ or 5) contains a copy of $\text{Aut}(H)/\{\pm 1\} \cong (A_5 \times A_5).2$. So the arrays can be described in terms of the orthogonal groups $O_4^+(q)$; the case $q = 4$ is described in §7 and $q = 5$ in [10].

5 The Geometry of E_8

In this section we collect some results about the E_8 lattice ([3], p. 120)—or rather, some results about the 8-space $E_8/2E_8$ over \mathbf{F}_2 —that will enable us in §7 to view the ± 60 vertices and the 25 24-cells of a 600-cell in a new way. We assume some familiarity with the basic results and terminology for vector spaces and quadratic forms over \mathbf{F}_2 .

From [1, p. 2], the 270 isotropic 4-spaces of $E_8/2E_8$ are permuted in two orbits of 135 each by $O_8^+(2)$, and given a pair $\{V_1, V_2\}$ of disjoint isotropic 4-spaces (disjoint other than the zero vector, of course) there are 28 others disjoint from both V_1 and V_2 . These 28 can be labeled by duads on eight letters. Since two of the 28 4-spaces are disjoint when their duads share a letter, there are two different ways to complete the pair $\{V_1, V_2\}$ into a maximal set of mutually disjoint 4-spaces (where the next two 4-spaces added towards a maximal set may be labeled by duads (ab) and (ac), and where any two duads that we add must share a letter): (i) we can add the seven 4-spaces $\{(ab), (ac), (ad), \dots, (ah)\}$ and complete the original pair $\{V_1, V_2\}$ into a set of nine 4-spaces, or (ii) we can add the three 4-spaces $\{(ab), (ac), (bc)\}$ and complete the pair into a set of five 4-spaces.

The automorphism group of the set of nine 4-spaces is the maximal subgroup A_9 of $O_8^+(2)$ [7, p. 540]. If we relabel the nine 4-spaces $V_1, V_2, (ab), (ac), (ad), \dots, (ah)$ by $1, 2, \dots, 9$, respectively, then the remaining 126 4-spaces in the 135-orbit can be labeled by the $\binom{9}{5} = 126$ pentads of the nine symbols, according to which five of the nine 4-spaces they intersect: any pair of the 135 4-spaces are either disjoint or intersect in a 2-space (which contains three non-zero vectors), and since our nine disjoint 4-spaces contain all $135 = 9 \cdot 15$ isotropic vectors among them, each of the 126 4-spaces has its 15 non-zero vectors allocated among exactly five of the nine. But then the five disjoint 4-spaces of case (ii) above are $\{1, 2, 3, 4, (56789)\}$, and the subgroup of A_9 that stabilizes this pentad of 4-spaces is $(A_4 \times A_5)2$.

The only 4-spaces that intersect all four of the 4-spaces $\{1, 2, 3, 4\}$ are $\{(12345), (12346), (12347), (12348), (12349)\}$. But these five 4-spaces are also mutually disjoint³, and each one intersects all five 4-spaces $\{1, 2, 3, 4, (56789)\}$ in a 2-space of three (non-zero) vectors; in other words, the 15 vectors in any of the ten 4-spaces consist of three vectors from each 4-space in the other pentad, and so both pentads contain the same $5 \cdot 15 = 75$ isotropic vectors. Since $O_8^+(2)$ is transitive on tetrads of disjoint 4-spaces there must exist a group element that interchanges the two pentads, so the automorphism group of the two pentads is the maximal subgroup $(A_5 \times A_5)2^2$ of $O_8^+(2)$ [2, p. 85].

Furthermore, the intersection pattern of the two pentads creates the same 5×5 array as the 25 24-cells of the 600-cell. We can arrange the 75 isotropic vectors common to both pentads into a 5×5 array of 25 2-spaces (each having three non-zero vectors), where the five rows are headed by the first pentad $\{V_i\}$, the five columns are headed by the second pentad $\{W_j\}$, and the $(ij)^{th}$ entry is the 2-space ℓ_{ij} that contains the three vectors that lie in both V_i and W_j , as in Figure 1.

	W_1	W_2	.	.	W_5
V_1	ℓ_{11}	ℓ_{12}	.	.	ℓ_{15}

V_5	ℓ_{51}	ℓ_{52}	.	.	ℓ_{55}

Figure 1. Two pentads of isotropic 4-spaces and their 25 lines (2-spaces) of intersection

The identification of the 25 isotropic 2-spaces in Figure 1 with the 25 24-cells of a 600-cell will be made apparent in §7.

6 Changing the Norms of Vectors

In this section we define maps from quadratic fields to \mathbb{Q} in a variety of ways, which leads a variety of embedded H_4 polytopes. Our main objective will be to

³Two 4-spaces are disjoint when their labels (by one of 9 monads or one of 126 pentads) intersect with even cardinality.

embed a 600-cell in the E_8 lattice.

The coordinates and inner products for H_4 polytopes lie in the golden field $\mathbb{Q}(\sqrt{5})$, making it impossible to (legitimately) embed H_4 polytopes in the E_8 lattice, whose inner products lie in \mathbb{Z} . But it was shown in [15] that a “reduced norm” for the 120 vertices of a 600-cell H identifies them as 120 of the 240 root vectors of E_8 , and the remaining 120 root vectors come from the same norm-reducing formula applied to the scaled 600-cell φH . The reduction formula is obtained by composing the standard inner product in $\mathbb{Q}(\sqrt{5})$ with a modified trace map from $\mathbb{Q}(\sqrt{5})$ to \mathbb{Q} , and the trace map of [15] identifies a vertex v in H or φH of norm $a + b\sqrt{5}$ ($a, b \in \mathbb{Q}$) with a vector $v \in E_8$ of reduced norm $a - b$. ([15], p. 57)

But [16] uses a different trace map to embed the 600-cell in E_8 ([16], p. 158), which raises the question of how many different norm reductions are possible. The answer is as follows:

Proposition: Let $\mathbb{Q}(\sqrt{n})$ ($n > 0$) be a real quadratic field. Then for $m \in \mathbb{Q}$ there exists a norm reduction from $\mathbb{Q}(\sqrt{n})$ to \mathbb{Q} that maps \sqrt{n} to m , i.e.,

$$a + b\sqrt{n} \rightarrow a + bm \quad (a, b \in \mathbb{Q})$$

if and only if $|m| < \sqrt{n}$.

Proof: The norm of a typical scalar $x + y\sqrt{n}$ in $\mathbb{Q}(\sqrt{n})$ (where $x, y \in \mathbb{Q}$) must reduce to a positive number in \mathbb{Q} . But under the map that reduces \sqrt{n} to m , this scalar’s norm of

$$(x + y\sqrt{n})^2 = x^2 + 2xy\sqrt{n} + ny^2 \quad \text{reduces to} \\ x^2 + 2xym + ny^2 = (x+my)^2 + (n-m^2)y^2.$$

When $(x, y) = (-m, 1)$ this norm is $n - m^2$, so we must have $|m| < \sqrt{n}$. But in that case—whenever $|m| < \sqrt{n}$ —if a vector in a space over $\mathbb{Q}(\sqrt{n})$ has coordinates $\{(x_i + y_i\sqrt{n})\}$, we can split each coordinate into two coordinates $(x_i + my_i, \sqrt{n - m^2}y_i)$ to obtain a norm reduction that maps $\sqrt{n} \rightarrow m$. QED

In the particular case of lattices over $\mathbb{Q}(\sqrt{5})$ we obtain \mathbb{Q} -lattices under the norm-reduction map

$$a + b\sqrt{5} \rightarrow a + bm \quad (m \in \mathbb{Q})$$

for any $|m| < \sqrt{5}$, and for an integral lattice over $\mathbb{Z}[\varphi]$ we obtain an integral \mathbb{Z} -lattice for $m \in \{0, \pm 1, \pm 2\}$, where the coordinates for these values of m are given by $(x_i + y_i\sqrt{5}) \rightarrow \{(x_i, y_i\sqrt{5}), (x_i \pm y_i, 2y_i), \text{ or } (x_i \pm 2y_i, y_i)\}$ respectively. Note that the natural norm may need to be doubled when $m \in \{0, \pm 2\}$ since $\varphi = \frac{1}{2}(1 + \sqrt{5})$ is an integer in the golden field that reduces to $\frac{1}{2}, \frac{3}{2}, -\frac{1}{2}$ for $m = 0, 2, -2$, respectively.

The following are a few examples of how the Proposition may be used to transform H_4 polytopes into lattices over \mathbb{Q} .

Example 1: We begin with the 600-cell. Then the norm-changing maps

$$a + b\sqrt{5} \rightarrow a - b, \quad a + b\sqrt{5} \rightarrow a + b \quad (m = -1, 1, \text{ resp.})$$

both embed the 600-cell as 120 of the 240 norm 2 vectors in E_8 . Note that [15] used $m = -1$ (and mapped $\varphi = \frac{1}{2}(1 + \sqrt{5})$ to 0) whereas [16] used $m = 1$ (and mapped $\varphi^{-1} = \frac{1}{2}(-1 + \sqrt{5})$ to 0.)

Example 2: Using the 600-cell again, we can choose $m = 0$ instead, so that $a + b\sqrt{5} \rightarrow a$. In this case, if we double the natural inner product then we obtain a rank 8 \mathbb{Z} -lattice L generated by ± 60 norm 4 vectors, each of which has inner product

$$\begin{array}{ll} \pm 4 & \text{with 1 pair of vectors;} \\ \pm 2 & 20; \\ \pm 1 & 24; \text{ and} \\ 0 & 15. \end{array}$$

Now, we can find four vertices $\{v_1, v_2, v_3, v_4\}$ of the 600-cell H so that all ± 60 vertices of H are $\mathbb{Z}[\varphi]$ -linear combinations of the v_i 's. It's easily verified that the $\mathbb{Z}[\varphi]$ -lattice H generated by these four vertices is self-dual, so if we let $\{w_1, w_2, w_3, w_4\}$ denote the dual basis of H (i.e., inner products satisfy $v_i \cdot w_j = \delta_{ij}$) then the w_j 's are $\mathbb{Z}[\varphi]$ -linear combinations of the v_i 's. When we apply our norm reduction procedure to the lattice H (still using $m = 0$) each coordinate $\{a + b\sqrt{5}\}$ of a vector v in H splits into two coordinates $\{a, b\sqrt{5}\}$ of the norm-reduced vector v_L of the \mathbb{Z} -lattice L , as above.

But L is generated by the eight vectors $\{(v_i)_L, (\varphi v_i)_L\}$, and since we have $(v_i)_L \cdot (w_j)_L = 2\delta_{ij}$ (recall that the natural inner product has been doubled), and since we also have $(\varphi v_i)_L \cdot (w_j)_L = \delta_{ij} = (v_i)_L \cdot (\varphi w_j)_L$ and $(\varphi v_i)_L \cdot (\varphi w_j)_L = 3\delta_{ij}$, the basis dual to $\{(v_i)_L, (\varphi v_i)_L\}$ is given by $\{\frac{1}{5}(3w_i - \varphi w_i)_L, \frac{1}{5}(-w_i + 2\varphi w_i)_L\}$.

Thus $d(L) = 5^4$, and L must be the unique 8-dimensional rootless even lattice Λ with $d(\Lambda) = 5^4$ [8, p. 3].

Example 3: The 720 vertices of the rectified 600-cell (whose vertices are the midpoints of each of the 720 edges of the 600-cell) can be given coordinates of shapes $(0, 0, 2\varphi, 2\varphi^2)$, $(1, 1, \varphi^3, \varphi^3)$, $(0, 1, \varphi, 1+3\varphi)$, $(0, \varphi^2, \varphi^3, 2+\varphi)$, $(1, \varphi, 2\varphi^2, \varphi^2)$ and $(\varphi, \varphi^2, 2\varphi, \varphi^3)$. These vertices have norm $20 + 8\sqrt{5}$, and by choosing $m = -2$ (which maps $a + b\sqrt{5} \rightarrow a - 2b$) the rectified 600-cell is embedded as 720 norm 4 vectors of E_8 . In fact, it's easy to show that the 2160 norm 4 vectors of E_8 consist of five embedded H_4 polytopes: two 600-cells (of 120 vertices each); two 120-cells (of 600 vertices each); and a rectified 600-cell of 720 vertices.

7 $\text{Aut}(H_4) \rightarrow O_4^+(4).2$

The variety of norm-reducing maps (and their resulting embeddings) means that a lattice—or polytope—can be viewed in a number of new ways. This is the case when we embed 600-cells H and φH in E_8 and project them to $E_8/2E_8$:

the 8-space over \mathbf{F}_2 employs the geometry of the 600-cell to become a 4-space over \mathbf{F}_4 (**Fact 10**), as we now demonstrate.

Let $\varepsilon: H \rightarrow E_8$ denote the embedding of H in E_8 used in [15] (i.e., the norm reduction map of §6, Example 1 that uses $m = -1$), and similarly let $\varepsilon_\varphi: \varphi H \rightarrow E_8$ denote the embedding of φH . We now construct a map $\Phi: E_8 \rightarrow E_8$ that sends the each of the 120 root vectors embedded from H onto its counterpart—the corresponding root vector embedded from φH —by defining $\Phi(\varepsilon(v)) = \varepsilon_\varphi(\varphi v)$ for any vertex $v \in H$. Note that since $\text{Aut}(H) \cong 2(A_5 \times A_5)2$ permutes H and φH exactly the same way, Φ commutes with all of $2(A_5 \times A_5)2 \leq \text{Aut}(E_8)$.

In terms of coordinates, if $\alpha = a + b\sqrt{5}$ ($a, b \in \mathbb{Q}$) is one of the four coordinates of a vertex $v \in H$ then α splits into two coordinates $(a - b, 2b)$ when H is embedded in E_8 , as in §6. Likewise, the scaled coordinate $\varphi\alpha = \frac{1}{2}[(a + 5b) + (a + b)\sqrt{5}]$ (the corresponding coordinate of φv) splits into two coordinates $(2b, a + b)$ under the embedding ε_φ . Setting $x = a - b$ and $y = 2b$, we find that Φ maps a split coordinate pair (x, y) to $(y, x + y)$.

Since the vertices $v_0 = (2, 0, 0, 0)$ and $\varphi v_0 = (1 + \sqrt{5}, 0, 0, 0)$ of H embed in E_8 as the orthogonal root vectors $(2, 0, \dots, 0)$ and $(0, 2, 0, \dots, 0)$ and since Φ commutes with all of $2(A_5 \times A_5)2 \leq \text{Aut}(E_8)$, every root vector $\varepsilon(v)$ embedded from H is orthogonal to its counterpart $\varepsilon_\varphi(\varphi v)$. Thus $v + \Phi(v)$ is a norm 4 vector of E_8 whenever $v \in \text{Im}(\varepsilon)$.

In addition, since the 120 root vectors embedded from H generate E_8 , Φ extends to a (well-defined) endomorphism of E_8 , and so we now have $\Phi(g(v)) = g(\Phi(v))$ for all $g \in 2(A_5 \times A_5)2$ and all vectors $v \in E_8$. Since Φ maps a pair (x, y) to $(y, x + y)$, Φ^2 maps (x, y) to $(x + y, x + 2y)$; thus Φ retains the defining property of $\varphi = \frac{1}{2}(1 + \sqrt{5})$:

$$(i): \Phi^2 = \Phi + 1, \quad \text{so that} \quad (ii): \Phi^3 = 2\Phi + 1$$

If $\overline{\Phi}$ denotes the endomorphism of $E_8/2E_8$ induced by Φ , then by (ii), $\overline{\Phi}^3 = 1$, and by (i) the orbit of any vector $v \in E_8/2E_8$ under $\overline{\Phi}$ is the set $\{v, \overline{\Phi}(v), v + \overline{\Phi}(v)\}$ —that is, the three non-zero vectors in a 2-space over \mathbf{F}_2 . But then $\overline{\Phi}$ is an automorphism that transforms our 8-space over \mathbf{F}_2 into a 4-space over \mathbf{F}_4 : we have an element $\omega (= \overline{\Phi})$ where $\omega^3 = 1$, and the orbit of any vector v under ω is the set $\{v, \omega v, \overline{\omega} v (= v + \omega v)\}$. In addition, as we'll see below, the quadratic form on the 8-space over \mathbf{F}_2 gives rise to a quadratic form on our 4-space over \mathbf{F}_4 , and since Φ commutes with $2(A_5 \times A_5)2$ we also have $g(\omega v) = \omega g(v)$ for all $g \in (A_5 \times A_5)2 \cong O_4^+(4).2$.

If v is the image in $E_8/2E_8$ of a root vector embedded from H , then the three (non-zero) vectors $\{v, \overline{\Phi}(v), v + \overline{\Phi}(v)\}$ of a 2-space consist of the non-isotropic vector v that came from H , its (non-isotropic) counterpart $\overline{\Phi}(v)$ embedded from φH , and the isotropic vector $v + \overline{\Phi}(v)$ (recall that $\varepsilon(v_0) + \varepsilon_\varphi(\varphi v_0)$ is a norm 4 vector of E_8 for any vertex v_0 of H , as above.) Thus 180 of the 255 non-zero vectors of $E_8/2E_8$ form 60 2-spaces that can be labeled by the ± 60 vertices $\{\pm v\}$ of H , and these 2-spaces account for all 60 + 60 non-isotropic vectors and 60 of the 135 isotropic vectors of $E_8/2E_8$.

The 75 remaining isotropic vectors form the 25 2-spaces of Figure 1 (§5), where each 2-space ℓ_{ij} has a distinct stabilizer S_{ij} of index 25 in $(A_5 \times A_5)2 \leq \text{Aut}(E_8/2E_8)$. Since $\bar{\Phi}$ commutes with all of $(A_5 \times A_5)2$, we have $S_{ij}(\bar{\Phi}(\ell_{ij})) = \bar{\Phi}(S_{ij}(\ell_{ij})) = \bar{\Phi}(\ell_{ij})$, so that $\bar{\Phi}$ stabilizes each 2-space and therefore permutes its three (non-zero) vectors.

But the isomorphism between subgroups $(A_5 \times A_5)2$ of both $\text{Aut}(E_8/2E_8)$ and $\text{Aut}(H)/\{\pm 1\}$ means that each stabilizer $S_{ij} \leq \text{Aut}(E_8/2E_8)$ corresponds to a specific subgroup of index 25 in $\text{Aut}(H)/\{\pm 1\}$. Since these subgroups lift in $\text{Aut}(H)$ to the stabilizers of the 25 24-cells of the 600-cell, each 2-space of Figure 1 corresponds to a specific 24-cell of H .

So we have obtained **Fact 10**: embedding the 600-cell H in E_8 turns the 8-space over \mathbf{F}_2 into a 4-space over \mathbf{F}_4 whose 85 (projective) points are labeled by the 60 vertex pairs and the 25 24-cells of H .

Our 4-space over \mathbf{F}_4 derives a quadratic form Q_ω from the quadratic form Q on the 8-space over \mathbf{F}_2 . If v is any vector of $E_8/2E_8$ then we define $Q_\omega(v) = Q(v) + \omega Q(\omega v) + \bar{\omega} Q(\bar{\omega} v)$; a simple calculation shows that Q_ω is a quadratic form. If the vectors $\{v, \omega v, \bar{\omega} v\}$ are all isotropic then $Q_\omega(v) = 0$ (so the 25 points labeled by 24-cells are singular); if v is isotropic but ωv and $\bar{\omega} v$ are non-isotropic then $Q_\omega(v) = 1$; and if v is non-isotropic then $Q_\omega(v) = \bar{\omega}$ if v is the image in $E_8/2E_8$ of a vector embedded from H and $Q_\omega(v) = \omega$ when v came from φH .

Naturally the \mathbf{F}_2 quadratic form Q can be recovered from Q_ω using a trace formula: we have $Q(v) =$

$$\text{Tr}_{\mathbf{F}_4/\mathbf{F}_2}(Q_\omega(v)) = \begin{cases} 0, & \text{if } Q_\omega(v) \in \{0, 1\} \\ 1, & \text{if } Q_\omega(v) \in \{\omega, \bar{\omega}\} \end{cases}$$

The isomorphism $\text{Aut}(H_4)/\{\pm 1\} \cong O_4^+(4).2$ means that certain types of geometric objects are favored in the 600-cell; these correspond to the $\frac{4^4-1}{4-1} = 85$ 1-spaces ("points")—and by duality the 85 3-spaces ("planes")—plus the $\frac{85 \cdot 84}{5 \cdot 4} = 357$ 2-spaces ("lines") in a 4-space over \mathbf{F}_4 . Each line contains $\frac{4^2-1}{4-1} = 5$ points and each plane contains $\frac{4^3-1}{4-1} = 21$ points.

The following proposition will help determine the structure of the lines and planes in terms of their constituent points.

Proposition: The subgroup S_C of $\text{Aut}(H)$ that stabilizes a 24-cell C in a 600-cell H is isomorphic to $2(A_4 \times A_4)2$, and the subgroup S_v that fixes a vertex v of H is isomorphic to $2 \times A_5$ (**Fact 4**).

Proof: $\text{Aut}(H) \cong 2(A_5 \times A_5)2$ is transitive on the 25 24-cells in their 5×5 array, so the first statement is obvious.

Since $\text{Aut}(H)$ is transitive on the 120 vertices of the 600-cell, S_v has order $14,400/120 = 120$. Furthermore, since v is negated by r_v (the reflection that uses v^\perp as its hyperplane), $-r_v$ is an element of order 2 that fixes v . But $-r_v$

is an odd permutation of $\{1, 2, \dots, 9, X\}$ that lies in the center of S_v , so that $S_v \cong \langle -r_v \rangle \times S_v^e$, where S_v^e is the even subgroup of index 2 in S_v .

Recall (§2) that the vertex v is labeled by five duads $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$, and that left multiplications A_{5L} achieve all even permutations of $\{1, 2, 3, 4, 5\}$ while right multiplications A_{5R} achieve all even permutations of $\{6, 7, 8, 9, X\}$. Thus every even permutation of the five duads (that is, every even permutation of the five 24-cells that contain the vertex v) can be achieved by an element $w_L w'_R$ of $\text{Aut}(H)$, where w is chosen so that w_L permutes $\{1, 2, 3, 4, 5\}$ exactly as the duads are to be permuted, and w' is chosen so that w'_R sends each of $\{6, 7, 8, 9, X\}$ back to its original duad partner. Since $w_L w'_R$ preserves the set of five duads, v is either fixed or negated. If v is negated we replace $w_L w'_R$ by $-w_L w'_R$, so that $w_L w'_R \in S_v^e$ and thus S_v^e maps onto A_5 . Since the order of $\langle -r_v \rangle \times S_v^e$ is 120, we must have $S_v^e \cong A_5$ and thus $S_v \cong 2 \times A_5$. QED

The identification of the 85 points as 25 24-cells (ij) and ± 60 vertices $(1\ j_1)(2\ j_2)(3\ j_3)(4\ j_4)(5\ j_5)$ makes it easy to determine the lines and planes of our 4-space, simply by observing how S_C and S_v permute these 85 objects via their labels. For example, the stabilizer S_C of a 24-cell (ij) permutes the other 24-cells in orbits of sizes 8 and 16, where the orbit of size 8 consists of the 24-cells whose labels contain either i or j . Furthermore, this subgroup permutes the 60 vertices in orbits of sizes 12 and 48, according to whether or not (ij) is one of the five duads of the vertex in question—that is, whether or not the vertex is contained in the 24-cell C .

The vertex stabilizer S_v permutes the 25 24-cells in orbits of sizes 5 and 20, where the orbit of size 5 consists of the five duads that label v (i.e., the five 24-cells that contain v .) And S_v permutes the 60 vertices in orbits of sizes $1 + 15 + 20 + 12 + 12$, depending on whether the inner product with v (modulo $\{\pm 1\}$) is 2, 0, 1, φ or φ^{-1} , respectively, as in the proof of the Theorem in §2. (Equivalently, the orbits are determined by the conjugacy class of the A_5 permutation of $\{6, 7, 8, 9, X\}$ that takes the five duads of v to the five duads of the other vertex.)

It's now a simple exercise to determine the different types of lines and planes in our 4-space over \mathbf{F}_4 . For example, let ℓ be a line (of five points) generated by a vertex-labeled point and a 24-cell-labeled point, where the vertex v is contained in the 24-cell C . The stabilizer S_v of the vertex v is isomorphic to $2 \times A_5$, so the stabilizer of both v and the 24-cell C is a subgroup $S_{C,v} = S_C \cap S_v \cong 2 \times A_4$. As above, S_C permutes the 25 24-cells in orbits of sizes $1 + 8 + 16$, and S_v permutes the 25 in orbits of sizes 5 and 20.

The 5-orbit of S_v decomposes under $S_{C,v}$ as $1 + 4$ since $S_{C,v}$ fixes C and is transitive on the four other 24-cells that contain v . The 20-orbit decomposes as $8 + 12$ since $S_{C,v}$ is transitive on the 8 of 20 duads (24-cells) that share one symbol with C , and transitive on the 12 that contain neither symbol. In particular, the line ℓ cannot contain another point labeled by a 24-cell, since $S_{C,v}$ already fixes two of the line's five points (the ones labeled by v and by C) and so must permute the other three; but $S_{C,v}$ permutes the 25 24-cells as $1 + 4 + 8 + 12$, so (other than the fixed point of C itself) the orbit sizes are too

large.

So the other three points on the line must all be labeled by vertices, and we now consider the orbits of $S_{C,v}$ on the 60 vertices. The orbit of size 15 under S_v consists of the 15 vertices orthogonal to v , three of which lie in the 24-cell C and 12 of which lie in the other four 24-cells that contain v . Since $S_{C,v} \cong 2 \times A_4$ is transitive on these four 24-cells, the orbit of size 15 splits as $3 + 12$.

The orbit of size 20 under S_v splits into orbits of sizes $8 + 12$ under S_C : the 20 vertices in this orbit have inner product 1 with v , and 8 of these 20 lie in C (and are permuted transitively by $S_{C,v}$) while the other 12 are split among the other four 24-cells that contain v . These 12 are also permuted transitively.

Finally, the orbits of size 12 under S_v must remain intact under $S_{C,v}$, since the stabilizer in S_v of one of the 12 vertices has order $120/12 = 10$; but the order of $S_{C,v}$ isn't divisible by 5, so its stabilizer subgroup can have order at most 2, resulting in the same orbit size of $24/2 = 12$.

So only one orbit of $S_{C,v}$ contains as few as three points, and these points are labeled by the three vertices of C that are orthogonal to v . Thus the line ℓ consists of a single point labeled by a 24-cell C and four points labeled by the vertices of a 16-cell in C , and so there are $75 = 25 \cdot 3$ such lines in our geometry.

Similar arguments determine the other three types of lines. The two types of planes are especially simple to identify, since by duality their stabilizers are the $25 + 60$ subgroups S_C and S_v .

Thus we find our geometry has:

- (A) $85 = 60 + 25$ points that correspond to:
 - (i) The ± 60 vertices of a 600-cell
 - (ii) The 25 24-cells of a 600-cell
- (B) $85 = 60 + 25$ planes of 21 points each, corresponding to:
 - (i)' The ± 60 vertices. The $21 = 1 + 15 + 5$ points correspond to a vertex, the 15 vertices orthogonal to it, and the 5 24-cells that contain it.
 - (ii)' The 25 24-cells. The $21 = 1 + 8 + 12$ points correspond to a 24-cell, the 8 24-cells disjoint from it, and the 12 vertices it contains.

Finally, the 357 lines honor Schoute's ten partitions alongside the three planar shapes (§4) that a 600-cell's vertices create:

- (C) $357 = 10 + 72 + 75 + 200$ lines of 5 points each, corresponding to:
 - (iii) Schoute's ten partitions of the 120 vertices of a 600-cell. The $0+5$ points correspond to a set of five disjoint 24-cells. (Schoute's ten lines are totally singular.)
 - (iv) ± 72 pentagons of 5 vertices each. The $5+0$ points correspond to the 5 vertices of a pentagon. ($\langle v.w \rangle = \varphi$ or φ^{-1} , $72 = \frac{60 \cdot 24}{5 \cdot 4}$).
 - (v) 75 16-cells of 4 mutually orthogonal vertices each. The $4+1$ points correspond to the four vertices of a 16-cell, and the 24-cell that contains it. ($\langle v.w \rangle = 0$, $75 = \frac{60 \cdot 15}{4 \cdot 3}$).
 - (vi) ± 200 triangles of 3 vertices each. The $3+2$ points correspond to the triangle of vertices whose labels contain both duads $(i \ j)$ and $(i' \ j')$, plus the

two related 24-cells $(i\ j')$ and $(i'\ j)$. ($\langle v.w \rangle = 1$, $200 = \frac{60 \cdot 20}{3 \cdot 2}$).

References

- [1] Butler, M. et al, *The Unknown Subgroup of $\text{Aut}(E_8)$* , ArXiv 1709.05532, 2017
- [2] Conway, J.H.C. et al., **ATLAS of Finite Groups**, Clarendon Press, Oxford, 1985
- [3] Conway, J.H.C. & Sloane, N.J.A, **Sphere Packings, Lattices and Groups**, 3rd edition, Springer-Verlag, 1999
- [4] Coxeter, H.S.M., *Regular Compound Polytopes in More than Four Dimensions*, **J. Math Phys.** **12** (1933), 334-345
- [5] Coxeter, H.S.M., **Regular Polytopes**, 3rd edition, Dover, 1973
- [6] Du Val, P., **Homographies, Quaternions and Rotations**, Clarendon Press, Oxford, 1964
- [7] Dye, R.H., *The Simple Group $FH(8, 2)$ of Order $2^{12}3^55^{27}$ and the Associated Geometry of Triality*, **Proc LMS** **18** (1968), 521-562
- [8] Griess, R.L. & Lam, C.H., *A Moonshine Path for $5A$ and Associated Lattices of Ranks 8 and 16*, **Journal of Algebra** **331** (2011), 338-361; ArXiv 1006.390
- [9] Jacobson, N., **Basic Algebra I**, 2nd edition, Dover, 2005
- [10] Johnson, D., *H_4 over \mathbf{F}_5* , ArXiv
- [11] Littlewood, D.E., *The Groups of the Regular Solids in n Dimensions*, **Proc LMS** **s2-32** (1931), 10-20
- [12] Schläfli, L., *Réduction d'une intégrale multiple, qui comprend l'arc de cercle et l'aire du triangle sphérique comme cas particuliers*, **Journal de Mathématiques Pures et Appliquées (1)**, **20** (1855), 359-394
- [13] Schoute, P.H., **Mehrdimensionale Geometrie**, vol. 2, Leipzig 1905
- [14] Somerville, D.M.Y., **Introduction to the Geometry of n Dimensions**, Methuen & Co., London, 1929
- [15] Tits, J., *Quaternions over $\mathbb{Q}(\sqrt{5})$, Leech's Lattice and the Sporadic Group of Hall-Janko*, **Journal of Algebra** **63** (1980), 56-75
- [16] Wilson, R.A., *The Geometry of the Hall-Janko Group as a Quaternionic Reflection Group*, **Geometriae Dedicata** **20** (1986), 157-173

Contacts: tomme.denney32@gmail.com, babyshay12@icloud.com, jdejaneke@yahoo.com, tiannarobinson71401@gmail.com, majidhbutler3509@gmail.com, nisha.claiborne@gmail.com