

# HANKEL DETERMINANTS OF SECOND AND THIRD ORDER FOR THE CLASS $\mathcal{S}$ OF UNIVALENT FUNCTIONS

MILUTIN OBRADOVIĆ AND NIKOLA TUNESKI

**ABSTRACT.** In this paper we give the upper bounds of the Hankel determinants of the second and third order for the class  $\mathcal{S}$  of univalent functions in the unit disc.

Let  $\mathcal{A}$  be the class of functions  $f$  that are analytic in the open unit disc  $\mathbb{D} = \{z : |z| < 1\}$  of the form  $f(z) = z + a_2 z^2 + a_3 z^3 + \dots$  and let  $\mathcal{S}$  be the class of univalent functions in the unit disc  $\mathbb{D}$ . Let  $\mathcal{S}^*$  and  $\mathcal{K}$  denote the subclasses of  $\mathcal{A}$  which are starlike and convex in  $\mathbb{D}$ , respectively, and let  $\mathcal{U}$  denote the set of all  $f \in \mathcal{A}$  in  $\mathbb{D}$  satisfying the condition

$$\left| \left( \frac{z}{f(z)} \right)^2 f'(z) - 1 \right| < 1 \quad (z \in \mathbb{D}).$$

(see [4], [5], [6]).

The  $q$ th Hankel determinant for a function  $f$  from  $\mathcal{A}$  is defined for  $q \geq 1$ , and  $n \geq 1$  by

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \dots & a_{n+q} \\ \vdots & \vdots & & \vdots \\ a_{n+q-1} & a_{n+q} & \dots & a_{n+2q-2} \end{vmatrix}.$$

Thus, the second Hankel determinant is

$$(1) \quad H_2(2) = a_2 a_4 - a_3^2$$

and the third is

$$H_3(1) = \begin{vmatrix} 1 & a_2 & a_3 \\ a_2 & a_3 & a_4 \\ a_3 & a_4 & a_5 \end{vmatrix} = a_3(a_2 a_4 - a_3^2) - a_4(a_4 - a_2 a_3) + a_5(a_3 - a_2^2).$$

The concept of Hankel determinant finds its application in the theory of singularities (see [9]) and in the study of power series with integral coefficients.

For some subclasses of the class  $\mathcal{S}$  of univalent functions the sharp estimation of  $|H_2(2)|$  are known. For example, for the classes  $\mathcal{S}^*$  and  $\mathcal{U}$  we have that  $|H_2(2)| \leq 1$  (see [2, 7]), while  $|H_2(2)| \leq \frac{1}{8}$  for the class  $\mathcal{K}$  ([2]). Finding sharp estimates of the third order Hankel determinant turns out to be more complicated, so very few are known. An overview of results on the upper bound of  $|H_3(1)|$  can be found in [10],

---

2000 *Mathematics Subject Classification.* 30C45, 30C50, 30C55.

*Key words and phrases.* univalent, Hankel determinant of second order, Hankel determinant of third order.

while new non-sharp upper bounds for different classes and conjectures about the sharp ones are given in [8].

In this paper we give an upper bound of  $|H_2(2)|$  and  $|H_3(1)|$  for the class  $\mathcal{S}$ . Namely, we have

**Theorem 1.** *For the class  $\mathcal{S}$  we have*

$$|H_2(2)| \leq A, \quad \text{where} \quad 1 \leq A \leq \frac{11}{3} = 3,66\dots$$

and

$$|H_3(1)| \leq B, \quad \text{where} \quad \frac{4}{9} \leq B \leq \frac{32 + \sqrt{285}}{15} = 3.258796\dots$$

*Proof.* In the proof of this theorem we will use mainly the notations and results given in the book of N. A. Lebedev ([3]).

Let  $f \in \mathcal{S}$  and let

$$\log \frac{f(t) - f(z)}{t - z} = \sum_{p,q=0}^{\infty} \omega_{p,q} t^p z^q,$$

where  $\omega_{p,q}$  are called Grunsky's coefficients with property  $\omega_{p,q} = \omega_{q,p}$ . For those coefficients we have the next Grunsky's inequality ([1, 3]):

$$(2) \quad \sum_{q=1}^{\infty} q \left| \sum_{p=1}^{\infty} \omega_{p,q} x_p \right|^2 \leq \sum_{p=1}^{\infty} \frac{|x_p|^2}{p},$$

where  $x_p$  are arbitrary complex numbers such that last series converges.

Further, it is well-known that if

$$(3) \quad f(z) = z + a_2 z^2 + a_3 z^3 + \dots$$

belongs to  $\mathcal{S}$ , then also

$$f_2(z) = \sqrt{f(z^2)} = z + c_3 + c_5 z^5 + \dots$$

belongs to the class  $\mathcal{S}$ . Then for the function  $f_2$  we have the appropriate Grunsky's coefficients of the form  $\omega_{2p-1,2q-1}^{(2)}$  and the inequality (2) has the form

$$(4) \quad \sum_{q=1}^{\infty} (2q-1) \left| \sum_{p=1}^{\infty} \omega_{2p-1,2q-1}^{(2)} x_{2p-1} \right|^2 \leq \sum_{p=1}^{\infty} \frac{|x_{2p-1}|^2}{2p-1}.$$

As it has been shown in [3, p.57], if  $f$  is given by (3) then the coefficients  $a_2, a_3, a_4$  and  $a_5$  are expressed by Grunsky's coefficients  $\omega_{2p-1,2q-1}^{(2)}$  of the function  $f_2$  given by (3) in the following way (in the next text we omit upper index 2 in  $\omega_{2p-1,2q-1}^{(2)}$ ):

$$(5) \quad \begin{aligned} a_2 &= 2\omega_{11}, \\ a_3 &= 2\omega_{13} + 3\omega_{11}^2, \\ a_4 &= 2\omega_{33} + 8\omega_{11}\omega_{13} + \frac{10}{3}\omega_{11}^3, \\ a_5 &= 2\omega_{35} + 8\omega_{11}\omega_{33} + 5\omega_{15}^2 + 18\omega_{11}^2\omega_{13} + \frac{7}{3}\omega_{11}^4, \\ 0 &= 3\omega_{15} - 3\omega_{11}\omega_{13} + \omega_{11}^3 - 3\omega_{33}. \end{aligned}$$

Now, from (1) and (5) we have

$$\begin{aligned} H_2(2) &= 4\omega_{11}\omega_{33} + 4\omega_{11}^2\omega_{13} - 4\omega_{13}^2 - \frac{7}{3}\omega_{11}^4 \\ &= 4\omega_{11}\omega_{33} - \frac{4}{3}\omega_{11}^4 - (2\omega_{13} - \omega_{11}^2)^2, \end{aligned}$$

and from here

$$(6) \quad |H_2(2)| \leq 4|\omega_{11}||\omega_{33}| + \frac{4}{3}|\omega_{11}|^4 + |2\omega_{13} - \omega_{11}^2|^2.$$

Since for the class  $\mathcal{S}$  we have  $|a_3 - a_2^2| \leq 1$  (see [1]) and since from (5)

$$|2\omega_{13} - \omega_{11}^2| = |a_3 - a_2^2|,$$

then

$$(7) \quad |2\omega_{13} - \omega_{11}^2| \leq 1.$$

On the other hand, from (4) for  $x_{2p-1} = 0$ ,  $p = 3, 4, \dots$  we have

$$(8) \quad |\omega_{11}x_1 + \omega_{31}x_3|^2 + 3|\omega_{13}x_1 + \omega_{33}x_3|^2 \leq |x_1|^2 + \frac{|x_3|^2}{3}.$$

From (8) for  $x_1 = 1$ ,  $x_3 = 0$  and since  $\omega_{31} = \omega_{13}$ , we have

$$|\omega_{11}|^2 + 3|\omega_{13}|^2 \leq 1,$$

which implies

$$(9) \quad |\omega_{13}|^2 \leq \frac{1}{3}(1 - |\omega_{11}|^2).$$

Also, for  $x_1 = 0$ ,  $x_3 = 1$  we obtain

$$|\omega_{31}|^2 + 3|\omega_{33}|^2 \leq \frac{1}{3}$$

and so

$$(10) \quad |\omega_{33}| \leq \frac{1}{3}\sqrt{1 - 3|\omega_{31}|^2} \leq \frac{1}{3}.$$

Finally, from (6), (7), (9) and (10) we have

$$|H_2(2)| \leq \frac{4}{3}|\omega_{11}| + \frac{4}{3}|\omega_{11}|^4 + 1 \leq \frac{11}{3},$$

because from (5) we have that

$$|a_2| = |2\omega_{11}| \leq 2 \quad \Rightarrow \quad |\omega_{11}| \leq 1.$$

Since  $\mathcal{S}^*$  and  $\mathcal{U}$  are both subsets of  $\mathcal{S}$  with 1 as a sharp upper bound of  $|H_2(2)|$ , we have that on the class  $\mathcal{S}$ ,  $|H_2(2)| \geq 1$ .

As for Hankel determinant of the third order, by using (5), we can write

$$\begin{aligned} H_3(1) &= a_3(a_2a_4 - a_3^2) - a_4(a_4 - a_2a_3) + a_5(a_3 - a_2^2) \\ &= -8\omega_{13}^3 + 2\omega_{11}^4\omega_{13} + \frac{8}{3}\omega_{11}^3\omega_{33} - 4\omega_{33}^2 - \frac{4}{9}\omega_{11}^6 \\ &\quad + 4\omega_{13}\omega_{35} + 10\omega_{13}\omega_{15}^2 - 5\omega_{11}^2\omega_{15}^2 - 2\omega_{11}^2\omega_{35} \\ &= -2\omega_{13}(4\omega_{13}^2 - \omega_{11}^4) - \left(2\omega_{33} - \frac{2}{3}\omega_{11}^3\right)^2 + (2\omega_{35} + 5\omega_{15}^2)(2\omega_{13} - \omega_{11}^2), \end{aligned}$$

and from here

$$\begin{aligned}
|H_3(1)| &\leq \underbrace{2|\omega_{13}| |4\omega_{13}^2 - \omega_{11}^4|}_{B_1} + \underbrace{\left|2\omega_{33} - \frac{2}{3}\omega_{11}^3\right|^2}_{B_2} + \underbrace{|2\omega_{35} + 5\omega_{15}^2| |(2\omega_{13} - \omega_{11}^2)|}_{B_3} \\
&= B_1 + B_2 + B_3.
\end{aligned}$$

By using the relations (7) and (9), we obtain

$$\begin{aligned}
B_1 &= 2 \cdot |\omega_{13}| \cdot |2\omega_{13} - \omega_{11}^2| \cdot |2\omega_{13} + \omega_{11}^2| \\
&\leq 2 \cdot |\omega_{13}| \cdot |2\omega_{13} + \omega_{11}^2| \\
&\leq 2 \cdot |\omega_{13}| \cdot (2|\omega_{13}| + |\omega_{11}|^2) \\
&= 4|\omega_{13}|^2 + 2|\omega_{13}| \cdot |\omega_{11}|^2 \\
&\leq \frac{2}{3} \cdot \left[ 2(1 - |\omega_{11}|^2) + \sqrt{3} \cdot |\omega_{11}|^2 \cdot \sqrt{1 - |\omega_{11}|^2} \right] \\
&=: \frac{2}{3} \cdot \varphi(|\omega_{11}|^2),
\end{aligned}$$

where

$$\varphi(t) = 2(1 - t) + \sqrt{3}t\sqrt{1 - t}, \quad 0 \leq t \leq 1.$$

It is easily to show that the function  $\varphi$  decreases on  $(0, 1)$  and has maximum  $\varphi(0) = 2$ , which implies

$$(11) \quad B_1 \leq \frac{2}{3} \cdot \varphi(0) = \frac{4}{3}.$$

From the last equation in the relation (5), we have

$$2\omega_{33} - \frac{2}{3}\omega_{11}^3 = 2\omega_{15} - 2\omega_{11}\omega_{13},$$

and from here

$$(12) \quad \left| 2\omega_{33} - \frac{2}{3}\omega_{11}^3 \right| \leq 2|\omega_{15}| + 2|\omega_{11}||\omega_{13}|.$$

Similarly as in (8), we have

$$(13) \quad |\omega_{11}x_1 + \omega_{31}x_3|^2 + 3|\omega_{13}x_1 + \omega_{33}x_3|^2 + 5|\omega_{15}x_1 + \omega_{35}x_3|^2 \leq |x_1|^2 + \frac{|x_3|^2}{3}.$$

If we put  $x_1 = 1$  and  $x_3 = 0$ , then we get

$$|\omega_{11}|^2 + 3|\omega_{13}|^2 + 5|\omega_{15}|^2 \leq 1,$$

and so

$$(14) \quad |\omega_{15}| \leq \frac{1}{\sqrt{5}} \sqrt{1 - |\omega_{11}|^2 - 3|\omega_{13}|^2}.$$

From (12) and (14), we have

$$\begin{aligned}
\left| 2\omega_{33} - \frac{2}{3}\omega_{11}^3 \right| &\leq \frac{2}{\sqrt{5}} \left( \sqrt{1 - |\omega_{11}|^2 - 3|\omega_{13}|^2} + \sqrt{5}|\omega_{11}| \cdot |\omega_{13}| \right) \\
&=: \frac{2}{\sqrt{5}} \cdot \psi(|\omega_{11}|, |\omega_{13}|),
\end{aligned}$$

where

$$\psi(t, s) = \sqrt{1 - t^2 - 3s^2} + \sqrt{5}ts, \quad 0 \leq t \leq 1, \quad 0 \leq s \leq \frac{1}{\sqrt{3}}\sqrt{1 - t^2}.$$

It is an elementary fact to find that in cited domain  $\max \psi = 1$  attained for  $t = s = 0$ , which implies

$$(15) \quad B_2 = \left| 2\omega_{33} - \frac{2}{3}\omega_{11}^3 \right|^2 \leq \left( \frac{2}{\sqrt{5}} \right)^2 = \frac{4}{5}.$$

From relation (13) we also have

$$5|\omega_{15}x_1 + \omega_{35}x_3|^2 \leq |x_1|^2 + \frac{|x_3|^2}{3}.$$

If we put in the previous relation  $x_1 = 5\omega_{15}$ ,  $x_3 = 2$ , and then use (14) we receive

$$|2\omega_{35} + 5\omega_{15}^2|^2 \leq 5|\omega_{15}|^2 + \frac{4}{15} \leq 1 - |\omega_{11}|^2 - 3|\omega_{13}|^2 + \frac{4}{15} \leq \frac{19}{15},$$

which finally gives

$$(16) \quad B_3 = |2\omega_{35} + 5\omega_{15}^2| \cdot |2\omega_{13} - \omega_1^2| \leq \sqrt{\frac{19}{15}}$$

(in the last step we have used the relation (7)). By using relations (11), (15) and (16), we obtained

$$|H_3(1)| \leq B_1 + B_2 + B_3 \leq \frac{4}{3} + \frac{4}{5} + \sqrt{\frac{19}{15}} = \frac{32 + \sqrt{285}}{15}.$$

The function defined by  $\frac{zf'(z)}{f(z)} = \frac{1+z^3}{1-z^3}$  where  $a_2 = a_3 = a_5 = 0$ ,  $a_4 = \frac{2}{3}$  is starlike (thus univalent) and  $H_3(1) = -\frac{4}{9}$ . Therefore on the class  $\mathcal{S}$ ,

$$|H_3(1)| \geq \frac{4}{9}.$$

□

## REFERENCES

1. P.L. Duren, Univalent function, Springer-Verlag, New York, 1983.
2. A. Janteng, S.A. Halim and M. Darus, Hankel determinant for starlike and convex functions, Int. J. Math. Analysis, Vol. 1, No. 13, 2007, 619–225.
3. N.A. Lebedev, Area principle in the theory of univalent functions, Published by "Nauka", Moscow, 1975 (in Russian).
4. M. Obradović and S. Ponnusamy, New criteria and distortion theorems for univalent functions, *Complex Variables Theory Appl.* **44**(2001), 173–191. (Also Reports of the Department of Mathematics, Preprint 190, June 1998, University of Helsinki, Finland).
5. M. Obradović and S. Ponnusamy, On the class  $\mathcal{U}$ , Proc. 21st Annual Conference of the Jammu Math. Soc. and a National Seminar on Analysis and its Application, 11–26, 2011.
6. M. Obradović, S. Ponnusamy and K. J. Wirths, Geometric studies on the class  $\mathcal{U}(\lambda)$ , Bull. Malays. Math. Sci. Soc. **39** (2016), no. 3, 1259–1284.
7. M. Obradović and N. Tuneski, Some properties of the class  $\mathcal{U}$ , Ann. Univ. Mariae Curie-Skłodowska, accepted.
8. M. Obradović and N. Tuneski, New upper bounds of the third Hankel determinant for some classes of univalent functions, submitted. (<https://arxiv.org/abs/1911.10770>)
9. Dienes P., The Taylor Series: An Introduction to the Theory of Functions of a Complex Variable; NewYork-Dover: Mineola, NY, USA, 1957.

10. Shi L., Srivastava H.M., Arif M., Hussain S., Khan H., An Investigation of the Third Hankel Determinant Problem for Certain Subfamilies of Univalent Functions Involving the Exponential Function, *Symmetry* 2019, 11, 598.

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF BELGRADE,  
BULEVAR KRALJA ALEKSANDRA 73, 11000, BELGRADE, SERBIA

*E-mail address:* obrad@grf.bg.ac.rs

DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF MECHANICAL ENGINEERING,  
SS. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, KARPOŠ II B.B., 1000 SKOPJE, REPUBLIC OF  
NORTH MACEDONIA.

*E-mail address:* nikola.tuneski@mf.edu.mk