

EXTREMES, EXTREMAL INDEX ESTIMATION, RECORDS, MOMENT PROBLEM FOR THE PSEUDO-LINDLEY DISTRIBUTION AND APPLICATIONS

ABSTRACT. The pseudo-Lindley distribution which was introduced in Zeghdoudi and Nedjar (2016) is studied with regards to its upper tail. In that regard, and when the underlying distribution function follows the Pseudo-Lindley law, we investigate the behavior of its values, the asymptotic normality of the Hill estimator and the double-indexed generalized Hill statistic process (Ngom and Lo), the asymptotic normality of the records values and the moment problem.

[†] Gane Samb LO.

LERSTAD, Gaston Berger University, Saint-Louis, Sénégal (main affiliation).

LSTA, Pierre and Marie Curie University, Paris VI, France.

AUST - African University of Sciences and Technology, Abuja, Nigeria

gane-samb.lo@edu.ugb.sn, gslo@aust.edu.ng, ganesamblo@ganesamblo.net

Permanent address : 1178 Evanston Dr NW T3P 0J9, Calgary, Alberta, Canada.

[†] Dr. Modou NGOM

Work Affiliation : Ministry of High School (SENEGAL)

LERSTAD, Gaston Berger University, Saint-Louis, Sénégal.

ngomodoungom@gmail.com

[†] Dr. Moumouni Diallo

Université des Sciences Sociale et de Gestion de Bamako (USSGB)

Faculté des Sciences Économiques et de Gestion (FSEG)

Email: moudiallo1@gmail.com.

Keywords. Lindley's distribution; Pseudo-Lindley distribution; Extreme value theory; record values; Hill's estimator; asymptotic laws **AMS 2010 Mathematics Subject Classification:** 60G70; 62G20; 62H10; 62H15

1. INTRODUCTION

1. General facts.

The following probability distribution function (*pdf*), named as the Pseudo-Lindley *pdf*,

$$(1.1) \quad f(x) = f(x, \theta, \beta) = \frac{\theta(\beta - 1 + \theta x)e^{-\theta x}}{\beta} 1_{(x \geq 0)}$$

with parameters $\theta > 0$ and $\beta > 1$, has been introduced by [Zeghdoudi \(2016\)](#) as a generalization of the Lindley *pdf* :

$$(1.2) \quad \ell(x) = \frac{\theta^2(1+x)e^{-\theta x}}{1+\theta} 1_{(x \geq 0)}.$$

in the sense that for $\beta = 1 + \theta$, $f(\circ)$ is identical to $\ell(\circ)$.

Actually, f derives from ℓ by a mixture of a Lindley distributed random variable and an independent $\Gamma(2, \theta)$ random variables with mixture coefficients $r_1 = (\beta - 1)/\beta$ and $r_2 = 1/\beta$, where $1 < r_1$, $r_2 < 1$ and $r_1 + r_2 = 1$.

The cumulative distribution *cdf* function is given by

$$1 - F(x) = (\beta^{-1}(\beta + \theta x)e^{-\theta x}) 1_{(x \geq 0)}.$$

The Lindley original distribution is an important law that been used and still being used in Reliability, in Survival analysis and other important disciplines. Because of its original remarkable qualities, it kicked off a considerable number generalizations as pointed out by [Zeghdoudi \(2016\)](#). The current generalization (1.1) has been tested on real data and simulated and shows real interest in survival analysis, on the Guinean Ebola data for example ([Zeghdoudi \(2016\)](#)). The paper of focused on asymptotic tests of that law based on moments estimators. The interest that distribution demonstrated in real data modeling motivated us to give some asymptotic theories on it, in view of statistical tests. In this paper, we deal with the properties of the upper tail, the extreme value distribution and the record values. etc., which of them providing statistical tests.

Throughout the paper, X, X_1, X_2, \dots is a sequence independent real-valued random variables (*rv*), defined on the same probability space $(\Omega, \mathcal{A}, \mathbb{P})$, with common cumulative distribution function F , with the first asymptotic moment function and the generalized inverse function defined by

$$R(x, F) = \frac{1}{1 - F(x)} \int_x^{+\infty} (1 - F(y)) dy, \quad x \in]0, +\infty[$$

and

$$F^{-1}(u) = \inf\{x \in \mathbb{R}, F(x) \geq u\} \text{ for } u \in]0, 1[\text{ and } F^{-1}(0) = F^{-1}(0+).$$

For each $n \geq 1$, we denote the ordered statistics of the sample X_1, \dots, X_n by

$$X_{1,n} \leq \dots \leq X_{n,n}.$$

Usually, in extreme value theory, we focus on upper extreme and the hypothesis $X > 0$ and the log-transform $Y = \log X$ is instrumental in all major results in that field. We denote $G(x) = F(e^x)$, $x \in \mathbb{R}_+$. The Renyi representation is also of common use in the following form. The sequence is replaced as follows

$$(1.3) \quad \{\{X_{1,n} \leq \dots \leq X_{n,n}\}, n \geq 1\} =_d \{\{F^{-1}(1 - U_{n-j+1,n}), 1 \leq j \leq n\}, n \geq 1\},$$

where $=_d$ stands for the equality in distribution. Finally, the following Malmquist representation (see [Shorack and Wellner \(1995\)](#), also [Lo et al. \(2016\)](#), page ...) is also used : for each $n \geq 1$, there exist a finite sequence of standard independent exponential random variables $E_{1,n}, \dots, E_{n,n}$ such that

$$(1.4) \quad \left\{ \left(\frac{U_{i+1,n}}{U_{i,n}} \right)^i, 1 \leq i \leq n \right\} =_d \{E_{i,n}, 1 \leq i \leq n\}.$$

2. EXTREMES

We can directly see that F is the Gumbel distribution G_0 by three different arguments. First, by using the Von Mises' argument (see [de Haan \(1970\)](#) or [Lo et al. \(2018\)](#), Proposition 24, page 184)

$$(2.1) \quad \lim_{x \rightarrow 0} \frac{f'(x)(1 - F(x))}{f^2(x)} = -1.$$

A second argument comes from that $Y = \exp(X)$ has the distribution $G(x) = F(\log x) = \beta^{-1}(\beta + \theta \log x)x^{-\theta x}$. Since

$$(2.2) \quad \lim_{x \rightarrow 0} \frac{1 - G(\lambda x)}{1 - G(x)} = \lambda^{-\theta},$$

$G \in D(G_{1/\theta})$ and since $F(x) = G(e^x)$ for $x \geq 1$, by Theorem (in Lo), $F \in D(G_0)$.

A third argument is relation to the development of the quantile function. In the appendix (page 16), we give a number of expansions of that quantile that could be used for different purposes. For example we have (see page 19),

$$(2.3) \quad F^{-1}(1-u) = \theta^{-1}(\log(1/u) - \log \log(1/u)) + \theta^{-1}K(u)$$

with $K(u) = O(\log 1/u)^{-2}$. By using it we get

$$\frac{F^{-1}(1-\lambda u) - F^{-1}(1-u)}{(1/\theta)} \rightarrow -\log \lambda \text{ as } u \rightarrow 0.$$

By the π -variation criteria of [de Haan \(1970\)](#) (See [Lo et al. \(2016\)](#), Proposition 11, page 88), we have $F \in D(G_0)$ and $R(x, F) \rightarrow \gamma = 1/\theta$ as $x \rightarrow +\infty$. Formula (2.3) is actually a second-order condition for the quantile function (see [de Haan \(1970\)](#)). We apply it right to get a rate of convergence of the maximum observations. Put $\gamma = 1/\theta$.

2. Expansion of the maximum values.

By the Renyi representation and by denoting $Z_n = -\log(nU_{1,n})$, we have that $\log(1 + Z_n/(\log n)) \rightarrow_{\mathbb{P}} 0$ and since $\log U_{1,n} = O_{\mathbb{P}}(\log n)^{-1}$

$$\begin{aligned} X_{n,n} - F^{-1}(1 - 1/n) &= \gamma Z_n + \gamma \log(1 + Z_n/(\log n)) \\ &\quad + O((\log n)^{-2}) + O((\log U_{1,n})^{-2}). \end{aligned}$$

and hence

$$(2.4) \quad \frac{X_{n,n} - F^{-1}(1 - 1/n)}{\gamma} = Z_n + O_{\mathbb{P}}(\log n)^{-1} = \Gamma + O_{\mathbb{P}}(1)$$

It is easy to see that Z_n converge to Gumbel law Λ with *cdf*

$$G_0(x) = \exp(-\exp(-x)), \quad x \in \mathbb{R}$$

So we have that $X_{n,n}$ converges to a Λ law. But we obtain the random rate of convergence $Z_n/\log n$ since

$$\frac{\log Z_n}{\log n} \left(\frac{X_{n,n} - F^{-1}(1 - 1/n)}{\gamma} - Z_n \right) = 1$$

As well for $k = k(n) \rightarrow +\infty$ such that $k(n)/n \rightarrow 0$, and by taking $T_n = \log(nU_{k,n}/k)$ and $q_n = n/k(n)$ which goes to $+\infty$, we have

$$(2.5) \quad \frac{X_{n-k,n} - F^{-1}(1 - k/n)}{\gamma} = T_n + \log(1 + T_n/\log q_n) + O_{\mathbb{P}}((\log q_n)^{-2}).$$

3. Estimating the extreme value index $\gamma = 1/\theta$.

The [Hill \(1975\)](#)'s estimator

$$(2.6) \quad H_n = \frac{1}{k(n)} \sum_{j=1}^{k(n)} j (X_{n-j+1,n} - X_{n-j,n}),$$

is the most celebrated estimator the extreme value index $\gamma = 1/\theta$ of $Z = \exp(X)$. Among a significant number of generalizations, the [Ngom and Lo \(2016\)](#)'s generalization, called the Double-indexed function Hill estimator, is one the sharpest one. It is defined as

$$H_n(f, s) = \left(\sum_{j=1}^{k(n)} f(j) (X_{n-j+1,n} - X_{n-j,n})^s / (a_n(f, s)) \right)^{1/s},$$

where $f : \mathbb{N} \setminus \{0\} \rightarrow \mathbb{R}_+ \setminus \{0\}$ is a measurable mapping and $s > 0$. Let us define

$$a_n(f, s) = \Gamma(s+1) \sum_{j=1}^{k(n)} f(j) j^{-s}, \quad C^2(s) = \Gamma(2s+1) - \Gamma(s+1)^2, \quad s_n^2(f, s) = C^2(s) \sum_{j=1}^{k(n)} f(j)^2 j^{-2s},$$

and

$$B_n(f, s) = \max\{f(j)j^{-s}/s_n(f, s), \ 1 \leq j \leq k(n)\}.$$

We simply notice that the classical Hill's estimator is $H_n(I_d, 1)$ where I_d is the identity function on $\mathbb{N} \setminus \{0\}$. Let us give asymptotic normality for

Double-indexed function Hill estimator.

(a) Extreme Limit Theorem.

We begin with the simple Hill's estimator.

Theorem 1. *For $]0, n] \ni k(n) \rightarrow +\infty$ such that*

$$(K1) \quad k(n)^{3/4} / \log n \rightarrow 0.$$

we have, as $n \rightarrow +\infty$,

$$(2.7) \quad \sqrt{k(n)} (H_n - \gamma) \rightsquigarrow \mathcal{N}(0, \gamma^2).$$

We want to establish the random rate of convergence associated with the convergence 2.7 in the part (a) of the following corollary. In the part (b), we want to share that we need any other condition on top of $k(n)/n \rightarrow 0$ to have the central limit theorem if F^{-1} is reduced to

$$(2.8) \quad F_*^{-1}(1 - u) = \gamma \log u - C(\gamma) \log \log(1/u), \quad u \in]0, 1[, \quad C(\gamma) \geq 1.$$

Corollary 1. *We have the following results.*

(a) Here again F is the cdf of the Pseudo-Lindley distribution with parameters $\theta > 0$ and $\beta > 0$ and the notation above. Let $k(n)/\log n \rightarrow 0$.

Let $W(1)$ is a standard Gaussian random variable. Then we have

$$\frac{\log n}{\gamma \sqrt{k(n)}} \left(\sqrt{k(n)} (H_n - \gamma) - \gamma W(1) \right) \rightarrow_{\mathbb{P}} 1,$$

(b) If F^{-1} were reduced as in Formula (2), we have the asymptotic normality

$$\sqrt{k(n)} (H_n - \gamma) \rightarrow \mathcal{N}(0, \gamma^2)$$

whenever $k(n)/n \rightarrow 0$ and

$$\log n \left(\sqrt{k(n)} (H_n - \gamma) - \gamma W(1) \right) = O_{\mathbb{P}}(1). \quad \diamond$$

Proof of Theorem 1. By Malmquist representations (See [Shorack and Wellner \(1995\)](#) or [Lo et al. \(2016\)](#), Proposition 32, page 135), by Formula (4.18), we have for any $1 \leq j \leq k$,

$$(2.9) \quad \begin{aligned} X_{n-j+1,n} - X_{n-j,n} &= F^{-1}(1 - U_{j,n}) - F^{-1}(1 - U_{j+1,n}) \\ &= \gamma j^{-1} E_{j,n} - \gamma \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} + O_{\mathbb{P}}((\log n)^{-2}) \end{aligned}$$

and next

$$j(X_{n-j+1,n} - X_{n-j,n}) = \gamma E_{j,n} - \gamma j \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} + O_{\mathbb{P}}(k(\log n)^{-2}).$$

So for $Z_n = \log n U_{1,n}$ (which converges in law to Λ) and

$$(2.10) \quad \left| \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} \right| \leq \frac{j^{-1} E_{j,n}}{|\log n - Z_n|}.$$

Hence

$$\frac{1}{k(n)} \left| \sum_{j=1}^{k(n)} j \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} \right| \leq \frac{S_{k(n)}^*}{k} O_{\mathbb{P}}((\log n)^{-1}).$$

where $S_{k(n)}^* = E_{j,n} + \dots + E_{k,n}$. We finally get

$$\sqrt{k(n)}(H_n - \gamma) = \gamma \frac{S_{k(n)}^* - k}{\sqrt{k(n)}} + O_{\mathbb{P}}\left(\frac{1}{\log n}, \frac{k^{3/2}}{(\log n)^2}\right)$$

We conclude that, whenever (K1) holds, we have

$$\sqrt{k(n)}(H_n - \gamma) = \gamma \frac{S_{k(n)}^* - k}{\sqrt{n}} + o_{\mathbb{P}}(1). \quad \square$$

Proof of the Corollary 1. The proof of Part (b) is the conclusion of the proof of Theorem 1 up to the formula (2.10). If holds, further steps are dismissed. And we need only $k(n)/n \rightarrow 0$ to conclude. Let us set

$$Z_n^* = \frac{1}{\sqrt{k(n)}} \sum_{j=1}^{k(n)} j \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)}, \quad n \geq 1.$$

For the first part, we already knew that $Z_n^* = O_{\mathbb{P}}(1/\log n)$. We denoted by $W(1)$ a standard Gaussian random variable. By the classical Kómlós-Màjor-Tusnàdy (KMT) approximation, we have

$$\left| \frac{S_k^*(n) - k(n)}{\sqrt{k(n)}} - \gamma W(1) \right| = O_{\mathbb{P}} \left(\frac{\log k(n)}{\sqrt{k(n)}} \right).$$

Straightforward expansions using the different rates of convergence lead to

$$\frac{\sqrt{k(n)}(H_n - \gamma) - \gamma W(1)}{\gamma Z_n^*} \rightarrow_{\mathbb{P}} 1,$$

whenever $k(n)/n \rightarrow 0$. Now we apply Proposition in **?**, page 22. Since the function $\log(1/u)$ is slowly varying and that $U_{1,n}/U_{k+1,n}$ and $U_{k+1,n}/U_{1,n}$ are both asymptotically bounded in probability, we have

$$t_n = \sup_{1 \leq j \leq k(n)} \sup_{s \in [U_{j,n}, U_{j+1,n}]} \left| \frac{\log(1/s)}{\log n} - 1 \right| \rightarrow_{\mathbb{P}} 0.$$

It comes that

$$Z_n^* = \frac{\sqrt{k(n)}}{\log n} (k_{(n)}^{-1} S_{k(n)}^*) (1 + O(t_n)) = \frac{\sqrt{k(n)}}{\log n} (1 + o(1)),$$

which gives the desired result. ■

We have the following convergence of the Double-indexed functional Hill statistics.

Theorem 2. *We have the following two results.*

(a) *If the following conditions hold, as $n \rightarrow +\infty$*

$$s_n(f, 1)/(s_n(f, s) \log n) \rightarrow 0 \text{ and } B_n(f, s) \rightarrow 0,$$

then

$$\frac{T_n(f, s) - \gamma^s a_n(f, s)}{s_n(f, s)} \rightsquigarrow \mathcal{N}(0, \gamma^{2s}).$$

(b) Furthermore, if $a_n(f, s)/a_n(f, s) \rightarrow +\infty$, then

$$\frac{s_n(f, s)}{a_n(f, s)} \left(\left(\frac{T_n(f, s)}{a_n(f, s)} \right)^{1/s} - \gamma \right) \rightsquigarrow \mathcal{N}(0, s^{-2} \gamma^2). \blacksquare.$$

Proof. Let us exploit the proof of Theorem 1. we have for $j \in \{1, \dots, k(n)\}$, $s \geq 1$,

$$\begin{aligned} A_{i,n} &= f(j) (X_{n-j+1,n} - X_{n-j,n})^s \\ &= f(j) \left(\gamma j^{-1} E_{j,n} - \gamma \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} + O_{\mathbb{P}}(F_{k(n)}(\log n)^{-2}) \right)^s \\ &=: f(j) (\gamma j^{-1} E_{j,n} - R_{j,n} + C_{j,n})^s, \end{aligned}$$

with

$$C_{j,n} = O_{\mathbb{P}}((\log n)^{-2}) \text{ (uniformly in } j), \left| \gamma \int_{U_{j,n}}^{U_{j+1,n}} \frac{du}{u \log(1/u)} \right| \leq \frac{\gamma j^{-1} E_{j,n} b(n)}{|\log n - Z_n|}.$$

We get, by the mean value theorem, $j \in \{1, \dots, k(n)\}$, $s \geq 1$,

$$\begin{aligned} &A_{i,n} - \gamma^s f(j) j^{-s} E_{j,n}^s \\ &\leq s f(j) |R_{j,n} + C_{j,n}| (\gamma j^{-1} E_{j,n} + |R_{j,n}| + |C_{j,n}|)^{s-1} \\ &\leq \left(\frac{s \gamma f(j) j^{-1} E_{j,n}}{|\log n - Z_n|} \right) (\gamma j^{-1} E_{j,n} + |R_{j,n}| + |C_{j,n}|)^{s-1}. \end{aligned}$$

In the lines below, we will bound the term with the power $s - 1$. But if $s = 1$, there will be nothing to bound. So formulas regarding that term are dismissed for $s = 1$ and are used only for $s > 2$. For $s \geq 1$, we will use the C_{s-1} inequality (for $s \leq 2$, with $|a + b|^{s-1} \leq 2^{s-2} |a|^{s-1} + |b|^{s-1}$, $C_{s-1} = 2^{s-2}$). For $0 < r < 1$, it can be easily checked that, for $u > 0$ fixed, the function $g(v) = (u + v)^r - u^r - v^r$ of $v \geq 0$ takes the value $g(v) = 0$ and has a non-positive derivative function, so that $g(v) \leq g(0) = 0$ for any $v \geq 0$, which is

equivalent to $(u+v)^r \leq u^r + v^r$. We finally have that $|a+b|^{s-1} \leq D_s |a|^{s-1} + |b|^{s-1}$ with $D_s = 1$ for $1 < s < 2$ and $D_s = C_{s-1}$ for $s \geq 2$. Applying that inequality leads, $j \in \{1, \dots, k(n)\}$, $s \geq 1$, to

$$\begin{aligned} & A_{i,n} - \gamma^s f(j) j^{-s} E_{j,n}^s \\ & \leq \left(\frac{s \gamma f(j) j^{-1} E_{j,n}}{|\log n - Z_n|} \right) \left(D_s \gamma^{s-1} j^{s-1} E_{j,n}^{s-1} + \frac{D_s^2 \gamma^{s-1} j^{s-1} E_{j,n}^{s-1}}{(|\log n - X_n|^{s-1})} + O_{\mathbb{P}} \left(\frac{D_s^2}{(\log n)^{2(s-1)}} \right) \right). \end{aligned} \quad (A)$$

Let us denote

$$S_n(f, s) = \sum_{j=1}^{k(n)} f(j) j^{-s} E_{j,n}^s$$

and

$$T_n(f, s) = \sum_{j=1}^{k(n)} f(j) (X_{n-j+1,n} - X_{n-j,n})^s,$$

By combining the results above, we arrive at

$$\begin{aligned} & \left| T_n(f, s) - \gamma^s S_n(f, s) \right| \\ & \leq \left(\frac{s \gamma S_n(f, 1)}{|\log n - Z_n|} \right) \left(D_s \gamma^{s-1} S_n(Id, s-1) + \frac{D_s^2 \gamma^{s-1} S_n(Id, s-1)}{(|\log n - Z_n|^{s-1})} + O_{\mathbb{P}} \left(\frac{D_s^2}{(\log n)^{2(s-1)}} \right) \right). \end{aligned} \quad (B)$$

Let us study $S_n(f, s)$. As a sequence of partial sums of real-value independent random variables indexed by $j \in \{1, \dots, k(n)\}$ with first and second moments

$$\Gamma(s+1) f(j) j^{-s} \text{ and } (\Gamma(2s+1) - \Gamma(s+1)^2) f(j)^2 j^{-2s},$$

the asymptotic normality is given by the the theorem of Levy-Feller-Linderberg (See Theorem 20 in [Lo et al. \(2018\)](#) we apply to the centered rrv 's $\xi_j = f(j) j^{-s} (E_{j,n}^s - \Gamma(s+1))$, after remarking that

$$\left\{ \frac{\text{Var}(\xi_j)}{\sum_{j=1}^{k(n)} \text{Var}(\xi_j)}, 1 \leq j \leq k(n) \right\} = C(s) B_n(f, s).$$

So, as $n \rightarrow +\infty$,

$$\left(\frac{1}{s_n(f, s)} \sum_{j=1}^{k(n)} (f(j)j^{-s}(E_{j,n}^s - \Gamma(s+1))) \rightsquigarrow \mathcal{N}(0, 1) \right) \text{ and } B_n(f, s) \rightarrow 0$$

and the Lynderberg condition holds, that is, for any $\varepsilon > 0$,

$$g(n, \varepsilon) = \frac{1}{s_n(f, s)} \sum_{j=1}^{k(n)} \int_{(|\xi_j| > \varepsilon s_n(f, s))} \xi_j^2 d\mathbb{P} \rightarrow 0.$$

But, for $K^2(s) = \Gamma(4s+1) - 4\Gamma(3s+1)\Gamma(s+1) + \Gamma(2s+1)\Gamma(s+1)2 - 3\Gamma(3s+1)^4$,

$$\mathbb{E}\xi^4 = K(s)f(s)^4j^{-4s}$$

and, by the Cauchy-Schwarz inequality

$$\begin{aligned} & \int_{(|\xi_j| > \varepsilon s_n(f, s))} \xi_j^2 d\mathbb{P} \\ & \leq \left(\int \xi_j^4 d\mathbb{P} \right)^{1/2} \left(\int 1_{(|\xi_j| > \varepsilon s_n(f, s))} d\mathbb{P} \right)^{1/2} \\ & = Kf(j)^2j^{2s} \left(\int 1_{(|\xi_j| > \varepsilon s_n(f, s))} d\mathbb{P} \right)^{1/2} \\ & = Kf(j)^2j^{2s} \mathbb{P}(|\xi_j| > \varepsilon s_n(f, s))^{1/2} \\ & \leq Kf(j)^2j^{2s} \left(\frac{K(s)^2f(j)^4j^{-4s}}{\varepsilon^4 s_n^4(f, s)} \right)^{1/2} \\ & = K(s)^2 (f(j)^2j^{2s})^2 (s_n^{-2}(f, s))^2 \\ & = \frac{C(s)}{K(s)} B_n(f, s) \frac{\mathbb{V}ar(\xi_j)}{s^2(f, s)} \end{aligned}$$

So

$$g(n, \varepsilon) = \left(\frac{K(s)}{C(s)} \right)^2 B_n(f, s) \rightarrow 0.$$

Our hypothesis $B_n(f, s) \rightarrow 0$ makes the Lynderberg hold and the central limit theorem holds for $S_n(f, s)$, that is

$$\frac{S_n(f, s) - \gamma^s a_n(f, s)}{s_n(f, s)} \rightsquigarrow \mathcal{N}(0, 1)$$

Now, let us return to the approximation (B) at page 10. We have that for $s = 1$, the expression denoted as C_n between the pair of big parentheses should be equal to one as explained before. If $s > 1$, we have $\sigma^2(s) = \sum_{j \geq 1} j^{-2(s-1)} < +\infty$, we apply a theorem of Kolmogorov (see Lo et al. (2018), Proposition 25, page 233), $S_n(Id, s-1)$ weakly converges to the random variable $W(s)$ with variance $\sigma^2(s)$. Hence $C_n = O_{\mathbb{P}}(1)$. We arrive at

$$(2.11) \quad \left| \frac{T_n(f, s) - a_n(f, s)}{s_n(f, s)} - \frac{\gamma^s (S_n(f, s) - a_n(f, s))}{s_n(f, s)} \right| \leq O_{\mathbb{P}} \left(\frac{S_n(f, 1)}{s_n(f, s) \log n} \right).$$

The later bound goes to zero in probability if and only if $s_n(f, 1)/(s_n(f, s) \log n) \rightarrow 0$. Now, we have

$$\frac{s_n(f, s)}{a_n(f, s)} \left(\frac{T_n(f, s)}{a_n(f, s)} - \gamma^s \right) = Z_n + o_{\mathbb{P}}(1).$$

If $s_n(f, s)/a_n(f, s) \rightarrow +\infty$, we can use the δ -method applied to $g(t) = t^{1/s}$ to get

$$\frac{a_n(f, s)}{s_n(f, s)} \left(\left(\frac{T_n(f, s)}{a_n(f, s)} \right)^{1/s} - \gamma \right) \rightsquigarrow \mathcal{N}(0, s^{-2} \gamma^2). \blacksquare.$$

Remark. In Ngom and Lo (2016), we gave a direct proof of the asymptotic normality of $S_n(f, s)$ by using the two hypotheses $B_n(f, s) \rightarrow 0$ and $s_n(f, s) \rightarrow +\infty$. Here, it seems that we only used the first one. But that one could not hold if $S_n(f, s)$ contains a sub-sequence converging to a finite and positive number. That remark should be recalled in interpreting the results in Ngom and Lo (2016).

3. UPPER RECORDS VALUES

The main result is :

Theorem 3. *If, for each $n \geq 1$, $X^{(n)}$ stands for n -th record value, we have as $n \rightarrow +\infty$,*

$$\frac{X^{(n)} - \gamma n}{\gamma \sqrt{n}} \rightsquigarrow \mathcal{N}(0, 1).$$

Remark. We refer the reader to [Lo and Ahsanullah \(2019\)](#) for a simple introduction to records theory.

Proof. We already noticed that $Z = \exp(X)$ is the extremal domain of attraction of $G_\gamma(x) = \exp(-(1 + \gamma x))$, for $\gamma x > -1$. From Part (b) of Theorem 1 in [Lo and Ahsanullah \(2019\)](#), the n -th record $Z^{(n)} = \exp(X^{(n)})$ have the representation

$$(3.1) \quad \left(\frac{\exp(X^{(n)})}{H^{-1}(1 - e^{-n})} \right)^{1/\sqrt{n}} = \exp(\gamma S_n^*) + o_{\mathbb{P}}(1)$$

where S_n^* has the same law as $\gamma^{-1}(T_n - n)/\sqrt{n}$ with T_n denoting a γ law with parameters n and 1. Since $H^{-1}(1 - u) = \exp(F^{-1}(1 - u))$, we have

$$(3.2) \quad \frac{X^{(n)} - F^{-1}(1 - e^{-n})}{\gamma \sqrt{n}} = S_n^* + o_{\mathbb{P}}(1)$$

By the central limit theorem, it comes that

$$(3.3) \quad \frac{X^{(n)} - F^{-1}(1 - e^{-n})}{\gamma \sqrt{n}} = \mathcal{N}(0, 1) + o_{\mathbb{P}}(1).$$

By using Formula (2.3), we get

$$(3.4) \quad \frac{X^{(n)} - \gamma n}{\gamma \sqrt{n}} = S_n^* + o_{\mathbb{P}}(1)$$

$$(3.5) \quad \frac{X^{(n)} - \gamma n}{\gamma \sqrt{n}} = \mathcal{N}(0, 1) + o_{\mathbb{P}}(1).$$

The proof is over. \square

4. THE MOMENT PROBLEM

Typically, the moment problem on \mathbb{R} (see [Shohat and Tamarkin \(1943\)](#)) is the following. Given a sequences real numbers $(m_n)_{n \geq 1}$, can we find a distribution (not necessarily a *cdf*) F on \mathbb{R} as the unique solution of the moments equations.

$$\forall n \geq 1, m_n = \int x^n dF(x).$$

This is a nice but difficult mathematical question treated in [Shohat and Tamarkin \(1943\)](#). But in the context of probability theory on \mathbb{R} , we may have a fixed *cdf* F of random variable X having moments

$$\forall n \geq 1, \mathbb{E}X^n = m_n \text{ finite.}$$

The moment problem becomes : Is the sequence of moments $(m_n)_{n \geq 1}$ characterize the probability law of X . In that regard, we have

Theorem 4. *The moments of the pseudo-Lindely probability law are the following*

$$\forall n \geq 1, m_n = \frac{n!(\beta + n)}{\theta^n \beta}.$$

Any real-valued random variable have the moments $(m_n)_{n \geq 1}$ follows the pseudo-Lindley law.

Proof. At the place of a simple proof, we proceed to slight round-up of the moment problem and explain how to find a simple criteria based on Analysis. A possible tool is the characteristic function which characterize its associated probability law. We have the following expansion of any characteristic function of X (see [loeve \(1997\)](#) or [Lo et al. \(2018\)](#), Lemma 5, page 255), we have

$$(4.1) \quad \mathbb{E}e^{iuX} = 1 + \sum_{k=1}^n \frac{(iu)^k m_k}{k!} + \theta 2^{1-\delta} \mu^{n+\delta} \frac{|u|^{n+\delta}}{(n+1)!}.$$

By usual analysis tools, the series in Formula (4.1) converges in the $] -R, R[$ where R is found according the Cauchy rule

$$\limsup_{n \rightarrow +\infty} (m_n)^{1/n} = R > 0.$$

The conclusion is that two random variables have the same moments of all orders have characteristic functions coinciding on $] - R, R[$. Finally, (see [loeve \(1997\)](#), page 225, Part B.; see also [Billingsley \(1995\)](#)) two characteristic function coinciding on an interval $] - R, R[$ coincide everywhere and thus, are associated to the same probability law.

Let us apply to the pseudo-Lindley law. In [Zeghdoudi \(2016\)](#), the moments are given by

$$\forall n \geq 1, m_n = \frac{n!(\beta + n)}{\theta^n \beta}.$$

Straightforward computation based on the Stirling formula leads to $R = 1/\theta$. This is enough to prove the claim of the theorem.

Appendix . Let $R = \beta/\theta$. In the computations below, $u \in (0, 1)$ and $x \geq 0$ are linked by $u = 1 - F(x)$. So $u \rightarrow 0$ if and only if $x \rightarrow +\infty$. Also, below, functions of x are functions of u actually. We denote $A(u) = \log(1 + R/x)$. We have $A(u) \rightarrow 0$ as $u \rightarrow 0$. By writing

$$\log(\beta + \theta x) = \log(\beta + \theta x) - \log \theta x + \log \theta x = \log \theta x + A(u),$$

we see that $u = 1 - F(x)$ gives

$$(4.2) \quad \theta x = \log(1/u) + \log R + \log x + A(u).$$

So, we have

$$(4.3) \quad F^{-1}(1 - u) = \theta^{-1} \log(1/u)(1 + o(1)).$$

and

$$(4.4) \quad \log x = \log \log(1/u)(1 + o(1)).$$

Now, we wish to develop that asymptotic equivalence with rates of convergence. Let $B(u) = \log R + \log x + A(u)$. From Formula 4.2, we have

$$(4.5) \quad \frac{x}{\theta^{-1} \log(1/u)} - 1 = \frac{B(u)}{\log(1/u)}.$$

By Formula (4), we notice that

$$(4.6) \quad B(u) = \log R + \log x + (R/x) - (R/x)^2/2 + O(\log(1/u)^{-3}) = O(\log x) = (\log \log u)(1 + o(1)),$$

and hence, for $D(u) = \log R + A(u)$,

$$(4.7) \quad \frac{\log(1/u)}{\log x} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) = 1 + \frac{D(u)}{\log x}.$$

Also

$$\frac{D(u)}{\log x} = \frac{\log R + (R/x) - (R/x)^2/2 + O(x^{-3})}{\log x}$$

Next, we have

$$(4.8) \quad \begin{aligned} & \frac{\log x}{-\log R} \left(\frac{\log(1/u)}{\log x} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) - 1 \right) \\ &= 1 + \frac{R}{x \log R} - \frac{R^2}{2x^2 \log R} + O(x^{-3}) \end{aligned}$$

and finally

$$(4.9) \quad \begin{aligned} & \frac{x \log R}{R} \left(\frac{\log x}{\log R} \left(\frac{\log(1/u)}{\log x} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) - 1 \right) - 1 \right) \\ &= 1 - \frac{R}{2x} + O(x^{-2}). \end{aligned}$$

Now we want to do the same for $\log x$. From Formula , we have

$$(4.10) \quad \log(\theta x) = \log \log(1/u) + \log(1 + B(u)/\log(1/u))$$

from which we get

$$(4.11) \quad \log x - \log \log(1/u) = -\log \theta + (B(u)/\log(1/u)) + O((B(u)/\log(1/u))^2).$$

From Formula (4.7), we have

$$\begin{aligned} & \frac{\log(1/u)}{\log x} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) - \frac{\log(1/u)}{\log \log 1/u} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) \\ &= \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) \frac{-(\log(1/u))(\log x - \log \log 1/u)}{(\log x)(\log \log 1/u)} \\ &= (1 + D(u)/\log x) \left(\frac{1}{(\log x)(\log \log 1/u)} (-\log \theta + (B(u)/\log(1/u)) + O(B(u)/\log(1/u)^2)) \right) \\ &= O((\log \log 1/u)^2) \end{aligned}$$

Formula (4.7) becomes

$$(4.12) \quad \frac{\log(1/u)}{\log \log 1/u} \left(\frac{x}{\theta^{-1} \log(1/u)} - 1 \right) = 1 + \frac{D(u)}{\log x} + O((\log \log 1/u)^2).$$

That formula will be used with Formula 4.11 and

$$(4.13) \quad \begin{aligned} \frac{B(u)}{\log 1/u} &= \frac{\log R}{\log 1/u} + \frac{\log \log 1/u}{\log 1/u} (1 + o(1)) \\ &+ \frac{(R/x) - (R/x)^2/2}{\log 1/u} + O((\log 1/u)^{-4}). \end{aligned}$$

From 4.2, and from the following formula we can check by using differentiation methods to establish monotonicity

$$x - x^2/2 \leq \log(1+x) \leq x$$

we have

$$(4.14) \quad (R/x) - R^2/(2x^2) + \log R + \log x \leq \theta x - \log(1/u) \leq (R/x) + \log R + \log x.$$

But we also have

$$x = \log(1/u) \left(1 + \frac{\log \beta^{-1} + \log x + A(u)}{\log(1/u)} \right)$$

which implies

$$\log x = \log \log(1/u) + \log \left(1 + \frac{\log \beta^{-1} + \log x + A(u)}{\log(1/u)} \right)$$

By putting

$$H(u) = \frac{\log \beta^{-1} + \log x + A(u)}{\log(1/u)},$$

we finally get

$$(4.15) \quad H(u) - H(u)^2/2 \leq \log x - \log \log(1/u) \leq H(u).$$

By combining Formulas (4.14) and (4.15), we get

$$(4.16) \quad |\theta x - \log(1/u) - \log \log(1/u)| \leq \frac{1}{2} \left(\frac{R^2}{x^2} + H(u)^2 \right).$$

Since (R/x^2) and $H(u)^2$ are both $O(\log 1/u)^{-2}$, we have

$$(4.17) \quad F^{-1}(1-u) = \theta^{-1}(\log(1/u) - \log \log(1/u)) + O(\log 1/u)^{-2}.$$

But since the derivative $\log \log(1/u)$ is $(-u \log(1/u))^{-1}$, we have for $d = -\log \log 2$,

$$\forall u \in]0, 1[, \log \log(1/u) - = \int_u^{1/2} \frac{1}{u \log(1/u)} du,$$

and finally

$$(4.18) \quad F^{-1}(1-u) = d + \theta^{-1}(\log(1/u) - \int_u^{1/2} \frac{1}{u \log(1/u)} du + O((\log 1/u)^{-2})).$$

REFERENCES

- Patrick Billingsley (1995). *Probability and Measure*. Wiley. Third Edition.
- Lo G.S. (2019). A series of statistical packages using VB6. Arxiv:
- Lo, G.S.(2016). Weak Convergence (IA). Sequences of random vectors. SPAS Books Series. Saint-Louis, Senegal - Calgary, Canada. Doi : 10.16929/sbs/2016.0001. Arxiv : 1610.05415. ISBN : 978-2-9559183-1-9
- Lo G. S.(2010). A simple note on some empirical stochastic process as a tool in uniform L-statistics weak laws. *Afrika Statistika*, **5**, pp. 437-446. (MR2920301)
- Lo G. S.(2016). How to use the functional empirical process for deriving asymptotic laws for functions of the sample. Arxiv : 1607.02745 (<https://arxiv.org/pdf/1607.02745>)
- Zeghdoudi H and Nedjar S. (2016). A pseudo Lindley-distribution and its application. *Afrika Statistika* Volume 11 (1), pp. 923-932. DOI: <http://dx.doi.org/10.16929/as/2016.923.83>
- Lo G.S., K. T. A. Ngom M. and Diallo M.(2018). Weak Convergence (IIA) - Functional and Random Aspects of the Univariate Extreme Value Theory. Arxiv : 1810.01625
- Ngom and Lo G.S. (2016) A double-indexed functional Hill Process and Applications. *Journal of Mathematical Research* (e-ISSN 1916-9809). Vol 8 (4) , pp. 144-165, Doi : 105539/jmr/v8n4p144

- Shorack G.R. and Wellner J.A. (1986). *Empirical Processes with Applications to Statistics*, Wiley-Interscience, New-York.
- de Haan, L. (1970). *On regular variation and its application to the weak convergence of sample extremes*. Mathematical Centre Tracts, **32**, Amsterdam. (MR0286156)
- de Haan, L. and Ferreira A. (2006). *Extreme value theory: An introduction*. Springer. (MR2234156)
- Lo, G.S. and AhnSanullah M.(2019). Asymptotic Laws of the strong Upper Records Values in the extreme domain of attraction and beyond. arXiv:1905.03380
- Shohat J.A. and Tamarkin J.D (1943) The problem of moment, Mathematical Surveys and monographs. Volume I. American Society of Mathematics (Re-edited in 1950, 1963 and 1970).
- Hill, B.(1975). A simple general approach to the inference about the tail index of a distribution. *Ann Statist.* 3, 1163-1174.
- Loève, Michel.(1997). *Probability Theory I*. Springer-Verlag, 4th Edition.