GLOBAL C^{∞} REGULARITY OF THE STEADY PRANDTL EQUATION

YUE WANG AND ZHIFEI ZHANG

ABSTRACT. In this paper, we prove the global C^{∞} regularity of the Oleinik's solution for the steady Prandtl equation with favorable pressure.

1. INTRODUCTION

In this paper, we study the steady Prandtl equation:

(1.1)
$$\begin{cases} u\partial_x u + v\partial_y u - \partial_y^2 u = -\frac{dp}{dx}(x), & x \ge 0, \ y \ge 0, \\ \partial_x u + \partial_y v = 0, \\ u|_{y=0} = v|_{y=0} = 0 \quad \text{and} \quad \lim_{y \to +\infty} u(x, y) = U(x), \end{cases}$$

where the outer flow (U(x), p(x)) satisfies

(1.2)
$$U(x)U'(x) + p'(x) = 0$$

This system derived by Prandtl could be used to describe the behavior of the solution near y = 0 for the steady Navier-Stokes equations when the viscosity coefficient is small.

The existence and regularity of solution for the system (1.1) was proved by Oleinik [8] for a class of positive data $u_0(y)$ prescribed at x = 0. Let us make it precise. We denote by \mathcal{K} a class of functions, which satisfy

$$u \in C_b^{2,\alpha}([0,+\infty))(\alpha > 0), \quad u(0) = 0, \ u_y(0) > 0, \ u(y) > 0 \text{ for } y \in (0,+\infty),$$
$$\lim_{y \to +\infty} u(y) = U(0) > 0, \quad u_{yy}(y) - \frac{dp}{dx}(0) = O(y^2).$$

Oleinik proved the following classical result(see Theorem 2.1.1 in [8]).

Theorem 1.1. If $u_0 \in \mathcal{K}$ and $\frac{dp}{dx}(x)$ is smooth, then there exists X > 0 such that the steady Prandtl equation (1.1) admits a solution $u \in C^1([0, X) \times \mathbf{R}_+)$ with the following properties:

- 1. regularity: u is bounded and continuous in $[0, X] \times \mathbf{R}_+$; u_y, u_{yy} are bounded and continuous in $[0, X) \times \mathbf{R}_+$; and v, v_y, u_x are locally bounded and continuous in $[0, X) \times \mathbf{R}_+$.
- 2. non-degeneracy: u(x,y) > 0 in $[0,X) \times (0,+\infty)$ and for all $\bar{x} < X$, there exists $y_0 > 0, m > 0$ so that

$$\partial_y u(x,y) \ge m$$
 in $[0,\bar{x}] \times [0,y_0]$.

3. global existence: if $p'(x) \leq 0$, then the solution is global in x.

Then the following problems are natural and important:

• Problem 1. Boundary layer separation in the case of unfavorable pressure, i.e., p'(x) > 0.

Date: March 2, 2022.

- Problem 2. Global C^{∞} regularity of Oleinik's solution in the case of favorable pressure, i.e., $p'(x) \leq 0$.
- Problem 3. Vanishing viscosity limit of the steady Navier-Stokes equations.

For Problem 1, Dalibard and Masmoudi [1] proved the boundary layer separation for a class of special data and p'(x) = 1, and show that the solution behaves near the separation:

$$\partial_y u(x,0) \sim (x^* - x)^{\frac{1}{2}}, \quad x < x^*$$

Shen, Wang and Zhang [9] proved the boundary layer separation for the Oleinik's type data in \mathcal{K} and p'(x) > 0, and studied the local behavior of the solution near the separation point. See also [2] for an unpublished result.

For Problem 3, there are some important progress on the stability for some special boundary layer flows such as the Blasius flow and shear flow [4, 3].

Global C^{∞} regularity of Oleinik's solution is also a long-standing problem. The main challenge is to study the regularity of the solution for a degenerate parabolic equation. Indeed, Oleinik's proof is based on the Von Mises transformation:

(1.3)
$$\psi(x,y) = \int_0^y u(x,z)dz, \quad w = u^2$$

A direct calculation shows that

(1.4)
$$2\partial_y u = \partial_\psi w, \quad 2\partial_y^2 u = \sqrt{w}\partial_\psi^2 w.$$

Then the new unknown $w(x, \psi)$ satisfies

(1.5)
$$\partial_x w - \sqrt{w} \partial_\psi^2 w = -2 \frac{dp}{dx}$$
 in $[0, X) \times \mathbf{R}_+,$

together with

(1.6)
$$w(0,\psi) = w_0(\psi) = u_0(y)^2, \quad w(x,0) = 0, \quad \lim_{\psi \to +\infty} w(x,\psi) = U(x)^2.$$

Now (1.5) is a degenerate parabolic equation due to w = 0 at $\psi = 0$.

Recently, Guo and Iyer [5] proved the higher regularity of the solution in a local time, i.e., $0 < x < X \ll 1$. The goal of this paper is to give an affirmative answer to **Problem 2.** The following is our main result.

Theorem 1.2. Let u be a global solution to (1.1) constructed in Theorem 1.1 with $u_0 \in \mathcal{K}$ and $\frac{dp(x)}{dx} \leq 0$ smooth. For any positive integers m, k and any positive constants X, Y with $\epsilon < X$, there exists a positive constant C depending only on $\epsilon, X, Y, u_0, p, k, m$ so that

$$|\partial_x^k \partial_y^m u(x,y)| \le C \quad for \quad (x,y) \in [\epsilon, X] \times [0,Y].$$

For $y \in [\delta, +\infty)$ and $\delta > 0$, (1.5) is an uniform parabolic equation due to w(x, y) > 0 for $y \ge \delta$ and (1.2) with favorable pressure and U(0) > 0. Thus, in the domain $[\epsilon, +\infty) \times [\delta, +\infty)$, the C^{∞} regularity of the solution is a direct consequence of classical parabolic regularity theory [7]. Therefore, we will focus on the domain $[\epsilon, X] \times [0, \delta]$ for some small $\delta > 0$.

Our result does not give the uniform bounds of $|\partial_x^k \partial_y^m u(x, y)|$ in x, y. In particular, the large time behavior of the solution is a very interesting problem (see also [6]), which will be left to our future work. In the unsteady case, Xin and Zhang [10] proved the global existence of weak solution under the favorable pressure. However, the global C^{∞} regularity up to the boundary remains open.

2. Lower order regularity estimates

In the sequel, we assume that u is a global solution to (1.1) constructed in Theorem 1.1 with $u_0 \in \mathcal{K}$ and $\frac{dp(x)}{dx} \leq 0$ smooth. First of all, we improve the growth estimate of $|\partial_x w|$.

Lemma 2.1. Let $0 < \epsilon < X$, there exists $\delta_1 > 0$ so that

$$|\partial_x w(x,\psi)| < C\psi$$
 for $(x,\psi) \in [\epsilon, X] \times [0,\delta_1].$

Here C is a constant independent of ψ .

Proof. Thanks to Theorem 2.1.14 in [8], it holds in $[0, X] \times [0, \delta_1]$ that

(2.1)
$$|\partial_x w| \le C\psi^{\frac{1}{2}+\alpha}, \quad 0 < m < \partial_\psi w < M \Rightarrow m\psi < w < M\psi,$$

for some $\alpha \in (0, \frac{1}{2})$, and positive constants m, M depending only on X, u_0, p . Without loss of generality, we may assume $\delta_1 < 1$.

Take a smooth cutoff function $0 \leq \zeta(x) \leq 1$ so that $\zeta(x) \equiv 1$ in $[\epsilon, X]$ and $\zeta(x) \equiv 0$ in $[0, \frac{\epsilon}{2}]$. Since w is smooth in the interior, we have, in $(0, X] \times (0, \delta_1]$,

$$\partial_x \partial_x w - \sqrt{w} \partial_\psi^2 \partial_x w = \frac{\partial_x w}{2\sqrt{w}} \partial_\psi^2 w - 2\frac{d^2 p}{dx^2} = \frac{(\partial_x w)^2}{2w} + 2\frac{\partial_x w \frac{dp}{dx}}{2w} - 2\frac{d^2 p}{dx^2},$$

which gives

$$\partial_x \left[\partial_x w \zeta(x) \right] - \sqrt{w} \partial_\psi^2 \left[\partial_x w \zeta(x) \right] = \frac{(\partial_x w)^2}{2w} \zeta(x) + 2 \frac{\partial_x w \frac{dp}{dx}}{2w} \zeta(x) - 2 \frac{d^2 p}{dx^2} \zeta(x) \\ + \partial_x w \partial_x \zeta(x) := I.$$

From (2.1), we know that

(2.2)
$$|I| \le C\psi^{2\alpha} + C\psi^{\alpha - \frac{1}{2}} + C + C\psi^{\frac{1}{2} + \alpha} \le C\psi^{\alpha - \frac{1}{2}}.$$

Next we take $\varphi(\psi) = A_1\psi - A_2\psi^{1+\beta}$ with constants A_1, A_2 big and $\beta \in (0,1)$ to be determined. Then by (2.1) and (2.2), we get

$$\partial_x [\partial_x w \zeta(x) - \varphi] - \sqrt{w} \partial_{\psi}^2 [\partial_x w \zeta(x) - \varphi] \leq |I| - A_2 \sqrt{w} \beta (1+\beta) \psi^{\beta-1}$$
$$\leq C \psi^{\alpha - \frac{1}{2}} - A_2 \sqrt{m} \beta (1+\beta) \psi^{\beta - \frac{1}{2}}.$$

Taking $\beta = \alpha$ and A_2 large enough, we conclude

$$\partial_x \left[\partial_x w \zeta(x) - \varphi \right] - \sqrt{w} \partial_{\psi}^2 \left[\partial_x w \zeta(x) - \varphi \right] < 0 \quad in \quad (0, X] \times (0, \delta_1).$$

We get by (2.1) that for $x \leq X$,

$$(\partial_x w\zeta - \varphi)(x, 0) = 0,$$

$$(\partial_x w\zeta - \varphi)(x, \delta_1) \le M \delta_1^{\frac{1}{2} + \alpha} - A_1 \delta_1 + A_2 \delta_1^{1+\beta},$$

Requiring $A_1 \ge A_2$, by the definition of $\zeta(x)$, we have

$$(\partial_x w \zeta - \varphi)(0, \psi) \le 0$$
 in $[0, \delta_1],$

and take A_1 large enough depending on M, δ_1, A_2 so that

$$(\partial_x w\zeta - \varphi)(x, \delta_1) \le 0.$$

Then the maximum principle ensures that

$$(\partial_x w \zeta - \varphi)(x, \psi) \le 0$$
 in $[0, X] \times [0, \delta_1],$

which implies

$$\partial_x w(x,\psi) \le A_1 \psi - A_2 \psi^{1+\beta} \le \frac{A_1}{2} \psi$$
 in $[\epsilon, X] \times [0, \delta_1],$

if δ_1 is chosen suitably small.

The fact that $-\partial_x w \leq \frac{A_1}{2}\psi$ in $[\epsilon, X] \times [0, \delta_1]$ could be similarly proved by considering $-\partial_x w\zeta - \varphi$.

Proposition 2.2. Let $0 < \epsilon < X$, there exists $\delta_2 > 0$ so that for $(x, \psi) \in [\epsilon, X] \times (0, \delta_2]$,

$$|\partial_{\psi}\partial_{x}w(x,\psi)| \leq C, \quad |\partial_{x}^{2}w(x,\psi)| \leq C\psi^{-\frac{1}{2}}, \quad |\partial_{\psi}^{2}\partial_{x}w(x,\psi)| \leq C\psi^{-1}.$$

Here C is a constant independent of ψ .

Proof. By Lemma 2.1, there exists $\delta_1 > 0$ so that

(2.3)
$$|\partial_x w(x,\psi)| \le C\psi \quad \text{in} \quad \left[\frac{\epsilon}{2}, X\right] \times [0,\delta_1].$$

For any $(x_3, \psi_3) \in [\epsilon, X] \times (0, \delta_0]$ where $\delta_0 = \min\left\{\frac{2}{3}\delta_1, \frac{\epsilon}{2}\right\}$, we denote

$$Q = \left\{ x_3 - \psi_3^{\frac{1}{2}} \le x \le x_3 \right\} \times \left\{ \frac{1}{2} \psi_3 \le \psi \le \frac{3}{2} \psi_3 \right\}.$$

By (2.3) and the definition of δ_0 , we have

$$(2.4) |\partial_x w| \le C\psi \quad \text{in} \quad Q$$

Now we make a transformation

$$T: \quad Q \longrightarrow [-1,0]_{\tilde{x}} \times \left[-\frac{1}{2},\frac{1}{2}\right]_{\tilde{\psi}} := \widetilde{Q},$$
$$(x,\psi) \mapsto (\tilde{x},\tilde{\psi}),$$

where

$$x - x_3 = \psi_3^{\frac{3}{2}} \tilde{x}, \quad \psi - \psi_3 = \psi_3 \tilde{\psi}.$$

Thanks to $\partial_x = \frac{1}{\psi_3^3} \partial_{\tilde{x}}, \partial_{\psi} = \frac{1}{\psi_3} \partial_{\tilde{\psi}}$, we find that

$$\partial_{\tilde{x}}(\psi_3^{-1}w) - \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}} \partial_{\tilde{\psi}}^2(\psi_3^{-1}w) = -2\frac{d}{d\tilde{x}}p\,\psi_3^{-1} \quad \text{in} \quad \widetilde{Q}.$$

It follows from (2.1) that

$$0 < c \le \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}} \le C, \quad |\psi_3^{-1}w| \le C \quad \text{in} \quad \widetilde{Q},$$

and for any $z_1, z_2 \in Q$ (i.e., $\tilde{z}_1 = Tz_1, \tilde{z}_2 = Tz_2 \in \widetilde{Q}$),

$$\left| \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}}(\tilde{z}_1) - \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}}(\tilde{z}_2) \right| = \frac{1}{\psi_3^{\frac{1}{2}}} \frac{|w(z_1) - w(z_2)|}{\sqrt{w}(z_1) + \sqrt{w}(z_2)}$$
$$\leq C \frac{\psi_3 |\tilde{z}_1 - \tilde{z}_2|}{\psi_3} = C |\tilde{z}_1 - \tilde{z}_2|$$

which implies

$$\frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}}\Big|_{\mathcal{C}^{\alpha}(\widetilde{Q})} \le C$$

for any $\alpha \in (0, 1)$. On the other hand, since p is smooth, we have

$$\left|\frac{d}{d\tilde{x}}p\,\psi_3^{-1}\right|_{\mathcal{C}^{0,1}([-1,0]_{\tilde{x}})} \le C.$$

By standard interior a priori estimates (see Theorem 8.11.1 in [7]), we have

(2.5)
$$|\partial_{\tilde{\psi}}^2 w \psi_3^{-1}|_{\mathcal{C}^{\alpha}\left(\left[-\frac{1}{2},0\right]_{\tilde{x}} \times \left[-\frac{1}{4},\frac{1}{4}\right]_{\tilde{\psi}}\right)} \le C$$

Let $f = \partial_x w \psi_3^{-1}$, which satisfies

$$\partial_{\tilde{x}}f - \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}} \partial_{\tilde{\psi}}^2 f - \frac{\partial_{\tilde{\psi}}^2 w}{2\sqrt{w}\psi_3^{\frac{1}{2}}} f = -2\frac{d}{d\tilde{x}}\frac{d}{dx}p\,\psi_3^{-1}.$$

By (2.4), we have

$$|f| \le C \quad \text{in} \quad \widetilde{Q}$$

We write

$$\frac{\partial_{\tilde{\psi}}^2 w}{2\sqrt{w}\psi_3^{\frac{1}{2}}} = \partial_{\tilde{\psi}}^2 w \psi_3^{-1} \frac{\psi_3^{\frac{1}{2}}}{2\sqrt{w}}.$$

Thanks to

$$\frac{\psi_3^{\frac{1}{2}}}{\sqrt{w}}(\tilde{z}_1) - \frac{\psi_3^{\frac{1}{2}}}{\sqrt{w}}(\tilde{z}_2) \Big| = \psi_3^{\frac{1}{2}} \frac{\left|\frac{w(z_1) - w(z_2)}{w(z_1)w(z_2)}\right|}{\frac{1}{\sqrt{w}}(z_1) + \frac{1}{\sqrt{w}}(z_2)} \le C|\tilde{z}_1 - \tilde{z}_2|,$$

we have

(2.6)
$$\left|\frac{\psi_3^{\frac{1}{2}}}{\sqrt{w}}\right|_{\mathcal{C}^{\alpha}(\widetilde{Q})} \le C,$$

which along with (2.5) gives

$$\Big|\frac{\partial_{\tilde{\psi}}^2 w}{2\sqrt{w}\psi_3^{\frac{1}{2}}}\Big|_{\mathcal{C}^{\alpha}([-\frac{1}{2},0]_{\tilde{x}}\times[-\frac{1}{4},\frac{1}{4}]_{\tilde{\psi}})} \le C.$$

As before, since $\frac{dp}{dx}$ is smooth, we have

$$\left| \frac{d}{d\tilde{x}} \frac{d}{dx} p \, \psi_3^{-1} \right|_{\mathcal{C}^{0,1}([-1,0]_{\tilde{x}})} \le C.$$

Then standard interior a priori estimates yield that

 $|\partial_{\tilde{x}}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} + |\partial_{\tilde{\psi}}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} + |\partial_{\tilde{\psi}}^{2}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} \leq C.$

Especially, it holds that

$$|\partial_x^2 w(x_3,\psi_3)| \le C \psi_3^{-\frac{1}{2}}, \quad |\partial_\psi \partial_x w(x_3,\psi_3)| \le C, \quad |\partial_\psi^2 \partial_x w(x_3,\psi_3)| \le C \psi_3^{-1},$$

which give our results.

3. Higher order regularity estimates

In this section, we study the higher order regularity of the solution constructed in Theorem 1.1 with $u_0 \in \mathcal{K}$ and $\frac{dp(x)}{dx} \leq 0$ smooth. The following is one of our main results.

Proposition 3.1. Let $0 < \epsilon < X$ and $k \ge 2$, there exists $\delta > 0$ so that in $[\epsilon, X] \times [0, \delta]$,

 $|\partial_x^k w| \leq C \psi, \quad |\partial_\psi \partial_x^k w| \leq C, \quad |\partial_\psi^2 \partial_x^k w| \leq C \psi^{-1}.$

Here C is a constant independent of ψ .

Proof. The proof is based on the induction argument. Thanks to Lemma 2.1 and Proposition 2.2, we may inductively assume that for $0 \le j \le k - 1$, there holds that in $\left[\frac{\epsilon}{2}, X\right] \times [0, \delta_3]$,

$$(3.1)\left|\partial_{\psi}\partial_{x}^{j}w\right| \leq C, \ \left|\partial_{\psi}^{2}\partial_{x}^{j}w\right| \leq C\psi^{-1}, \ \left|\partial_{x}^{j}w\right| \leq C\psi, \ \left|\partial_{x}^{j}\sqrt{w}\right| \leq C\psi^{\frac{1}{2}}, \ \left|\partial_{x}^{k}w\right| \leq C\psi^{-\frac{1}{2}}.$$

Without loss of generality, assume $\delta_3 \ll 1$. Our goal is to show that there exists $\delta_4 \ll \delta_3$ so that in $[\epsilon, X] \times [0, \delta_4]$,

$$(3.2)|\partial_{\psi}\partial_x^k w| \le C, \ |\partial_{\psi}^2 \partial_x^k w| \le C\psi^{-1}, \ |\partial_x^k w| \le C\psi, \ |\partial_x^k \sqrt{w}| \le C\psi^{\frac{1}{2}}, \ |\partial_x^{k+1} w| \le C\psi^{-\frac{1}{2}}.$$

This will follow from the following Lemma 3.2, Lemma 3.3 and Lemma 3.4.

Lemma 3.2. Assume that (3.1) holds. Then it holds that in $[\frac{7\epsilon}{8}, X] \times [0, \delta_3]$,

$$|\partial_x^k w| < M_1 \psi^{1-\beta}, \quad |\partial_x^k \sqrt{w}| \le M_1 \psi^{\frac{1}{2}-\beta}$$

for any $0 < \beta \ll 1$.

Proof. Fix any $h < \frac{\epsilon}{8}$. Set $D = (0, X] \times (0, \delta_3)$ and let

$$g = \begin{cases} \frac{\partial_x^{k-1}w(x-h,\psi) - \partial_x^{k-1}w(x,\psi)}{-h}\zeta + M\psi\ln\psi & \frac{5\epsilon}{8} \le x \le X, \psi \ge 0, \\ M\psi\ln\psi & 0 \le x < \frac{5\epsilon}{8}, \psi \ge 0, \end{cases}$$

where *M* is a big positive constant to be determined, and $\zeta(x)$ is a smooth cutoff function so that $0 \leq \zeta(x) \leq 1$, $\zeta(x) \equiv 1$ in $[\frac{7}{8}\epsilon, X]$, $\zeta(x) \equiv 0$ in $[0, \frac{5\epsilon}{8}]$.

By the definition of g and continuity of $\partial_x^{k-1}w$ up to $\{\psi = 0\}$, we know that g is smooth in D and continuous in \overline{D} (Growth estimate implies continuity, as pointed out on P38 in [8]). By the assumption, we have g(x,0) = 0. By the definition of ζ , we have $g(0,\psi) \leq 0$. Taking M large enough, we have for $x \in [0, X]$,

$$g(x,\delta_3) \le C(\delta_3)^{-\frac{1}{2}} + M\delta_3 \ln \delta_3 \le 0.$$

Next we show that by a proper choice of M, the positive maximum of g cannot be achieved in the interior. We argue by the contradiction. Assume that there exists a point $(x_0, \psi_0) = p_0 \in D$ such that

$$g(p_0) = \max_{\overline{D}} g > 0,$$

which implies $x_0 > \frac{5\epsilon}{8}$ and

(3.3)
$$\frac{1}{-h} \left(\partial_x^{k-1} w(x_0 - h, \psi_0) - \partial_x^{k-1} w(x_0, \psi_0) \right) > 0.$$

By (2.1), we have

(3.4)
$$-\sqrt{w}\partial_{\psi}^{2}(M\psi\ln\psi) = -M\sqrt{w}\psi^{-1} \leq -\gamma M\psi^{-\frac{1}{2}}$$

$$\Box$$

for $\gamma = \sqrt{m}$. By (1.5), we have in D,

$$\begin{aligned} \partial_x \partial_x^{k-1} w - \sqrt{w} \partial_\psi^2 \partial_x^{k-1} w &= -2 \frac{d^k p}{dx^k} + \sum_{m=1}^{k-2} C_{k-1}^m (\partial_x^{k-1-m} \sqrt{w}) \partial_\psi^2 \partial_x^m w + (\partial_x^{k-1} \sqrt{w}) \partial_\psi^2 u \\ &= -2 \frac{d^k p}{dx^k} + \sum_{m=1}^{k-2} C_{k-1}^m (\partial_x^{k-1-m} \sqrt{w}) \partial_\psi^2 \partial_x^m w + \frac{\partial_x^{k-1} w}{2\sqrt{w}} \frac{\partial_x w}{\sqrt{w}} \\ &+ \left(\frac{\partial_x^{k-1} w}{2\sqrt{w}}\right) \frac{2}{\sqrt{w}} \frac{dp}{dx} + \sum_{m=0}^{k-3} C_{k-2}^m \partial_\psi^2 w \partial_x^{m+1} w \partial_x^{k-2-m} \frac{1}{2\sqrt{w}}.\end{aligned}$$

We denote

$$I_{1} = -2\frac{d^{k}p}{dx^{k}} + \sum_{m=1}^{k-2} C_{k-1}^{m} (\partial_{x}^{k-1-m}\sqrt{w}) \partial_{\psi}^{2} \partial_{x}^{m} w + \frac{\partial_{x}^{k-1}w \partial_{x}w}{2w},$$

$$I_{2} = \left(\frac{\partial_{x}^{k-1}w}{2\sqrt{w}}\right) \frac{2}{\sqrt{w}} \frac{dp}{dx},$$

$$I_{3} = \sum_{m=0}^{k-3} C_{k-2}^{m} \partial_{\psi}^{2} w \partial_{x}^{m+1} w \partial_{x}^{k-2-m} \frac{1}{2\sqrt{w}}.$$

Then for $x \ge \frac{5\epsilon}{8}$,

(3.5)
$$\partial_x g_1 - \sqrt{w(p_h)} \partial_{\psi}^2 g_1 = \frac{\sqrt{w(p_h)} - \sqrt{w(p)}}{-h} \partial_{\psi}^2 \partial_x^{k-1} w(p) + \sum_{i=1}^3 \frac{1}{-h} (I_i(p_h) - I_i(p)).$$

where $g_1 = \frac{1}{-h} \left(\partial_x^{k-1} w(p_h) - \partial_x^{k-1} w(p) \right)$ with $p_h = (x - h, \psi), \ p = (x, \psi).$ Using the induction assumption (3.1), it is easy to verify that for $x \ge \frac{5\epsilon}{8}$,

(3.6)
$$\left|\frac{1}{-h}(\sqrt{w}(p_h) - \sqrt{w}(p))\partial_{\psi}^2 \partial_x^{k-1} w(p)\right| \le C\psi^{-\frac{1}{2}},$$

(3.7)
$$\left| \frac{1}{-h} (I_1(p_h) - I_1(p)) \right| \le C \psi^{-\frac{1}{2}},$$

(3.8)
$$\left| \frac{1}{-h} (I_3(p_h) - I_3(p)) \right| \le C \psi^{-\frac{1}{2}}.$$

We have

$$\frac{1}{-h}(I_2(p_h) - I_2(p)) = g_1 \cdot \left[\frac{1}{w}\frac{dp}{dx}(p_h)\right] + \partial_x^{k-1}w(p)\frac{1}{-h}\left[\frac{1}{w}\frac{dp}{dx}(p_h) - \frac{1}{w}\frac{dp}{dx}(p)\right].$$

By the induction assumption (3.1) and $g_1(p_0) > 0, \frac{dp}{dx} \le 0$, we have

(3.9)
$$\frac{1}{-h}(I_2(p_h) - I_2(p)) \le C \text{ at } p = p_0.$$

Summing up (3.6), (3.7), (3.9), (3.8), we conclude that at $p = p_0$,

$$\partial_x g_1 - \sqrt{w} \partial_{\psi}^2 g_1 \le C_2 \psi^{-\frac{1}{2}}.$$

This along with (3.4) shows that for $x \ge \frac{5\epsilon}{8}$, we have at $p = p_0$,

(3.10)
$$\partial_x g - \sqrt{w} \partial_{\psi}^2 g \le C \psi^{-\frac{1}{2}} - \gamma M \psi^{-\frac{1}{2}}.$$

Take *M* large enough. Then we have $\partial_x g(p_0) - \sqrt{w} \partial_{\psi}^2 g(p_0) < 0$. However, if p_0 is the interior positive maximum point of *g*, we have

$$\partial_x g(p_0) - \sqrt{w} \partial_{\psi}^2 g(p_0) \ge 0,$$

which leads to a contradiction. Hence, for M chosen as above and independent of h, we have

$$\max_{\overline{D}} g \le 0.$$

Similarly, we can show that $\min_{\overline{D}} g_2 \ge 0$, where

$$g_2 = \begin{cases} \frac{\partial_x^{k-1}w(x-h,\psi) - \partial_x^{k-1}w(x,\psi)}{-h} \zeta - M\psi \ln \psi & \frac{5\epsilon}{8} \le x \le X, \psi \ge 0, \\ -M\psi \ln \psi & 0 \le x < \frac{5\epsilon}{8}, \psi \ge 0. \end{cases}$$

Since M is independent of h and h is arbitrary, we have

$$|\partial_x^k w| \le -M\psi \ln \psi$$
 in $(\frac{7}{8}\epsilon, X] \times (0, \delta_3].$

Thanks to

(3.11)
$$2\sqrt{w}\partial_x^k\sqrt{w} + \sum_{m=1}^{k-1} C_k^m(\partial_x^m\sqrt{w}\partial_x^{k-m}\sqrt{w}) = \partial_x^k(\sqrt{w}\sqrt{w}) = \partial_x^kw_x^k$$

we get by (3.1) that in $(\frac{7}{8}\epsilon, X] \times (0, \delta_3]$,

$$|\sqrt{w}\partial_x^k \sqrt{w}| \le -C\psi \ln \psi.$$

Then we have the desired result.

Lemma 3.3. Assume that (3.1) holds. Then it holds that in $\left[\frac{15}{16}\epsilon, X\right] \times [0, \delta_3]$,

$$|\partial_x^k w| \le C\psi, \quad |\partial_x^k \sqrt{w}| \le C\psi^{\frac{1}{2}}$$

Proof. Set

$$g = \partial_x^k w \zeta - A_1 \psi + A_2 \psi^{\frac{3}{2} - \beta},$$

where $\zeta(x)$ is a smooth cutoff function so that $0 \leq \zeta(x) \leq 1$, $\zeta(x) \equiv 1$ in $[\frac{15}{16}\epsilon, X]$, $\zeta(x) \equiv 0$ in $[0, \frac{7\epsilon}{8}]$, and A_1, A_2 are large constants to be determined.

Take $\beta = \frac{1}{200}$ in Lemma 3.2. Then in $[\frac{7\epsilon}{8}, X] \times [0, \delta_3]$,

(3.12)
$$|\partial_x^k w| \le C\psi^{1-\beta}, \quad |\partial_x^k \sqrt{w}| \le C\psi^{\frac{1}{2}-\beta}.$$

Set $D = (0, X] \times (0, \delta_3)$. Then g is smooth in D and continuous in \overline{D} . By (3.12), we have g(x, 0) = 0. Requiring $A_1 \ge A_2$, by the definition of ζ , we have $g(0, \psi) \le 0$. Taking A_1 large depending on A_2 so that for $x \in [0, A]$, $g(x, \delta_3) \le 0$.

Next we show that by a proper choice of A_1, A_2 , the maximum of g cannot be achieved in the interior. By (2.1), we have

(3.13)
$$-\sqrt{w}\partial_{\psi}^{2}(-A_{1}\psi+A_{2}\psi^{\frac{3}{2}-\beta}) \leq -\gamma A_{2}\psi^{-\beta},$$

where $\gamma = (\frac{3}{2} - \beta)(\frac{1}{2} - \beta)\sqrt{m} > 0$. By (1.5), we have in D,

$$\partial_x \partial_x^k w - \sqrt{w} \partial_\psi^2 \partial_x^k w = -2 \frac{d^{k+1}p}{dx^{k+1}} + \sum_{m=0}^{k-1} C_k^m (\partial_x^{k-m} \sqrt{w}) \partial_\psi^2 \partial_x^m w.$$

By (1.5), we have

$$\partial_{\psi}^2 \partial_x^m w = \partial_x^m \partial_{\psi}^2 w = \partial_x^m \Big(\frac{\partial_x w}{\sqrt{w}} + \frac{2}{\sqrt{w}} \frac{dp}{dx} \Big).$$

By (3.12) and (3.1), we have, for $x \ge \frac{7\epsilon}{8}$ and $0 \le j \le k-1$, $|\partial_r^j w| \le C\psi, \quad |\partial_r^k w| \le C\psi^{1-\beta}.$

Then for $0 \le m \le k-1$ and $x \ge \frac{7\epsilon}{8}$,

$$|\partial_{\psi}^2 \partial_x^m w| \le C \psi^{-\frac{1}{2}} + C \psi^{\frac{1}{2}-\beta} \le C \psi^{-\frac{1}{2}},$$

where we note $\beta \ll \frac{1}{2}$. On the other hand, by (3.12) and (3.1), we have for $0 \leq m \leq k-1$,

$$|\partial_x^{k-m}\sqrt{w}| \le C\psi^{\frac{1}{2}-\beta}$$

Hence, we conclude that for $x \ge \frac{7\epsilon}{8}$,

$$\partial_x \partial_x^k w - \sqrt{w} \partial_\psi^2 \partial_x^k w \le C \psi^{-\beta},$$

from which and (3.13), we deduce that at $p = p_0$,

$$\partial_x g - \sqrt{w} \partial_{\psi}^2 g \le C_2 \psi^{-\beta} - \gamma A_2 \psi^{-\beta}.$$

Now take A_2 large depending on C_2 . Then we have $\partial_x g - \sqrt{w} \partial_{\psi}^2 g < 0$ in D, which implies that the maximum of g cannot be achieved in the interior. Hence,

$$\max_{\overline{D}} g \le 0.$$

Similarly, we can prove that $\max_{\overline{D}} -\partial_x^k w\zeta - A_1\psi + A_2\psi^{\frac{3}{2}-\beta} \leq 0$. Hence, we have

$$|\partial_x^k w| \le A_1 \psi - A_2 \psi^{\frac{3}{2}-\beta} \le A_1 \psi$$
 in $\left[\frac{15}{16}\epsilon, X\right] \times [0, \delta_3].$

Moreover, by (3.11) and (3.1), we have that in $\left[\frac{15}{16}\epsilon, X\right] \times [0, \delta_3]$,

$$|\partial_x^k \sqrt{w}| \le C\psi^{\frac{1}{2}}$$

This proves our desired result.

Lemma 3.4. Assume that (3.1) holds. Then it holds that in $[\epsilon, X] \times [0, \delta_4]$,

$$|\partial_{\psi}\partial_x^k w| \le C, \quad |\partial_{\psi}^2 \partial_x^k w| \le C\psi^{-1}, \quad |\partial_x^{k+1} w| \le C\psi^{-\frac{1}{2}}$$

Proof. By Lemma 3.3 and (3.1), it holds that in $\left[\frac{15}{16}\epsilon, X\right] \times [0, \delta_3]$,

(3.14)
$$|\partial_x^j w| \le C\psi, \quad |\partial_x^j \sqrt{w}| \le C\psi^{\frac{1}{2}}, \ 0 \le j \le k.$$

For $(x_3, \psi_3) \in [\epsilon, X] \times (0, \delta_0]$ with $\delta_0 = \min\left\{\frac{2}{3}\delta_3, \frac{\epsilon}{16}\right\}$, we denote

$$Q = \left\{ x_3 - \psi_3^{\frac{3}{2}} \le x \le x_3 \right\} \times \left\{ \frac{1}{2} \psi_3 \le \psi \le \frac{3}{2} \psi_3 \right\}.$$

By straight calculations, we obtain

$$\begin{aligned} \partial_x \partial_x^k w - \sqrt{w} \partial_\psi^2 \partial_x^k w &= -2 \frac{d^{k+1}p}{dx^{k+1}} + \partial_x^k \sqrt{w} \partial_\psi^2 w + \sum_{m=1}^{k-2} C_k^m (\partial_x^{k-m} \sqrt{w}) \partial_\psi^2 \partial_x^m w \\ &+ C_k^{k-1} \frac{\partial_x w}{2\sqrt{w}} \partial_\psi^2 \partial_x^{k-1} w. \end{aligned}$$

Since

$$\begin{split} \partial_{\psi}^{2} \partial_{x}^{m} w &= \partial_{x}^{m} \partial_{\psi}^{2} w = \partial_{x}^{m} \Big(\frac{\partial_{x} w}{\sqrt{w}} + \frac{2}{\sqrt{w}} \frac{dp}{dx} \Big) \\ &= \frac{\partial_{x}^{m+1} w}{\sqrt{w}} + \sum_{l=1}^{m} C_{m}^{l} \partial_{x}^{m-l+1} w \partial_{x}^{l} \frac{1}{\sqrt{w}} + \partial_{x}^{m} \Big(\frac{2}{\sqrt{w}} \frac{dp}{dx} \Big) \end{split}$$

and

$$\partial_x^k \sqrt{w} = \partial_x^{k-1} \frac{\partial_x w}{2\sqrt{w}} = \frac{\partial_x^k w}{2\sqrt{w}} + \sum_{l=1}^{k-1} C_{k-1}^l \partial_x^{k-1-l+1} w \partial_x^l \frac{1}{2\sqrt{w}},$$

we obtain

$$\begin{split} \partial_x \partial_x^k w &- \sqrt{w} \partial_\psi^2 \partial_x^k w = -2 \frac{d^{k+1}p}{dx^{k+1}} + \sum_{m=1}^{k-2} C_k^m (\partial_x^{k-m} \sqrt{w}) \partial_\psi^2 \partial_x^m w \\ &+ \frac{\partial_x^k w}{2\sqrt{w}} \partial_\psi^2 w + \sum_{l=1}^{k-1} C_{k-1}^l \partial_x^{k-l-l+1} w \Big(\partial_x^l \frac{1}{2\sqrt{w}} \Big) \partial_\psi^2 w \\ &+ C_k^{k-1} \frac{\partial_x w \partial_x^k w}{2w} + C_k^{k-1} \frac{\partial_x w}{2\sqrt{w}} \Big[\sum_{l=1}^{k-1} C_{k-1}^l \partial_x^{k-l} w \partial_x^l \frac{1}{\sqrt{w}} + \partial_x^{k-1} \Big(\frac{2}{\sqrt{w}} \frac{dp}{dx} \Big) \Big]. \end{split}$$

Make a transformation $T:Q\to \widetilde{Q}=[-1,0]_{\widetilde{x}}\times [-\frac{1}{2},\frac{1}{2}]_{\widetilde{\psi}}$ via

$$x - x_3 = \psi_3^{\frac{3}{2}} \tilde{x}, \quad \psi - \psi_3 = \psi_3 \tilde{\psi}.$$

Let $f = \partial_x^k w \psi_3^{-1}$, which satisfies

$$\begin{split} \partial_{\tilde{x}}f - \frac{\sqrt{w}}{\psi_{3}^{\frac{1}{2}}}\partial_{\tilde{\psi}}^{2}f + cf &= -2\frac{d^{k+1}p}{dx^{k+1}}\psi_{3}^{\frac{1}{2}} + \psi_{3}^{\frac{1}{2}}\sum_{m=1}^{k-2}C_{k}^{m}(\partial_{x}^{k-m}\sqrt{w})\partial_{\psi}^{2}\partial_{x}^{m}w \\ &+ \psi_{3}^{\frac{1}{2}}\sum_{l=1}^{k-1}C_{k-1}^{l}\partial_{x}^{k-l}w\Big(\partial_{x}^{l}\frac{1}{2\sqrt{w}}\Big)\partial_{\psi}^{2}w \\ &+ \psi_{3}^{\frac{1}{2}}\frac{\partial_{x}w}{2\sqrt{w}}\Big[\sum_{l=1}^{k-1}C_{k-1}^{l}\partial_{x}^{k-l}w\partial_{x}^{l}\frac{1}{\sqrt{w}} + \partial_{x}^{k-1}\Big(\frac{2}{\sqrt{w}}\frac{dp}{dx}\Big)\Big] = F, \end{split}$$

where $c = -\frac{1}{2\sqrt{w}}\partial_{\psi}^2 w \psi_3^{\frac{3}{2}} - \frac{\partial_x w}{2w} \psi_3^{\frac{3}{2}}$. By (3.14), we have

$$|f| \le C \quad \text{in} \quad [-1,0]_{\tilde{x}} \times \left[-\frac{1}{2}, \frac{1}{2}\right]_{\tilde{\psi}}.$$

From the proof of Proposition 2.2, we know that for $\alpha \in (0, 1)$,

$$0 < c \le \frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}} \le C, \quad \left|\frac{\sqrt{w}}{\psi_3^{\frac{1}{2}}}\right|_{\mathcal{C}^{\alpha}([-1,0]_{\tilde{x}} \times [-\frac{1}{2},\frac{1}{2}]_{\tilde{\psi}})} \le C.$$

Using the induction assumption (3.1) and (3.14), we can deduce that for $j \leq k-1$ and $m \leq k-2$,

$$|\nabla_{\tilde{x},\tilde{\psi}}\partial_x^j\sqrt{w}| \le C\psi_3^{\frac{1}{2}}, \quad \left|\nabla_{\tilde{x},\tilde{\psi}}\partial_x^j(\frac{1}{\sqrt{w}})\right| \le C\psi_3^{-\frac{1}{2}}, \quad \left|\nabla_{\tilde{x},\tilde{\psi}}\partial_\psi^2\partial_x^m w\right| \le C\psi_3^{-\frac{1}{2}},$$

where in the third inequality we need to use the fact that

$$\begin{aligned} \partial_{\psi}(\partial_{\psi}^{2}\partial_{x}^{m}w) = & \frac{\partial_{\psi}\partial_{x}^{m+1}w}{\sqrt{w}} - \frac{\partial_{\psi}w\partial_{x}^{m+1}w}{2(\sqrt{w})^{3}} + \sum_{l=1}^{m}C_{m}^{l}\partial_{x}^{m-l+1}\partial_{\psi}w\partial_{x}^{l}\frac{1}{\sqrt{w}} \\ &+ \sum_{l=1}^{m}C_{m}^{l}\partial_{x}^{m-l+1}w\partial_{x}^{l}\frac{\partial_{\psi}w}{-2(\sqrt{w})^{3}} + \partial_{x}^{m}\Big(\frac{\partial_{\psi}w}{-(\sqrt{w})^{3}}\frac{dp}{dx}\Big). \end{aligned}$$

These along with (2.5) and (2.6) imply hat

$$|c|_{\mathcal{C}^{\alpha}(\widetilde{Q})} + |F|_{\mathcal{C}^{\alpha}(\widetilde{Q})} \le C.$$

Then standard interior a priori estimates yield that

$$\partial_{\tilde{x}}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} + |\partial_{\tilde{\psi}}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} + |\partial_{\tilde{\psi}}^{2}f|_{L^{\infty}([-\frac{1}{4},0]_{\tilde{x}}\times[-\frac{1}{8},\frac{1}{8}]_{\tilde{\psi}})} \leq C.$$

Especially, this implies

$$|\partial_x^{k+1}w(x_3,\psi_3)| \le C\psi_3^{-\frac{1}{2}}, \quad |\partial_\psi\partial_x^k w(x_3,\psi_3)| \le C, \quad |\partial_\psi^2\partial_x^k w(x_3,\psi_3)| \le C\psi_3^{-1}.$$

Since (x_3, ψ_3) is arbitrarily, we deduce the desired result.

Corollary 3.5. Let $0 < \epsilon < X$ and integer $k \ge 0$, there exists $\delta > 0$ so that in $[\epsilon, X] \times [0, \delta]$, (3.15) $|\partial_{\psi}^{m} \partial_{x}^{k} w| \le C \psi^{1-m}$.

Proof. By Lemma 2.1, (2.1), Proposition 2.2 and Proposition 3.1, we know that (3.15) holds for m = 0, 1, 2. Moreover, we have

$$\left|\partial_x^k \frac{1}{\sqrt{w}}\right| \le C\psi^{-\frac{1}{2}}, \quad \left|\partial_x^k \partial_\psi \frac{1}{\sqrt{w}}\right| \le C\psi^{-\frac{3}{2}}, \quad \left|\partial_x^k \partial_\psi^2 \frac{1}{\sqrt{w}}\right| \le C\psi^{-\frac{5}{2}}.$$

Let us inductively assume that (3.15) and

(3.16)
$$\left|\partial_x^k \partial_\psi^m \frac{1}{\sqrt{w}}\right| \le C\psi^{-\frac{1}{2}-m}$$

hold for $0 \le m \le l$ with $l \ge 1$. Next we show that for m = l + 1, (3.15) and (3.16) still hold. Thanks to

$$\partial_{\psi}^{l+1} \partial_x^k w = \partial_{\psi}^{l-1} \partial_x^k \partial_{\psi}^2 w = \partial_x^k \partial_{\psi}^{l-1} \left(\frac{\partial_x w}{\sqrt{w}} + \frac{2}{\sqrt{w}} \frac{dp}{dx} \right)$$
$$= \partial_x^k \left(\sum_{a=0}^{l-1} C_{l-1}^a \partial_{\psi}^{l-1-a} \partial_x w \partial_{\psi}^a \frac{1}{\sqrt{w}} + 2 \frac{dp}{dx} \partial_{\psi}^{l-1} \frac{1}{\sqrt{w}} \right)$$

and the induction assumption, we have

$$(3.17) |\partial_{\psi}^{l+1}\partial_x^k w| \le C\psi^{\frac{1}{2}-l}$$

A direct calculation gives

$$\begin{split} 0 = &\partial_x^k \partial_{\psi}^{l+1} \Big(\frac{1}{\sqrt{w}} \frac{1}{\sqrt{w}} w \Big) \\ = &\partial_x^k \Big[2\sqrt{w} \partial_{\psi}^{l+1} \frac{1}{\sqrt{w}} + \sum_{a=1}^l \sum_{j=0}^{l+1-a} C_{l+1}^a C_{l+1-a}^j (\partial_{\psi}^a \frac{1}{\sqrt{w}}) (\partial_{\psi}^j \frac{1}{\sqrt{w}}) \partial_{\psi}^{l+1-j-a} w \\ &+ \sum_{j=0}^l C_{l+1}^j \frac{1}{\sqrt{w}} (\partial_{\psi}^j \frac{1}{\sqrt{w}}) \partial_{\psi}^{l+1-j} w \Big], \end{split}$$

from which and the induction assumption, we can deduce that

$$\left|\partial_x^k \partial_\psi^{l+1} \frac{1}{\sqrt{w}}\right| \le C\psi^{-\frac{3}{2}-l}.$$

This along with (3.17) gives the desired result.

4. Global C^{∞} regularity

In this section, we prove Theorem 1.2. As we remark below Theorem 1.2, it is enough to prove the regularity of the solution on the domain $[\epsilon, X] \times [0, \delta]$ for some small $\delta > 0$.

Proof. To make the notation clear, we write the Von Mises transformation as follows

$$(\tilde{x},\psi) = \left(x, \int_0^y u \, dy\right).$$

A direct calculation (or See P25 in [8]) shows that

$$\partial_y = \sqrt{w} \partial_\psi, \quad \partial_x = \partial_{\tilde{x}} + \partial_x \psi(x, y) \partial_\psi,$$
$$\partial_x \psi = \frac{1}{2} \sqrt{w} \int_0^\psi w^{-\frac{3}{2}} \partial_{\tilde{x}} w \, d\psi.$$

By (2.1) and Proposition 3.1, we have

$$(4.1) |\partial_x \psi| \le C \psi.$$

Thanks to $\partial_y = \sqrt{w} \partial_{\psi}$, we find that

$$\begin{aligned} \partial_x^k 2\partial_y u &= (\partial_{\tilde{x}} + \partial_x \psi \partial_{\psi})^k \partial_{\psi} w, \\ \partial_x^k 2\partial_y^2 u &= (\partial_{\tilde{x}} + \partial_x \psi \partial_{\psi})^k \left(\partial_{\tilde{x}} w + 2\frac{dp}{dx} \right) \\ &= (\partial_{\tilde{x}} + \partial_x \psi \partial_{\psi})^k (\partial_{\tilde{x}} w) + 2\frac{d^{k+1}p}{dx^{k+1}}. \end{aligned}$$

Using (4.1) and Corollary 3.5, we can deduce that

$$(4.2) |\partial_x^k \partial_y u| + |\partial_x^k \partial_y^2 u| \le C.$$

Assume that for $m \ge 1$ and any integer k,

$$(4.3) |\partial_x^k \partial_y^j u| \le C, \quad j \le m$$

Using the fact that

$$\begin{split} \partial_x^k \partial_y^{m+1} u = &\partial_x^k \partial_y^{m-1} \partial_y^2 u = \partial_x^k \partial_y^{m-1} \Big(u \partial_x u + \partial_y u \int_0^y -\partial_x u \, dy \Big) \\ = &\partial_x^k \Big(\sum_{a=0}^{m-1} C_{m-1}^a \partial_y^{m-1-a} u \partial_y^a \partial_x u - \sum_{a=0}^{m-2} C_{m-1}^{a+1} \partial_y^{m-1-a} u \partial_y^a \partial_x u + \partial_y^m u \int_0^y -\partial_x u \, dy \Big), \end{split}$$

we can deduce from (4.2) and (4.3) that

$$|\partial_x^k \partial_y^j u| \le C, \quad j \le m+1.$$

This proves the theorem.

12

г		_

Acknowledgments

The authors thank Professor Yan Guo for helpful discussions. Y. Wang is partially supported by Beijing Advanced Innovation Center for Imaging Theory and Technology and key research project of the Academy for Multidisciplinary Studies, Capital Normal University.

References

- A. Dalibard and N. Masmoudi, Separation for the stationary Prandtl equation, Publ. Math. Inst. Hautes Études Sci., 130 (2019), 187-297.
- [2] W. E, Boundary layer theory and the zero-viscosity limit of the Navier-Stokes equation, Acta Math. Sin., 16(2000), 207-218.
- [3] D. Gerard-Varet and Y. Maekawa, Sobolev stability of Prandtl expansions for the steady Navier-Stokes equations, Arch. Ration. Mech. Anal., 233 (2019), 1319-1382.
- [4] Y. Guo and S. Iyer, Validity of steady Prandtl layer expansions, arXiv:1805.05891.
- [5] Y. Guo and S. Iyer, Regularity and expansion for steady Prandtl equations, arXiv:1903.08086.
- [6] S. Iyer, On global-in-x stability of Blasius profiles, arXiv:1812.03906.
- [7] Krylov, N. V. Lectures on elliptic and parabolic equations in Hölder spaces, vol. 12 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1996.
- [8] O. A. Oleinik and V. N. Samokhin, Mathematical models in boundary layer theory, Applied Mathematics and Mathematical Computation 15. Chapman & Hall/CRC, Boca Raton, FL, 1999.
- W. Shen, Y. Wang and Z. Zhang, Boundary layer separation and local behavior for the steady Prandtl equation, arXiv:1904.08055.
- [10] Z. Xin and L. Zhang, On the global existence of solutions to the Prandtl's system, Adv. Math., 181 (2004), 88-133.

SCHOOL OF MATHEMATICAL SCIENCES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, CHINA *E-mail address:* yuewang37@pku.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, 100871, BEIJING, P. R. CHINA *E-mail address:* zfzhang@math.pku.edu.cn