

Bayesian high-dimensional linear regression with generic spike-and-slab priors

Bai Jiang^{*} and Qiang Sun[†]

Abstract

Spike-and-slab priors are popular Bayesian solutions for high-dimensional linear regression problems. Previous theoretical studies on spike-and-slab methods focus on specific prior formulations and use prior-dependent conditions and analyses, and thus can not be generalized directly. In this paper, we propose a class of generic spike-and-slab priors and develop a unified framework to rigorously assess their theoretical properties. Technically, we provide general conditions under which generic spike-and-slab priors can achieve the nearly-optimal posterior contraction rate and the model selection consistency. Our results include those of [Narisetty and He \(2014\)](#) and [Castillo et al. \(2015\)](#) as special cases.

Keywords: high-dimensional linear regression, generic spike-and-slab prior, model selection, posterior contraction.

1 Introduction

Consider the linear regression model

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta}^* + \sigma_*\boldsymbol{\varepsilon}, \quad (1.1)$$

where $\mathbf{X} \in \mathbb{R}^{n \times p}$ is a deterministic design matrix, $\boldsymbol{\beta}^* \in \mathbb{R}^p$ is a vector of unknown regression coefficients, $\sigma_* > 0$ is unknown standard deviation, and $\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ is a standard normal vector. We are interested in parameter estimation and model selection in the high-dimensional regime where $p \gg n$ and a small number s of covariates contribute to the response. Formally, we assume the index set of non-zero regression coefficients $\xi_* := \{j : \beta_j^* \neq 0\}$ is s -sparse. The goals are to estimate unknown parameters $(\sigma_*, \boldsymbol{\beta}^*)$ and to identify the true sparse model (index set) ξ_* .

For this high-dimensional linear regression problem, many methods have been proposed from the Bayesian perspective. They commonly encourage sparsity of the regression coefficients by adopting suitable priors ([Park and Casella, 2008](#); [Hans, 2009](#); [Carvalho et al.,](#)

^{*}ByteDance AI Lab, 250 Bryant Street, Mountain View, CA 94041, USA; Email: bai.jiang@bytedance.com.

[†]Department of Statistical Sciences, University of Toronto, 100 St. George Street, Toronto, ON M5S 3G3, Canada; Email: qsun@utstat.toronto.edu.

2010; Griffin and Brown, 2012; Armagan et al., 2013; Narisetty and He, 2014; Castillo et al., 2015; Bhattacharya et al., 2015; Martin et al., 2017; Rockova and George, 2018). These priors can be mainly categorized into two categories: shrinkage priors and spike-and-slab priors.

The shrinkage priors are directly motivated by the equivalence between regularized maximum likelihood estimators in the frequentist framework (among others, Tibshirani, 1996; Fan and Li, 2001; Zou, 2006; Candes and Tao, 2007; Zhang, 2010) and *maximum a posteriori* estimators in the Bayesian framework. Examples include the Bayesian lasso prior (Park and Casella, 2008; Hans, 2009), the horseshoe prior (Carvalho et al., 2010; Polson and Scott, 2012), the correlated normal-gamma prior (Griffin and Brown, 2012), the double Pareto prior (Armagan et al., 2013), the Dirichlet-Laplace prior (Bhattacharya et al., 2015) and the spike-and-slab lasso prior (Rockova and George, 2018), to name a few. Recently Song and Liang (2017) provide sufficient conditions for generic shrinkage priors to achieve the nearly-optimal parameter estimation rate and the model selection consistency.

The spike-and-slab priors are hierarchical priors which naturally arise from the probabilistic consideration of the high-dimensional linear regression model (1.1). A generic spike-and-slab prior $\Pi(\sigma, \xi, \beta)$ takes the form of

$$\begin{aligned}\sigma^2 &\sim g(\sigma^2), \\ \xi &\sim \pi(\xi), \quad \xi \subseteq \{1, \dots, p\}, \\ \beta_j | \sigma^2 &\sim h_0(\beta_j / \sigma) / \sigma, \quad \forall j \notin \xi, \\ \beta_j | \sigma^2 &\sim h_1(\beta_j / \sigma) / \sigma, \quad \forall j \in \xi,\end{aligned}\tag{1.2}$$

where g is a density function over $(0, \infty)$, $\xi \subseteq \{1, \dots, p\}$ indexes all possible 2^p subset models, $\pi(\xi)$ is a model selection prior introducing model sparsity, h_0 is a “spike” distribution for modeling negligible coefficients (e.g., the Dirac measure at 0) and h_1 is a “slab” distribution for modeling significant coefficients. This generic spike-and-slab prior of form (1.2) dates back to the Dirac-and-slab priors (Mitchell and Beauchamp, 1988; George and McCulloch, 1993; Johnson and Rossell, 2012) and mixture Gaussian priors (Ishwaran and Rao, 2005) in the small- p -large- n setup. Later, in the high-dimensional regime, Narisetty and He (2014) showed that a mixture Gaussian prior with shrinking and diffusing scale hyperparameters consistently selects the true sparse model. Castillo et al. (2015) studied both the parameter estimation rate and the model selection consistency for Dirac-and-Laplace priors. Other recent works also consider the correlated Gaussian distribution as the slab prior (Yang et al., 2016; Martin et al., 2017).

Despite of these works, theoretical properties for generic spike-and-slab priors of form (1.2) remain unclear. Previous works usually narrow their focuses down to specific choices of spike-and-slab formulations, such as the mixture Gaussian prior (Narisetty and He, 2014) and the Dirac-and-Laplace prior (Castillo et al., 2015), and conduct theoretical assessments under conditions that cope with their choices of formulations. Consequently, their analyses rely on various conditions and are not generalizable to other spike-and-slab priors.

Regarding the model selection prior $\pi(\xi)$, a popular choice is the independently and identically distributed (i.i.d.) Bernoulli prior (Narisetty and He, 2014), in which each covariate j is independently selected into the model ξ with probability $1/p$. Castillo et al. (2015, Assumption 1) considered a class of model selection priors exponentially decreasing on model size, which we refer to as the Castillo-Schmidt-Vaart priors or simply the CSV priors. Note that the i.i.d. Bernoulli prior does not belong to the class of CSV priors. Regarding the spike and slab distributions, popular choices include the Laplace or the Gaussian distribution and the Dirac measure at 0 as the spike distribution only (Narisetty and He, 2014; Castillo et al., 2015; Rockova and George, 2018). Although some combinations of the above-mentioned model selection priors, spike priors and slab priors have been recognized, the potential of other combinations for Bayesian high-dimensional linear regression has been overlooked.

On the other hand, different conditions on the eigen-structure of the Gram matrix $\mathbf{X}^T \mathbf{X}$ have been proposed and assumed. Examples include conditions on the minimum non-zero eigenvalue (Narisetty and He, 2014, MNEV), the minimum restricted eigenvalue (Castillo et al., 2015, MREV), and the minimum sparse eigenvalue (Song and Liang, 2015, MSEV). The MREV condition has been widely assumed for frequentist methods (Bickel et al., 2009; Raskutti et al., 2010; Fan et al., 2018). As for Bayesian methods, however, it is unclear how these different conditions relate to each other and whether the results built on one of them can transfer to the other.

In this paper, we develop a unified framework to analyze Bayesian methods with generic spike-and-slab priors. This framework could not only facilitate theoretical assessments of a broad class of spike-and-slab priors, but also unifies seemingly different conditions for existing spike-and-slab methods.

First, for the parameter estimation task, we give a high-level condition for model selection prior $\pi(\xi)$, which is satisfied by both CSV priors and the i.i.d. Bernoulli prior. Another interesting finding is that MNEV, MREV and MSEV are lower bounds of a quantity, which we call the *minimum united eigenvalue* (MUEV) and denote by λ . A positive λ suffices for the Bayesian spike-and-slab methods to succeed. Second, for the model selection task, we show that, under the commonly-seen beta-min condition (Bühlmann and Van De Geer, 2011, Corollary 7.6), the generic spike-and-slab prior selects overfitted models that overshoot the true model size by no more than a constant factor. Finally, we identify two more technical conditions, which enable eliminating false discoveries in the overfitted models and consistently selecting true sparse model. Conditions for previous spike-and-slab methods (Castillo et al., 2015; Narisetty and He, 2014) are shown to be special cases of our conditions tailored to their specific spike-and-slab priors.

The rest of the paper proceeds as follows. Section 2 introduces a class of generic spike-and-slab priors. Section 3 builds up the posterior contraction of parameters (σ, β) upon the new local eigenvalue condition relating to MUEV λ . Section 4 presents additional conditions and theorems for the model selection task. Section 5 sketches the proofs of theorems, with

proofs of technical lemmas deferred to the appendix. Section 6 concludes the paper with a discussion.

Notation

For the high-dimensional linear regression model (1.1), both p and s should be understood as sequences of n , i.e., $p = p_n$ and $s = s_n$, although their subscripts n are omitted. Similarly, for the spike-and-slab prior $\Pi(\sigma, \xi, \beta)$ specified by (1.2), π , h_0 and h_1 should be understood as sequences of distributions π_n , h_{0n} and h_{1n} . Let

$$\epsilon_n := \sqrt{s \log p/n},$$

be the nearly-optimal¹ ℓ_2 -estimation error rate for estimating β^* .

For $\beta \in \mathbb{R}^p$, let β_j denote its j -th component and β_ξ denote its sub-vector consisting of coordinates in the subset $\xi \subseteq \{1, \dots, p\}$. We also call the index set ξ a model in the context of model selection. For a vector v , let $\|v\|_q$ with $q \in [1, \infty]$ denote its ℓ_q -norm. For ℓ_2 -norm, we omit the subscript 2 and write $\|v\|$ for simplicity.

We write \mathbf{X}_j to denote the j -th column of \mathbf{X} and \mathbf{X}_ξ to denote the sub-matrix consisting of columns indexed by $\xi \in \{1, \dots, p\}$. For a model ξ , let $|\xi|$ be its cardinality, and $\text{rank}(\xi)$ be the column rank of \mathbf{X}_ξ . The model ξ is said to be of full rank if $\text{rank}(\xi) = |\xi|$. Let \mathcal{F} denote the set of all full-rank models. Formally

$$\mathcal{F} := \{\xi \subseteq \{1, \dots, p\} : \text{rank}(\xi) = |\xi|\}.$$

Let \mathbf{P}_ξ denote the projection matrix onto the column space of \mathbf{X}_ξ . Note that, in case of $\xi \in \mathcal{F}$, $\mathbf{P}_\xi = \mathbf{X}_\xi (\mathbf{X}_\xi^\top \mathbf{X}_\xi)^{-1} \mathbf{X}_\xi^\top$. For $\xi \in \mathcal{F}$, let $\mathbf{X}_\xi^\dagger = (\mathbf{X}_\xi^\top \mathbf{X}_\xi)^{-1} \mathbf{X}_\xi^\top$ be the left pseudo-inverse of \mathbf{X}_ξ .

For a symmetric matrix \mathbf{A} , we write its largest eigenvalue as $\lambda_{\max}(\mathbf{A})$ and its smallest eigenvalue as $\lambda_{\min}(\mathbf{A})$. For two symmetric matrices \mathbf{A} and \mathbf{B} , $\mathbf{A} \geq \mathbf{B}$ or $\mathbf{B} \leq \mathbf{A}$ means $\mathbf{A} - \mathbf{B}$ is positive semi-definite. For two positive sequences a_n and b_n , $a_n \prec b_n$ or $b_n \succ a_n$ means $\limsup_{n \rightarrow \infty} a_n/b_n = 0$; $a_n \preceq b_n$ or $b_n \succeq a_n$ means $\limsup_{n \rightarrow \infty} a_n/b_n < \infty$; $a_n \asymp b_n$ means $a_n \preceq b_n$ and $a_n \succeq b_n$; $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ means $a_n > b_n$ for sufficiently large n . We write $o(1)$ to denote an arbitrarily small positive constant.

Let $\mathbb{P}_{(\sigma, \xi, \beta)}$ and $\mathbb{E}_{(\sigma, \xi, \beta)}$ denote the measure and expectation associated with model (1.1) with parameters (σ, ξ, β) . Write $\mathbb{P}_{(\sigma_*, \xi_*, \beta_*)}$ and $\mathbb{E}_{(\sigma_*, \xi_*, \beta_*)}$ as \mathbb{P}_* and \mathbb{E}_* for simplicity. When the probability $\mathbb{P}_{(\sigma, \xi, \beta)}(E)$ of an event E does not depend on (σ, ξ, β) , we write $\mathbb{P}_{(\sigma, \xi, \beta)}(E)$ as $\mathbb{P}(E)$.

¹The optimal ℓ_2 -estimation error rate is $\sqrt{s \log(p/s)/n}$; see [Raskutti et al. \(2011\)](#) and [Su and Candes \(2016\)](#) among others.

2 A Class of Generic Spike-and-slab Priors

Throughout the paper, we tacitly assume that the response vector \mathbf{Y} has been centered at 0, and thus include no intercept term in the linear regression model (1.1). Each covariate \mathbf{X}_j is centered and standardized such that $\|\mathbf{X}_j\| = \sqrt{n}$. The true standard deviation $\sigma_\star \in (0, \infty)$ is fixed. The true model ξ_\star is of full rank. We focus on the asymptotic regime where $p > n$ but $\epsilon_n = \sqrt{s \log p/n} \rightarrow 0$.

Our following assumption specifies a class of generic spike-and-slab priors.

Assumption 2.1 (On Prior).

- (a) Variance prior: The density function of variance $g(\sigma^2)$ is continuous and positive for any $\sigma^2 \in (0, +\infty)$.
- (b) Model selection prior: The model ξ is selected *a priori* with probability being proportional to $\pi(\xi)1\{\xi \in \mathcal{F}\}$; And, $\pi(\xi)$ satisfies $\pi(\emptyset) \asymp 1$, and with constants $A_1, A_2 > 0$,

$$\pi(\xi_\star) \geq p^{-A_1 s}, \quad \sum_{\xi: |\xi| > t} \pi(\xi) \leq p^{-A_2 t}, \quad \forall t \geq 1$$

for sufficiently large n^2 .

- (c) Spike prior: The sequence z_{0n} such that $\int 1\{|z| > z_{0n}\} h_0(z) dz = e^{-n}$ satisfies

$$z_{0n} \prec \frac{1}{p} \sqrt{\frac{\log p}{n}}.$$

- (d) Slab prior: For $z_{1n} := \max_{j \in \xi_\star} |\beta_j^\star / \sigma_\star| + \epsilon_n$ and some constant $A_3 > 0$, the slab density function $h_1(z)$ satisfies

$$\inf_{z: |z| \leq z_{1n}} h_1(z) \succeq p^{-A_3}.$$

Assumption 2.1(a) is satisfied when g is the inverse-gamma density function. If σ_\star^2 is known to take values in an interval, we could set g to be a continuous and positive density function over that interval, e.g., the truncated inverse-gamma density function.

Assumption 2.1(b) requires the model selection prior $\pi(\xi)$ to assign a sufficient mass to the true sparse model ξ_\star , and to downweight large models. The following examples show that this requirement is met by the commonly-used i.i.d. Bernoulli prior (Narisetty and He, 2014), and the CSV priors including the complexity prior and the Binomial-Beta prior (Castillo et al., 2012). Appendix S.1 collects the detailed proofs.

Example 2.2 (I.I.D. Bernoulli Prior). The i.i.d. Bernoulli prior $\pi(\xi)$ selects each index j into ξ with probability $1/p$. This prior meets the condition for π in Assumption 2.1(b) with $\pi(\emptyset) \approx e^{-1}$, and any $A_1 > 1$ and $A_2 = 1$. The deduction of $A_2 = 1$ needs a novel tail probability inequality due to Pelekis (2016, Theorem 1).

²Recall that both $p = p_n$ and $s = s_n$ are sequences of n .

Example 2.3 (CSV Prior). Suppose $w(t)$ is a discrete distribution over possible model sizes $t \in \{0, \dots, p\}$, and

$$B_1 p^{-B_3} w(t-1) \leq w(t) \leq B_2 p^{-B_4} w(t-1), \quad t = 1, \dots, p,$$

with constants $B_1, B_2, B_3, B_4 > 0$. The prior $\pi(\xi)$ for model selection given by

$$\pi(\xi) = w(|\xi|) \binom{p}{|\xi|}^{-1}$$

meets Assumption 2.1(b) with any $A_1 > B_3 + 1$ and any $A_2 < B_4$.

Assumption 2.1(c) requires the spike distribution to be the Dirac measure at zero or degenerate to it at an appropriate rate. This rate would ensure the aggregated signal of inactive covariates with *a priori* regression coefficients to be negligible

$$\max_{\xi \supseteq \xi_\star} \frac{\|\mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c}\|}{\sqrt{n}} \prec \sqrt{\frac{\log p}{n}} \leq \epsilon_n.$$

Assumption 2.1(d) avoids excessive thinness of the slab distribution around the true regression coefficients, which would otherwise cause the slab prior to miss true non-zero regression coefficients. In case of the Laplace slab distribution $h_1(z) = (\rho_n/2) \exp(-\rho_n|z|)$, the choice of the inverse-scale hyper-parameter $p^{-A'_3} \preceq \rho_n \preceq A''_3 \log p / z_{1n}$ fulfills Assumption 2.1(d) with $A_3 = A'_3 + A''_3$. In case of the Gaussian slab distribution $h_1(z) = (2\pi\tau_{1n}^2)^{-1/2} \exp[-z^2/(2\tau_{1n}^2)]$, the choice of the variance hyper-parameter $A'_3 z_{1n}^2 / \log p \preceq \tau_{1n}^2 \preceq p^{A''_3}$ fulfills Assumption 2.1(d) with $A_3 = 1/A'_3 + A''_3/2$.

3 Posterior Contraction

It is impossible to estimate the coefficients $\boldsymbol{\beta}_\star$ in the high-dimensional linear regression model (1.1) without conditions on the Gram matrix $\mathbf{X}^\top \mathbf{X}$. Indeed, the Gram matrix is non-invertible in the high-dimensional regime, rendering an unidentifiability issue. A common remedy is to assume some kind of “local invertibility” of the Gram matrix. We formalize this idea in the following definition and assumption.

Definition 3.1 (Minimum United Eigenvalue (MUEV)). The minimum united eigenvalue of order t for the design matrix \mathbf{X} is defined as

$$\text{MUEV}(t) := \min_{\xi \in \mathcal{F}: |\xi \cup \xi_\star| \leq t} \lambda_{\min}(\mathbf{X}_{\xi \cup \xi_\star}^\top \mathbf{X}_{\xi \cup \xi_\star} / n),$$

where \mathcal{F} is the set of all full-rank models ξ .

Assumption 3.2 (MUEV Condition). There exists constant $K > 0$ such that

$$\lambda = \lambda(K) := \text{MUEV}((K+1)s) > 0.$$

We also collect other local eigenvalues used in the literature.

Definition 3.3 (Minimum Restricted Eigenvalue (MREV)). The minimum restricted eigenvalue of order t (with parameter $\alpha \geq 1$) is defined as

$$\text{MREV}(t) := \min_{\xi: |\xi| \leq t} \inf_{\beta} \left\{ \frac{\beta^T \mathbf{X}^T \mathbf{X} \beta}{n \|\beta\|^2} : \beta \neq \mathbf{0}, \|\beta_{\xi^c}\|_1 \leq \alpha \|\beta_{\xi}\|_1 \right\}.$$

Definition 3.4 (Minimum Sparse Eigenvalue (MSEV)). The minimum sparse eigenvalue of order t is defined as

$$\text{MSEV}(t) := \min_{\xi: |\xi| \leq t} \lambda_{\min}(\mathbf{X}_{\xi}^T \mathbf{X}_{\xi} / n).$$

Definition 3.5 (Minimum Non-zero Eigenvalue (MNEV)). The minimum non-zero eigenvalue of order t is defined as

$$\text{MNEV}(t) := \min_{\xi: |\xi| \leq t} \lambda_{\min}^N(\mathbf{X}_{\xi}^T \mathbf{X}_{\xi} / n),$$

where $\lambda_{\min}^N(\mathbf{A})$ denotes the minimum non-zero eigenvalue of a symmetric matrix \mathbf{A} .

The next lemma discusses the relation of MUEV to other local eigenvalues of the Gram matrix. It states that the MUEV condition is the weakest among other possible conditions defined upon MSEV, MREV and MNEV. Note that the premise for $\text{MUEV} \geq \text{MNEV}$ in the lemma was assumed in the original paper of MNEV; see (Narisetty and He, 2014, Condition 4.5).

Lemma 3.6. For any $t > s$,

$$\text{MUEV}(t) \geq \text{MSEV}(t) \geq \text{MREV}(t).$$

If $(\mathbf{I} - \mathbf{P}_{\xi})\mathbf{X}_{\xi_{\star}} \neq \mathbf{0}$ for any $\xi \not\supseteq \xi_{\star}$ of size $|\xi| \leq t$, then

$$\text{MUEV}(t) \geq \text{MNEV}(t).$$

Now we are ready to present our main results regarding the posterior contraction rate of β in terms of ℓ_2 -norm.

Theorem 3.7 (Posterior Contraction). Suppose Assumption 2.1 and Assumption 3.2 hold with $A_1 + A_3 + 1 < A_2 K$. For any constants $M_1, M_2 > \sqrt{8 \max\{A_2, 1\} K}$, the posterior distribution $\Pi(\sigma, \xi, \beta | \mathbf{X}, \mathbf{Y})$ concentrates on the subset of the parameter space

$$\hat{\Theta} = \left\{ (\sigma, \xi, \beta) : \begin{array}{l} \frac{\sigma^2}{\sigma_{\star}^2} \in \left[\frac{1 - M_1 \epsilon_n}{1 + M_1 \epsilon_n}, \frac{1 + M_1 \epsilon_n}{1 - M_1 \epsilon_n} \right], \\ |\xi \setminus \xi_{\star}| \leq Ks, \xi \in \mathcal{F}, \\ \max_{j \notin \xi} |\beta_j| \leq \sigma z_{0n}, \\ \|\beta - \beta^{\star}\| \leq M_2 \sigma_{\star} \epsilon_n / \sqrt{\lambda} \end{array} \right\}$$

in the sense that, with some constants C_1, C_2 ,

$$\mathbb{P}_{\star} \left(\Pi(\hat{\Theta} | \mathbf{X}, \mathbf{Y}) \geq 1 - e^{-C_1 s \log p} \right) \gtrsim 1 - e^{-C_2 s \log p}.$$

This theorem establishes the posterior contraction rate of regression coefficients β in terms of ℓ_2 -norm. Roughly speaking, the posterior distribution puts almost all mass in an ϵ_n -ball centering at true coefficients β^* , with high probability. The posterior contraction rate is a standard metric to evaluate estimation accuracy of Bayesian approaches (Ghosal et al., 2000; Shen and Wasserman, 2001).

Two appealing byproducts of Theorem 3.7 are the adaptivity of the posterior distribution to the unknown variance σ_\star^2 , and the unknown sparsity level s . The working variance σ^2 accurately estimates σ_\star^2 up to a relative error of order ϵ_n . The working model ξ does not overshoot the true model size $s = |\xi_\star|$ by more than a constant factor K .

Additionally, Theorem 3.7 allows $\lambda = \lambda(K)$ in Assumption 3.2 to decrease to zero as $n \rightarrow \infty$, providing broader applicabilities. In this case, Theorem 3.7 gives the posterior contraction rate $\epsilon_n/\sqrt{\lambda}$. When λ is of constant order, this rate is near optimal.

4 Model Selection

The task of consistent model selection is harder than the task of parameter estimation, and thus requires more assumptions. Indeed, if some coordinates of β^* are too close to zero, then no method can detect these nearly-zero coordinates as being non-zero. In such case, the posterior distribution may select only a subset of the true model and possibly other coordinates (with nearly-zero coefficients). At the same time, the parameter estimation could be still accurate due to the negligible coefficient values of these false discoveries. To avoid these extreme cases that cause the model selection task to fail, we need some kind of “beta-min” condition (Bühlmann and Van De Geer, 2011) on the minimal value of the true coefficients β^* .

Assumption 4.1 (Beta-min Condition). There exists a constant $M_3 > 0$ such that

$$\min_{j \in \xi_\star} |\beta_j^*| \geq \frac{M_3 \sigma_\star \epsilon_n}{\sqrt{\lambda}}.$$

Under this assumption, the posterior distribution would select all active covariates. This insight is formalized in the following theorem.

Theorem 4.2 (Overfitted Model Selection). If assumptions in Theorem 3.7 as well as Assumption 4.1 with $M_3 > \sqrt{8A_3K}$ hold, then the posterior distribution $\Pi(\sigma, \xi, \beta | \mathbf{X}, \mathbf{Y})$ concentrates on the subset of the parameter space

$$\tilde{\Theta} = \left\{ (\sigma, \xi, \beta) : \begin{array}{l} \frac{\sigma^2}{\sigma_\star^2} \in \left[\frac{1 - M_1 \epsilon_n}{1 + M_1 \epsilon_n}, \frac{1 + M_1 \epsilon_n}{1 - M_1 \epsilon_n} \right], \\ |\xi \setminus \xi_\star| \leq Ks, \xi \in \mathcal{F}, \xi \supseteq \xi_\star, \\ \max_{j \notin \xi} |\beta_j| \leq \sigma z_{0n}, \\ \|\beta_\xi - \beta_\xi^*\| \leq M_2 \sigma_\star \epsilon_n / \sqrt{\lambda} \end{array} \right\}.$$

in the sense that, with some constants C_3, C_4 ,

$$\mathbb{P}_\star \left(\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}) \geq 1 - e^{-C_3 s \log p} \right) \gtrsim 1 - e^{-C_4 s \log p}.$$

However, Theorem 4.2 does not guarantee the elimination of false discoveries in the overfitted models. To achieve the true sparse model, we need to bound

$$\underbrace{\frac{\Pi(\tilde{\Theta} \cap \{\xi = \gamma\}|\mathbf{X}, \mathbf{Y})}{\Pi(\tilde{\Theta} \cap \{\xi = \xi_\star\}|\mathbf{X}, \mathbf{Y})}}_{\text{posterior ratio}} = \underbrace{\frac{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi = \gamma)}{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi = \xi_\star)}}_{\text{conditional posterior ratio}} \times \underbrace{\frac{\pi(\gamma)}{\pi(\xi_\star)}}_{\text{prior ratio}}$$

for each overfitted model $\gamma \supset \xi_\star$ with $|\gamma| \leq (K+1)s$. The two terms on the right-hand side of the last display are the conditional posterior and prior ratios between models γ and ξ_\star respectively. Given some continuity condition of the slab distribution h_1 , the posterior ratio can be bounded as

$$\text{conditional posterior ratio} \leq 2 \left(\frac{\sqrt{2\pi} \sup_z h_1(z) p^{1+o(1)}}{\sqrt{n\lambda}} \right)^{|\gamma|-s}.$$

For many diffusing slab distributions that have diminishing maximum values $\sup_z h_1(z) \rightarrow 0$, the conditional posterior ratio is upper bounded by the diffusing rate of $h_1(z)$ after proper normalization. The prior ratio is solely determined by the model selection prior $\pi(\xi)$. Combining these pieces together yields the following theorem for model selection consistency of spike-and-slab methods.

Theorem 4.3 (Consistent Model Selection). Suppose assumptions in Theorem 4.2 and the following two conditions hold.

- (a) The log-slab function $\log h_1(z)$ is L_n -Lipschitz continuous on $[-z_{1n}, +z_{1n}]$ with $sL_n \prec \sqrt{n \log p}$.
- (b) With some small constant $\eta > 0$,

$$r_n := \max_{j=1}^{Ks} \left[\frac{\pi(|\xi| = s+j)}{\pi(\xi_\star)} \right]^{1/j} \times \frac{\sup_z h_1(z) p^{1+\eta}}{\sqrt{n\lambda}} < 1.$$

Then, with constants C_5, C_6 ,

$$\mathbb{P}_\star \left(\Pi(\xi = \xi_\star|\mathbf{X}, \mathbf{Y}) \geq 1 - e^{-C_5 s \log p} - r_n \right) \gtrsim 1 - p^{-C_6}.$$

Consequently, if $r_n \prec 1$ then $\Pi(\xi = \xi_\star|\mathbf{X}, \mathbf{Y})$ converges to 1 in expectation and in probability.

The expression of rate r_n precisely characterizes the roles of the model selection prior $\pi(\xi)$ and the diffusing slab distribution $h_1(z)$ in a successful Bayesian model selection procedure. Conditions of Narisetty and He (2014); Castillo et al. (2015) are special cases of our general conditions in Theorem 4.3.

Example 4.4 (Narisetty and He, 2014). For the Gaussian slab distribution with the variance parameter τ_{1n}^2 and the i.i.d. Bernoulli prior of model selection (see Example 2.2),

$$L_n \asymp \tau_{1n}^{-2}, \quad \max_{j=1}^{Ks} \left[\frac{\pi(|\xi| = s + j)}{\pi(\xi_*)} \right]^{1/j} \leq 1.$$

(as they assume fixed true coefficients β^* , and thus $z_{1n} \asymp 1$). In this setup, conditions (a) and (b) in Theorem 4.3 turn to be

$$n\tau_{1n}^2\lambda \prec p^{2+2\eta}.$$

This is the rate assumed by Narisetty and He (2014, the first display of Section 2.1).

Example 4.5 (Castillo et al., 2015). For the Laplace slab distribution with the inverse-scale parameter ρ_n and the CSV prior of model selection (see Example 2.3),

$$L_n = \rho_n, \quad \max_{j=1}^{Ks} \left[\frac{\pi(|\xi| = s + j)}{\pi(\xi_*)} \right]^{1/j} \leq (K + 1)s \times B_3 p^{-B_4}.$$

In this setup, conditions (a) and (b) in Theorem 4.3 are satisfied if

$$s\rho_n\sqrt{\log p/n} \prec 1, \quad s < p^\eta, \quad \rho_n \leq 4\sqrt{n \log p}$$

for some $\eta < B_4 - 1$. These sufficient conditions have been used by Castillo et al. (2015, Corollary 1).

5 Proofs of Theorems

In this section, we sketch proofs of Theorem 3.7, Theorem 4.2 and Theorem 4.3. Technical lemmas and their proofs are collected in the appendix.

5.1 Proof of Theorem 3.7

The claimed inequality of Theorem 3.7 is equivalent to

$$\mathbb{P}_\star \left(\Pi(\hat{\Theta}^c | \mathbf{X}, \mathbf{Y}) \geq e^{-C_1 s \log p} \right) \lesssim e^{-C_2 s \log p}. \quad (5.1)$$

Our central technique to prove (5.1) is the following lemma, which is borrowed from Barron (1998, Lemma 6) and Song and Liang (2017, Lemma A4).

Lemma 5.1. Consider a parametric model $\{P_\theta\}_{\theta \in \Theta}$, and a data generation \mathbf{D} from the true parameter $\theta_\star \in \Theta$. Let $\Pi(\theta)$ be a prior distribution over Θ . If

- (a) $\Pi(\Theta_0) \leq \delta_0$,
- (b) there exists a test function $\phi(\mathbf{D})$ such that

$$\sup_{\theta \in \Theta_{\text{test}}} \mathbb{E}_\theta[1 - \phi(\mathbf{D})] \leq \delta_1, \quad \mathbb{E}_{\theta_\star}[\phi(\mathbf{D})] \leq \delta'_1,$$

(c) and

$$\mathbb{P}_{\theta_*} \left(\int_{\Theta} \frac{P_{\theta}(\mathbf{D})}{P_{\theta_*}(\mathbf{D})} d\Pi(\theta) \leq \delta_2 \right) \leq \delta'_2,$$

then for any δ_3 ,

$$\mathbb{P}_{\theta_*} \left(\Pi(\Theta_0 \cup \Theta_{\text{test}} | \mathbf{D}) \geq \frac{\delta_0 + \delta_1}{\delta_2 \delta_3} \right) \leq \delta'_1 + \delta'_2 + \delta_3.$$

Specifically, in the linear regression model (1.1),

$$\mathbf{D} = (\mathbf{X}, \mathbf{Y}), \quad \theta = (\sigma, \xi, \beta), \quad P_{\theta}(\mathbf{D}) = \mathcal{N}(\mathbf{Y} | \mathbf{X}\beta, \sigma^2 \mathbf{I}).$$

The high-level idea is that, to get the posterior concentration on a desired subset $\hat{\Theta}$ of the parameter space, one can split the set of undesired parameter values $\hat{\Theta}^c$ as two subsets Θ_0 and Θ_{test} , with parameter values in Θ_0 received negligible *a priori* probability mass and parameter values in Θ_{test} distinguished from θ_* by a uniformly powerful test ϕ .

We first construct Θ_0 , Θ_{test} and ϕ , which will be used to prove (5.1) in the framework described by Lemma 5.1. For Θ_0 and Θ_{test} , let

$$\begin{aligned} \Theta_0 &= \{(\sigma, \xi, \beta) \in \Theta : \xi \notin \mathcal{F}\} \cup \{(\sigma, \xi, \beta) \in \Theta : |\xi \setminus \xi_*| > Ks\} \\ &\quad \cup \left\{ (\sigma, \xi, \beta) \in \Theta : \max_{j \notin \xi} |\beta_j| > \sigma z_{0n} \right\}, \\ \Theta_{\text{test}} &= \Theta_1 \cup \Theta_2, \end{aligned}$$

with

$$\begin{aligned} \Theta_1 &= \left\{ (\sigma, \xi, \beta) \in \Theta_0^c : \frac{\sigma^2}{\sigma_*^2} \notin \left[\frac{1 - M_1 \epsilon_n}{1 + M_1 \epsilon_n}, \frac{1 + M_1 \epsilon_n}{1 - M_1 \epsilon_n} \right] \right\}, \\ \Theta_2 &= \left\{ (\sigma, \xi, \beta) \in \Theta_0^c \cap \Theta_1^c : \|\beta - \beta^*\| > M_2 \sigma_* \epsilon_n / \sqrt{\lambda} \right\}. \end{aligned}$$

We take the test function as

$$\phi = \max\{\phi_1, \phi_2\},$$

with

$$\begin{aligned} \phi_1 &= 1 \left\{ \max_{\xi \in \mathcal{F}: |\xi \setminus \xi_*| \leq Ks} \left| \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \mathbf{Y}}{n \sigma_*^2} - 1 \right| > M_1 \epsilon_n \right\}, \\ \phi_2 &= 1 \left\{ \max_{\xi \in \mathcal{F}: |\xi \setminus \xi_*| \leq Ks} \left\| \mathbf{X}_{\xi \cup \xi_*}^\dagger \mathbf{Y} - \beta_{\xi \cup \xi_*}^* \right\| > M_2 \sigma_* \epsilon_n / 2\sqrt{\lambda} \right\}. \end{aligned}$$

Next, by the following lemmas, the conditions required by Lemma 5.1 are verified with the above-defined Θ_0 , Θ_{test} and ϕ .

Lemma 5.2. Suppose Assumption 2.1(b)(c) hold. Then

$$\Pi(\Theta_0) \lesssim e^{-[A_2 K - o(1)]s \log p}.$$

Lemma 5.3.

$$\mathbb{E}_\star \phi_1 \lesssim e^{-[M_1^2/8-K-o(1)]n\epsilon_n^2}, \quad \sup_{(\sigma,\xi,\beta) \in \Theta_1} \mathbb{E}_{(\sigma,\xi,\beta)}(1 - \phi_1) \lesssim e^{-[M_1^2/8-o(1)]n\epsilon_n^2}.$$

Lemma 5.4. Suppose Assumption 3.2 holds. Then

$$\mathbb{E}_\star \phi_2 \lesssim e^{-[M_2^2/8-K-o(1)]n\epsilon_n^2}, \quad \sup_{(\sigma,\xi,\beta) \in \Theta_2} \mathbb{E}_{(\sigma,\xi,\beta)}(1 - \phi_2) \lesssim e^{-[M_2^2/8-o(1)]n\epsilon_n^2}.$$

Lemma 5.5. Suppose Assumption 2.1 holds. For any small constant $\eta > 0$,

$$\mathbb{P}_\star \left(\int \frac{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I})}{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta^\star, \sigma_\star^2 \mathbf{I})} d\Pi(\sigma, \xi, \beta) \leq e^{-(A_1+A_3+1-\eta)s \log p} \right) \lesssim e^{-C_\eta s \log p},$$

with constant $C_\eta > 0$ depending on η .

In particular, Lemma 5.2 verifies condition (a) of Lemma 5.1 with

$$\delta_0 = e^{-[A_2 K - o(1)]s \log p}.$$

Lemmas 5.3-5.4 verifies condition (b) of Lemma 5.1 as follows:

$$\begin{aligned} \sup_{(\sigma,\xi,\beta) \in \Theta_{\text{test}}} \mathbb{E}_{(\sigma,\xi,\beta)}(1 - \phi) &\leq \max_{j=1}^2 \left\{ \sup_{(\sigma,\xi,\beta) \in \Theta_j} \mathbb{E}_{(\sigma,\xi,\beta)}(1 - \phi_j) \right\} \\ &\lesssim e^{-[\min\{M_1, M_2\}^2/8-o(1)]s \log p} =: \delta_1, \\ \mathbb{E}_\star \phi &\leq \sum_{j=1}^2 \mathbb{E}_\star \phi_j \lesssim e^{-[\min\{M_1, M_2\}^2/8-K-o(1)]s \log p} =: \delta'_1. \end{aligned}$$

Lemma 5.5 verifies condition (c) of Lemma 5.1 with

$$\delta_2 = e^{-[A_1+A_3+1-o(1)]s \log p}, \quad \delta'_2 = e^{-C_\eta s \log p}.$$

Finally, we note that $\delta_0 > \delta_1$, because $M_1, M_2 > \sqrt{8A_2 K}$. Choosing suitable $\delta_3 = e^{-C s \log p}$ such that

$$A_1 + A_3 + 1 - o(1) < C < A_2 K - o(1)$$

completes the proof.

5.2 Proof of Theorem 4.2

Since

$$\widehat{\Theta} \cap \{(\sigma, \xi, \beta) : \xi \supseteq \xi_\star\} \subseteq \widetilde{\Theta},$$

it suffices to show that

$$\mathbb{P}_\star \left(\Pi((\sigma, \xi, \beta) \in \widehat{\Theta}^c \text{ or } \xi \not\supseteq \xi_\star | \mathbf{X}, \mathbf{Y}) \geq e^{-C_3 s \log p} \right) \lesssim e^{-C_4 s \log p}. \quad (5.2)$$

To this end, we use the technique developed from Lemma 5.1 again. Recall notation Θ_0 , Θ_1 , Θ_2 , ϕ_1 and ϕ_2 in the proof of Theorem 3.7 and redefine (with a little abuse of notation)

$$\Theta_{\text{test}} = \Theta_1 \cup \Theta_2 \cup \Theta_3, \quad \phi = \max\{\phi_1, \phi_2, \phi_3\},$$

where

$$\begin{aligned} \Theta_3 &= \{(\sigma, \xi, \beta) \in \Theta_0^c \cap \Theta_1^c : \xi \not\supseteq \xi_\star\}, \\ \phi_3 &= 1 \left\{ \min_{\xi \in \mathcal{F}: |\xi \setminus \xi_\star| \leq Ks, \xi \not\supseteq \xi_\star} \|(\mathbf{P}_{\xi \cup \xi_\star} - \mathbf{P}_\xi) \mathbf{Y}\| < M_3 \sigma_\star \sqrt{n} \epsilon_n / 2 \right\}. \end{aligned}$$

With $M_3 = \sqrt{8A_2K}$,

$$\Theta_0 \cup \Theta_{\text{test}} = \widehat{\Theta}^c \cup \{(\sigma, \xi, \beta) : \xi \not\supseteq \xi_\star\}.$$

We proceed to verify conditions of Lemma 5.1. As in the proof of Theorem 3.7, Lemma 5.2 and Lemma 5.5 have verified conditions (a) and (c). It is only left to verify condition (b) for redefined Θ_{test} and ϕ . The following lemma serves for this purpose.

Lemma 5.6. Suppose Assumption 3.2 and Assumption 4.1 hold with $M_3 = \sqrt{8A_2K}$. Then

$$\mathbb{E}_\star \phi_3 \lesssim e^{-[M_3^2/8 - o(1)]n\epsilon_n^2}, \quad \sup_{(\sigma, \xi, \beta) \in \Theta_3} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_3) \lesssim e^{-[M_3^2/8 - o(1)]n\epsilon_n^2}.$$

Next, combining Lemmas 5.3-5.4 and Lemma 5.6 yields

$$\begin{aligned} \sup_{(\sigma, \xi, \beta) \in \Theta_{\text{test}}} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi) &\leq \max_{j=1}^3 \left\{ \sup_{(\sigma, \xi, \beta) \in \Theta_j} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_j) \right\} \\ &\lesssim e^{-[\min\{M_1, M_2, M_3\}^2/8 - o(1)]s \log p} =: \delta_1, \\ \mathbb{E}_\star \phi &\leq \sum_{j=1}^3 \mathbb{E}_\star \phi_j \lesssim e^{-[\min\{M_1^2/8 - K, M_2^2/8 - K, M_3^2\} - o(1)]s \log p} =: \delta'_1. \end{aligned}$$

5.3 Proof of Theorem 4.3

The proof of Theorem 3.7 uses two technical lemmas, which are stated as follows.

Lemma 5.7. For any constant $\eta > 0$, let

$$\Omega_1(\eta) = \cup_{t=s+1}^{(K+1)s} \cup_{\xi \in \mathcal{F}: \xi \supseteq \xi_\star, |\xi|=t} \{\|(\mathbf{P}_\xi - \mathbf{P}_{\xi_\star})\boldsymbol{\epsilon}\|^2 \geq (2 + \eta)(t - s) \log p\}$$

and $\Omega_2 = \{\|\boldsymbol{\epsilon}\| \geq 2\sqrt{n}\}$. Then $\Omega(\eta) := \Omega_1(\eta) \cup \Omega_2$ satisfies

$$\mathbb{P}(\Omega(\eta)) \lesssim e^{-C_\eta s \log p},$$

for some constant C_η depending on η .

Lemma 5.8. Suppose Assumption 2.1(c), Assumption 3.2 and condition (b) of Theorem 4.3 hold. Conditional on event $\Omega(\eta)^c$ (with $\Omega(\eta)$ defined in Lemma 5.7),

$$\sup_{\xi \in \mathcal{F}: \xi \supseteq \xi_*, |\xi|=t} \frac{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi)}{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi_*)} \leq 2 \left(\sqrt{\frac{2\pi}{n\lambda}} \times \sup_z h_1(z) \times p^{1+\eta} \right)^{t-s}$$

for any $t = s + 1, \dots, (K + 1)s$, for sufficiently large $n > N$ (where N does not depends on t).

Proof of Theorem 4.3. From Lemma 5.8 and condition (b), it follows that

$$\begin{aligned} & \sum_{\gamma \in \mathcal{F}: \gamma \supseteq \xi_*, |\gamma|=t} \frac{\Pi(\tilde{\Theta} \cap \{\xi = \gamma\}|\mathbf{X}, \mathbf{Y})}{\Pi(\tilde{\Theta} \cap \{\xi = \xi_*\}|\mathbf{X}, \mathbf{Y})} \\ &= \sum_{\gamma \in \mathcal{F}: \gamma \supseteq \xi_*, |\gamma|=t} \frac{\pi(\gamma)}{\pi(\xi_*)} \times \frac{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi = \gamma)}{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi = \xi_*)} \\ &\lesssim \frac{\pi(|\xi| = t)}{\pi(\xi_*)} \times \left(\sqrt{\frac{2\pi}{n\lambda}} \times \sup_z h_1(z) \times p^{1+\eta} \right)^{t-s} \\ &\leq r_n^{t-s} \end{aligned}$$

Then

$$\sum_{\substack{\gamma \in \mathcal{F}: \gamma \supseteq \xi_*, \\ s+1 \leq |\gamma| \leq (K+1)s}} \frac{\Pi(\tilde{\Theta} \cap \{\xi = \gamma\}|\mathbf{X}, \mathbf{Y})}{\Pi(\tilde{\Theta} \cap \{\xi = \xi_*\}|\mathbf{X}, \mathbf{Y})} \lesssim \sum_{t=s+1}^{(K+1)s} r_n^{t-s} \leq \frac{r_n}{1 - r_n}.$$

Combining this result with Theorem 4.2 concludes the proof. \square

6 Discussion

In this paper, we identify a class of generic spike-and-slab priors and then develop a unified theoretical framework to analyze these spike-and-slab methodologies. Comparing with the literature, we characterize the weakest conditions to guarantee near optimal posterior contraction rate and consistent model selection property. Our conditions and results are general and include previous works as special cases.

Acknowledgement

We would like to thank Professor Edward George, Professor Faming Liang, Professor Qifan Song and Professor Jianqing Fan for helpful discussions in the initial stage of this project.

References

ARMAGAN, A., DUNSON, D. B. and LEE, J. (2013). Generalized double pareto shrinkage. *Statistica Sinica* **23** 119.

- BARRON, A. R. (1998). Information-theoretic characterization of bayes performance and the choice of priors in parametric and nonparametric problems. *Bayesian Statistics* **6** 27–52.
- BHATTACHARYA, A., PATI, D., PILLAI, N. S. and DUNSON, D. B. (2015). Dirichlet-Laplace priors for optimal shrinkage. *Journal of the American Statistical Association* **110** 1479–1490.
- BICKEL, P. J., RITOV, Y. and TSYBAKOV, A. B. (2009). Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics* **37** 1705–1732.
- BÜHLMANN, P. and VAN DE GEER, S. (2011). *Statistics for high-dimensional data: Methods, theory and applications*. Springer, New York, US.
- CANDES, E. and TAO, T. (2007). The Dantzig selector: Statistical estimation when p is much larger than n . *The Annals of Statistics* **35** 2313–2351.
- CARVALHO, C. M., POLSON, N. G. and SCOTT, J. G. (2010). The horseshoe estimator for sparse signals. *Biometrika* **97** 465–480.
- CASTILLO, I., SCHMIDT-HIEBER, J. and VAN DER VAART, A. (2015). Bayesian linear regression with sparse priors. *Annals of Statistics* **43** 1986–2018.
- CASTILLO, I., VAN DER VAART, A. ET AL. (2012). Needles and straw in a haystack: Posterior concentration for possibly sparse sequences. *The Annals of Statistics* **40** 2069–2101.
- FAN, J. and LI, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association* **96** 1348–1360.
- FAN, J., LIU, H., SUN, Q. and ZHANG, T. (2018). I-LAMM for sparse learning: Simultaneous control of algorithmic complexity and statistical error. *The Annals of statistics* **46** 814.
- GEORGE, E. I. and MCCULLOCH, R. E. (1993). Variable selection via gibbs sampling. *Journal of the American Statistical Association* **88** 881–889.
- GHOSAL, S., GHOSH, J. K. and VAN DER VAART, A. W. (2000). Convergence rates of posterior distributions. *The Annals of Statistics* **28** 500–531.
- GRIFFIN, J. E. and BROWN, P. J. (2012). Structuring shrinkage: Some correlated priors for regression. *Biometrika* **99** 481–487.
- HANS, C. (2009). Bayesian Lasso regression. *Biometrika* **96** 835–845.
- ISHWARAN, H. and RAO, J. S. (2005). Spike and slab gene selection for multigroup microarray data. *Journal of the American Statistical Association* **100** 764–780.

- JOHNSON, V. E. and ROSSELL, D. (2012). Bayesian model selection in high-dimensional settings. *Journal of the American Statistical Association* **107** 649–660.
- LAURENT, B. and MASSART, P. (2000). Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics* **28** 1302–1338.
- LIU, J. (2005). Eigenvalue and singular value inequalities of schur complements. In *The Schur Complement and Its Applications*, chap. 2. Springer, New York, US, 47–82.
- MARTIN, R., MESS, R., WALKER, S. G. ET AL. (2017). Empirical bayes posterior concentration in sparse high-dimensional linear models. *Bernoulli* **23** 1822–1847.
- MITCHELL, T. J. and BEAUCHAMP, J. J. (1988). Bayesian variable selection in linear regression. *Journal of the American Statistical Association* **83** 1023–1032.
- NARISSETTY, N. N. and HE, X. (2014). Bayesian variable selection with shrinking and diffusing priors. *The Annals of Statistics* **42** 789–817.
- PARK, T. and CASELLA, G. (2008). The Bayesian Lasso. *Journal of the American Statistical Association* **103** 681–686.
- PELEKIS, C. (2016). A lower bound on binomial tails: an approach via tail conditional expectations. *arXiv preprint arXiv:1609.06651* .
- POLSON, N. G. and SCOTT, J. G. (2012). Local shrinkage rules, lévy processes and regularized regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **74** 287–311.
- RASKUTTI, G., WAINWRIGHT, M. J. and YU, B. (2010). Restricted eigenvalue properties for correlated gaussian designs. *Journal of Machine Learning Research* **11** 2241–2259.
- RASKUTTI, G., WAINWRIGHT, M. J. and YU, B. (2011). Minimax rates of estimation for high-dimensional linear regression over ℓ_q -balls. *IEEE Transactions on Information Theory* **57** 6976–6994.
- ROCKOVA, V. and GEORGE, E. I. (2018). The spike-and-slab Lasso. *Journal of the American Statistical Association* **113** 431–444.
- SHEN, X. and WASSERMAN, L. (2001). Rates of convergence of posterior distributions. *The Annals of Statistics* **29** 687–714.
- SONG, Q. and LIANG, F. (2015). High-dimensional variable selection with reciprocal ℓ_1 -regularization. *Journal of the American Statistical Association* **110** 1607–1620.
- SONG, Q. and LIANG, F. (2017). Nearly optimal bayesian shrinkage for high dimensional regression. *arXiv preprint arXiv:1712.08964* .

- SU, W. and CANDÈS, E. (2016). Slope is adaptive to unknown sparsity and asymptotically minimax. *The Annals of Statistics* **44** 1038–1068.
- TIBSHIRANI, R. (1996). Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **58** 267–288.
- YANG, Y., WAINWRIGHT, M. J. and JORDAN, M. I. (2016). On the computational complexity of high-dimensional bayesian variable selection. *The Annals of Statistics* **44** 2497–2532.
- ZHANG, C.-H. (2010). Nearly unbiased variable selection under minimax concave penalty. *The Annals of statistics* **38** 894–942.
- ZOU, H. (2006). The adaptive Lasso and its oracle properties. *Journal of the American Statistical Association* **101** 1418–1429.

Appendix

S.1 Proofs for Example 2.2 and Example 2.3

We collect proofs for Example 2.2 and Example 2.3 by showing both the Bernoulli prior and the CSV priors satisfy Assumption 2.1 (b).

Proof for Example 2.2. It is elementary that

$$\pi(\emptyset) = \left(1 - \frac{1}{p}\right)^p \rightarrow e^{-1}$$

as $p \rightarrow \infty$. Next, let $T \sim \text{Binomial}(p, 1/p)$. Due to Lemma S.3.1,

$$\begin{aligned} \sum_{\xi: |\xi| > t} \pi(\xi) &= \mathbb{P}(T > t) \leq \mathbb{P}(T \geq t+1) \\ &\leq \frac{(1/p)^{2(t+1)}}{2} \times \binom{p}{t+1} \bigg/ \binom{t+1}{t+1} \leq p^{-t}. \end{aligned}$$

Finally,

$$\pi(\xi_*) = \left(\frac{1}{p}\right)^s \left(1 - \frac{1}{p}\right)^{p-s} \gtrsim p^{-s} \times \frac{e^{-1}}{2}.$$

□

Proof for Example 2.3. Recall that both $p = p_n$ and $s = s_n$ are assumed to be sequences of n . First, find large integer N_1 such that, for any $n > N_1$, $B_2 p^{-B_4} \leq 1/2$. Then

$$1 = \sum_{t=0}^p w(t) \leq w(0) (1 + B_2 p^{-B_4} + B_2^2 p^{-2B_4} + \dots) \leq \frac{w(0)}{1 - B_2 p^{-B_4}} \leq 2w(0)$$

implying $w(0) \geq 1/2$. Next, find large integer N_2 such that, for any $n > N_2$, $\log(2B_2) < (B_4 - A_2) \log p$. Then, for any $n > \max\{N_1, N_2\}$ and any $t \geq 0$,

$$\begin{aligned} \sum_{\xi: |\xi| > t} \pi(\xi) &\leq \sum_{j=t}^p w(j) \leq w(t) (1 + B_2 p^{-B_4} + B_2^2 p^{-2B_4} + \dots) \\ &\leq \frac{w(t)}{1 - B_2 p^{-B_4}} \leq 2w(t) \leq 2w(0) B_2^t p^{-B_4 t} \leq (2B_2)^t p^{-B_4 t} \leq p^{-A_2 t}. \end{aligned}$$

Third, find large integer N_3 such that, for any $n > N_3$, $\log(B_3/2) < (A_1 - B_3 - 1) \log p$. Then, for any $n > \max\{N_1, N_3\}$,

$$\begin{aligned} \pi(\xi_*) &= w(s) \binom{p}{s}^{-1} \geq w(0) B_1^s p^{-B_3 s} \binom{p}{s}^{-1} \\ &\geq w(0) B_1^s p^{-B_3 s} p^{-s} \geq (B_1/2)^s p^{-(B_3+1)s} \geq p^{-A_1 s} \end{aligned}$$

Therefore, $N = \max\{N_1, N_2, N_3\}$ is the desideratum.

□

S.2 Proofs of Technical Lemmas

Proof of Lemma 3.6. The first inequality is trivial. The second inequality follows from the facts that

$$\lambda_{\min}(\mathbf{X}_\xi^\top \mathbf{X}_\xi) = \inf_{\boldsymbol{\beta}} \left\{ \frac{\boldsymbol{\beta}^\top \mathbf{X}^\top \mathbf{X} \boldsymbol{\beta}}{\|\boldsymbol{\beta}\|^2} : \boldsymbol{\beta}_{\xi^c} = \mathbf{0}, \boldsymbol{\beta}_\xi \neq \mathbf{0} \right\}$$

and that

$$\boldsymbol{\beta}_{\xi^c} = \mathbf{0}, \boldsymbol{\beta}_\xi \neq \mathbf{0} \implies \boldsymbol{\beta} \neq \mathbf{0}, \|\boldsymbol{\beta}_{\xi^c}\|_1 \leq \alpha \|\boldsymbol{\beta}_\xi\|_1.$$

For the third inequality, it suffices to show upon the identifiability condition that any full-rank model ξ such that $|\xi \cup \xi_\star| \leq t$ results in a full-rank model $\xi \cup \xi_\star$. This is obvious for cases of underfitted models $\xi \subseteq \xi_\star$. For other cases, we prove by contradiction. Suppose for the sake of contradiction that $\xi \cup \xi_\star$ with $\xi \not\subseteq \xi_\star$ is not of full rank. We construct a vector basis of model $\xi_\star \cup \xi$ by merging all vectors of $\xi \setminus \xi_\star$ and some selected vectors of ξ_\star . Let $\gamma \subseteq \xi_\star$ denote the index set of the selected vectors. We must have $\gamma \subset \xi_\star$ since $\xi \cup \xi_\star$ is not of full rank. Therefore, $(\xi \setminus \xi_\star) \uplus \gamma \not\subseteq \xi_\star$, but $(\mathbf{I} - \mathbf{P}_{(\xi \setminus \xi_\star) \uplus \gamma}) \mathbf{X}_{\xi_\star} = \mathbf{0}$, which contradict to the premise. \square

Proof of Lemma 5.2. Due to Assumption 2.1(b),

$$\begin{aligned} \Pi(|\xi \setminus \xi_\star| > Ks) &= \frac{\sum_{\xi \in \mathcal{F}: |\xi \setminus \xi_\star| > Ks} \pi(\xi)}{\sum_{\xi \in \mathcal{F}} \pi(\xi)} \\ &\leq \frac{\sum_{\xi: |\xi| > Ks} \pi(\xi)}{\pi(\emptyset)} \\ &\lesssim p^{-[A_2 K - o(1)]s}. \end{aligned}$$

Due to Assumption 2.1(c), for any $\xi \subseteq \{1, \dots, p\}$ any $\sigma > 0$,

$$\Pi\left(\max_{j \notin \xi} |\beta_j| > \sigma z_{0n} \mid \xi, \sigma\right) \leq p \int 1\{|z| > z_{0n}\} h_0(z) dz \leq e^{-n + \log p}.$$

Putting these pieces together completes the proof. \square

Proof of Lemma 5.3, part(a). Write

$$\begin{aligned} \phi_1 &= 1 \left\{ \max_{\xi \in \mathcal{F}: |\xi \setminus \xi_\star| \leq Ks} \left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_\star}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n \right\} \\ &\leq 1 \left\{ \max_{\xi: |\xi \setminus \xi_\star| \leq Ks} \left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_\star}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n \right\}. \end{aligned}$$

Since projection matrices $\mathbf{P}_{\xi_1 \cup \xi_\star} \leq \mathbf{P}_{\xi_2 \cup \xi_\star}$ for nested models $\xi_1 \subseteq \xi_2$, the quantity $\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_\star}) \boldsymbol{\varepsilon}$ achieves its minimum value at some ξ of size Ks and its maximum value at $\xi = \emptyset$.

$$\begin{aligned} \phi_1 &\leq 1 \left\{ \max_{\xi: |\xi| = Ks \text{ or } 0} \left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_\star}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n \right\} \\ &\leq \sum_{\xi: |\xi| = Ks \text{ or } 0} 1 \left\{ \left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_\star}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n \right\}. \end{aligned}$$

For each ξ of size Ks or 0, write

$$\begin{aligned} & \mathbb{P}_* \left(\left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n \right) \\ & \leq \mathbb{P} \left(\frac{\chi_{n-s}^2}{n} > 1 + M_1 \epsilon_n \right) + \mathbb{P} \left(\frac{\chi_{n-(K+1)s}^2}{n} < 1 - M_1 \epsilon_n \right). \end{aligned}$$

Putting the last two displays together with the probability bound of the chi-square distribution (Lemma S.3.2, part (b)) yields

$$\mathbb{E}_* \phi_1 \leq \left[1 + \binom{p}{Ks} \right] \times e^{-[M_1^2/8 - o(1)]n\epsilon_n^2} \leq e^{-[M_1^2/8 - K - o(1)]n\epsilon_n^2}.$$

□

Proof of Lemma 5.3, part(b). Define

$$\begin{aligned} \phi_{1,\gamma} &= 1 \left\{ \left| \frac{\mathbf{Y}^\top (\mathbf{I} - \mathbf{P}_{\gamma \cup \xi_*}) \mathbf{Y}}{n\sigma_*^2} - 1 \right| \leq M_1 \epsilon_n \right\}, \\ \Theta_{1,\gamma} &= \{(\sigma, \xi, \beta) \in \Theta_1 : \xi = \gamma\}. \end{aligned}$$

Then

$$\phi_1 = \max_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_*| \leq Ks} \phi_{1,\gamma}, \quad \Theta_1 = \cup_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_*| \leq Ks} \Theta_{1,\gamma}.$$

Applying Lemma S.3.3 yields

$$\begin{aligned} \sup_{(\sigma, \xi, \beta) \in \Theta_1} \mathbb{E}_{(\sigma, \xi, \beta)} (1 - \phi_1) &\leq \max_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_*| \leq Ks} \sup_{(\sigma, \xi, \beta) \in \Theta_{1,\gamma}} \mathbb{E}_{(\sigma, \xi, \beta)} (1 - \phi_{1,\gamma}) \\ &= \sup_{(\sigma, \xi, \beta) \in \Theta_1} \mathbb{E}_{(\sigma, \xi, \beta)} (1 - \phi_{1,\xi}). \end{aligned}$$

We proceed to bound, for any $(\sigma, \xi, \beta) \in \Theta_1$,

$$\mathbb{E}_{(\sigma, \xi, \beta)} (1 - \phi_{1,\xi}) = \mathbb{P}_{(\sigma, \xi, \beta)} \left(\left| \frac{\mathbf{Y}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \mathbf{Y}}{n\sigma_*^2} - 1 \right| \leq M_1 \epsilon_n \right).$$

To this end, the restriction $\frac{\sigma^2}{\sigma_*^2} \notin \left[\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}, \frac{1+M_1\epsilon_n}{1-M_1\epsilon_n} \right]$ of $\Theta_1 \subseteq \Theta_0^c$ implies that

$$\left| \frac{\mathbf{Y}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \mathbf{Y}}{n\sigma_*^2} - 1 \right| \leq M_1 \epsilon_n \implies \left| \frac{\mathbf{Y}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \mathbf{Y}}{n\sigma^2} - 1 \right| > M_1 \epsilon_n.$$

Another restriction $\max_{j \notin \xi} |\beta_j| \leq \sigma z_{0n}$ of $\Theta_1 \subseteq \Theta_0^c$ implies that $\|\mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c}\| \leq \sigma \sqrt{np} z_{0n}$.

As we will show later, under $\mathbb{P}_{(\sigma, \xi, \beta)}$,

$$\left| \frac{\mathbf{Y}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \mathbf{Y}}{n\sigma^2} - 1 \right| > M_1 \epsilon_n \implies \left| \frac{\boldsymbol{\varepsilon}^\top (\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}) \boldsymbol{\varepsilon}}{n} - 1 \right| > M_1 \epsilon_n - 3p z_{0n}. \quad (\text{S.1})$$

Then

$$\begin{aligned} \mathbb{P}_{(\sigma, \xi, \beta)}(1 - \phi_{1, \xi}) &\leq \mathbb{P}\left(\left|\frac{\boldsymbol{\varepsilon}^T(\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*})\boldsymbol{\varepsilon}}{n} - 1\right| > (M_1\epsilon_n - 3pz_{0n})\right) \\ &\leq \mathbb{P}\left(\frac{\chi_{n-s}^2}{n} > 1 + (M_1\epsilon_n - 3pz_{0n})\right) + \mathbb{P}\left(\frac{\chi_{n-(K+1)s}^2}{n} < 1 - (M_1\epsilon_n - 3pz_{0n})\right). \end{aligned}$$

Note that this bound holds uniformly for all $(\sigma, \xi, \beta) \in \Theta_1$, and that Assumption 2.1(c) derives that $pz_{0n}/\epsilon_n \rightarrow 0$. Therefore, the probability bound of the chi-square distribution (Lemma S.3.2, part (a)) concludes the proof.

It is only left to show (S.1). For simplicity of notation, we write $\mathbf{R} = \mathbf{I} - \mathbf{P}_{\xi \cup \xi_*}$, $\mathbf{b} = \mathbf{X}_{\xi^c} \beta_{\xi^c} / \sigma \sqrt{n}$, then $\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_{\xi \cup \xi_*})\mathbf{Y} / n\sigma^2 = \|\mathbf{R}(\boldsymbol{\varepsilon} + \mathbf{b})\|^2$ under $\mathbb{P}_{(\sigma, \xi, \beta)}$. Since

$$\|\mathbf{R}\boldsymbol{\varepsilon}\|^2 - 2\|\mathbf{R}\boldsymbol{\varepsilon}\|\|\mathbf{b}\| \leq \|\mathbf{R}(\boldsymbol{\varepsilon} + \mathbf{b})\|^2 \leq (\|\mathbf{R}\boldsymbol{\varepsilon}\| + \|\mathbf{b}\|)^2,$$

we have for small $\|\mathbf{b}\|$,

$$\begin{aligned} \|\mathbf{R}(\boldsymbol{\varepsilon} + \mathbf{b})\|^2 > 1 + M_1\epsilon_n &\implies \|\mathbf{R}\boldsymbol{\varepsilon}\|^2 > \left(\sqrt{1 + M_1\epsilon_n} - \|\mathbf{b}\|\right)^2 \\ &\implies \|\mathbf{R}\boldsymbol{\varepsilon}\|^2 > 1 + M_1\epsilon_n - 3\|\mathbf{b}\| \end{aligned}$$

and

$$\begin{aligned} \|\mathbf{R}(\boldsymbol{\varepsilon} + \mathbf{b})\|^2 < 1 - M_1\epsilon_n &\implies \|\mathbf{R}\boldsymbol{\varepsilon}\|^2 < \left(\|\mathbf{b}\| + \sqrt{\|\mathbf{b}\|^2 + 1 - M_1\epsilon_n}\right)^2 \\ &\implies \|\mathbf{R}\boldsymbol{\varepsilon}\|^2 < 1 - M_1\epsilon_n + 3\|\mathbf{b}\|. \end{aligned}$$

□

Proof of Lemma 5.4, part (a). For any $\xi \in \mathcal{F}$ such that $|\xi \setminus \xi_*| \leq Ks$, write

$$\begin{aligned} \frac{\|\mathbf{X}_{\xi \cup \xi_*}^\dagger \mathbf{Y} - \beta_{\xi \cup \xi_*}^*\|^2}{\sigma_\star^2} &= \boldsymbol{\varepsilon}^T \mathbf{X}_{\xi \cup \xi_*} (\mathbf{X}_{\xi \cup \xi_*}^T \mathbf{X}_{\xi \cup \xi_*})^{-2} \mathbf{X}_{\xi \cup \xi_*}^T \boldsymbol{\varepsilon} \\ &\leq \frac{\boldsymbol{\varepsilon}^T \mathbf{X}_{\xi \cup \xi_*} (\mathbf{X}_{\xi \cup \xi_*}^T \mathbf{X}_{\xi \cup \xi_*})^{-1} \mathbf{X}_{\xi \cup \xi_*}^T \boldsymbol{\varepsilon}}{\lambda_{\min}(\mathbf{X}_{\xi \cup \xi_*}^T \mathbf{X}_{\xi \cup \xi_*})} \leq \frac{\boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon}}{n\lambda}, \end{aligned}$$

implying

$$\begin{aligned} \phi_2 &\leq 1 \left\{ \max_{\xi \in \mathcal{F}: |\xi \setminus \xi_*| \leq Ks} \boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon} > M_2^2 n \epsilon_n^2 / 4 \right\} \\ &\leq 1 \left\{ \max_{\xi: |\xi \setminus \xi_*| \leq Ks} \boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon} > M_2^2 n \epsilon_n^2 / 4 \right\}. \end{aligned}$$

Since projection matrices $\mathbf{P}_{\xi_1 \cup \xi_*} \leq \mathbf{P}_{\xi_2 \cup \xi_*}$ for nested models $\xi_1 \subseteq \xi_2$, the quantity $\boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon}$ achieves its maximum value at some ξ of size Ks .

$$\begin{aligned} \phi_2 &\leq 1 \left\{ \max_{\xi: |\xi| = Ks} \boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon} > M_2^2 n \epsilon_n^2 / 4 \right\} \\ &\leq \sum_{\xi: |\xi| = Ks} 1 \left\{ \boldsymbol{\varepsilon}^T \mathbf{P}_{\xi \cup \xi_*} \boldsymbol{\varepsilon} > M_2^2 n \epsilon_n^2 / 4 \right\}. \end{aligned}$$

For each ξ of size Ks , we note that $\text{rank}(\xi \cup \xi_\star) \leq (K+1)s$ and write

$$\mathbb{P}_\star(\boldsymbol{\varepsilon}^\top \mathbf{P}_{\xi \cup \xi_\star} \boldsymbol{\varepsilon} > M_2^2 n \epsilon_n^2 / 4) \leq \mathbb{P}\left(\chi_{(K+1)s}^2 > M_2^2 n \epsilon_n^2 / 4\right).$$

Putting the last two displays together with the probability bound of the chi-square distribution (Lemma S.3.2, part (b)) yields

$$\mathbb{E}_\star \phi_2 \leq \binom{p}{Ks} \times e^{-[M_2^2/8 - o(1)]n\epsilon_n^2} \leq e^{-[M_2^2/8 - K - o(1)]n\epsilon_n^2}.$$

□

Proof of Lemma 5.4, part(b). Define

$$\begin{aligned} \phi_{2,\gamma} &= 1 \left\{ \|\mathbf{X}_{\gamma \cup \xi_\star}^\dagger \mathbf{Y} - \boldsymbol{\beta}_{\gamma \cup \xi_\star}^\star\| > \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \right\}, \\ \Theta_{2,\gamma} &= \{(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_2 : \xi = \gamma\}, \end{aligned}$$

then

$$\begin{aligned} \phi_2 &= \max_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_\star| \leq Ks} \phi_{2,\gamma} \\ \Theta_2 &= \cup_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_\star| \leq Ks} \Theta_{2,\gamma}. \end{aligned}$$

Applying Lemma S.3.3 yields

$$\begin{aligned} \sup_{(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_2} \mathbb{E}_{(\sigma, \xi, \boldsymbol{\beta})}(1 - \phi_2) &\leq \max_{\gamma \in \mathcal{F}: |\gamma \setminus \xi_\star| \leq Ks} \mathbb{E}_{(\sigma, \xi, \boldsymbol{\beta})}(1 - \phi_{2,\gamma}) \\ &= \sup_{(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_2} \mathbb{E}_{(\sigma, \xi, \boldsymbol{\beta})}(1 - \phi_{2,\xi}). \end{aligned}$$

We proceed to bound, for any $(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_2$,

$$\mathbb{E}_{(\sigma, \xi, \boldsymbol{\beta})}(1 - \phi_{2,\xi}) = \mathbb{P}_{(\sigma, \xi, \boldsymbol{\beta})} \left(\|\mathbf{X}_{\xi \cup \xi_\star}^\dagger \mathbf{Y} - \boldsymbol{\beta}_{\xi \cup \xi_\star}^\star\| \leq \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \right).$$

Under $\mathbb{P}_{(\sigma, \xi, \boldsymbol{\beta})}$, restrictions $\frac{\sigma^2}{\sigma_\star^2} \notin \left[\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}, \frac{1+M_1\epsilon_n}{1-M_1\epsilon_n} \right]$ and $\max_{j \notin \xi} |\beta_j| \leq \sigma z_{0n}$ of $\Theta_1 \subseteq \Theta_0^c$ imply

$$\begin{aligned} \|\mathbf{X}_{\xi \cup \xi_\star}^\dagger \mathbf{Y} - \boldsymbol{\beta}_{\xi \cup \xi_\star}^\star\| &\leq \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \\ \implies \|\boldsymbol{\beta}_{\xi \cup \xi_\star} - \boldsymbol{\beta}_{\xi \cup \xi_\star}^\star + \mathbf{X}_{\xi \cup \xi_\star}^\dagger (\mathbf{X}_{\xi^c \cap \xi_\star^c} \boldsymbol{\beta}_{\xi^c \cap \xi_\star^c} + \sigma \boldsymbol{\varepsilon})\| &\leq \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \\ \implies \|\mathbf{X}_{\xi \cup \xi_\star}^\dagger (\mathbf{X}_{\xi^c \cap \xi_\star^c} \boldsymbol{\beta}_{\xi^c \cap \xi_\star^c} + \sigma \boldsymbol{\varepsilon})\| &\geq \|\boldsymbol{\beta} - \boldsymbol{\beta}^\star\| - \|\boldsymbol{\beta}_{\xi^c}\| - \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \\ \implies \frac{1}{\sqrt{n\lambda}} \|\mathbf{P}_{\xi \cup \xi_\star} (\mathbf{X}_{\xi^c \cap \xi_\star^c} \boldsymbol{\beta}_{\xi^c \cap \xi_\star^c} + \sigma \boldsymbol{\varepsilon})\| &\geq \|\boldsymbol{\beta} - \boldsymbol{\beta}^\star\| - \|\boldsymbol{\beta}_{\xi^c}\| - \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \\ \implies \frac{1}{\sqrt{n\lambda}} (\sqrt{np} \|\boldsymbol{\beta}_{\xi^c}\| + \sigma \|\mathbf{P}_{\xi \cup \xi_\star} \boldsymbol{\varepsilon}\|) &\geq \|\boldsymbol{\beta} - \boldsymbol{\beta}^\star\| - \|\boldsymbol{\beta}_{\xi^c}\| - \frac{M_2 \sigma_\star \epsilon_n}{2\sqrt{\lambda}} \\ \implies \|\mathbf{P}_{\xi \cup \xi_\star} \boldsymbol{\varepsilon}\| &\geq \left(\frac{M_2}{2} \sqrt{\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}} - \frac{pz_{0n}}{\epsilon_n} - \frac{pz_{0n}\sqrt{\lambda}}{\epsilon_n\sqrt{p}} \right) \sqrt{n}\epsilon_n \\ \implies \|\mathbf{P}_{\xi \cup \xi_\star} \boldsymbol{\varepsilon}\| &\geq \left(\frac{M_2}{2} \sqrt{\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}} - \frac{2pz_{0n}}{\epsilon_n} \right) \sqrt{n}\epsilon_n. \end{aligned}$$

Thus,

$$\mathbb{P}_{(\sigma, \xi, \beta)}(1 - \phi_{2, \xi}) \leq \mathbb{P} \left(\chi_{(K+1)s}^2 \geq \left(\frac{M_2}{2} \sqrt{\frac{1 - M_1 \epsilon_n}{1 + M_1 \epsilon_n}} - \frac{2pz_{0n}}{\epsilon_n} \right)^2 n \epsilon_n^2 \right).$$

Note that this bound holds uniformly for all $(\sigma, \xi, \beta) \in \Theta_2$. Assumption 2.1(c) deriving that $pz_{0n}/\epsilon_n \rightarrow 0$ and the probability bound of the chi-square distribution (Lemma S.3.2, part (b)) conclude the proof. \square

Proof of Lemma 5.5. The proof consists of four steps.

(i) Let

$$\Theta_\star = \Theta_\star(\eta_1, \eta_2) := \left\{ (\sigma, \xi, \beta) : \begin{array}{l} \sigma^2/\sigma_\star^2 \in [1, 1 + \eta_1 \epsilon_n^2], \\ \xi = \xi_\star, \\ |\beta_j| \leq \sigma z_{0n}, j \notin \xi_\star, \\ |\beta_j - \beta_j^\star| \leq \eta_2 \sigma \epsilon_n / s, j \in \xi_\star \end{array} \right\}.$$

It is obvious that

$$\int \frac{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I})}{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta_\star, \sigma_\star^2 \mathbf{I})} d\Pi \geq \Pi(\Theta_\star) \inf_{(\sigma, \xi, \beta) \in \Theta_\star} \frac{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I})}{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta_\star, \sigma_\star^2 \mathbf{I})},$$

(ii) Prove that if $\eta_2 < 1$ then

$$\Pi(\Theta_\star) \gtrsim e^{-[A_1 + A_3 + 1 - o(1)]s \log p / 2}.$$

(iii) Prove the contrapositive of the implication statement

$$\begin{aligned} \inf_{(\sigma, \xi, \beta) \in \Theta_\star} \frac{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I})}{\mathcal{N}(\mathbf{Y}|\mathbf{X}\beta_\star, \sigma_\star^2 \mathbf{I})} &\leq e^{-[\eta_1 + \eta_2^2/2 + \sqrt{C}\eta_2 + \eta_3 + o(1)]s \log p} \\ \implies \|\mathbf{P}_{\xi_\star} \boldsymbol{\varepsilon}\|^2 &\geq Cn\epsilon_n^2 \text{ or } \|\boldsymbol{\varepsilon}\|^2 \geq 4n, \end{aligned}$$

for any small $\eta_3 > 0$.

(iv) Prove

$$\mathbb{P}_\star \left(\|\mathbf{X}_{\xi_\star} \boldsymbol{\varepsilon}\|^2 \geq Cn\epsilon_n^2 \text{ or } \|\boldsymbol{\varepsilon}\|^2 \geq 4n \right) \lesssim e^{-[C/2 - o(1)]n\epsilon_n^2}.$$

Setting sufficiently small η_1, η_2, η_3 and suitable C such that

$$\eta > \eta_1 + \eta_2^2/2 + \sqrt{C}\eta_2 + \eta_3, \quad C_\eta > C/2$$

completes the proof.

Proof of (ii):

$$\begin{aligned} \Pi(\Theta_\star) &= \int_{\sigma_\star^2}^{(1+\eta_1 \epsilon_n^2)\sigma_\star^2} g(\sigma^2) d\sigma^2 \times \pi(\xi_\star) \\ &\times \Pi \left(\max_{j \notin \xi_\star} |\beta_j| \leq \sigma z_{0n} \mid \xi = \xi_\star, \sigma \right) \\ &\times \prod_{j \in \xi_\star} \int_{\beta_j^\star - \eta_2 \sigma \epsilon_n / s}^{\beta_j^\star + \eta_2 \sigma \epsilon_n / s} h_1 \left(\frac{\beta_j}{\sigma} \right) d \left(\frac{\beta_j}{\sigma} \right). \end{aligned}$$

The first term, due to Assumption 2.1(a), is bounded from below as

$$\int_{\sigma_*^2}^{\sigma_*^2(1+\eta_1\epsilon_n^2)} g(\sigma^2) d\sigma^2 \geq \eta_1 \sigma_*^2 \epsilon_n^2 g(\sigma_*^2)/2 \gtrsim e^{-cs \log p},$$

for any small $c > 0$. The second term, due to Assumption 2.1(b), is bounded from below as

$$\pi(\xi_*) \succeq p^{-A_1 s}.$$

The third term, due to Assumption 2.1(c), is bounded from below as

$$\Pi \left(\max_{j \in \xi^c} |\beta_j| \leq \sigma z_{0n} \middle| \xi = \xi_*, \sigma \right) \geq 1 - e^{-n + \log p}.$$

The fourth term, due to Assumption 2.1(d), is bounded from below as

$$\begin{aligned} & \prod_{j \in \xi_*} \int_{\beta_j^*/\sigma - \eta_2 \epsilon_n/s}^{\beta_j^*/\sigma + \eta_2 \epsilon_n/s} h_1(z) dz \\ & \geq \left(2\eta_2 \sqrt{\frac{\log p}{ns}} \times \inf \left\{ h_1(z) : |z| \leq \max_{j \in \xi_*} |\beta_j^*/\sigma| + \eta_2 \epsilon_n/s \right\} \right)^s \\ & \geq \left(2\eta_2 \sqrt{\frac{\log p}{ns}} \times \inf \{ h_1(z) : |z| \leq z_{1n} \} \right)^s \gtrsim p^{-(A_3+1)s}. \end{aligned}$$

Proof of (iii): we are going to show that, given $\|\mathbf{P}_{\xi_*} \boldsymbol{\varepsilon}\|^2 < Cn\epsilon_n^2$ and $\|\boldsymbol{\varepsilon}\|^2 < 4n$, the density ratio

$$\Lambda = \frac{\mathcal{N}(\mathbf{Y}|\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{I})}{\mathcal{N}(\mathbf{Y}|\mathbf{X}\boldsymbol{\beta}_*, \sigma_*^2 \mathbf{I})} \gtrsim e^{-(\eta_1 + \eta_2^2/2 + \sqrt{C}\eta_2 + \eta_3)s \log p}$$

for any $(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_*$ and any small constant $\eta_3 > 0$. Write

$$\begin{aligned} -2 \log \Lambda &= \|\sigma_* \boldsymbol{\varepsilon} + \mathbf{X}(\boldsymbol{\beta}^* - \boldsymbol{\beta})\|^2 / \sigma^2 - \|\boldsymbol{\varepsilon}\|^2 + 2n \log(\sigma^2 / \sigma_*^2) \\ &= (\sigma_*^2 / \sigma^2 - 1) \|\boldsymbol{\varepsilon}\|^2 + \|\mathbf{X}(\boldsymbol{\beta}^* - \boldsymbol{\beta}) / \sigma\|^2 + 2\sigma_* \boldsymbol{\varepsilon}^T \mathbf{X}_{\xi_*} (\boldsymbol{\beta}_{\xi_*} - \boldsymbol{\beta}) / \sigma^2 \\ &\quad + 2\sigma_* \boldsymbol{\varepsilon}^T \mathbf{X}_{\xi_*^c} \boldsymbol{\beta}_{\xi_*^c} / \sigma^2 + 2n \log(\sigma^2 / \sigma_*^2) \\ &\leq \|\mathbf{X}(\boldsymbol{\beta}^* - \boldsymbol{\beta}) / \sigma\|^2 + 2\|\mathbf{X}_{\xi_*}^T \boldsymbol{\varepsilon}\| \|(\boldsymbol{\beta}_{\xi_*} - \boldsymbol{\beta}_{\xi_*}) / \sigma\| \\ &\quad + 2\|\boldsymbol{\varepsilon}\| \|\mathbf{X}_{\xi_*^c} \boldsymbol{\beta}_{\xi_*^c} / \sigma\| + 2n \log(\sigma^2 / \sigma_*^2). \end{aligned}$$

where the definition of Θ_* enforces restrictions that

$$\begin{aligned} \log(\sigma^2 / \sigma_*^2) &\leq \eta_1 \epsilon_n^2, \quad \|\boldsymbol{\beta}_{\xi_*}^* - \boldsymbol{\beta}_{\xi_*}\| \leq \eta_2 \sigma \epsilon_n / \sqrt{s}, \quad \|\mathbf{X}_{\xi_*^c} \boldsymbol{\beta}_{\xi_*^c}\| \leq \sigma \sqrt{np} z_{0n}, \\ \|\mathbf{X}(\boldsymbol{\beta}^* - \boldsymbol{\beta})\| &\leq \|\mathbf{X}_{\xi_*^c} \boldsymbol{\beta}_{\xi_*^c}\| + \|\mathbf{X}_{\xi_*} (\boldsymbol{\beta}_{\xi_*}^* - \boldsymbol{\beta}_{\xi_*})\| \leq \sigma \sqrt{np} z_{0n} + \sqrt{ns} \times \frac{\eta_2 \sigma \epsilon_n}{\sqrt{s}}, \end{aligned}$$

and the premise derives that $\|\boldsymbol{\varepsilon}\| < 2\sqrt{n}$ and that

$$\|\mathbf{X}_{\xi_*}^T \boldsymbol{\varepsilon}\|^2 \leq \lambda_{\max}(\mathbf{X}_{\xi_*}^T \mathbf{X}_{\xi_*}) \|\mathbf{P}_{\xi_*} \boldsymbol{\varepsilon}\|^2 < ns \times Cn\epsilon_n^2.$$

Collecting these pieces together yields the desired lower bound for the density ratio.

Proof of (iv): it follows from the facts that $\|\mathbf{P}_{\xi_*} \boldsymbol{\varepsilon}\|^2 \sim \chi_s^2$, $\|\boldsymbol{\varepsilon}\|^2 \sim \chi_n^2$ and the probability bounds of the chi-squared distribution (Lemma S.3.2). \square

Proof of Lemma 5.6, part (a). We first show that

$$\min_{\xi \not\supseteq \xi_*: |\xi \setminus \xi_*| \leq Ks} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^*\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n. \quad (\text{S.2})$$

Indeed, for any $\xi \not\supseteq \xi_*$ with set difference $|\xi \setminus \xi_*| \leq Ks$,

$$\begin{aligned} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^*\|^2 &= \|(\mathbf{I} - \mathbf{P}_\xi) \mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^*\|^2 \\ &= \|(\mathbf{I} - \mathbf{P}_\xi) \mathbf{X}_{\xi_* \setminus \xi} \boldsymbol{\beta}_{\xi_* \setminus \xi}^*\|^2 \\ &= \boldsymbol{\beta}_{\xi_* \setminus \xi}^T \mathbf{X}_{\xi_* \setminus \xi}^T (\mathbf{I} - \mathbf{P}_\xi) \mathbf{X}_{\xi_* \setminus \xi} \boldsymbol{\beta}_{\xi_* \setminus \xi} \end{aligned}$$

Note that $\mathbf{X}_{\xi_* \setminus \xi}^T (\mathbf{I} - \mathbf{P}_\xi) \mathbf{X}_{\xi_* \setminus \xi}$ is the Schur complement of the principal submatrix $\mathbf{X}_{\xi_* \setminus \xi}^T \mathbf{X}_{\xi_* \setminus \xi}$ in the matrix $\mathbf{X}_{\xi \cup \xi_*}^T \mathbf{X}_{\xi \cup \xi_*}$. Thus, by Lemma S.3.4,

$$\lambda_{\min} \left(\mathbf{X}_{\xi_* \setminus \xi}^T (\mathbf{I} - \mathbf{P}_\xi) \mathbf{X}_{\xi_* \setminus \xi} \right) \geq \lambda_{\min} \left(\mathbf{X}_{\xi \cup \xi_*}^T \mathbf{X}_{\xi \cup \xi_*} \right) \geq n\lambda.$$

It further leads to (S.2) as

$$\|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^*\|^2 \geq n\lambda \|\boldsymbol{\beta}_{\xi_* \setminus \xi}^*\|^2 \geq n\lambda |\xi_* \setminus \xi| \min_{j \in \xi_*} |\beta_j^*|^2 \geq M_3^2 \sigma_*^2 n \epsilon_n^2.$$

From the fact that $\mathbf{P}_{\xi \cup \xi_*} \leq \mathbf{P}_\xi + \mathbf{P}_{\xi_*}$, it follows that

$$\begin{aligned} \phi_3 &= 1 \left\{ \min_{\xi \not\supseteq \xi_*: |\xi \setminus \xi_*| \leq Ks} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) (\mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^* + \sigma_* \boldsymbol{\epsilon})\| < M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \right\} \\ &\leq 1 \left\{ \min_{\xi \not\supseteq \xi_*: |\xi \setminus \xi_*| \leq Ks} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{X}_{\xi_*} \boldsymbol{\beta}_{\xi_*}^*\| \right. \\ &\quad \left. - \max_{\xi \not\supseteq \xi_*: |\xi \setminus \xi_*| \leq Ks} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \sigma_* \boldsymbol{\epsilon}\| < M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \right\} \\ &\leq 1 \left\{ \max_{\xi \not\supseteq \xi_*: |\xi \setminus \xi_*| \leq Ks} \|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \sigma_* \boldsymbol{\epsilon}\| > M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \right\} \\ &\leq 1 \left\{ \|\mathbf{P}_{\xi_*} \boldsymbol{\epsilon}\| > M_3 \sqrt{n} \epsilon_n / 2 \right\}. \end{aligned}$$

Putting it together with the tail probability bound of the chi-square distribution (Lemma S.3.2) yields

$$\mathbb{E}_* \phi_3 \leq \mathbb{P} \left(\chi_s^2 > M_3 n \epsilon_n^2 / 4 \right) \leq e^{-[M_3^2/8 - o(1)] n \epsilon_n^2}.$$

□

Proof of Lemma 5.6, part(b). Define

$$\begin{aligned} \phi_{3,\gamma} &= 1 \left\{ \|(\mathbf{P}_{\gamma \cup \xi_*} - \mathbf{P}_\gamma) \mathbf{Y}\| < M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \right\}, \\ \Theta_{3,\gamma} &= \{(\sigma, \xi, \boldsymbol{\beta}) \in \Theta_3 : \xi = \gamma\}. \end{aligned}$$

Then

$$\phi_3 = \max_{\gamma \not\supseteq \xi_*: |\gamma \setminus \xi_*| \leq Ks} \phi_{3,\gamma}, \quad \Theta_3 = \cup_{\gamma \not\supseteq \xi_*: |\gamma \setminus \xi_*| \leq Ks} \Theta_{3,\gamma}.$$

Applying Lemma S.3.3 yields

$$\begin{aligned} \sup_{(\sigma, \xi, \beta) \in \Theta_3} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_3) &\leq \max_{\gamma \notin \xi_* : |\gamma \setminus \xi_*| \leq Ks} \sup_{(\sigma, \xi, \beta) \in \Theta_{3, \gamma}} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_{3, \gamma}) \\ &= \sup_{(\sigma, \xi, \beta) \in \Theta_3} \mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_{3, \xi}). \end{aligned}$$

We proceed to bound, for any $(\sigma, \xi, \beta) \in \Theta_3$,

$$\mathbb{E}_{(\sigma, \xi, \beta)}(1 - \phi_{3, \xi}) = \mathbb{P}_{(\sigma, \xi, \beta)}(\|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{Y}\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n / 2).$$

Under $\mathbb{P}_{(\sigma, \xi, \beta)}$, restrictions $\frac{\sigma^2}{\sigma_*^2} \notin \left[\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}, \frac{1+M_1\epsilon_n}{1-M_1\epsilon_n} \right]$ and $\max_{j \notin \xi} |\beta_j| \leq \sigma z_{0n}$ of $\Theta_1 \subseteq \Theta_0^c$ and the fact that $\mathbf{P}_{\xi \cup \xi_*} \leq \mathbf{P}_\xi + \mathbf{P}_{\xi_*}$ imply that

$$\begin{aligned} &\|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) \mathbf{Y}\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \\ \implies &\|(\mathbf{P}_{\xi \cup \xi_*} - \mathbf{P}_\xi) (\mathbf{X}_{\xi^c} \beta_{\xi^c} + \sigma \boldsymbol{\epsilon})\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \\ \implies &\|\mathbf{X}_{\xi^c} \beta_{\xi^c}\| + \sigma \|\mathbf{P}_{\xi_*} \boldsymbol{\epsilon}\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \\ \implies &\sigma \epsilon_n + \sigma \|\mathbf{P}_{\xi_*} \boldsymbol{\epsilon}\| \geq M_3 \sigma_* \sqrt{n} \epsilon_n / 2 \\ \implies &\|\mathbf{P}_{\xi_*} \boldsymbol{\epsilon}\| \geq \left(\frac{M}{2} \sqrt{\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}} - \frac{pz_{0n}}{\epsilon_n} \right) \sqrt{n} \epsilon_n. \end{aligned}$$

Thus,

$$\mathbb{P}_{(\sigma, \xi, \beta)}(1 - \phi_{3, \xi}) \leq \mathbb{P} \left(\chi_s^2 \geq \left(\frac{M_3}{2} \sqrt{\frac{1-M_1\epsilon_n}{1+M_1\epsilon_n}} - \frac{pz_{0n}}{\epsilon_n} \right)^2 n \epsilon_n^2 \right).$$

Note that this bound holds uniformly for all $(\sigma, \xi, \beta) \in \Theta_3$ and that Assumption 2.1(c) derives that $p z_{0n} / \epsilon_n \rightarrow 0$. Therefore, the probability bound of the chi-square distribution (Lemma S.3.2, part (b)) concludes the proof. \square

Proof of Lemma 5.7. First, for any full-rank overfitted model ξ of size $t \leq (K+1)s$,

$$\begin{aligned} \mathbb{P}(\|(\mathbf{P}_\xi - \mathbf{P}_{\xi_*}) \boldsymbol{\epsilon}\|^2 \geq 2(1+\eta)(t-s) \log p) &\leq \mathbb{P}(\chi_{t-s}^2 \geq 2(1+\eta)(t-s) \log p) \\ &\lesssim p^{-(1+\eta/2)(t-s)}, \end{aligned}$$

due to the probability bound of the chi-squared distribution (Lemma S.3.2(b)). It follows that

$$\mathbb{P}(\Omega_1(\eta)) \leq \sum_{t=s+1}^{(K+1)s} p^{t-s} \times p^{-(1+\eta/2)(t-s)} \leq \sum_{j=1}^{Ks} p^{-\eta j/2} \asymp p^{-\eta/2}.$$

On the other hand, due to Lemma S.3.2(a), $\mathbb{P}(\Omega_2) \leq e^{-3n/8}$. Collecting these two pieces together completes the proof. \square

Proof of Lemma 5.8. For simplicity of notation, define

$$\begin{aligned}\mathcal{A} &:= \left[\sigma_\star^2 \times \frac{1 - M_1 \epsilon_n}{1 + M_1 \epsilon_n}, \sigma_\star^2 \times \frac{1 + M_1 \epsilon_n}{1 - M_1 \epsilon_n} \right], \\ \mathcal{B} &:= [-\sigma z_{0n}, +\sigma z_{0n}], \\ \mathcal{C} &:= \{\mathbf{v} \in \mathbb{R}^s : \|\mathbf{v} - \beta_{\xi_\star}^\star\| \leq M_2 \sigma_\star \epsilon_n\}, \\ \mathcal{C}' &:= \left\{ \mathbf{v} \in \mathbb{R}^s : \|\mathbf{v} - \beta_{\xi_\star}^\star\| \leq \sqrt{M_2^2 \sigma_\star^2 \epsilon_n^2 - (t-s)\sigma^2 z_{0n}^2} \right\} \subseteq \mathcal{C}.\end{aligned}$$

Write

$$\begin{aligned}\frac{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi)}{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \xi_\star)} &= \frac{\int_{\mathcal{A}} \int_{\mathcal{B}^{p-t}} \Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \sigma, \xi, \beta_{\xi^c}) h_0(\beta_{\xi^c}/\sigma) d(\beta_{\xi^c}/\sigma) g(\sigma^2) d\sigma^2}{\int_{\mathcal{A}} \int_{\mathcal{B}^{p-t}} \Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \sigma, \xi_\star, \beta_{\xi^c}) h_0(\beta_{\xi^c}/\sigma) d(\beta_{\xi^c}/\sigma) g(\sigma^2) d\sigma^2} \\ &\leq \sup_{\sigma^2 \in \mathcal{A}, \beta_{\xi^c} \in \mathcal{B}^{p-t}} \frac{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \sigma, \xi, \beta_{\xi^c})}{\Pi(\tilde{\Theta}|\mathbf{X}, \mathbf{Y}, \sigma, \xi_\star, \beta_{\xi^c})}.\end{aligned}\tag{S.3}$$

For any $\sigma^2 \in \mathcal{A}$, $\beta_{\xi^c} \in \mathcal{B}^{p-t}$, the numerator of (S.3) is bounded from above as

$$\begin{aligned}&= \int_{\beta_{\xi^c} : \|\beta_{\xi^c} - \beta_{\xi_\star}^\star\| \leq M_2 \sigma_\star \epsilon_n} \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) h_1(\beta_{\xi^c}/\sigma) d(\beta_{\xi^c}/\sigma) \\ &\leq \left[\sup_{\beta_{\xi^c} : \|\beta_{\xi^c} - \beta_{\xi_\star}^\star\| \leq M_2 \sigma_\star \epsilon_n} h_1(\beta_{\xi^c}/\sigma) \right] \int \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) d(\beta_{\xi^c}/\sigma) \\ &\leq \sup_{\mathbf{z} \in \mathcal{C}/\sigma} h_1(\mathbf{z}) \times \left[\sup_{\mathbf{z}} h_1(\mathbf{z}) \right]^{t-s} \times \int \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) d(\beta_{\xi^c}/\sigma).\end{aligned}$$

On the other hand, the denominator of (S.3) is bounded from below as

$$\begin{aligned}&\geq \int_{\beta_{\xi^c} : \beta_{\xi^c} \setminus \xi_\star \in \mathcal{B}^{t-s}, \beta_{\xi_\star} \in \mathcal{C}'} \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) h_1(\beta_{\xi^c}/\sigma) h_0(\beta_{\xi^c \setminus \xi_\star}/\sigma) d(\beta_{\xi^c}/\sigma) \\ &\geq \inf_{\beta_{\xi^c \setminus \xi_\star} \in \mathcal{B}^{t-s}} \int_{\mathcal{C}'} \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) h_1(\beta_{\xi^c}/\sigma) d(\beta_{\xi^c}/\sigma) \\ &\quad \times \int_{\mathcal{B}^{t-s}} h_0(\beta_{\xi^c \setminus \xi_\star}/\sigma) d(\beta_{\xi^c \setminus \xi_\star}/\sigma) \\ &\geq \inf_{\beta_{\xi^c \setminus \xi_\star} \in \mathcal{B}^{t-s}} \left[\inf_{\beta_{\xi_\star} \in \mathcal{C}'} h_1(\beta_{\xi_\star}/\sigma) \times \int \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) d(\beta_{\xi^c}/\sigma) \right] \\ &\quad \times \int_{\mathcal{B}^{t-s}} h_0(\beta_{\xi^c \setminus \xi_\star}/\sigma) d(\beta_{\xi^c \setminus \xi_\star}/\sigma) \\ &\geq \inf_{\mathbf{z} \in \mathcal{C}/\sigma} h_1(\mathbf{z}) \times \left[\int_{\mathcal{B}/\sigma} h_0(\mathbf{z}) d\mathbf{z} \right]^{t-s} \times \inf_{\beta_{\xi^c \setminus \xi_\star} \in \mathcal{B}^{t-s}} \int \mathcal{N}(\mathbf{Y}|\mathbf{X}\beta, \sigma^2 \mathbf{I}) d(\beta_{\xi^c}/\sigma).\end{aligned}$$

Comparison of the last two displays reveals that (S.3) is upper bounded by the product of three terms, which are bounded as follows.

(i) Due to condition (a) of Theorem 4.3,

$$\begin{aligned}\frac{\sup_{\mathbf{z} \in \mathcal{C}/\sigma} h_1(\mathbf{z})}{\inf_{\mathbf{z} \in \mathcal{C}/\sigma} h_1(\mathbf{z})} &\leq \frac{\sup_{\mathbf{z} : \|\mathbf{z} - \beta_{\xi_\star}^\star/\sigma\|_1 \leq M_2 \sqrt{s} \sigma_\star \epsilon_n/\sigma} h_1(\mathbf{z})}{\inf_{\mathbf{z} : \|\mathbf{z} - \beta_{\xi_\star}^\star/\sigma\|_1 \leq M_2 \sqrt{s} \sigma_\star \epsilon_n/\sigma} h_1(\mathbf{z})} \\ &\leq e^{L_n \times 2M_2 \sqrt{s} \sigma_\star \epsilon_n/\sigma} \lesssim e^{\eta \log p/2}.\end{aligned}$$

(ii) Due to Assumption 2.1(c),

$$\left(\frac{\sup_z h_1(z)}{\int_{\mathcal{B}/\sigma} h_0(z) dz} \right)^{t-s} \leq \frac{\sup_z h_1(z)}{1 - (t-s)e^{-n}} \leq 2 \left(\sup_z h_1(z) \right)^{t-s},$$

for sufficiently large n such that $n \geq \log(2Ks)$.

(iii) Some elementary calculus gives that

$$\frac{\int \mathcal{N}(\mathbf{Y}|\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{I}) d(\boldsymbol{\beta}_\xi/\sigma)}{\int \mathcal{N}(\mathbf{Y}|\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{I}) d(\boldsymbol{\beta}_{\xi_*}/\sigma)} = \frac{\det \left| 2\pi(\mathbf{X}_\xi^\top \mathbf{X}_\xi)^{-1} \right|^{1/2}}{\det \left| 2\pi(\mathbf{X}_{\xi_*}^\top \mathbf{X}_{\xi_*})^{-1} \right|^{1/2}} \times \exp \left\{ \frac{R_{\xi_*}^2 - R_\xi^2}{2\sigma^2} \right\},$$

with $R_\gamma^2 = \|(\mathbf{I} - \mathbf{P}_\gamma)(\sigma_* \boldsymbol{\varepsilon} - \mathbf{X}_{\gamma^c} \boldsymbol{\beta}_{\gamma^c})\|^2$. By Lemma S.3.4 and Assumption 3.2,

$$\begin{aligned} \frac{\det \left| \mathbf{X}_\xi^\top \mathbf{X}_\xi \right|}{\det \left| \mathbf{X}_{\xi_*}^\top \mathbf{X}_{\xi_*} \right|} &= \det \left| \mathbf{X}_{\xi \setminus \xi_*}^\top (\mathbf{I} - \mathbf{P}_{\xi_*}) \mathbf{X}_{\xi \setminus \xi_*} \right| \\ &\geq \left[\lambda_{\min} \left(\mathbf{X}_{\xi \setminus \xi_*}^\top (\mathbf{I} - \mathbf{P}_{\xi_*}) \mathbf{X}_{\xi \setminus \xi_*} \right) \right]^{t-s} \\ &\geq \left[\lambda_{\min} \left(\mathbf{X}_\xi^\top \mathbf{X}_\xi \right) \right]^{t-s} \geq (n\lambda)^{t-s}. \end{aligned}$$

Conditionally on $\Omega(\eta)$ defined in Lemma 5.7,

$$\begin{aligned} R_{\xi_*}^2 - R_\xi^2 &= \|(\mathbf{P}_\xi - \mathbf{P}_{\xi_*})(\sigma_* \boldsymbol{\varepsilon} - \mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c})\|^2 + \|(\mathbf{I} - \mathbf{P}_{\xi_*}) \mathbf{X}_{\xi \setminus \xi_*} \boldsymbol{\beta}_{\xi \setminus \xi_*}\|^2 \\ &\quad - 2(\sigma_* \boldsymbol{\varepsilon} - \mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c})^\top (\mathbf{I} - \mathbf{P}_{\xi_*}) \mathbf{X}_{\xi \setminus \xi_*} \boldsymbol{\beta}_{\xi \setminus \xi_*} \\ &\leq [\sigma_* \|(\mathbf{P}_\xi - \mathbf{P}_{\xi_*}) \boldsymbol{\varepsilon}\| + \|\mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c}\|]^2 + \|\mathbf{X}_{\xi \setminus \xi_*} \boldsymbol{\beta}_{\xi \setminus \xi_*}\|^2 \\ &\quad + 2\sigma_* \|\boldsymbol{\varepsilon}\| \|\mathbf{X}_{\xi \setminus \xi_*} \boldsymbol{\beta}_{\xi \setminus \xi_*}\| + 2\|\mathbf{X}_{\xi^c} \boldsymbol{\beta}_{\xi^c}\| \|\mathbf{X}_{\xi \setminus \xi_*} \boldsymbol{\beta}_{\xi \setminus \xi_*}\| \\ &\leq [\sigma_* \sqrt{(2+\eta)(t-s) \log p} + \sigma \sqrt{n} p z_{0n}]^2 + [\sigma \sqrt{n}(t-s) z_{0n}]^2 \\ &\quad + 4\sigma_* \sqrt{n} \times \sigma \sqrt{n}(t-s) z_{0n} + 2\sigma \sqrt{n} p z_{0n} \times \sigma \sqrt{n}(t-s) z_{0n} \\ &\leq (2 + 3\eta/2)(t-s) \log p, \end{aligned}$$

for sufficiently large $n \geq N$, with N not depending on ξ .

Combining (i)-(iii) with (S.3) concludes the proof. \square

S.3 Preliminary Lemmas

Lemma S.3.1 (Pelekis (2016, Theorem 1.1)). For a Binomial distributed random variable $T \sim \text{Binomial}(p, \mu)$, if $p\mu < t \leq p-1$ then

$$\mathbb{P}(T \geq t) \leq \frac{\mu^{2(\tilde{t}+1)}}{2} \binom{p}{\tilde{t}+1} \bigg/ \binom{t}{\tilde{t}+1},$$

where $\tilde{t} = \lfloor (t - p\mu)/(1 - \mu) \rfloor < t$.

Lemma S.3.2 (Probability Bounds of Chi-squared Distribution). Let χ_d^2 be a chi-squared random variable of degree d .

(a) For any ϵ_n such that $n\epsilon_n > d_n$,

$$\begin{aligned}\mathbb{P}(\chi_{n-d_n}^2/n \geq 1 + \epsilon_n) &\leq e^{-\min\left\{\frac{(n\epsilon_n + d_n)^2}{8(n-d_n)}, \frac{n\epsilon_n + d_n}{8}\right\}}, \\ \mathbb{P}(\chi_{n-d_n}^2/n \leq 1 - \epsilon_n) &\leq e^{-\min\left\{\frac{(n\epsilon_n - d_n)^2}{8(n-d_n)}, \frac{n\epsilon_n - d_n}{8}\right\}}.\end{aligned}$$

In addition, if $\epsilon_n \rightarrow 0$ but $n\epsilon_n \succ d_n$,

$$\begin{aligned}\mathbb{P}(\chi_{n-d_n}^2/n \geq 1 + \epsilon_n) &\preceq e^{-(1/8-o(1))n\epsilon_n^2}, \\ \mathbb{P}(\chi_{n-d_n}^2/n \leq 1 - \epsilon_n) &\preceq e^{-(1/8-o(1))n\epsilon_n^2}.\end{aligned}$$

(b)

$$\mathbb{P}(\chi_{d_n}^2 \geq t_n) \leq e^{-(\sqrt{2t_n - d_n} - \sqrt{d_n})^2/4}.$$

In addition, if $t_n \succ d_n$ then for any \tilde{t}_n such that $\tilde{t}_n/t_n \rightarrow 1$

$$\mathbb{P}(\chi_{d_n}^2 \geq t_n) \preceq e^{-[1/2-o(1)]\tilde{t}_n}.$$

Proof. For part (a), the first assertion follows from the sub-exponential tail of chi-squared distribution, and the second assertion is due to

$$\begin{aligned}[1/8 - o(1)]n\epsilon_n^2 &\preceq \frac{(n\epsilon_n + d_n)^2}{8(n-d_n)} \preceq \frac{n\epsilon_n + d_n}{8}, \\ [1/8 - o(1)]n\epsilon_n^2 &\preceq \frac{(n\epsilon_n - d_n)^2}{8(n-d_n)} \preceq \frac{n\epsilon_n - d_n}{8}.\end{aligned}$$

For part (b), the first assertion is a corollary of [Laurent and Massart \(2000, Lemma 1\)](#), and the second assertion follows from

$$[1/2 - o(1)]\tilde{t}_n \preceq \left(\sqrt{2t_n - d_n} - \sqrt{d_n}\right)^2/4.$$

□

Lemma S.3.3. For a collection of subspace $\{\Theta_j\}_{j=1}^m$ and a collection of test functions $\{\varphi_j\}_{j=1}^m$

$$\sup_{\theta \in \cup_{j=1}^m \Theta_j} \mathbb{E}_\theta \left(1 - \max_{j=1}^m \varphi_j\right) \leq \max_{j=1}^m \left\{ \sup_{\theta \in \Theta_j} \mathbb{E}_\theta (1 - \varphi_j) \right\}.$$

Proof.

$$\begin{aligned}\sup_{\theta \in \cup_{j=1}^m \Theta_j} \mathbb{E}_\theta \left(1 - \max_{j=1}^m \varphi_j\right) &= \max_{j=1}^m \left\{ \sup_{\theta \in \Theta_j} \mathbb{E}_\theta \left(1 - \max_{k=1}^m \varphi_k\right) \right\} \\ &= \max_{j=1}^m \left\{ \sup_{\theta \in \Theta_j} \mathbb{E}_\theta \left(\min_{k=1}^m (1 - \varphi_k)\right) \right\} \\ &\leq \max_{j=1}^m \left\{ \sup_{\theta \in \Theta_j} \mathbb{E}_\theta (1 - \varphi_j) \right\}.\end{aligned}$$

□

Lemma S.3.4 ([Liu \(2005, Part of Corollary 2.4\)](#)). Let

$$\mathbf{S} = \begin{bmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} \\ \mathbf{S}_{21} & \mathbf{S}_{22} \end{bmatrix}$$

be a $p \times p$ positive semi-definite matrix with $q \times q$ non-singular principal sub-matrix \mathbf{S}_{11} then

$$\lambda_{\min}(\mathbf{S}_{22} - \mathbf{S}_{21}\mathbf{S}_{11}^{-1}\mathbf{S}_{12}) \geq \lambda_{\min}(\mathbf{S}).$$