

Reflexive sheaves, Hermitian-Yang-Mills connections, and tangent cones

Xuemiao Chen*, Song Sun†

December 1, 2021

Abstract

In this paper we give a complete algebro-geometric characterization of analytic tangent cones of admissible Hermitian-Yang-Mills connections over any reflexive sheaves.

Contents

1	Introduction	2
2	Preliminaries	5
2.1	Analytic tangent cones	5
2.2	Algebraic tangent cones	10
2.3	Moduli of semi-stable sheaves	13
3	Optimal algebraic tangent cones from admissible HYM	16
3.1	Properties of the degree function	17
3.2	Construction of torsion-free sheaves on D	21
3.3	The main construction	28
3.4	Optimality	37
4	Proof of the main theorem	43
4.1	The limiting sheaf and connection	43
4.2	The analytic blow-up cycle	48

*University of Maryland; xmchen@umd.edu.

†University of California, Berkeley; sosun@berkeley.edu.

1 Introduction

Let ω be a smooth Kähler metric on the ball $B = \{|z| < 1\}$ in \mathbb{C}^n , and let \mathcal{E} be a reflexive coherent sheaf defined on a neighborhood of \overline{B} . Let A be an admissible Hermitian-Yang-Mills (HYM) connection on \mathcal{E} with respect to the Kähler metric ω . This means that A is the Chern connection of a smooth Hermitian-Yang-Mills (HYM) metric H on the locally free locus of \mathcal{E} , and the curvature F_A has finite L^2 norm on any compact subset of B . Denote by $\text{Sing}(\mathcal{E})$ the singular locus of \mathcal{E} . It is known that $\text{Sing}(\mathcal{E})$ is a complex analytic subset of complex co-dimension at least 3. Our main goal in this paper is to understand the singular behavior of A at 0 in terms of the underlying sheaf \mathcal{E} .

We first briefly review the notions of analytic and algebraic tangent cones, and more details will be provided in Section 2. The notion of *analytic tangent cones* is first studied by Tian [18]. Consider the dilation map

$$\lambda : B_{\lambda^{-1}} \rightarrow B, z \mapsto \lambda \cdot z$$

where $\lambda > 0$, and the family of rescaled connections

$$A_\lambda = \lambda^* A.$$

Letting $\lambda \rightarrow 0$, passing to a subsequence and applying gauge transforms, A_λ converges smoothly to a connection A_∞ on $\mathbb{C}^n \setminus (\Sigma \cup Z(\mathcal{E}))$, and A_∞ is HYM with respect to the standard flat metric. Here Σ is the *bubbling set*, i.e., the subset of $\mathbb{C}^n \setminus Z(\mathcal{E})$ where the convergence is not smooth, and $Z(\mathcal{E})$ is the Zariski tangent cone of $\text{Sing}(\mathcal{E})$ at 0. It is known by [1, 18] that A_∞ extends to an admissible HYM connection on a reflexive sheaf \mathcal{E}_∞ on \mathbb{C}^n , and $\Sigma \cup Z(\mathcal{E})$ is a closed affine algebraic subvariety in \mathbb{C}^n . Moreover, passing to a further subsequence, there is a complex codimension 2 algebraic cycle on \mathbb{CP}^{n-1} , called the *analytic blow-up cycle*, which is of the form

$$\Sigma_b^{an} = \sum_k m_k^{an} \cdot [\underline{\Sigma}_b^k],$$

such that the affine cone over the support $\cup_k \underline{\Sigma}_b^k$ of Σ_b^{an} is precisely the pure complex codimension two part of $\Sigma \cup Z(\mathcal{E})$, and the *analytic multiplicity* m_k^{an} is a positive integer characterizing the blow-up of Yang-Mills energy transverse to $\underline{\Sigma}_b^k$ at a generic point. On the other hand, the rest of Σ is contained in $\text{Sing}(A_\infty) = \text{Sing}(\mathcal{E}_\infty)$. We call the pair $(A_\infty, \Sigma_b^{an})$ an analytic tangent cone.

The terminology “tangent cone” is justified by the fact that A_∞ is a HYM cone connection in the sense of [3] (see Section 2.1). The underlying sheaf \mathcal{E}_∞ is of the form $\psi_* \pi^* \underline{\mathcal{E}}_\infty$, where

$$\pi : \mathbb{C}^n \setminus \{0\} \rightarrow \mathbb{CP}^{n-1}$$

is the natural projection map and

$$\psi : \mathbb{C}^n \setminus \{0\} \rightarrow \mathbb{C}^n$$

is the inclusion map, and $\underline{\mathcal{E}}_\infty$ is a reflexive sheaf on \mathbb{CP}^{n-1} which is a direct sum of polystable¹ sheaves. Moreover, up to gauge equivalence A_∞ is uniquely determined by the sheaf $\underline{\mathcal{E}}_\infty$. So the information of an analytic tangent cone is completely encoded in the algebraic data $(\underline{\mathcal{E}}_\infty, \underline{\Sigma}_b^{an})$. We point out that a priori from the definition analytic tangent cones at 0 depend on not only the initial connection A , but also on the choice of subsequences as $\lambda \rightarrow 0$. Uniqueness of tangent cones independent of subsequences is in general a difficult question in many geometric analytic problems.

Recall from [3, 5] we introduced the notion of an *algebraic tangent cone* at a singularity of a reflexive coherent analytic sheaf \mathcal{E} . This is defined to be a torsion-free sheaf $\widehat{\underline{\mathcal{E}}}$ on the exceptional divisor $D = \mathbb{CP}^{n-1}$ that is given by the restriction of a reflexive extension $\widehat{\mathcal{E}}$ of $p^*(\mathcal{E}|_{B \setminus \{0\}})$ across D , where

$$p : \widehat{B} \rightarrow B$$

is the blowup at 0. In general algebraic tangent cones at 0 are not necessarily unique, due to the fact that the exceptional divisor has complex co-dimension exactly 1. We say an algebraic tangent cone $\widehat{\underline{\mathcal{E}}}$ is *optimal* if

$$\Phi(\widehat{\underline{\mathcal{E}}}) := \mu(\underline{\mathcal{E}}_1) - \mu(\underline{\mathcal{E}}_m / \underline{\mathcal{E}}_{m-1}) \in [0, 1),$$

where

$$0 = \underline{\mathcal{E}}_0 \subset \underline{\mathcal{E}}_1 \subset \cdots \subset \underline{\mathcal{E}}_m = \widehat{\underline{\mathcal{E}}}$$

is the Harder-Narasimhan filtration of $\widehat{\underline{\mathcal{E}}}$, and $\mu(\cdot)$ denotes the slope of a torsion-free sheaf on \mathbb{CP}^{n-1} with respect to the standard polarization. The function Φ measures how far an algebraic tangent cone is from being semistable.

Given a torsion-free sheaf $\underline{\mathcal{E}}$ on \mathbb{CP}^{n-1} , we denote by $Gr^{HN}(\underline{\mathcal{E}})$ (resp. $Gr^{HNS}(\underline{\mathcal{E}})$) the graded sheaf associated to the Harder-Narasimhan (resp. Harder-Narasimhan-Seshadri) filtration of $\underline{\mathcal{E}}$. In [5] it is proved that an optimal algebraic tangent cone always exists and it is unique up to certain natural transforms. In particular, the isomorphism class of the corresponding graded torsion-free sheaf $Gr^{HN}(\widehat{\underline{\mathcal{E}}})$, up to tensoring each factor by some $\mathcal{O}(k)$, does not depend on the choice of optimal algebraic tangent cones. For our purpose, we need to consider instead $Gr^{HNS}(\widehat{\underline{\mathcal{E}}})$. The latter is not unique in general but certain algebraic invariants can be extracted. More specifically, we define a reflexive sheaf over \mathbb{C}^n

$$\mathcal{G}^{alg} := \psi_* \pi^*(Gr^{HNS}(\widehat{\underline{\mathcal{E}}}))^{**}, \quad (1.1)$$

¹Throughout this paper, when we talk about stability of sheaves on the projective space, we always mean *slope* stability with respect to the standard polarization.

and a complex codimension 2 algebraic cycle

$$\Sigma_b^{alg} := \Sigma_b^{alg}(\widehat{\mathcal{E}})$$

on \mathbb{CP}^{n-1} (c.f. Definition 2.18). We call Σ_b^{alg} the *algebraic blow-up cycle* of \mathcal{E} at 0. It is a fact that both \mathcal{G}^{alg} and Σ_b^{alg} do not depend on the specific choice of optimal algebraic tangent cones, so they are local algebraic invariants of the stalk of \mathcal{E} at 0 (see Section 2.2 for more details).

In [5] we made a conjecture relating the analytic and algebraic tangent cones, motivated by the results in [3, 4]. In this paper we give a proof of this conjecture in complete generality, based on the techniques introduced in [3, 4] and a new approach. Simply put, the algebraic data underlying the analytic tangent cones matches exactly with the above algebraic invariants of optimal algebraic tangent cones. More precisely, we have

Theorem 1.1. *Given \mathcal{E} and A as above, then there is a unique analytic tangent cone $(A_\infty, \Sigma_b^{an})$ at 0, which is completely determined by the stalk of \mathcal{E} at 0:*

- (I). *A_∞ is gauge equivalent to the HYM cone on \mathcal{G}^{alg} (see Section 2.1 for the definition of a HYM cone). In particular, \mathcal{E}_∞ is isomorphic to \mathcal{G}^{alg} , and*

$$\text{Sing}(A_\infty) = \text{Sing}(\mathcal{G}^{alg}).$$

- (II) *The analytic blow-up cycle Σ_b^{an} equals the algebraic blow-up cycle Σ_b^{alg} .*

In [3, 4] we proved this result under the extra assumption that 0 is an isolated and homogeneous singularity of \mathcal{E} , or that there is an algebraic tangent cone $\widehat{\mathcal{E}}$ which is locally free and stable (see also [13] for a special case). The arguments there are more analytical and involve PDE estimates based on explicit construction of good background Hermitian metrics. In the general setting the previous approach meets severe difficulties. In this paper, we proceed using a different idea which is more intrinsic and is of more algebraic nature. The key new input in this paper is that, as we will show in Section 3, an admissible HYM connection A naturally recovers all the equivalence classes of optimal algebraic tangent cones (see Section 3.)

In Section 2 we include some background material. In Section 4 we finish the proof of Theorem 1.1, using recent results on moduli compactification of admissible HYM connections and semi-stable sheaves over projective manifolds (see [9] and [10]).

Remark on notations: In this paper we need to introduce many notations regarding sheaves on different spaces. The convention is that a coherent sheaf \mathcal{E} is denoted by a letter with calligraphic font; a coherent sheaf $\underline{\mathcal{E}}$ on the projective space \mathbb{CP}^{n-1} comes with an extra underline; a coherent sheaf $\widehat{\mathcal{E}}$ on the blowup \widehat{B} of the ball B comes with an extra widehat.

Acknowledgements: Both authors were partially supported by the Simons Collaboration Grant on Special Holonomy in Geometry, Analysis and Physics (No. 488633, S.S.), and the second author was partially supported by NSF grant DMS-1916520. The first author would like to thank Santai Qu for helpful discussions. He also thanks UC Berkeley for hospitality during his visit between January 2018 and May 2019. The second author thanks Princeton University for hospitality during Fall 2019. We also thank the anonymous referees for valuable comments which helped improve the exposition of the paper.

2 Preliminaries

2.1 Analytic tangent cones

In this subsection, we recall some backgrounds about analytic tangent cones of Hermitian-Yang-Mills connections. For details see [3]. As in the introduction we let \mathcal{E} be a reflexive sheaf defined over a neighborhood of \overline{B} in \mathbb{C}^n , where $\overline{B} = \{z \in \mathbb{C}^n : |z| \leq 1\}$. We assume that under the standard holomorphic coordinates $\{z_1, \dots, z_n\}$ on \mathbb{C}^n

$$\omega = \omega_0 + O(|z|^2), \quad (2.1)$$

where

$$\omega_0 := \frac{\sqrt{-1}}{2} \partial \bar{\partial} |z|^2$$

is the standard flat metric. Let H be an admissible HYM metric on \mathcal{E} . This means that H is a smooth Hermitian metric on \mathcal{E} outside $\text{Sing}(\mathcal{E})$, and the associated Chern connection A satisfies the following conditions

- the HYM equation

$$\sqrt{-1} \Lambda_\omega F_A = c \cdot \text{Id} \quad (2.2)$$

holds on $B \setminus \text{Sing}(\mathcal{E})$, for some constant $c \in \mathbb{R}$, called the *Einstein constant*;

- A has locally finite Yang-Mills energy:

$$\int_{K \setminus \text{Sing}(\mathcal{E})} |F_A|^2 \, d\text{Vol}_\omega < \infty$$

for any compact subset $K \subset B$.

Notice that a Hermitian-Yang-Mills connection is a projectively unitary Ω -anti-self-dual instanton in the sense of Tian [18], so is *stationary* by [18],

Proposition 5.1.2. Hence by Proposition 5.1.1 in [18], Price's monotonicity formula holds. In particular, we have

$$\sup_{r \in (0,1]} r^{4-2n} \int_{B_r \setminus \text{Sing}(\mathcal{E})} |F_A|^2 d\text{Vol}_\omega < \infty. \quad (2.3)$$

For any $\lambda > 0$, we denote the rescaling map

$$\lambda : B_{\lambda^{-1}} \rightarrow B; z \mapsto \lambda \cdot z,$$

where in this paper B_r always denotes the ball $\{|z| < r\}$ in \mathbb{C}^n . For any sequence of positive numbers $\lambda_j \rightarrow 0$, we get a sequence of admissible HYM connections

$$A_j := \lambda_j^* A$$

with respect to the Kähler metric $\omega_j := \lambda_j^{-2} \cdot \lambda_j^* \omega$. By (2.3), this sequence has uniformly bounded Yang-Mills energy over any compact subset $K \subset \mathbb{C}^n$. Notice by (2.1) obviously ω_j converges smoothly to ω_0 as $j \rightarrow \infty$.

We denote by $Z(\mathcal{E})$ the \mathbb{C}^* invariant reduced subvariety in \mathbb{C}^n underlying the Zariski tangent cone of $\text{Sing}(\mathcal{E})$, i.e., $Z(\mathcal{E})$ is the Hausdorff limit of $\text{Sing}(\mathcal{E})$ in \mathbb{C}^n under the above rescaling as $\lambda \rightarrow 0$. It is well-known that $Z(\mathcal{E})$ has the same complex dimension as $\text{Sing}(\mathcal{E})$ (see for example [19]), so is of complex codimension at least 3 in \mathbb{C}^n .

The associated *bubbling set* of this sequence is defined as

$$\Sigma = \{z \in \mathbb{C}^n \setminus Z(\mathcal{E}) \mid \lim_{r \rightarrow 0} \liminf_{j \rightarrow \infty} r^{4-2n} \int_{B_z(r)} |F_{A_j}|^2 d\text{Vol}_{\omega_j} \geq \epsilon_0\}, \quad (2.4)$$

where ϵ_0 denote the ϵ -regularity constant for Yang-Mills connections over the flat \mathbb{C}^n (see Theorem 2.2.1 in [18] for example). Applying Uhlenbeck's ϵ -regularity theorem and standard analytic results on the convergence of Yang-Mills connections, by passing to a subsequence we may assume there is a smooth connection A_∞ on some Hermitian vector bundle $(\mathcal{E}_\infty, H_\infty)$ defined over $\mathbb{C}^n \setminus (\Sigma \cup Z(\mathcal{E}))$, which is HYM with respect to the flat metric ω_0 . Moreover, for $j \gg 1$, there exist Hermitian isomorphisms

$$P_j : (\lambda_j^* \mathcal{E}, \lambda_j^* H) \rightarrow (\mathcal{E}_\infty, H_\infty)$$

such that $(P_j^{-1})^*(A_j)$ converges smoothly to A_∞ ². The connection A_∞ can be viewed as an admissible HYM connection on \mathbb{C}^n , hence by Bando-Siu we know [1] \mathcal{E}_∞ extends to a reflexive sheaf on \mathbb{C}^n and A_∞ extends smoothly outside $\text{Sing}(\mathcal{E}_\infty) \subset \Sigma \cup Z(\mathcal{E})$, so that the set of essential singularities of A_∞ is given by $\text{Sing}(A_\infty) = \text{Sing}(\mathcal{E}_\infty)$.

²In this paper, when we talk about convergence of a sequence of objects, we often need to pass to a subsequence. We will abuse notation and not re-label the new subsequence, if no confusion arises.

It is proved by Tian (see Theorem 4.3.3 in [18]) that Σ is a complex-analytic set in $\mathbb{C}^n \setminus Z(\mathcal{E})$, and the complement $\Sigma \setminus \text{Sing}(A_\infty)$ has pure complex codimension 2. Since $Z(\mathcal{E})$ is of complex codimension at least 3, by the Remmert-Stein-Shiffman extension theorem (see [16] for example) we know the closure Σ_b of the pure complex codimension 2 part of Σ in \mathbb{C}^n is also a pure codimension 2 complex analytic set in \mathbb{C}^n . Let $\{\Sigma_b^k\}$ be the irreducible components of Σ_b . Then by Theorem 4.3.3 in [18], passing to a subsequence we may assume the convergence of Radon measures on $\mathbb{C}^n \setminus Z(\mathcal{E})$

$$\frac{1}{8\pi^2} |F_{A_j}|^2 d\text{Vol}_{\omega_j} \rightharpoonup \frac{1}{8\pi^2} |F_{A_\infty}|^2 d\text{Vol}_{\omega_0} + \nu, \quad (2.5)$$

with

$$\nu = \sum_k m_k^{an} \cdot \mathcal{H}^{2n-4} \llcorner (\Sigma_b^k \setminus Z(\mathcal{E})),$$

where \mathcal{H}^{2n-4} denotes the $2n - 4$ dimensional Hausdorff measure, and m_k^{an} are positive integers called the analytic multiplicities. Now by definition we have

$$(\cup_k \Sigma_b^k) \cup \text{Sing}(A_\infty) \subset \Sigma \cup Z(\mathcal{E}).$$

Although not needed in this paper, one can show that the two sets above are indeed identical, using Lemma 3.18 in [9].

Fixing an irreducible component Σ_b^k , and taking a generic complex 2-dimensional slice Δ which intersects Σ_b^k transversely, we have the following formula computing analytic multiplicities

Lemma 2.1 ([17], Lemma 4.1).

$$m_k^{an} = \lim_{j \rightarrow \infty} \frac{1}{8\pi^2} \int_{\Delta} \text{Tr}(F_{A_j} \wedge F_{A_j}) - \text{Tr}(F_{A_\infty} \wedge F_{A_\infty}).$$

In our special setting, one can say more about the structure of the limiting data (See [18, 3, 4])

- (1). Σ is invariant under the natural action of \mathbb{C}^* on \mathbb{C}^n .
- (2). $F_{A_\infty}(\partial_r, \cdot) = 0$, where ∂_r is the radial vector field on \mathbb{C}^n ;
- (3). $\sqrt{-1}\Lambda_{\omega_0} F_{A_\infty} = 0$, away from $\text{Sing}(\mathcal{E}_\infty)$.

In particular, each Σ_b^k is an affine cone over a pure codimension 2 algebraic subvariety $\underline{\Sigma}_b^k$ in \mathbb{CP}^{n-1} .

Definition 2.2. The *analytic blow-up cycle* is the codimension 2 algebraic cycle on \mathbb{CP}^{n-1} given by

$$\Sigma_b^{an} := \sum_k m_k^{an} \cdot [\underline{\Sigma}_b^k]$$

Definition 2.3. We call the pair $(A_\infty, \Sigma_b^{an})$ an *analytic tangent cone* of A (or (\mathcal{E}, H)) at 0.

Remark 2.4. The definition here is different from [4], where an analytic tangent cone is defined to be the triple (A_∞, Σ, ν) . But it is easy to see that these two definitions contain exactly the same information. The above definition is more convenient to use in this paper.

Remark 2.5. We emphasize here that the analytic tangent cones are *not* a priori unique, and may depend on the choice of subsequences as $\lambda \rightarrow 0$. But it is not difficult to see that any analytic tangent cone can be obtained by taking the rescaled limit corresponding to a subsequence of the fixed sequence $\{\lambda_j := 2^{-j}\}$. We shall use this fact in Section 3.

The above extra properties in our setting imply that A_∞ is a HYM cone in the following sense. Let $\underline{\mathcal{F}}$ be a polystable reflexive sheaf over \mathbb{CP}^{n-1} with slope μ . By Theorem 3 in [1], there exists an admissible HYM metric \underline{H} on $\underline{\mathcal{F}}$ with respect to the Fubini-Study metric. Now on the reflexive sheaf $\mathcal{F} = \psi_* \pi^* \underline{\mathcal{F}}$, the metric $|z|^{2\mu} \pi^* \underline{H}$ is again an admissible HYM metric, with respect to the flat metric ω_0 , and with vanishing Einstein constant. We let $A_{\mathcal{F}}$ be the associated Chern connection which is an admissible HYM connection. Such $(\mathcal{F}, A_{\mathcal{F}})$ is called a *simple HYM cone*. It should be noted here that tensoring $\underline{\mathcal{F}}$ with some $\mathcal{O}(k)$ does not change the resulting simple HYM cone.

Definition 2.6. A direct sum of simple HYM cones is called a HYM cone. In particular, it is determined by a direct sum of polystable reflexive sheaves on \mathbb{CP}^{n-1} .

The above properties (2) and (3) implies that

Lemma 2.7 ([3], Theorem 2.23). *The tangent cone connection A_∞ is a HYM cone on \mathcal{E}_∞ . More precisely, we can write $\mathcal{E}_\infty = \psi_* \pi^* \underline{\mathcal{E}}_\infty$, where*

$$\underline{\mathcal{E}}_\infty = \bigoplus_i \underline{\mathcal{E}}_\infty^{\mu_i} \quad (2.6)$$

so that each $\underline{\mathcal{E}}_\infty^{\mu_i}$ is a polystable reflexive sheaf with slope given by $\mu_i \in [0, 1)$, with $\mu_i \neq \mu_j$ if $i \neq j$, and A_∞ is gauge equivalent to $\bigoplus_i A_i$ where A_i is the simple HYM cone determined by $\underline{\mathcal{E}}_\infty^{\mu_i}$.

Remark 2.8. It follows from the discussion in [3] that given A_∞ , such $\underline{\mathcal{E}}_\infty$ and the above decomposition are unique up to isomorphism, under the normalization condition that $\mu_i \in [0, 1)$.

To sum up, an analytic tangent cone $(A_\infty, \Sigma_b^{an})$ is uniquely determined by the corresponding algebraic data \mathcal{E}_∞ and Σ_b^{an} up to isomorphisms. Our main result Theorem 1.1 thus gives an algebro-geometric characterization of \mathcal{E}_∞ and Σ_b^{an} in terms of \mathcal{E} itself.

For our purpose later, we also need to discuss the notion of convergence of holomorphic sections. Given a sequence of holomorphic sections σ_j of

$$\mathcal{E}_j := \lambda_j^* \mathcal{E}$$

with uniformly bounded L^2 norm over B and a holomorphic section σ_∞ on some analytic tangent cone \mathcal{E}_∞ over B , then

Definition 2.9. We say σ_j converges to σ_∞ if $P_j(\sigma_j)$ converges to σ_∞ smoothly outside $\Sigma \cup Z(\mathcal{E})$.

The following is essentially a consequence of [1]. For the convenience of readers we include a sketch of proof here.

Lemma 2.10. *For any compact subset $K \subset B$, there is a constant $C = C(K)$ independent of j such that*

$$|\sigma_j|_{L^\infty(K)}^2 \leq C \int_B |\sigma_j|^2 d\text{Vol}_{\omega_j} \quad (2.7)$$

Remark 2.11. Here and later in this paper, the norm of a holomorphic section is always meant to be the one defined by the natural Hermitian metric, and the integral is always taken on the complement of the singular set of the sheaf.

Proof. From Theorem 2 (b) in [1] we know $|\sigma_j|^2$ is locally bounded in B and is smooth away from $\text{Sing}(\mathcal{E}_j)$. By the HYM equation one computes

$$\Delta|\sigma_j|^2 = 2|\nabla\sigma_j|^2 - 2c \cdot \lambda_j^2 |\sigma_j|^2 \geq -2c \cdot \lambda_j^2 |\sigma_j|^2$$

on $B \setminus \text{Sing}(\mathcal{E}_j)$, where c is the Einstein constant of the original admissible HYM metric on \mathcal{E} (c.f. (2.2)). By the fact that $\text{Sing}(\mathcal{E}_j)$ has complex codimension at least 3 and $|\sigma_j|^2$ is locally bounded, using a cut-off function and integration by parts one then sees that $|\sigma_j|^2 \in W_{loc}^{1,2}(B)$, and the inequality $\Delta|\sigma_j|^2 \geq -c \cdot \lambda_j^2 |\sigma_j|^2$ holds in the weak sense on B . Then the conclusion follows from local Moser iteration. \square

It follows easily from (2.7) and the Hartogs's extension theorem (see [16]) for holomorphic sections of reflexive sheaves, that given a sequence $\{\sigma_j\}$ with $\int_B |\sigma_j|^2$ uniformly bounded, one can always extract a convergent subsequence. Moreover since the set $\Sigma \cup Z(\mathcal{E})$ has vanishing Lebesgue measure, we have

$$\int_B |\sigma_\infty|^2 d\text{Vol}_{\omega_0} \leq \liminf_{j \rightarrow \infty} \int_B |\sigma_j|^2 d\text{Vol}_{\omega_j}. \quad (2.8)$$

We may also refer to the above convergence as *weak* convergence.

Definition 2.12. We say σ_j *strongly* converges to σ_∞ if $\{\sigma_j\}$ converges to σ_∞ , and furthermore

$$\int_B |\sigma_\infty|^2 d\text{Vol}_{\omega_0} = \lim_{j \rightarrow \infty} \int_B |\sigma_j|^2 d\text{Vol}_{\omega_j}.$$

Again since $\Sigma \cup Z(\mathcal{E})$ has vanishing Lebesgue measure, it is clear that strong convergence follows from convergence if one can establish an a priori bound

$$\int_B |\sigma_j|^{2+\epsilon} d\text{Vol}_{\omega_j} \leq C$$

for some $\epsilon > 0$. In reality we will indeed derive a uniform L^∞ bound to guarantee strong convergence, see [3] and Section 3.1. In view of (2.7), the key point is to rule out the blowing up of L^∞ norm near ∂B . The following fact will be used in Section 3.2.

Lemma 2.13. *Suppose σ_j and σ'_j converge strongly to σ_∞ and σ'_∞ respectively, and f is a fixed holomorphic function on B , then*

- $\sigma_j + \sigma'_j$ converges strongly to $\sigma_\infty + \sigma'_\infty$;
- $f \cdot \sigma_j$ converges strongly to $f \cdot \sigma_\infty$.

Proof. It suffices to notice that if σ_j converges to σ_∞ , then the convergence is strong if and only if

$$\lim_{r \rightarrow 1^-} \limsup_{j \rightarrow \infty} \int_{B \setminus B_r} |\sigma_j|^2 d\text{Vol}_{\omega_j} = 0.$$

This is itself a consequence of (2.7) and (2.8). □

2.2 Algebraic tangent cones

In this subsection, we collect the results on algebraic tangent cones of reflexive sheaves. For details see [5]. We fix a reflexive sheaf \mathcal{E} over $B \subset \mathbb{C}^n$. Let

$$p : \widehat{B} \rightarrow B$$

denote the blowup of B at $0 \in B$ and denote by $D = \mathbb{CP}^{n-1}$ the exceptional divisor. Then we define \mathcal{A} to be the set of isomorphism classes of reflexive sheaves $\widehat{\mathcal{E}}$ over \widehat{B} so that $\widehat{\mathcal{E}}|_{\widehat{B} \setminus D}$ is isomorphic to $p^*\mathcal{E}|_{\widehat{B} \setminus 0}$. An element $\widehat{\mathcal{E}} \in \mathcal{A}$ is called an *extension* of \mathcal{E} at $0 \in B$ and the torsion-free sheaf

$$\underline{\widehat{\mathcal{E}}} := \iota_D^* \widehat{\mathcal{E}}$$

is called an algebraic tangent cone of \mathcal{E} at 0, where

$$\iota_D : D \rightarrow \widehat{B}$$

denotes the obvious inclusion map. We define a function

$$\Phi : \mathcal{A} \rightarrow \mathbb{Q}_{\geq 0}; \quad \widehat{\mathcal{E}} \mapsto \mu(\underline{\mathcal{E}}_1) - \mu(\underline{\mathcal{E}}_m/\underline{\mathcal{E}}_{m-1}),$$

where

$$0 = \underline{\mathcal{E}}_0 \subset \underline{\mathcal{E}}_1 \subset \cdots \subset \underline{\mathcal{E}}_m = \widehat{\underline{\mathcal{E}}}$$

is the Harder-Narasimhan filtration of $\underline{\mathcal{E}}$.

Definition 2.14. $\widehat{\mathcal{E}} \in \mathcal{A}$ is called an optimal extension of \mathcal{E} at 0 if $\Phi(\widehat{\mathcal{E}}) \in [0, 1)$. In this case we also call $\widehat{\underline{\mathcal{E}}}$ an optimal algebraic tangent cone of \mathcal{E} at 0.

Notice $\widehat{\mathcal{E}}$ is optimal if $\widehat{\underline{\mathcal{E}}}$ is semi-stable, i.e., $\Phi(\widehat{\underline{\mathcal{E}}}) = 0$. But simple examples (see [5]) show that we can not always achieve semistability and this is the reason for introducing the weaker notion of being optimal.

Now given an optimal extension $\widehat{\mathcal{E}}$, and a subsheaf $\underline{\mathcal{E}}_i$ occurring in the Harder-Narasimhan filtration of \mathcal{E} , we define the Hecke transform (or elementary transform) of $\widehat{\mathcal{E}}$ along $\underline{\mathcal{E}}_i$ to be the reflexive sheaf $\widehat{\mathcal{E}}^i$ which is given by the following natural exact sequence

$$0 \rightarrow \widehat{\mathcal{E}}^i \rightarrow \widehat{\mathcal{E}} \rightarrow (\iota_D)_*(\widehat{\underline{\mathcal{E}}}/\underline{\mathcal{E}}_i) \rightarrow 0$$

Then by Corollary 3.3 in [5], we know that $\widehat{\mathcal{E}}^i$ is again an optimal extension of \mathcal{E} at 0. We also say $\widehat{\mathcal{E}}^i$ and $\widehat{\mathcal{E}}$ differ by a *Hecke transform of special type*. It is also shown that the graded sheaves $Gr^{HN}(\widehat{\underline{\mathcal{E}}})$ and $Gr^{HN}(\widehat{\underline{\mathcal{E}}})$ are isomorphic up to tensoring each factor by some $\mathcal{O}(k)$ on D .

Next it is easy to see that if $\widehat{\mathcal{E}} \in \mathcal{A}$, then for any $k \in \mathbb{Z}$, the sheaf $\widehat{\mathcal{E}}' := \widehat{\mathcal{E}} \otimes [D]^k$ is again an extension, where $[D]$ denotes the line bundle on \widehat{B} defined by the divisor D . In this case we say $\widehat{\mathcal{E}}'$ and $\widehat{\mathcal{E}}$ are *equivalent* extensions. Restricting to D , we have $\widehat{\underline{\mathcal{E}}} = \underline{\mathcal{E}} \otimes \mathcal{O}(-k)$, so in particular $\widehat{\mathcal{E}}$ is optimal if and only if $\widehat{\mathcal{E}}'$ is optimal.

Theorem 2.15 ([5]). *Given a reflexive sheaf \mathcal{E} over B , we have*

- *An optimal extension of \mathcal{E} at $0 \in B$ always exists and up to equivalence, two optimal extensions differ by a Hecke transform of special type. In particular, there are exactly m different optimal extensions up to equivalence, where m is the length of the Harder-Narasimhan filtration of $\widehat{\underline{\mathcal{E}}}$.*
- *The graded sheaf $Gr^{HN}(\widehat{\underline{\mathcal{E}}})$ is uniquely determined by \mathcal{E} up to tensoring each factor with some $\mathcal{O}(k)$. In particular the sheaf $\psi_*\pi^*(Gr^{HN}(\widehat{\underline{\mathcal{E}}}))$ on \mathbb{C}^n is uniquely determined by \mathcal{E} .*

Remark 2.16. If we want strict *uniqueness* of optimal extensions, we can impose the normalizing condition that $\mu(\underline{\mathcal{E}}_1)$ and $\mu(\underline{\mathcal{E}}_m/\underline{\mathcal{E}}_{m-1})$ are both

in the interval $[0, 1)$. This will remove the freedom of performing Hecke transform of special type or tensoring with $\mathcal{O}(k)$. For our purpose in this paper, the statement in Theorem 2.15 is more suitable since it implies that each factor of $Gr^{HN}(\widehat{\mathcal{E}})$ can be viewed as the maximal destabilizing subsheaf of *some* optimal algebraic tangent cone.

For our purpose, we need to consider the sheaf $Gr^{HNS}(\widehat{\mathcal{E}})$ which in general depends on the choice of the Harder-Narasihman-Seshadri filtration of $\widehat{\mathcal{E}}$ (see Example 3.1 in [2]). Nonetheless, we can still extract algebraic invariants from $Gr^{HNS}(\widehat{\mathcal{E}})$ that suffice for the need in this paper.

We first introduce a general definition following [2].

Definition 2.17. Given a torsion sheaf \mathcal{T} on \mathbb{CP}^{n-1} with support in codimension at least 2, we define the codimension 2 *support cycle* of \mathcal{T} to be the algebraic cycle

$$\mathcal{C}(\mathcal{T}) := \sum_k m_k^{alg} \cdot [\underline{\Sigma}_k]$$

where $\underline{\Sigma}_k$ are irreducible codimension 2 components of the support of \mathcal{T} , and the *algebraic multiplicity*

$$m_k^{alg} = h^0(\underline{\Delta}, \mathcal{T}|_{\underline{\Delta}})$$

for a complex 2 dimensional slice $\underline{\Delta}$ which intersects $\underline{\Sigma}_k$ transversely at a generic point.

Definition 2.18. Let $\underline{\mathcal{F}}$ be a torsion-free coherent sheaf on \mathbb{CP}^{n-1} . We define the pure codimension 2 algebraic cycle $\Sigma_b^{alg}(\underline{\mathcal{F}})$ of $\underline{\mathcal{F}}$ to be

$$\Sigma_b^{alg}(\underline{\mathcal{F}}) := \mathcal{C}(\mathcal{T}),$$

where $\mathcal{T} = (Gr^{HNS}(\underline{\mathcal{F}}))^{**}/Gr^{HNS}(\underline{\mathcal{F}})$.

Remark 2.19. It follows from the definition that $\Sigma_b^{alg}(\underline{\mathcal{F}})$ is supported on the pure codimension 2 part of the support of \mathcal{T} . When $\underline{\mathcal{F}}$ is locally free, we know by Proposition 2.3 in [17] that away from $\text{Sing}((Gr^{HNS}(\underline{\mathcal{F}}))^{**})$, the support of \mathcal{T} has pure codimension 2.

We have

Proposition 2.20 ([2] Proposition 2.1). *Given a semistable torsion free sheaf $\underline{\mathcal{F}}$ on \mathbb{CP}^{n-1} , the reflexive sheaf $(Gr^{HNS}(\underline{\mathcal{F}}))^{**}$ and the codimension 2 algebraic cycle $\Sigma_b^{alg}(\underline{\mathcal{F}})$ do not depend on the choice of the Harder-Narasihman-Seshadri filtration of $\underline{\mathcal{F}}$.*

An immediate corollary is

Corollary 2.21. *For a reflexive sheaf \mathcal{E} over B , let $\widehat{\mathcal{E}}$ be any optimal extension of \mathcal{E} at $0 \in B$, then the sheaf*

$$\mathcal{G}^{alg} := \psi_* \pi^* (Gr^{HNS}(\widehat{\mathcal{E}}))^{**} \quad (2.9)$$

and the algebraic blow-up cycle

$$\Sigma_b^{alg} := \Sigma_b^{alg}(\widehat{\mathcal{E}}) \quad (2.10)$$

do not depend on the choice of the optimal extension $\widehat{\mathcal{E}}$ at 0 , hence are algebraic invariants of the stalk of \mathcal{E} at 0 .

Remark 2.22. Since $(Gr^{HNS}(\widehat{\mathcal{E}}))^{**}$ is a direct sum of polystable sheaves, we can apply the construction in Section 2.1 to obtain a HYM cone on the sheaf \mathcal{G}^{alg} . By Corollary 2.21, we know that up to gauge equivalence such a HYM cone is *canonically* associated to the stalk of \mathcal{E} at 0 .

2.3 Moduli of semi-stable sheaves

In this subsection, we will review some algebro-geometric results from [9, 10]. We mention that these results will not be used until Section 4.1. The results are proved on general polarized projective manifolds but for our purpose we will only consider the case when the base manifold is \mathbb{CP}^{n-1} with the standard polarization $\mathcal{O}(1)$.

Let $\underline{\mathcal{E}}$ be a semi-stable torsion-free sheaf on \mathbb{CP}^{n-1} . Denote by r the rank of $\underline{\mathcal{E}}$, and denote by τ the Hilbert polynomial of $\underline{\mathcal{E}}$. Throughout this paper we shall denote

$$\underline{\mathcal{E}}(k) := \underline{\mathcal{E}} \otimes \mathcal{O}(k).$$

Since the set of semi-stable torsion-free sheaves having the same Hilbert polynomials as $\underline{\mathcal{E}}$ forms a bounded family (see [15]), we may fix k large so that for any such $\underline{\mathcal{E}}'$, $\underline{\mathcal{E}}'(k)$ is globally generated, and

$$H^i(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}'(k)) = 0$$

for all $i > 0$. Denote the sheaf

$$\mathcal{H} = \mathbb{C}^{\oplus \tau(k)} \otimes \mathcal{O}(-k).$$

Choosing a basis of $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}(k))$ gives an exact sequence

$$\mathcal{H} \xrightarrow{q} \underline{\mathcal{E}} \rightarrow 0,$$

hence yields a point in the Quot scheme $\mathbf{Quot}(\mathcal{H}, \tau)$. Here $\mathbf{Quot}(\mathcal{H}, \tau)$ is by definition the set of equivalence classes of quotients

$$q : \mathcal{H} \rightarrow \underline{\mathcal{E}}' \rightarrow 0,$$

where $\underline{\mathcal{E}}'$ is a coherent sheaf over \mathbb{CP}^{n-1} with Hilbert polynomial equal to τ . Two quotients $\underline{q}_1 : \mathcal{H} \rightarrow \underline{\mathcal{E}}_1$ and $\underline{q}_2 : \mathcal{H} \rightarrow \underline{\mathcal{E}}_2$ are equivalent if $\text{Ker}(\underline{q}_1) = \text{Ker}(\underline{q}_2)$; this is the same as saying that there exists an isomorphism $\underline{\rho} : \underline{\mathcal{E}}_1 \rightarrow \underline{\mathcal{E}}_2$ so that

$$\underline{\rho} \circ \underline{q}_1 = \underline{q}_2.$$

Notice there is a natural action of $GL(\tau(k), \mathbb{C})$ on $\mathbf{Quot}(\mathcal{H}, \tau)$ given by

$$M.\underline{p}' = \underline{p}' \circ M$$

for any $M \in GL(\tau(k), \mathbb{C})$ and any quotient \underline{p}' in $\mathbf{Quot}(\mathcal{H}, \tau)$.

By [11], we know $\mathbf{Quot}(\mathcal{H}, \tau)$ is a projective scheme which admits a decomposition

$$\mathbf{Quot}(\mathcal{H}, \tau) = \coprod \mathbf{Quot}(\mathcal{H}, (c_1, \dots, c_{\min(r, n-1)})),$$

where $\mathbf{Quot}(\mathcal{H}, (c_1, \dots, c_{\min(r, n-1)}))$ consists of those quotients with fixed Chern classes $(c_1, \dots, c_{\min(r, n-1)})$. Let

$$\mathbf{Quot}(\mathcal{H}, c(\underline{\mathcal{E}})) = \mathbf{Quot}(\mathcal{H}, c_1(\underline{\mathcal{E}}), \dots, c_{\min(r, n-1)}(\underline{\mathcal{E}})).$$

Now we denote by $R^{\mu ss} \subset \mathbf{Quot}(\mathcal{H}, c(E))$ the subscheme of quotients $q : \mathcal{H} \rightarrow \underline{\mathcal{E}}' \rightarrow 0$ so that

- $\underline{\mathcal{E}}'$ is torsion-free;
- $\det(\underline{\mathcal{E}}') \cong \det(\underline{\mathcal{E}})$;
- $\underline{\mathcal{E}}'$ is semi-stable;
- q induces an isomorphism $\mathbb{C}^{\oplus \tau(k)} \cong H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}'(k))$.

Let \mathcal{Z} denote the reduced weak normalization of $R^{\mu ss}$ as a complex analytic space. This means that we first take the underlying reduced complex analytic space of $R^{\mu ss}$, and then take its weak normalization, so that every locally defined *continuous* function on \mathcal{Z} which is holomorphic on the smooth part \mathcal{Z}_{reg} of \mathcal{Z} is in fact holomorphic. Notice the weak normal property is “weaker” than being normal, in the sense that we impose the extension property only for continuous functions. As a simple example, in complex dimension 1, a nodal singularity is weakly normal but not normal, and a cusp singularity is not weakly normal.

Proposition 2.23 ([10], Definition 4.4. and Theorem 5.5). *There exists a compact complex analytic space $M^{\mu ss}$, with a natural continuous map*

$$\phi : \mathcal{Z} \rightarrow M^{\mu ss}$$

so that

- The image of a fixed $GL(\tau(k), \mathbb{C})$ orbit in \mathcal{Z} is a point;
- If two quotients $q_i : \mathcal{H} \rightarrow \underline{\mathcal{E}}_i, i = 1, 2$ in \mathcal{Z} have the same image in $M^{\mu ss}$, then

$$(Gr^{HNS}(\underline{\mathcal{E}}_1))^{**} \cong (Gr^{HNS}(\underline{\mathcal{E}}_2))^{**},$$

and

$$\Sigma_b^{alg}(\underline{\mathcal{E}}_1) = \Sigma_b^{alg}(\underline{\mathcal{E}}_2).$$

For our purpose in this paper, we also need the following fact regarding the convergence of a sequence in the space $\mathbf{Quot}(\mathcal{H}, \tau)$ in the analytic topology. We fix a smooth Hermitian metric on \mathcal{H} . Given any sequence of $M_i \in GL(\tau(k), \mathbb{C})$, we define

$$q_i = q \circ M_i : \mathcal{H} \rightarrow \underline{\mathcal{E}} \rightarrow 0.$$

Furthermore, we assume q_i converge to

$$q_\infty : \mathcal{H} \rightarrow \underline{\mathcal{E}}_\infty$$

in the analytic topology of $\mathbf{Quot}(\mathcal{H}, \tau)$.

The maps q_i (resp. q_∞) induce smooth bundle endomorphisms $\pi_i : \mathcal{H} \rightarrow \mathcal{H}$ (resp. $\pi_\infty : \mathcal{H} \rightarrow \mathcal{H}$), which are given by projection onto the orthogonal complement of $\text{Ker}(q_i)$ (resp. $\text{Ker}(q_\infty)$) in \mathcal{H} , and are defined away from $\text{Sing}(\underline{\mathcal{E}})$ (resp. $\text{Sing}(\underline{\mathcal{E}}_\infty)$). We have

Lemma 2.24 ([9]). *Away from $\text{Sing}(\underline{\mathcal{E}}) \cup \text{Sing}(\underline{\mathcal{E}}_\infty)$, π_i converges to π_∞ smoothly.*

This follows exactly the same as the proof of Lemma 2.17 in [9]. For the convenience of readers we reproduce the arguments here. The key point is a geometric interpretation of the abstract convergence of q_i to q_∞ in \mathbf{Quot} . To see this we use the fact that for some fixed m large, the \mathbf{Quot} scheme \mathbf{Quot} admits an embedding into a fixed Grassmannian $Gr(W, r)$ where $W = H^0(\mathbb{CP}^{n-1}, \mathcal{H}(m))$ and $r = \dim W - \tau(m)$. Now by fixing a Hermitian metric on W , q_i (resp. q_∞) can be viewed as a vector space endomorphism P_{q_i} (resp. P_{q_∞}) of W given by orthogonal projection onto the subspace $H^0(\mathbb{CP}^{n-1}, \text{Ker}(q_i)(m))$ (resp. $H^0(\mathbb{CP}^{n-1}, \text{Ker}(q_\infty)(m))$). The convergence of q_i to q_∞ implies that P_{q_i} converges to P_{q_∞} . On the other hand, we can also view P_{q_i} (resp. P_{q_∞}) as a bundle endomorphism of the holomorphic vector bundle corresponding to the sheaf $W \otimes \mathcal{O}_{\mathbb{CP}^{n-1}}$, and the image of the natural composition

$$W \otimes \mathcal{O}_{\mathbb{CP}^{n-1}} \xrightarrow{P_{q_i} \text{ (resp. } P_{q_\infty})} W \otimes \mathcal{O}_{\mathbb{CP}^{n-1}} \rightarrow \mathcal{H}(m)$$

is exactly given by $\text{Ker}(q_i)(m)$ (resp. $\text{Ker}(q_\infty)(m)$) away from $\text{Sing}(\underline{\mathcal{E}})$ (resp. $\text{Sing}(\underline{\mathcal{E}}_\infty)$). This yields that away from $\text{Sing}(\underline{\mathcal{E}}) \cup \text{Sing}(\underline{\mathcal{E}}_\infty)$, the sub-bundles

$\text{Ker}(q_i)$ converge smoothly to $\text{Ker}(q_\infty)$. This clearly implies the convergence of π_i .

Combining with the continuity of the map ϕ in Proposition 2.23 we obtain

Corollary 2.25. *If $\underline{\mathcal{E}}_\infty$ is a torsion-free and semi-stable, then*

$$(Gr^{HNS}(\underline{\mathcal{E}}_\infty))^{**} = (Gr^{HNS}(\underline{\mathcal{E}}))^{**}$$

and

$$\Sigma_b^{alg}(\underline{\mathcal{E}}_\infty^{alg}) = \Sigma_b^{alg}(\underline{\mathcal{E}}).$$

Furthermore, if $\underline{\mathcal{E}}_\infty^{**}$ is polystable, then

$$\Sigma_b^{alg}(\underline{\mathcal{E}}) = \mathcal{C}(\underline{\mathcal{E}}_\infty^{**}/\underline{\mathcal{E}}_\infty).$$

Proof. The first part follows from Proposition 2.23 directly since q_i stays in a fixed $GL(\tau(k), \mathbb{C})$ orbit. It remains to prove the second part. Indeed, we take a Seshadri filtration of $\underline{\mathcal{E}}_\infty$ as

$$0 \subset \underline{\mathcal{E}}_1 \subset \cdots \subset \underline{\mathcal{E}}_m = \underline{\mathcal{E}}_\infty$$

where $\underline{\mathcal{E}}_i$ are saturated in $\underline{\mathcal{E}}_\infty$. Since $\underline{\mathcal{E}}_\infty^{**}$ is polystable, we have a canonical isomorphism $\underline{\mathcal{E}}_\infty^{**}/\underline{\mathcal{E}}_1^{**} \cong (\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1)^{**}$. Consequently, using the fact that $\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1$ is torsion-free, we obtain the following exact sequence

$$0 \rightarrow (\underline{\mathcal{E}}_1)^{**}/\underline{\mathcal{E}}_1 \rightarrow \underline{\mathcal{E}}_\infty^{**}/\underline{\mathcal{E}}_\infty \rightarrow (\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1)^{**}/(\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1) \rightarrow 0.$$

Since each term has support in codimension at least 2, it follows from Lemma 2.15 in [9] that

$$\mathcal{C}(\underline{\mathcal{E}}_\infty^{**}/\underline{\mathcal{E}}_\infty) = \mathcal{C}((\underline{\mathcal{E}}_1)^{**}/\underline{\mathcal{E}}_1) + \mathcal{C}((\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1)^{**}/(\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1)).$$

Since by assumption $\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1$ is again a torsion-free sheaf whose double dual is polystable, the conclusion follows from induction by repeating the process for $\underline{\mathcal{E}}_\infty/\underline{\mathcal{E}}_1$. \square

3 Optimal algebraic tangent cones from admissible HYM

The goal of this subsection is to show that an admissible HYM connection naturally gives rise to optimal algebraic tangent cones. More precisely, in Section 3.1 we study general properties of the degree function introduced in [3]; in Section 3.2 we define certain canonical torsion-free sheaves on the exceptional divisor D of the blowup $p : \widehat{B} \rightarrow B$; in Section 3.3 we show these torsion-free sheaves do arise as algebraic tangent cones; in Section 3.4 we show these algebraic tangent cones are optimal.

3.1 Properties of the degree function

We first recall the definition of the degree function in [3]. Given (\mathcal{E}, H, A) as in the introduction, we denote by \mathcal{E}_0 the stalk of \mathcal{E} at 0. Then the degree function

$$d : \mathcal{E}_0 \rightarrow \mathbb{R} \cup \{\infty\} \quad (3.1)$$

is defined by setting that $d(s) = \infty$ if $s = 0$ is the zero section, and that for a non-zero holomorphic section s defined in a neighborhood of 0,

$$d(s) := \lim_{r \rightarrow 0^+} \frac{\log \int_{B_r} |s|^2 d\text{Vol}_\omega}{2 \log r} - n. \quad (3.2)$$

Lemma 3.1. *$d(s)$ is well-defined and lies in $(\text{rank}(\mathcal{E})!)^{-1}\mathbb{Z} \cup \{\infty\}$*

Proof. This is proved in Corollary 3.7 in [3] in the setting when 0 is an isolated singularity, but this assumption is not essentially used there. Two key points are the interior estimate (2.7), and the Hartogs's extension theorem. \square

The same definition applies to an analytic tangent cone \mathcal{E}_∞ . Because of the cone structure we have a notion of *homogeneous holomorphic sections* on \mathcal{E}_∞ . We say a holomorphic section s of \mathcal{E}_∞ is homogeneous of degree β if away from $\text{Sing}(\mathcal{E}_\infty)$, we have

$$\nabla_{\partial_r} s = \beta r^{-1} s.$$

It is easy to see for such s , $d(s) = \beta$ using the above definition of the degree function. The following Lemma will be used in the next section.

Lemma 3.2. *Given any analytic tangent cone \mathcal{E}_∞ , then for any fixed $\beta \in \mathbb{R}$, the space V_β of homogeneous holomorphic sections on \mathcal{E}_∞ of degree β is finite dimensional.*

Proof. We define a norm on V_β by setting

$$\|s\|_{L^2(B)}^2 := \int_B |s|^2 d\text{Vol}_{\omega_0}.$$

It suffices to show that the unit sphere in V_β is compact. Given a sequence $s_j \in V_\beta$ with $\|s_j\|_{L^2(B)} = 1$, then after passing to a subsequence we obtain a weak limit s_∞ with $\|s_\infty\|_{L^2(B)} \leq 1$. We need to show the equality holds. Notice that away from $\text{Sing}(\mathcal{E}_\infty)$, s_j converges smoothly to s_∞ . So it suffices to prevent mass concentration near ∂B . The key point is that by homogeneity we have for all j

$$\int_{B_2} |s_j|^2 d\text{Vol}_{\omega_0} = 2^{2n+2\beta} \quad (3.3)$$

So by (2.7) we obtain $\|s_j\|_{L^\infty(B)} \leq C$ for a uniform $C > 0$. Then it is easy to conclude. \square

The understanding of the above degree function is crucial in studying analytic tangent cones. This is first used in [6] when studying singularities of Kähler-Einstein metrics and then was introduced in [3] to study singularities of HYM connections. It is proved in [3] that if $d(s) < \infty$ then s gives rise to non-trivial limit homogeneous sections of degree $d(s)$ on all the analytic tangent cones, hence it provides a basic link between \mathcal{E} and \mathcal{E}_∞ . Notice as pointed out in Remark 2.5 when studying analytic tangent cones we may restrict to a fixed sequence $\lambda_j \rightarrow 0$ given by

$$\lambda_j := 2^{-j}.$$

If we denote

$$[s]_j := \frac{\lambda_j^* s}{\|s\|_j},$$

and

$$\|s\|_j := \|\lambda_j^* s\|_{L^2(B)} = \sqrt{\frac{1}{\text{Vol } B_{2^{-j}}} \int_{B_{2^{-j}}} |s|^2 d\text{Vol}_\omega}, \quad (3.4)$$

then passing to subsequences we get strong convergence of $[s]_j$ to homogeneous holomorphic sections of degree $d(s)$ on analytic tangent cones. Any such limit s_∞ satisfies $\|s_\infty\|_{L^2(B)} = 1$. Again the arguments in [3] were written in the setting of an isolated singularity of \mathcal{E} but tracing the proof one sees that this assumption is not used. Notice if $d(s) = \infty$, which a priori could be the case, then we will not be able to obtain anything interesting. Therefore it is important to find sections s with $d(s)$ finite.

In [3, 4] our idea to study the degree function was to compare the unknown Hermitian metric H with certain explicitly constructed background Hermitian metric. This allows us to compute the degree explicitly when \mathcal{E} is homogeneous, i.e., $\mathcal{E} = \psi_* \pi^* \underline{\mathcal{E}}$ for some locally free sheaf $\underline{\mathcal{E}}$ on \mathbb{CP}^{n-1} . In general when \mathcal{E} is non-homogeneous or when \mathcal{E} has non-isolated singularities this approach seems to involve very complicated difficulties. In this paper a crucial new observation is that one can directly show finiteness of $d(s)$ for a non-zero s , and use this to perform abstract studies without explicit computation of $d(s)$. The main result of this subsection is

Theorem 3.3. *The following hold*

- (1). *For all $s \in \mathcal{E}_0$, we have $d(s) = \infty$ if and only if $s = 0$;*
- (2). *For all $s \in \mathcal{E}_0$, we have $d(s) \geq 0$;*
- (3). *Given $s, s' \in \mathcal{E}_0$, we have*

$$d(s + s') \geq \min\{d(s), d(s')\}; \quad (3.5)$$

(4). Suppose there is another admissible HYM connection (A', H') on a reflexive sheaf \mathcal{E}' over B , then for $s \in \mathcal{E}_0$ and $s' \in \mathcal{E}'_0$, we have

$$d(s \otimes s') = d(s) + d(s'), \quad (3.6)$$

where in each term the degree function has the obvious meaning.

Proof. The key is Item (1). First we recall that by Cartan's Theorem A any coherent sheaf over a Stein manifold is generated by global sections. Now by assumption the dual sheaf \mathcal{E}^* is defined on a neighborhood Ω of \overline{B} , so it is generated by finitely many global sections over a slightly smaller neighborhood Ω' of \overline{B} . In other words, over Ω' we have a surjective sheaf homomorphism $\mathcal{O}^{\oplus n_1} \rightarrow \mathcal{E}^*$ for some n_1 . Applying this again to the kernel sheaf, we then obtain an exact sequence on B of the form

$$\mathcal{O}^{\oplus n_2} \rightarrow \mathcal{O}^{\oplus n_1} \rightarrow \mathcal{E}^* \rightarrow 0. \quad (3.7)$$

Taking dual we obtain

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{O}^{\oplus n_1} \xrightarrow{\rho} \mathcal{O}^{\oplus n_2} \quad (3.8)$$

We can now endow the natural flat Hermitian metric on $\mathcal{O}^{\oplus n_1}$, then we also get induced Hermitian metrics H_0 on \mathcal{E} and H_0^* on \mathcal{E}^* away from $\text{Sing}(\mathcal{E}) = \text{Sing}(\mathcal{E}^*)$.

Lemma 3.4. *There exists a constant $C > 0$ such that on $B_{1/2} \setminus \text{Sing}(\mathcal{E})$ we have*

$$H \geq C \cdot H_0 \quad (3.9)$$

Proof. By basic linear algebra it suffices to show $H^* \leq CH_0^*$. This then follows from the fact that any holomorphic section ζ of \mathcal{E}^* over B has $|\zeta|_{H^*}$ uniformly bounded on $B_{1/2}$ (see Theorem 2 in [1]). \square

For a holomorphic function f defined on a neighborhood of 0, we denote by $\deg(f)$ the vanishing order of f at 0. Notice if f is not identically zero, then

$$\deg(f) = \frac{1}{2} \lim_{r \rightarrow 0} \frac{\log \int_{B_r} |f|^2 d\text{Vol}_{\omega_0}}{\log r} - n.$$

This is an easy consequence using Taylor expansion of f at 0. We also make the convention that $\deg(f) = \infty$ if f is identically zero.

Given any non-zero $s \in \mathcal{E}_0$, we define

$$d^0(s) = \frac{1}{2} \lim_{r \rightarrow 0} \frac{\log \int_{B_r} |s|_{H_0}^2 d\text{Vol}_{\omega_0}}{\log r} - n.$$

Here we emphasize that d^0 is defined with respect to the metric H_0 . If we view s as a tuple of holomorphic functions (F_1, \dots, F_{n_1}) using (3.8), then it is easy to see that if $s \neq 0$, then

$$d^0(s) = \min_j \deg(F_j).$$

Lemma 3.4 then shows that for nonzero s ,

$$d(s) \leq d_0(s) < \infty.$$

This proves Item (1) of Theorem 3.3.

Item (2) follows from the fact that for any holomorphic section s of \mathcal{E} , $|s|_H$ is locally bounded (c.f. [1], Theorem 2). Item (3) follows easily from the definition.

Now we prove Item (4). Given nonzero s and s' , by passing to a subsequence we may assume the rescaled sequences $[s]_j, [s']_j, [s \otimes s']_j$ converges strongly to nonzero homogeneous limit sections $s_\infty, s'_\infty, s''_\infty$ respectively. It follows from definition that $[s]_j \otimes [s']_j$ converges strongly to $s_\infty \otimes s'_\infty$. On the other hand, we have

$$[s \otimes s']_j = C_j \cdot [s]_j \otimes [s']_j$$

for some $C_j \in \mathbb{C}^*$. Since $s_\infty, s'_\infty, s''_\infty$ are all nonzero, it follows that $|C_j|$ and $|C_j|^{-1}$ are uniformly bounded as j tends to infinity. Passing to a further subsequence we may assume

$$s''_\infty = C_\infty s_\infty \otimes s'_\infty$$

for some $C_\infty \neq 0$. It then follows that

$$d(s \otimes s') = d(s''_\infty) = d(s_\infty) + d(s'_\infty) = d(s) + d(s').$$

This finishes the proof of Theorem 3.3. \square

Corollary 3.5. *Denote by \mathcal{O}_0 the stalk at 0 of the sheaf of holomorphic functions on B . Then for all $s \in \mathcal{E}_0$ and $f \in \mathcal{O}_0$, we have*

$$d(fs) = \deg(f) + d(s). \quad (3.10)$$

Proof. This follows from Item (4) above, applied to the case when \mathcal{E}' is the trivial Hermitian line bundle on B . \square

For our purpose later, we also need the following semi-continuity property of degrees under taking analytic tangent cones. This property will be crucial in a few places later in this section.

Proposition 3.6. *Let s_j be a sequence of holomorphic sections of \mathcal{E} over a fixed neighborhood B' of 0. Suppose $d(s_j) \geq \mu$ for all j , and the rescaled sequence $[s_j]_j$ converges to a nonzero limit section s_∞ on some analytic tangent cone as $j \rightarrow \infty$, then s_∞ (which is not necessarily homogeneous) has degree at least μ .*

Proof. We first make the following

Claim 3.7. *For any $\epsilon > 0$ small enough so that $\mu - \epsilon \notin ((\text{rank } \mathcal{E})!)^{-1}\mathbb{Z}$, there exists an $i_0 = i_0(\epsilon)$, so that for any $i \geq i_0$ and $s \in H^0(B', \mathcal{E})$ with $d(s) \geq \mu$, we have $\|s\|_i \leq 2^{-(\mu-\epsilon)}\|s\|_{i-1}$.*

Given this Claim, it follows that

$$\|[s_j]_j\|_i \leq 2^{-(\mu-\epsilon)}\|[s_j]_j\|_{i-1}$$

for all $i \geq i_0$. Taking limit as $j \rightarrow \infty$ we obtain

$$\|s_\infty\|_i \leq 2^{-(\mu-\epsilon)}\|s_\infty\|_{i-1}$$

for all $i \geq i_0 + 1$. It follows that $d(s_\infty) \geq \mu - \epsilon$. Letting $\epsilon \rightarrow 0$ we obtain the conclusion. \square

Proof of Claim 3.7. Otherwise, there exists a subsequence j_i and $s_{j_i} \in H^0(B', \mathcal{E})$ with $d(s_{j_i}) \geq \mu$, so that for all i large we have

$$\|s_{j_i}\|_i > 2^{-(\mu-\epsilon)}\|s_{j_i}\|_{i-1}.$$

On the other hand, by Proposition 3.15 in [3] (again the proof extends trivially to our general setting), we know that there exists some i'_0 (depending on ϵ) so that for any $i \geq i'_0$ and $s \in H^0(B_{2^{-i}}, \mathcal{E})$ if

$$\|s\|_i \geq 2^{-(\mu-\epsilon)}\|s\|_{i-1}$$

then

$$\|s\|_{i+1} \geq 2^{-(\mu-\epsilon)}\|s\|_i.$$

Then for any i so that $j_i \geq i'_0$, we know s_{j_i} must have degree smaller than $\mu - \epsilon$ which is a contradiction. \square

3.2 Construction of torsion-free sheaves on D

In this subsection we define certain canonical torsion-free coherent sheaves on the exceptional divisor $D = \mathbb{CP}^{n-1}$ of the blowup $p : \widehat{B} \rightarrow B$, which are intrinsically associated to the HYM connection A on \mathcal{E} .

The construction of this subsection can be done using only the stalk \mathcal{E}_0 , but for the discussion in the next subsection it is more convenient that we

work with global sections over B instead of the stalk \mathcal{E}_0 . Clearly the degree function defined previously induces a degree function

$$d : H^0(B, \mathcal{E}) \rightarrow (\text{rank}(\mathcal{E})!)^{-1} \mathbb{Z}_{\geq 0} \cup \{\infty\}, \quad (3.11)$$

which satisfies the same properties as those listed in Theorem 3.3.

Let $\mathcal{S} = \text{Im}(d)$. We list the nonnegative numbers in $\mathcal{S} + \mathbb{Z}$ as

$$0 \leq \mu_1 < \cdots < \mu_k < \mu_{k+1} < \cdots. \quad (3.12)$$

We denote by m the biggest integer such that $\mu_m - \mu_1 < 1$. It follows from Corollary 3.5 that for $i = 1, \dots, m$ and $l \in \mathbb{Z}_{\geq 0}$, we have $\mu_{i+ml} = \mu_i + l$.

Definition 3.8. For $k \geq 1$, we define

$$M_k := \{s \in H^0(B, \mathcal{E}) \mid d(s) \geq \mu_k\}. \quad (3.13)$$

It follows from Theorem 3.3 (3) that M_k is a \mathbb{C} -vector space. Since any holomorphic section $s \in H^0(B, \mathcal{E})$ with $d(s) < \infty$ gives rise to nonzero homogeneous sections on the analytic tangent cones with degree $d(s)$, it follows that on any analytic tangent cone \mathcal{E}_∞ , for each $i = 1, \dots, m$, there is a nontrivial direct summand $\underline{\mathcal{E}}_\infty^{\mu_i}$ of $\underline{\mathcal{E}}_\infty$ (see Lemma 2.7). However at this moment we do not know if there are possibly other direct summands of $\underline{\mathcal{E}}_\infty$ since we have not shown how to construct local holomorphic sections of \mathcal{E} from a homogeneous section of \mathcal{E}_∞ . Later we will indeed show there are no extra direct summands, see Remark 3.32.

Proposition 3.9. For each k , M_k/M_{k+1} is finite dimensional as a \mathbb{C} -vector space.

Proof. Suppose M_k/M_{k+1} is nontrivial. We choose a sequence of elements $s_{k,1}, s_{k,2}, \dots$ in M_k in the following way. We first choose $s_{k,1}$ such that the induced element $\tilde{s}_{k,1}$ in M_k/M_{k+1} is not zero. Suppose $s_{k,1}, \dots, s_{k,l}$ are chosen. If $\tilde{s}_{k,1}, \dots, \tilde{s}_{k,l}$ span M_k/M_{k+1} then we stop. Otherwise we choose $s_{k,l+1}$ such that $\tilde{s}_{k,1}, \dots, \tilde{s}_{k,l+1}$ are linearly independent in M_k/M_{k+1} . This process a priori may be infinite. In any case if $l \leq \dim_{\mathbb{C}} M_k/M_{k+1}$ then we denote by $\mathcal{H}_{k,l}$ the \mathbb{C} -vector space spanned by $s_{k,1}, \dots, s_{k,l}$. Then by definition $\mathcal{H}_{k,l} \cap M_{k+1} = 0$, hence $d(s) = \mu_k$ for all $s \in \mathcal{H}_{k,l} \setminus \{0\}$.

From these we also construct a sequence of L^2 orthonormal sections $\{\sigma_{k,i}^j\}$ over B for j sufficiently large (depending on k and i), as follows. First we define for $j \gg 1$

$$\sigma_{k,1}^j := \frac{\lambda_j^* s_{k,1}}{\|\lambda_j^* s_{k,1}\|_{L^2(B)}}.$$

Suppose $\sigma_{k,1}^j, \dots, \sigma_{k,l}^j$ are defined for $j \geq j_0$. Then for $j \gg j_0$ we define $\sigma_{k,l+1}^j$ to be the L^2 orthonormal projection of $\lambda_j^* s_{k,l+1}^j$ to the complement of

the space spanned by $\sigma_{k,1}^j, \dots, \sigma_{k,l}^j$, and then normalized to have L^2 norm 1. This is the standard Gram-Schmidt process.

Now fix an analytic tangent cone \mathcal{E}_∞ . After passing to a subsequence we may assume for each $i = 1, \dots, l$, $\sigma_{k,i}^j$ converges to a holomorphic section $\sigma_{k,i}^\infty$ on \mathcal{E}_∞ with $\|\sigma_{k,i}^\infty\|_{L^2(B)} \leq 1$, and they are L^2 orthogonal over B . Now for each i , $\sigma_{k,i}^\infty$ is homogeneous of degree μ_k and $\|\sigma_{k,i}^\infty\|_{L^2(B)} = 1$. Indeed, for $i = 1$ this is simply the fact that $d(s_{k,1}) = \mu_k$; for $i \geq 2$ this follows from the same induction argument as Proposition 3.12 in [3]. Given this, it follows that l can not be bigger than the dimension of homogeneous holomorphic sections of degree μ_k on \mathcal{E}_∞ , which is finite by Lemma 3.2. \square

We denote

$$n_k = \dim_{\mathbb{C}} M_k/M_{k+1},$$

then the above process stops with $l = n_k$. For any $\mu \in \{\mu_1, \dots, \mu_m\}$, we define

$$N^\mu := \bigoplus_{\mu_k \equiv \mu \pmod{\mathbb{Z}}} M_k/M_{k+1}. \quad (3.14)$$

We define a \mathbb{Z} -grading on N^μ by setting the degree of $[s]$ to be $\mu_k - \mu$ for $0 \neq [s] \in M_k/M_{k+1}$. As a direct corollary of Theorem 3.3, we know N^μ is a graded module over $\mathbb{C}[z_1, \dots, z_n]$.

The following is the main result of this subsection.

Theorem 3.10. *For each $\mu \in \{\mu_1, \dots, \mu_m\}$, N^μ is a finitely generated torsion-free module over $\mathbb{C}[z_1, \dots, z_n]$.*

Before proving this we need some preparation. We fix a given analytic tangent cone \mathcal{E}_∞ . Let $\{\sigma_{k,i}^j\}$ be the elements constructed as in the proof of Proposition 3.9. By passing to a subsequence, we may assume for each k and i , $\{\sigma_{k,i}^j\}$ strongly converges to a set of L^2 orthonormal homogeneous sections $\{\sigma_{k,i}^\infty\}$ of \mathcal{E}_∞ with degree μ_k . Suppose $\mu_k = \mu + e$ for $e \in \mathbb{Z}_{\geq 0}$. Then $\{\sigma_{k,i}^\infty\}$ can be viewed as sections in $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}_\infty^\mu(e))$, where $\underline{\mathcal{E}}_\infty^\mu$ is a direct summand of $\underline{\mathcal{E}}_\infty$ of slope μ .

Denote by S_k the \mathbb{C} -vector space spanned by $\{\sigma_{k,1}^\infty, \dots, \sigma_{k,n_k}^\infty\}$. It can be viewed as a subspace of $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}_\infty^\mu(e))$. By definition we have

$$\dim_{\mathbb{C}} S_k = n_k = \dim_{\mathbb{C}} M_k/M_{k+1}. \quad (3.15)$$

Define

$$N_\infty^\mu = \bigoplus_{\mu_k \equiv \mu \pmod{\mathbb{Z}}} S_k. \quad (3.16)$$

This is also a graded vector space over \mathbb{C} with natural grading given by $\mu_k - \mu$.

Proposition 3.11. N_∞^μ is a graded submodule of $\bigoplus_e H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}_\infty^\mu(e))$.

Proof. It suffices to show that for any k fixed, given any homogeneous polynomial $f \in \mathbb{C}[z_1, \dots, z_n]$ and $\sigma_{k,j}^\infty \in S_k$ for any j , we have $f\sigma_{k,j}^\infty \in S_{k'}$ where $\mu_{k'} = \mu_k + \deg(f)$. By assumption, $f\sigma_{k,i}^\infty$ is the limit of $f\sigma_{k,i}^j$ as $j \rightarrow \infty$. However, for each j we have

$$f\sigma_{k,i}^j - \sum_{i'=1}^{n_{k'}} a_{i'}^j \sigma_{k',i'}^j = s_{k'+1}^j \quad (3.17)$$

where $\mu_{k'} = \mu_k + \deg(f)$, $a_{i'}^j \in \mathbb{C}$, and $s_{k'+1}^j \in M_{k'+1}$.

We claim $s_{k'+1}^j$ converges to zero in L^2 and then the result follows. Otherwise, suppose the L^2 norm of $\{s_{k'+1}^j\}_j$ has a positive lower bound after passing to a subsequence. Then rescale by factors $A_j \leq C$ we can assume both sides of (3.17) have $L^2(B)$ norm exactly 1 for all j . Then it follows that $a_{i'}^j$ is also uniformly bounded for all i' . Passing to a subsequence we may assume $a_{i'}^j$ converges to $a_{i'}^\infty$ for all i' . Since $\sigma_{k,i}^j$ and $\sigma_{k',i'}^j$ converge strongly, and f is fixed, by Lemma 2.13 we then obtain that both sides of (3.17) strongly converge to a holomorphic section s_∞ on \mathcal{E}_∞ with $\|s_\infty\|_{L^2(B)} = 1$.

Notice by definition for each j , $s_{k'+1}^j$ comes from the rescaling of a holomorphic section defined over the fixed ball B . Then by Proposition 3.6 s_∞ has degree at least $\mu_{k'+1}$. On the other hand, we know $f\sigma_{k,j}^\infty$ is homogeneous of degree $\mu_{k'}$ and by definition the limit $\sigma_{k',i'}^\infty$ is also homogeneous of degree $\mu_{k'}$. This is a contradiction. \square

Definition 3.12. We define $\underline{\mathcal{N}}_\infty^\mu$ to be the subsheaf of $\underline{\mathcal{E}}_\infty^\mu$ generated by N_∞^μ .

In particular, $\underline{\mathcal{N}}_\infty^\mu$ is torsion-free, and for $k \gg 1$, $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{N}}_\infty^\mu(k))$ can be identified with S_k . But notice it depends on various choices made above.

Corollary 3.13. For $k \gg 1$, we have

$$S_{k'} = \sum_{l=1}^n z_l S_k,$$

where k' is such that $\mu_{k'} = \mu_k + 1$.

Proof. This follows from the exact sequence

$$0 \rightarrow \underline{\mathcal{F}} \rightarrow (\underline{\mathcal{N}}_\infty^\mu)^{\oplus n} \xrightarrow{(z_1, \dots, z_n)} \underline{\mathcal{N}}_\infty^\mu(1) \rightarrow 0 \quad (3.18)$$

for some sheaf $\underline{\mathcal{F}}$ on \mathbb{CP}^{n-1} . Tensoring with $\mathcal{O}(e)$, and noticing that for $e \gg 1$, $H^1(\mathbb{CP}^{n-1}, \mathcal{F}(e)) = 0$, we obtain the conclusion. \square

Now we prove Theorem 3.10.

Proof of Theorem 3.10. We first prove the torsion-free property, this follows directly from Corollary 3.5. Indeed, suppose $[s] \in N^\mu$ is non-zero, then we can write $[s] = \sum_{i \geq i_1} [s_i]$, where $[s_i] \in M_i/M_{i+1}$ and $s_{i_1} \neq 0$. For any nonzero $f \in \mathbb{C}[z_1, \dots, z_n]$, we write

$$f = \sum_{j \geq j_1} f_j,$$

where each f_j is homogeneous and $f_{j_1} \neq 0$. Let i' be the unique integer such that $\mu_{i'} - \mu_{i_1} = j_1$. Then by Corollary 3.5 we know $d(fs) = d(f_{j_1}s_{i_1}) = \mu_{i'}$, and the component $[f_{j_1}s_{i_1}]$ of $[fs]$ in $M_{i'}/M_{i'+1}$ is nonzero.

Now we prove N^μ is finitely generated. It suffices to show that for k large,

$$M_{k'}/M_{k'+1} = \sum_l z_l(M_k/M_{k+1}),$$

where k' is such that $\mu_{k'} = \mu_k + 1$.

Claim 3.14. $\dim_{\mathbb{C}} \sum_l z_l(M_k/M_{k+1}) \geq \dim_{\mathbb{C}} \sum_l z_l S_k$.

Given this Claim, using Corollary 3.13 we have

$$\begin{aligned} \dim_{\mathbb{C}} \sum_l z_l(M_k/M_{k+1}) &\geq \dim_{\mathbb{C}} \sum_l z_l S_k \\ &= \dim_{\mathbb{C}} S_{k'} \\ &= \dim_{\mathbb{C}} M_{k'}/M_{k'+1} \\ &\geq \dim_{\mathbb{C}} \sum_k z_l(M_k/M_{k+1}), \end{aligned}$$

where the third equation follows by (3.15). This forces

$$\dim_{\mathbb{C}} \sum_k z_l(M_k/M_{k+1}) = \dim_{\mathbb{C}} M_{k'}/M_{k'+1}.$$

Combining this with the fact that

$$\sum_k z_l(M_k/M_{k+1}) \subset M_{k'}/M_{k'+1},$$

we obtain

$$\sum_k z_l(M_k/M_{k+1}) = M_{k'}/M_{k'+1}.$$

This finishes the proof of Theorem 3.10. □

Proof of Claim 3.14. Let $\{\sigma_{\infty}^{k',i}\}$ be an L^2 orthonormal basis for $S_{k'}$ constructed as above. Since we know

$$\sum_l z_l S_k = S_{k'},$$

there exists a sequence of sections $\tau_{k',i}^j$ in the \mathbb{C} -linear span of $z_l \sigma_{k,i}^j$ which converge strongly to $\sigma_{k',i}^{\infty}$. In particular, we have

$$\lim_{j \rightarrow \infty} \int_B \langle \tau_{k',i}^j, \tau_{k',l}^j \rangle = \delta_{il} \quad (3.19)$$

for all $1 \leq i, l \leq n_{k'}$.

It suffices to show that $\{\tau_{k',i}^j : 1 \leq i \leq n_{k'}\}$ are linearly independent in $M_{k'}/M_{k'+1}$ for j large. We argue by contradiction. Otherwise by passing to a subsequence we can assume for j large there are constants $a_i^j \in \mathbb{C}$, with

$$\sum_i a_i^j \tau_{k',i}^j = s^j \in M_{k'+1}.$$

We normalize $\|s^j\|_{L^2(B)} = 1$ which implies a_i^j are all uniformly bounded in j since $\{\tau_{k+1,i}^j\}$ are approximately L^2 -orthonormal for j large enough. In particular, passing to a subsequence we can assume $\{a_i^j : 1 \leq i \leq n_{k'}\}$ converge to $\{a_i^{\infty} : 1 \leq i \leq n_{k'}\}$ and there exists some i such that a_i^{∞} is nonzero. By Lemma 2.13, we can assume s^j converges strongly to some non-zero holomorphic section $s^{\infty} = \sum_i a_i^{\infty} \sigma_{k',i}^{\infty}$. In particular $d(s^{\infty}) = \mu_{k'}$. On the other hand, Proposition 3.6 implies that $d(s^{\infty}) \geq \mu_{k'+1}$. This is a contradiction. \square

Theorem 3.10 allows us to make the following

Definition 3.15. For $\mu \in \{\mu_1, \dots, \mu_m\}$, we define $\underline{\mathcal{E}}^{\mu}$ to be the torsion-free sheaf on \mathbb{CP}^{n-1} associated to the module N^{μ} .

By definition, for k large $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}^{\mu}(k))$ can be identified with M_k/M_{k+1} . On the other hand, we also know that by definition for k large $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{N}}_{\infty}^{\mu}(k))$ can be identified with S_k . So by (3.15) we have

$$\dim_{\mathbb{C}} H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{E}}^{\mu}(k)) = \dim_{\mathbb{C}} H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{N}}_{\infty}^{\mu}(k)). \quad (3.20)$$

It follows that $\underline{\mathcal{E}}^{\mu}$ and $\underline{\mathcal{N}}_{\infty}^{\mu}$ have the same Hilbert polynomial.

Now we recall the asymptotic Riemann-Roch theorem (see Page 189 in [14])

Lemma 3.16 (Asymptotic Riemann-Roch Theorem). *Let $\underline{\mathcal{F}}$ be a torsion-free coherent sheaf over \mathbb{CP}^{n-1} . Then*

$$\chi(\underline{\mathcal{F}}(k)) = r \cdot \frac{k^{n-1}}{(n-1)!} + r(\mu(\underline{\mathcal{F}}) + \frac{n}{2}) \frac{k^{n-2}}{(n-2)!} + O(k^{n-3}).$$

where χ denotes the holomorphic Euler characteristic, r denotes the rank of $\underline{\mathcal{F}}$, and $\mu(\underline{\mathcal{F}})$ denotes the slope of $\underline{\mathcal{F}}$.

An immediate consequence is

Corollary 3.17. *We have*

$$\text{rank}(\underline{\mathcal{E}}^\mu) = \text{rank}(\underline{\mathcal{N}}_\infty^\mu) \quad (3.21)$$

and

$$\mu(\underline{\mathcal{E}}^\mu) = \mu(\underline{\mathcal{N}}_\infty^\mu). \quad (3.22)$$

Remark 3.18. In the above discussion we work on a fixed analytic tangent cone \mathcal{E}_∞ , but it is clear that given an analytic tangent cone, by passing to a further subsequence one can extract the a subsheaf $\underline{\mathcal{N}}_\infty^\mu$ of $\underline{\mathcal{E}}_\infty^\mu$ as above. Here “passing to a further subsequence” is necessary in general since we need the convergence of the chosen holomorphic sections. Notice $\underline{\mathcal{E}}^\mu$ does not depend on the choice of the analytic tangent cone, and the equalities above hold for all such $\underline{\mathcal{N}}_\infty^\mu$.

Now we realize the above defined sheave $\underline{\mathcal{E}}^\mu$ naturally as a factor of the graded sheaf associated to a filtration of sheaf on \mathbb{CP}^{n-1} . For $q \in \mathbb{Z}_{\geq 0}$, $i \geq 1$, we denote

$$M_q^{an,i} := \{s \in H^0(B, \mathcal{E}) | d(s) \geq \mu_i + q\}.$$

For $i = 1, \dots, m$, we then obtain a filtration of $H^0(B, \mathcal{E})$ given by $\{M_q^{an,i}\}_{q \geq 0}$. We denote the associated graded module by

$$N^{an,i} := \bigoplus_{q \geq 0} M_q^{an,i} / M_{q+1}^{an,i}.$$

For $i = 1, \dots, m$ and $l = 0, \dots, m$, we also denote the graded module

$$N_l^{an,i} := \bigoplus_{q \geq 0} M_q^{an,m-l+i} / M_{q+1}^{an,i}.$$

Then we obtain a filtration of graded modules

$$0 = N_0^{an,i} \subset N_1^{an,i} \subset \dots \subset N_m^{an,i} = N^{an,i}.$$

By definition we also know that for $l \geq 1$ the quotient module $N_l^{an,i} / N_{l-1}^{an,i}$ is isomorphic to the graded module $N^{\mu[m-l+i]}$ (given by (3.14)) except possibly the lowest degree component. Here we use the notation that

$$[m-l+i] := \begin{cases} m-l+i & \text{if } i \leq l; \\ -l+i & \text{if } i > l. \end{cases}$$

Then by Theorem 3.10 and a simple induction on l it follows that all the graded modules $N_l^{an,i}$ are finitely generated and torsion-free. This enables us to make the following

Definition 3.19. For $i = 1, \dots, m$, we define $\underline{\mathcal{E}}^{an,i}$ to be the torsion-free coherent sheaf on \mathbb{CP}^{n-1} associated to the graded module $N^{an,i}$.

Definition 3.20. For $i, l = 1, \dots, m$ we define $\underline{\mathcal{E}}_l^i$ to be the torsion-free coherent sheaf on \mathbb{CP}^{n-1} associated to the graded module $N_l^{an,i}$.

By definition we have a natural filtration of sheaves given by

$$0 = \underline{\mathcal{E}}_0^i \subset \underline{\mathcal{E}}_1^i \subset \dots \subset \underline{\mathcal{E}}_m^i = \underline{\mathcal{E}}^{an,i}. \quad (3.23)$$

Furthermore, for $i, l = 1, \dots, m$ we have the isomorphism

$$\underline{\mathcal{E}}_l^i / \underline{\mathcal{E}}_{l-1}^i \simeq \underline{\mathcal{E}}^{\mu_{[m-l+i]}}. \quad (3.24)$$

Here we use the fact that two finite generated graded module over $\mathbb{C}[z_1, \dots, z_n]$ define isomorphic coherent sheaves on \mathbb{CP}^{n-1} if and only if they are isomorphic in sufficiently large degrees (see Exercise 5.9 in [12]).

From the above definition we see that the sheaves $\underline{\mathcal{E}}^{an,i}$ are put into equal footing. In our later discussion when we prove properties of these sheaves we will often restrict to the case $i = 1$, and the other cases are just the same up to change of notation. To make notational convenience we also set the following

Notation 3.21. We make the convention that when we omit the upper script i , we always mean $i = 1$. So in particular $\underline{\mathcal{E}}^{an} := \underline{\mathcal{E}}^{an,1}$, $M_k^{an} := M_k^{an,1}$, etc.

Notice by definition the sheaves $\underline{\mathcal{E}}^{an,i}$ on $D = \mathbb{CP}^{n-1}$ depend only on the degree function d , and do not depend on choice of analytic tangent cones. However at this point we can not say much about the geometric properties of either $\underline{\mathcal{E}}^{an,i}$, or the filtration (3.23). We have only compared the dimension of the space of sections of the quotients associated to the filtration with that of a subsheaf of any analytic tangent cone $\underline{\mathcal{E}}_\infty$. Also the construction of $\underline{\mathcal{N}}_\infty^\mu$ depends not only on the analytic tangent cone \mathcal{E}_∞ , but also on the choice of holomorphic sections $\{s_{k,i}\}$ at the beginning of this subsection, and it is not a priori clear why this is an intrinsic object.

In the remainder of this section, we will show that $\underline{\mathcal{E}}^{an,i}$ is an optimal algebraic tangent cone, i.e., it is isomorphic to the restriction to D of some optimal extension of $p^*(\mathcal{E})|_{\widehat{B} \setminus D}$ across D and the filtration in (3.23) is precisely the Harder-Narasimhan filtration of $\underline{\mathcal{E}}^{an,i}$.

3.3 The main construction

The main goal of this subsection is to prove

Theorem 3.22. For $i = 1, \dots, m$, there exists a reflexive sheaf $\widehat{\mathcal{E}}^i$ on \widehat{B} , such that $\widehat{\mathcal{E}}^i|_{\widehat{B} \setminus D}$ is isomorphic to $p^*\mathcal{E}|_{\widehat{B} \setminus D}$, $\widehat{\mathcal{E}}^i|_D$ is isomorphic to $\underline{\mathcal{E}}^{an,i}$, and $H^0(\widehat{B}, \widehat{\mathcal{E}}^i(-kD))$ is naturally identified with $M_k^{an,i}$ for $k \gg 1$.

Remark 3.23. A priori from the proof below the construction of $\widehat{\mathcal{E}}^i$ depends on various choices, but later in the next subsection we shall prove each such $\widehat{\mathcal{E}}^i$ is an optimal extension, so is indeed unique up to isomorphism by Theorem 2.15.

In the following we shall only prove the case $i = 1$, and the arguments for $i \neq 1$ are similar. So we shall omit the superscript i throughout this subsection.

The main difficulty in proving such a statement is that we are working on a mixed situation between algebraic geometry and complex analytic geometry. The exceptional divisor D is algebraic so we can describe sheaves over D in terms of graded modules as in the last subsection. On the other hand, \widehat{B} is not algebraic so it seems not easy to describe $\widehat{\mathcal{E}}$ in terms of purely algebraic objects. To overcome this issue we define an auxiliary sheaf first and then define $\widehat{\mathcal{E}}$ as a subsheaf.

For our purpose, we need the following vanishing theorem of A. Fujiki (see Theorem N' in [8])

Lemma 3.24. *Given any coherent analytic sheaf $\widehat{\mathcal{F}}$ over \widehat{B} , $H^1(\widehat{B}, \widehat{\mathcal{F}}(-kD)) = 0$ for $k \gg 1$.*

As a direct corollary of this, we have

Lemma 3.25. *Given any coherent sheaf $\widehat{\mathcal{F}}$ over \widehat{B} , $\widehat{\mathcal{F}}|_D$ is isomorphic to the sheaf associated to the graded module*

$$\bigoplus_{k \geq 0} H^0(\widehat{B}, \widehat{\mathcal{F}}(-kD)) / H^0(\widehat{B}, \widehat{\mathcal{F}}(-(k+1)D)).$$

Proof. It is a general result in algebraic geometry (see Proposition 5.15 in [12]) that $\widehat{\mathcal{F}}|_D$ is the sheaf associated to the graded module

$$\bigoplus_{k \geq 0} H^0(D, \widehat{\mathcal{F}}|_D(k)).$$

By Exercise 5.9 in [12], it suffices to show that for $k \gg 1$

$$H^0(D, \widehat{\mathcal{F}}|_D(k)) = H^0(\widehat{B}, \widehat{\mathcal{F}}(-kD)) / H^0(\widehat{B}, \widehat{\mathcal{F}}(-(k+1)D)).$$

To see this, we use the natural short exact sequence

$$0 \rightarrow \widehat{\mathcal{F}}(-(k+1)D) \rightarrow \widehat{\mathcal{F}}(-kD) \rightarrow \widehat{\mathcal{F}}|_D(k) \rightarrow 0,$$

and the fact that $H^1(\widehat{B}, \widehat{\mathcal{F}}(-(k+1)D)) = 0$ for $k \gg 1$ which follows from Lemma 3.24. \square

Now we fix a short exact sequence given in (3.8):

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{O}^{\oplus n_1} \xrightarrow{\rho} \mathcal{O}^{\oplus n_2}. \quad (3.25)$$

We denote by

$$R := \mathcal{O}(B)$$

the ring of holomorphic functions over B , and we denote by \mathfrak{m} the maximal ideal of R consisting of those functions vanishing at 0. Notice R is not Noetherian. Pulling back (3.25) to \widehat{B} , we have

$$0 \rightarrow \widehat{\mathcal{E}}_0 \rightarrow \mathcal{O}^{\oplus n_1} \xrightarrow{\widehat{\rho}} \mathcal{O}^{\oplus n_2}, \quad (3.26)$$

where we define $\widehat{\mathcal{E}}_0$ to be the kernel of $\widehat{\rho}$.

Following the discussion in Section 3.1, we define for $k \in \mathbb{Z}_{\geq 0}$,

$$M_k^0 := \{s \in H^0(B, \mathcal{E}) : d^0(s) \geq k\},$$

where d^0 is the degree function defined with respect to the fixed induced metric H_0 . Then $\{M_k^0\}_k$ forms an \mathfrak{m} -filtration of $H^0(B, \mathcal{E})$, that is to say,

$$\mathfrak{m} \cdot M_k^0 \subset M_{k+1}^0$$

for all k . Similarly by previous discussion we know $\{M_k^{an}\}_k$ also forms an \mathfrak{m} -filtration of $H^0(B, \mathcal{E})$. Since $d(s) \leq d_0(s)$ for all s , we have $M_k^{an} \subset M_l^0$ for $l \leq \mu_1 + k$.

We define the blowup ring

$$\widehat{R} := \bigoplus_{k \geq 0} \mathfrak{m}^k$$

and the graded modules over \widehat{R} given by

$$\widehat{M}^0 := \bigoplus_{k \geq 0} M_k^0,$$

and

$$\widehat{M}^{an} := \bigoplus_{k \geq 0} M_k^{an}. \quad (3.27)$$

Now we make the identification

$$M_k^0 = H^0(\widehat{B}, \text{Ker}(\widehat{\rho})(-kD))$$

as follows. Given $s \in M_k^0$, then we can view s via (3.25) as a vector-valued holomorphic function over B which has vanishing order at least k at 0. Then p^*s is a section of $\text{Ker}(\widehat{\rho})$ with vanishing order at least k along D . It is easy to see that the converse also holds by Hartog's extension theorem.

Lemma 3.26. $\{M_k^0\}_k$ is a stable \mathfrak{m} -filtration of $H^0(B, \mathcal{E})$, i.e.,

$$M_{k+1}^0 = \mathfrak{m} \cdot M_k^0$$

for $k \gg 1$.

Proof. It suffices to show $M_{k+1}^0 \subset \mathfrak{m} \cdot M_k^0$. Let $\{z_1, \dots, z_n\}$ denote the coordinate functions on B . Then $\{p^* z_i\}_i$ forms a set of global generators of $\mathcal{O}(-D)$. In particular, we have an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \text{Ker}(\hat{\rho})^{\oplus n} \xrightarrow{(p^* z_1, \dots, p^* z_n)} \text{Ker}(\hat{\rho})(-D) \rightarrow 0.$$

Tensoring with $\mathcal{O}(-kD)$, we have the following exact sequence

$$0 \rightarrow \mathcal{F}(-kD) \rightarrow (\text{Ker}(\hat{\rho})(-kD))^{\oplus n} \xrightarrow{(p^* z_1, \dots, p^* z_n)} \text{Ker}(\hat{\rho})(-(k+1)D) \rightarrow 0.$$

By Lemma 3.24 above, we have a surjective map

$$(H^0(\hat{B}, (\text{Ker}(\hat{\rho})(-kD))))^{\oplus n} \xrightarrow{(p^* z_1, \dots, p^* z_n)} H^0(\hat{B}, \text{Ker}(\hat{\rho})(-(k+1)D))$$

for k large. In particular, by the identification above we know $M_{k+1}^0 \subset \mathfrak{m} \cdot M_k^0$ and thus $M_{k+1}^0 = \mathfrak{m} \cdot M_k^0$. \square

Proposition 3.27. $\{M_k^{an}\}_k$ is a stable \mathfrak{m} -filtration of $H^0(B, \mathcal{E})$.

Proof. By Theorem 3.10, we know that for k large,

$$\mathfrak{m} \cdot M_k^{an} \subset M_{k+1}^{an} \subset \mathfrak{m} \cdot M_k^{an} + M_{k+2}^{an}.$$

So it suffices to show for any fixed k large, there exists $l_0 = l_0(k)$ such that

$$M_l^{an} \subset \mathfrak{m} \cdot M_k^{an}$$

for $l \geq l_0$. Since

$$M_l^{an} \subset M_l^0 = \mathfrak{m} \cdot M_{l-1}^0,$$

it suffices to show $M_{l-1}^0 \subset M_k^{an}$ for $l \gg 1$. Notice by definition we can find l' large such that

$$\mathfrak{m}^{l'} \cdot H^0(B, \mathcal{E}) \subset M_k^{an}.$$

By Lemma 3.26 we may also assume $M_{l+1}^0 = \mathfrak{m} \cdot M_l^0$ for all $l \geq l'$. Now for $l > 2l' + 1$, we have

$$M_{l-1}^0 = \mathfrak{m}^{l'} \cdot M_{l-1-l'}^0 \subset \mathfrak{m}^{l'} \cdot M_0^0 \subset M_k^{an}.$$

\square

Proposition 3.28. For all k , M_k^0 and M_k^{an} are finitely generated R modules.

Proof. We first show $H^0(B, \mathcal{E})$ is finitely generated over R . By assumption, we know \mathcal{E} is globally generated, so there exists a finite global resolution of \mathcal{E} by sections $s_1, \dots, s_N \in H^0(B, \mathcal{E})$ given as

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{O}^{\oplus N} \rightarrow \mathcal{E} \rightarrow 0 \quad (3.28)$$

for some coherent sheaf \mathcal{F} . Since B is Stein, we know $H^1(B, \mathcal{F}) = 0$ and thus we have a surjective map

$$H^0(B, \mathcal{O})^{\oplus N} \xrightarrow{(s_1, \dots, s_N)} H^0(B, \mathcal{E}) \rightarrow 0.$$

In particular, we know $H^0(B, \mathcal{E})$ is finitely generated over R .

Now we consider M_k^0 . Let $\mathcal{E}^k \subset \mathcal{E}$ be the coherent subsheaf generated by M_k^0 in \mathcal{E} . By definition, for all k , we have

$$\mathfrak{m}^k \cdot H^0(B, \mathcal{E}) \subset M_k^0.$$

This implies that $\mathcal{I}_0^k \cdot \mathcal{E} \subset \mathcal{E}^k$. In particular, $\{z_i^k s_j : 1 \leq i \leq n, 1 \leq j \leq N\}$ globally generates \mathcal{E}^k away from 0. Now take finitely many sections of M_k^0 which generate the stalk of \mathcal{E}^k at 0, then combined with $\{z_i^k s_j : 1 \leq i \leq n, 1 \leq j \leq N\}$, they globally generate \mathcal{E}^k . So it follows that for any section s of \mathcal{E}^k we have $d^0(s) \geq k$. Therefore $H^0(M, \mathcal{E}^k) = M_k^0$. It follows from the above argument using vanishing of H^1 that M_k^0 is finitely generated over R .

Again the same argument also works with M_k^{an} , noticing that for all k , $M_k^{an} \supset \mathfrak{m}^l \cdot H^0(B, \mathcal{E})$ for $l = k + \mu_1 + 1$. \square

An immediate corollary is

Corollary 3.29. *Both \widehat{M}^0 and \widehat{M}^{an} are finitely generated over \widehat{R} .*

Proof. This follows from Proposition 5.3 in Eisenbud [7]. Notice this result does not require the ring \widehat{R} to be Noetherian. \square

Now we finish the proof of Theorem 3.22. We divide it into a few pieces.

Existence:

By Corollary 3.29, \widehat{M}^{an} is finitely generated over \widehat{R} . We can choose a set of finitely many homogeneous generators $\{[s_{k_i}] \in M_{k_i}^{an}/M_{k_i+1}^{an}\}_{i=1}^l$, and the corresponding representatives $s_{k_i} \in M_{k_i}^{an} \setminus M_{k_i+1}^{an}$. In particular, we have

$$d(s_{k_i}) = k_i + \mu_q$$

for some $q \in \{1, \dots, m\}$. Given $s \in M_{k_i}^{an}$, we know

$$d^0(s) \geq d(s) \geq k_i + \mu_1 \geq k_i,$$

thus $s \in M_{k_i}^0$. In particular, we have

$$M_{k_i}^{an} \subset M_{k_i}^0$$

and

$$s_{k_i} \in H^0(\widehat{B}, \text{Ker}(\widehat{\rho})(-k_i D)),$$

hence s_{k_i} defines a map $\mathcal{O}(k_i D) \rightarrow \text{Ker}(\widehat{\rho})$. We define $\widehat{\mathcal{E}}$ to be the image sheaf of the natural map

$$\bigoplus_{i=1}^l \mathcal{O}(k_i D) \xrightarrow{(s_{k_1}, \dots, s_{k_l})} \text{Ker}(\widehat{\rho})$$

In particular, $\widehat{\mathcal{E}}$ is coherent and it lies in the following exact sequence

$$0 \rightarrow \widehat{\mathcal{F}} \rightarrow \bigoplus_{i=1}^l \mathcal{O}(k_i D) \rightarrow \widehat{\mathcal{E}} \rightarrow 0, \quad (3.29)$$

for some sheaf $\widehat{\mathcal{F}}$.

Global sections:

We show that $H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) = M_k^{an}$ for k sufficiently large. Notice both are naturally subspaces of $H^0(B, \mathcal{E})$. We first show that

$$H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) \subset M_k^{an}.$$

By using the short exact sequence (3.29), since $H^1(\widehat{B}, \widehat{\mathcal{F}}(-kD)) = 0$ for $k \gg 1$, we have a surjective map

$$\bigoplus_{i=1}^l H^0(\widehat{B}, \mathcal{O}((k_i - k)D)) \xrightarrow{(s_{k_1}, \dots, s_{k_l})} H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) \rightarrow 0$$

for k large. In particular, for any section $s \in H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD))$,

$$s = \sum_i f_{k-k_i} \cdot s_{k_i}$$

for some $f_{k-k_i} \in \mathfrak{m}^{k-k_i}$. This implies

$$d(s) \geq \mu_1 + k,$$

so $s \in M_k^{an}$. It remains to show that $M_k^{an} \subset H^0(\widehat{B}, \widehat{\mathcal{E}}(-k))$. By our choices of $\{s_{k_i}\}$ it follows that given any $s' \in M_k^{an}$, we can write

$$s' = \sum_i g_{k-k_i} \cdot s_{k_i}$$

where $g_{k-k_i} \in \mathfrak{m}^{k-k_i}$. In particular, we know $s' \in H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD))$.

Restriction:

$\widehat{\mathcal{E}}|_D = \underline{\mathcal{E}}^{an}$ this follows from Lemma 3.25 and the previous item.

Reflexivity:

By definition $\widehat{\mathcal{E}}$ is a subsheaf of $\text{Ker}(\widehat{\rho})$ and we have a natural inclusion

$$0 \rightarrow \widehat{\mathcal{E}}^{**}/\widehat{\mathcal{E}} \rightarrow \text{Ker}(\widehat{\rho})/\widehat{\mathcal{E}}.$$

The above inclusion induces an inclusion of global sections

$$0 \rightarrow H^0(\widehat{B}, (\widehat{\mathcal{E}}^{**}/\widehat{\mathcal{E}})(-kD)) \rightarrow H^0(\widehat{B}, (\text{Ker}(\widehat{\rho})/\widehat{\mathcal{E}})(-kD))$$

for any k . By the discussion above, we have

$$H^0(\widehat{B}, \text{Ker}(\widehat{\rho})(-kD)) = M_k^0$$

and

$$H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) = M_k^{an}$$

for k large. In particular, by Lemma 3.24 we have

$$H^0(\widehat{B}, (\text{Ker}(\widehat{\rho})/\widehat{\mathcal{E}})(-kD)) = M_k^0/M_k^{an}$$

for k large. Similarly,

$$H^0(\widehat{B}, (\widehat{\mathcal{E}}^{**}/\widehat{\mathcal{E}})(-kD)) = H^0(\widehat{B}, \widehat{\mathcal{E}}^{**}(-kD))/M_k^{an}. \quad (3.30)$$

Claim 3.30. *For any fixed $k \gg 1$, given any $s \in H^0(\widehat{B}, \widehat{\mathcal{E}}^{**}(-kD))$, there exists a homogeneous polynomial function P over \mathbb{C}^n so that*

$$p^*P \cdot s \in H^0(\widehat{B}, \mathcal{I}_D^d \cdot \widehat{\mathcal{E}}(-kD)),$$

where $d = \deg(P)$, and \mathcal{I}_D denotes the ideal sheaf of D on \widehat{B} .

Given this, fix any $s \in H^0(\widehat{B}, \widehat{\mathcal{E}}^{**}(-kD))$, we have

$$p^*P \cdot s \in H^0(\widehat{B}, \mathcal{I}_D^d \cdot \widehat{\mathcal{E}}(-kD)) = M_{k+d}^{an}.$$

Notice s naturally induces a section of \mathcal{E} on B , and we have then

$$d(P \cdot s) \geq \mu_1 + k + d.$$

Thus by Corollary 3.5 we get

$$d(s) \geq \mu_1 + k,$$

hence $s \in M_k^{an}$. By (3.30), for any k large, we have

$$H^0(\widehat{B}, (\widehat{\mathcal{E}}^{**}/\widehat{\mathcal{E}})(-kD)) = 0.$$

This implies $\widehat{\mathcal{E}} = \widehat{\mathcal{E}}^{**}$, in other words, $\widehat{\mathcal{E}}$ is reflexive. This finishes the proof of Theorem 3.22.

Proof of Claim 3.30. Let $\tau_k = (\widehat{\mathcal{E}}^{**}/\widehat{\mathcal{E}})(-kD)$. Since $\widehat{\mathcal{E}}$ is torsion-free, the support $V \subset D \subset \widehat{B}$ of τ_k is of complex codimension at least 2 in \widehat{B} . We have the following observation

At any point $z \in V$, there exists a meromorphic function over \widehat{B} of the form

$$f = \frac{p^*P}{p^*Q}$$

which vanishes at z and $f \cdot (\tau_k)_z = 0$, where $(\tau_k)_z$ denotes the stalk of τ_k at z , and P, Q are homogeneous polynomials on \mathbb{C}^n .

In particular, we see

$$(p^*P \cdot s)_z \in (\mathcal{I}_D^{\deg(P)} \cdot \widehat{\mathcal{E}}(-kD))_z,$$

Since V is compact, we can find finitely many such p^*P_1, \dots, p^*P_l so that for any $z \in D$, there exists some P_i so that

$$(p^*P_i \cdot s)_z \in (\mathcal{I}_D^{d_i} \cdot \widehat{\mathcal{E}}(-kD))_z.$$

Here $d_i = \deg(P_i)$. If we denote $\widehat{P} = p^*(P_1 \cdots P_l)$, it follows that for any $z \in D$, there exists some i such that

$$\begin{aligned} (\widehat{P} \cdot s)_z &\in p^*(P_1 \cdots P_{i-1} P_{i+1} \cdots P_l)_z (p^*P_i \cdot \widehat{\mathcal{E}}(-kD))_z \\ &\subset p^*(P_1 \cdots P_{i-1} P_{i+1} \cdots P_l)_z (\mathcal{I}_D^{d_i} \cdot \widehat{\mathcal{E}}(-kD))_z \\ &\subset (\mathcal{I}_D^d \cdot \widehat{\mathcal{E}}(-kD))_z. \end{aligned}$$

Here $d = d_1 + \cdots + d_l$. This finishes the proof of Claim 3.30.

Now we justify the observation above. By definition, there exists some N so that the annihilator ideal sheaf $\text{Ann}(\tau_k)$ satisfies

$$(\mathcal{I}_V)^N \subset \text{Ann}(\tau_k) \subset \mathcal{I}_V,$$

where \mathcal{I}_V denotes the ideal sheaf of V on \widehat{B} . It suffices to show that there exists a meromorphic function over \widehat{B} of the form $f = \frac{p^*P}{p^*Q}$ so that f vanishes along V near z . To see this, we let $\mathcal{I}_{V,D}$ denote the ideal sheaf associated to V in D . We know $\mathcal{I}_{V,D}(l)$ is globally generated for l large. Fix such an l , since V has complex codimension at least 1 in D , there exists some $P' \in H^0(D, \mathcal{I}_{V,D}(l))$ and $Q' \in H^0(D, \mathcal{O}(l))$ so that P' is not identically zero and $Q'(z) \neq 0$. Then $\phi^*(\frac{P'}{Q'})$ will be what we need. Here $\phi : \widehat{B} \rightarrow D$ denote the restriction of the projection map $\mathcal{O}(-1) \rightarrow D$. Indeed, it is direct to check that $(\phi^*(\frac{P'}{Q'})) = p^*(\frac{P}{Q})$ where P, Q are the homogeneous polynomials on \mathbb{C}^n corresponding to P' and Q' respectively. \square

For each $i = 1, \dots, m$, we define the sheaf on D

$$\underline{\mathcal{T}}^{\mu_i} := \underline{\mathcal{E}}_{\infty}^{\mu_i} / \underline{\mathcal{N}}_{\infty}^{\mu_i}$$

and we denote

$$\underline{\mathcal{T}} := \bigoplus_{i=1}^m \underline{\mathcal{T}}^{\mu_i}.$$

Corollary 3.31. *$\underline{\mathcal{T}}$ is a torsion sheaf. Moreover, for all $i = 1, \dots, m$, we have*

$$\mu(\underline{\mathcal{E}}^{\mu_i}) \leq \mu_i,$$

and the equality holds if and only if the support of $\underline{\mathcal{T}}^{\mu_i}$ has complex codimension at least 2.

Proof. By Lemma 3.16 and Corollary 3.17, we already know that

$$\text{rank}(\underline{\mathcal{E}}^{\mu_i}) = \text{rank}(\underline{\mathcal{N}}_{\infty}^{\mu_i}) \leq \text{rank}(\underline{\mathcal{E}}_{\infty}^{\mu_i}).$$

On the other hand, by Theorem 3.22 and the filtration (3.23), we have

$$\sum_{i=1}^m \text{rank}(\underline{\mathcal{E}}^{\mu_i}) = \text{rank}(\underline{\mathcal{E}}^{an}) = \text{rank}(\mathcal{E}) \geq \sum_{i=1}^m \text{rank}(\underline{\mathcal{E}}_{\infty}^{\mu_i}).$$

This forces the inequality above to be an equality, so $\underline{\mathcal{T}}$ is a torsion sheaf. By definition, we also have

$$0 \rightarrow \underline{\mathcal{N}}_{\infty}^{\mu_i} \rightarrow \underline{\mathcal{E}}_{\infty}^{\mu_i} \rightarrow \underline{\mathcal{T}}_{\infty}^{\mu_i} \rightarrow 0,$$

which implies

$$c_1(\underline{\mathcal{E}}_{\infty}^{\mu_i}) = c_1(\underline{\mathcal{N}}_{\infty}^{\mu_i}) + c_1(\underline{\mathcal{T}}_{\infty}^{\mu_i}).$$

Let V be the closure of the codimension 1 part of the support of $\underline{\mathcal{T}}_{\infty}^{\mu_i}$. We know $c_1(\underline{\mathcal{T}}_{\infty}^{\mu_i})$ is equal to the Poincaré dual of V (see Proposition 3.1 in [17]). In particular, we have $\mu(\underline{\mathcal{E}}^{\mu_i}) \leq \mu_i$ and the equality holds if and only if $c_1(\underline{\mathcal{T}}_{\infty}^{\mu_i}) = 0$, i.e., the support of $\underline{\mathcal{T}}^{\mu_i}$ has complex codimension at least 2. \square

Remark 3.32. In particular, we also know that

$$\underline{\mathcal{E}}_{\infty} = \bigoplus_{i=1}^m \underline{\mathcal{E}}_{\infty}^{\mu_i}.$$

In other words, if we write

$$\underline{\mathcal{N}}_{\infty} := \bigoplus_{i=1}^m \underline{\mathcal{N}}_{\infty}^{\mu_i}$$

then

$$\mathcal{T} = \underline{\mathcal{E}}_\infty / \underline{\mathcal{N}}_\infty.$$

Furthermore, we know

$$\underline{\mathcal{N}}_\infty^{**} = \underline{\mathcal{E}}_\infty \quad (3.31)$$

which is an important fact to be used later.

In the following subsection, we will show that the support of \mathcal{T} has complex codimension at least 2, and as a consequence we see that $\widehat{\mathcal{E}}$ is an optimal extension.

3.4 Optimality

For $i = 1, \dots, m$, let $\widehat{\mathcal{E}}^i$ be the reflexive sheaf constructed in Theorem 3.22. The goal of this subsection is to prove

Theorem 3.33. *$\widehat{\mathcal{E}}^i$ is an optimal extension of \mathcal{E} at 0. Moreover, the Harder-Narasimhan filtration of $\underline{\mathcal{E}}^{an,i}$ is given by (3.23):*

$$0 = \underline{\mathcal{E}}_0^i \subset \underline{\mathcal{E}}_1^i \subset \dots \subset \underline{\mathcal{E}}_m^i = \underline{\mathcal{E}}^{an,i}, \quad (3.32)$$

and the associated graded sheaf $Gr^{HN}(\underline{\mathcal{E}}^{an,i})$ is isomorphic to $\bigoplus_{i=1}^m \underline{\mathcal{E}}^{\mu_i}$.

Let \mathcal{G}^{alg} and Σ_b^{alg} be defined in Corollary 2.21. Then an immediate consequence is

Corollary 3.34. *We have*

$$\mathcal{G}^{alg} = \psi_* \pi^* \bigoplus_{i=1}^m (Gr^{HNS}(\underline{\mathcal{E}}^{\mu_i}))^{**}, \quad (3.33)$$

and

$$\Sigma_b^{alg} = \sum_{i=1}^m \Sigma_b^{alg}(\underline{\mathcal{E}}^{\mu_i}). \quad (3.34)$$

To prove Theorem 3.33, again we only prove the case $i = 1$ and the other cases are similar. So we shall omit the superscript i throughout this subsection. By (3.24) we know for $l = 1, \dots, m$

$$\underline{\mathcal{E}}^l / \underline{\mathcal{E}}^{l-1} = \underline{\mathcal{E}}^{\mu_{m-l+1}}. \quad (3.35)$$

So Theorem 3.33 follows from the definition and the following two Propositions.

Proposition 3.35. *For $i = 1, \dots, m$, we have $\mu(\underline{\mathcal{E}}^{\mu_i}) = \mu_i$. In particular, the support of the torsion sheaf \mathcal{T} has codimension at least 2.*

Proposition 3.36. *For $i = 1, \dots, m$, $\underline{\mathcal{E}}^{\mu_i}$ is semi-stable.*

In the following we shall prove these two results. By Corollary 3.31 we already know $\mu(\underline{\mathcal{E}}^{\mu_i}) \leq \mu_i$. To prove Proposition 3.35 it suffices to prove the reversed inequality. The key idea is to make use of the dual sheaf \mathcal{E}^* , which is endowed with the induced HYM connection. We can apply the previous construction in this section to obtain a natural algebraic tangent cone of \mathcal{E}^* . We will show this algebraic tangent cone is naturally dual to the algebraic tangent cone $\widehat{\mathcal{E}}$ of \mathcal{E} , and applying 3.31 yields the desired reversed inequality. This duality should be viewed as a manifestation of the fact that taking dual is an intrinsic operation, both algebraically and analytically. Below we give the detailed arguments.

By construction we know $\widehat{\mathcal{E}}$ satisfies the following

- (1) $\widehat{\mathcal{E}}|_{\widehat{B} \setminus D} \simeq (p^*\mathcal{E})|_{\widehat{B} \setminus D}$, and $H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD))$ is naturally identified with $M_k^{an} \subset H^0(B, \widehat{\mathcal{E}})$ for k large;
- (2) there exist finitely many sections $s_i \in M_{k_i}^{an} \setminus M_{k_i+1}^{an}$, $i = 1, \dots, l$, which globally generate $\widehat{\mathcal{E}}$ in the following sense

$$\bigoplus_{1 \leq i \leq l} \mathcal{O}(k_i D) \xrightarrow{(s_1, \dots, s_l)} \widehat{\mathcal{E}} \rightarrow 0 \quad (3.36)$$

where s_i is naturally viewed as an element in $H^0(\widehat{B}, \widehat{\mathcal{E}}(-k_i D))$;

- (3) there exists a filtration of $\underline{\mathcal{E}}^{an} := \widehat{\mathcal{E}}|_D$

$$0 = \underline{\mathcal{E}}_0 \subset \underline{\mathcal{E}}_1 \subset \dots \subset \underline{\mathcal{E}}_m = \underline{\mathcal{E}}^{an}$$

so that $\mu(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i) \leq \mu_{m-i}$ and $\text{rank}(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i) = \text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{m-i}})$;

- (4) the support of the torsion sheaf $\underline{\mathcal{T}}$ has complex codimension at least 2 if and only if $\mu(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i) = \mu_{m-i}$ for any $1 \leq i \leq m$.

Similar construction applies to the dual admissible HYM connection A^* on $\mathcal{F} := \mathcal{E}^*$. Abusing notation, we still denote the degree function associated to \mathcal{F} by d . It is clear that given an analytic tangent cone \mathcal{E}_∞ of A at 0, then $\mathcal{F}_\infty := \mathcal{E}_\infty^*$ is an analytic tangent cone of A^* . It follows that for any nonzero section $s^* \in H^0(B, \mathcal{F})$, $d(s^*) \equiv -\mu_i \pmod{\mathbb{Z}}$ for some $i \in \{1, \dots, m\}$. For any $k' \in \mathbb{Z}$, we denote

$$L_{k'}^{an} = \{s^* \in H^0(B, \mathcal{F}) | d(s^*) \geq k' - \mu_1\}.$$

Then $\widehat{\mathcal{F}}$ satisfies

- (1') $\widehat{\mathcal{F}}|_{\widehat{B} \setminus D} \simeq (p^*\mathcal{F})|_{\widehat{B} \setminus D}$, and $H^0(\widehat{B}, \widehat{\mathcal{F}}(-k'D))$ is naturally identified with $L_{k'}^{an} \subset H^0(B, \widehat{\mathcal{F}})$ for all k' large;

- (2') there exist finitely many sections $s_{i'}^* \in L_{k_{i'}}^{an} \setminus L_{k_{i'}+1}^{an}$, $i' = 1, \dots, l'$, which globally generate $\widehat{\mathcal{F}}$ in the following sense

$$\bigoplus_{1 \leq i \leq l'} \mathcal{O}(k_{i'} D) \xrightarrow{(s_1^*, \dots, s_{l'}^*)} \widehat{\mathcal{F}} \rightarrow 0 \quad (3.37)$$

where $s_{i'}^*$ is naturally viewed as an element in $H^0(\widehat{B}, \widehat{\mathcal{F}}(-k_{i'} D))$;

- (3') There exists a filtration of $\underline{\mathcal{F}}^{an} := \widehat{\mathcal{F}}|_D$

$$0 = \underline{\mathcal{F}}_0 \subset \underline{\mathcal{F}}_1 \subset \dots \subset \underline{\mathcal{F}}_m = \underline{\mathcal{F}}^{an}$$

so that $\mu(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'}) \leq -\mu_{i'}$ with $\text{rank}(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'}) = \text{rank}(\underline{\mathcal{F}}_\infty^{-\mu_{i'}}) = \text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{i'}})$;

- (4') the support of the torsion sheaf $\underline{\mathcal{T}}'$ has complex codimension at least 2 if and only if $\mu(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'}) = -\mu_{i'}$ for any $1 \leq i' \leq m$.

Lemma 3.37. *We have*

$$\mu(\underline{\mathcal{E}}^{an}) + \mu(\underline{\mathcal{F}}^{an}) \leq 0,$$

and the equality holds if and only if the support of $\underline{\mathcal{T}}$ and $\underline{\mathcal{T}}'$ have complex codimension at least 2.

Proof. By (3) and (3') above, we have

$$\begin{aligned} & \mu(\underline{\mathcal{E}}^{an}) + \mu(\underline{\mathcal{F}}^{an}) \\ &= \sum_{i=1}^m \frac{\text{rank}(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i) \mu(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i)}{\text{rank}(\underline{\mathcal{E}})} + \sum_{i'=1}^m \frac{\text{rank}(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'}) \mu(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'})}{\text{rank}(\underline{\mathcal{E}})} \\ &= \sum_{i=1}^m \frac{\text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{m+1-i}}) \mu(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i)}{\text{rank}(\underline{\mathcal{E}})} + \sum_{i'=1}^m \frac{\text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{i'}}) \mu(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'})}{\text{rank}(\underline{\mathcal{E}})} \\ &\leq \sum_{i=1}^m \frac{\text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{m+1-i}}) \mu_{m+1-i}}{\text{rank}(\underline{\mathcal{E}})} - \sum_{i'=1}^m \frac{\text{rank}(\underline{\mathcal{E}}_\infty^{\mu_{i'}}) \mu_{i'}}{\text{rank}(\underline{\mathcal{E}})} \\ &= 0. \end{aligned}$$

The equality holds if and only if $\mu(\underline{\mathcal{E}}_{i+1}/\underline{\mathcal{E}}_i) = \mu_{m+1-i}$ and $\mu(\underline{\mathcal{F}}_{i'+1}/\underline{\mathcal{F}}_{i'}) = -\mu_{i'}$ for all $1 \leq i, i' \leq m$. By Corollary 3.31, this holds if and only if the support of $\underline{\mathcal{T}}$ and $\underline{\mathcal{T}}'$ has complex codimension at least 2. \square

Lemma 3.38. *There exists a natural paring*

$$L_{k'}^{an}/L_{k'+1}^{an} \times M_k^{an}/M_{k+1}^{an} \rightarrow \mathfrak{m}^{k+k'}/\mathfrak{m}^{k+k'+1}$$

which sends $([s_{k'}^*], [s_k])$ to $[s_{k'}^*(s_k)]$. Furthermore, this paring is non-degenerate for k and k' both large.

Proof. The existence of the paring follows from Theorem 3.3, Item (4). It remains to prove this paring is non-degenerate. Suppose there exists some nonzero $[s^*] \in L_{k'}^{an}/L_{k'+1}^{an}$ so that $s^*(t) \in \mathfrak{m}^{k'+k+1}$ for all $t \in M_k^{an}$ for k large. Let s_∞^* be the nonzero rescaled limit section of \mathcal{F}_∞ given by s^* and t_∞ be the nonzero rescaled limit section of \mathcal{E}_∞ given by t . Taking the limit of the fact $s^*(t) \in \mathfrak{m}^{k'+k+1}$, we know $s_\infty^*(t_\infty) = 0$. By Corollary 3.31, for k large, we know that such limiting homogeneous sections t_∞ generate the fiber of \mathcal{E}_∞ at a generic point of \mathbb{C}^n . In particular, we know s_∞^* has to vanish at a generic point hence $s_\infty^* = 0$. Contradiction. \square

Lemma 3.39. *There exists a natural paring $\widehat{\mathcal{E}} \otimes \widehat{\mathcal{F}} \rightarrow \mathcal{O}_{\widehat{B}}$ induced by the paring $\mathcal{E} \otimes \mathcal{F} \rightarrow \mathcal{O}_B$. Furthermore, this paring induces an isomorphism between $\widehat{\mathcal{E}}^*$ and $\widehat{\mathcal{F}}$.*

Proof. We first define the natural paring $\widehat{\mathcal{E}} \otimes \widehat{\mathcal{F}} \rightarrow \mathcal{O}_{\widehat{B}}$. This follows from (3.36) and (3.37). More precisely, we first define a paring

$$P : \bigoplus_i \mathcal{O}(k_i) \times \bigoplus_{i'} \mathcal{O}(k_{i'}) \rightarrow \mathcal{O}_{\widehat{B}}$$

which sends $((f_1 \cdots, f_l), (g_1 \cdots g_{l'}))$ to $\sum_{i,i'} f_i g_{i'} s_i^*(s_{i'})$. Here $s_i^*(s_{i'}) \in \mathfrak{m}^{k_i+k_{i'}}$ by Theorem 3.3, Item (4), and we naturally view it as a holomorphic function defined over \widehat{B} . In particular, $s_i^*(s_{i'})$ has a vanishing order at least $k_i + k_{i'}$ along D and thus $f_i g_{i'} s_i^*(s_{i'})$ is a well-defined local homomorphic function. It is now easy to see that this paring descends to be a paring between $\widehat{\mathcal{E}}$ and $\widehat{\mathcal{F}}$, which we also denote as P . Furthermore, by definition, away from D , this paring is naturally isomorphic to the paring between \mathcal{E} and \mathcal{F} over B^* .

Claim 3.40. *P induces a vector bundle isomorphism at a generic point when restricting to the exceptional divisor D .*

Given this claim, we know P induces an isomorphism between $\widehat{\mathcal{E}}$ and $\widehat{\mathcal{F}}$ away from a codimension 2 analytic subvariety of \widehat{B} . Since $\widehat{\mathcal{E}}$ and $\widehat{\mathcal{F}}$ are reflexive, P actually induces a global isomorphism. \square

Proof of Claim 3.40. For any k and k' , P induces a paring

$$\widehat{\mathcal{E}}(-kD) \times \widehat{\mathcal{F}}(-k'D) \rightarrow \mathcal{O}(-(k+k')D).$$

Let \underline{P} denote the induced paring on D

$$\underline{\mathcal{E}}^{an}(k) \times \underline{\mathcal{F}}^{an}(k') \rightarrow \mathcal{O}(k+k').$$

By definition, \underline{P} induces the paring of global sections

$$L_{k'}^{an}/L_{k'+1}^{an} \times M_k^{an}/M_{k+1}^{an} \rightarrow \mathfrak{m}^{k+k'}/\mathfrak{m}^{k+k'+1}$$

which sends $([s_{k'}^*], [s_k])$ to $[s_{k'}^*(s_k)]$. Indeed, this follows from the construction that

$$H^0(D, \underline{\mathcal{E}}^{an}(k)) = M_k^{an}/M_{k+1}^{an}$$

and

$$H^0(D, \underline{\mathcal{F}}^{an}(k')) = L_{k'}^{an}/L_{k'+1}^{an}$$

for k and k' large. Furthermore, this pairing is non-degenerate for k and k' large. Let

$$\underline{P}^* : \underline{\mathcal{E}}^{an}(k) \rightarrow (\underline{\mathcal{F}}^{an}(k'))^* \otimes \mathcal{O}((k' + k)D) = (\underline{\mathcal{F}}^{an})^*(k)$$

be the map induced by \underline{P} . We know that \underline{P}^* induces an injective map of global sections

$$\underline{P}^* : H^0(D, \underline{\mathcal{E}}^{an}(k)) \rightarrow H^0(D, (\underline{\mathcal{F}}^{an})^*(k))$$

for any k large. In particular, by Lemma 3.16, the rank of the image sheaf $\text{Im}(\underline{P}^*)$ has rank equal to $\text{rank}(\underline{\mathcal{E}}^{an})$ which implies \underline{P} induces an isomorphism at a generic point. \square

Proof of Proposition 3.35. By Lemma 3.39 we have $(\widehat{\mathcal{E}})^* \simeq \widehat{\mathcal{F}}$, so away from a codimension 2 subvariety in D , we have $(\underline{\mathcal{E}}^{an})^* \simeq \underline{\mathcal{F}}^{an}$. It follows that $\mu(\underline{\mathcal{E}}^{an}) + \mu(\underline{\mathcal{F}}^{an}) = 0$. Then the conclusion follows from Lemma 3.37 and Corollary 3.31. \square

Proof of Proposition 3.36. We argue by contradiction. Suppose $\underline{\mathcal{E}}^{\mu_i}$ is not semistable, then there exists a subsheaf $\underline{\mathcal{F}}$ of $\underline{\mathcal{E}}^{\mu_i}$ which is stable and satisfies $\mu(\underline{\mathcal{F}}) > \mu_i$. Take a basis $\{[s_k]\}_{k=1}^N$ of $H^0(\mathbb{CP}^{n-1}, \underline{\mathcal{F}}(i_0)) \subset M_{i'_0}/M_{i'_0+1}$ where $s_k \in M_{i'_0} \setminus M_{i'_0+1}$ for $k = 1, \dots, N$. Here i_0 is chosen large enough so that $\underline{\mathcal{F}}(i_0)$ is globally generated and $\mu_{i'_0} = i_0 + \mu_i$. In particular, we have an exact sequence

$$\mathcal{O}^{\oplus N} \xrightarrow{([s_1], \dots, [s_N])} \underline{\mathcal{F}}(i_0) \rightarrow 0.$$

Let

$$m_j = \max_k \|s_k\|_j.$$

Fix an analytic tangent cone \mathcal{E}_∞ , and passing to a further subsequence of $\{j\}$ if necessary we may assume $m_j^{-1} \cdot \lambda_j^* s_k$ converges strongly to a homogeneous section s_k^∞ of \mathcal{E}_∞ for all k , and at least one of the s_k^∞ is non-zero. We define a nontrivial sheaf homomorphism

$$Q : \mathcal{O}^{\oplus N} \rightarrow \mathcal{E}_\infty$$

by sending (a_1, \dots, a_N) to $\sum a_k s_k^\infty$. Also we know $d(s_k^\infty) = \mu_i + i_0$ for any k with $s_k^\infty \neq 0$. We want to show Q descends to be a nontrivial sheaf homomorphism from $\underline{\mathcal{F}}(i_0)$ to $\underline{\mathcal{E}}_\infty^{\mu_i}(i_0)$. It suffices to show that if

$$\sum_k a_k [s_k]([z]) = 0$$

then

$$\sum_k a_k s_k^\infty|_{\mathbb{C}^*.z} = 0.$$

Claim 3.41. *Given $[z] \notin \text{Sing}(\underline{\mathcal{E}}^{an}) \cup \Sigma \cup Z(\mathcal{E})$, where Σ is the bubbling set of the convergence to the analytic tangent cone \mathcal{E}_∞ , if*

$$\sum_k a_k[s_k]([z]) = 0,$$

then

$$\sum_k a_k s_k|_{\mathbb{C}^*.z \cap B} = s|_{\mathbb{C}^*.z \cap B}$$

for some $s \in H^0(B, \mathcal{E})$ with $d(s) > i_0 + \mu_i$.

Given this Claim, we have

$$\frac{\sum_k a_k (\lambda_j^* s_k)(z)}{m_j} = \frac{(\lambda_j)^* s(z)}{m_j} = \frac{\|s\|_j}{m_j} \cdot \frac{(\lambda_j)^* s(z)}{\|s\|_j}$$

By definition

$$\lim_{j \rightarrow \infty} \frac{\log m_j}{-j \log 2} = 2(i_0 + \mu_i + n)$$

whereas

$$\lim_{j \rightarrow \infty} \frac{\log \|s\|_j}{-j \log 2} = 2(d(s) + n)$$

In particular,

$$\lim_{j \rightarrow \infty} \frac{\|s\|_j}{m_j} = 0$$

It follows that $\sum_k a_k s_k^\infty|_{\mathbb{C}^*.z} = 0$. So Q descends to be a nontrivial sheaf homomorphism from $\underline{\mathcal{F}}(i_0)$ to $\underline{\mathcal{E}}_\infty^{\mu_i}(i_0)$ away from $\pi(\Sigma \cup Z(\mathcal{E})) \cup \text{Sing}(\underline{\mathcal{E}}^{an})$ which is of complex codimension at least 2. It then extends to a nontrivial map from $\underline{\mathcal{F}}(i_0)$ to $\underline{\mathcal{E}}_\infty^{\mu_i}(i_0)$ over the entire D . However, since $\underline{\mathcal{F}}(i_0)$ is stable and $\underline{\mathcal{E}}_\infty^{\mu_i}(i_0)$ is polystable with $\mu(\underline{\mathcal{F}}(i_0)) > \mu_i + i_0 = \mu(\underline{\mathcal{E}}_\infty^{\mu_i}(i_0))$, such a map can not be non-trivial. This is a contradiction. \square

Proof of Claim 3.41. Without loss of generality we may assume

$$[z] = ((0, 0, \dots, 0), [1, 0, \dots, 0]) \in \widehat{B} \subset B \times \mathbb{CP}^{n-1}.$$

Then the map p is locally given by

$$p(z_1, \omega_2, \dots, \omega_n) = (z_1, z_1 \omega_2, \dots, z_1 \omega_n).$$

By choosing a local trivialization of $\widehat{\mathcal{E}}(-i_0 D)$ near $[z]$ given by *global* sections, we can view the sections locally as vector valued holomorphic functions. Near $[z]$, by Taylor expansion, we have

$$\sum_k a_k s_k(z) = z_1 f_1 + \omega_2 f_2 + \dots \omega_n f_n.$$

Let $\mathbb{C}_{[z]}$ be the line given by $\omega_2 = \cdots \omega_n = 0$, which can be identified with the line $z_2 = \cdots = z_n = 0$ in \mathbb{C}^n through the projection map p . Then

$$\sum_k a_k s_k(z)|_{\mathbb{C}_{[z]}} = z_1 f_1|_{\mathbb{C}_{[z]}}.$$

Let f be the pull-back of $f_1|_{\mathbb{C}_{[z]}}$ under the composition map $\widehat{B} \rightarrow B \xrightarrow{\rho} \mathbb{C}_{[z]}$, where ρ is the natural orthogonal projection with respect to the flat metric. By our choice of local trivialization we may view f as a section s' of $\widehat{\mathcal{E}}(-i_0 D)$. Let $s = z_1 s'$, then we have

$$\sum_k a_k s_k|_{\mathbb{C}^*.z \cap B} = s|_{\mathbb{C}^*.z \cap B}.$$

Also

$$d(s) = d(s') + 1 \geq i_0 + \mu_1 + 1 > i_0 + \mu_i$$

because $\mu_i - \mu_1 < 1$. □

Remark 3.42. From Remark 3.23 a priori the definition of $\widehat{\mathcal{E}}^i$ is not canonical, but since we know it is optimal and the restriction $\widehat{\mathcal{E}}^i|_D$ is isomorphic to $\underline{\mathcal{E}}^{an,i}$ which is intrinsically defined, by Theorem 2.15 we know that the isomorphism class of $\widehat{\mathcal{E}}^i$ is uniquely determined. Moreover, we have obtained m optimal extensions $\widehat{\mathcal{E}}^i (i = 1, \dots, m)$. By Theorem 2.15, we know they are related by Hecke transforms of special type, and our constructions can recover all the different optimal algebraic tangent cones up to equivalence.

4 Proof of the main theorem

In this Section we prove Theorem 1.1. In Section 4.1 we prove part (I), and in Section 4.2 we prove Part (II). Throughout this section we fix an analytic tangent cone $(A_\infty, \Sigma_b^{an})$ with underlying reflexive sheaf \mathcal{E}_∞ , which arises as the rescaled limit corresponding to a subsequence of the fixed sequence $\{\lambda_j = 2^{-j}\}$. We will use the algebro-geometric results in [10] and [9], which is very different from the pointwise orthogonal projection technique developed in [4]. The overall discussion is very similar to the arguments in [9].

4.1 The limiting sheaf and connection

By Lemma 2.7 and Corollary 3.34, Part (I) of Theorem 1.1 is a consequence of the following

Proposition 4.1. *For $l = 1, \dots, m$, we have*

$$\underline{\mathcal{E}}_\infty^{\mu_l} \simeq (Gr^{HNS}(\underline{\mathcal{E}}_\infty^{\mu_l}))^{**},$$

and

$$\Sigma_b^{alg}(\underline{\mathcal{E}}_\infty^{\mu_l}) = \mathcal{C}(\underline{\mathcal{E}}_\infty^{\mu_l} / \underline{\mathcal{N}}_\infty^{\mu_l}).$$

Notice by (3.24) in Section 3.2 we know for $i = 1, \dots, m$, $\underline{\mathcal{E}}^{\mu_{m-i+1}}$ is identified with $\underline{\mathcal{E}}_1^i$, which is the first term of in the Harder-Narasimhan filtration of $\underline{\mathcal{E}}^{an,i}$. We shall only prove the case $i = 1$, which corresponds to the case $l = m$ in the above Proposition. For general l one can get the conclusion by repeating the argument with $\underline{\mathcal{E}}^{an,i}$ replaced by $\underline{\mathcal{E}}^{an,m-l+1}$. As before we shall omit the superscript i and only consider $\underline{\mathcal{E}}^{an} = \underline{\mathcal{E}}^{an,1}$.

Denote by $\widehat{\mathcal{E}}$ the corresponding optimal extension constructed in Section 3.3. Then we know for $k \gg 1$ there is a natural identification

$$H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) \simeq M_k^{an} = \{s \in H^0(B, \mathcal{E}) : d(s) \geq \mu_1 + k\}.$$

Suppose the Harder-Narasimhan filtration of $\underline{\mathcal{E}}^{an} = \widehat{\mathcal{E}}|_D$ is given by

$$0 = \underline{\mathcal{E}}_0 \subset \underline{\mathcal{E}}_1 \subset \dots \subset \underline{\mathcal{E}}_m = \underline{\mathcal{E}}^{an},$$

where $\mu(\underline{\mathcal{E}}_l / \underline{\mathcal{E}}_{l-1}) = \mu_{m+1-l}$. Then

$$\underline{\mathcal{E}}_l / \underline{\mathcal{E}}_{l-1} = \underline{\mathcal{E}}^{\mu_{m+1-l}}$$

is the sheaf associated to the graded module $N^{\mu_{m+1-l}}$ (see (3.14)).

Now for $k \gg 1$ we know $\widehat{\mathcal{E}}(-kD)$ is globally generated and we have the following exact sequence

$$H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) \rightarrow H^0(D, \underline{\mathcal{E}}^{an} \otimes \mathcal{O}_D(k)) \rightarrow 0$$

given by restriction to D . We may also assume the sheaf $\underline{\mathcal{E}}_1(k)$ is globally generated. Choose sections $s_i \in H^0(\widehat{B}, \widehat{\mathcal{E}}(-kD)) = M_k^{an}$, $i = 1, \dots, N$, so that when restricting to D they form a basis $\{\underline{s}_i\}$ of the vector space $H^0(D, \underline{\mathcal{E}}_1(k))$, and they globally generate the sheaf $\underline{\mathcal{E}}_1(k)$.

As in section 3.2, for each j , we can perform the Gram-Schmidt process to the sections $\lambda_j^* s_i$ over B , and obtain sections σ_i^j , $i = 1, \dots, N$, which are L^2 -orthonormal over B . Passing to a further subsequence if necessary we may assume the sections σ_i^j converge strongly to holomorphic sections σ_i^∞ of \mathcal{E}_∞ . The latter are all homogeneous of degree $k + \mu_m$ and they induce sections $\{\underline{\sigma}_i^\infty\}_{i=1}^N$ of the sheaf $\underline{\mathcal{N}}_\infty^{\mu_m}(k)$ on D . By definition, for $k \gg 1$ we know $\underline{\mathcal{N}}_\infty^{\mu_m}(k)$ is globally generated by these sections.

Now we fix a $k \gg 1$ so that all the above named properties are satisfied. Notice $\{\sigma_i^j\}_{i=1}^N$ can also be viewed as sections of $\lambda_j^* \widehat{\mathcal{E}}(-kD)$ over \widehat{B} , and they differ from $\{\lambda_j^* s_i\}_{i=1}^N$ be an element in $GL(N; \mathbb{C})$. In particular we have a sequence of sheaf homomorphisms over \widehat{B} given by

$$\mathcal{O}_{\widehat{B}}^{\oplus N} \xrightarrow{q_j = (\sigma_1^j, \dots, \sigma_N^j)} \lambda_j^* \widehat{\mathcal{E}}(-kD). \quad (4.1)$$

Now we want to take limits of these as j tends to infinity. Over \widehat{B} there are no natural limits, but we can restrict to either D or $\widehat{B} \setminus D \simeq B \setminus \{0\}$

to take limits. The first limit is *algebraic* and the second limit is *analytic*. It turns out these two limits can be related, and we will use the algebraic results quoted in Section 2.3 to get the conclusion about the analytic limit.

On the one hand, by the choice of k , we know that when restricting to D , (4.1) yields an exact sequence

$$\mathcal{O}_D^{\oplus N} \xrightarrow{q_j := (\underline{\sigma}_1^j, \dots, \underline{\sigma}_N^j)} \underline{\mathcal{E}}_1(k) \rightarrow 0.$$

In particular we obtain a sequence of points in $\mathbf{Quot}(\mathcal{H}, \tau)$, where τ denotes the Hilbert polynomial of $\underline{\mathcal{E}}_1$. By passing to a subsequence, we can take an algebraic limit

$$\mathcal{O}_D^{\oplus N} \xrightarrow{q_\infty^{alg} = (\underline{\sigma}_1^{alg, \infty}, \dots, \underline{\sigma}_N^{alg, \infty})} \underline{\mathcal{E}}_\infty^{alg}(k) \rightarrow 0 \quad (4.2)$$

in $\mathbf{Quot}(\mathcal{H}, \tau)$. Notice the notation here is different from Section 2.3 in that we have tensored everything by $\mathcal{O}(k)$. Also notice a priori $\underline{\mathcal{E}}_\infty^{alg}$ may not be torsion-free, and may depend on the choice of subsequences.

Fixing a smooth Hermitian metric on $\mathcal{O}_D^{\oplus N}$, we can identify the map q_j with the projection map π_j to the orthogonal complement of $\text{Ker}(q_j)$ away from $\text{Sing}(\underline{\mathcal{E}}_1)$. We may similarly identify q_∞ with π_∞ outside $\text{Sing}(\underline{\mathcal{E}}_\infty^{alg})$. We also fix a smooth Hermitian metric on the locally free part of $\underline{\mathcal{E}}_1$, so that the adjoint q_j^* is well-defined. From Lemma 2.24, we have the following

Lemma 4.2. π_j converge to π_∞ smoothly away from $\text{Sing}(\underline{\mathcal{E}}_\infty^{alg}) \cup \text{Sing}(\underline{\mathcal{E}}_1)$.

For our purpose, we also note the following elementary formula

$$\pi_j = q_j^* (q_j q_j^*)^{-1} q_j$$

away from $\text{Sing}(\underline{\mathcal{E}}_1)$.

On the other hand, we can restrict the exact sequence (4.1) to $B \setminus \{0\}$,

$$\mathcal{O}_{B \setminus \{0\}}^{\oplus N} \xrightarrow{q_j = (\sigma_1^j, \dots, \sigma_N^j)} \lambda_j^* \mathcal{E}. \quad (4.3)$$

Now by our discussion above each σ_i^j converges to σ_i^∞ , which is a degree $\mu_m + k$ homogeneous section of \mathcal{E}_∞ . So we obtain the exact sequence

$$\mathcal{O}_D^{\oplus N} \xrightarrow{q_\infty^{an} := (\underline{\sigma}_1^\infty, \dots, \underline{\sigma}_N^\infty)} \underline{\mathcal{N}}_\infty^{\mu_m}(k) \rightarrow 0. \quad (4.4)$$

Our goal is to compare (4.2) with (4.4). In the arguments below we need to consider the dual sheaf \mathcal{E}^* , endowed with the dual HYM metric H^* . Let $\widehat{\mathcal{E}}^*$ be the optimal extension of \mathcal{E}^* at 0 constructed in Section 3.4. Then there exists a non-degenerate pairing between $\widehat{\mathcal{E}}$ and $\widehat{\mathcal{E}}^*$ induced by the pairing between \mathcal{E} and \mathcal{E}^* over B . For $k' \gg 1$, we may find a set of sections $\{\overline{\sigma}_i^j\}_{i=1}^{N'}$

in $H^0(\widehat{B}_j, (\widehat{\mathcal{E}^*})_j(-k'D))$ (where $(\widehat{\mathcal{E}^*})_j := \lambda_j^*(\widehat{\mathcal{E}^*})$), which are L^2 orthonormal over B with respect to the rescaled metric $\lambda_j^* H^*$. Moreover, they converge strongly to homogeneous limits $\{\overline{\sigma}_i^\infty\}_{i=1}^{N'}$, which globally generate $\mathcal{F}_\infty = \mathcal{E}_\infty^*$ away from a complex codimension 2 subvariety \mathcal{S} (by Proposition 3.35).

Proposition 4.3. *There exists an isomorphism*

$$\underline{\rho}_\infty : \underline{\mathcal{E}}_\infty^{alg}(k) \rightarrow \underline{\mathcal{N}}_\infty^{\mu_m}(k)$$

induced by $\underline{q}_\infty^{alg}$ and $\underline{q}_\infty^{an}$, i.e., $\underline{\rho}_\infty$ sends $\sum_j a_j \underline{\sigma}_j^{alg, \infty}$ to $\sum_j a_j \underline{\sigma}_j^\infty$. In particular, $\underline{\mathcal{E}}_\infty^{alg}$ is torsion-free.

Proof. It suffices to show that there exists a sheaf inclusion

$$\text{Ker}(\underline{q}_\infty^{alg}) \subset \text{Ker}(\underline{q}_\infty^{an})$$

away from a proper subvariety of D . Indeed, given this, $\underline{q}_\infty^{an}(\text{Ker}(\underline{q}_\infty^{alg}))$ will be a torsion subsheaf of $\underline{\mathcal{N}}_\infty^{\mu_m}$ and has to be zero since $\underline{\mathcal{N}}_\infty^{\mu_m}$ is torsion-free. This shows that $\underline{\rho}_\infty$ is well-defined. By definition $\underline{\rho}_\infty$ is surjective, so must be an isomorphism. This is because the two sheaves $\underline{\mathcal{E}}_\infty^{alg}$ and $\underline{\mathcal{N}}_\infty^{\mu_m}$ have the same Hilbert polynomial by (3.20) and the definition of the Quot scheme. Take a generic point $[z] \in D$ such that the following hold

- $\underline{\mathcal{E}}_1$ and $\underline{\mathcal{E}}_\infty^{alg}$ are locally free near $[z]$ and the convergence $\underline{\pi}_j \rightarrow \underline{\pi}_\infty$ is smooth.
- there exists a neighborhood U of the line $\mathbb{C}_{[z]} \subset \mathbb{C}^n \cap B$, such that for all $\delta > 0$, \mathcal{E} and \mathcal{E}_∞ are locally free on $U \setminus B_\delta$ and $U \cap (\Sigma \cup Z(\mathcal{E})) = \{0\}$ where Σ is the bubbling set (see (2.4)) and $Z(\mathcal{E})$ is the \mathbb{C}^* invariant reduced subvariety in \mathbb{C}^n underlying the Zariski tangent cone of $\text{Sing}(\mathcal{E})$ at 0.
- $\widehat{\mathcal{E}}$ is locally free in a neighborhood of $[z]$.
- $\underline{\mathcal{N}}_\infty^{\mu_m}$ is locally free in a neighborhood of $[z]$, and its fiber over $[z]$ is generated by the limit sections $\underline{\sigma}_1^\infty, \dots, \underline{\sigma}_N^\infty$.
- $\mathbb{C}_{[z]} \cap \mathcal{S} = \{0\}$.

Over the point $[z]$, it suffices to show the inclusion $\text{Ker}(\underline{q}_\infty^{alg})|_{[z]} \subset \text{Ker}(\underline{q}_\infty^{an})|_{[z]}$ as vector subspaces of \mathbb{C}^N (the fiber of $\mathcal{O}_D^{\oplus N}$ at $[z]$). Suppose $a \in \mathbb{C}^{\oplus N}$ is given so that $\underline{q}_\infty^{alg}(a) = 0$. Then by definition we have $\underline{\pi}_\infty(a)|_{[z]} = 0$. Hence by Lemma 4.2

$$\lim_{j \rightarrow \infty} \underline{\pi}_j(a)|_{[z]} = 0.$$

Fix a nonzero $z \in \mathbb{C}^n$ on the complex line $\mathbb{C}_{[z]}$. We want to show

$$\lim_{j \rightarrow \infty} q_j(a)(z) = 0,$$

where this limit is taken in the analytic sense.

Now we let

$$\sigma^j = q_j(a - \pi_j(a)|_{[z]}).$$

It follows that $\sigma^j([z]) = 0$. Moreover, the analytic limit on the analytic tangent cone \mathcal{E}_∞ is given by

$$\sigma^\infty := \lim_{j \rightarrow \infty} \sigma^j = \lim_{j \rightarrow \infty} q_j(a).$$

It suffices to show that σ^∞ vanishes at any $z \neq 0$ on the line $\mathbb{C}_{[z]}$.

To show this we make use of the dual sheaf \mathcal{E}^* . Notice by our choice of $[z]$, \mathcal{F}_∞ is locally free in a neighborhood of $[z]$, so we may choose $r = \text{rank}(\mathcal{E})$ sections, say $\bar{\sigma}_1^\infty, \dots, \bar{\sigma}_r^\infty$, which generate the fibers of \mathcal{F}_∞ over $\mathbb{C}_{[z]} \setminus \{0\}$.

For each i , we have

$$\bar{\sigma}_i^j(\sigma^j) = f_i^j \in H^0(\widehat{B}, \mathcal{I}_{[z]}(-(k+k')D)). \quad (4.5)$$

In particular, when viewed as a holomorphic function over B , $f_i^j|_{\mathbb{C}_{[z]}}$ has a vanishing order at least $k+k'+1$. If we take limits to the analytic tangent cones, we know that $\bar{\sigma}_i^\infty$ has degree k' and σ^∞ has degree k , so the function

$$f_i^\infty = \bar{\sigma}_i^\infty(\sigma^\infty)$$

is a homogeneous polynomial which either is identically zero, or has degree exactly equal to $k+k'$. In particular, restricting to the line $\mathbb{C}_{[z]}$, $f_i^\infty|_{\mathbb{C}_{[z]}}$ either vanishes identically or has a vanishing order exactly equal to $k+k'$ at 0. However, since for each j , $f_i^j|_{\mathbb{C}_{[z]}}$ vanishes at 0 up to order at least $k'+k+1$, and they converge uniformly to f_i^∞ outside any small neighborhood of 0, by convergence of residues, we know $f_i^\infty|_{\mathbb{C}_{[z]}}$ vanishes at 0 up to order at least $k'+k+1$. In particular, for any i , f_i^∞ is identically zero on the line $\mathbb{C}_{[z]}$. So

$$\bar{\sigma}_i^\infty(\sigma^\infty)|_{\mathbb{C}_{[z]}} = 0.$$

By assumption, $\{\bar{\sigma}_i^\infty\}_{i=1}^r$ generate the fibers of $\mathcal{E}_\infty^*|_{\mathbb{C}_{[z]}}$, so we conclude that $\sigma^\infty|_{\mathbb{C}_{[z]}} = 0$. □

Now we need to use the results from Section 2.3 and finish the proof. Indeed, since $\underline{\mathcal{E}}_\infty^{\mu_m} = (\underline{\mathcal{N}}_\infty^{\mu_m})^{**}$ admits an admissible HYM connection, it is polystable, Proposition 4.1 follows directly from Corollary 2.25.

Remark 4.4. In the proof above, to find each factor $\underline{\mathcal{E}}_\infty^{\mu_i}$, we can choose a set of global sections $\{s_{i,l} \in H^0(B, \mathcal{E})\}_{l=1}^{N_i}$ with $d(s_{i,l}) = k + \mu_i$ (here we can assume that $k \gg 1$ is uniform for all i), such that the homogeneous limits given by those sections generate $\underline{\mathcal{N}}_\infty^{\mu_i}(k)$. Furthermore, by assumption, the union of all $s_{i,l}$ ($i = 1, \dots, m, l = 1, \dots, N_i$) spans M_k^{an}/M_{k+1}^{an} over \mathbb{C} , and they globally generate $\widehat{\mathcal{E}}(-kD)$ over \widehat{B} . For simplicity we shall denote these sections by $\sigma_1, \dots, \sigma_N$, and the induced sections of $\lambda_j^* \mathcal{E}$ by $\sigma_1^j, \dots, \sigma_N^j$.

4.2 The analytic blow-up cycle

By Corollary 3.34 and Proposition 4.1, to prove Part (II) of Theorem 1.1, it suffices to prove

$$\sum_{l=1}^m \mathcal{C}(\underline{\mathcal{E}}_\infty^{\mu_l} / \underline{\mathcal{N}}_\infty^{\mu_l}) = \Sigma_b^{an}. \quad (4.6)$$

By definition we have

$$\underline{\mathcal{E}}_\infty = \bigoplus_{l=1}^m \underline{\mathcal{E}}_\infty^{\mu_l}.$$

As in Remark 3.32, we denote

$$\underline{\mathcal{N}}_\infty := \bigoplus_{l=1}^m \underline{\mathcal{N}}_\infty^{\mu_l},$$

and

$$\mathcal{N}_\infty := \psi_* \pi^* \underline{\mathcal{N}}_\infty.$$

By Remark 4.4, one can find a sequence of global resolutions of $\mathcal{E}_j = \lambda_j^* \mathcal{E}$ over $B \setminus \{0\}$, which are of the form

$$0 \rightarrow \mathcal{K}_j \rightarrow \mathcal{O}_{B \setminus \{0\}}^{\oplus N} \xrightarrow{p_j = (\sigma_1^j, \dots, \sigma_N^j)} \mathcal{E}_j \rightarrow 0.$$

By passing to a subsequence we may assume the above converges to a global resolution of $\mathcal{N}_\infty \subset \underline{\mathcal{E}}_\infty$:

$$0 \rightarrow \mathcal{K}_\infty \rightarrow \mathcal{O}_{B \setminus \{0\}}^{\oplus N} \xrightarrow{p_\infty = (\sigma_1^\infty, \dots, \sigma_N^\infty)} \mathcal{N}_\infty \rightarrow 0. \quad (4.7)$$

Here the crucial fact for us is that by homogeneity Equation (4.7) naturally descends to the projective space \mathbb{CP}^{n-1} as a short exact sequence

$$0 \rightarrow \underline{\mathcal{K}}_\infty \rightarrow \mathcal{O}_{\mathbb{CP}^{n-1}}^{\oplus N} \xrightarrow{\underline{p}_\infty = (\underline{\sigma}_1^\infty, \dots, \underline{\sigma}_N^\infty)} \underline{\mathcal{N}}_\infty(k) \rightarrow 0. \quad (4.8)$$

This enables us to use the global Bott-Chern formula in [17, 9] to calculate the algebraic multiplicity of the torsion sheaf $\underline{\mathcal{T}} = \underline{\mathcal{E}}_\infty / \underline{\mathcal{N}}_\infty$ along a codimension two irreducible subvariety in \mathbb{CP}^{n-1} .

In the following we shall fix a flat Hermitian metric on $\mathcal{O}_D^{\oplus N}$ which induces a flat metric on $\mathcal{O}_{B \setminus \{0\}}^{\oplus N}$. Let \mathcal{K}_j^\perp denote the orthogonal complement \mathcal{K}_j^\perp of \mathcal{K}_j in $\mathcal{O}^{\oplus N}$ with respect to the flat metric. It is a smooth sub-bundle of $\mathcal{O}^{\oplus N}$ away from the $\text{Sing}(\mathcal{E}_j)$. We denote by $p_j^* A_j$ the induced connection on \mathcal{K}_j^\perp through the smooth isomorphism $\mathcal{K}_j^\perp \simeq \mathcal{E}_j$. Similarly we define $p_\infty^* A_\infty, \underline{p}_\infty^* \underline{A}_\infty$ etc.

Given *any* codimension two irreducible subvariety $\underline{\Gamma} \subset \mathbb{CP}^{n-1}$, the algebraic multiplicity $m^{alg}(\underline{\Gamma})$ of $\underline{\mathcal{E}}_\infty/\underline{\mathcal{N}}_\infty$ along $\underline{\Gamma}$ is defined to be

$$m^{alg}(\underline{\Gamma}) := \dim H^0(\underline{\Delta}, \mathcal{T}|_{\underline{\Delta}})$$

where $\underline{\Delta}$ is a holomorphic transverse slice of $\underline{\Gamma}$ at a generic point. Then by definition we have

$$\mathcal{C}(\underline{\mathcal{T}}) = \sum_{\underline{\Gamma}} m^{alg}(\underline{\Gamma}) \cdot [\underline{\Gamma}].$$

Take any such $\underline{\Gamma}$. We can compute $m^{alg}(\underline{\Gamma})$ using the exact sequence (4.8) (see [17, 9]). To explain this, we fix a flat unitary connection \underline{A}_f on $\mathcal{O}_D^{\oplus N}$ hence a flat unitary connection A_f on $\mathcal{O}_{B \setminus \{0\}}^{\oplus N}$. Let G_j be the induced connection on \mathcal{K}_j and G_∞ be the pull-back of a fixed admissible connection \underline{G}_∞ on $\underline{\mathcal{K}}_\infty$. Here being admissible means that \underline{G}_∞ is a smooth connection on the locally free locus of $\underline{\mathcal{K}}_\infty$ which has finite Yang-Mills energy and bounded $\Lambda_{\omega_{FS}} F_{\underline{G}_\infty}$. Notice since \mathcal{N}_∞ is torsion-free, we know \mathcal{K}_∞ hence $\underline{\mathcal{K}}_\infty$ is reflexive, which implies its singular set has complex codimension at least 3. So we can make the connection \underline{G}_∞ smooth at a generic point of $\underline{\Gamma}$.

We know that A_∞ descends to a HYM connection on $\underline{\mathcal{E}}_\infty(k)$, which is the direct sum of admissible HYM connections. We denote it by \underline{A}_∞ . Choose a two dimensional smooth subvariety $\underline{\Delta}$ of \mathbb{CP}^{n-1} so that $\underline{\Delta}$ intersects $\underline{\Gamma}$ transversely and positively at $\underline{z}_1, \dots, \underline{z}_l$. Then we have (see Page 59 in [9])

$$\begin{aligned} & l \cdot m^{alg}(\underline{\Gamma}) \\ &= \frac{1}{8\pi^2} \sum_i \left\{ \int_{\underline{\Delta} \cap B_\epsilon(\underline{z}_i)} \text{Tr}(F_{\underline{A}_f} \wedge F_{\underline{A}_f}) - \text{Tr}(F_{\underline{G}_\infty \oplus (\underline{p}_\infty)^* \underline{A}_\infty} \wedge F_{\underline{G}_\infty \oplus (\underline{p}_\infty)^* \underline{A}_\infty}) \right. \\ & \quad \left. - \int_{\partial(\underline{\Delta} \cap B_\epsilon(\underline{z}_i))} CS(\underline{A}_f, \underline{G}_\infty \oplus (\underline{p}_\infty)^* \underline{A}_\infty) \right\} \end{aligned} \tag{4.9}$$

where $B_\epsilon(\underline{z}_i)$ denotes a small ball of radius ϵ in \mathbb{CP}^{n-1} , and $CS(\cdot, \cdot)$ is the Chern-Simons functional defined using the obvious trivialization of $\mathcal{O}_{\mathbb{CP}^{n-1}}^{\oplus N}$ over the boundary of each $\underline{\Delta} \cap B_\epsilon(\underline{z}_i)$ (see [4], Section 4.2). Here we may assume each $\underline{\Delta} \cap B_\epsilon(\underline{z}_i)$ only intersects $\underline{\Gamma}$ at one point and also

$$(B_\epsilon(\underline{z}_i) \setminus \underline{\Gamma}) \cap (\text{supp}(\Sigma_b^{an}) \cup \text{Sing}(\underline{A}_\infty)) = \emptyset$$

Notice (4.9) was stated in [9] when $\underline{\Gamma}$ is contained in the support of $\underline{\mathcal{T}}$. But if $\underline{\Gamma}$ is not contained in the support of $\underline{\mathcal{T}}$, then $m^{alg}(\underline{\Gamma}) = 0$ and (4.9) obviously holds.

For each i , we choose a holomorphic lift $\Delta_i \subset B$ of $\underline{\Delta} \cap B_\epsilon(\underline{z}_i)$. Then as a direct corollary of Equation (4.9), we obtain (see similar calculation in

Section 4.2 in [4])

$$\begin{aligned}
l \cdot m^{alg}(\underline{\Gamma}) &= \frac{1}{8\pi^2} \sum_i \left\{ \int_{\Delta_i} \text{Tr}(F_{A_f} \wedge F_{A_f}) - \text{Tr}(F_{G_\infty \oplus p_\infty^* A_\infty} \wedge F_{G_\infty \oplus p_\infty^* A_\infty}) \right. \\
&\quad \left. - \int_{\partial \Delta_i} CS(A_f, G_\infty \oplus p_\infty^* A_\infty) \right\}
\end{aligned} \tag{4.10}$$

On the other hand, let $\Gamma \subset \mathbb{C}^n$ be the affine cone over $\underline{\Gamma}$, then the analytic multiplicity along Γ is given by

$$l \cdot m^{an}(\Gamma) := \lim_{j \rightarrow \infty} \frac{1}{8\pi^2} \sum_i \left\{ \int_{\Delta_i} \text{Tr}(F_{A_j} \wedge F_{A_j}) - \text{Tr}(F_{A_\infty} \wedge F_{A_\infty}) \right\}.$$

So we have

$$\Sigma_b^{an} = \sum_{\underline{\Gamma}} m^{an}(\Gamma) \cdot [\underline{\Gamma}].$$

Now the equality (4.6) follows from

Proposition 4.5. *For any codimension two irreducible subvariety $\underline{\Gamma} \subset \mathbb{CP}^{n-1}$, we have $m^{alg}(\underline{\Gamma}) = m^{an}(\Gamma)$.*

Proof. By Equation (4.10) and Lemma 2.1, a direct computation yields

$$\begin{aligned}
&l \cdot m^{alg}(\underline{\Gamma}) - l \cdot m^{an}(\Gamma) \\
&= \sum_i \left\{ \int_{\partial \Delta_i} CS(G_\infty \oplus p_\infty^* A_\infty, G_j \oplus p_j^* A_j) \right. \\
&\quad \left. + \frac{1}{8\pi^2} \int_{\Delta_i} \text{Tr}(F_{G_j} \wedge F_{G_j}) - \text{Tr}(F_{G_\infty} \wedge F_{G_\infty}) \right\}.
\end{aligned}$$

It suffices to show that for each i

$$\lim_{j \rightarrow \infty} \left\{ \int_{\partial \Delta_i} CS(G_\infty \oplus p_\infty^* A_\infty, G_j \oplus p_j^* A_j) + \int_{\Delta_i} \text{Tr}(F_{G_j} \wedge F_{G_j}) - \text{Tr}(F_{G_\infty} \wedge F_{G_\infty}) \right\} = 0. \tag{4.11}$$

By assumption, using the given flat metric, we obtain an orthogonal splitting

$$\mathcal{O}_{B \setminus \{0\}}^{\oplus N} = \mathcal{K}_j \oplus \mathcal{K}_j^\perp.$$

Then the orthogonal projection π_j from $\mathcal{O}_{B \setminus \{0\}}^{\oplus N}$ to the orthogonal complement of \mathcal{K}_j converges to π_∞ away from the center of Δ_i . Now by assumption \mathcal{K}_∞ is locally free over Δ_i so we may fix a smooth trivialization of \mathcal{K}_∞ over Δ_i . In particular this yields another trivialization of \mathcal{K}_∞ over $\partial \Delta_i$. Fix an arbitrary smooth trivialization of \mathcal{K}_∞^\perp over $\partial \Delta_i$, so together we get a new trivialization of

$$\mathcal{O}_{B \setminus \{0\}}^{\oplus N} = \mathcal{K}_\infty \oplus \mathcal{K}_\infty^\perp$$

over $\partial\Delta_i$. Using the smooth convergence above we may for $j \gg 1$ find smooth trivializations of \mathcal{K}_j and \mathcal{K}_j^\perp so that they are identified with \mathcal{K}_∞ and \mathcal{K}_∞^\perp (as complex vector bundles). Notice by [4], Lemma 4.1, the relative Chern-Simons integral does not depend on the choice of the trivialization of the bundle, so using the new trivialization we may write

$$CS(G_\infty \oplus p_\infty^* A_\infty, G_j \oplus p_j^* A_j) = CS(G_\infty, G_j) + CS(p_\infty^* A_\infty, p_j^* A_j).$$

Now since the above trivializations of \mathcal{K}_∞ and \mathcal{K}_j on $\partial\Delta_i$ extend over Δ_i , it follows that (see Lemma 4.1 in [4])

$$CS(G_j, G_\infty) = \int_{\Delta_i} \text{Tr}(F_{G_j} \wedge F_{G_j}) - \text{Tr}(F_{G_\infty} \wedge F_{G_\infty}).$$

So it suffices to prove

$$\lim_{j \rightarrow \infty} \int_{\partial\Delta_i} CS(p_\infty^* A_\infty, p_j^* A_j) = 0.$$

Essentially, this follows from the fact that p_∞ comes from the limit of maps which are initially globally defined over Δ_i . By assumption, there exists a sequence of gauge transforms g_j and Hermitian isomorphisms $P_j : \mathcal{E}_j \rightarrow \mathcal{E}_\infty$, both defined away from Γ , so that $(P_j^{-1})^*(g_j \cdot A_j)$ converges smoothly to A_∞ away from Γ . Furthermore, $P_j g_j p_j$ converge to p_∞ smoothly away from Γ . Given this, we have

$$\begin{aligned} & \int_{\partial\Delta_i} CS(p_\infty^* A_\infty, p_j^* A_j) \\ &= \lim_{j' \rightarrow \infty} \int_{\partial\Delta_i} CS((P_{j'} g_{j'} p_{j'})^* (P_{j'}^{-1})^* g_{j'} \cdot A_{j'}, p_j^* A_j) \\ &= \lim_{j' \rightarrow \infty} \int_{\partial\Delta_i} CS((p_{j'})^* A_{j'}, p_j^* A_j) \\ &= \lim_{j' \rightarrow \infty} \int_{\Delta_i} \text{Tr}(F_{A_{j'}} \wedge F_{A_{j'}}) - \text{Tr}(F_{A_j} \wedge F_{A_j}), \end{aligned}$$

where the last equality follows from the fact that p_j and p'_j are globally defined over Δ_i and Lemma 4.1 in [4]. Now let j tend to infinity, the last term converges to zero by Lemma 2.1. □

References

- [1] S. Bando, Y-T. Siu. *Stable sheaves and Einstein-Hermitian metrics*, Geometry and Analysis on Complex Manifolds, T. Mabuchi, J. Noguchi, and T. Ochiai, eds., World Scientific, River Edge, NJ, 1994, 39–50.

- [2] N. Buchdahl, A. Teleman, M. Toma. *A continuity theorem for families of sheaves on complex surfaces*. Journal of Topology 10.4 (2017): 995-1028.
- [3] X.M. Chen, and S. Sun. *Singularities of Hermitian-Yang-Mills connections and Harder-Narasimhan-Seshadri filtrations*. Duke Mathematical Journal 169, no. 14 (2020): 2629-2695.
- [4] X.M. Chen, S. Sun. *Analytic tangent cones of admissible Hermitian-Yang-Mills connections*. To appear in Geometry and Topology arXiv:1806.11247 (2018).
- [5] X.M. Chen, S. Sun. *Algebraic Tangent Cones of Reflexive Sheave*. International Mathematics Research Notices, rny276, <https://doi.org/10.1093/imrn/rny276>
- [6] S. Donaldson, S. Sun. *Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, II*. Journal of Differential Geometry 107, no. 2 (2017): 327-371.
- [7] D. Eisenbud. *Commutative Algebra: with a view toward algebraic geometry*. Vol. 150. Springer Science Business Media, 2013.
- [8] A. Fujiki. *On the blowing down of analytic spaces*. Publ. Res. Inst. Math. Sci. 10 (1974/75), 473-507.
- [9] D. Greb, B. Sibley, M. Toma, R. Wentworth. *Complex algebraic compactifications of the moduli space of Hermitian-Yang-Mills connections on a projective manifold*. arXiv:1810.00025.(2018).
- [10] D. Greb and M. Toma, *Compact moduli spaces for slope-semistable sheaves*. Algebr. Geom. 4 (2017), no. 1, 40-78. MR 3592465
- [11] A. Grothendieck. *Techniques de construction et théorèmes d'existence en géométrie algébrique. IV. Les schémas de Hilbert*. Séminaire Bourbaki 6.221 (1960): 249-276.
- [12] R. Hartshorne, *Algebraic geometry*. Vol. 52. Springer Science & Business Media, 2013.
- [13] A. Jacob, H. Sá Earp, and T. Walpuski. *Tangent cones of Hermitian Yang-Mills connections with isolated singularities*. Mathematical Research Letter 25.5 (2018): 1429.
- [14] S. Kobayashi. *Differential geometry of complex vector bundles*. Publications of the Mathematical Society of Japan, 15. Kano Memorial Lectures, 5. Princeton University Press, Princeton, NJ (1987).
- [15] M. Maruyama. *On boundedness of families of torsion free sheaves*. J. Math. Kyoto Univ. 21 (1981), no. 4, 673-701.

- [16] B. Shiffman. *On the removal of singularities of analytic sets*. Michigan Math. J 15.1 (1968): 1.
- [17] B. Sibley, R. Wentworth. *Analytic cycles, Bott-Chern forms, and singular sets for the Yang-Mills flow on Kähler manifolds*. Advances in mathematics 279 (2015), 501-531.
- [18] G. Tian. *Gauge theory and calibrated geometry*. I. Ann. of Math. (2) 151.1 (2000), 193–268
- [19] H. Whitney. *Tangents to an analytic variety*. Annals of mathematics (1965): 496-549.