# Classifying spaces of infinity-sheaves

Daniel Berwick-Evans Department of Mathematics, University of Illinois at Urbana-Champaign\*

## Pedro Boavida de Brito

Department of Mathematics, Instituto Superior Técnico, University of Lisbon<sup>†</sup>

#### Dmitri Pavlov

Department of Mathematics and Statistics, Texas Tech University<sup>‡</sup>

#### Abstract

We prove that the set of concordance classes of sections of an  $\infty$ -sheaf on a manifold is representable, extending a theorem of Madsen and Weiss. This is reminiscent of an h-principle in which the role of isotopy is played by concordance. As an application, we offer an answer to the question: what does the classifying space of a Segal space classify?

## Contents

1	Introduction	2
2	The concordance resolution is concordance-invariant 2.a Concordance-invariant $\infty$ -sheaves on manifolds are representable 2.b Homotopy groups of BF	
3	Weak Kan fibrations 3.a Kan fibrations and weak Kan fibrations of simplicial spaces	
4	Weak Kan fibrancy of the concordance resolution  4.a The sheaf associated to a simplicial set	15 16
<b>5 6</b>	The shape functor preserves the $\infty$ -sheaf property 5.a The finite case	
A	Simplicial sets and spaces  A.a Weak equivalences of simplicial sets	

<sup>\*</sup>https://faculty.math.illinois.edu/~danbe/; danbe@illinois.edu

<sup>†</sup>https://www.math.tecnico.ulisboa.pt/~pbrito/; pedrobbrito@tecnico.ulisboa.pt

<sup>&</sup>lt;sup>‡</sup>https://dmitripavlov.org/

### 1 Introduction

Let F be an  $\infty$ -sheaf (alias homotopy sheaf, see Definition 2.15) on Man, the site of finite-dimensional smooth manifolds without boundary and smooth maps. For a manifold M, an element of  $F(M \times \mathbb{R})$  is called a *concordance*. Two elements  $\sigma_0$ ,  $\sigma_1$  in F(M) are said to be *concordant* (and we write  $\sigma_0 \sim_c \sigma_1$ ) if there exists a concordance whose restriction to  $M \times \{k\}$  is  $\sigma_k$  for k = 0, 1.

Concordance is an equivalence relation, and a familiar one in many situations. Here are three examples. When  $F = C^{\infty}(-, N)$ , maps are concordant if and only if they are smoothly homotopic. For the sheaf of closed differential n-forms, two sections (i.e., closed n-forms) are concordant if and only if they are cohomologous. For the stack of vector bundles, a pair of vector bundles are concordant if and only if they are isomorphic. In these three cases, concordance classes have a well-known description in terms of homotopy classes of maps into a space, namely the space underlying N, the Eilenberg–MacLane space  $K(\mathbb{R}, n)$ , and the space BO(n), respectively. In this paper we generalize these classical representability results: concordance classes of sections of  $any \infty$ -sheaf F is represented by a space BF that we call the classifying space of F.

We now assemble the ingredients to state our results precisely. We denote by  $\mathbb{A}^n$  the smooth extended simplex, that is, the subspace of  $\mathbb{R}^{n+1}$  whose coordinates sum to one. By varying n, this defines a cosimplicial object in Man. Define a presheaf

$$\mathbf{B}F(M) := |[k] \mapsto F(M \times \mathbb{A}^k)|$$

with values in spaces (i.e., simplicial sets), where the vertical bars denote the homotopy colimit of the simplicial space. The construction **B** is a form of localization; it is the universal way to render the maps  $F(M) \to F(M \times \mathbb{A}^1)$  invertible for all M and F. It is a familiar construction in the motivic literature, for example in the work of Morel-Voevodsky [15] (who call it Sing), but it has also appeared in the context of geometric topology in the papers of Waldhausen [21], Weiss-Williams [24], and Madsen-Weiss [13]. The link between  $\mathbf{B}F$  and the concordance relation  $\sim_c$  is the bijection  $\pi_0\mathbf{B}F(M) \cong \pi_0F(M)/\sim_c$ .

Define the classifying space BF as a Kan complex replacement of  $\mathbf{B}F(*)$ . The main result is:

**Theorem 1.1.** Let F be an  $\infty$ -sheaf on Man. There is an evaluation map

$$\mathbf{B}F(M) \to \mathrm{map}(M, \mathrm{B}F)$$

which is a natural weak equivalence of spaces for every manifold M.

It is not difficult to show—essentially by a variant of Brown's representability theorem, see Section 2—that Theorem 1.1 is equivalent to the following.

**Theorem 1.2.** If F is an  $\infty$ -sheaf, then  $\mathbf{B}F$  is an  $\infty$ -sheaf.

These statements may be regarded as analogues of the h-principle, where the usual relation of *isotopy* is replaced by that of *concordance*. Here we have in mind the strand of the h-principle that gives conditions (e.g., microflexibility) which guarantee that an isotopy invariant functor (e.g., a sheaf) is an  $\infty$ -sheaf. The relation of concordance is more severe than that of isotopy, and this explains why the hypotheses are less restrictive than those of typical h-principles, e.g., there are no dimension restrictions, open versus closed manifolds, etc.

Just as with the h-principle, the key step in our proof involves verifying certain fibration properties. As such, a significant part of the paper is a study of weak lifting properties for maps of simplicial spaces. We introduce the notion of weak Kan fibration of simplicial spaces and simplicial sets. A crucial result shows that weak Kan fibrations are realization fibrations (see Definition 3.14 and Theorem 3.17); this implies that geometric realization is stable under homotopy pullback along weak Kan fibrations.

We emphasize that these results—and hence Theorem 1.2—do not follow from formal considerations. There are simple counterexamples in the category of schemes as in the  $\mathbb{A}^1$ -homotopy theory of Morel–Voevodsky [15, §3, Example 2.7]. The  $\infty$ -sheaf property is a homotopy limit condition whereas  $\mathbf{B}$  involves geometric realization, a homotopy colimit. Commuting these is a subtle issue. This is where we use the weak Kan property to prove Theorem 1.2. The verification that  $\mathbf{B}F$  is weak Kan and certain restriction maps are weak Kan fibrations follows from a geometric argument about smooth maps (Lemma 4.20). We also rely on the existence of partitions of unity.

Our main results improve on prior work of others. The  $\pi_0$ -statement of Theorem 1.1 is due to Madsen–Weiss [13, Appendix A] when F is a sheaf of sets (or of discrete categories). Improving Madsen–Weiss' result from  $\pi_0$  to  $\pi_n$  is not a great achievement. Improving their result from sheaves of sets to  $\infty$ -sheaves of spaces requires new ideas and is our main contribution. Bunke–Nikolaus–Völkl [3, §7] have proved a version of Theorem 1.1 for  $\infty$ -sheaves on compact manifolds with values in spectra. From the point of view of Theorem 1.2, this case is essentially formal since in a stable setting homotopy pullbacks squares are homotopy pushouts squares, and so (for finite covers) the problem of commuting homotopy pullbacks with geometric realization disappears. It has also been pointed out to us by John Francis that [1] contains related results that are proved in the context of stratified spaces. As we understand it, these results [1, 2.3.16, 2.4.5] are both more general (they apply to stratified spaces) and less general than ours and than Madsen–Weiss' (they apply to a certain class of stratified spaces) and less general than ours and than Madsen–Weiss' (they apply to a certain class of stratified spaces). This restricted class of sheaves is, from the point of view of Theorem 1.1, too severe as it excludes many  $\infty$ -sheaves, even set-valued ones.

Finally, unlike Madsen-Weiss' approach, our arguments apply in the topological or PL category too: Theorem 1.1 remains true if we consider topological or PL manifolds instead of smooth manifolds. In fact, our arguments simplify significantly in those cases (see Section 4 for explanations).

Applications of Theorem 1.1 abound. This stems from the fact that we not only prove abstract representability, but also give a formula for the representing space. This formula can be identified with classical constructions. Two illustrative examples, connecting back to the beginning of this introduction, are the classical de Rham theorem and the classification of vector bundles (with or without connection). In Section 6, we discuss a further application: a classification of **C**-bundles, where **C** is a Segal space.

We mention here another consequence of Theorem 1.1. Let  $\mathbb{D}$  denote the full subcategory of Man spanned by  $\mathbb{R}^n$ ,  $n \geq 0$ . The  $\infty$ -categories  $\mathsf{Sh}(\mathsf{Man})$  and  $\mathsf{Sh}(\mathbb{D})$  of  $\infty$ -sheaves on Man and  $\mathbb{D}$ , respectively (with respect to the usual open covers by codimension zero embeddings), are examples of  $\infty$ -topoi, as are the slice  $\infty$ -categories  $\mathsf{Sh}(\mathsf{Man})/F$  for an  $\infty$ -sheaf F on Man, and  $\mathsf{Sh}(\mathbb{D})/F$  for an  $\infty$ -sheaf F on  $\mathbb{D}$ .

**Proposition 1.3.** The functor B from  $\mathsf{Sh}(\mathsf{Man})$  to spaces is homotopy left adjoint to the functor which sends a space to the constant  $\infty$ -sheaf on that space. Moreover, B is homotopy left adjoint to the inclusion of the  $\infty$ -category of constant  $\infty$ -sheaves on  $\mathsf{Man}$  into  $\mathsf{Sh}(\mathsf{Man})$ . These two statements also hold if  $\mathsf{Man}$  is replaced by  $\mathbb{D}$ .

In other words, BF is the shape of Sh(Man)/F as in Lurie [11, Chapter 7], or equivalently, the fundamental  $\infty$ -groupoid of F in the sense of Schreiber [19, Section 3.4], and **B**F is the shape modality of F in the sense of Schreiber [19, Definition 3.4.4].

*Proof.* By a constant  $\infty$ -sheaf, we mean the homotopy sheafification of a constant presheaf. Every constant presheaf on  $\mathbb D$  is an  $\infty$ -sheaf, and for a space K the canonical functor from the constant  $\infty$ -sheaf to the mapping space  $\infty$ -sheaf const<sub>K</sub>  $\to$  map(-,K) is an objectwise weak equivalence. On the other hand, a constant presheaf on Man is in general not an  $\infty$ -sheaf. However, since every open cover can be refined by a good open cover, the homotopy sheafification of a presheaf on Man is determined by its restriction to  $\mathbb D$ . Therefore const<sub>K</sub>  $\to$  map(-,K) is also a weak equivalence in Sh(Man).

If F is a representable presheaf, represented by a manifold M, then  $BF \simeq M$ . From this it follows that

$$map(F, const_K) \simeq map(BF, K)$$

for F a representable, and extending by colimits, the same is true for any presheaf F, and then for any  $\infty$ -sheaf (the mapping space on the left is computed in the  $\infty$ -category of  $\infty$ -sheaves or equivalently, since  $\text{const}_K$  is an  $\infty$ -sheaf, in the  $\infty$ -category of presheaves).

As for the second statement, we have by Theorem 1.1 that  $\mathbf{B}F$  is a constant  $\infty$ -sheaf on Man, for any  $\infty$ -sheaf F. Then, by Yoneda,  $\operatorname{map}(\mathbf{B}F,\operatorname{const}_K) \simeq \operatorname{map}(BF,K)$ , which by the first part implies the statement.

Another application of this work is a construction of classifying spaces of field theories. This has recently been done by the third author and Grady [9], see in particular Theorem 7.2.9 there. Stolz and Teichner have conjectured that concordance classes of particular classes of field theories determine cohomology theories [20]. By Brown representability, this conjecture requires concordance classes of field theories to define a

representable functor. In brief, they define field theories as functors out of a category of bordisms equipped with a smooth map to a fixed manifold M. When the relevant bordism category is fully extended, field theories are an  $\infty$ -sheaf evaluated on M. The main result of this paper then shows that concordance classes of fully extended field theories are representable. Furthermore, we identify a formula for the classifying space of field theories.

#### Acknowledgements

We would like to thank Peter Teichner for asking the question that started this project, and for his encouragement and patient feedback throughout the process. We also thank Michael Weiss for generously sharing his insight into this problem (particularly in regard to Lemma 4.18). This project has been going on for many years, and during that time we have benefited from discussions with many colleagues and hosting of various institutions. In particular, we thank Ulrich Bunke, Jacob Lurie, Gustavo Granja, Aaron Mazel-Gee and Thomas Nikolaus for helpful discussions and comments, and the referee for helpful remarks that simplified an argument in a previous version of this paper. We are grateful to the MSRI, the MPI and the HIM in Bonn, and the universities of Münster and Stanford for their hospitality. D.P.'s gratitude extends to the nLab for being a wonderful resource. P.B. was supported by FCT through grant SFRH/BPD/99841/2014.

### 2 The concordance resolution is concordance-invariant

**Definition 2.1.** Throughout, *space* will mean simplicial set. The category of such is denoted by S. A *simplicial space* is a simplicial object in spaces, and the category of such is denoted S. Of course, this is the same as a bisimplicial set, though the terminology emphasizes that there is a preferred simplicial direction. A simplicial set is often viewed as a simplicial discrete space, by regarding a set as a discrete (or constant) simplicial set. We denote the diagonal of a bisimplicial set S by S, this is our preferred model for homotopy colimits of simplicial spaces.

**Definition 2.2.** We write  $\Delta^n$  for the representable simplicial set and  $\Delta[n]$  for the corresponding simplicial discrete space. Similarly, we write  $\partial \Delta^n$  and  $\Lambda_k^n$  for the simplicial set boundary and kth horn respectively, and  $\partial \Delta[n]$  and  $\Lambda_k[n]$  for the corresponding simplicial spaces.

**Definition 2.3.** We denote by Man the site of smooth manifolds and smooth maps, equipped with the Grothendieck topology of open covers.

**Definition 2.4.** A presheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  is *concordance invariant* if for all manifolds M the map  $F(M) \to F(M \times \mathbb{R})$  induced by the projection  $M \times \mathbb{R} \to M$  is a weak equivalence.

**Definition 2.5.** Set

$$\mathbb{A}: \Delta \to \mathsf{Man}, \qquad \mathbb{A}^n = \Big\{ x \in \mathbb{R}^{n+1} \; \Big| \; \sum_i x_i = 1 \Big\}.$$

Given a presheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$ , denote by  $\mathbf{B}F$  the presheaf

$$\mathbf{B} F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}, \qquad \mathbf{B} F(M) := \big| [k] \mapsto F(M \times \mathbb{A}^k) \big|,$$

where |-| denotes the diagonal of a bisimplicial set.

In this section we show that  $\mathbf{B}F$  is always concordance invariant. Furthermore, if  $\mathbf{B}F$  is an  $\infty$ -sheaf, then it is representable. The arguments are largely formal, so to make this structure more transparent we begin the discussion for an arbitrary category enriched over S and later specialize to the category of manifolds. These results are mostly a repackaging of Morel-Voevodsky [15], see also Herrmann-Strunk [10].

Notation 2.6. Let **D** be an essentially small category enriched in spaces. Let  $\mathbf{D}_0$  denote the underlying (discrete) category of **D** and  $i: \mathbf{D}_0 \hookrightarrow \mathbf{D}$  the inclusion. We require **D** to have finite products and that the product functor  $\times$  makes **D** tensored over  $\Delta$ . That is, for each n there exists an object  $X \times \mathbb{A}^n \in \mathbf{D}_0$  characterized by the property that the set of n-simplices of the mapping space  $\max(X, Y)$  is identified with  $\max_{\mathbf{D}_0} (X \times \mathbb{A}^n, Y)$ , natural in  $X, Y \in \mathbf{D}$  and  $n \in \Delta$ .

In this setting it makes sense to talk about concordance invariant presheaves on  $\mathbf{D}_0$ .

**Definition 2.7.** Given a category  $\mathbf{D}_0$  as in Notation 2.6, a presheaf  $F: \mathbf{D}_0^{\mathrm{op}} \to \mathsf{S}$  is concordance invariant if the map induced by the projection

$$F(X) \to F(X \times \mathbb{A}^1)$$

is a weak equivalence of spaces for all  $X \in \mathbf{D}_0$ .

**Definition 2.8.** Given a category  $\mathbf{D}_0$  as in Notation 2.6, the *concordance resolution* of a functor  $F:(\mathbf{D}_0)^{\mathrm{op}}\to\mathsf{S}$  is the functor

$$F(-\times \mathbb{A}^{\bullet}): (\mathbf{D}_0)^{\mathrm{op}} \to \mathsf{sS}, \qquad X \mapsto F(X \times \mathbb{A}^{\bullet}).$$

We denote the homotopy colimit of  $F(-\times \mathbb{A}^{\bullet})$  by

$$\mathbf{B} F(X) := \operatornamewithlimits{hocolim}_{[n] \in \Delta^{\operatorname{op}}} F(X \times \mathbb{A}^n) = \big| F(X \times \mathbb{A}^{\bullet}) \big|.$$

Here we use the diagonal of a bisimplicial set as a model for the homotopy colimit over  $\Delta^{op}$ .

For the case  $\mathbf{D}_0 = \mathsf{Man}$  this definition of  $\mathbf{B}F$  coincides with Definition 2.5.

**Proposition 2.9.** Given a category  $\mathbf{D}_0$  as in Notation 2.6, for any presheaf  $F: \mathbf{D}_0^{\mathrm{op}} \to \mathsf{S}$ , the functor  $F(-\times \mathbb{A}^{\bullet})$  lifts to an enriched functor on  $\mathbf{D}$ .

*Proof.* We will define a simplicial map

$$\operatorname{map}(X,Y) \to \operatorname{map}(F(Y \times \mathbb{A}^{\bullet}), F(X \times \mathbb{A}^{\bullet})).$$

Let  $g: X \times \mathbb{A}^n \to Y$  be a morphism in  $\mathbf{D}_0$ . Given a morphism  $\alpha: [k] \to [n]$  in  $\Delta$ , consider the composition

$$F(Y \times \mathbb{A}^k) \xrightarrow{F(g \times \mathrm{id}_{\mathbb{A}^k})} F(X \times \mathbb{A}^n \times \mathbb{A}^k) \xrightarrow{F(\mathrm{id}_X \times \mathbb{A}^\alpha \times \mathrm{id}_{\mathbb{A}^k})} F(X \times \mathbb{A}^k \times \mathbb{A}^k) \xrightarrow{F(\mathrm{id}_X \times d)} F(X \times \mathbb{A}^k),$$

where  $d: \mathbb{A}^k \to \mathbb{A}^k \times \mathbb{A}^k$  is the diagonal map. This is functorial in g and  $\alpha$  and so defines a map between hom-sets

$$hom(X \times \mathbb{A}^n, Y) \to hom(F(Y \times \mathbb{A}^{\bullet}) \times \Delta[n], F(X \times \mathbb{A}^{\bullet}))$$
 (2.10)

for each  $n \geq 0$ . Therefore  $F(-\times \mathbb{A}^{\bullet})$  is enriched over spaces.

Corollary 2.11. Given a category  $\mathbf{D}_0$  as in Notation 2.6, for any presheaf  $F: \mathbf{D}_0^{\mathrm{op}} \to \mathsf{S}$ , the presheaf  $\mathbf{B}F$  is enriched over spaces and is concordance invariant (Definition 2.7).

*Proof.* To see that  $\mathbf{B}F$  is enriched, post-compose (2.10) with the homotopy colimit functor (alias geometric realization or diagonal) and use the fact that it commutes with products. In particular,  $\mathbf{B}F$  sends smooth homotopies to simplicial homotopies and smooth homotopy equivalences to simplicial homotopy equivalences. Hence it is concordance invariant (Definition 2.7).

Remark 2.12. The functor B is homotopy left adjoint to the discretization functor

$$i^* : \mathsf{PSh}(\mathbf{D}) \to \mathsf{PSh}(\mathbf{D}_0),$$

given by the restriction along the inclusion  $i : \mathbf{D}_0 \to \mathbf{D}$ . This follows from the fact that **B** is a left adjoint functor whose value on a representable presheaf on  $X \in \mathbf{D}_0$  is the representable presheaf on  $i(X) \in \mathbf{D}$ . In Corollary 2.14, we refine this statement to a Quillen equivalence of model categories.

The following proposition implies that the category of enriched presheaves on  $\mathbf{D}$  and the category of concordance-invariant presheaves (Definition 2.7) on  $\mathbf{D}_0$  have equivalent homotopy theories.

**Proposition 2.13.** Given a category  $\mathbf{D}_0$  as in Notation 2.6, a presheaf  $F: \mathbf{D}_0^{\mathrm{op}} \to \mathsf{S}$  is concordance invariant (Definition 2.7) if and only if the map  $F(X) \to i^* \mathbf{B} F(X)$  is a weak equivalence for all X.

*Proof.* If F is concordance invariant then the simplicial object  $F(X \times \mathbb{A}^{\bullet})$  is homotopically constant with value F(X). For the converse, consider the diagram

$$F(X) \xrightarrow{} i^* \mathbf{B} F(X)$$

$$\downarrow \qquad \qquad \downarrow$$

$$F(X \times \mathbb{A}^1) \xrightarrow{} i^* \mathbf{B} F(X \times \mathbb{A}^1)$$

The horizontal maps are weak equivalences by assumption. The vertical map on the right is a weak equivalence since  $\mathbf{B}F$  is concordance invariant. Thus, by the 2-out-of-3 property the left map is a weak equivalence.

Corollary 2.14. The restriction functor

$$i^*: \mathsf{PSh}(\mathbf{D}) o \mathsf{PSh}(\mathbf{D}_0)$$

is a right Quillen equivalence, where  $\mathsf{PSh}(\mathbf{D})$  is equipped with the projective model structure and  $\mathsf{PSh}(\mathbf{D}_0)$  is equipped with the  $\mathbb{A}^1$ -invariant projective model structure, i.e., the left Bousfield localization of the projective model structure with respect to the map  $\mathbb{A}^1 \to \mathbb{A}^0$ .

#### 2.a Concordance-invariant ∞-sheaves on manifolds are representable

**Definition 2.15.** A presheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  is an  $\infty$ -sheaf if for every manifold M and open cover  $\{U_i \to M\}_{i \in I}$ , the canonical map from F(M) to the homotopy limit (over  $\Delta$ ) of the cosimplicial space

$$\prod_{i_0 \in I} F(U_{i_0}) := \prod_{i_0, i_1 \in I} F(U_{i_0} \cap U_{i_1}) \cdots$$

is a weak equivalence of spaces.

Any  $\infty$ -sheaf F satisfies  $F(\emptyset) \simeq *$ . This is implied by the descent condition for the empty cover of the empty manifold.

**Remark 2.16.** A set-valued sheaf is an  $\infty$ -sheaf. A *stack is* a groupoid-valued  $\infty$ -sheaf. Common alternative terminologies for  $\infty$ -sheaves include  $\infty$ -stacks and homotopy sheaves.

The following proposition is due to Morel-Voevodsky [15] and Dugger [6].

**Proposition 2.17.** Given a presheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$ , the presheaf  $\mathbf{B}F$  is an  $\infty$ -sheaf if and only if the evaluation map

$$\mathbf{B}F(M) \to \operatorname{map}(\operatorname{map}(*, M), \operatorname{B}F)$$

is a weak equivalence for every M, where the evaluation map is the adjoint to the simplicial map

$$\operatorname{map}(*, M) \to \operatorname{map}(\mathbf{B}F(M), \mathbf{B}F(*)) \to \operatorname{map}(\mathbf{B}F(M), \mathbf{B}F(*)) \to \operatorname{map}(\mathbf{B}F(M), \mathbf{B}F)$$

gotten by the enrichment afforded by Corollary 2.11 and the map  $\mathbf{B}F(*) \to \mathbf{B}F$  being the Kan complex replacement of  $\mathbf{B}F(*)$ . Here  $\max(*, M)$  is isomorphic to  $\mathrm{Sing}\,M$ , the singular simplicial set of M.

*Proof.* Take a good open cover  $\{U_i\}_{i\in I}$  of M and let  $\mathbf{U}_{\bullet}:\Delta^{\mathrm{op}}\to\mathsf{Man}$  denote its Čech nerve. There is a commutative square:

$$\mathbf{B}F(M) \xrightarrow{} \mathbf{R}\operatorname{map}(\operatorname{Sing} M, \mathbf{B}F(*))$$

$$\downarrow \qquad \qquad \downarrow$$

$$\operatorname{holim}_{n \in \Delta} \mathbf{B}F(\mathbf{U}_n) \xrightarrow{} \operatorname{holim}_{n \in \Delta} \mathbf{R}\operatorname{map}(\operatorname{Sing} \mathbf{U}_n, \mathbf{B}F(*)),$$

where Sing denotes the singular simplicial set functor. The right-hand vertical arrow is an equivalence since  $\operatorname{\mathbf{B}} F(V) \simeq \operatorname{\mathbf{B}} F(*)$  for V contractible (by concordance invariance of  $\operatorname{\mathbf{B}} F$ ). The statement now follows by the 2-out-of-3 property.

### 2.b Homotopy groups of BF

**Proposition 2.18.** For an  $\infty$ -sheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  (Definition 2.15) with a point  $x \in F(*)$ , the map

$$\pi_0 \mathbf{B} F(S^d)_* \to \pi_d \mathbf{B} F(*) = \pi_d(\mathbf{B} F)$$

is an isomorphism. Here  $S^d = \{x \in \mathbb{R}^{d+1} \mid \sum_i x_i^2 = 1\} \in \mathsf{Man}$  is the smooth d-dimensional sphere, with a base point  $b \in S^d$ , and  $\mathbf{B}F(S^d)_*$  denotes the homotopy fiber of the map  $\mathbf{B}F(b) : \mathbf{B}F(S^d) \to \mathbf{B}F(*)$  over the image of x in  $\mathbf{B}F(*)$ .

*Proof.* By Theorem 1.1, the top map in the commutative square

$$\mathbf{B}F(S^d) \longrightarrow \mathbf{R}\operatorname{map}(S^d, \mathbf{B}F(*))$$

$$\mathbf{B}F(b) \downarrow \qquad \qquad \downarrow \mathbf{R}\operatorname{map}(b, \mathbf{B}F(*))$$

$$\mathbf{B}F(*) \longrightarrow \mathbf{R}\operatorname{map}(*, \mathbf{B}F(*))$$

is a weak equivalence and the bottom map is a weak equivalence by construction. Thus the induced map of vertical homotopy fibers over a point  $x \in F(*)$  is a weak equivalence. Taking  $\pi_0$  of the map between homotopy fibers, we obtain the desired claim.

**Remark 2.19.** Elements in  $\pi_0 \mathbf{B} F(S^d)_*$  are concordance classes of sections of F over  $S^d$  which restrict to x on  $b \in S^d$ . We postpone the explanation to Remark 4.12.

### 3 Weak Kan fibrations

As a warm-up to the ideas in this section, we will prove that the concordance relation  $\sim_c$  is an equivalence relation when F is an  $\infty$ -sheaf. This generalizes the standard fact that smooth homotopy is an equivalence relation, but the core of the argument is identical: gluing a pair of smooth maps along an open submanifold yields a smooth map.

**Lemma 3.1.** If  $F : \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  is an  $\infty$ -sheaf (Definition 2.15), then  $\sim_c$  is an equivalence relation on  $F(M)_0$ , the set of 0-simplices in F(M).

Proof. Reflexivity and symmetry are obvious. To establish transitivity, suppose  $\sigma_0$ ,  $\sigma_1$  and  $\sigma_2$  are such that  $\sigma_0 \sim_c \sigma_1$  and  $\sigma_1 \sim_c \sigma_2$ . Let  $i_k$  denote the inclusion of  $M \times \{k\}$  into  $M \times \mathbb{A}^1$  and pick sections  $\sigma_{01}$  and  $\sigma_{12}$  over  $M \times \mathbb{A}^1$  such that  $i_0^*\sigma_{01} = \sigma_0$ ,  $i_1^*\sigma_{01} = i_0^*\sigma_{12} = \sigma_1$  and  $i_1^*\sigma_{12} = \sigma_2$ . Take a smooth map  $r : \mathbb{A}^1 \to \mathbb{A}^1$  which fixes 0 and 1 and maps the complement of a small neighborhood of 1/2 to  $\{0,1\}$ . The sections  $r^*\sigma_{01}$  and  $r^*\sigma_{12}$  over  $M \times \mathbb{A}^1$  are such that the restriction of  $r^*\sigma_{01}$  to an open neighborhood of  $[1,\infty)$  agrees with the restriction of  $r^*\sigma_{12}$  to an open neighborhood of  $[-\infty,0]$ . So, using the sheaf property and reparametrizing, we may glue these sections to obtain a section  $\sigma_{012}$  over  $M \times \mathbb{A}^1$  with  $i_0^*\sigma_{012} = \sigma_0$  and  $i_1^*\sigma_{012} = \sigma_2$ , i.e.,  $\sigma_0 \sim_c \sigma_2$ .

The fact that  $\sim_c$  is an equivalence relation is a shadow of an important property possessed by the concordance resolution: it is a  $\theta$ -weak Kan complex (Definition 3.9). Informally, the weak nature can be seen in the proof of transitivity at the point where sections  $\sigma_{01}$  and  $\sigma_{12}$  are replaced by  $r^*\sigma_{01}$  and  $r^*\sigma_{12}$ . This step is essential since sections cannot be glued along closed sets. The failure of gluing along closed sets also means that concordance resolution does not satisfy the usual Kan condition as it does not have the right lifting property with respect to  $\Lambda_1^2 \to \Delta^2$ . Similar features of smooth geometry allow us to show that certain restriction maps for the concordance resolution have analogous weak fibrancy properties. The key definition formalizing this property is that of a weak Kan fibration.

## 3.a Kan fibrations and weak Kan fibrations of simplicial spaces

In this section, we define and investigate weak Kan fibrations of simplicial spaces (or sets). These generalize Kan fibrations and are related to (and inspired by) Dold fibrations of topological spaces [4]. We refer the reader to Appendix A for background on simplicial spaces.

**Definition 3.2.** Let  $f: X \to Y$  be a Reedy fibration between Reedy fibrant simplicial spaces. We say that f is a *Kan fibration* if it has the right lifting property with respect to all horn inclusions (Definition 2.2). That is, if for every solid square

$$\Lambda_k[n] \xrightarrow{X} X$$

$$\downarrow \qquad \qquad \downarrow f$$

$$\Delta[n] \xrightarrow{X} Y$$

there is a lift as pictured, where  $n \ge 1$  and  $0 \le k \le n$ . Similarly, we say that f is a trivial Kan fibration if it has the right lifting property with respect to  $\partial \Delta[n] \to \Delta[n]$  for all  $n \ge 0$ .

**Definition 3.3.** Let  $f: X \to Y$  be a Reedy fibration between Reedy fibrant simplicial spaces. We say that f has the weak right lifting property (weak RLP) with respect to a map  $i: A \hookrightarrow B$  (and i has the weak LLP with respect to f) if for every commutative square


$$\begin{array}{ccc}
A & \xrightarrow{\beta} X \\
\downarrow i & \stackrel{\widetilde{\alpha}}{\searrow} \stackrel{\nearrow}{\searrow} & \downarrow f \\
B & \xrightarrow{\alpha} Y
\end{array}$$

there is a lift  $\tilde{\alpha}$  as pictured, making the lower triangle commute strictly and the upper triangle commute up to a specified vertical homotopy. Such a homotopy consists of a map of simplicial spaces

$$H: A \times \Delta[1] \to X$$

subject to the requirement that  $H_0 = \beta$ ,  $H_1 = \widetilde{\alpha} \circ i$ , and  $f \circ H = \alpha \circ i \circ \pi$ , where  $\pi$  denotes the projection of  $A \times \Delta[1]$  onto A.

**Remark 3.4.** It will be useful to have some reformulations of Definition 3.3. Under the Reedy fibrancy hypothesis above, the map  $f: X \to Y$  has the weak RLP with respect to  $i: A \to B$  if and only if for every commutative square in the background:



there exists a map from the mapping cylinder  $M(i) = B \sqcup_i (A \times \Delta[1])$  to X making the diagram commute strictly. Here the map in the foreground  $M(i) \to B$  collapses  $A \times \Delta[1]$  to A – we denote it by  $\pi(i)$ .

To put it differently, consider the category  $sS^{[1]} = Fun(0 \to 1, sS)$  whose objects are maps of simplicial spaces and morphisms are commutative squares. Then the requirement above is that the map induced by precomposition

$$\operatorname{map}\begin{pmatrix} M(i) & X \\ \downarrow & , \downarrow \\ B & Y \end{pmatrix} \to \operatorname{map}\begin{pmatrix} A & X \\ \downarrow & , \downarrow \\ B & Y \end{pmatrix}$$

(with the square on the left in the diagram above) is surjective. Here map denotes the space of morphisms in  $\mathsf{sS}^{[1]}$ .

The Reedy model structure appears in the previous definitions as an artifact that guarantees homotopy invariance with respect to degreewise weak homotopy equivalences of simplicial spaces. But it is possible (and worthwhile) to formulate a more homotopical definition of the weak RLP.

**Definition 3.5.** A map  $f: X \to Y$  between arbitrary simplicial spaces satisfies the *weak right lifting* property (weak RLP) with respect to a map  $i: A \to B$  if

$$\mathbf{R}$$
map $(\pi(i), f) \to \mathbf{R}$ map $(i, f)$ 

is surjective on  $\pi_0$ , where **R**map refers to the derived mapping space computed in the category  $sS^{[1]}$  with objectwise weak equivalences.

**Remark 3.6.** We emphasize the homotopy invariance properties of this definition: a map f has the weak RLP with respect to an map i if and only if it has the weak RLP with respect to any map (degreewise) weakly equivalent to i. Also, if a map f has the weak RLP with respect to i then so does any map (degreewise) weakly equivalent to f.

**Proposition 3.7.** A Reedy fibration f satisfies the weak RLP in the sense of Definition 3.3 if and only if it satisfies the weak RLP in the sense of Definition 3.5.

*Proof.* Equip sS with the Reedy (= injective) model structure and, relative to it, also equip  $sS^{[1]}$  with the injective model structure. In this model structure on  $sS^{[1]}$ , all objects are cofibrant. Cofibrations are morphisms which are objectwise Reedy cofibrations of simplicial spaces, i.e., degreewise injections. Fibrant objects are Reedy fibrations between Reedy fibrant simplicial spaces.

The morphism  $i \to \pi(i)$ , i.e., the commutative diagram

$$\begin{array}{ccc}
A & \longrightarrow & M(i) \\
\downarrow & & \downarrow \\
B & \longrightarrow & B
\end{array}$$

is a cofibration since the horizontal maps are degreewise injections. Also, the map f is a fibrant object in  $\mathsf{sS}^{[1]}$ . Therefore, the induced map

$$map(\pi(i), f) \to map(i, f)$$

is a fibration between Kan complexes, which is weakly equivalent to

$$\mathbf{R}$$
map $(\pi(i), f) \to \mathbf{R}$ map $(i, f)$ .

The result now follows, since a fibration of simplicial sets is surjective on  $\pi_0$  if and only if it is surjective on 0-simplices.

**Definition 3.8.** A map  $X \to Y$  between simplicial spaces is a *weak Kan fibration* if it has the weak right lifting property with respect to the maps

$$h_i: \mathrm{Sd}^i(\Lambda_k[n]) \hookrightarrow \mathrm{Sd}^i(\Delta[n])$$

for each  $i \ge 0$ ,  $n \ge 1$  and  $0 \le k \le n$ . Here the functor Sd:  $sS \to sS$  is as in Definition A.12.

The definition also makes sense for maps  $X \to Y$  of simplicial sets by regarding them as simplicial discrete spaces.

Note that every simplicial space is a weak Kan complex, in the sense that the map  $X \to *$  is a weak Kan fibration. This may seem odd at first, but it makes sense in light of Section 3.b, as every space is tautologically quasi-fibrant. A more interesting variation is

**Definition 3.9.** Given  $\ell \geq -1$ , a  $\ell$ -weak Kan complex is a simplicial space X such that the terminal map  $X \to *$  is a weak Kan fibration in which the vertical homotopies preserve the  $\ell$ -skeleton of the horn inclusions  $h_i$ .

So a (-1)-weak Kan complex is simply a weak Kan complex, i.e., a simplicial space, and a  $\infty$ -weak Kan complex is a Kan complex (in the usual sense, as in Definition 3.2). For a 0-weak Kan complex X, the image of  $\pi_0 X_1 \to \pi_0 X_0 \times \pi_0 X_0$  is an equivalence relation (c.f. Lemma 3.1). We will not make use of the notion of  $\ell$ -weak Kan complex for  $\ell > 0$ , and for  $\ell = 0$  we will use it more as an illustration of arguments.

**Example 3.10.** A Kan fibration (Definition 3.2) is a weak Kan fibration. Of course, the usual definition of a Kan fibration does not mention subdivisions; this is because a map satisfying the strict RLP against all horn inclusions  $\Lambda_k[n] \to \Delta[n]$  automatically satisfies the same property against all subdivisions of those, since these subdivided horn inclusions can be presented using cobase changes and the strict LLP is stable under cobase change, so the strict LLP for horn inclusions implies the strict LLP for subdivided horn inclusions. The same is *not* true for the weak RLP, so we need to take subdivisions seriously.

Likewise, if X is a  $\ell$ -weak Kan complex and  $K \hookrightarrow L$  is an inclusion of simplicial sets, it does not follow automatically that  $X^L \to X^K$  is a weak Kan fibration.

**Lemma 3.11.** Kan's Ex functor (Definition A.12) preserves weak Kan fibrations.

*Proof.* The functor Ex is right adjoint to the subdivision functor, so we are investigating a square of the form

$$\operatorname{Sd}^{i+1} \Lambda_{k}[n] \xrightarrow{X} X$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$\operatorname{Sd}^{i+1} \Delta[n] \xrightarrow{Y} Y$$

$$(3.12)$$

Since f is a weak Kan fibration, there is a lift as shown together with a homotopy  $H: \mathrm{Sd}^{i+1} \Lambda_k[n] \times \Delta[1] \to X$ . Since there is always a map from the subdivision of the product to the product of the subdivisions, we can precompose H with

$$\operatorname{Sd}(\operatorname{Sd}^i \Lambda_k[n] \times \Delta[1]) \to \operatorname{Sd}^{i+1} \Lambda_k[n] \times \operatorname{Sd} \Delta[1] \to \operatorname{Sd}^{i+1} \Lambda_k[n] \times \Delta[1].$$

This gives the required homotopy for the upper triangle which is vertical over Y.

**Remark 3.13.** Weak Kan fibrations are stable under various operations. They are stable under *homotopy* base change (with respect to degreewise weak equivalences of simplicial spaces). In other words, weak Kan fibrations that are moreover Reedy fibrations between Reedy fibrant objects are stable under pullback.

Weak Kan fibrations are also stable under fiberwise homotopy retracts (that is, if  $g:W\to Y$  is a homotopy retract over Y of a weak Kan fibration  $f:X\to Y$  then g is a weak Kan fibration). In particular, weak Kan fibrations are stable under fiberwise homotopy equivalences. Moreover, if we allow subdivisions of the vertical homotopies in the definition of the weak lifting property, i.e., if we replace  $\Delta[1]$  by  $\mathrm{Sd}^i\,\Delta[1]$  for  $i\geq 0$  in Definition 3.3, then a composition of two weak Kan fibrations is also a weak Kan fibration. Since these properties will not be used in what follows, and the proofs are not particularly difficult, we omit further explanations.

#### 3.b Weak Kan fibrations are realization fibrations

**Definition 3.14** (Rezk [18]). A map  $f: X \to Y$  of simplicial spaces is a realization fibration if for every  $Z \to Y$  the induced map

$$|X \times_Y^h Z| \to |X| \times_{|Y|}^h |Z| \tag{3.15}$$

is a weak equivalence of spaces. The vertical bars refer to the diagonal simplicial set, which models the homotopy colimit over  $\Delta^{op}$ .

**Remark 3.16.** Realization fibrations are related to quasi-fibrations in the sense of Dold–Thom [5]. For example, if a map  $f: X \to Y$  between simplicial *sets* is a realization fibration, then the map (3.15) with Z a point is identified with the inclusion of the fiber of f into the homotopy fiber. That is, f is a quasi-fibration. On the other hand, not all quasi-fibrations are realization fibrations: realization fibrations are stable under homotopy pullback, whereas quasi-fibrations need not be.

We now turn to the main technical result of the section.

**Theorem 3.17.** A weak Kan fibration is a realization fibration.

As we already emphasized, the subdivisions of simplices and horns in the definition of a weak Kan fibration are important for a number of reasons. Lemma 3.11 is one such reason, that will be exploited later on. Below is another.

**Example 3.18.** This is an example of a map which has the weak right lifting property against the (non-subdivided) map  $(h_0)$  but is not a realization fibration. Suppose X is the union of three non-degenerate 1-simplices as in the picture:



and  $Y = \Delta[1]$ . Let  $f: X \to Y$  be the projection in the vertical direction. This is not a Kan fibration: there are 2-horns  $\Lambda^k[2]$  in X that cannot be filled. On the other hand, f has the weak right lifting property with respect to the map  $(h_0)$ . But f is not a quasi-fibration, since at one point the fiber is disconnected while the homotopy fiber is not. Hence f cannot be a realization fibration. This is not a contradiction: f is not a weak Kan fibration as it does not have the weak right lifting property for  $(h_2)$ , the second subdivision of the horn inclusion.

**Proposition 3.19.** Suppose  $f: X \to Y$  is a Reedy fibration between Reedy fibrant simplicial spaces. Then f is a weak Kan fibration of simplicial spaces if and only if  $|f|:|X|\to |Y|$  is a weak Kan fibration of simplicial sets. The vertical bars denote the diagonal functor.

*Proof.* Let us denote by  $\kappa$  the map  $\operatorname{sd}^i \Lambda_k^n \to \operatorname{sd}^i \Delta^n$  for  $i \geq 0$ . Write  $\delta$  for the functor which sends a simplicial set K to the corresponding simplicial discrete space  $[n] \mapsto K_n$ . Recall the commutative square of spaces  $\kappa \to \pi(\kappa)$  where  $\pi(\kappa)$  is the projection of mapping cylinder  $M(\kappa)$  onto  $\operatorname{sd}^i \Delta^n$ . By Definition 3.8, the map f is a weak Kan fibration if and only if

$$\mathbf{R}$$
map $(\delta(\pi(\kappa)), f) \to \mathbf{R}$ map $(\delta(\kappa), f)$ 

is surjective on  $\pi_0$ .

The realization (i.e., diagonal) functor has a left adjoint  $d_!: S \to sS$ , and there is a natural transformation  $d_! \to \delta$  which is a weak equivalence (Lemma A.8). Therefore, f is a weak Kan fibration if and only if

$$\mathbf{R}$$
map $(d_!\pi(\kappa), f) \to \mathbf{R}$ map $(d_!\kappa, f)$ .

is surjective on  $\pi_0$ . Regarding  $sS^{[1]}$  with the injective model structure, the map  $d_!\kappa \to d_!\pi(\kappa)$  is a cofibration between cofibrant objects (since  $d_!$  sends monomorphisms to monomorphisms), and the target f is fibrant by hypothesis. As such, the above holds if and only if the Kan fibration between Kan simplicial sets

$$\operatorname{map}(d_!\pi(\kappa), f) \to \operatorname{map}(d_!(\kappa), f)$$

is surjective. By adjunction, this holds if and only if |f| has the weak RLP with respect to  $\kappa$ . (Note that |f|, being a map between simplicial discrete spaces, is automatically a Reedy fibration between Reedy fibrant objects, and so a fibrant object in  $sS^{[1]}$ .)

**Remark 3.20.** The same proof, with  $\kappa$  of the form  $\Lambda_k^n \to \Delta^n$  and with  $\pi(\kappa)$  replaced by the identity  $\Delta^n \to \Delta^n$ , shows that a Reedy fibration between Reedy fibrant simplicial spaces f is a (trivial) Kan fibration if and only if |f| is a (trivial) Kan fibration.

In order to prove that weak Kan fibrations are realizations fibrations, we will use the following criterion due to Rezk.

**Theorem 3.21** (Rezk [18]). A map  $f: X \to Y$  of simplicial spaces is a realization fibration if and only if for all maps  $\Delta[m] \to Y$  and  $\Delta[0] \to \Delta[m]$ , the induced map on pullbacks

$$X \times_Y^h \Delta[0] \to X \times_Y^h \Delta[m]$$


is a weak equivalence after realization.

Proof of Theorem 3.17. Let  $f: X \to Y$  be a weak Kan fibration. We will verify that f satisfies the condition in Theorem 3.21. We may assume, without loss of generality, that f is a Reedy fibration between Reedy fibrant simplicial spaces. Then the homotopy pullbacks above become pullbacks. Using the fact that realization and  $\operatorname{Ex}^{\infty}$  commute with finite limits, our task is then to show that

$$\operatorname{Ex}^{\infty} |X| \times_{\operatorname{Ex}^{\infty} |Y|} \operatorname{Ex}^{\infty} \Delta^{0} \to \operatorname{Ex}^{\infty} |X| \times_{\operatorname{Ex}^{\infty} |Y|} \operatorname{Ex}^{\infty} \Delta^{m}$$

is a weak equivalence of Kan complexes. Since f is a weak Kan fibration, the same can be said of |f| by Proposition 3.19 and of  $\operatorname{Ex}^{\infty}|f|$  by Lemma 3.11. To simplify the notation, let us write  $g:U\to V$  for  $\operatorname{Ex}^{\infty}|f|:\operatorname{Ex}^{\infty}|X|\to\operatorname{Ex}^{\infty}|Y|$ .

In view of Proposition A.1 (and Corollary A.2 and Example A.5), we will show that for every solid diagram



there are dashed maps as pictured. Let us write  $A \to B$  for the middle vertical arrow. Consider the map

$$\Lambda^{n+1} \sqcup_{\partial \Delta^n \times \{0\}} A \to \operatorname{Ex}^{\infty} \Delta^m \tag{3.22}$$

determined by the lower horizontal map in the diagram above and by the map  $A \to \Delta^0 \to \operatorname{Ex}^\infty \Delta^m$ . Since the terminal map  $\operatorname{Ex}^\infty \Delta^m \to \Delta^0$  is a trivial Kan fibration, the map (3.22) extends along the inclusion

$$\Lambda^{n+1} \sqcup_{\partial \Delta^n \times \{0\}} A \hookrightarrow B.$$

Next, we want to define a map  $B \to U$  which is compatible with the composition

$$B \to \operatorname{Ex}^{\infty} \Delta^m \to V$$

of the map we have just constructed. This will give us the lower dashed map in the diagram above. Consider the diagram

$$\Lambda^{n+1} \longrightarrow U \\
\downarrow \qquad \qquad \downarrow g \\
\Delta^{n+1} \longrightarrow V$$

where the lower map is the composition  $\Delta^{n+1} \to B \to \operatorname{Ex}^{\infty} \Delta^m \to V$ . Since g is a weak Kan fibration, we obtain a lift and vertical homotopy, i.e., the required map  $B \to U$ . Therefore, we have defined a map  $B \to U \times_V \operatorname{Ex}^{\infty} \Delta^m$ , whose restriction to A factors through  $U \times_V \Delta^0$ . This completes the proof.

# 4 Weak Kan fibrancy of the concordance resolution

Let  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  be an  $\infty$ -sheaf. In this section and the next, we apply the general theory developed in the previous sections to prove Theorem 1.2. The goal of this section is to prove the following three propositions.

**Proposition 4.1.** For any  $\infty$ -sheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  (Definition 2.15), the simplicial space  $[n] \to F(\mathbb{A}^n)$  is a 0-weak Kan complex (Definition 3.9).

**Proposition 4.2.** For any  $\infty$ -sheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  (Definition 2.15), the map of simplicial spaces

$$F(\mathbb{A}^{\bullet} \times \mathbb{A}^1) \to F(\mathbb{A}^{\bullet} \times \partial \mathbb{A}^1)$$

is a weak Kan fibration (Definition 3.8). Here  $\partial \mathbb{A}^1 = \mathbb{A}^0 \sqcup \mathbb{A}^0$  is the disjoint union of two points.

**Proposition 4.3.** Let  $* \hookrightarrow S^n$  be an inclusion of a basepoint into the smooth *n*-dimensional sphere. For any  $\infty$ -sheaf  $F : \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  (Definition 2.15), the induced map of simplicial spaces

$$F(\mathbb{A}^{\bullet} \times S^n) \to F(\mathbb{A}^{\bullet})$$

is a weak Kan fibration (Definition 3.8).

We use these to prove Theorem 1.2 in Section 5, and the reader may wish to jump directly to that section to see these propositions in action. In fact, in that section and in the remainder of the paper, we will only need Propositions 4.2 and 4.3. We include Proposition 4.1 because its proof anticipates much of the proof of the other two propositions.

The proofs of these propositions are based on the following simple observation. If one were to try to prove that  $F(\mathbb{A}^{\bullet})$  is a *strict* Kan complex, an obvious approach would be to construct a deformation retraction from the simplex  $\mathbb{A}^n$  to its horn. This is possible in the topological and PL settings, but not possible smoothly. However, the basic idea can be salvaged if one only asks for  $F(\mathbb{A}^{\bullet})$  to be 0-weak Kan, which roughly translates into asking for a retraction up to a suitable homotopy. This parallels the proof that concordance is an equivalence relation (Lemma 3.1): we modify smooth maps between manifolds (via homotopies) to achieve certain constancy properties. A relative version of this line of reasoning applies to the maps in Propositions 4.2 and 4.3.

#### 4.a The sheaf associated to a simplicial set

**Definition 4.4.** Denote by

$$\|-\|_{\mathrm{pre}}: \mathsf{sS} \to \mathsf{PSh}(\mathsf{Man},\mathsf{S})$$

the simplicial left adjoint functor that sends  $\Delta[n]$  to the representable presheaf of  $\mathbb{A}^n$ . The corresponding simplicial right adjoint functor is

$$\mathsf{PSh}(\mathsf{Man},\mathsf{S})\to \mathsf{sS}, \qquad F\mapsto (n\mapsto F(\mathbb{A}^n)).$$

Namely, for a simplicial set K, the presheaf  $||K||_{\text{pre}} : \mathsf{Man}^{\text{op}} \to \mathsf{Sets}$  is given by the coend

$$K_n \otimes_{[n] \in \Delta} \operatorname{map}(-, \mathbb{A}^n).$$

**Remark 4.5.** The adjunction of Definition 4.4 is Quillen if both categories are equipped with the projective model structure or with the injective model structure. Therefore, there is a weak equivalence

$$\mathbf{R}$$
map $(\|K\|_{\mathrm{pre}}, F) \to \mathbf{R}$ map $(K, F(\mathbb{A}^{\bullet}))$ 

natural in the simplicial space K and the presheaf F.

**Remark 4.6.** Note that  $||K||_{\text{pre}}$  is usually not a sheaf of sets. For a simple example illustrating this, take  $||\Lambda_1^2||$  which is the pushout

$$\operatorname{map}(-,\mathbb{A}^1) \bigsqcup_{\operatorname{map}(-,\mathbb{A}^0)} \operatorname{map}(-,\mathbb{A}^1)$$

and pick an open cover of  $\mathbb{R}^1$  by two open sets and compatible sections over each that do not lift to a section over  $\mathbb{R}^1$ .

**Definition 4.7.** Given a simplicial set K, denote by ||K|| the associated sheaf of sets of  $||K||_{pre}$ .

Recall that for presheaves of *sets*, the notions of a sheaf and  $\infty$ -sheaf, as well as sheafification and  $\infty$ -sheafification, coincide. Thus, in the above definition, the sheafification functor can be replaced by  $\infty$ -sheafification. If we tried to define ||K|| for simplicial *spaces* K, we would have to use  $\infty$ -sheafification from the start.

**Remark 4.8.** Since sheafification is left adjoint to the inclusion of sheaves into presheaves, for F an  $\infty$ -sheaf, the weak equivalence above lifts to a weak equivalence

$$\mathbf{R}$$
map $(\|K\|, F) \xrightarrow{\simeq} \mathbf{R}$ map $(K, F(\mathbb{A}^{\bullet}))$ 

natural in the simplicial space K and the  $\infty$ -sheaf F.

Remark 4.9. To be more concrete, suppose N is a smooth manifold and K is a smooth triangulation of N. From the description above, we see that a section of ||K|| over a manifold M can be described as a smooth map  $M \to N$  which factors locally through some simplex of the triangulation. Alternatively, such a section is the data of an open cover  $\{U_i\}$  of M and sections of  $||K||_{\text{pre}}$  over each  $U_i$ , i.e., smooth maps whose image is in some extended simplex of K, which are compatible on overlaps  $U_i \cap U_j$ . Two such sets of data are identified if after passing to some refined open cover the individual sections of  $||K||_{\text{pre}}$  become equal.

**Proposition 4.10.** Let F be an injectively fibrant object of PSh(Man, S). Then the maps in Propositions 4.2 and 4.3 are Reedy fibrations between Reedy fibrant simplicial spaces.

*Proof.* We show that the map in Proposition 4.2 is a Reedy fibration. The argument for the one in Proposition 4.3 is similar. Let  $A \to B$  be a trivial Reedy cofibration of simplicial spaces, i.e., a map of simplicial spaces which is a degreewise monomorphism and a degreewise weak equivalence. Using the adjunction of Definition 4.4, denote by Q the following pushout of presheaves


$$||A||_{\text{pre}} \times \partial \mathbb{A}^{1} \longrightarrow ||B||_{\text{pre}} \times \partial \mathbb{A}^{1}$$

$$\downarrow \qquad \qquad \downarrow$$

$$||A||_{\text{pre}} \times \mathbb{A}^{1} \longrightarrow Q$$

$$(4.11)$$

By adjunction,  $F(\mathbb{A}^{\bullet} \times \mathbb{A}^{1}) \to F(\mathbb{A}^{\bullet} \times \partial \mathbb{A}^{1})$  has the right lifting property with respect to  $A \to B$  if and only if every solid diagram of presheaves



has a lift as pictured. The top horizontal map in square (4.11) is a trivial cofibration by hypothesis. Since acyclic cofibrations are stable under cobase change, it follows that the lower horizontal map is also a trivial cofibration. But the composition

$$||A||_{\mathrm{pre}} \times \mathbb{A}^1 \to Q \to ||B||_{\mathrm{pre}} \times \mathbb{A}^1$$

is also a trivial cofibration by hypothesis, and so by two-out-of-three the right-hand map is a weak equivalence. The right-hand map is clearly an injective cofibration. Since F is injectively fibrant, we conclude that the dashed map exists.

**Remark 4.12.** Let  $* \hookrightarrow S^n$  be an inclusion of a basepoint into the smooth *n*-dimensional sphere. Recall (Proposition 4.3) that the induced map

$$F(\mathbb{A}^{\bullet} \times S^n) \to F(\mathbb{A}^{\bullet})$$

is a weak Kan fibration. Applying the homotopy colimit functor to this map and then taking its homotopy fiber over a given basepoint  $b \in F(\mathbb{A}^0)$  yields the space  $\mathbf{B}F(S^d)_*$  of Proposition 2.18. Since weak Kan fibrations are realization fibrations (Theorem 3.17), the space  $\mathbf{B}F(S^d)_*$  is weakly equivalent to the homotopy colimit functor applied to the map

$$Z_{\bullet} := \mathrm{fiber}_b(F(\mathbb{A}^{\bullet} \times S^n) \to F(\mathbb{A}^{\bullet})),$$

where we assume F to be injectively fibrant so that fiber<sub>b</sub> computes the homotopy fiber over b (and the simplicial degenerations of b). Weak Kan fibrations are stable under base change, so the map  $Z_{\bullet} \to *$  is a weak Kan fibration and the relation on  $\pi_0 Z_0$  determined by the map  $\pi_0 Z_1 \to \pi_0 Z_0 \times \pi_0 Z_0$  is an equivalence relation (c.f. Lemma 3.1). Hence, the set of path components of  $\mathbf{B}F(S^d)_*$  is identified with the quotient of  $\pi_0 Z_0$  by this equivalence relation.

## 4.b Closed simplices

**Definition 4.13.** Denote by  $\Delta^n : \mathsf{Man}^{\mathrm{op}} \to \mathsf{Sets}$  the subsheaf of the representable sheaf of sets  $\mathrm{map}(-, \mathbb{A}^n)$  (where  $\mathbb{A}^n \in \mathsf{Man}$  is the *n*-dimensional extended smooth simplex), consisting of sections  $X \to \mathbb{A}^n$  whose image is contained in the closed simplex  $\mathbb{R}^{n+1}_{>0} \cap \mathbb{A}^n \subset \mathbb{A}^n$ .

**Definition 4.14.** Given a simplicial set K, denote by  $\overline{||K||}$ :  $\mathsf{Man}^{\mathrm{op}} \to \mathsf{Sets}$  the subsheaf of sets of ||K|| consisting of smooth maps  $U \to M$  that factor locally through some *closed* simplex of K in M. In formulas,

$$\overline{\|K\|} = L(\overline{\|K\|}_{\mathrm{pre}}) = L(K_n \otimes_{[n] \in \Delta} \Delta^n)$$

where  $\Delta^n$  is as in Definition 4.13, L(W) denotes the associated sheaf of a presheaf of sets W, and  $\overline{\|K\|}_{pre}$  denotes the left adjoint functor of  $F \mapsto \text{map}(\Delta^{\bullet}, F)$  applied to the simplicial set K.

**Lemma 4.15.** Suppose K is a simplicial set induced from a finite-dimensional semisimplicial set (given by a presheaf of sets on the category  $(\Delta_{\rm inj})_{\leq N}$  of simplices of dimension  $\leq N$  and their injective maps) by the left adjoint functor to the forgetful (restriction) functor from simplicial sets to semisimplicial sets. Then the inclusion  $\overline{\|K\|} \to \|K\|$  has a homotopy inverse, in the sense that there is a map  $\|K\| \times \mathbb{A}^1 \to \|K\|$  whose restriction to  $\|K\| \times \{0\}$  is the identity and whose restriction to  $\|K\| \times \{1\}$  factors through  $\overline{\|K\|}$ . An analogous statement holds for the inclusion  $\overline{\|K\|}_{\rm pre} \to \|K\|_{\rm pre}$ .

*Proof.* The condition on K allows us to replace the indexing category  $\Delta$  for coends with the category  $\Delta_{\rm inj}$ . The claimed homotopy inverse comes from homotopy inverses  $\mathbb{A}^n \to \mathbf{\Delta}^n$  that are functorial with respect to  $[n] \in \Delta_{\rm inj}$ . Namely, consider a homotopy

$$\lambda^n: \mathbb{A}^n \times [0,1] \to \mathbb{A}^n$$

that in barycentric coordinates is constructed as follows. Fix  $c: \mathbb{A}^1 \to \mathbb{A}^1$  a smooth function with  $c \equiv 0$  in a small neighborhood of  $(\infty, 0]$  and  $c \equiv 1$  in a small neighborhood of  $[1, \infty)$ . Then we set

$$\lambda^n((x_0,\ldots,x_n),t)=(y_0/C_t,\ldots,y_n/C_t)$$

where  $y_i = tc(x_i) + (1-t)x_i$  and  $C_t = \sum_{i=0}^n y_i$ . Extend  $\lambda^n$  to  $\mathbb{A}^n \times \mathbb{A}^1$  by precomposing  $\lambda^n$  with id  $\times c$ :  $\mathbb{A}^n \times \mathbb{A}^1 \to \mathbb{A}^n \times [0,1]$ . It is clear that this gives a homotopy equivalence for each fixed n. Moreover,  $\lambda$  is functorial with respect to injections  $[n] \to [m]$ . Hence it defines a homotopy equivalence  $\overline{\|K\|} \to \|K\|$ . The case of  $\overline{\|K\|}_{pre} \to \|K\|_{pre}$  is treated in the same manner.

**Proposition 4.16.** For any manifold M and any presheaf  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$ , the restriction map

$$\operatorname{Rmap}(M \times \mathbb{A}^{\bullet}, F) \to \operatorname{Rmap}(M \times \Delta^{\bullet}, F)$$

is a weak equivalence after realization.

*Proof.* We can assume F to be injectively fibrant, so **R**map can be replaced with map. By replacing F with map(M, F) we can assume  $M = \mathbb{R}^0$ . We will verify the conditions of Proposition A.9 and show that every square

$$\partial \Delta[n] \longrightarrow \operatorname{Ex}^{\infty} F(\mathbb{A}^{\bullet})$$

$$\downarrow \qquad \qquad \downarrow$$

$$\Delta[n] \longrightarrow \operatorname{Ex}^{\infty} \operatorname{map}(\Delta^{\bullet}, F)$$

admits a lift making the upper and lower triangles commute up to homotopy, and such that these two homotopies are compatible on  $\partial \Delta[n]$ . For a simplicial compact space K, a map  $K \to \operatorname{Ex}^{\infty} Y$  factors through some finite stage, and so it corresponds to a map  $\operatorname{Sd}^i K \to Y$ . Then, by adjunction, the square above amounts to a map  $P \to F$  where

$$P := \|\operatorname{Sd}^i \partial \Delta[n]\|_{\operatorname{pre}} \bigsqcup_{\overline{\|\operatorname{Sd}^i \partial \Delta[n]\|_{\operatorname{pre}}}} \overline{\|\operatorname{Sd}^i \Delta[n]\|_{\operatorname{pre}}}.$$

By Lemma 4.15, there is a self-homotopy H of  $\|\operatorname{Sd}^i \Delta[n]\|_{\operatorname{pre}}$  such that  $H_0 = \operatorname{id}$ ,  $H_1$  factors through

$$\overline{\|\operatorname{Sd}^i\Delta[n]\|}_{\operatorname{pre}} \to \|\operatorname{Sd}^i\Delta[n]\|_{\operatorname{pre}}$$

and H preserves  $\|\operatorname{Sd}^i\partial\Delta[n]\|_{\operatorname{pre}}$ ,  $\overline{\|\operatorname{Sd}^i\partial\Delta[n]\|_{\operatorname{pre}}}$  and  $\overline{\|\operatorname{Sd}^i\Delta[n]\|_{\operatorname{pre}}}$ . This defines a map

$$\|\operatorname{Sd}^i \Delta[n]\|_{\operatorname{pre}} \to P \to F$$

and a self homotopy of P, giving the lift and homotopies that were needed.

### 4.c Smooth maps with prescribed constancy conditions

**Proposition 4.17.** Suppose K is a simplicial set and  $J: ||K|| \to \Delta^n$  is a subdivision of  $\Delta^n$  (meaning K has finitely many nondegenerate simplices, the map J sends any nondegenerate simplex of K to an injective affine map into  $\underline{\Delta}^n$ , and the union of interiors of images of simplices in K is equal to the interior of  $\underline{\Delta}^n$ ). Denote by  $j: ||K|| \to \Delta^n$  the restriction of J along the inclusion  $||K|| \to \Delta^n$ . Then the map J has a homotopy inverse. More precisely, there is a map  $T: \underline{\Delta}^n \to ||K||$  and a smooth homotopy

$${h_t: \mathbb{A}^n \to \mathbb{A}^n}_{t \in [0,1]}$$

which restricts to a homotopy  $\Delta^n \times \mathbb{A}^1 \to \Delta^n$  between the identity and jr, and to a homotopy  $\overline{\|K\|} \times \mathbb{A}^1 \to \overline{\|K\|}$  between the identity and rj.

This is a consequence of the lemma below.

**Lemma 4.18.** Given K and J as in Proposition 4.17, there exists a smooth homotopy  $\{h_t : \mathbb{A}^n \to \mathbb{A}^n\}_{t \in [0,1]}$  such that

- (i)  $h_0$  is the identity,
- (ii)  $h_t$  maps each closed simplex  $\Delta^n \subset \overline{\|K\|}$  to itself for all t, and
- (iii) each closed simplex  $\Delta^n \subset \overline{\|K\|} \subset \mathbb{A}^n$  has a neighborhood in  $\mathbb{A}^n$  which gets mapped to that same simplex by  $h_1$ .

*Proof.* We use the following terminology during this proof: for a simplex  $\sigma$  of K, a homotopy of maps  $(f_t : \mathbb{A}^n \to \mathbb{A}^n)_{t \in [c,d]}$  satisfies property (iii) $_{\sigma}$  if  $\|\sigma\|$  has a neighborhood in  $\mathbb{A}^n$  which gets mapped to  $\|\sigma\|$  by  $f_d$ .

Fix some k with  $-1 \le k \le n$  and suppose per induction that we have already constructed a smooth homotopy  $(h_t : \mathbb{A}^n \to \mathbb{A}^n)_{t \in [0,a]}$  for some a < 1 satisfying conditions (i), (ii) and (iii) $_{\sigma}$  for every simplex  $\sigma$  of dimension at most k.

We want to extend this to a smooth homotopy  $(h_t : \mathbb{A}^n \to \mathbb{A}^n)_{t \in [0,b]}$ , where b > a, that satisfies conditions (i), (ii) and (iii) $_{\sigma}$  for every simplex  $\sigma$  of dimension at most k+1.

For a closed k-simplex  $\tau$ , let  $W_{\tau}$  be a neighborhood of  $\tau$  in  $\mathbb{A}^n$  which gets mapped to  $\tau$  by  $h_a$ . This exists by the inductive assumption. We shall define a homotopy

$$(g_t: \mathbb{A}^n \to \mathbb{A}^n)_{t \in [0,b-a]}$$

where  $g_0 = \operatorname{id}$ ,  $g_t$  maps each simplex of  $\overline{\|K\|}$  to itself for all t, and  $g_{b-a}$  maps a neighborhood in  $\mathbb{A}^n$  of a large enough subset of the interior of each (k+1)-simplex to that same simplex. Large enough means it should intersect  $W_{\tau}$  for each boundary face  $\tau \subset \|\sigma\|$ . Then define  $(h_t)_{t \in [0,b]}$  as the concatenation of  $(g_{t-a}h_a)_{t \in [a,b]}$  with  $(h_t)_{t \in [0,a]}$ . (In order for the concatenation to be smooth, we may arrange so that the homotopy  $(h_t)_{t \in [0,a]}$  is stationary for t close to a and the homotopy  $(g_t)$  is stationary for t close to a.)

For each (k+1)-simplex  $\|\sigma\|$ , choose a small tubular neighborhood  $U(\sigma)$  of  $\operatorname{int}(\|\sigma\|)$  in  $\mathbb{A}^n$  such that for each point  $x \in \operatorname{int}(\|\sigma\|)$  and every closed simplex  $\tau$  of  $\|K\|$ , the intersection  $U(\sigma)_x \cap \tau$  is a linear cone in  $\mathbb{A}^n$ . That is, there exist linearly independent vectors  $v_1, \ldots, v_\ell$  such that points in  $U(\sigma)_x \cap \tau$  are of the form

 $c_0v_0 + c_1v_1 + \cdots + c_\ell v_\ell$  where  $c_i \geq 0$ . By shrinking if necessary, we may also assume that  $U(\sigma) \cap U(\sigma')$  is empty if  $\sigma$  and  $\sigma'$  are distinct (k+1)-simplices.

Now we fix a  $\sigma$ , and take a map

$$\psi: U(\sigma) \to U(\sigma)$$

over  $\operatorname{int}(\sigma)$  satisfying the following conditions: for x close to the boundary of  $\sigma$ ,  $\psi_x$  is the identity; for x close to the barycenter,  $\psi_x(v) = 0$  for  $v \in U(\sigma)_x$  and |v| small and  $\psi_x(v) = v$  for |v| large. Extend by the identity to obtain a map  $g_1^{\sigma} : \mathbb{A}^n \to \mathbb{A}^n$ . Linearly interpolate between the identity and  $h_1^{\sigma}$  to get a homotopy  $(h_t^{\sigma})$  satisfying properties (i), (ii) and (iii) $\sigma$ . Now concatenate all  $(h_t^{\sigma})$ 's to obtain the required homotopy  $(h_t)$ .

Proof of Proposition 4.17. The lemma gives us a smooth homotopy h on  $\mathbb{A}^n$ . Condition (ii) implies that h restricts to a homotopy  $\{h_t : \overline{\|K\|} \to \overline{\|K\|}\}$ . Condition (iii) gives the required factorization of  $h_1$  as

$$\mathbf{\Delta}^n \xrightarrow{r} \overline{\|K\|} \xrightarrow{j} \mathbf{\Delta}^n$$

in the category of sheaves of sets, where the factorization of  $h_1$  through  $\overline{\|K\|}$  defines r.

**Remark 4.19.** Lemma 4.18 admits a more general version which applies to arbitrary manifolds M equipped with a suitable triangulation, though we will not require that level of generality. This is claimed in Madsen–Weiss [13, Appendix A.1].

An inclusion of simplicial complexes  $L \hookrightarrow K$  is called a *relative horn inclusion* if K is obtained from L by attaching a simplex along a horn in L. The following lemma will be crucial in the proof of Proposition 4.1.

**Lemma 4.20.** Let B be a subdivision of  $\Delta^n$  as in Proposition 4.17. Given a sequence of relative horn inclusions

$$A = A_0 \hookrightarrow A_1 \hookrightarrow \cdots \hookrightarrow A_\ell = B$$


there exists a homotopy  $\underline{H}: \overline{\|B\|} \times \mathbb{A}^1 \to \overline{\|B\|}$  such that H restricts to a homotopy  $\overline{\|A\|} \times \mathbb{A}^1 \to \overline{\|A\|}$ ,  $H_0 = \mathrm{id}$  and  $H_1$  factors through  $\overline{\|A\|}$ .

We introduce some terminology in preparation for the proof of this lemma. Let  $0 < k \le \ell$ . A homotopy

$$H^{(k)}: \overline{\|B\|} \times \mathbb{A}^1 \to \overline{\|B\|}$$

is said to have property  $(\sigma_k)$  if  $H^{(k)}$  restricts to a homotopy  $\overline{\|A_k\|} \times \mathbb{A}^1 \to \overline{\|A_k\|}$ ,  $H_0 = \mathrm{id}$  and  $H_1$  factors through  $\overline{\|A_k\|}$ .

Proof. Before tackling the lemma in full generality, we prove it for the easy case of  $A_0 \hookrightarrow A_1 = B$  for a single horn inclusion  $\Lambda \hookrightarrow \Delta^n$ . Choose a subdivision T of  $\Delta^n$  which is the simplicial cone on a subdivision of  $\Delta^{n-1}$ . For concreteness, we take T to be the simplicial cone of  $\mathrm{sd}\,\Delta^{n-1}$ , the first barycentric subdivision of  $\Delta^{n-1}$ . (That is, T is the nerve of the category obtained by adjoining a terminal object v to the poset of non-degenerate simplices of the standard (n-1)-simplex.) We call T the cone-subdivision of the n-simplex. Here is a picture for n=3:



By Proposition 4.17, we have a homotopy

$$h: \mathbf{\Delta}^n \times \mathbb{A}^1 \to \mathbf{\Delta}^n$$

which factors through ||T|| at time 1. Now, let T' be the simplicial complex obtained from T by discarding the vertex  $c \in T$  corresponding to the top simplex in  $\Delta^{n-1}$ . (To obtain a simplicial complex, we must also discard all the simplices in T that have c as a face.) Then T' is a subdivision of the n-dimensional horn. The inclusion  $i_T: T' \hookrightarrow T$  admits a retraction  $r_T: T \to T'$ , essentially given by collapsing c onto v. This is a simplicial map, it is the application of the appropriate degeneracy map on each simplex of T. Moreover, we can construct a homotopy on each simplex of T between the identity and the aforementioned map. This can be done by linear interpolation, for example. Thus we obtain a homotopy

$$h': \overline{\|T\|} \times \mathbb{A}^1 \to \overline{\|T\|}$$

which factors through ||T'|| (and hence through  $||\Lambda||$ ) at time 1. Clearly, the composition (concatenation) of h and h' gives a homotopy H satisfying the conditions of the lemma, i.e., having property  $(\sigma_0)$ .

With this special case in hand we proceed to the general one, arguing by induction. Suppose we have constructed a homotopy  $H^{(k)}$  having property  $(\sigma_k)$ . We now construct a homotopy  $H^{(k-1)}$  having property  $(\sigma_{k-1})$  as follows. Firstly, take a subdivision K of  $A_k$  (and hence a subdivision of  $A_{k-1}$ ) that restricts to the cone triangulation on the simplex attached to  $A_{k-1}$ . Lemma 4.18 gives us a homotopy on  $\Delta^n$  that restricts to a homotopy

$$f: \overline{\|A_k\|} \times \mathbb{A}^1 \to \overline{\|A_k\|}$$

with  $f_0 = id$  and which factors through  $\overline{||K||}$  at time 1.

Now, by collapsing the cone subdivision of the attached simplex to the (subdivided) horn using the simplicial map from the case of a single horn inclusion, we obtain a homotopy

$$g: \overline{\|K\|} \times \mathbb{A}^1 \to \overline{\|K\|} \subset \overline{\|A_k\|}$$

with  $g_0 = \text{id}$  and which factors through  $\overline{\|A_{k-1}\|}$  at time 1. Compose (concatenate) the homotopies f and g to obtain a homotopy h on  $\|A_k\|$ . Then define  $H^{(k-1)}$  to be composition of  $H^{(k)}$  and h. For this composition to be smooth, we emphasize that it is important to first apply Lemma 4.18 to the whole of  $\overline{\|A_k\|}$ , not just the on the simplex that we are collapsing. This completes the induction.

Corollary 4.21. The extended-simplices version of Lemma 4.20 holds. Namely, under the conditions of that lemma, there exists a homotopy  $G: ||B|| \times \mathbb{A}^1 \to ||B||$  such that G restricts to a homotopy  $||A|| \times \mathbb{A}^1 \to ||A||$ ,  $G_0 = \operatorname{id}$  and  $G_1$  factors through ||A||.

*Proof.* Starting from the homotopy  $H: \overline{\|B\|} \times \mathbb{A}^1 \to \overline{\|B\|}$  of Lemma 4.20, we obtain a homotopy

$$\widetilde{H}: \|B\| \times \mathbb{A}^1 \xrightarrow{\lambda} \overline{\|B\|} \times \mathbb{A}^1 \xrightarrow{H} \overline{\|B\|} \xrightarrow{i} \|B\|$$

where i is the inclusion and  $\lambda$  is the map constructed in Lemma 4.15. It is clear that  $\widetilde{H}$  restricts to a homotopy on ||A||,  $\widetilde{H}_0 = \lambda i$  and  $\widetilde{H}_1$  factors through ||A||. Now define G as the concatenation of the homotopy on ||B|| from Lemma 4.15 (between the identity and  $\lambda i$ ) with the homotopy  $\widetilde{H}$ .

## 4.d Proof of Propositions 4.1, 4.2, and 4.3

For concreteness, we assume F to be injectively fibrant (by replacing it if necessary) so that Proposition 4.10 applies.

Proof of Proposition 4.1. Let  $i: A \hookrightarrow B$  denote the map  $h_i$ :

$$h_i: \mathrm{Sd}^i(\Lambda_k[n]) \hookrightarrow \mathrm{Sd}^i(\Delta[n])$$

 $(i \ge 0, n \ge 1, \text{ and } 0 \le k \le n)$ , as in Definition 3.8. By adjunction of Definition 4.4, it suffices to find weak liftings

$$||A||_{\text{pre}} \xrightarrow{f} F$$

$$\downarrow \qquad \qquad \widetilde{\alpha} \qquad \qquad \uparrow$$

$$||B||_{\text{pre}}$$

together with a homotopy  $H: \|A\|_{\text{pre}} \times \|\Delta^1\|_{\text{pre}} \to F$  between  $\widetilde{\alpha}\iota$  and f. Indeed, this gives us the required homotopy  $A \times \Delta[1] \to F(\mathbb{A}^{\bullet})$  by precomposing H with  $\|A \times \Delta^1\|_{\text{pre}} \to \|A\|_{\text{pre}} \times \|\Delta^1\|_{\text{pre}}$  and applying the adjunction again.

We observe that it suffices to solve the above problem with  $\|-\|_{\text{pre}}$  replaced by  $\|-\|$  everywhere and the map  $f:\|A\|_{\text{pre}}\to F$  replaced by some arbitrary map  $\hat{f}:\|A\|\to F$  such that  $\hat{f}\circ\iota_A=f$ . Indeed, by precomposing the resulting maps with the natural maps  $\iota_{(-)}:\|-\|_{\text{pre}}\to\|-\|$  we obtain the desired data.

The strategy is to find a homotopy retraction of  $\iota$ , i.e., a map  $r:\|B\|\to \|A\|$  together with a homotopy  $\|A\|\times \|\Delta^1\|\to \|A\|$  between  $r\iota$  and the identity. In fact, we construct this homotopy on  $\|A\|$  as the restriction of a homotopy on  $\|B\|$ , and therefore prove that  $\iota$  is a homotopy equivalence. (We will need this stronger statement in the proof of Proposition 4.2.) This is achieved by a direct application of Corollary 4.21. More precisely, we choose a sequence of relative horn inclusions from  $A=\operatorname{sd}^i\Lambda^n_k$  to  $B=\operatorname{sd}^i\Lambda^n$ , for each  $i\geq 0$ , n>0 and  $0\leq k\leq n$ , and apply Corollary 4.21. That the resulting homotopy preserves the 0-simplices of each horn follows directly from its construction.

Proof of Proposition 4.2. We keep the notation  $i:A\hookrightarrow B$  for the map  $h_i$  as in Definition 3.8. As in the proof of Proposition 4.1, we use the adjunction of Definition 4.4 and the natural transformation  $\|-\|_{\text{pre}}\to\|-\|$  to reduce the problem to constructing certain maps of sheaves of sets.

Let P denote the pushout of sheaves of sets

$$\|A\| \times \mathbb{A}^1 \bigsqcup_{\|A\| \times \partial \mathbb{A}^1} \|B\| \times \partial \mathbb{A}^1.$$

To verify the weak RLP with respect to i it suffices to prove that for any map  $\alpha: P \to F$  there is a dashed map  $\widetilde{\alpha}$ :

making the diagram commute up to a homotopy  $H: P \times \mathbb{A}^1 \to F$  from  $\alpha$  to  $\widetilde{\alpha}\iota$ , which is fixed on  $||B|| \times \partial \mathbb{A}^1$  pointwise. Being fixed pointwise means that the restriction of H to  $(||B|| \times \partial \mathbb{A}^1) \times \mathbb{A}^1$  factors as the projection to  $||B|| \times \partial \mathbb{A}^1$  followed by  $\alpha$ . The result now follows from the lemma below.

**Lemma 4.22.** Suppose  $i: A \hookrightarrow B$  is the map  $h_i$  as in Definition 3.8. There is a homotopy  $H: (\|B\| \times \mathbb{A}^1) \times \mathbb{A}^1 \to \|B\| \times \mathbb{A}^1$  such that

- 1. H restricts to a homotopy  $P \times \mathbb{A}^1 \to P$  which fixes  $||B|| \times \partial \mathbb{A}^1$  pointwise, where P denotes the pushout of sheaves of sets  $||A|| \times \mathbb{A}^1 \sqcup_{||A|| \times \partial \mathbb{A}^1} ||B|| \times \partial \mathbb{A}^1$ .
- 2.  $H_0$  is the identity and  $H_1$  factors through P.

Proof. Choose a bump function  $c: \mathbb{A}^1 \to [0,1] \subset \mathbb{A}^1$  with  $c \equiv 0$  in an open neighborhood of  $(-\infty,0] \cup [1,\infty)$ , and  $c \equiv 1$  in a open neighborhood J of 1/2, c(t) increasing for  $t \leq 1/2$  and decreasing for  $t \geq 1/2$ . Then choose a function  $f: \mathbb{A}^1 \to \mathbb{A}^1$  with  $f(t) \equiv 0$  when  $c \neq 1$  and  $t \leq 1/2$  and  $f(t) \equiv 1$  when  $c \neq 1$  and t > 1/2. Let  $H_B: ||B|| \times \mathbb{A}^1 \to ||B||$  be the map constructed in the proof of Proposition 4.1. Define maps  $R_1, R_2: (||B|| \times \mathbb{A}^1) \times \mathbb{A}^1 \to ||B|| \times \mathbb{A}^1$  as

$$R_1(x,t,s) = (H_B(x,s \cdot c(t)),t)$$

and

$$R_2(x, t, s) = (x, s \cdot f(t) + (1 - s)t).$$

for  $x \in \overline{\|B\|}$ , and  $t, s \in \mathbb{A}^1$ . Then define H as


$$(x,t,s) \mapsto \begin{cases} R_1(x,t,2s) & \text{for } s \le 1/2\\ R_2(x,t,2s-1) \circ R_1(x,t,1) & \text{for } s > 1/2 \end{cases}$$

Now  $R_1$  and  $R_2$  separately satisfy condition (1) in the lemma, so H does as well. As for property (2), the map induced by H at s=1/2 has image in  $||A|| \times J$ , and the homotopy  $R_2$  preserves this subspace. Where  $c \neq 1$ , the map induced by H at s=1 has image in  $||B|| \times \partial \mathbb{A}^1$  (which follows from how we chose f). So altogether, this shows we get the claimed factorization.

Proof of Proposition 4.3. The proof is of the same sort as that of Proposition 4.2, using the adjunction of Definition 4.4 and the natural transformation  $\|-\|_{\text{pre}} \to \|-\|$  to reduce the problem to constructing certain maps of sheaves of sets. Let  $i: A \hookrightarrow B$  be the map  $h_i$  as in Definition 3.8. Consider the pushout

$$P:=S^n\times \|A\|\bigsqcup_{*\times \|A\|}*\times \|B\|.$$

The same manipulations as before show that to verify the weak RLP with respect to i it suffices to prove that for any map  $\alpha: P \to F$  there is a map  $\widetilde{\alpha}$ :



making the diagram commute up to a homotopy  $P \times \mathbb{A}^1 \to F$  from  $\alpha$  to  $\tilde{\alpha}\iota$  which is fixed on  $* \times ||B||$  pointwise. The result now follows from a lemma analogous to Lemma 4.22, below.

**Lemma 4.23.** Suppose  $i: A \hookrightarrow B$  is the map  $h_i$  as in Definition 3.8. There is a homotopy  $H: (\|B\| \times S^n) \times \mathbb{A}^1 \to \|B\| \times S^n$  such that

- 1. H restricts to a homotopy  $P \times \mathbb{A}^1 \to P$  which fixes  $||B|| \times *$  pointwise, where P is the pushout of sheaves of sets  $S^n \times ||A|| \sqcup_{* \times ||A||} * \times ||B||$ .
- 2.  $H_0$  is the identity and  $H_1$  factors through P.

*Proof.* Let  $* \in D_{\epsilon} \subset D_{\delta} \subset S^n$  be open disk neighborhoods of radius  $\epsilon$  and  $\delta$  with  $\epsilon < \delta$ . Choose a smooth function  $c: S^n \to [0,1] \subset \mathbb{A}^1$  such that (i)  $c|_{S^n \setminus D_{\epsilon}} \equiv 0$  and (ii) c(\*) = 1. For example, we can choose c to be a bump function with support in  $D_{\epsilon}$  that is 1 at the basepoint.

Also choose a homotopy  $h: \mathbb{A}^1 \times S^n \to S^n$  such that (i)  $h(1,-)|_{D_{\epsilon}}$  is constant to  $* \in S^n$  and (ii)  $h(t,-)|_{S^n \setminus D_{\delta}} = \mathrm{id}$ . In words, this homotopy collapses a neighborhood of the basepoint to itself.

Let  $H_B: ||B|| \times \mathbb{A}^1 \to ||B||$  be the map constructed in Proposition 4.1. We construct H as the composition of two homotopies. Define

$$R_1: S^n \times ||B|| \times \mathbb{A}^1 \to S^n \times ||B|| \qquad R_1(z, x, t) = (z, H_B(x, t(1 - c(z))))$$

and

$$R_2: S^n \times ||B|| \times \mathbb{A}^1 \to S^n \times ||B|| \qquad R_2(z, x, t) = (h(t, z), x).$$

The first homotopy collapses  $S^n \times \|B\|$  onto  $S^n \times \|A\|$  outside a neighborhood  $N \subset D_{\epsilon}$  of the basepoint. The second collapses  $D_{\epsilon} \times \|B\|$  to  $* \times \|B\|$ . The composition of these homotopies satisfies the claimed properties.

## 5 The shape functor preserves the $\infty$ -sheaf property

In this section we assemble the previous results to prove Theorem 1.2. Our approach uses the following characterization of  $\infty$ -sheaves.

**Theorem 5.1.** A presheaf  $F:\mathsf{Man}^\mathrm{op}\to\mathsf{S}$  is an  $\infty$ -sheaf (Definition 2.15) if and only if  $F(\varnothing)\simeq *$  and

(1) for all manifolds M and open covers of M with two elements  $\{U, V\}$ , the commutative square

$$F(M) \longrightarrow F(V)$$

$$\downarrow \qquad \qquad \downarrow$$

$$F(U) \longrightarrow F(U \cap V)$$

is a homotopy pullback square; and

(2) for all manifolds M and an open cover  $\{V_i\}_{i\geq 0}$  by a nested sequence of open sets with  $\overline{V_i} \subset V_{i+1}$ , the canonical map

$$F(M) = F\left(\bigcup_{i} V_{i}\right) \to \underset{i}{\operatorname{holim}} F(V_{i})$$

is a weak homotopy equivalence.

This is probably well-known and is similar to a special case of [22, Theorem 5.2] and [2, Theorem 7.2]. For completeness, we provide a proof below.

In preparation, note that by cofinality there is an identification of homotopy limits,

$$\operatorname{holim}_{[n] \in \Delta} \prod_{i_0, \dots, i_n \in I} F(U_{i_0} \cap \dots \cap U_{i_n}) \simeq \operatorname{holim}_{S \subset I} F(U_S)$$

where the homotopy limit on the right ranges over all finite, non-empty subsets  $S \subset I$  and  $U_S$  is notation for  $\bigcap_{i \in S} U_i$ .

Proof of Theorem 5.1. Suppose first that M is a compact manifold and take an open cover  $\{U_i\}_{i\in I}$  of M. Then  $\{U_i\}$  has a finite subcover  $\{U_i\}_{i\in J}$ ,  $J\subset I$ , and so

$$\operatorname{holim}_{S \subset I} F(U_S) \simeq \operatorname{holim}_{S \subset J} F(U_S).$$

The homotopy limit on the right-hand side is indexed over a finite category (a cube) and so it is equivalent to an iterated homotopy pullback. Condition (1) applied inductively shows that this iterated homotopy pullback is F(M).

If M is non-compact, take an exhaustion of M by interiors of compact manifolds  $V_0 \subset V_1 \subset \cdots$  with  $M = \bigcup V_i$ . Such an exhaustion can be obtained by picking a smooth proper map  $f: M \to \mathbb{R}$  and setting  $V_i$  to be the interior of  $f^{-1}((-\infty,i])$ . Then, for an open cover  $\{U_i \to M\}_{i \in I}$ , we have that

$$\underset{S \subset I}{\text{holim}} F(U_S) \simeq \underset{S \subset I}{\text{holim}} \underset{j>0}{\text{holim}} F(V_j \cap U_S) \tag{5.2}$$

by (2) applied to the covers  $\{U_S \cap V_j \to U_S\}_j$  for each S. Now commute the homotopy limits and use that the cover  $\{V_j \cap U_i \to V_j\}_i$  (for a fixed j) has a finite subcover to conclude using (1) that (5.2) is weakly equivalent to

$$\operatorname{holim}_{j\geq 0} F(V_j).$$

By invoking (2) again, this homotopy limit is weakly equivalent to F(M).

Remark 5.3. In a previous iteration of this paper, condition (2) did not include the closure condition on the covers. The referee kindly pointed out that our proof implied the new (weaker) statement and suggested a simple argument that replaces condition (2) by the following weaker condition (2)': if M is a (possibly infinite) disjoint union of submanifolds  $U_i$  then  $F(M) \to \prod_i^h F(U_i)$  is a weak equivalence. The argument goes as follows. Suppose  $\{V_i\}$  is an open cover as in (2). Let  $W_0$  be the disjoint union of  $V_{k+2} \setminus \overline{V_k}$ , taken over k even, and let  $W_1$  be the disjoint union of  $V_{k+2} \setminus \overline{V_k}$ , taken over k odd. Then  $W_0$  and  $W_1$  form an open cover of M so, by (1), we have

$$F(M) \to \operatorname{holim}(F(W_0) \to F(W_0 \cap W_1) \leftarrow F(W_1)).$$

By (2)', the target is equivalent to

$$\operatorname{holim}(\operatorname{holim}_i F(W_0 \cap V_i) \to \operatorname{holim}_i F(W_0 \cap W_1 \cap V_i) \leftarrow \operatorname{holim}_i F(W_1 \cap V_i)),$$

which, by commuting homotopy limits and using (1), is equivalent to holim  $F(V_i)$ .

**Remark 5.4.** The same proof works for topological manifolds. The main observation for the non-compact case is that there exists a proper map  $M \to \mathbb{R}$  for M a topological manifold (the requirement is that partitions of unity exist). For generalizations of these statements, see Pavlov [16].

We shall tackle properties (1) and (2) for  $\mathbf{B}F$  separately below. We call them the *finite* and *non-compact* cases, respectively.

#### 5.a The finite case

**Theorem 5.5.** Let  $F: \mathsf{Man}^{\mathrm{op}} \to \mathsf{S}$  be an  $\infty$ -sheaf and M a smooth manifold with U, V two open subsets of M such that  $U \cup V = M$ . Then the commutative square

$$\begin{array}{ccc}
\mathbf{B}F(M) & \longrightarrow & \mathbf{B}F(V) \\
\downarrow & & \downarrow \\
\mathbf{B}F(U) & \longrightarrow & \mathbf{B}F(U \cap V)
\end{array}$$

is homotopy cartesian.

*Proof.* The (homotopy) pullback

$$(\mathbf{B}F(U) \times \mathbf{B}F(V)) \times_{\mathrm{map}(\partial \Delta^1, \mathbf{B}F(U \cap V))} \mathrm{map}(\Delta^1, \mathbf{B}F(U \cap V))$$

is identified with

$$(\mathbf{B}F(U) \times \mathbf{B}F(V)) \times_{\mathbf{B}F(U \cap V \times \partial \mathbb{A}^1)}^{h} \mathbf{B}F(U \cap V \times \mathbb{A}^1)$$
(5.6)

since  $\mathbf{B}F$  is concordance invariant. By Proposition 4.2, we may commute the homotopy pullback with geometric realization, and thus (5.6) is identified with the geometric realization of the simplicial space

$$(F(U \times \mathbb{A}^{\bullet}) \times F(V \times \mathbb{A}^{\bullet})) \times_{F(U \cap V \times \partial \mathbb{A}^{1} \times \mathbb{A}^{\bullet})}^{h} F(U \cap V \times \mathbb{A}^{1} \times \mathbb{A}^{\bullet}). \tag{5.7}$$

To prove that the map from  $F(M \times \mathbb{A}^{\bullet})$  to (5.7) is a weak equivalence after realization we first refine the cover in a convenient way using a partition of unity subordinate to  $\{U,V\}$ . So let  $f_U:M\to [0,1]$  and  $f_V:M\to [0,1]$  with  $f_U+f_V\equiv 1$  and  $\operatorname{supp}(f_U)\subset U$ ,  $\operatorname{supp}(f_V)\subset V$ . Take  $U'=f_U^{-1}(2/3,1]$  and  $V'=f_V^{-1}(2/3,1]$ . Notice that  $U'\cap V'=\emptyset$ , and  $\{U',V',U\cap V\}$  covers M. Let  $c:\mathbb{A}^1\to\mathbb{A}^1$  be a cutoff function with  $c|_{(-\infty,1/3)}\equiv 0$  and  $c|_{(2/3,\infty)}\equiv 1$ , and define  $f:=c\circ f_V|_{U\cap V}$ , so that  $f:U\cap V\to\mathbb{A}^1$ .

Rearrange (5.7) as an iterated homotopy pullback and consider the maps:

$$F(U \times \mathbb{A}^{\bullet}) \times_{F(U \cap V \times \mathbb{A}^{\bullet})}^{h} F(U \cap V \times \mathbb{A}^{1} \times \mathbb{A}^{\bullet}) \times_{F(U \cap V \times \mathbb{A}^{\bullet})}^{h} F(V \times \mathbb{A}^{\bullet})$$

$$\downarrow \text{res}$$

$$F(U' \times \mathbb{A}^{\bullet}) \times_{F(U' \cap V \times \mathbb{A}^{\bullet})}^{h} F(U \cap V \times \mathbb{A}^{1} \times \mathbb{A}^{\bullet}) \times_{F(U \cap V' \times \mathbb{A}^{\bullet})}^{h} F(V' \times \mathbb{A}^{\bullet})$$

$$\text{pr}^{*} \Big[ \Big] f^{*}$$

$$F(U' \times \mathbb{A}^{\bullet}) \times_{F(U' \cap V \times \mathbb{A}^{\bullet})}^{h} F(U \cap V \times \mathbb{A}^{\bullet}) \times_{F^{\mathbb{A}^{\bullet}} F(U \cap V' \times \mathbb{A}^{\bullet})}^{h} F(V' \times \mathbb{A}^{\bullet}).$$

The restriction map from  $F(M \times \mathbb{A}^{\bullet})$  to the last space is a weak equivalence since  $\{U', U \cap V, V'\}$  cover M and F is an  $\infty$ -sheaf. Similarly, the map res is a levelwise weak equivalence since  $\{U', U \cap V\}$  covers U and  $\{V', U \cap V\}$  covers V. The arrow pr\* is induced by the projection  $\operatorname{pr}: (U \cap V) \times \mathbb{A}^1 \to U \cap V$ . We obtain a map  $U \cap V \to (U \cap V) \times \mathbb{A}^1$  from  $f: U \cap V \to \mathbb{A}^1$  defined in the previous paragraph. By construction,  $f|_{U'\cap V}=0$  and  $f|_{U\cap V'}=1$ , which is precisely the compatibility condition required to extend to a map on sections in the fibered product which we denote by  $f^*$ . Notice that since  $\operatorname{pr} \circ (\operatorname{id}_{U\cap V}, f) = \operatorname{id}_{U\cap V}$ , we have  $(\operatorname{id}_{U\cap V}, f)^* \circ \operatorname{pr}^* = \operatorname{id}$ . It remains to show that  $\operatorname{pr}^* \circ (\operatorname{id}_{U\cap V}, f)^*$  is homotopic to the identity.

We consider the interpolation  $h:((U\cap V)\times\mathbb{A}^1)\times\mathbb{A}^1\to\mathbb{A}^1$  between  $f\circ \operatorname{pr}$  and the projection map  $q:(U\cap V)\times\mathbb{A}^1\to\mathbb{A}^1$ , given by

$$h(t) = (1 - t) \cdot q + t \cdot (f \circ pr)$$

and extend it to a smooth homotopy

$$H = (\mathrm{id}_{U \cap V}, h) : ((U \cap V) \times \mathbb{A}^1) \times \mathbb{A}^1 \to ((U \cap V) \times \mathbb{A}^1).$$

Since  $F(-\times \mathbb{A}^{\bullet})$  sends smooth homotopies to simplicial homotopies (Proposition 2.9), and the map H fixes  $(U \cap V) \times \partial \mathbb{A}^1$  pointwise, the map H induces the required simplicial homotopy from  $\operatorname{pr}^* \circ (\operatorname{id}_{U \cap V}, f)^*$  to id.

**Remark 5.8.** The beginning of the above proof has the following obvious generalization. For a diagram  $F \to G \leftarrow H$  of  $\infty$ -sheaves, define the *geometric* homotopy pullback to be the  $\infty$ -sheaf whose value at a manifold M is the homotopy pullback of the diagram

$$G(M imes \mathbb{A}^1)$$
 $\downarrow$  endpoints
 $F(M) imes H(M) \longrightarrow G(M) imes G(M)$ 

Then the classifying space functor  ${\bf B}$  sends geometric homotopy pullbacks of  $\infty$ -sheaves to homotopy pullbacks of spaces.

#### 5.b The non-compact case

**Theorem 5.9.** Let  $\{U_i\}_{i\geq 0}$  be a collection of manifolds (possibly non-compact). For an  $\infty$ -sheaf F: Man<sup>op</sup>  $\to S$ , the natural map

$$\mathbf{B}F\left(\bigsqcup_{i}U_{i}\right)\to\prod_{i}^{h}\mathbf{B}F(U_{i})$$

is a weak equivalence.

We deduce this from the lemma below, by setting  $F_i = F(U_i \times -)$ .

**Lemma 5.10.** Let  $\{F_i\}_{i\in I}$  be a collection of  $\infty$ -sheaves indexed over a possibly infinite set I. Then the map  $|\prod_i^h F_i(\mathbb{A}^{\bullet})| \to \prod_i^h |F_i(\mathbb{A}^{\bullet})|$  is a weak equivalence of spaces.

We use the symbol  $\prod$  for the homotopy product, i.e., the derived functor of the product. Note this has a different meaning in simplicial spaces (with degreewise weak equivalences) and simplicial sets (with the usual weak equivalences). In the simplicial space case, it means: replace each factor by a degreewise fibrant simplicial space and then compute the product; in the simplicial set case, it means: replace each factor by a Kan complex and then compute the product. The homotopy product (of spaces or of simplicial spaces) agrees with the non-derived product when the indexing set is finite. In general, they do not agree when the set is infinite but Lemma 5.10 says they agree for the concordance resolution.

To prove Lemma 5.10, we develop a geometric analogue of the  $\mathsf{Ex}^\infty$  functor. Recall (Definition A.12) the adjunction  $\mathsf{Sd} \dashv \mathsf{Ex}$ , where  $\mathsf{Sd}$  and  $\mathsf{Ex}$  are functors of the form  $\mathsf{sS} \to \mathsf{sS}$ . Recall also (Definition 4.4) the adjunction  $\|-\|_{\mathrm{pre}} \dashv -(\mathbb{A}^\bullet)$  between functors  $\|-\|_{\mathrm{pre}} : \mathsf{sS} \to \mathsf{PSh}(\mathsf{Man},\mathsf{S})$  and  $-(\mathbb{A}^\bullet) : \mathsf{PSh}(\mathsf{Man},\mathsf{S}) \to \mathsf{sS}$ .

#### **Definition 5.11.** Denote by

$$\mathrm{Ex}_{\mathsf{Man}}:\mathsf{PSh}(\mathsf{Man},\mathsf{S}) o \mathsf{sS}$$

the functor which assigns to a presheaf F the simplicial space

$$[n] \mapsto \mathbf{R} \operatorname{map}(||\operatorname{Sd} \Delta^n||_{\operatorname{pre}}, F).$$

We define a natural transformation

$$h^*: -(\mathbb{A}^{\bullet}) \to \operatorname{Ex}_{\mathsf{Man}}$$

of functors  $PSh(Man, S) \rightarrow sS$  which on a presheaf F is

$$F(\mathbb{A}^{\bullet}) \xrightarrow{\sim} \mathbf{R} \operatorname{map}(||\Delta[\bullet]||_{\operatorname{pre}}, F) \to \mathbf{R} \operatorname{map}(||\operatorname{Sd}\Delta[\bullet]||_{\operatorname{pre}}, F),$$

where the second map is given by precomposition with  $h: \|\operatorname{Sd}\Delta[n]\|_{\operatorname{pre}} \hookrightarrow \|\Delta[n]\|_{\operatorname{pre}}$ , which is the affine extension of the map that sends each vertex of  $\operatorname{Sd}\Delta[n]$ , i.e., a non-degenerate simplex of  $\Delta[n]$ , to the corresponding barycenter in  $\|\Delta[n]\|$  (as in Section 4.c).

**Definition 5.12.** Given  $F \in \mathsf{PSh}(\mathsf{Man},\mathsf{S})$ , denote by  $\mathsf{Ex}^\infty_{\mathsf{Man}}F(\mathbb{A}^\bullet)$  the colimit (which is also a homotopy colimit) of

$$F(\mathbb{A}^{\bullet}) \xrightarrow{h^*} \operatorname{Ex}(F(\mathbb{A}^{\bullet})) \xrightarrow{h^*} \operatorname{Ex}^2(F(\mathbb{A}^{\bullet})) \to \cdots$$

The object  $\mathsf{Ex}^\infty_{\mathsf{Man}} F(\mathbb{A}^\bullet)$  is not the usual Kan  $\mathsf{Ex}^\infty$  simplicial space, since the map h uses geometric barycenters, whereas its usual analogue uses the last vertex operator.

**Proposition 5.13.** For an  $\infty$ -sheaf  $F \in \mathsf{PSh}(\mathsf{Man},\mathsf{S})$ , the simplicial space  $\mathsf{Ex}^\infty_{\mathsf{Man}} F(\mathbb{A}^\bullet)$  is a Kan complex. Moreover, the map  $F(\mathbb{A}^\bullet) \to \mathsf{Ex}^\infty_{\mathsf{Man}} F(\mathbb{A}^\bullet)$  satisfies the conditions of Corollary A.11.

*Proof.* We assume, without loss of generality, that F is injectively fibrant (so that the derived mapping spaces may be replaced by honest mapping spaces). The assertion that  $\mathsf{Ex}^\infty_{\mathsf{Man}} F(\mathbb{A}^\bullet)$  is a Kan complex is a consequence of the following observation: for an n-dimensional horn  $\Lambda$ , the map  $h: \|\operatorname{Sd} \Lambda\|_{\operatorname{pre}} \to \|\Lambda\|_{\operatorname{pre}}$  factors as

$$\|\operatorname{Sd}\Lambda\|_{\operatorname{pre}} \hookrightarrow \|\operatorname{Sd}\Delta[n]\|_{\operatorname{pre}} \to \|\Lambda\|_{\operatorname{pre}}$$

where the last map is the collapse map. Such a factorization gives a factorization of the map

$$\|\operatorname{Sd}^{i+1}\Lambda\|_{\operatorname{pre}} \to \|\operatorname{Sd}^i\Lambda\|_{\operatorname{pre}} \xrightarrow{\alpha} F$$

through  $\|\operatorname{Sd}^{i+1}\Delta[n]\|_{\operatorname{pre}}$ , for any choice of  $\alpha$ . In adjoint form, this guarantees that the diagram

$$\Lambda \xrightarrow{\alpha} \operatorname{Ex}^{i}(F(\mathbb{A}^{\bullet})) 
\downarrow h^{*} 
\Delta[n] --- > \operatorname{Ex}^{i+1}(F(\mathbb{A}^{\bullet}))$$

has a lift as pictured, and hence that  $\mathsf{Ex}^\infty_\mathsf{Man} F(\mathbb{A}^\bullet)$  is a Kan complex.

We now prove the second claim. We will show that, for  $j \geq 0$ , every diagram

$$\operatorname{Sd}^{j} \partial \Delta[n] \longrightarrow F(\mathbb{A}^{\bullet})$$

$$\downarrow \qquad \qquad \downarrow$$

$$\operatorname{Sd}^{j} \Delta[n] \longrightarrow \operatorname{Ex}_{\mathsf{Man}}^{\infty} F(\mathbb{A}^{\bullet})$$

has a lift  $\operatorname{Sd}^j \Delta[n] \to F(\mathbb{A}^{\bullet})$  together with homotopies of the upper and lower triangles such that the two resulting maps  $\operatorname{Sd}^j \partial \Delta[n] \times \Delta[1] \to \operatorname{\mathsf{Ex}}^\infty_{\mathsf{Man}} F(\mathbb{A}^{\bullet})$  agree. In adjoint form, the data of this commutative square is a map

$$\|\operatorname{Sd}^{j+i}\Delta[n]\|_{\operatorname{pre}}\bigsqcup_{\|\operatorname{Sd}^{j+i}\partial\Delta[n]\|_{\operatorname{pre}}}\|\operatorname{Sd}^{j}\partial\Delta[n]\|_{\operatorname{pre}}\to F$$

for some  $i \geq 0$ . The weak lift exists since there are homotopies

$$H: \|\operatorname{Sd}^{j+i}\Delta[n]\|_{\operatorname{pre}} \times \mathbb{A}^1 \to F$$
 and  $G: \|\operatorname{Sd}^j\partial\Delta[n]\|_{\operatorname{pre}} \times \mathbb{A}^1 \to F$ 

which agree on  $\|\operatorname{Sd}^{j+i}\partial\Delta[n]\|_{\operatorname{pre}} \times \mathbb{A}^1$  and both factor at time 1 through a map  $\|\operatorname{Sd}^j\Delta[n]\|_{\operatorname{pre}} \to F$ . These homotopies arise from a homotopy retraction of the map induced by h,

$$P \to \|\operatorname{Sd}^j \Delta[n]\|_{\operatorname{pre}}$$

where P denotes the pushout in the display. The existence of such a homotopy retraction follows by an application of Proposition 4.17. In more detail, setting  $K = \operatorname{Sd}^{i+j} \Delta[n]$  in that proposition, we obtain a self homotopy of  $\mathbb{A}^n$  which maps each simplex of  $\operatorname{Sd}^{i+j} \Delta[n]$  (embedded affinely in  $\mathbb{A}^n$  via h) to itself. But that homotopy necessarily maps each simplex of  $\operatorname{Sd}^j \Delta[n]$  (embedded affinely in  $\mathbb{A}^n$  via h) to itself. As such, the restriction of the homotopy on  $\mathbb{A}^n$  gives the horizontal maps in a commutative diagram

$$\|\operatorname{Sd}^{i+j}\Delta[n]\|_{\operatorname{pre}} \times \mathbb{A}^{1} \longrightarrow \|\operatorname{Sd}^{i+j}\Delta[n]\|_{\operatorname{pre}}$$

$$\downarrow h \qquad \qquad \downarrow h$$

$$\|\operatorname{Sd}^{j}\Delta[n]\|_{\operatorname{pre}} \times \mathbb{A}^{1} \longrightarrow \|\operatorname{Sd}^{j}\Delta[n]\|_{\operatorname{pre}}$$

and, evaluating at time 1, a map  $r: \|\operatorname{Sd}^j \Delta[n]\|_{\operatorname{pre}} \to \|\operatorname{Sd}^{i+j} \Delta[n]\|_{\operatorname{pre}}$  from the lower left corner to the upper right corner making the resulting diagram commute. This determines a self-homotopy of P which at time 0 is the identity and at time 1 factors through  $\|\operatorname{Sd}^j \Delta[n]\|_{\operatorname{pre}} \to P$  via r, as claimed.

*Proof.* Consider the composite

$$\Bigl|\prod_i^h F_i(\mathbb{A}^\bullet)\Bigr| \to \Bigl|\prod_i^h \operatorname{Ex}_{\operatorname{Man}}^\infty F_i(\mathbb{A}^\bullet)\Bigr| \to \prod_i^h \bigl|\operatorname{Ex}_{\operatorname{Man}}^\infty F_i(\mathbb{A}^\bullet)\bigr|.$$

If a collection of maps satisfies the conditions of Corollary A.11 then so does its product. It follows, using Proposition 5.13, that the first map is a weak equivalence. The second map is a weak equivalence since  $\mathsf{Ex}^\infty_{\mathsf{Man}}\,F_i(\mathbb{A}^\bullet)$  is Kan and the realization of a Kan simplicial space is a Kan simplicial set.  $\square$ 

# 6 What does the classifying space of an $\infty$ -category classify?

In this section, we suggest an answer to the question in the title. This expands on earlier questions and earlier answers in Moerdijk [14] and Weiss [23].

Let  $\mathbf{C}$  be a Segal space. For convenience, we assume that  $\mathbf{C}$  is Reedy fibrant as a simplicial space, otherwise the mapping spaces below need to be derived. (For definitions and more explanations, see Rezk [17].) For example,  $\mathbf{C}$  could be the (Reedy fibrant replacement of the) nerve of a (topological) category. Informally, the following data should produce something deserving the name of a  $\mathbf{C}$ -bundle on a manifold M:

- an open cover  $\mathcal{U} = \{U_i\}$  of M and a total order on its indexing set I
- maps  $\{\phi_i: U_i \to \mathbf{C}_0 = \mathbf{ob}(\mathbf{C})\}$
- maps  $\{\phi_{i < j} : U_i \cap U_j \to \mathbf{C}_1 = \mathbf{mor}(\mathbf{C})\}$
- ...

These data are then required to satisfy compatibility conditions; e.g., for a point  $x \in U_i \cap U_j$ ,  $\phi_{i < j}(x)$  is a morphism in  $\mathbb{C}$  from  $\phi_i(x)$  to  $\phi_j(x)$ , etc. As everywhere else in this paper, space means simplicial set, so in the above a map from  $U_i$  is taken to mean a map of simplicial sets from the singular simplices of  $U_i$  to a given simplicial set.

We make the above informal description a C-bundle precise as follows.

**Definition 6.1.** A C-bundle is an open cover  $\mathcal{U} = \{U_i \to M\}_{i \in I}$  (we stress that here we do not require that I is totally ordered) with a simplicial space map  $N\mathcal{U} \to \mathbf{C}$ , where  $N\mathcal{U}$  denotes the nerve of the following topological category. The space of objects is

$$\bigsqcup_{\varnothing \neq S \subset I} U_S$$

where the coproduct runs over non-empty finite subsets S of I and  $U_S := \cap_{s \in S} U_s$ . Given objects (R, x) and (S, y), with  $x \in U_R$  and  $y \in U_S$ , there is a morphism  $(R, x) \to (S, y)$  if and only if  $R \subset S$  (so that  $U_S \subset U_R$ ) and x = y. Therefore, the space of morphisms is

$$\bigsqcup_{\varnothing \neq R \subset S} U_S.$$

We view  $N\mathcal{U}$  as a simplicial space. Since  $N\mathcal{U}$  is Reedy cofibrant, the mapping space map $(N\mathcal{U}, \mathbf{C})$  agrees with the derived mapping space  $\mathbf{R}$ map $(N\mathcal{U}, \mathbf{C})$ .

**Remark 6.2.** We note that the informal description can be viewed as a special case of the definition by setting the image of certain morphisms—prescribed according to the ordering of I—to be identities. Conversely, given a **C**-bundle it is sometimes possible to construct a **C**-bundle as in the informal description above by adding to the collection  $\mathcal{U}$  all finite intersections of open sets in the original cover and choosing a total ordering on the resulting collection.

We now build a space of **C**-bundles. First, a definition:

**Definition 6.3.** For a manifold M, we define a simplicially enriched category Cov(M) of open covers  $\mathcal{U}$  of M and their refinements. Recall that a refinement  $\mathcal{U} \to \mathcal{V}$  is a choice of function  $\alpha : I \to J$  between the indexing sets of the covers such that  $U_i \subset V_{\alpha(i)}$  for each  $i \in I$ . We define a k-simplex in the space of morphisms of Cov(M),

$$map(\mathcal{U}, \mathcal{V})$$

to be a (k+1)-tuple of refinements  $\alpha_0, \ldots, \alpha_k : \mathcal{U} \to \mathcal{V}$ . The face and degeneracy maps are clear.

The space  $\operatorname{map}(\mathcal{U}, \mathcal{V})$  may of course be empty. If it is non-empty, it is the nerve of a groupoid, and for every pair of objects  $\alpha_0, \alpha_1$  there is by construction a unique morphism  $\alpha_0 \to \alpha_1$ . It follows that every k-sphere in  $\operatorname{map}(\mathcal{U}, \mathcal{V})$  has an unique filler, for every  $k \geq 0$ . Therefore,  $\operatorname{map}(\mathcal{U}, \mathcal{V})$  is either empty or contractible. As such,  $\operatorname{Cov}(M)$  is equivalent (as a simplicially enriched category) to the preorder of open covers  $\mathcal{U}$  of M with order relation  $\mathcal{U} \leq \mathcal{V}$  if  $\mathcal{U}$  refines  $\mathcal{V}$ .

The assignment  $\mathcal{U} \mapsto N\mathcal{U}$  defines a simplicially enriched functor from  $\mathsf{Cov}(M)$  to the category of simplicial spaces, since  $\mathsf{map}(\mathcal{U}, \mathcal{V})$  is a subspace of the space  $\mathsf{map}(N\mathcal{U}, N\mathcal{V})$  of simplicial space maps. Indeed, each refinement  $\mathcal{U} \to \mathcal{V}$  defines a map of simplicial spaces  $N\mathcal{U} \to N\mathcal{V}$ . For each pair of refinements  $\alpha_0, \alpha_1 : \mathcal{U} \to \mathcal{V}$ , the relations  $U_i \subset V_{\alpha_0(i)}$  and  $U_i \subset V_{\alpha_1(i)}$  imply that  $U_i \subset V_{\alpha_0(i)} \cap V_{\alpha_1(i)}$  and, as such, define a simplicial map  $N\mathcal{U} \times \Delta[1] \to N\mathcal{V}$ . More generally, a choice of refinements  $\alpha_0, \ldots, \alpha_k : \mathcal{U} \to \mathcal{V}$  implies the relation  $U_i \subset V_{\alpha_0(i)} \cap \cdots \cap V_{\alpha_k(i)}$  and so defines a map  $N\mathcal{U} \times \Delta[k] \to N\mathcal{V}$ .

**Definition 6.4.** The  $\infty$ -sheaf of C-bundles is the functor which to a manifold M associates the space

$$\mathbf{C}(M) := \underset{\mathcal{U} \to M}{\operatorname{hocolim}} \operatorname{map}(N\mathcal{U}, \mathbf{C})$$

given by the homotopy colimit of the enriched functor  $\mathcal{U} \mapsto \text{map}(N\mathcal{U}, \mathbf{C})$  on Cov(M). This functor is indeed enriched since on morphisms it is the restriction of the canonical map of spaces

$$\operatorname{map}(N\mathcal{U}, N\mathcal{V}) \to \operatorname{map}_{S}(\operatorname{map}(N\mathcal{V}, \mathbf{C}), \operatorname{map}(N\mathcal{U}, \mathbf{C}))$$

to  $\operatorname{map}_{\mathsf{Cov}(M)}(\mathcal{U}, \mathcal{V})$ .

The formula in this definition applies even if M has corners. So we may view  $\mathbf{C}$  as a functor on the larger category of manifolds with corners and smooth maps. In this setting, the subsheaf of sets  $\mathbf{\Delta}^n \subset \mathbb{A}^n$  of Section 4.b is representable.

**Proposition 6.5.** For each n, the space  $C(\Delta^n)$  is identified with the space of n-simplices of  $Ex^{\infty}$  C.

*Proof.* Let Cov denote the category  $Cov(\Delta^n)$  of open covers of  $\Delta^n$  and refinements. Let  $Cov_{sd}$  be the full subcategory of Cov spanned by open covers by open stars of the vertices of some barycentric subdivision of  $\Delta^n \subset \mathbb{A}^n$ .

The set of objects of  $\operatorname{Cov}_{\operatorname{sd}}$  is therefore identified with the non-negative integers: for each  $i \geq 0$ , the corresponding open cover  $\mathcal{U}(i)$  of  $\Delta^n$  is indexed by the set of vertices of the i-th barycentric subdivision of  $\Delta^n$ . The simplicial space  $N\mathcal{U}(i)$  is degreewise weakly equivalent to the simplicial discrete space  $\operatorname{Sd}^{i+1}\Delta[n]$ . To see this, note that for a subset  $S \subset \operatorname{Sd}^i\Delta[n]_0$  the space  $U_S$  is the open star of the unique non-degenerate simplex in  $\operatorname{Sd}^i\Delta[n]$  with vertex set S, if that simplex exists, and otherwise is empty; and the 0-simplices of  $\operatorname{Sd}^{i+1}\Delta[n]$  are by definition the non-degenerate simplices of  $\operatorname{Sd}^i\Delta[n]$ .

For each  $i \geq 0$ , there is a contractible choice of morphisms  $\mathcal{U}(i+1) \to \mathcal{U}(i)$  in  $\mathsf{Cov}_{\mathsf{sd}}$ . Among these, we are interested in a specific morphism, namely the one whose underlying function between indexing sets  $\mathsf{Sd}^{i+1} \, \Delta[n]_0 = \mathsf{Sd}(\mathsf{Sd}^i \, \Delta[n])_0 \to \mathsf{Sd}^i \, \Delta[n]_0$  is the last vertex map. The corresponding functor  $\mathbb{N} \to \mathsf{Cov}_{\mathsf{sd}}$  that selects these morphisms is an equivalence of simplicial categories.

Write  $j: \mathsf{Cov}_{\mathsf{sd}} \hookrightarrow \mathsf{Cov}$  for the inclusion. Clearly, every open cover of  $\Delta^n$  can be refined by one in  $\mathsf{Cov}_{\mathsf{sd}}$ . That is to say, for every open cover  $\mathcal V$  of  $\Delta^n$ , the comma category  $j/\mathcal V$  is non-empty. The category  $j/\mathcal V$  is equivalent to the discrete category (preorder) of open covers  $\mathcal U(i)$  in  $\mathsf{Cov}_{\mathsf{sd}}$  such that  $\mathcal U(i) \leq \mathcal V$  with refinement relation  $\leq$ . Clearly,  $\mathcal U(i) \leq \mathcal U(i')$  if and only if  $i \geq i'$ . From this description it is clear that  $j/\mathcal V$  is contractible. This shows that j is homotopy final, i.e., that the homotopy colimit defining  $\mathbf C(\Delta^n)$  may be indexed by the smaller  $\mathsf{Cov}_{\mathsf{sd}}$ . Therefore,

$$\mathbf{C}(\boldsymbol{\Delta}^n) \simeq \operatornamewithlimits{hocolim}_{\mathcal{U} \in \mathsf{Cov}_{\mathsf{sd}}} \operatorname{map}(N\mathcal{U}, \mathbf{C}) \simeq \operatornamewithlimits{hocolim}_{i>0} \operatorname{map}(\operatorname{Sd}^i \boldsymbol{\Delta}[n], \mathbf{C})$$

and so the result follows.

**Theorem 6.6.** For every smooth manifold M, the natural map

$$\mathbf{BC}(M) \to \mathbf{R}\mathrm{map}(M, B\mathbf{C})$$

is a weak equivalence. Here BC denotes the classifying space of C, i.e., the geometric realization of C viewed as a simplicial space, and BC is the functor B applied to the  $\infty$ -sheaf in Definition 6.4.

*Proof.* This is immediate from Theorem 1.1, together with the identification of  $|\mathbf{C}(\Delta^{\bullet})|$  with  $|\mathbf{Ex}^{\infty}\mathbf{C}|$  from Proposition 6.5, and  $|\mathbf{Ex}^{\infty}\mathbf{C}|$  with  $B\mathbf{C} = |\mathbf{C}|$  from Proposition A.13.

**Remark 6.7.** Clearly, if  $\mathbf{C} \to \mathbf{D}$  is a map inducing a weak equivalence between classifying spaces  $B\mathbf{C} \to B\mathbf{D}$ , then  $\mathbf{BC}(M) \to \mathbf{BD}(M)$  is a weak equivalence for every manifold M. This is the case, for example, if  $\mathbf{C} \to \mathbf{D}$  is a Dwyer–Kan equivalence of Segal spaces.

## A Simplicial sets and spaces

## A.a Weak equivalences of simplicial sets

**Proposition A.1** (Dugger–Isaksen [7, Proposition 4.1]). A map  $f: X \to Y$  between Kan complexes is a weak equivalence if and only if for every  $n \ge 0$  and every commutative square

$$\begin{array}{ccc} \partial \Delta^n & \longrightarrow X \\ \downarrow & \downarrow & \uparrow \\ \Delta^n & \longrightarrow Y \end{array}$$

there is a lift as pictured making the upper triangle commute and the lower triangle commute up to a homotopy  $H: \Delta^n \times \Delta^1 \to Y$  which is fixed on  $\partial \Delta^n$ .

In Dugger–Isaksen's terminology, a map f solving the lifting problem of this proposition is said to have the relative homotopy lifting property (RHLP) with respect to  $\partial \Delta^n \to \Delta^n$ .

It will be useful to think of these lifting properties in the following way. Let  $S^{[1]}$  denote the category whose objects are maps of simplicial sets and morphisms are commutative squares. Let  $\tau$  denote the following morphism in  $S^{[1]}$ 

(source i, target j). Proposition A.1 then reads: a map f between Kan complexes is a weak equivalence if and only if  $\tau^* : \text{map}(j, f) \to \text{map}(i, f)$  is a surjection.

Corollary A.2. Let  $\tau': i' \to j'$  be a commutative square weakly equivalent to  $\tau$ . (That is,  $\tau'$  is related to  $\tau$  by a zigzag of weak equivalences of squares.) A map  $f: X \to Y$  of simplicial sets is a weak equivalence if and only if

$$\mathbf{R}$$
map $(j', f) \to \mathbf{R}$ map $(i', f)$ 

is a surjection on  $\pi_0$ , where **R**map refers to the homotopy function complex in  $S^{[1]}$  relative to objectwise weak equivalences.

*Proof.* Since derived mapping spaces are invariant by weak equivalences by definition or construction, it suffices to prove that f is a weak equivalence if and only if

$$\operatorname{Rmap}(j, f) \to \operatorname{Rmap}(i, f)$$
 (A.3)

is a surjection on  $\pi_0$ . To interpret the derived mapping spaces, let us equip  $S^{[1]}$  with the *projective* model structure. In this model structure, an object (i.e., map) is fibrant if source and target are Kan simplicial sets. Without loss of generality, we may assume that f is fibrant. Cofibrant objects are simplicial maps that are cofibrations (between cofibrant objects, which is no condition here). Cofibrations are commutative squares

$$\begin{array}{ccc}
A & \longrightarrow & A' \\
\downarrow i & & \downarrow j \\
B & \longrightarrow & B'
\end{array}$$

(source i; target j) where the top map and the map

$$A' \bigsqcup_A B \to B'$$

is a cofibration of simplicial sets. It is not difficult to see that the morphism  $\tau$  is then a cofibration between cofibrant objects. It follows that

$$\tau^* : \operatorname{map}(j, f) \to \operatorname{map}(i, f)$$

is identified with (A.3) and is a Kan fibration. Since a Kan fibration is surjective if and only if it is surjective on  $\pi_0$ , the result follows.

Below are three examples which give rise to equivalent lifting problems:

**Example A.4.** Let  $\tau'$  be the following morphism in  $S^{[1]}$ :

$$\begin{array}{ccc} \partial \Delta^n & \longrightarrow \partial \Delta^n \times \Delta^1 \sqcup_{\partial \Delta^n \times \{1\}} \Delta^n \\ \downarrow^{i'} & & \downarrow^{j'} \\ \Delta^n & \longrightarrow \Delta^n \times \Delta^1 \end{array}$$

Then  $\tau'$  is weakly equivalent to  $\tau$  and is a projective cofibration.

**Example A.5.** Let  $\tau'$  be the following morphism in  $S^{[1]}$ :

$$\begin{array}{ccc} \partial \Delta^{n} & \longrightarrow & \partial \Delta^{n} \times \Delta^{1} \sqcup_{\partial \Delta^{n} \times \{1\}} \Delta^{n} \\ \downarrow \downarrow & & \downarrow j' \\ \Lambda^{n+1} & \longrightarrow & \Lambda^{n+1} \times \Delta^{1} \sqcup_{\Lambda^{n+1} \times \{1\}} \Delta^{n+1} \end{array}$$

Then  $\tau'$  is weakly equivalent to  $\tau$  and is a projective cofibration.

**Example A.6.** Let D be the simplicial set defined as the quotient  $\Delta^2/d_0$  where  $d_0: \Delta^1 \to \Delta^2$  is the face that misses 0. The two remaining faces  $d_1, d_2$  give two inclusions  $\Delta^1 \to D$ . Let  $\tau'$  be the following morphism in  $S^{[1]}$ :

Then  $\tau'$  is weakly equivalent to  $\tau$  and is a projective cofibration.

So, in view of the previous result, a map  $f: X \to Y$  between Kan complexes is a weak equivalence if and only if

$$(\tau')^*: \operatorname{map}(j',f) \to \operatorname{map}(i',f)$$

is surjective for  $\tau':i'\to j'$  as in the examples above.

#### A.b Simplicial spaces

**Definition A.7.** A simplicial space is a contravariant functor from  $\Delta$  to spaces (alias simplicial sets).

A simplicial space  $[m] \mapsto X_m$  may be viewed as a bisimplicial set, i.e., a contravariant functor from  $\Delta \times \Delta$  to Sets. However, the two  $\Delta$  directions play different roles and it is important to distinguish them.

A map  $X \to Y$  between simplicial spaces is a (degreewise) weak equivalence if for each  $m \ge 0$  the map  $X_m \to Y_m$  is a weak equivalence of spaces. We write  $\mathbf{R} \operatorname{map}(X,Y)$  for the homotopy function complex with respect to degreewise weak equivalences. This may be computed as  $\operatorname{map}(X^c,Y^f)$  in a model structure on simplicial spaces with levelwise weak equivalences, for a cofibrant replacement  $X^c \to X$  and a fibrant replacement  $Y \to Y^f$ . There are two canonical choices for such a model structure: the Reedy (= injective) model structure and the projective model structure.

The diagonal functor  $d: sS \to S$  has a left adjoint  $d_!$  which is the unique colimit-preserving functor with  $d_!(\Delta^n) = \Delta^n \otimes \Delta[n]$ . (For a simplicial set K and a simplicial space X, the tensor  $K \otimes X$  is the simplicial space with n-simplices  $K \times X_n$ .)

There is another colimit-preserving functor  $\delta: \mathsf{S} \to \mathsf{sS}$  defined by  $\delta(\Delta^n) = \Delta[n]$ , i.e., pullback along the projection onto the first factor  $\Delta \times \Delta \to \Delta$ . The projection  $\Delta^n \otimes \Delta[n] \to \Delta^0 \otimes \Delta[n]$  induces a natural transformation  $d_! \to \delta$ .

**Lemma A.8.** For each simplicial set X, the natural map  $d_!(X) \to \delta(X)$  is a degreewise weak equivalence of simplicial spaces.

*Proof.* For representables, this is clear. A general simplicial set X is a filtered colimit of finite-dimensional simplicial sets  $X_i$  and filtered colimits of simplicial spaces are homotopy colimits, so it is enough to prove the statement for finite-dimensional simplicial sets. Suppose that we have proved the statement for all simplicial sets of dimension < n. We want to prove the statement for a simplicial set X of dimension n. Let  $\mathrm{sk}_{n-1}X$  denote the  $(n-1)^{th}$  skeleton of X, so that we have a pushout

$$\bigcup_{X_n} \partial \Delta^n \longrightarrow \operatorname{sk}_{n-1} X$$

$$\downarrow \qquad \qquad \downarrow$$

$$\bigcup_{X_n} \Delta^n \longrightarrow X$$

Since  $d_!$  and  $\delta$  are colimit-preserving, the result follows by induction and the case of representables.

**Proposition A.9.** Let  $f: X \to Y$  be a map between Reedy fibrant simplicial spaces which satisfy the Kan condition. Then  $|f|: |X| \to |Y|$  is a weak equivalence if and only if every square

$$\partial \Delta[n] \xrightarrow{\gamma} X$$

$$\downarrow \qquad \qquad \downarrow$$

$$\Delta[n] \xrightarrow{\gamma} Y$$

has a lift as pictured making the lower triangle commute up to a given homotopy  $\Delta[n] \times \Delta[1] \to Y$  and making the upper triangle commute up to a given homotopy  $\partial \Delta[n] \times \Delta[1] \to X$ . These two homotopies are required to be homotopic as maps  $\partial \Delta[n] \times \Delta[1] \to Y$  and the homotopy should be constant on  $\partial \Delta[n] \times \partial \Delta[1]$ .

*Proof.* Since X and Y are Kan complexes and d preserves Kan fibrations (Remark 3.20), |X| and |Y| are Kan complexes. In view of Proposition A.1, Example A.6 and the remarks that follow it, |f| is a weak equivalence if and only if the map

$${\tau'}^*: \operatorname{map}(i', |f|) \to \operatorname{map}(j', |f|)$$

is surjective (using the notation from Example A.6). By adjunction, this is equivalent to saying that

$$(d_i \tau')^* : \operatorname{map}(d_i i', f) \to \operatorname{map}(d_i j', f) \tag{A.10}$$

is surjective. Since f is a map between Reedy fibrant simplicial spaces, it is a fibrant object in  $\mathsf{sS}^{[1]}$  with the *projective* model structure on the category of functors  $[1] \to \mathsf{sS}$ , where  $\mathsf{sS}$  is equipped with the Reedy model structure. Since  $d_!$  preserves monomorphisms,  $d_!\tau$  is a cofibration between cofibrant objects in that same model structure (see the proof of Corollary A.2). Therefore, the map (A.10) is a fibration and as such it is surjective if and only if it is surjective on  $\pi_0$ . These considerations also lead us to identify (A.10) with the map on derived mapping spaces

$$(d_1\tau')^*$$
:  $\mathbf{R}$ map $(d_1i', f) \to \mathbf{R}$ map $(d_1j', f)$ ,

which by Lemma A.8 is identified with

$$\delta(\tau')^* : \mathbf{R} \operatorname{map}(\delta(i'), f) \to \mathbf{R} \operatorname{map}(\delta(j'), f).$$

Since  $\delta(\tau')$  is also a cofibration between cofibrant objects, this map is identified with

$$\delta(\tau')^* : \operatorname{map}(\delta(i'), f) \to \operatorname{map}(\delta(j'), f).$$

The surjectivity of this last map is equivalent to the existence of the lift in the statement of the proposition.

Corollary A.11. Let  $f: X \to Y$  be a map between Reedy fibrant simplicial spaces. Suppose that for every  $j \ge 0$  and every square

$$\operatorname{Sd}^{j} \partial \Delta[n] \xrightarrow{\qquad} X$$

$$\downarrow \qquad \qquad \downarrow$$

$$\operatorname{Sd}^{j} \Delta[n] \xrightarrow{\qquad} Y$$

there is a lift as pictured making the lower triangle commute up to a given homotopy  $\mathrm{Sd}^j \Delta[n] \times \Delta[1] \to Y$  and making the upper triangle commute up to a given homotopy  $\mathrm{Sd}^j \partial \Delta[n] \times \Delta[1] \to X$ . These two homotopies are required to be homotopic as maps  $\mathrm{Sd}^j \partial \Delta[n] \times \Delta[1] \to Y$  and the homotopy should be constant on  $\mathrm{Sd}^j \partial \Delta[n] \times \partial \Delta[1]$ . Then  $|f|: |X| \to |Y|$  is a weak equivalence.

*Proof.* Apply Proposition A.9 replacing X and Y by the simplicial spaces  $\operatorname{Ex}^{\infty} X$  and  $\operatorname{Ex}^{\infty} Y$ , which satisfy the Kan condition by Proposition A.13.

#### A.c Subdivisions and Ex

Recall the simplicial subdivision sd  $\Delta^n$ , i.e., the nerve of the poset of non-empty subsets of  $[n] = \{0, \dots, n\}$ .

**Definition A.12.** Denote by

$$\mathrm{Sd}:\mathsf{sS}\to\mathsf{sS}$$

the simplicial left adjoint functor that sends  $\Delta[n]$  to sd  $\Delta^n$  viewed as a simplicial discrete space. Denote by

$$\mathrm{Ex}:\mathsf{sS}\to\mathsf{sS}$$

the simplicial right adjoint functor of Sd.

Every simplicial discrete space is Reedy cofibrant, so replacing X by a Reedy fibrant simplicial space  $X^f$ , we may write the right derived functor of Ex evaluated at X as the (honest) mapping space map(Sd  $\Delta[n], X^f$ ).

There is a natural map  $\gamma: \operatorname{Sd}\Delta[n] \to \Delta[n]$ , sending a subset  $\{i_0, \ldots, i_k\} \subset [n]$  to  $i_k$  (the last vertex). The colimit

$$X \xrightarrow{\gamma^*} \operatorname{Ex} X \xrightarrow{\gamma^*} \operatorname{Ex}^2 X \to \cdots$$

is denoted by  $\operatorname{Ex}^{\infty} X$ . The map  $\gamma$  has a section  $\Delta[n] \to \operatorname{Sd} \Delta[n]$  so if X is Reedy fibrant, all the maps in the tower are degreewise cofibrations and so the colimit computes the homotopy colimit.

We collect the important properties of the  $\operatorname{Ex}^{\infty}$  endofunctor below. These parallel (or, rather, include) the well-known ones for simplicial *sets*.

**Proposition A.13.** For a simplicial space X,


- 1.  $\operatorname{Ex}^{\infty} X$  is a Kan simplicial space,
- 2.  $X \to \operatorname{Ex}^{\infty} X$  is a weak equivalence after geometric realization,
- 3. for each i, including  $i = \infty$ ,  $\operatorname{Ex}^i$  preserves (trivial) Kan fibrations, zero simplices and finite homotopy limits

*Proof.* By construction, the functor  $\operatorname{Ex}^i$ , for  $0 \le i \le \infty$ , sends weak equivalences of simplicial spaces to weak equivalences. If X is Reedy fibrant then

$$\operatorname{map}(\operatorname{Sd}\Delta[n],X) \to \operatorname{map}(\operatorname{Sd}\partial\Delta[n],X)$$

is a fibration (since  $\operatorname{Sd} \partial \Delta[n] \to \operatorname{Sd} \Delta[n]$  is a degreewise monomorphism, hence a cofibration). Therefore,  $\operatorname{Ex} X$  is Reedy fibrant. By standard compactness arguments, it follows that  $\operatorname{Ex}^{\infty} X$  is also Reedy fibrant. Hence, in proving (1), (2) and (3), we may assume from the outset that X is Reedy fibrant.

The arguments to prove (1) and (3) are identical to the classical ones for simplicial sets, so we do not reproduce them here. As for (2), take a trivial Kan fibration  $X' \to X$  where X' is a simplicial set (see Lurie [12, Proposition 7]), and consider the square



Since the diagonal preserves trivial Kan fibrations, the vertical maps are weak equivalences after applying the diagonal (for the right-hand one, use part (3)). The top horizontal map is a weak equivalence, e.g., see Goerss–Jardine [8, III.4.6]. We conclude that the diagonal of the lower map is a weak equivalence.

## References

- [1] David Ayala, John Francis, Nick Rozenblyum. A stratified homotopy hypothesis. Journal of the European Mathematical Society 21:4 (2019), 1071–1178. arXiv:1502.01713v4, doi:10.4171/jems/856. 1
- [2] Pedro Boavida de Brito, Michael Weiss. Manifold calculus and homotopy sheaves. Homology, Homotopy and Applications 15:2 (2013), 361–383. arXiv:1202.1305v2, doi:10.4310/hha.2013.v15.n2.a20. 5
- [3] Ulrich Bunke, Thomas Nikolaus, Michael Völkl. Differential cohomology theories as sheaves of spectra. Journal of Homotopy and Related Structures 11:1 (2016), 1–66. arXiv:1311.3188v1, doi:10.1007/s40062-014-0092-5.
- [4] Albrecht Dold. Partitions of Unity in the Theory of Fibrations. The Annals of Mathematics 78:2 (1963), 223–255. doi:10.2307/1970341. 3.a
- [5] Albrecht Dold, René Thom. Quasifaserungen und Unendliche Symmetrische Produkte. The Annals of Mathematics 67:2 (1958), 239–281. doi:10.2307/1970005. 3.16
- [6] Daniel Dugger. Universal homotopy theories. Advances in Mathematics 164:1 (2001), 144–176. arXiv:math/0007070v1, doi:10.1006/aima.2001.2014. 2.a
- [7] Daniel Dugger, Daniel C. Isaksen. Weak equivalences of simplicial presheaves. (Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-Theory.) Contemporary Mathematics 346 (2004), 97–113. arXiv:math/0205025v1, doi:10.1090/conm/346/06292. A.1
- [8] Paul G. Goerss, John F. Jardine. Simplicial homotopy theory. Progress in Mathematics 174 (1999). doi:10.1007/978-3-0346-0189-4. A.c
- [9] Daniel Grady, Dmitri Pavlov. Extended field theories are local and have classifying spaces. 2020. arXiv:2011.01208. 1
- [10] Philip Herrmann, Florian Strunk. Enriched simplicial presheaves and the motivic homotopy category. Journal of Pure and Applied Algebra 215:7 (2011), 1663–1668. arXiv:1007.3153v1, doi:10.1016/j.jpaa.2010.10.002. 2
- [11] Jacob Lurie. Higher Topos Theory. April 9, 2017. https://www.math.ias.edu/~lurie/papers/HTT.pdf. 1

- [12] Jacob Lurie. Simplicial spaces. February 6, 2011. https://www.math.ias.edu/~lurie/287xnotes/Lecture7.pdf. A.c
- [13] Ib Madsen, Michael Weiss. The stable moduli space of Riemann surfaces: Mumford's conjecture. Annals of Mathematics 165:3 (2007), 843–941. arXiv:math/0212321v3, doi:10.4007/annals.2007.165.843. 1, 1, 4.19
- [14] Izak Moerdijk. Classifying Spaces and Classifying Topoi. Lecture Notes in Mathematics 1616 (1995). doi:10.1007/bfb0094441. 6
- [15] Fabien Morel, Vladimir Voevodsky. A¹-homotopy theory of schemes. Publications mathématiques de l'IHÉS 90:1 (1999), 45–143. doi:10.1007/bf02698831. 1, 1, 2, 2.a
- [16] Dmitri Pavlov. Numerable open covers and representability of topological stacks. 2022. arXiv:2203.03120. 5.4
- [17] Charles Rezk. A model for the homotopy theory of homotopy theory. Transactions of the American Mathematical Society 353:3 (2001), 973–1007. arXiv:math/9811037v3, doi:10.1090/s0002-9947-00-02653-2. 6
- [18] Charles Rezk. When are homotopy colimits compatible with homotopy base change? February 2, 2014. https://faculty.math.illinois.edu/~rezk/i-hate-the-pi-star-kan-condition.pdf, 3.14, 3.21
- [19] Urs Schreiber. Differential cohomology in a cohesive ∞-topos. arXiv:1310.7930v1. 1
- [20] Stephan Stolz, Peter Teichner. Supersymmetric field theories and generalized cohomology. (Mathematical Foundations of Quantum Field Theory and Perturbative String Theory.) Proceedings of Symposia in Pure Mathematics 83 (2011), 279–340. arXiv:1108.0189v1, doi:10.1090/pspum/083/2742432. 1
- [21] Friedhelm Waldhausen. Algebraic K-theory of spaces. (Algebraic and Geometric Topology.) Lecture Notes in Mathematics 1126 (1985), 318–419. doi:10.1007/bfb0074449. 1
- [22] Michael Weiss. Embeddings from the point of view of immersion theory: Part I. Geometry & Topology 3:1 (1999), 67–101. arXiv:math/9905202, doi:10.2140/gt.1999.3.67. 5
- [23] Michael Weiss. What does the classifying space of a category classify? Homology, Homotopy and Applications 7:1 (2005), 185–195. doi:10.4310/hha.2005.v7.n1.a10. 6
- [24] Michael Weiss, Bruce Williams. Assembly. (Novikov Conjectures, Index Theorems, and Rigidity.) London Mathematical Society Lecture Notes 227 (1995), 332–352. doi:10.1017/cbo9780511629365.014.