

RANKS OF ABELIAN VARIETIES AND THE FULL MORDELL-LANG CONJECTURE IN DIMENSION ONE

ARNO FEHM AND SEBASTIAN PETERSEN

ABSTRACT. Let A be a non-zero abelian variety over a field F that is not algebraic over a finite field. We prove that the rational rank of the abelian group $A(F)$ is infinite when F is large in the sense of Pop (also called ample). The main ingredient is a deduction of the 1-dimensional case of the relative Mordell-Lang conjecture from a result of Rössler.

1. INTRODUCTION

A considerable amount of work has been done to show that over certain fields, for example over certain infinite algebraic extensions of global fields, all abelian varieties have infinite rank. Here, the **rank** of an abelian variety A over a field F is the rank of the Mordell-Weil group $A(F)$, i.e. $\dim_{\mathbb{Q}}(A(F) \otimes \mathbb{Q})$. See [Ros73, FJ74, RW02, Lar03, Im06, GJ06, Pet06, IL08, IL13] to mention a few examples.

In all cases known to us where one actually succeeded in proving that over a field every abelian variety has infinite rank, these fields turn out to be ample or are conjectured to be ample, where a field F is called **ample** (or **large** following [Pop96]) if every smooth F -curve C satisfies $C(F) = \emptyset$ or $|C(F)| = \infty$. For example, every separably closed, real closed, or Henselian valued field is ample. See [Pop96, Jar03, Pop10, FP11] for many more examples of ample fields, and [BSF12, Pop14] for surveys on the importance and ubiquity of ample fields in contemporary Galois theory and other areas.

In this work, we prove (see Section 5) our conjecture from [FP10]:

Theorem 1.1. *Let F be an ample field that is not algebraic over a finite field and let A/F be a non-zero abelian variety. Then the rank of $A(F)$ is infinite.*

In [FP10] we proved several special cases of Theorem 1.1. In particular, we prove the case when F has characteristic zero. In this work, we complete the case of positive characteristic. In the proof of Theorem 1.1, we use and strengthen results of Ghioca and Moosa [GM06] on the Mordell-Lang conjecture in positive characteristic: The 1-dimensional case of the Mordell-Lang conjecture is reduced to a special case, which is then solved by recent work of Rössler [Rös19]. What we get (see Section 4) is the following:

Theorem 1.2. *Let K be an algebraically closed field of positive characteristic p , A/K an abelian variety, C a closed subcurve of A , and $\Lambda \subseteq A(K)$ a subgroup of finite rank. If $\Lambda \cap C(K)$ is infinite, then C is elliptic or birational to a curve defined over the algebraic closure of $\mathbb{F}_p$.*

Theorem 1.1 reproves and generalizes several previous infinite rank results. In Section 6 we give a few concrete applications of Theorem 1.1.

NOTATION

Let K be a field. We denote by $\tilde{K}$ the algebraic closure of K , by K_{sep} the maximal separable, and by K_{ins} the maximal purely inseparable extension of K in $\tilde{K}$. If V is a K -scheme and F/K is a field extension, then $V_F := V \times_{\text{Spec}(K)} \text{Spec}(F)$. If V and W are K -schemes and $f: V \rightarrow W$ is a K -morphism, then $f_F: V_F \rightarrow W_F$ stands for the base extension of f . Throughout this paper, a K -variety is a separated algebraic K -scheme which is geometrically integral, and a K -curve is a K -variety of dimension 1. The absolute genus of a K -curve C is the genus $g(C')$ of a smooth projective model C' of $C_{\tilde{K}}$. We denote by $J_{C'}$ the Jacobian variety of C' . Subvarieties are always understood to be closed in the ambient variety. If $f: V \rightarrow W$ is a morphism and $X \subseteq W$, then we define $f^{-1}(X) := (V \times_W X)_{\text{red}}$, the reduced induced scheme structure on the fiber product. If $f: G \rightarrow H$ is a homomorphism of group schemes over K , then $\ker(f) := G \times_{H,e} \text{Spec}(K)$ is the scheme theoretic fibre of the unit section e .

2. ISOTRIVIAL AND SPECIAL CURVES

In this section we prove some preliminary results about isotrivial curves and special curves in the sense of Hrushovski, cf. [Hru96]. The main conclusion here, probably known to experts, is that a subcurve of an abelian variety is special if and only if it is elliptic or isotrivial.

Throughout this section, let K/K_0 be an extension of algebraically closed fields.

A K -variety V is **K/K_0 -isotrivial** (or simply isotrivial) if there exists a K_0 -variety V_0 and a K -birational map $V \dashrightarrow V_{0,K}$. Note that if V is a smooth projective curve, then V is K/K_0 -isotrivial if and only if there exists a K_0 -variety V_0 and a K -isomorphism $V \rightarrow V_{0,K}$, i.e. V descends to K_0 . If F is a subfield of K , then we say that an F -variety V is K/K_0 -isotrivial if V_K is K/K_0 -isotrivial.

The following lemma also appears as [BBP14, Fact 2.15, Remark 2.17], but we include a full proof for the convenience of the reader:

Lemma 2.1. *Let A/K be an abelian variety. Assume that there exists an abelian variety B_0/K_0 and an isogeny $\varphi: B_{0,K} \rightarrow A$. If $\dim(A) = 1$ or if φ is separable, then there exists an abelian variety A_0/K_0 with $A_{0,K} \cong A$.*

Proof. The kernel $N := \ker(\varphi)$ is a finite subgroup scheme of $B_{0,K}$. If $\dim(A) = 1$, then $\dim(B_0) = 1$ and thus there exists a finite subgroup scheme N_0/K_0 of B_0 such that $N = N_{0,K}$, see [Con06, Theorem 2.3]. If φ is separable, then N/K is a finite étale subgroup scheme of $B_{0,K}$ and thus there exists a finite subgroup scheme N_0/K_0 of B_0 such that $N = N_{0,K}$, see [Con06, Lemma 3.11]. In both cases it follows that $A_0 := B_0/N_0$ is an abelian variety over K_0 (see [Con06, 3.7] for the existence of the quotient) and $A_{0,K} = B_{0,K}/N_{0,K} = B_{0,K}/N \cong A$, as desired. $\square$

Lemma 2.2. *Let A/K be an abelian variety. Assume that there exist an abelian variety B_0/K_0 and a surjective homomorphism $B_{0,K} \rightarrow A$. Then there exists an abelian variety A_0/K_0 and a purely inseparable isogeny $A \rightarrow A_{0,K}$.*

Proof. By the Poincare reducibility theorem [Mil86, 12.1] there exists an abelian subvariety B' of $B_{0,K}$ such that A is K -isogenous to B' . By [Con06, 3.21] there exists an abelian subvariety B'_0 of B_0 such that $B' \cong B'_{0,K}$. After replacing B_0 by B'_0 we can assume that there exists an isogeny $\varphi : B_{0,K} \rightarrow A$. We put $B := B_{0,K}$.

There exists $n \in \mathbb{N}$ such that $\ker(\varphi)$ is a subgroup scheme of $\ker([n]_B)$. We define $N_0 := \ker([n]_{B_0})$ and $N := N_{0,K}$. The projection $p : B \rightarrow B/N$ induces an isogeny $f : A \rightarrow B/N$ such that $f \circ \varphi = p$ (see [Con06, 3.8]). We denote by $\ker(f)^\circ$ the connected component of the unit element of $\ker(f)$ and by $f_i : A \rightarrow A' := A/\ker(f)^\circ$ the canonical projection. Then f factors through A' , i.e. there exists an isogeny $f_s : A' \rightarrow B/N$ such that $f_s \circ f_i = f$ (see [Con06, 3.7]). We now have constructed isogenies

$$B \xrightarrow{\varphi} A \xrightarrow{f_i} A' \xrightarrow{f_s} B/N$$

such that $f_s \circ f_i \circ \varphi = p$. The restriction $f_i \circ \varphi|_{N^\circ} \rightarrow \ker(f_s)$ is trivial because N° is connected and $\ker(f_s) \cong \ker(f)/\ker(f)^\circ$ is étale (see [Con06, 3.10]). It follows that $\ker(f_i \circ \varphi) \supset N^\circ$ and thus $f_i \circ \varphi$ factors through an isogeny $g : B/N^\circ \rightarrow A'$ by [Con06, 3.7]. The kernel of g is a closed subgroup scheme of the finite étale group scheme N/N° , hence $\ker(g)$ is étale and $g : B/N^\circ \rightarrow A'$ is a *separable* isogeny. But $B/N^\circ = (B_0/(N_0)^\circ)_K$ and thus Lemma 2.1 implies that there exists an abelian variety A_0/K_0 such that $A' \cong A_{0,K}$. $\square$

Lemma 2.3. *Let $\varphi : D \dashrightarrow C$ be a dominant rational map of K -curves. If D is K/K_0 -isotrivial, then C is K/K_0 -isotrivial.*

Proof. Since K and K_0 are algebraically closed it suffices to consider the case that K/K_0 is transcendental. Without loss of generality we can assume that C and D are smooth projective and that $\varphi : D \rightarrow C$ is a non-constant morphism. Let D_0 be a smooth projective K_0 -curve with an isomorphism $\psi : D_{0,K} \rightarrow D$.

If $g(C) = 0$, then C is isomorphic to $\mathbb{P}_K^1$, hence C is K/K_0 -isotrivial.

If $g(C) = 1$, then C is an elliptic curve. By Albanese functoriality there is a surjective homomorphism of abelian varieties $(J_{D_0})_K \cong J_D \rightarrow C$, and therefore Lemma 2.2 implies that there exists an elliptic curve C_0/K_0 such that C is K -isogenous to $C_{0,K}$. Lemma 2.1 then shows that C is K/K_0 -isotrivial.

We may thus assume that $g(C) \geq 2$. Let $K_0 \subseteq F \subseteq K$ be an intermediate field such that F/K_0 is finitely generated and transcendental, and C, D, φ and ψ descend to F , i.e. there exist smooth projective F -curves C_1 and D_1 with $C_{1,K} = C$ and $D_{1,K} = D$, a morphism $\varphi_1 : D_1 \rightarrow C_1$ such that $\varphi_{1,K}$ corresponds to φ , and an isomorphism $D_{0,F} \cong D_1$. Since K_0 is algebraically closed, $|D_1(F)| = |D_0(F)| \geq |D_0(K_0)| = \infty$. Since $\varphi_1 : D_1(F) \rightarrow C_1(F)$ has finite fibres, this implies that $|C_1(F)| = \infty$. Thus, since C_1 is of genus at least 2, the Grauert-Manin theorem (see [Sam66] or [FJ08, Prop. 11.7.1.]) implies that C_1 is $\tilde{F}/K_0$ -isotrivial, and hence C is K/K_0 -isotrivial. $\square$

Lemma 2.4. *Let K_1 be an algebraically closed intermediate field of K/K_0 . Let C be a K_1 -curve. If C is K/K_0 -isotrivial, then C is K_1/K_0 -isotrivial.*

Proof. We can assume that C is smooth and projective. As C is K/K_0 -isotrivial, there exists a K_0 -curve C_0 and a K -isomorphism $C_K \rightarrow C_{0,K}$. Therefore there exists a finitely generated extension F/K_1 inside K and an F -isomorphism $f : C_F \rightarrow C_{0,F}$. Choose a K_1 -variety S with function field F . By [GD66, 8.8.2, 8.10.5] there exists a non-empty open subscheme S' of S such that f extends to an S -isomorphism $C \times_{K_1} S' \rightarrow C_{0,K_1} \times_{K_1} S'$. As K_1 is algebraically closed there exists $s \in S'(K_1)$, and specializing at s gives a K_1 -isomorphism $C \rightarrow C_{0,K_1}$. Hence C is K_1/K_0 -isotrivial. $\square$

Let A/K be an abelian variety. A subvariety $X \subseteq A$ is K/K_0 -**special** if there exist

- an abelian subvariety A' of A ,
- an abelian variety B_0/K_0 ,
- a K_0 -subvariety $X_0 \subseteq B_0$,
- a surjective K -homomorphism $h : A' \rightarrow B_{0,K}$,
- and an element $g \in A(K)$

such that $X = g + h^{-1}(X_{0,K})$. We write t_g for the translation by g on A .

The situation is as follows:

$$\begin{array}{ccccc} A & \xleftarrow{\quad \supset \quad} & A' & \xrightarrow{h} & B_{0,K} \\ \uparrow & & \uparrow & & \uparrow \\ X & \xleftarrow{t_g} & h^{-1}(X_{0,K}) & \xrightarrow{h} & X_{0,K} \end{array}$$

Proposition 2.5. *Let A/K be an abelian variety and $C \subseteq A$ a K -subcurve. Then the following statements are equivalent.*

- (1) *C is a K/K_0 -special subvariety of A .*
- (2) *C is of absolute genus 1 or K/K_0 -isotrivial.*

Proof. PROOF OF (1) $\Rightarrow$ (2).

If C is special, then there exist an abelian subvariety $A' \subseteq A$, an abelian variety B_0/K_0 , a K_0 -subvariety $C_0 \subseteq B_0$, a surjective K -homomorphism $h : A' \rightarrow B_{0,K}$, and an element $g \in A(K)$ such that $C = g + h^{-1}(C_{0,K})$. Without loss of generality assume that $0_{B_0} \in C_0$, $0_A \in C$, and $g = 0_A$. Then $C \subseteq A'$, so assume without loss of generality that $A = A'$.

If $\dim(C_0) = 0$, then $C = \ker(h)$ is an elliptic curve, hence of absolute genus 1.

If $\dim(C_0) \geq 1$, then $\dim(C_0) = \dim(C) = 1$, so h has finite fibers over $C_{0,K}$, and hence all fibers of h are finite. Consequently, h is an isogeny. Let $\hat{h} : B_{0,K} \rightarrow A$ be an isogeny with $h \circ \hat{h} = [n]_{B_{0,K}}$ for some $n \in \mathbb{N}$, and let C'_0 be an irreducible component of $[n]_{B_0}^{-1}(C_0)$ such that $[n]_{B_0}(C'_0) = C_0$. Now we have a non-constant morphism

$$C'_{0,K} \xrightarrow{\hat{h}} C = h^{-1}(C_{0,K}) \xrightarrow{h} C_{0,K}$$

and Lemma 2.3 implies that C is K/K_0 -isotrivial.

PROOF OF (2) $\Rightarrow$ (1).

If C is of absolute genus 1, then the normalization $\lambda : \bar{C} \rightarrow C$ of C is an elliptic curve. By [Mil86, Cor. 2.2] there exists $g \in A(K)$ and a homomorphism $\lambda' : \bar{C} \rightarrow A$ of abelian varieties such that $\lambda = t_g \circ \lambda'$. It follows that $A' := \text{im}(\lambda')$ is an abelian subvariety of A and $C = g + A'$. If B_0 is the zero abelian variety over K_0 and $h : A' \rightarrow B_{0,K}$ the zero homomorphism, then $A' = h^{-1}(0_{B_{0,K}})$ and thus $C = g + h^{-1}(0_{B_{0,K}})$ is special.

From now on assume that C is isotrivial. We have to prove that C is special. There exists a smooth projective K_0 -curve C_0 and a finite birational morphism $f : C_{0,K} \rightarrow C$. Let $P_0 \in C_0(K_0)$ be a point and let $\alpha_0 : C_0 \rightarrow J_{C_0}$ be the corresponding Albanese map (sending P_0 to 0). It is a closed immersion because the genus of C_0 must be at least 1 by [Mil86, 3.8]. By applying a translation on A we can assume without loss of generality that $f(P_0) = 0_A$. By Albanese functoriality, the morphism f induces a homomorphism $f' : J_{C_{0,K}} \rightarrow A$ with $f' \circ \alpha_{0,K} = f$. Thus C lies on the abelian subvariety $A' := f'(J_{C_{0,K}})$, and $C = f'(\alpha_{0,K}(C_{0,K}))$. Lemma 2.2 gives an abelian variety B_0/K_0 and a purely inseparable isogeny $h : A' \rightarrow B_{0,K}$. Now consider the morphisms

$$C_{0,K} \xrightarrow{\alpha_{0,K}} J_{C_{0,K}} \xrightarrow{f'} A' \xrightarrow{h} B_{0,K}$$

By [Mil86, 20.4(b)] the homomorphism $h \circ f'$ is defined over K_0 , that is, there exists a homomorphism $\beta_0 : J_{C_0} \rightarrow B_0$ such that $h \circ f' = \beta_{0,K}$. Now $C'_0 := \beta_0(\alpha_0(C_0))$ is a subvariety of B_0 such that $C'_{0,K} = h(f'(\alpha_{0,K}(C_{0,K})))$. As h is a purely inseparable isogeny, the topological space underlying each fibre is a singleton. Hence $h : A' \rightarrow B_{0,K}$ is a homeomorphism on the underlying topological spaces of A' and $B_{0,K}$. It follows that

$$h^{-1}(C'_{0,K}) = (C'_{0,K} \times_{B_{0,K}} A')_{\text{red}} = f'(\alpha_{0,K}(C_{0,K})) = C.$$

Thus C is special. $\square$

3. RÖSSLER'S THEOREM ON INSEPARABLE POINTS ON CURVES

In this section we recall and slightly generalize a recent result of Rössler about inseparable points on curves, which strengthens the aforementioned Grauert-Manin theorem.

Theorem 3.1. *Let F/F_0 be a finitely generated extension of fields of positive characteristic, and let C be an F -curve of absolute genus $g \geq 2$. If $C(F_{\text{ins}})$ is infinite, then C is $\tilde{F}/\tilde{F}_0$ -isotrivial.*

In the following proof we deduce Theorem 3.1 from Rössler's [Rös19, Thm. 1.1], where this theorem is proven in the special case where F_0 is algebraically closed, $\text{trdeg}(F/F_0) = 1$ and C is smooth and projective.

Proof. There exists a regular projective curve over F_{ins} that is birational to $C_{F_{\text{ins}}}$. This curve is automatically geometrically regular because F_{ins} is perfect (cf. [GD65, 6.7.7]), hence smooth over F_{ins} by [GD67, 17.5.2]. It follows that there exists a finite extension E/F inside F_{ins} and a smooth projective E -curve $\bar{C}$ that is birational to C_E . Thus $\bar{C}(E_{\text{ins}})$ is infinite and it suffice to prove that

$\bar{C}$ is $\tilde{F}/\tilde{F}_0$ -isotrivial. We can thus assume for the rest of the proof that C is smooth and projective.

Let $\mathcal{F}$ be the set of all algebraically closed intermediate fields F' of $\tilde{F}/\tilde{F}_0$ with $\text{trdeg}(\tilde{F}/F') = 1$. Then $\tilde{F}_0 = \bigcap_{F' \in \mathcal{F}} F'$: Indeed, every $x \in \tilde{F} \setminus \tilde{F}_0$ is an element of a transcendence base $(x, x_2, \dots, x_n)$ of $\tilde{F}/\tilde{F}_0$ and x is not contained in the algebraic closure F' of $F_0(x_2, \dots, x_n)$.

For every $F' \in \mathcal{F}$, $C((F'F)_{\text{ins}}) \supseteq C(F_{\text{ins}})$ is infinite, so [Rös19, Thm. 1.1] gives that C is $\tilde{F}/F'$ -isotrivial. Therefore, C is in fact $\tilde{F}/\bigcap_{F' \in \mathcal{F}} F'$ -isotrivial: This can be seen by looking at the point $c \in M_g(\tilde{F})$ corresponding to C on the coarse moduli space M_g for smooth proper curves of genus g , see [MFK92, 5.6, 7.14], or explicitly by [Koi72, Corollary 3.2.2]. $\square$

Remark 3.2. *We note that this theorem could also be deduced from the older [Kim97] instead of [Rös19]: Let F be a function field of one variable over a field F_0 of positive characteristic p . Consider an affine plane curve*

$$C = \text{Spec}(F[X, Y]/(f(X, Y)))$$

where $f(X, Y) \in F[X, Y]$ is absolutely irreducible. We denote by $f^{(n)}(X, Y)$ the polynomial obtained from $f(X, Y)$ by raising the coefficients of $f(X, Y)$ to the p^n -th power and by

$$C^{(n)} := \text{Spec}(F[X, Y]/(f^{(n)}(X, Y)))$$

the corresponding curve. For $n \geq 1$ there is the Frobenius injection

$$\text{Fr}: C^{(n-1)}(F) \rightarrow C^{(n)}(F), (x, y) \mapsto (x^p, y^p)$$

and we define

$$C^{(n)}(F)_{\text{new}} := C^{(n)}(F) \setminus \text{Fr}(C^{(n-1)}(F)).$$

Finally, $C^{(0)}(F)_{\text{new}} := C(F)$. Note that the Frobenius gives a bijection between $C^{(n)}(F)$ and $C(F^{p^{-n}})$, and hence between $C^{(n)}(F)_{\text{new}}$ and $C(F^{p^{-n}}) \setminus C(F^{p^{-n+1}})$. Then [Kim97, Theorem 1] claims that if the absolute genus of C is at least 2 and $C^{(n)}(F)_{\text{new}} \neq \emptyset$ for infinitely many n , then C is $\tilde{F}/\tilde{\mathbb{F}}_p$ -isotrivial.

This, however, is incorrect, as the following example shows: Let F_0 be any non-perfect transcendental ample field (e.g. $F_0 = \mathbb{F}_p(t)_{\text{sep}}$), F any function field of one variable over F_0 (e.g. $F = F_0(X)$), and C any F_0 -curve of absolute genus at least 2 which is not $\tilde{F}_0/\tilde{\mathbb{F}}_p$ -isotrivial and has a smooth F_0 -rational point (see e.g. Lemma 5.1 below for the existence of such a curve). Then [Kim97, Theorem 1] would imply that there exists an n such that $C(F_{\text{ins}}) = C(F^{p^{-n}})$, hence $C(F_{0, \text{ins}}) = C(F_{0, \text{ins}}) \cap C(F^{p^{-n}}) = C(F_0^{p^{-n}})$, contradicting for example [Feh10, Corollary 5].

4. THE MORDELL-LANG CONJECTURE

The main ingredient in the proof of Theorem 1.1 is a partial result towards the general Mordell-Lang conjecture for function fields.

Conjecture 4.1 (General Mordell-Lang conjecture for function fields). *Let K/K_0 be an extension of algebraically closed fields, A/K an abelian variety,*

$X \subseteq A$ a subvariety, and $\Lambda \subseteq A(K)$ a subgroup of finite rank such that $\Lambda \cap X(K)$ is Zariski-dense in X . Then X is K/K_0 -special.

A slightly weaker result was proven by Hrushovski in [Hru96], where the condition that Λ has finite rank is replaced by the stronger condition that $\Lambda \otimes \mathbb{Z}_{(p)}$ is a finitely generated $\mathbb{Z}_{(p)}$ -module, where p is the characteristic of K .

In [GM06], Ghioca and Moosa use work of Scanlon and the theory of generic difference fields to reduce the general Mordell-Lang conjecture to the special case where $\Lambda \subseteq A(F_{\text{ins}})$ for some finitely generated field F . This reduction also works if on both sides one adds the condition that X is a curve. The following result is a slight generalization of [GM06, Corollary 2.6]:

Proposition 4.2. *Let K be an algebraically closed field of positive characteristic p , and let $K_0 = \mathbb{F}_p$. Let A/K be an abelian variety, C a subcurve of A , and $\Lambda \subseteq A(K)$ a subgroup of finite rank. If $\Lambda \cap C(K)$ is infinite, then the absolute genus of C is 1 or C is K/K_0 -isotrivial, i.e. C is K/K_0 -special.*

Proof. First of all, A and C are defined over a finitely generated subfield F of K . By the reduction [GM06, Theorem 2.2], we may assume that $\Lambda \subseteq A(F_{\text{ins}})$. This implies that $C(F_{\text{ins}})$ is infinite.

Since C is a subcurve of an abelian variety, the absolute genus of C is at least 1, [Mil86, 3.8]. If the absolute genus of C is 1, then C is K/K_0 -special by Proposition 2.5. If the absolute genus of C is at least 2, then C is K/K_0 -isotrivial by Theorem 3.1, and hence K/K_0 -special by Proposition 2.5. $\square$

This proves Theorem 1.2.

Remark 4.3. *Proposition 4.2 is stated in [GM06, Corollary 2.6] in the special case $A = J_C$. The proof there builds on [Kim97, Theorem 1], cf. Remark 3.2.*

5. PROOF OF THEOREM 1.1

The idea of the proof of Theorem 1.1 is to construct a curve on A that is non-special and has a smooth F -rational point. To achieve this, we ‘lift’ a non-isotrivial curve from $\mathbb{A}^2$ to A .

Lemma 5.1. *If F/F_0 is a transcendental field extension, then there exists a smooth F -curve C of absolute genus at least 2 with $C(F) \neq \emptyset$ such that C is not $\tilde{F}/\tilde{F}_0$ -isotrivial.*

Proof. Let $t \in F \setminus \tilde{F}_0$. First assume that $\text{char}(F) \neq 2$. Let

$$E : y^2 = f(x) = (x-1)(x+1)(x-t^2)$$

be the elliptic curve over F with j -invariant

$$j(E) = \frac{(16t^4 + 48)^3}{64t^8 - 128t^4 + 64}.$$

Since $j(E) \in F \setminus \tilde{F}_0$, it follows that E is not $\tilde{F}/\tilde{F}_0$ -isotrivial. The map $(x, y) \mapsto (x^2, y)$ defines a separable morphism from the curve

$$C : y^2 = f(x^2)$$

to E which is ramified over $(0, t) \in E(F)$. Therefore, C is a smooth F -curve of absolute genus at least 2 by the Riemann-Hurwitz formula, and $C(F) \neq \emptyset$. By Lemma 2.3, C is not $\tilde{F}/\tilde{F}_0$ -isotrivial.

If $\text{char}(F) = 2$, we can proceed similarly and let $E : y^2 + xy = f(x) = x^3 + tx + 1$ be the elliptic curve with j -invariant $j(E) = (t^2 + 1)^{-1}$. The map $(x, y) \mapsto (x^3, y)$ defines a separable morphism from the curve $C : y^2 + x^3y = f(x^3)$ to E that is ramified over $(0, 1) \in E(F)$. $\square$

Lemma 5.2. *Let F be a field. Let X be an F -variety of dimension n and Y an F -variety of dimension $m < n$. Let $x \in X(F)$ be a smooth point of X and let $y \in Y(F)$ a smooth point of Y . Then there exists an F -subvariety $V \subseteq X$ of dimension $\dim(V) = m$ such that x is a smooth point of V , and a dominant F -rational map $V \dashrightarrow Y$ mapping x to y .*

Proof. There exists a smooth open neighbourhood Y' of y in Y and an étale morphism $f : Y' \rightarrow \mathbb{A}^m$, see [GD67, 17.11.4]. By [Mil80, I.3.14] this morphism f is locally standard étale. Hence, after replacing Y' by a smaller open neighbourhood of y if necessary, there exists an affine open subscheme $U = \text{Spec}(R) \subseteq \mathbb{A}^m$ such that $f(Y') \subseteq U$ and $f|_{Y'} : Y' \rightarrow U$ is standard étale, i.e. such that there exist a polynomial $Q \in R[T]$ and a U -isomorphism of Y' with an open subscheme of $\text{Spec}(R[T]/(Q))$. In particular there exists an immersion

$$i : Y' \rightarrow \text{Spec}(R[T]) = U \times \mathbb{A}^1 \rightarrow \mathbb{A}^{m+1} \rightarrow \mathbb{A}^n.$$

There exists a smooth open neighbourhood X' of x in X and an étale morphism $g : X' \rightarrow \mathbb{A}^n$, again by [GD67, 17.11.4], and we can assume that $g(x) = i(y)$. We form the fibre product $Z = X' \times_{\mathbb{A}^n, g, i} Y' = g^{-1}(Y')$ and consider the diagram

$$\begin{array}{ccc} Z & \xrightarrow{j} & X' \subset X \\ \downarrow p & & \downarrow g \\ Y \supset Y' & \xrightarrow{i} & \mathbb{A}^n. \end{array}$$

The projection p is étale, Y' is smooth and $\dim(Y') = m$. Hence Z is smooth of dimension m . Furthermore j is an immersion. Hence we can identify Z with a closed subscheme of an open subscheme X'' of X' . We have $x \in Z(F)$. Let Z_0 be the connected component of Z passing through x and let V be the closure of Z_0 in X (with the reduced induced subscheme structure). Then Z_0 is smooth and connected and $Z_0(F) \neq \emptyset$, hence it is geometrically integral, so also V is geometrically integral, see [GD65, 4.6.3]. Hence V is a *subvariety* of X with $\dim(V) = m$. Finally the étale morphism $p|_{Z_0} : Z_0 \rightarrow Y'$ induces a dominant rational map $V \dashrightarrow Y$ mapping x to y . $\square$

Theorem 5.3. *Let F_1 be a field which is not algebraic over a finite field, and let A/F_1 be a non-zero abelian variety. Then there exists a smooth F_1 -curve C_A with $C_A(F_1) \neq \emptyset$ and the following property: If F is an extension of F_1 with $|C_A(F)| = \infty$, then the rank of $A(F)$ is infinite.*

Proof. Replace A by $A \times A$ to assume without loss of generality that A is of dimension at least 2. Take any smooth F_1 -curve C of absolute genus at least 2 with $C(F_1) \neq \emptyset$. By Lemma 5.2 there exist an F_1 -curve $D \subseteq A$ with a dominant F_1 -rational map $\varphi: D \dashrightarrow C$ such that D has a smooth F_1 -rational point. By the Riemann-Hurwitz formula, D is of absolute genus at least 2.

If F_1 is of characteristic zero and F is an extension of F_1 with $|D(F)| = \infty$, then the Mordell-Lang conjecture in characteristic zero implies that the rank of $A(F)$ is infinite, cf. [FP10, Theorem 2.6(a)].

If F_1 is of characteristic $p > 0$, then $K_1 = \tilde{F}_1$ is transcendental over $K_0 = \tilde{\mathbb{F}}_p$ and so we can choose the curve C to be not K_1/K_0 -isotrivial, see Lemma 5.1, and then also D is not K_1/K_0 -isotrivial by Lemma 2.3. If F is an extension of F_1 with $|D(F)| = \infty$, let $K = \tilde{F}$ and $\Lambda = A(F)$, and suppose that the rank of Λ is finite. Since $\Lambda \cap D(K) = D(F)$ is infinite and the absolute genus of D is at least 2, Proposition 4.2 implies that D is K/K_0 -isotrivial, and so Lemma 2.4 implies that D is K_1/K_0 -isotrivial, a contradiction. Thus the rank of $A(F)$ must be infinite.

Therefore, in both cases D contains a smooth open subcurve C_A that satisfies the requirements. $\square$

Theorem 1.1 is an immediate consequence of Theorem 5.3 for $F_1 = F$, since if F is ample, then $|C_A(F)| = \infty$.

6. COROLLARIES AND APPLICATIONS

6.1. Uncountable rank. It turns out that in the situation of Theorem 1.1, not only is the rank of $A(F)$ infinite, but it is as large as possible.

Corollary 6.1. *Let F be an ample field that is not algebraic over a finite field and let A/F be a non-zero abelian variety. Then the rank of $A(F)$ equals the cardinality of F .*

Proof. If F is countable, then the claim follows from Theorem 1.1. If F is uncountable, then $|A(F)| = |F|$ by [Har09, Prop. 3.3]. Since $A_{\text{tor}}(\tilde{F})$ is countable, this implies that $A(F) \otimes \mathbb{Q}$ cannot have a basis of less than $|F|$ elements. $\square$

6.2. Large extensions of Hilbertian fields. In [Pet06, Theorem 1.4], the second author generalizes [FJ74, Theorem 9.1] from finitely generated fields to arbitrary Hilbertian fields. This generalization also follows easily from our results. In the following, **almost all** is to be understood in the sense of Haar measure, and for an e -tuple $\sigma \in \text{Gal}(F)^e$, where $\text{Gal}(F) = \text{Aut}(F_{\text{sep}}/F)$ is the absolute Galois group of F , $F_{\text{sep}}(\sigma)$ is the fixed field in F_{sep} of the group generated by $\sigma_1, \dots, \sigma_e$.

Corollary 6.2. *If F is a Hilbertian field, A/F a non-zero abelian variety, and e a non-negative integer, then $A(F_{\text{sep}}(\sigma))$ has infinite rank for almost all $\sigma \in \text{Gal}(F)^e$.*

Proof. Let C_A be the F -curve associated to A by Theorem 5.3. Since F is Hilbertian, there exists a linearly disjoint sequence $F_1, F_2, \dots$ of extensions of F of fixed degree d such that $C_A(F_i) \setminus C_A(F) \neq \emptyset$ for all i , cf. [FJ08, Proof

of Theorem 18.6.1, Part A]. By Borel-Cantelli, the set of $\sigma \in \text{Gal}(F)^e$ for which $F_{\text{sep}}(\sigma)$ contains infinitely many of the F_i has measure one, cf. [FJ08, 18.5.3(b)]. For those σ we have that $|C_A(F_{\text{sep}}(\sigma))| = \infty$, and hence that the rank of $A(F_{\text{sep}}(\sigma))$ is infinite by Theorem 5.3. $\square$

We do not know if almost all $F_{\text{sep}}(\sigma)$ are ample if F is not countable, therefore we had to apply Theorem 5.3, and not Theorem 1.1 directly.

6.3. Fields of totally S -adic numbers. We give a simplified proof and a generalization of [FP10, Theorem 1.3]. If S is a finite set of local primes of a field F (see [FP10, p.2]), the field $F_{\text{tot},S}$ of **totally S -adic numbers** over F is defined as the maximal Galois extension of F contained in all completions $\hat{F}_{\mathfrak{p}}$, $\mathfrak{p} \in S$. For $\sigma \in \text{Gal}(F)^e$, let $F_{\text{sep}}[\sigma]$ be the maximal Galois extension of F in the fixed field $F_{\text{sep}}(\sigma)$.

Corollary 6.3. *Let F be a countable Hilbertian field, S a finite set of local primes of F , and e a non-negative integer. If A/F is a non-zero abelian variety, then $A(F_{\text{tot},S} \cap F_{\text{sep}}[\sigma])$ has infinite rank for almost all $\sigma \in \text{Gal}(F)^e$.*

Proof. Geyer and Jarden prove in [GJ02] that the field $F_{\text{tot},S} \cap F_{\text{sep}}[\sigma]$ is ample for almost all $\sigma \in \text{Gal}(F)^e$. Hence, Theorem 1.1 implies the claim. $\square$

This also generalizes [FJ74, Theorem 2.1], [GJ06, Theorem B], and a special case of [Pet06, Theorem A.1].

6.4. Larsen's conjecture. Our last application of Theorem 1.1 is a conditional result. Among the numerous conjectures about which fields are ample, the following conjecture (see e.g. [JK10], [BSF12, §4.2]) is particularly interesting for the questions considered in this work.

Conjecture 6.4 (Koenigsmann). *If F is an infinite field with finitely generated absolute Galois group, then F is ample.*

One of the consequences of Theorem 1.1 is that a proof of Conjecture 6.4 would immediately prove the following conjecture of Larsen, see [Lar03, Question 1], [Im06], [IL08], [IL19].

Conjecture 6.5 (Larsen). *Let F be an infinite finitely generated field, A/F a non-zero abelian variety, and e a non-negative integer. Then $A(F_{\text{sep}}(\sigma))$ has infinite rank for all $\sigma \in \text{Gal}(F)^e$.*

Corollary 6.6. *Conjecture 6.4 implies Conjecture 6.5.*

Proof. The absolute Galois group of $F_{\text{sep}}(\sigma)$ is generated by σ , and hence is finitely generated. Therefore, Theorem 1.1 implies the claim. $\square$

ACKNOWLEDGMENTS

We would like to thank Moshe Jarden for his constant encouragement regarding this paper. This work was supported by the European Commission under contract MRTN-CT-2006-035495, and the Minkowski Center for Geometry at

Tel Aviv University, established by the Minerva Foundation. S.P. was supported by the NCN grant OPUS16 UMO 2018/31/B/ST1/01474 of the Polish ministry of science.

REFERENCES

- [BBP14] Franck Benoist, Elisabeth Bouscaren, and Anand Pillay. Semiabelian varieties over separably closed fields, maximal divisible subgroups, and exact sequences. *J. Inst. Math. Jussieu*, 15(1):29–69, 2014.
- [BSF12] Lior Bary-Soroker and Arno Fehm. Open problems in the theory of ample fields. In J.-M. Couveignes D. Bertrand, Ph. Boalch and P. Dèbes, editors, *Geometric and differential Galois theory*, volume 27 of *Séminaires & Congrès*, pages 1–11, 2012.
- [Con06] Brian Conrad. Chow’s K/k -image and K/k -trace and the Lang-Néron theorem. *L’Enseignement Mathématique (2)*, 52:37–108, 2006.
- [Feh10] Arno Fehm. Subfields of ample fields. Rational maps and definability. *Journal of Algebra*, 323(5):1738–1744, 2010.
- [FJ74] Gerhard Frey and Moshe Jarden. Approximation theory and the rank of abelian varieties over large algebraic fields. *Proceedings of the London Mathematical Society*, 28:112–128, 1974.
- [FJ08] Michael D. Fried and Moshe Jarden. *Field arithmetic*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge 11. Berlin: Springer. xxii, 780 p., third edition, 2008.
- [FP10] Arno Fehm and Sebastian Petersen. On the rank of abelian varieties over ample fields. *International Journal of Number Theory*, 6(3):579–586, 2010.
- [FP11] Arno Fehm and Elad Paran. Galois theory over rings of arithmetic power series. *Advances in Mathematics*, 226:4183–4197, 2011.
- [GD65] Alexander Grothendieck and Jean Dieudonné. Éléments de géométrie algébrique IV. Étude locale des schémas et des morphismes des schémas, seconde partie. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (24):5–231, 1965.
- [GD66] Alexander Grothendieck and Jean Dieudonné. Éléments de géométrie algébrique IV. Étude locale des schémas et des morphismes des schémas, troisième partie. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (28):5–255, 1966.
- [GD67] Alexander Grothendieck and Jean Dieudonné. Éléments de géométrie algébrique IV. Étude locale des schémas et des morphismes des schémas, quatrième partie. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (32):5–361, 1967.
- [GJ02] Wulf-Dieter Geyer and Moshe Jarden. PSC Galois extensions of Hilbertian fields. *Mathematische Nachrichten*, 236:119–160, 2002.
- [GJ06] Wulf-Dieter Geyer and Moshe Jarden. The rank of abelian varieties over large algebraic fields. *Archiv der Mathematik*, 86(3):211–216, 2006.
- [GM06] Dragos Ghioca and Rahim Moosa. Division points on subvarieties of isotrivial semi-abelian varieties. *International Mathematics Research Notices*, pages 1–23, 2006.
- [Har09] David Harbater. On function fields with free absolute Galois groups. *Journal für die reine und angewandte Mathematik*, 632:85–103, 2009.
- [Hru96] Ehud Hrushovski. The Mordell-Lang conjecture for function fields. *Journal of the AMS*, 9(3):667–690, 1996.
- [IL08] Bo-Hae Im and Michael Larsen. Abelian varieties over cyclic fields. *American Journal of Mathematics*, 130(5):1195–1210, 2008.

- [IL13] Bo-Hae Im and Michael Larsen. Some applications of the hailes-jewett theorem to field arithmetic. *Israel Journal of Mathematics*, 198:35–47, 2013.
- [IL19] Bo-Hae Im and Michael Larsen. Abelian varieties and finitely generated galois groups. arXiv:1902.04011 [math.NT], 2019.
- [Im06] Bo-Hae Im. Mordell-Weil groups and the rank of elliptic curves over large fields. *Canadian Journal of Mathematics*, 58(4):796–819, 2006.
- [Jar03] Moshe Jarden. On ample fields. *Archiv der Mathematik*, 80:475–488, 2003.
- [JK10] Markus Junker and Jochen Koenigsmann. Schlanke Körper (Slim Fields). *Journal of Symbolic Logic*, 75(2):481–500, 2010.
- [Kim97] Minhyong Kim. Purely inseparable points on curves of higher genus. *Mathematical Research Letters*, 4:663–666, 1997.
- [Koi72] Shoji Koizumi. The fields of moduli for polarized abelian varieties and for curves. *Nagoya Math. J.*, 48:37–55, 1972.
- [Lar03] Michael Larsen. Rank of elliptic curves over almost separably closed fields. *Bulletin of the London Mathematical Society*, 35(6):817–820, 2003.
- [MFK92] David Mumford, John Fogarty, and Frances Kirwan. *Geometric Invariant Theory*. Springer, third edition, 1992.
- [Mil80] James Milne. *Etale Cohomology*. Princeton Mathematical Series 33. Princeton University Press, 1980.
- [Mil86] James Milne. Abelian Varieties. In G. Cornell and J. H. Silverman, editors, *Arithmetic Geometry, Proceedings of Storrs Conference*. Springer, 1986.
- [Pet06] Sebastian Petersen. On a question of Frey and Jarden about the rank of abelian varieties (with an appendix by Moshe Jarden). *Journal of Number Theory*, 120:287–302, 2006.
- [Pop96] Florian Pop. Embedding problems over large fields. *Annals of Mathematics*, 144(1):1–34, 1996.
- [Pop10] Florian Pop. Henselian implies large. *Annals of Mathematics*, 172(3):2183–2195, 2010.
- [Pop14] Florian Pop. Little survey on large fields. In Teissier Campillo, Kuhlmann, editor, *Valuation Theory in Interaction*, EMS Series of Congress Reports, pages 432–463, 2014.
- [Ros73] Michael Rosen. S-units and S-class group in algebraic function fields. *Journal of Algebra*, 26:98–108, 1973.
- [Rös19] Damian Rössler. Perfect points on curves. <http://people.maths.ox.ac.uk/rossler/mypage/pdf-file> 2019.
- [RW02] Michael Rosen and Siman Wong. The rank of abelian varieties over infinite Galois extensions. *Journal of Number Theory*, 92:182–196, 2002.
- [Sam66] Pierre Samuel. La conjecture de mordell pour les corps de fonctions. *Seminaire N. Bourbaki*, 287:149–167, 1966.