

LIEB-ROBINSON BOUNDS AND STRONGLY CONTINUOUS DYNAMICS FOR A CLASS OF MANY-BODY FERMION SYSTEMS IN \mathbb{R}^d .

MARTIN GEBERT, BRUNO NACHTERGAELE, JAKE RESCHKE, AND ROBERT SIMS

ABSTRACT. We introduce a class of UV-regularized two-body interactions for fermions in \mathbb{R}^d and prove a Lieb-Robinson estimate for the dynamics of this class of many-body systems. As a step toward this result, we also prove a propagation bound of Lieb-Robinson type for Schrödinger operators. We apply the propagation bound to prove the existence of infinite-volume dynamics as a strongly continuous group of automorphisms on the CAR algebra.

1. INTRODUCTION

The goal of this paper is to study propagation estimates for interacting fermion systems in \mathbb{R}^d , $d \geq 1$, and to apply them to construct the infinite-volume dynamics for a class of such systems as a strongly continuous one-parameter group of automorphisms of the standard CAR algebra. We introduce a class of short-range, UV-regularized two-body interactions for which this is possible. Without the use of a UV cut-off of some kind, such a result cannot be expected to hold. See, for example, the discussion in [BR97, Introduction to Section 6.3]. Nevertheless, as Sakai notes in the last paragraph of his book [Sak91], constructing the dynamics for interacting systems is one of the most important problems. To address this problem, a common approach is to consider the dynamics in representations of the algebra of observables associated with a class of sufficiently regular states. This is not our approach here. Instead we introduce a UV regularization of the interactions. This allow us to construct the infinite system dynamics as automorphisms of the CAR algebra of observables that depend continuously on time. A typical situation where it is advantageous to consider the dynamics on the observables algebra of the infinite system is in non-equilibrium statistical mechanics, where until now one would either use a quasi-free dynamics (as, e.g., in [FMU03]) or work in a lattice setting where UV regularization is provided by the lattice (see, e.g., [HM15, GR17, BBDRF18, BBDRF19], and [Rob68, Rue69] for the original and fundamental existence result for quantum spin systems.).

One broad class of models in which UV degrees of freedom are naturally absent are mean field models and related limiting regimes and the dynamics of such models have been studied including in infinite volume. For example, well-posedness for the Hartree equation in infinite volume, which describes the mean field limit [EESY04], has been proved by Lewin and Sabin in [LS15].

The regularization we adopt in this paper is smearing the interactions by Gaussians parameterized by $\sigma > 0$ in such a way that the pair interaction between point particles is recovered in the limit $\sigma \rightarrow 0$ (See Appendix A for a proof). Formally, in second quantization, this leads to a Hamiltonian of the form

$$(1.1) \quad H_\Lambda = \int_{\mathbb{R}^d} (\nabla a_x^+ \nabla a_x + V(x) a_x^+ a_x) dx + \frac{1}{2} \int_\Lambda \int_\Lambda W(x-y) a^+(\varphi_x^\sigma) a^+(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma) dx dy,$$

Date: June 27, 2022.

Based upon work supported by the National Science Foundation under Grant DMS-1813149.

where V is an external potential such as a smooth periodic function, and W is a short-range two-body interaction. We defer stating precise conditions on V and W until Section 2. The smearing is only needed in the interaction and one can take for φ_x^σ an L^1 -normalized Gaussian of width σ and centered at $x \in \mathbb{R}^d$. The parameter σ can be interpreted as the size of the particles and, as discussed in Appendix A, restricted to the N -particle Hilbert space, for any finite number of particles N , in either a finite or infinite volume, the dynamics converges to the standard Schrödinger dynamics generated by the self-adjoint Hamiltonian H_N given by

$$(1.2) \quad H_N = \sum_{k=1}^N (-\Delta_k + V(x_k)) + \sum_{1 \leq k < l \leq N} W(x_k - x_l).$$

Having a state-independent definition of the dynamics has both conceptual and practical advantages. From early on it was realized however that the subtle, non-robust, property of (thermodynamic) stability may be an obstacle to using perturbation series to define Heisenberg dynamics for infinite systems in the continuum [Dys52]. Therefore, it is not surprising that attempts were made to construct toy models of interacting theories for which stability could be proved. An early example is [Str68]. In [SW70] an infinite-volume dynamics for interacting fields is obtained using relativistic locality (Minkowski space). The only previous Euclidean construction of infinite-system dynamics on the CAR algebra over $L^2(\mathbb{R}^d)$ that explicitly considers a regularized pair interaction, as far as we are aware, is by Narnhofer and Thirring [NT90]. In that work the authors were motivated by the desire to preserve the Galilean invariance of the dynamics, which led them to employ a somewhat contrived UV regularization. The smearing of the form (1.1) used here is, we believe, more natural and likely to faithfully reproduce the low-energy physics.

Before summarizing our results, we point out that *defining* a dynamics on the CAR algebra over $L^2(\mathbb{R}^d)$ is by itself not the issue. Including pair interactions in a densely defined self-adjoint Hamiltonian on Fock space has been accomplished a long time ago. The corresponding one-parameter group of unitaries can be used to define a dynamics as a group of automorphisms on the bounded operators on Fock space, which includes the CAR algebra. This dynamics, however, is in general *not strongly continuous*. This is because the commutator of the unregularized interaction term with a creation or annihilation operator is unbounded.

Our proof of convergence of the thermodynamic limit of the infinite-volume dynamics hinges on a propagation estimate of Lieb-Robinson type [LR72] for systems in which the interaction is only active in a bounded volume Λ , with estimates that are uniform in Λ . Let $\tau_t^\Lambda(\cdot)$ denote the Heisenberg evolution with the interactions restricted to Λ (see (2.19) for the precise definition) and define the one-particle Schrödinger evolution in the usual way:

$$(1.3) \quad f_t = e^{-it(-\Delta+V)} f, \quad f \in L^2(\mathbb{R}^d), \quad t \in \mathbb{R}.$$

Lieb-Robinson Bound for Schrödinger operators. Let V be given as the Fourier transform of a finite Borel measure of compact support on \mathbb{R}^d . For $\sigma > 0$ and $x \in \mathbb{R}^d$, denote by φ_x^σ the L^1 normalized Gaussian on \mathbb{R}^d with mean x and variance σ . Then, there exist constants C_1, C_2, C_3 , such that for all $f \in L^2(\mathbb{R}^d)$ and $t \in \mathbb{R}$ one has

$$(1.4) \quad |\langle e^{-itH_1} f, \varphi_x^\sigma \rangle| \leq C_1 e^{C_2 |t| \ln |t|} \int_{\mathbb{R}^d} dy e^{-\frac{C_3}{t^2+1} |x-y|} |f(y)|.$$

A more detailed estimate and explicit constants are given in Proposition 3.2 and Corollary 3.3. For discrete Schrödinger operators on graphs a Lieb-Robinson type propagation estimate holds for any real-valued diagonal potential [AW12].

Let τ_t^Λ , be the Heisenberg dynamics generated by H_Λ in (1.1) for bounded $\Lambda \subset \mathbb{R}^d$, and $t \in \mathbb{R}$. We will prove the following result as Theorem 2.5.

Propagation Bound for many-body fermion dynamics. *Let $W \in L^\infty(\mathbb{R}^d)$ be real-valued and satisfying $W(-x) = W(x)$ and $|W(x)| \leq Ce^{-a|x|}$, for some $C, a > 0$. Then, there exist continuous functions $C(t), a(t) > 0$ such that for all bounded and measurable $\Lambda \subset \mathbb{R}^d$, and $f, g \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ one has the bound*

$$(1.5) \quad \|\{\tau_t^\Lambda(a(f)), a^\#(g)\}\| \leq \|f\|_1 \|g\|_1 e^{C(t)} e^{-a(t)d(\text{supp}(f), \text{supp}(g))},$$

where $d(\text{supp}(f), \text{supp}(g))$ denotes the distance between the essential supports of f and g , and $a^\#(\cdot)$ is either a creation or an annihilation operator.

Explicit forms of $C(t)$ and $a(t)$ are given in Theorem 2.5. This Lieb-Robinson type bound provides localization estimates for general elements in the CAR algebra by the usual algebraic relations in the same way as for lattice fermion systems as in [HK06, NSY18, BdSP17].

As an application of this propagation bound above, which is of independent interest, we then prove the existence and continuity of the infinite systems dynamics. See Theorem 2.7 for the precise statement. There are other approaches to proving the convergence of the dynamics in the thermodynamic limit. Using propagation bounds, however, yields a short and intuitive proof.

Strongly continuous infinite-volume dynamics. *There exists a strongly continuous one-parameter group of automorphism of the CAR algebra over $L^2(\mathbb{R}^d)$, $\{\tau_t\}_{t \in \mathbb{R}}$, such that*

$$(1.6) \quad \lim_{\Lambda \uparrow \mathbb{R}^d} \tau_t^\Lambda(a(f)) = \tau_t(a(f)), \text{ for all } f \in L^2(\mathbb{R}^d).$$

The strategy for proving existence of the thermodynamic limit of the Heisenberg dynamics using propagation bounds appears to be quite general and has been employed successfully for lattice systems [BR97, NSSSZ10, NSY19]. This method works whenever the interactions restricted to a bounded region are described by a bounded self-adjoint operator. It is worth noting that the free part of the dynamics does not require a cut-off for this result to hold. Due to its uniformity in Λ , the propagation bound (1.5) extends to the infinite system dynamics.

Several generalizations of the propagation bounds could be considered. For Schrödinger operators, we expect that the restrictions on V can be relaxed. The many-body bounds are derived here for regularized pair interactions only. Our approach can handle k -body terms with virtually no changes. A different type of extension of obvious interest would be to consider fermions in an external magnetic field. In contrast, constructing the many-body dynamics for boson systems one has to face an additional element of unboundedness that has long been understood to force one to consider a weaker topology to express the continuity in time [Ver11]. Already for boson lattice systems, such as oscillator lattices, Lieb-Robinson bounds can be derived but one finds bounds that are no longer in terms of the operator norm of the observables [ALBN10, NRSS09]. Such bounds can nevertheless still be used to prove the existence of infinite-systems dynamics [NSSSZ10]. Another approach to define the dynamics of infinite oscillator lattices was developed by Buchholz [Buc17], who constructs a strongly continuous dynamics on the Resolvent Algebra [BG08].

The existence of propagation bounds of Lieb-Robinson type and the strongly continuous infinite-volume dynamics for many-body systems with Hamiltonians of the form (1.1) provide a new avenue for applications. For example, if we choose for V a periodic potential, such that $-\Delta + V$ has a band structure with a gap, the non-interacting many-body ground state at suitable fermion density is gapped. We expect this gap to persist in the presence of interactions as in (1.1) with W sufficiently

small. Stability of the ground state gap has been proved for broad classes of lattice systems [BHM10, Kli10, BH11, SPGC11, MZ13, CMPGS13, DRS19, FP18, Has19, NSY20]. We believe that an analogous result for the continuum systems studied in this paper is now within reach.

2. MODEL AND STATEMENT OF MAIN RESULTS

Let $d \geq 1$ and take Δ to be the Laplace operator on \mathbb{R}^d . For any real-valued $V \in L^\infty(\mathbb{R}^d)$, we will denote by

$$(2.1) \quad H_1 = -\Delta + V$$

the corresponding (self-adjoint) Schrödinger operator with domain $H^2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d)$, see [RS75] for more details. As required, we will impose further conditions on V , e.g. see (2.11).

Our goal is to analyze a class of operators on the fermionic Fock space. We will follow closely the notation in [BR97], see specifically Section 5.2.1, and refer the reader there for more details. Let us denote by

$$(2.2) \quad \mathfrak{F}^- = \bigoplus_{n=0}^{\infty} (L^2(\mathbb{R}^d)^{\otimes n})^-$$

the anti-symmetric Fock space (Hilbert space) generated by $L^2(\mathbb{R}^d)$. In the above, $L^2(\mathbb{R}^d)^{\otimes n}$ is short for $\bigotimes_{k=1}^n L^2(\mathbb{R}^d)$ and $(\cdot)^-$ denotes anti-symmetrization. For each $f \in L^2(\mathbb{R}^d)$, take $a(f) \in \mathcal{B}(\mathfrak{F}^-)$, the bounded linear operators over \mathfrak{F}^- , to be the annihilation operator corresponding to f , and denote by $a^*(f)$, its adjoint, the corresponding creation operator. It is well-known that these creation and annihilation operators satisfy the canonical anti-commutation relations (CAR)

$$(2.3) \quad \{a(f), a(g)\} = 0 \quad \text{and} \quad \{a(f), a^*(g)\} = \langle f, g \rangle \mathbb{1} \quad \text{for all } f, g \in L^2(\mathbb{R}^d)$$

where $\{A, B\} = AB + BA$ denotes the anti-commutator, $\langle \cdot, \cdot \rangle$ the scalar product in $L^2(\mathbb{R}^d)$, and $\mathbb{1}$ is the identity acting on \mathfrak{F}^- . In addition, one has that

$$(2.4) \quad \|a^*(f)\| = \|a(f)\| = \|f\|_2 \quad \text{for all } f \in L^2(\mathbb{R}^d)$$

where here, and in the following, $\|\cdot\|_p$ will refer to the L^p -norm for $p \in [1, \infty]$ and $\|\cdot\|$ will denote the operator norm.

The models we will consider are defined in terms of a particular class of annihilation and creation operators. Let $\sigma > 0$, take $x \in \mathbb{R}^d$, and consider the Gaussian $\varphi_x^\sigma : \mathbb{R}^d \rightarrow \mathbb{R}$ with

$$(2.5) \quad \varphi_x^\sigma(y) = \frac{1}{(2\pi\sigma^2)^{d/2}} e^{-\frac{|y-x|^2}{2\sigma^2}} \quad \text{for all } y \in \mathbb{R}^d.$$

We say that φ_x^σ is centered at $x \in \mathbb{R}^d$ with variance σ^2 . We have chosen an L^1 -normalization, i.e. $\|\varphi_x^\sigma\|_1 = 1$ for all $x \in \mathbb{R}^d$. Given (2.4), it is clear that for any $x \in \mathbb{R}^d$,

$$(2.6) \quad \|a^*(\varphi_x^\sigma)\|^2 = \|a(\varphi_x^\sigma)\|^2 = \|\varphi_x^\sigma\|_2^2 = (4\pi\sigma^2)^{-d/2} =: C_\sigma$$

where we have introduced the notation $C_\sigma > 0$ as this quantity will enter our estimates frequently.

For any bounded and measurable set $\Lambda \subset \mathbb{R}^d$, we will analyze the operator

$$(2.7) \quad H_\Lambda^\sigma = d\Gamma(H_1) + W_\Lambda^\sigma$$

acting on \mathfrak{F}^- , where $d\Gamma(H_1)$ denotes the second quantization of H_1 , again see [BR97] for the definition, and the interaction W_Λ^σ is given by

$$(2.8) \quad W_\Lambda^\sigma = \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy W_\Lambda(x, y) a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma)$$

where $W_\Lambda : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ has the form $W_\Lambda(x, y) = \chi_{\Lambda \times \Lambda}(x, y)W(x - y)$ for some real-valued $W \in L^\infty(\mathbb{R}^d)$ and $\chi_{\Lambda \times \Lambda}(x, y)$ denotes the indicator function of the set $\Lambda \times \Lambda$. In this case, for each fixed $\sigma > 0$ and any $\Lambda \subset \mathbb{R}^d$ that is bounded and measurable, we have that

$$(2.9) \quad \|W_\Lambda^\sigma\| \leq \frac{1}{2}C_\sigma^2 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy |W_\Lambda(x, y)| \leq \frac{1}{2} \left(\frac{1}{4\pi\sigma^2} \right)^d \|W\|_\infty |\Lambda|^2$$

where $|\Lambda|$ denotes the Lebesgue measure of Λ . Thus for bounded W , $W_\Lambda^\sigma \in \mathcal{B}(\mathfrak{F}^-)$ for each choice of $\sigma > 0$ and $\Lambda \subset \mathbb{R}^d$ as above. We conclude that, in these cases, H_Λ^σ is a well-defined self-adjoint operator on the anti-symmetric Fock space \mathfrak{F}^- .

As we progress, more assumptions will need to be made on W . For ease of later reference, we state them here.

Assumption 2.1 (On W). *Let $W : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy*

- (i) $W \in L^\infty(\mathbb{R}^d)$ is real-valued;
- (ii) W is symmetric, i.e. $W(-x) = W(x)$ for almost every $x \in \mathbb{R}^d$;
- (iii) W is short-range, i.e. there are positive numbers a and c_W for which

$$(2.10) \quad |W(x)| \leq c_W e^{-a|x|} \quad \text{for almost every } x \in \mathbb{R}^d.$$

2.1. Bounds on the propagator of one-particle Schrödinger operators. In this section, we derive propagation bounds for one-particle Schrödinger operators with the form H_1 as in (2.1). To make a precise statement, we require the following from the potential V .

Assumption 2.2 (On V). *Let $V : \mathbb{R}^d \rightarrow \mathbb{C}$ have the form*

$$(2.11) \quad V(x) = \int_{\mathbb{R}^d} d\mu(k) e^{-ik \cdot x}$$

where $\mu : \text{Borel}(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a Borel measure on \mathbb{R}^d satisfying:

- (i) μ has support contained in a ball, i.e. there is some $M > 0$ and $\text{supp } \mu \subset B_M(0)$;
- (ii) $\mu = \mu^+ - \mu^-$ where μ^+ and μ^- are non-negative finite measures on $\text{Borel}(\mathbb{R}^d)$, i.e. $\mu^\pm(\mathbb{R}^d) < \infty$. We set $|\mu| = \mu^+ + \mu^-$;
- (iii) μ is even, i.e. $\mu(A) = \mu(-A)$ for all $A \in \text{Borel}(\mathbb{R}^d)$.

Under these assumptions, V is the Fourier transform of a signed, compactly supported, finite measure μ , which is real-valued and bounded. Two parameters that will appear in estimates are C_μ and M , to characterize V . They need not be chosen optimally but should satisfy

$$(2.12) \quad \int_{\mathbb{R}^d} d|\mu|(x) \leq C_\mu, \quad \sup\{|k| \mid k \in \text{supp}(|\mu|)\} \leq M.$$

Two classes of examples of potentials V satisfying Assumption 2.2 are the following.

- (i) Let V satisfy Assumption 2.2 and suppose the corresponding measure μ has a density $f \in L^1(\mathbb{R}^d)$. Then, these assumptions imply that f has compact support, that $f(x) = f(-x)$, and that

$$(2.13) \quad V(x) = \int_{\mathbb{R}^d} dk f(k) e^{-ik \cdot x}.$$

For example, our class of potentials V includes $V(x) = \text{sinc}^k(x)$ for all $k \in \mathbb{N}$ for which the density is the k -fold convolution of the indicator function $f(y) = 1_{[-1,1]}(y)$.

(ii) Let V satisfy Assumption 2.2 and suppose the corresponding measure μ satisfies: There is some $N \in \mathbb{N}$, points $\{a_n\}_{n=1}^N$ in \mathbb{R}^d , and numbers $\{b_n\}_{n=1}^N$ in \mathbb{R} for which

$$(2.14) \quad \mu(A) = \frac{1}{2} \sum_{n=1}^N b_n (\delta_{a_n}(A) + \delta_{-a_n}(A)) \quad \text{for any } A \in \text{Borel}(\mathbb{R}^d).$$

Here $\delta_{(\cdot)}$ denotes the Dirac measure. This form gives rise to potentials V with

$$(2.15) \quad V(x) = \sum_{n=1}^N b_n \cos(a_n \cdot x).$$

The main result of this section is:

Theorem 2.3 (Lieb-Robinson bound for Schrödinger operators). *Let V satisfy Assumption 2.2 and consider the Schrödinger operator $H_1 = -\Delta + V$ as defined in (2.1). Then there exist constants $C_1, C_2, C_3 > 0$ depending on d, μ , and σ such that the estimate*

$$(2.16) \quad |\langle e^{-itH_1} f, \varphi_x^\sigma \rangle| \leq C_1 e^{C_2 |t| \ln |t|} \int_{\mathbb{R}^d} dy e^{-\frac{C_3}{t^2+1} |x-y|} |f(y)|$$

holds for all $t \in \mathbb{R}$ and $f \in L^2(\mathbb{R}^d)$.

The constants C_1, C_2 , and C_3 are derived in the proof of Corollary 3.3.

Remarks 2.4. (i) *Theorem 2.3 relies, in general, on the smoothness of the class of test functions φ_y^σ that we used to probe the locality properties of the dynamics. For example, we can see from an explicit computation in the case of $V = 0$, that the exponential decay does generally not hold when the Gaussians is replaced by a non-smooth functions such as, for example a characteristic function. Using the formula*

$$(2.17) \quad (e^{-it(-\Delta)} \psi)(x) = \frac{1}{(4\pi it)^{d/2}} \int_{\mathbb{R}^d} dy e^{\frac{i|x-y|^2}{4t}} \psi(y),$$

valid for $t \neq 0$ and general $\psi \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$, a straightforward calculation, in the case $d = 1$, shows that the leading behavior of $|(e^{-it(-\Delta)} \chi_{[-1,1]})(x)|$, for $t > 0$ and $|x|$ large is given by

$$|(e^{-it(-\Delta)} \chi_{[-1,1]})(x)| \sim 2\sqrt{\frac{t}{\pi}} \frac{x}{x^2 - 1} \sin \frac{x}{2t}.$$

(ii) *In the case $V = 0$ and for the Gaussians φ_y^σ another explicit computation (see e.g. [Tes09, Sec. 7.3]), shows that*

$$(2.18) \quad |(e^{-it(-\Delta)} \varphi_y^\sigma)(x)| = \frac{1}{(2\pi)^{d/2}} \frac{e^{-\frac{\sigma^2 |x-y|^2}{8t^2 + 2\sigma^4}}}{(4t^2 + \sigma^4)^{d/4}}.$$

Using this form one immediately sees that an estimate analogous to (1.4) holds with a Gaussian distance dependence of the kernel.

(iii) *In view of the first two remarks it is clear that Theorem 2.3 is far from optimal. It is also likely that similar bounds hold for a broader class of potentials V .*

2.2. Lieb-Robinson bounds and the thermodynamic limit. Our next results concern Lieb-Robinson bounds for the Heisenberg dynamics associated to the operator H_Λ^σ defined in (2.7). We begin by recalling the notion of dynamics on Fock space.

As before, let $\mathcal{B}(\mathfrak{F}^-)$ denote the bounded linear operators on \mathfrak{F}^- . For each $\sigma > 0$ and any bounded, measurable $\Lambda \subset \mathbb{R}^d$, we define the Heisenberg dynamics associated to H_Λ^σ for each $t \in \mathbb{R}$ as the map $\tau_t^\Lambda : \mathcal{B}(\mathfrak{F}^-) \rightarrow \mathcal{B}(\mathfrak{F}^-)$ defined by

$$(2.19) \quad \tau_t^\Lambda(A) = e^{itH_\Lambda^\sigma} A e^{-itH_\Lambda^\sigma} \quad \text{for all } A \in \mathcal{B}(\mathfrak{F}^-).$$

We note that although $\tau_t^\Lambda(A)$ depends on σ , we have suppressed this in our notation.

We will analyze this dynamics on the CAR algebra generated by the set $\{a(f), a^*(f) : f \in L^2(\mathbb{R}^d)\}$. Again, we refer to [BR97, Sec 5.2] for more details. In particular, we will focus our attention to operators $A = a(f) \in \mathcal{B}(\mathfrak{F}^-)$ for some $f \in L^2(\mathbb{R}^d)$.

Let us also recall the free dynamics, i.e. the case where $W=0$ and there is no interaction. We will denote this well-studied, free dynamics of the CAR algebra by

$$(2.20) \quad \tau_t^\emptyset(a(f)) = e^{itd\Gamma(H_1)} a(f) e^{-itd\Gamma(H_1)}, \quad f \in L^2(\mathbb{R}^d), \quad t \in \mathbb{R}.$$

A straight forward calculation shows

$$(2.21) \quad \tau_t^\emptyset(a(f)) = a(f_t) \quad \text{where} \quad f_t = e^{-itH_1} f.$$

Our goal is to examine the behavior of τ_t^Λ as Λ tends to \mathbb{R}^d , and in particular, we wish to establish the existence of a dynamics in this thermodynamic limit. To do so, we regard τ_t^Λ as a perturbation of the infinite volume free dynamics τ_t^\emptyset on the finite volume Λ . In this case, the key to constructing the thermodynamic limit is an appropriate form of the Lieb-Robinson bound. To express the Lieb-Robinson bound for this model, we find it convenient to introduce the non-negative function

$$(2.22) \quad F_t^\Lambda(f, g) = \|\{\tau_t^\Lambda(a(f)), a^*(g)\} - \{\tau_t^\emptyset(a(f)), a^*(g)\}\| + \|\{\tau_t^\Lambda(a(f)), a(g)\}\|$$

Iteration is at the heart of most Lieb-Robinson bounds, and in the present context, our proof will show that the function F_t^Λ above iterates more simply than either term on the right-hand-side of (2.22). In any case, we find the following Lieb-Robinson bound.

Theorem 2.5 (Lieb-Robinson bound). *Fix $\sigma > 0$. Let V satisfy Assumption 2.2, W satisfy Assumption 2.1, and for each $t \in \mathbb{R}$ and any bounded, measurable set $\Lambda \subset \mathbb{R}^d$, denote by τ_t^Λ the dynamics associated to H_Λ^σ as defined in (2.19). For any $f, g \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$, the bound*

$$(2.23) \quad F_t^\Lambda(f, g) \leq D(t)(e^{P_3(t)} - 1) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy e^{-\frac{c_t|x-y|}{4}} |f(x)| |g(y)|$$

holds for functions $D(t) \sim e^{c|t|\ln|t|}$, P_3 a polynomial of degree $6d+1$ in $|t|$, and $c_t \sim \frac{1}{1+t^2}$. Explicit values for these functions are given in Section 4, see specifically Lemma 4.3 and Lemma 4.4.

Remark 2.6. *Here it is crucial to subtract the free time evolution*

$$(2.24) \quad \{\tau_t^\emptyset(a^*(f)), a(g)\} = \langle f_t, g \rangle \mathbb{1}$$

since $|\langle f_t, g \rangle|$ does not, in general, decay exponentially; see Remark 2.4 (i).

Our main application concerns the existence of a dynamics in the thermodynamic limit. In Section 5 we show how the following theorem is a consequence of the Λ -independent bounds proven in Theorem 2.5.

Theorem 2.7. *Under the assumptions of Theorem 2.5 there exists a strongly continuous one-parameter group of automorphisms of the CAR algebra over $L^2(\mathbb{R}^d)$, $\{\tau_t\}_{t \in \mathbb{R}}$, such that for all $f \in L^2(\mathbb{R}^d)$ and any increasing sequence (Λ_n) of bounded subsets of \mathbb{R}^d such that $\cup_n \Lambda_n = \mathbb{R}^d$,*

$$(2.25) \quad \lim_{n \rightarrow \infty} \tau_t^{\Lambda_n}(a(f)) = \tau_t(a(f))$$

in the operator norm topology, with convergence uniform in t in compact subsets of \mathbb{R} .

3. LIEB-ROBINSON BOUND FOR SCHRÖDINGER OPERATORS. PROOF OF THEOREM 2.3

In this section we use the notation $H_0 = -\Delta$ and $H_1 = -\Delta + V$. To prove Theorem 2.3 we will use a Dyson series expansion for e^{itH_1} :

$$(3.1) \quad e^{-itH_1} = e^{-itH_0} + (-i)^n \sum_{n=1}^{\infty} \int_0^t dt_n \cdots \int_0^{t_2} dt_1 e^{-i(t-t_n)H_0} V e^{-i(t_n-t_{n-1})H_0} V \cdots V e^{it_1 H_0}.$$

Since V is bounded (by Assumption 2.2), this series is absolutely convergent in norm. We are interested in estimating $|(e^{-itH_1} \varphi_y^\sigma)(x)|$, where φ_y^σ is the Gaussian function given in (2.5). Using the Fourier representation of V (2.11), the integrand of the n -th term in the expansion (3.1) applied to φ_y^σ can be expressed as follows:

$$(3.2) \quad \begin{aligned} & (e^{-i(t-t_n)H_0} V e^{-i(t_n-t_{n-1})H_0} V \cdots V e^{it_1 H_0} \varphi_y^\sigma)(x) \\ &= \int_{\mathbb{R}^d} d\mu(k_n) \cdots \int_{\mathbb{R}^d} d\mu(k_1) A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x). \end{aligned}$$

where

$$(3.3) \quad A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x) = (e^{-i(t-t_n)H_0} V_{k_n} e^{-i(t_n-t_{n-1})H_0} V_{k_{n-1}} \cdots V_{k_1} e^{it_1 H_0} \varphi_y^\sigma)(x).$$

Here V_k is the multiplication operator by $V_k(x) = e^{-ik \cdot x}$ and $|\mu| = \mu^+ + \mu^-$.

Lemma 3.1. *Let $n \in \mathbb{N}$. For all $k_1, \dots, k_n, y, x \in \mathbb{R}^d$, $t \geq t_n \geq \dots \geq t_1 \geq 0$ one has*

$$(3.4) \quad |A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x)| = \frac{1}{(2\pi)^{d/2}} \frac{e^{-\frac{\sigma^2}{8t^2+2\sigma^4} |(x-y)-2\sum_{l=0}^n (t_{l+1}-t_l) \sum_{j=1}^l k_j|^2}}{(4t^2 + \sigma^4)^{d/4}},$$

where we use the conventions $t_{n+1} = t$, $t_0 = 0$ and $\sum_{j=1}^0 = 0$. Furthermore,

$$(3.5) \quad |(e^{-itH_0} \varphi_y^\sigma)(x)| = \frac{1}{(2\pi)^{d/2}} \frac{e^{-\frac{\sigma^2 |x-y|^2}{8t^2+2\sigma^4}}}{(4t^2 + \sigma^4)^{d/4}}.$$

Proof. Let \mathcal{F} be the unitary Fourier transform on \mathbb{R}^d and \mathcal{F}^* be its inverse then we obtain

$$(3.6) \quad A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x) = (\mathcal{F}^* \mathcal{F} e^{-i(t-t_n)H_0} \mathcal{F}^* \mathcal{F} V_{k_n} \mathcal{F}^* \cdots \mathcal{F} V_{k_1} \mathcal{F}^* \mathcal{F} e^{it_1 H_0} \mathcal{F}^* \mathcal{F} \varphi_y^\sigma)(x).$$

Now, for all $t \in \mathbb{R}$, $k \in \mathbb{R}^d$ and $\psi \in L^2(\mathbb{R}^d)$ we have $\mathcal{F} e^{-itH_0} \mathcal{F}^* \psi = e^{-i(\cdot)^2 t} \psi$ and $\mathcal{F} V_k \mathcal{F}^* \psi = \psi(\cdot - k)$. Therefore,

$$A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} dk e^{ikx} \prod_{l=0}^n e^{-i(t_{l+1}-t_l) (k - \sum_{j=l+1}^n k_j)^2} (\mathcal{F} \varphi_y^\sigma)(k - \sum_{j=1}^n k_j),$$

where we use the convention that $\sum_{j=n+1}^n = 0$. Next we use that

$$(\mathcal{F} \varphi_y^\sigma)(k) = \frac{1}{(2\pi)^{d/2}} e^{-ik \cdot y} e^{-\frac{\sigma^2 |k|^2}{2}}$$

and perform a change of variables to obtain

$$(3.7) \quad A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x) = \frac{e^{i \sum_{j=1}^n k_j \cdot (x-y)}}{(2\pi)^d} \int_{\mathbb{R}^d} dk e^{ik \cdot (x-y)} \prod_{l=0}^n e^{-i(t_{l+1}-t_l) |k + \sum_{j=1}^l k_j|^2} e^{-\frac{\sigma^2 |k|^2}{2}},$$

where we use the convention $\sum_{j=1}^0 = 0$. Multiplying out $|k + \sum_{j=1}^l k_j|^2 = |k|^2 + 2k \cdot \sum_{j=1}^l k_j + |\sum_{j=1}^l k_j|^2$, using $\sum_{l=0}^n (t_{l+1} - t_l) = t_{n+1} - t_0 = t$ and taking the absolute value give

$$|A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x)| = \frac{1}{(2\pi)^d} \left| \int_{\mathbb{R}^d} dk e^{ik \cdot (x-y)} e^{-i|k|^2 t} e^{-2i \sum_{l=0}^n (t_{l+1}-t_l) \sum_{j=1}^l k \cdot k_j} e^{-\frac{\sigma^2 |k|^2}{2}} \right|.$$

Now, a calculation shows that for any $c > 0$ and $a, b \in \mathbb{R}$

$$(3.8) \quad \frac{1}{(2\pi)^{1/2}} \left| \int_{\mathbb{R}} dk e^{ikb} e^{-k^2(c+ia)} \right| = \frac{e^{-\frac{cb^2}{4(a^2+c^2)}}}{(4(a^2+c^2))^{1/4}}.$$

Hence,

$$(3.9) \quad |A(t_1, \dots, t_n, t, k_1, \dots, k_n, y, x)| = \frac{1}{(2\pi)^{d/2}} \frac{1}{(4t^2 + \sigma^4)^{d/4}} e^{-\frac{\sigma^2}{8t^2+2\sigma^4} |(x-y)-2 \sum_{l=0}^n (t_{l+1}-t_l) \sum_{j=1}^l k_j|^2},$$

which proves (3.4). Identity (3.5) follows from an explicit calculation using the integral kernel of e^{-itH_0} , see e.g. [Tes09, Sec. 7.3]. \square

With this lemma we have arrived at the following estimate. For $t \in \mathbb{R}$ and $x, y \in \mathbb{R}^d$, we have

$$(3.10) \quad |(e^{-itH_1} - e^{-itH_0})\varphi_y^\sigma(x)| \leq \frac{1}{(2\pi)^{d/2}} \frac{1}{(4t^2 + \sigma^4)^{d/4}} \sum_{n=1}^{\infty} \int_{\mathbb{R}^d} d|\mu|(k_1) \cdots \int_{\mathbb{R}^d} d|\mu|(k_n) \\ \times \int_0^{|t|} dt_n \cdots \int_0^{t_2} dt_1 e^{-\frac{\sigma^2}{8t_1^2+2\sigma^4} |(x-y)-2 \sum_{l=0}^n (t_{l+1}-t_l) \sum_{j=1}^l k_j|^2},$$

where, as before, we use the convention $\sum_{j=1}^0 = 0$. By estimating the RHS of this estimate, we will obtain the following proposition. Recall the definitions of C_μ and M in (2.12).

Proposition 3.2. *For all $t \in \mathbb{R}$ and $x, y \in \mathbb{R}^d$, we have*

$$|(e^{-itH_1} - e^{-itH_0})\varphi_y^\sigma(x)| \leq \frac{1}{(2\pi)^{d/2}} \frac{1}{(4t^2 + \sigma^4)^{d/4}} \left(e^{-\frac{\sigma^2 |x-y|^2}{32t^2+8\sigma^4}} (e^{C_\mu |t|} - 1) + \frac{1}{\sqrt{2\pi}} e^{-\frac{|x-y|}{4|t|M} (\ln \frac{|x-y|}{4MC_\mu t^2} - 1)} e^{C_\mu |t|} \right).$$

Using (3.5) to bound $|e^{-itH_0}\varphi_y^\sigma(x)|$, it is then straightforward to obtain an estimate for $|e^{-itH_1}\varphi_y^\sigma(x)|$. In our application, however, the following simplified estimate is easier to use.

Corollary 3.3. *There exist constants C_1, C_2 , and C_3 , such that for all $t \in \mathbb{R}$ and $x, y \in \mathbb{R}^d$, we have*

$$(3.11) \quad |e^{-itH_1}\varphi_y^\sigma(x)| \leq C_1 e^{C_2 |t| \ln |t| - C_3 \frac{|x-y|}{1+t^2}}.$$

Proof of Proposition 3.2. Note that, as expected from translation invariance, the RHS of (3.10) depends only on $x - y$. Therefore, w.l.o.g., we can assume $y = 0$. Let I_n , $n \geq 1$, denote the n -th term of the sum in the RHS of (3.10):

$$(3.12) \quad I_n = \int_{\mathbb{R}^d} d|\mu|(k_1) \cdots \int_{\mathbb{R}^d} d|\mu|(k_n) \int_0^{|t|} dt_n \cdots \int_0^{t_2} dt_1 e^{-\frac{\sigma^2}{8t_1^2+2\sigma^4} |(x-y)-2 \sum_{l=0}^n (t_{l+1}-t_l) \sum_{j=1}^l k_j|^2}.$$

Given $x \in \mathbb{R}^d$, $t \in \mathbb{R}$, we split the sum over n in (3.10) as follows:

$$(3.13) \quad B_0 = \sum_{n=1}^{N_0} I_n, \quad B_1 = \sum_{n=N_0+1}^{\infty} I_n, \quad \text{with } N_0 = \left\lfloor \frac{|x|}{(4|t|M)} \right\rfloor,$$

where $[a]$ denotes the integer part of a and M is as in (2.12).

First, if $\frac{|x|}{(4|t|M)} \geq 1$, the first sum is non-empty and we estimate its terms as follows. Let $n \leq |x|/(4|t|M)$, and note that for $k_j \in B_M(0)$ with $1 \leq j \leq l \leq n$ and $|t| = t_{n+1} \geq t_n \geq \dots \geq t_0 = 0$

$$(3.14) \quad \left| \sum_{l=0}^n (t_{l+1} - t_l) \sum_{j=1}^l k_j \right| \leq Mn \sum_{l=0}^n (t_{l+1} - t_l) = Mn|t|.$$

Therefore,

$$(3.15) \quad \left| x - 2 \sum_{l=0}^n (t_{l+1} - t_l) \sum_{j=1}^l k_j \right|^2 \geq \frac{1}{4} |x|^2.$$

and therefore

$$(3.16) \quad B_0 \leq e^{-\frac{\sigma^2|x|^2}{32t^2+8\sigma^4}} \sum_{n=1}^{N_0} \frac{C_\mu^n |t|^n}{n!} \leq e^{-\frac{\sigma^2|x|^2}{32t^2+8\sigma^4}} (e^{C_\mu|t|} - 1)$$

To estimate the terms in B_1 , note that the integrand in (3.12) is bounded by 1. Therefore, using C_μ defined in (2.12),

$$(3.17) \quad B_1 \leq \sum_{n=N_0+1}^{\infty} \frac{(C_\mu|t|)^n}{n!} \leq \frac{(C_\mu|t|)^{N_0+1}}{(N_0+1)!} e^{C_\mu|t|}.$$

Stirling's formula yields for all $m \geq 1$ the bound

$$(3.18) \quad \frac{1}{m!} \leq \frac{1}{\sqrt{2\pi}} e^{m-m \ln m}.$$

Using this and $N_0 + 1 \geq |x|/(4|t|M)$, we obtain

$$(3.19) \quad B_1 \leq \frac{1}{\sqrt{2\pi}} e^{-\frac{|x|}{4|t|M} (\ln \frac{|x|}{4MC_\mu t^2} - 1)} e^{C_\mu|t|}.$$

If $\frac{|x|}{(4|t|M)} < 1$, $B_0 = 0$ and we estimate B_1 as in (3.19) with $N_0 = 0$.

The proposition is proved by combining the estimates (3.16) and (3.19). \square

Proof of Corollary 3.3. We may again restrict ourselves to the case $y = 0$. Proposition 3.2 and (3.5) together immediately give the estimate

$$\begin{aligned} |e^{-itH_1} \varphi_0^\sigma(x)| &\leq \frac{1}{(2\pi)^{d/2}} \frac{1}{(4t^2 + \sigma^4)^{d/4}} \\ &\times \left[e^{-\frac{\sigma^2|x|^2}{8t^2+2\sigma^4}} + e^{-\frac{\sigma^2|x|^2}{32t^2+8\sigma^4}} (e^{C_\mu|t|} - 1) + \frac{1}{\sqrt{2\pi}} e^{-\frac{|x|^2}{4M|t|} (\ln \frac{|x|}{4MC_\mu t^2} - 1)} e^{C_\mu|t|} \right]. \end{aligned}$$

The last term in the square bracket is the estimate (3.19) for B_1 , which we can simplify by considering two cases for $(x, t) \in \mathbb{R}^{d+1}$, namely $\frac{|x|}{4MC_\mu t^2} \geq e^2$, and $\frac{|x|}{4MC_\mu t^2} < e^2$. In the first case, we have

$$(3.20) \quad B_1 \leq \frac{e^{-\frac{|x|}{4|t|M} + C_\mu|t|}}{\sqrt{2\pi}}.$$

On the other hand, if $\frac{|x|}{4MC_\mu t^2} < e^2$, we use the inequality $e^{-u \ln u} \leq e^{\frac{1}{e}}$ for all $u > 0$, to obtain

$$(3.21) \quad B_1 \leq \frac{1}{\sqrt{2\pi}} e^{\frac{1}{e}} e^{e^2 C_\mu |t| (\ln C_\mu |t| + 1) + C_\mu |t|} e^{-\frac{|x|}{4MC_\mu t^2} + e^2}.$$

By bounding B_1 by the sum of the RHSs of (3.20) and (3.21) and making a few more easy simplifications we arrive at the following bound:

$$\begin{aligned} |e^{-itH_1} \varphi_0^\sigma(x)| &\leq \frac{1}{(2\pi\sigma^2)^{d/2}} \\ &\times \left[e^{-\frac{\sigma^2|x|^2}{32t^2+8\sigma^4}} e^{C_\mu|t|} + \frac{1}{\sqrt{2\pi}} e^{-\frac{|x|}{4|t|M} + C_\mu|t|} + \frac{1}{\sqrt{2\pi}} e^{\frac{1}{e}} e^{e^2 C_\mu |t| (\ln C_\mu |t| + 1) + C_\mu |t|} e^{-\frac{|x|}{4MC_\mu t^2} + e^2} \right]. \end{aligned}$$

To estimate the Gaussian decay of the first term between the square brackets by a simple exponential, we use that for all $u \in \mathbb{R}$, $u^2 \geq u - 1/4$. Furthermore, since $u \ln u \geq u - 1$, we also have

$$e^{C_\mu|t|} \leq e^{1+C_\mu|t| \ln C_\mu|t|}.$$

Using this and replacing the constant prefactors by their maximum, we find

$$|e^{-itH_1} \varphi_0^\sigma(x)| \leq \frac{1}{(2\pi\sigma^2)^{d/2}} e^{\frac{1}{8\sigma^2} + \frac{1}{e} + e^2} e^{e^2 C_\mu |t| (\ln C_\mu |t| + 2)} \left(e^{-\frac{\sigma^2|x|}{32t^2+8\sigma^4}} + e^{-\frac{|x|}{4tM}} + e^{-\frac{|x|}{4MC_\mu t^2}} \right).$$

Finally, we use the estimate

$$\begin{aligned} \min \left\{ \frac{\sigma^2}{32t^2 + 8\sigma^4}, \frac{1}{4tM}, \frac{1}{4MC_\mu t^2} \right\} &\geq \frac{1}{t^2 \left(\frac{32}{\sigma^2} + 4MC_\mu \right) + 4tM + 8\sigma^2} \\ &\geq \frac{1}{t^2 \left(\frac{32}{\sigma^2} + 4MC_\mu + 2 \right) + 2M^2 + 8\sigma^2} \\ (3.22) \quad &\geq \frac{1}{\frac{32}{\sigma^2} + 4MC_\mu + 2(M^2 + 1) + 8\sigma^2} \frac{1}{t^2 + 1} \end{aligned}$$

to bound the sum of three exponentials by

$$3e^{-C_3 \frac{|x|}{t^2+1}}, \text{ with } C_3 = \frac{1}{\frac{32}{\sigma^2} + 4MC_\mu + 2(M^2 + 1) + 8\sigma^2}.$$

It is now straightforward to find suitable values for C_1 and C_2 for which the bound given in the corollary holds. \square

4. MANY-BODY LIEB-ROBINSON BOUND. PROOF OF THEOREM 2.5

The main goal of this section is to prove Theorem 2.5. We do so in three steps. First, in Section 4.1 we establish a basic estimate which facilitates an iteration scheme; this is the content of Lemma 4.1. Next, in Section 4.2, we estimate a kernel function which, among other things, ultimately justifies the convergence of our iteration, see Lemma 4.2 and Lemma 4.3. Finally, in Section 4.3 we perform the iteration and verify the bound claimed in Theorem 2.5.

4.1. A Preliminary Bound. In this section, we provide an estimate on the basic quantity of interest in Theorem 2.5. Let us briefly recall the set-up. We have fixed $\sigma > 0$, taken V and W satisfying Assumption 2.2 and Assumption 2.1 respectively, and introduced, see (2.22), the non-negative function

$$(4.1) \quad F_t^\Lambda(f, g) = \|\{\tau_t^\Lambda(a(f)), a^*(g)\} - \{\tau_t^\emptyset(a(f)), a^*(g)\}\| + \|\{\tau_t^\Lambda(a^*(f)), a^*(g)\}\|$$

for any bounded, measurable set $\Lambda \subset \mathbb{R}^d$, $t \in \mathbb{R}$, and functions $f, g \in L^2(\mathbb{R}^d)$. Our first estimate is as follows.

Lemma 4.1. *Under the assumptions described above, for any $t \geq 0$, we find that*

$$(4.2) \quad \begin{aligned} F_t^\Lambda(f, g) &\leq C_\sigma \int_0^t ds \int_{\mathbb{R}^d} dx K_{t-s}(f, x) |\langle e^{-isH_1} \varphi_x^\sigma, g \rangle| \\ &\quad + C_\sigma \int_0^t ds \int_{\mathbb{R}^d} dx K_{t-s}(f, x) F_s^\Lambda(\varphi_x^\sigma, g) \end{aligned}$$

where $C_\sigma > 0$ is as in (2.6) and with kernel function $K_t(f, x)$ given by

$$(4.3) \quad K_t(f, x) = \|W\|_1 |\langle e^{-itH_1} f, \varphi_x^\sigma \rangle| + 2 \left(|W| * |\langle e^{-itH_1} f, \varphi_{(\cdot)}^\sigma \rangle| \right)(x).$$

Proof. We begin by recalling a useful perturbation formula. Fix a bounded, measurable set $\Lambda \subset \mathbb{R}^d$ and take $\sigma > 0$, $t \geq 0$, and $f \in L^2(\mathbb{R}^d)$. In this case,

$$(4.4) \quad \tau_t^\Lambda(a(f)) = \tau_t^\emptyset(a(f)) + i \int_0^t ds \tau_s^\Lambda \left(\left[W_\Lambda^\sigma, \tau_{t-s}^\emptyset(a(f)) \right] \right)$$

a proof of which can be found in [BR97, Prop. 5.4.1]. Note that

$$(4.5) \quad \begin{aligned} \left[W_\Lambda^\sigma, \tau_{t-s}^\emptyset(a(f)) \right] &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy W_\Lambda(x, y) \left[a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma), a(f_{t-s}) \right] \\ &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy W_\Lambda(x, y) \left[a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma), a(f_{t-s}) \right] a(\varphi_y^\sigma) a(\varphi_x^\sigma) \end{aligned}$$

where we have set $f_t = e^{-itH_1} f$ and used (2.3). Further calculating, we find

$$(4.6) \quad \begin{aligned} \left[a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma), a(f_{t-s}) \right] &= a^*(\varphi_x^\sigma) \{a^*(\varphi_y^\sigma), a(f_{t-s})\} - \{a^*(\varphi_x^\sigma), a(f_{t-s})\} a^*(\varphi_y^\sigma) \\ &= \langle f_{t-s}, \varphi_y^\sigma \rangle a^*(\varphi_x^\sigma) - \langle f_{t-s}, \varphi_x^\sigma \rangle a^*(\varphi_y^\sigma). \end{aligned}$$

Using now the symmetry of W , we obtain

$$\begin{aligned} &\{\tau_t^\Lambda(a(f)), a^*(g)\} - \{\tau_t^\emptyset(a(f)), a^*(g)\} \\ &= i \int_0^t ds \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy W_\Lambda(x, y) \langle f_{t-s}, \varphi_y^\sigma \rangle \{ \tau_s^\Lambda(a^*(\varphi_x^\sigma)) \tau_s^\Lambda(a(\varphi_y^\sigma)) \tau_s^\Lambda(a(\varphi_x^\sigma)), a^*(g) \}. \end{aligned}$$

With the anti-commutator relation

$$(4.7) \quad \{ABC, D\} = \{A, D\}BC - A\{B, D\}C + AB\{C, D\},$$

the norm bound

$$(4.8) \quad \begin{aligned} &\left\| \{\tau_t^\Lambda(a(f)), a^*(g)\} - \{\tau_t^\emptyset(a(f)), a^*(g)\} \right\| \\ &\leq C_\sigma \int_0^t ds \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy |W(x-y)| |\langle f_{t-s}, \varphi_y^\sigma \rangle| \\ &\quad \times \left(\left\| \{\tau_s^\Lambda(a^*(\varphi_x^\sigma)), a^*(g)\} \right\| + \left\| \{\tau_s^\Lambda(a(\varphi_y^\sigma)), a^*(g)\} \right\| + \left\| \{\tau_s^\Lambda(a(\varphi_x^\sigma)), a^*(g)\} \right\| \right) \end{aligned}$$

readily follows. Here we have used the bound $|W_\Lambda(x, y)| \leq |W(x - y)|$ for all $x, y \in \mathbb{R}^d$. Similar arguments yield the bound

$$\begin{aligned}
& \|\{\tau_t^\Lambda(a^*(f)), a^*(g)\}\| \\
& \leq C_\sigma \int_0^t ds \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy |W(x - y)| |\langle f_{t-s}, \varphi_y^\sigma \rangle| \\
(4.9) \quad & \times (\|\{\tau_s^\Lambda(a^*(\varphi_x^\sigma)), a^*(g)\}\| + \|\{\tau_s^\Lambda(a^*(\varphi_y^\sigma)), a^*(g)\}\| + \|\{\tau_s^\Lambda(a(\varphi_x^\sigma)), a^*(g)\}\|).
\end{aligned}$$

Our goal, as in most Lieb-Robinson bounds, is to derive bounds which can be iterated. Since neither (4.8) nor (4.9) iterate separately, we bound their sum, i.e. the function $F_t^\Lambda(f, g)$ introduced in (4.1) above. Recalling that $\{\tau_t^\emptyset(a(f)), a^*(g)\} = \langle f_t, g \rangle \mathbb{1}$, we find

$$\begin{aligned}
F_t^\Lambda(f, g) & \leq C_\sigma \int_0^t ds \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy |W(x - y)| |\langle f_{t-s}, \varphi_y^\sigma \rangle| \\
& \quad \times (2F_s^\Lambda(\varphi_x^\sigma, g) + 2|\langle e^{-isH_1} \varphi_x^\sigma, g \rangle| + F_s^\Lambda(\varphi_y^\sigma, g) + |\langle e^{-isH_1} \varphi_y^\sigma, g \rangle|) \\
(4.10) \quad & = C_\sigma \int_0^t ds \int_{\mathbb{R}^d} dx K_{t-s}(f, x) (F_s^\Lambda(\varphi_x^\sigma, g) + |\langle e^{-isH_1} \varphi_x^\sigma, g \rangle|),
\end{aligned}$$

where we have introduced $K_t(f, x)$ as in (4.3). This is the claim in (4.2). \square

Looking at the bound proven in Lemma 4.1, in particular (4.2), it is natural to begin an iteration. To ensure convergence, we first provide an estimate on the kernel function $K_t(f, x)$.

4.2. Estimating the Kernel. In this section, we provide two useful apriori estimates on the kernel function $K_t(f, x)$ defined in (4.3) of Lemma 4.1. First, we estimate this function for general $f \in L^2(\mathbb{R}^d)$, this result is stated in Lemma 4.2 below. In Lemma 4.3, we provide a similar estimate in the special case that f is a Gaussian.

Before we prove our first estimate, it will be convenient to introduce the following notation. For any $r > 0$, set $G_r : \mathbb{R}^d \rightarrow \mathbb{R}$ to be

$$(4.11) \quad G_r(x) = e^{-r|x|} \quad \text{for all } x \in \mathbb{R}^d.$$

Lemma 4.2. *Let $t \geq 0$, $f \in L^2(\mathbb{R}^d)$, and $x \in \mathbb{R}^d$. We have*

$$(4.12) \quad K_t(f, x) \leq P_1(t) e^{C_2 t |\ln t|} (G_{b_t} * |f|)(x)$$

where G_{b_t} is as in (4.11) with

$$(4.13) \quad 4b_t = \frac{aC_3}{a(t^2 + 1) + C_3}$$

and $P_1 : \mathbb{R} \rightarrow (0, \infty)$ is the polynomial of degree 2d given by

$$(4.14) \quad P_1(t) = C_1(2c_W D_3^2 (4b_t)^{-d} + \|W\|_1).$$

Here $C_1, C_2, C_3 > 0$ are as in the proof of Theorem 2.3, a and c_W are as in (2.10), and $D_3 > 0$ is as in Lemma B.1.

Proof. We first note that

$$(4.15) \quad K_0(f, x) \leq \|W\|_1 (\varphi_0^\sigma * |f|)(x) + 2(|W| * \varphi_0^\sigma * |f|)(x)$$

which may be further estimated as in (4.12).

Now, for $t > 0$, we recall the bound in (1.4):

$$(4.16) \quad |\langle e^{-itH_1} f, \varphi_x^\sigma \rangle| \leq C_1 e^{C_2 t |\ln t|} \int_{\mathbb{R}^d} dy e^{-\frac{C_3}{t^2+1}|x-y|} |f(y)| = C_1 e^{C_2 t |\ln t|} (G_{a_t} * |f|)(x)$$

where we use the notation in (4.11) and set

$$(4.17) \quad a_t = \frac{C_3}{t^2 + 1}$$

In this case, the bound

$$(4.18) \quad \begin{aligned} K_t(f, x) &= \|W\|_1 |\langle e^{-itH_1} f, \varphi_x^\sigma \rangle| + 2(|W| * |\langle e^{-itH_1} f, \varphi_{(\cdot)}^\sigma \rangle|)(x) \\ &\leq C_1 e^{C_2 t |\ln t|} \|W\|_1 (G_{a_t} * |f|)(x) + 2C_1 e^{C_2 t |\ln t|} (|W| * (G_{a_t} * |f|))(x) \end{aligned}$$

is clear. Moreover, the exponential decay of W , see (2.10) in Assumption 2.1, implies

$$(4.19) \quad \begin{aligned} (|W| * (G_{a_t} * |f|))(x) &\leq c_W \int_{\mathbb{R}^d} dz e^{-a|x-z|} (G_{a_t} * |f|)(z) \\ &\leq c_W \int_{\mathbb{R}^d} dz \int_{\mathbb{R}^d} dy e^{-4b_t|x-z|} e^{-4b_t|z-y|} |f(y)| \end{aligned}$$

where we have used (4.13), and in particular, that $4b_t \leq \min\{a, a_t\}$. By Lemma B.1 there is $D_3 > 0$, depending only on d , such that

$$(4.20) \quad \int_{\mathbb{R}^d} dz e^{-4b_t|x-z|} e^{-4b_t|z-y|} \leq \frac{D_3^2}{(4b_t)^d} e^{-b_t|x-y|}.$$

We conclude that

$$(4.21) \quad (|W| * (G_{a_t} * |f|))(x) \leq c_W \frac{D_3^2}{(4b_t)^d} (G_{b_t} * |f|)(x)$$

and note that (4.12) follows from the point-wise bound $G_{a_t}(x) \leq G_{b_t}(x)$. \square

We now turn to the special case of a Gaussian.

Lemma 4.3. *Let $t \geq 0$, $\sigma > 0$, and $x, y \in \mathbb{R}^d$. We have*

$$(4.22) \quad K_t(\varphi_x^\sigma, y) \leq P_2(t) e^{C_2 t |\ln t|} G_{c_t}(x - y)$$

where G_{c_t} is as in (4.11) with

$$(4.23) \quad c_t = \frac{aC_3}{16(a(t^2 + 1) + C_3(1 + a\sigma^2))}$$

and $P_2 : \mathbb{R} \rightarrow \mathbb{R}$ is the polynomial of degree $4d$ given by

$$(4.24) \quad P_2(t) = \frac{e^{\frac{1}{8\sigma^2}}}{(2\pi\sigma^2)^{d/2}} P_1(t) \frac{D_3^2}{(4c_t)^d}.$$

Here we use freely the notation established in Lemma 4.2.

Proof. Applying Lemma 4.2 and the simple bound $u^2 \geq u - 1/4$, for all $u \in \mathbb{R}$, we find that

$$(4.25) \quad \begin{aligned} K_t(\varphi_x^\sigma, y) &\leq P_1(t) e^{C_2 t |\ln t|} (G_{b_t} * \varphi_x^\sigma)(y) \\ &= \frac{P_1(t) e^{\frac{1}{8\sigma^2}}}{(2\pi\sigma^2)^{d/2}} e^{C_2 t |\ln t|} \int_{\mathbb{R}^d} dz e^{-b_t|y-z|} e^{-\frac{|x-z|}{2\sigma^2}}. \end{aligned}$$

Recalling also Lemma B.1, the bound

$$(4.26) \quad \int_{\mathbb{R}^d} dz e^{-b_t|y-z|} e^{-\frac{|x-z|}{2\sigma^2}} \leq D_3^2 \left(\frac{1}{4c_t} \right)^d e^{-c_t|x-y|},$$

follows from the estimate

$$(4.27) \quad 4c_t \leq \min \left\{ b_t, \frac{1}{2\sigma^2} \right\}.$$

This proves the assertion. \square

4.3. Iterating the Bound. In this section, we will iterate the bound proven in Lemma 4.1, i.e. (4.2), and complete the proof of Theorem 2.5.

We first note that iteration of (4.2) produces, for any $N \in \mathbb{N}$, a bound of the form

$$(4.28) \quad F_t^\Lambda(f, g) \leq \sum_{n=1}^N a_n(t, f, g) + R_N(t, f, g)$$

where for each $1 \leq n \leq N$ the terms

$$(4.29) \quad \begin{aligned} a_n(t, f, g) = & C_\sigma^n \int_0^t dt_1 \int_{\mathbb{R}^d} dx_1 K_{t-t_1}(f, x_1) \int_0^{t_1} dt_2 \int_{\mathbb{R}^d} dx_2 K_{t_1-t_2}(\varphi_{x_1}^\sigma, x_2) \cdots \\ & \times \int_0^{t_{n-1}} dt_n \int_{\mathbb{R}^d} dx_n K_{t_{n-1}-t_n}(\varphi_{x_{n-1}}^\sigma, x_n) |\langle e^{-it_n H_1} \varphi_{x_n}^\sigma, g \rangle| \end{aligned}$$

and similarly, the remainder is given by

$$(4.30) \quad \begin{aligned} R_N(t, f, g) = & C_\sigma^N \int_0^t dt_1 \int_{\mathbb{R}^d} dx_1 K_{t-t_1}(f, x_1) \int_0^{t_1} dt_2 \int_{\mathbb{R}^d} dx_2 K_{t_1-t_2}(\varphi_{x_1}^\sigma, x_2) \cdots \\ & \times \int_0^{t_{N-1}} dt_N \int_{\mathbb{R}^d} dx_N K_{t_{N-1}-t_N}(\varphi_{x_{N-1}}^\sigma, x_N) F_{t_N}^\Lambda(\varphi_{x_N}^\sigma, g) \end{aligned}$$

Next, we estimate the terms $a_n(t, f, g)$.

Lemma 4.4. *Let $t > 0$ and $n \in \mathbb{N}$. We find the following bound*

$$(4.31) \quad a_n(t, f, g) \leq D(t) \frac{P_3(t)^n}{n!} \int_{\mathbb{R}^d} dx \int_{\mathbb{R}^d} dy e^{-\frac{c_t|x-y|}{4}} |f(x)| |g(y)|$$

where $P_3 : \mathbb{R} \rightarrow \mathbb{R}$ is a polynomial in t of degree $6d + 1$ with

$$(4.32) \quad P_3(t) = \frac{C_\sigma e^{C_2} D_3 t P_2(t)}{c_t^d} \quad \text{and} \quad D(t) = C_1 e^{C_2} D_3 \frac{P_1(t)}{P_2(t)} e^{C_2 t |\ln t|}.$$

All quantities appearing above are as in Lemma 4.2 and Lemma 4.3.

Proof. Fix $t > 0$. For our estimate, it will be convenient to recall some bounds from Section 4.2. First, let $f \in L^2(\mathbb{R}^d)$ and $x \in \mathbb{R}^d$. Lemma 4.2 shows that for $0 \leq t_1 \leq t$,

$$(4.33) \quad \begin{aligned} K_{t-t_1}(f, x) & \leq P_1(t-t_1) e^{C_2(t-t_1)|\ln(t-t_1)|} (G_{b_{t-t_1}} * |f|)(x) \\ & \leq P_1(t) e^{C_2(t-t_1)|\ln(t-t_1)|} (G_{c_t} * |f|)(x) \end{aligned}$$

where we have used that P_1 is increasing in t and that $b_{t-t_1} \geq b_t \geq c_t$. Next, an application of Lemma 4.3 shows that for any $0 \leq t_j \leq t_{j-1} \leq t$ and all $x, y \in \mathbb{R}^d$,

$$(4.34) \quad \begin{aligned} K_{t_{j-1}-t_j}(\varphi_x^\sigma, y) & \leq P_2(t_{j-1}-t_j) e^{C_2(t_{j-1}-t_j)|\ln(t_{j-1}-t_j)|} G_{c_{t_{j-1}-t_j}}(x-y) \\ & \leq P_2(t) e^{C_2(t_{j-1}-t_j)|\ln(t_{j-1}-t_j)|} G_{c_t}(x-y) \end{aligned}$$

Here we used that the polynomial $P_2(t)$ is increasing in t and the function c_t is decreasing in t . Similarly, arguing as in (4.16), we see that

$$(4.35) \quad |\langle e^{-it_n H_1} \varphi_{x_n}^\sigma, g \rangle| \leq C_1 e^{C_2 t_n |\ln t_n|} (G_{a_{t_n}} * |g|)(x_n) \leq C_1 e^{C_2 t_n |\ln t_n|} (G_{c_t} * |g|)(x_n)$$

for any $0 \leq t_n \leq t$ and $x_n \in \mathbb{R}^d$. As a final observation, note that for parameters $0 = t_{n+1} \leq t_n \leq \dots \leq t_j \leq t_{j-1} \leq \dots \leq t_1 \leq t_0 = t$, we have the bound

$$(4.36) \quad \begin{aligned} \sum_{j=1}^{n+1} (t_{j-1} - t_j) |\ln(t_{j-1} - t_j)| &\leq \sum_{j=1}^{n+1} (t_{j-1} - t_j) \ln(t_{j-1} - t_j) 1\{t_{j-1} - t_j \geq 1\} + n + 1 \\ &\leq \ln t 1\{t \geq 1\} t + n + 1 \\ &\leq t |\ln t| + n + 1, \end{aligned}$$

where $1\{\cdot\}$ stands for the indicator function, we used that $x |\ln x| \leq 1$ for all $x \in (0, 1]$, and in the argument of the \ln , we use the bound $t_{j-1} - t_j \leq t$, for $j = 1, \dots, n+1$, to produce a telescopic sum.

Putting all this together we find that

$$(4.37) \quad \begin{aligned} a_n(t, f, g) &\leq C_1 C_\sigma^n P_1(t) P_2(t)^{n-1} \int_0^t dt_1 \dots \int_0^{t_{n-1}} dt_n e^{C_2 \sum_{j=1}^{n+1} (t_{j-1} - t_j) |\ln(t_{j-1} - t_j)|} \\ &\quad \times \int_{\mathbb{R}^d} dx_1 \dots \int_{\mathbb{R}^d} dx_n (G_{c_t} * |f|)(x_1) G_{c_t}(x_1 - x_2) \dots G_{c_t}(x_{n-1} - x_n) (G_{c_t} * |g|)(x_n) \\ &\leq C_1 C_\sigma^n P_1(t) P_2(t)^{n-1} e^{C_2 t |\ln t|} e^{(n+1)C_2} \frac{t^n}{n!} \\ &\quad \times \int_{\mathbb{R}^d} dx_1 (G_{c_t} * |f|)(x_1) (G_{c_t} * \dots * G_{c_t} * |g|)(x_1) \end{aligned}$$

The latter integral can be further estimated as

$$(4.38) \quad \begin{aligned} \int_{\mathbb{R}^d} dx (G_{c_t} * |f|)(x) (G_{c_t} * \dots * G_{c_t} * |g|)(x) &= \int_{\mathbb{R}^d} dx |f(x)| (G_{c_t} * \dots * G_{c_t} * |g|)(x) \\ &\leq c_t^d \left(\frac{D_3}{c_t^d} \right)^{n+1} \int_{\mathbb{R}^d} dx \int_{\mathbb{R}^d} dy e^{-\frac{c_t |x-y|}{4}} |f(x)| |g(y)|. \end{aligned}$$

by using the point-wise estimate in Lemma B.1 on the $n+1$ -fold convolution of G_{c_t} with itself. This is the bound claimed in (4.31). \square

We can now complete the proof of Theorem 2.5.

Proof of Theorem 2.5. Given (4.28) and the estimate in Lemma 4.4, i.e. (4.31), it is clear that we need only show that the remainder term $R_N(t)$, see (4.30) goes to zero as $N \rightarrow \infty$. We will see that this is the case uniformly for t in compact sets.

Fix $T > 0$ and let $t \in [-T, T]$. We argue as in the proof of Lemma 4.4. The only difference between the term $a_N(t, f, g)$ and $R_N(t)$ is the final factor in the integrand: more precisely, $|\langle e^{-it_N H_1} \varphi_{x_N}^\sigma, g \rangle|$ is replaced with $F_{t_N}^\Lambda(\varphi_{x_N}^\sigma, g)$. In this case, the naive bound

$$(4.39) \quad F_{t_N}^\Lambda(\varphi_{x_N}^\sigma, g) \leq 6 \|\varphi_{x_N}^\sigma\|_2 \|g\|_2 = 6 \sqrt{C_\sigma} \|g\|_2$$

will suffice. In fact,

$$\begin{aligned}
R_N(t) &\leq 6\sqrt{C_\sigma}\|g\|_2 C_\sigma^N P_1(t) P_2(t)^{N-1} \int_0^t dt_1 \cdots \int_0^{t_{N-1}} dt_N e^{C_2 \sum_{j=1}^N (t_{j-1} - t_j) |\ln(t_{j-1} - t_j)|} \\
&\quad \times \int_{\mathbb{R}^d} dx_1 \cdots \int_{\mathbb{R}^d} dx_N (G_{c_t} * |f|)(x_1) G_{c_t}(x_1 - x_2) \cdots G_{c_t}(x_{N-1} - x_N) \\
&\leq 6\sqrt{C_\sigma}\|g\|_2 C_\sigma^N P_1(t) P_2(t)^{N-1} e^{C_2 t |\ln t|} e^{NC_2} \frac{t^N}{N!} \\
(4.40) \quad &\quad \times \int_{\mathbb{R}^d} dx_N (G_{c_t} * \cdots * G_{c_t} * |f|)(x_N).
\end{aligned}$$

Now, another application of Lemma B.1 demonstrates that

$$\begin{aligned}
\int_{\mathbb{R}^d} dx (G_{c_t} * \cdots * G_{c_t} * |f|)(x) &\leq \int_{\mathbb{R}^d} dx \int_{\mathbb{R}^d} dy (G_{c_t} * \cdots * G_{c_t})(x - y) |f(y)| \\
&\leq \left(\frac{D_3}{c_t^d}\right)^N c_t^d \int_{\mathbb{R}^d} dx \int_{\mathbb{R}^d} dy e^{-\frac{c_t |x-y|}{4}} |f(y)| \\
(4.41) \quad &= \left(\frac{D_3}{c_t^d}\right)^N c_t^d \|G_{c_t/4}\|_1 \|f\|_1
\end{aligned}$$

where here we have used that $f \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$. Since all the quantities in these estimates are explicit, it is clear that

$$(4.42) \quad \lim_{N \rightarrow 0} \sup_{t \in [-T, T]} R_N(t) = 0$$

which completes the proof of Theorem 2.5. \square

5. THE INFINITE-SYSTEM DYNAMICS. PROOF OF THEOREM 2.7

Our proof of Theorem 2.7 will make essential use of the following direct consequences of the propagation bounds of Theorem 2.5.

Let V and W satisfy Assumptions 2.2 and 2.1, respectively. Then, there exist continuous functions $\tilde{C}(\cdot), \tilde{a}(\cdot) > 0$, such that with φ_x^σ the Gaussians introduced in (2.5), and $f \in L^2(\mathbb{R}^d)$ of compact support, denoted by $\text{supp}(f)$, we have

$$(5.1) \quad \|\{\tau_t^\Lambda(a(f)), a^\#(\varphi_z^\sigma)\}\| \leq \|f\|_1 e^{\tilde{C}(|t|)} e^{-\tilde{a}(|t|)d(\text{supp}(f), z)}$$

where $a^\#(\cdot)$ refers to either an annihilation or creation operator, compare with (1.5). This estimate follows from Theorem 2.5 and Theorem 2.3 where we again use arguments as in (4.25) and (4.26).

Apart from this key estimate, the proof below uses a combination of several ideas introduced in [BR97, BHV06, NOS06, NSSSZ10].

Proof of Theorem 2.7. We first prove (2.25) for $f \in L^2(\mathbb{R}^d)$ of compact support, say $\text{supp} f \subset X \subset \mathbb{R}^d$, for some compact X . Let $(\Lambda_n)_{n \geq 1}$ be an increasing sequence of bounded, measurable sets such that $\bigcup_n \Lambda_n = \mathbb{R}^d$. To show that $(\tau_t^{\Lambda_n}(a(f)))_{n \geq 1}$ is Cauchy, uniformly in $t \in [-T, T]$ for $T > 0$, note that, for any $\Lambda_m \subseteq \Lambda_n$, the operator

$$W_{\Lambda_n}^\sigma - W_{\Lambda_m}^\sigma = \int_{\Lambda_n \times \Lambda_n \setminus \Lambda_m \times \Lambda_m} dx dy W(x - y) a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma)$$

is bounded by our assumptions. Hence, the generator of the strongly continuous dynamics $\tau_t^{\Lambda_n}$ is a bounded perturbation of the generator of $\tau_t^{\Lambda_m}$. In this case, we can apply the same perturbation formula as in (4.4) to compare the two dynamics. The following identity then holds

$$(5.2) \quad \begin{aligned} & \tau_t^{\Lambda_n}(a(f)) - \tau_t^{\Lambda_m}(a(f)) \\ &= \frac{i}{2} \int_0^t ds \int_{\Lambda_n \times \Lambda_n \setminus \Lambda_m \times \Lambda_m} dx dy W(x-y) \tau_t^{\Lambda_n} \left(\left[a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right] \right). \end{aligned}$$

Note the identity

$$(5.3) \quad \begin{aligned} & \left[a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right] = a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) \left\{ a(\varphi_x^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right\} \\ & \quad - a^*(\varphi_x^\sigma) a^*(\varphi_y^\sigma) \left\{ a(\varphi_y^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right\} a(\varphi_x^\sigma) \\ & \quad + a^*(\varphi_x^\sigma) \left\{ a^*(\varphi_y^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right\} a(\varphi_y^\sigma) a(\varphi_x^\sigma) \\ & \quad - \left\{ a^*(\varphi_x^\sigma), \tau_{t-s}^{\Lambda_m}(a(f)) \right\} a^*(\varphi_y^\sigma) a(\varphi_y^\sigma) a(\varphi_x^\sigma). \end{aligned}$$

Bounding (5.2) in norm, applying (5.1), and using the symmetry of W , we then find for any $T > 0$ constants C and $b > 0$, such that for every $t \in [-T, T]$ and every $n > m$,

$$(5.4) \quad \|\tau_t^{\Lambda_n}(a(f)) - \tau_t^{\Lambda_m}(a(f))\| \leq C \|\varphi_0^\sigma\|_2^3 \|f\|_1 \int_{\Lambda_n \times \Lambda_n \setminus \Lambda_m \times \Lambda_m} dx dy |W(x-y)| e^{-bd(X,x)},$$

which converges to 0 as $n > m \rightarrow \infty$ since $|W(x-y)| e^{-bd(X,x)} \in L^1(\mathbb{R}^{2d})$. This shows that for compactly supported f , the sequence $(\tau_t^{\Lambda_n}(a(f)))_{n \geq 1}$ is Cauchy (in norm) uniformly for $t \in [-T, T]$. Thus, the limit exists and gives rise to an isometry from \mathcal{P}_c , the set algebraically generated by $\{a(f), a^*(f) : f \in L^2(\mathbb{R}^d) \text{ of compact support}\}$, into $\mathcal{A}(L^2(\mathbb{R}^d))$. Equation (5.2) can be applied to see this limit is independent of the sequence (Λ_n) . As \mathcal{P}_c is dense in $\mathcal{A}(L^2(\mathbb{R}^d))$, this isometry extends uniquely to a homomorphism, τ_t , of $\mathcal{A}(L^2(\mathbb{R}^d))$. It is straightforward to verify that $\tau_t \circ \tau_s = \tau_{t+s}$ and, in particular, that τ_{-t} is the inverse of τ_t . Hence, $\{\tau_t \mid t \in \mathbb{R}\}$ is a one-parameter group of automorphisms of the CAR algebra.

To prove the strong continuity in t , it suffices to note that, for $f \in L^2(\mathbb{R}^d)$ and of compact support, the continuity of $t \mapsto \tau_t^{\Lambda_n}(a(f)) - \tau_t^\emptyset(a(f))$ carries over to the limiting function $t \mapsto \tau_t(a(f)) - \tau_t^\emptyset(a(f))$ due to the uniform convergence on compact intervals. Then, since τ_t^\emptyset is already known to be strongly continuous, $\tau_t(a(f))$ must be too. Finally, an $\varepsilon/3$ argument shows that the strong continuity extends to the full CAR algebra. \square

APPENDIX A. CONVERGENCE OF THE $\sigma \rightarrow 0$ LIMIT

We prove that, for any fixed finite number of fermions, the UV-regularized dynamics converges to the standard one as σ tends to 0. For this we consider arbitrary, not necessarily bounded, measurable $\Lambda \subset \mathbb{R}^d$. When Λ is not bounded, the interaction operator W_Λ^σ is generally unbounded. Therefore, we start by providing a more careful definition of H_Λ^σ . Note that (2.8) defines a bounded operator $W_{\Lambda;n}^\sigma$ on the n -particle subspace $(L^2(\mathbb{R}^d)^{\otimes n})^-$ for each n . Define the operator $W_{\Lambda;n}^0$ on $(L^2(\mathbb{R}^d)^{\otimes n})^-$ to be multiplication by the function $\sum_{1 \leq k < l \leq n} W_\Lambda(x_k - x_l)$. For $\sigma \geq 0$, define

$$(A.1) \quad H_{1;n} + W_{\Lambda;n}^\sigma = \sum_{k=1}^n (-\Delta_k + V(x_k)) + W_{\Lambda;n}^\sigma$$

acting on $(L^2(\mathbb{R}^d)^{\otimes n})^-$. For real-valued $V, W \in L^\infty(\mathbb{R}^d)$ this operator is self-adjoint on the domain $\mathcal{D}(H_{1;n}) = (H^2(\mathbb{R}^d)^{\otimes n})^-$. With $\sigma \geq 0$, let H_Λ^σ be the operator acting on Fock space by

$$H_\Lambda^\sigma = \bigoplus_{n=0}^{\infty} (H_{1;n} + W_{\Lambda;n}^\sigma)$$

with domain

$$\mathcal{D}(H_\Lambda^\sigma) = \{(\psi_n) \in \mathfrak{F}^- : \psi_n \in \mathcal{D}(H_{1;n}) \text{ and } \sum_{n=0}^{\infty} \|(H_{1;n} + W_{\Lambda;n}^\sigma)\psi_n\|_2^2 < \infty\}.$$

This operator is well-known to be self-adjoint, see e.g. [Wei80, Exercise 5.43].

Theorem A.1. *For real-valued $V, W \in L^\infty(\mathbb{R}^d)$, take H_1 as in (2.1). Then, for any measurable set $\Lambda \subset \mathbb{R}^d$*

$$H_\Lambda^\sigma \rightarrow H_\Lambda^0$$

in the strong resolvent sense as $\sigma \downarrow 0$.

Using a slight modification of [RS80, Thm VIII.20(a)], the above theorem readily implies:

Corollary A.2. *For $t \in \mathbb{R}$ we denote by $U_\Lambda^\sigma(t) = e^{-itH_\Lambda^\sigma}$ and $U_\Lambda^0(t) = e^{-itH_\Lambda^0}$ the unitary groups generated by H_Λ^σ and H_Λ^0 , respectively. Then*

$$\lim_{\sigma \downarrow 0} U_\Lambda^\sigma(t)\psi = U_\Lambda^0(t)\psi$$

for each $\psi \in \mathfrak{F}^-$, uniformly for t in compact subsets of \mathbb{R} .

Remark A.3. *Theorem A.1 and Corollary A.2 apply more generally to any self-adjoint operator H_1 .*

Proof of Theorem A.1. We start by reducing the proof Theorem A.1 to what is essentially the 2-particle situation. Recall that for $\sigma \geq 0$,

$$H_\Lambda^\sigma = \bigoplus_{n=0}^{\infty} (H_{1;n} + W_{\Lambda;n}^\sigma).$$

For each n , let $\mathcal{D}_n^- = (\mathcal{S}(\mathbb{R}^d)^{\otimes n})^-$ be the antisymmetrized n -fold tensor product of the Schwarz space $\mathcal{S}(\mathbb{R}^d)$, where $\mathcal{S}(\mathbb{R}^d)^{\otimes n} = \bigotimes_{j=1}^n \mathcal{S}(\mathbb{R}^d)$ is the set of finite linear combinations of functions ψ of the form $\psi(x_1, \dots, x_n) = \psi_1(x_1) \cdots \psi_n(x_n)$, with each $\psi_j \in \mathcal{S}(\mathbb{R}^d)$. Since $\mathcal{S}(\mathbb{R}^d)$ is a core for H_1 , \mathcal{D}_n^- is a core for $H_{1;n}$ by the Corollary to [RS80, Thm VIII.33]. It follows that \mathcal{D}_n^- is a core for $H_{1;n} + W_{\Lambda;n}^\sigma$ for every $\sigma \geq 0$. If

$$(A.2) \quad \mathcal{D}^- = \{(\psi_n) \in \mathfrak{F}^- : \psi_n \in \mathcal{D}_n^- \text{ and } \exists N \text{ with } \psi_n = 0 \forall n \geq N\},$$

then it is not difficult to see that \mathcal{D}^- is a core for H_Λ^σ for every $\sigma \geq 0$ (this is essentially Example 2 in [RS80, Sec VIII]).

As is well-known, strong resolvent convergence follows from strong convergence on a common core, see e.g. [RS80, Thm VIII.25(a)], and thus to prove Theorem A.1, it suffices to establish that

$$(A.3) \quad \lim_{\sigma \downarrow 0} H_\Lambda^\sigma \psi = H_\Lambda^0 \psi$$

for every $\psi \in \mathcal{D}^-$. Given the form of \mathcal{D}^- , see (A.2), it is clear that (A.3) will follow if

$$(A.4) \quad \lim_{\sigma \downarrow 0} W_{\Lambda;n}^\sigma \psi = W_{\Lambda;n}^0 \psi \text{ for every } \psi \in \mathcal{D}_n^-.$$

We need only prove (A.4). To this end, let $b^*(\cdot)$ and $b(\cdot)$ denote the creation and annihilation operators on the full Fock space $\mathfrak{F} = \bigoplus_{n=0}^{\infty} L^2(\mathbb{R}^d)^{\otimes n}$; for any $f \in L^2(\mathbb{R}^d)$ and $\psi \in L^2(\mathbb{R}^d)^{\otimes n}$,

$$(b(f)\psi)(x_1, \dots, x_{n-1}) = \sqrt{n} \int_{\mathbb{R}^d} dx f(x) \psi(x, x_1, \dots, x_{n-1})$$

and

$$(b^*(f)\psi)(x_1, \dots, x_{n+1}) = \sqrt{n+1} f(x_1) \psi(x_2, \dots, x_{n+1}).$$

For any $f, g \in L^2(\mathbb{R}^d)$, the operator $b^*(f)b^*(g)b(g)b(f)$ is reduced by the n -particle subspace, therefore for $\sigma > 0$ we may define

$$\tilde{W}_{\Lambda;n}^{\sigma} = \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} dx dy W_{\Lambda}(x, y) b^*(\varphi_x^{\sigma}) b^*(\varphi_y^{\sigma}) b(\varphi_y^{\sigma}) b(\varphi_x^{\sigma}) \Big|_{\bigotimes_{k=1}^n L^2(\mathbb{R}^d)},$$

which is a bounded operator on $L^2(\mathbb{R}^d)^{\otimes n}$. Also define $\tilde{W}_{\Lambda;n}^0$ by

$$(\tilde{W}_{\Lambda;n}^0 h)(x_1, \dots, x_n) = \frac{n(n-1)}{2} W_{\Lambda}(x_1, x_2) h(x_1, \dots, x_n)$$

for $h \in L^2(\mathbb{R}^d)^{\otimes n}$. With these definitions, for every $\sigma \geq 0$,

$$W_{\Lambda;n}^{\sigma} = A_n \tilde{W}_{\Lambda;n}^{\sigma} \Big|_{\big(\bigotimes_{k=1}^n L^2(\mathbb{R}^d)\big)^{-}}$$

where A_n is the antisymmetrization projection $L^2(\mathbb{R}^d)^{\otimes n} \rightarrow (L^2(\mathbb{R}^d)^{\otimes n})^{-}$. We conclude that: if we prove that $\lim_{\sigma \downarrow 0} \tilde{W}_{\Lambda;n}^{\sigma} \psi = \tilde{W}_{\Lambda;n}^0 \psi$ for every ψ of the form $\psi(x_1, \dots, x_n) = \psi_1(x_1) \cdots \psi_n(x_n)$, where $\psi_1, \dots, \psi_n \in \mathcal{S}(\mathbb{R}^d)$, then (A.4) follows. Moreover, since for every $\sigma \geq 0$ $\tilde{W}_{\Lambda;n}^{\sigma}$ acts nontrivially only on the first two particles, it will suffice to prove that $\tilde{W}_{\Lambda;2}^{\sigma} \rightarrow \tilde{W}_{\Lambda;2}^0$ strongly as $\sigma \downarrow 0$.

Let $\sigma > 0$ and introduce the function $\Phi^{\sigma} : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ by setting

$$(A.5) \quad \Phi^{\sigma}(x, y) = \varphi_0^{\sigma}(x) \varphi_0^{\sigma}(y) = \frac{1}{(2\pi\sigma^2)^d} e^{-\frac{1}{2\sigma^2}(|x|^2 + |y|^2)} \quad \text{for any } x, y \in \mathbb{R}^d.$$

It is clear that Φ^{σ} is L^1 -normalized, and moreover, a simple calculation shows that for $\psi \in L^2(\mathbb{R}^{2d})$,

$$\begin{aligned} (\tilde{W}_{\Lambda;2}^{\sigma} \psi)(x_1, x_2) &= \int_{\mathbb{R}^{4d}} dx dy dz_1 dz_2 W_{\Lambda}(x, y) \psi(z_1, z_2) \varphi_{z_1}^{\sigma}(x) \varphi_{x_1}^{\sigma}(x) \varphi_{z_2}^{\sigma}(y) \varphi_{x_2}^{\sigma}(y) \\ &= (\Phi^{\sigma} * (W_{\Lambda}(\Phi^{\sigma} * \psi)))(x_1, x_2). \end{aligned}$$

We are now ready to conclude the proof of the theorem.

Let $\psi \in \mathcal{S}(\mathbb{R}^d)^{\otimes 2}$. Then

$$\tilde{W}_{\Lambda;2}^{\sigma} \psi = \Phi^{\sigma} * (W_{\Lambda}(\Phi^{\sigma} * \psi)) = \Phi^{\sigma} * (W_{\Lambda} \psi) + \Phi^{\sigma} * (W_{\Lambda}(\Phi^{\sigma} * \psi - \psi))$$

The first term above converges to the desired limit. In fact, convolutions with appropriately scaled L^1 -functions converge in L^p -norm, see e.g. [Fol99, Thm 8.14 a)], and thus since $\psi \in L^2(\mathbb{R}^{2d})$,

$$\Phi^{\sigma} * (W_{\Lambda} \psi) \rightarrow W_{\Lambda} \psi \quad \text{in } L^2(\mathbb{R}^{2d}) \text{ as } \sigma \rightarrow 0.$$

We handle the remainder with Young's inequality, i.e. the bound

$$\|\Phi^{\sigma} * (W_{\Lambda}(\Phi^{\sigma} * \psi - \psi))\|_2 \leq \|\Phi^{\sigma}\|_1 \|W_{\Lambda}(\Phi^{\sigma} * \psi - \psi)\|_2 \leq \|W\|_{\infty} \|\Phi^{\sigma} * \psi - \psi\|_2.$$

A further application of [Fol99, Thm 8.14 a)] shows that

$$\lim_{\sigma \downarrow 0} \|\Phi^{\sigma} * \psi - \psi\|_2 = 0$$

which proves the result. \square

APPENDIX B. SEVERAL FOURIER TRANSFORMS

In this section we aim at proving the following Lemma:

Lemma B.1. *Let $a > 0$, $n \in \mathbb{N}$ with $n \geq 2$, and $x \in \mathbb{R}^d$. If $G_a(x) = e^{-a|x|}$, then there exists a constant $D_3 > 0$ such that*

$$(B.1) \quad \underbrace{(G_a * G_a * \cdots * G_a)}_{n-1 \text{ convolutions}}(x) \leq \left(\frac{D_3}{a^d}\right)^n a^d e^{-\frac{a|x|}{4}}.$$

To prove this, we first compute several Fourier transforms. We let \mathcal{F} denote the unitary Fourier transform and \mathcal{F}^* its inverse.

Lemma B.2. *Let $a > 0$ and $G_a(x) = e^{-a|x|}$. Then for $\xi \in \mathbb{R}^d$*

$$(B.2) \quad (\mathcal{F}G_a)(\xi) = \frac{2^{d/2}\Gamma(\frac{d+1}{2})}{\sqrt{\pi}} \frac{a}{(a^2 + \xi^2)^{\frac{d+1}{2}}},$$

where Γ denotes the Gamma function.

Proof. Let $\xi \in \mathbb{R}^d$. We compute using the spherical symmetry of G_a

$$(B.3) \quad \begin{aligned} (\mathcal{F}G_a)(\xi) &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} dx e^{-a|x|} e^{-ix\xi} \\ &= |\xi|^{\frac{2-d}{2}} \int_0^\infty dr r^{\frac{d}{2}} J_{\frac{d-2}{2}}(|\xi|r) e^{-ar} \end{aligned}$$

where $J_\nu(y)$ denotes the Bessel function of first kind. Computing this integral using [GR07, Sec. 6.621 eq. 1] gives

$$(B.4) \quad (\mathcal{F}G_a)(\xi) = \frac{2^{d/2}\Gamma(\frac{d+1}{2})}{\sqrt{\pi}} \frac{a}{(a^2 + |\xi|^2)^{\frac{d+1}{2}}},$$

which is the assertion. □

Lemma B.3. *Let $a > 0$, $\xi \in \mathbb{R}^d$ and*

$$(B.5) \quad H_a(\xi) = \frac{a}{(a^2 + |\xi|^2)^{\frac{d+1}{2}}}.$$

Then for $n \in \mathbb{N}$ we obtain

$$(B.6) \quad (\mathcal{F}^* H_a^n)(x) = \frac{2^{\frac{2-(d+1)n}{2}} a^{\frac{d-(d-1)n}{2}} |x|^{\frac{d(n-1)+n}{2}} K_{\frac{d(n-1)+n}{2}}(a|x|)}{\Gamma(\frac{(1+d)n}{2})},$$

where Γ denotes the Gamma function and $K_\nu(y)$ the modified Bessel function.

Proof. We compute, using again the Fourier transform for spherically symmetric functions,

$$(B.7) \quad \begin{aligned} (\mathcal{F}^* H_a^n)(x) &= \frac{a^n}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d\xi \left(\frac{1}{a^2 + |\xi|^2} \right)^{\frac{n(d+1)}{2}} e^{ix\xi} \\ &= a^n |x|^{\frac{2-d}{2}} \int_0^\infty dr r^{\frac{d}{2}} J_{\frac{d-2}{2}}(|x|r) \left(\frac{1}{a^2 + r^2} \right)^{\frac{n(d+1)}{2}}, \end{aligned}$$

where $J_\nu(y)$ is the Bessel function of first kind. Integrating the latter with the help of [GR07, Sec. 6.565 eq. 4], gives

$$(B.8) \quad (\mathcal{F}^* H_a^n)(x) = \frac{2^{1-\frac{(d+1)n}{2}} a^{\frac{d-(d-1)n}{2}} |x|^{\frac{d(n-1)+n}{2}} K_{\frac{d(n-1)+n}{2}}(a|x|)}{\Gamma\left(\frac{(1+d)n}{2}\right)}.$$

□

Hence, from Lemma B.2 and Lemma B.3 we obtain

$$(B.9) \quad \mathcal{F}^*((\mathcal{F}G_a)^n)(x) = \frac{(\Gamma(\frac{d+1}{2}))^n 2^{\frac{2-n}{2}} a^{\frac{d-(d-1)n}{2}} |x|^{\frac{d(n-1)+n}{2}} K_{\frac{d(n-1)+n}{2}}(a|x|)}{\pi^{n/2} \Gamma\left(\frac{(d+1)n}{2}\right)}.$$

Lemma B.4. *Let $\eta > 0$. The modified Bessel function K_η satisfies for $y > 0$ the bound*

$$(B.10) \quad 0 \leq K_\eta(y) \leq \frac{4^\eta}{y^\eta} e^{-\frac{y}{4}} \Gamma(\eta).$$

Proof. We write the modified Bessel function K_η as

$$(B.11) \quad K_\eta(y) = \int_0^\infty dt e^{-y \cosh(t)} \cosh(\eta t),$$

see [GR07, Sec. 8.432 eq. 1]. Using $\frac{1}{2}e^x \leq \cosh(x) \leq e^x$ valid for all $x \geq 0$, we obtain

$$(B.12) \quad \int_0^\infty dt e^{-y \cosh(t)} \cosh(\eta t) \leq \int_0^\infty dt e^{-\frac{y e^t}{2}} e^{\eta t} = \int_1^\infty du e^{-\frac{y u}{2}} u^{\eta-1},$$

where we performed the change of variables $u = e^t$ in the last line. Now, for $u \geq 1$ and $y > 0$ we have $e^{-\frac{y u}{2}} \leq e^{-\frac{y}{4}} e^{-\frac{y u}{4}}$ and therefore

$$(B.13) \quad \begin{aligned} \int_1^\infty du e^{-\frac{y u}{2}} u^{\eta-1} &\leq e^{-\frac{y}{4}} \int_0^\infty du e^{-\frac{y u}{4}} u^{\eta-1} \\ &= \frac{4^\eta}{y^\eta} e^{-\frac{y}{4}} \int_0^\infty du e^{-u} u^{\eta-1} = \frac{4^\eta}{y^\eta} e^{-\frac{y}{4}} \Gamma(\eta). \end{aligned}$$

□

Proof of Lemma B.1. Starting with (B.9) and using Lemma B.4 we obtain,

$$(B.14) \quad \begin{aligned} (G_a * G_a * \dots * G_a)(x) &= (2\pi)^{d(n-1)/2} \mathcal{F}^*((\mathcal{F}G_a)^n)(x) \\ &\leq \frac{(\Gamma(\frac{d+1}{2}))^n \pi^{\frac{d-1}{2}} 2^{\frac{3d+1}{2}} \pi^{-\frac{d}{2}} 2^{\frac{1}{2}} a^{-d(n-1)} \Gamma\left(\frac{d(n-1)+n}{2}\right) e^{-\frac{a|x|}{4}}}{\Gamma\left(\frac{(d+1)n}{2}\right)} \\ &\leq D_3^n a^{-d(n-1)} e^{-\frac{a|x|}{4}} \end{aligned}$$

for some explicit constant $D_3 > 0$ depending on d , where we used that $(2/\pi^d)^{\frac{1}{2}} \leq 1$ and

$$(B.15) \quad \frac{\Gamma\left(\frac{d(n-1)+n}{2}\right)}{\Gamma\left(\frac{(d+1)n}{2}\right)} \leq 1.$$

This gives the assertion. □

REFERENCES

- [AW12] M. Aizenman and S. Warzel, Absolutely continuous spectrum implies ballistic transport for quantum particles in a random potential on tree graphs, *J. Math. Phys.* **53**, 095205 (2012).
- [ALBN10] L. Amour, P. Levy-Bruhl and J. Nourrigat, Dynamics and Lieb-Robinson estimates for lattices of interacting anharmonic oscillators, *Colloq. Math.* **118**, 609–648 (2010).
- [BBDRF18] S. Bachmann, A. Bols, W. De Roeck and M. Fraas, Quantization of conductance in gapped interacting systems, *Ann. Henri Poincaré* **19**, 695–708 (2018), [arXiv:1707.06491](#).
- [BBDRF19] S. Bachmann, A. Bols, W. De Roeck and M. Fraas, A many-body index for quantum charge transport, [arXiv:1810.07351](#). To appear in *Commun. Math. Phys.*
- [BR97] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics*, 2nd ed., vol. 2, Springer Verlag, 1997.
- [BHM10] S. Bravyi, M. Hastings and S. Michalakis, Topological quantum order: stability under local perturbations, *J. Math. Phys.* **51**, 093512 (2010).
- [BH11] S. Bravyi and M. B. Hastings, A short proof of stability of topological order under local perturbations, *Commun. Math. Phys.* **307**, 609 (2011).
- [BHV06] S. Bravyi, M. B. Hastings and F. Verstraete, Lieb-Robinson bounds and the generation of correlations and topological quantum order, *Phys. Rev. Lett.* **97**, 050401 (2006).
- [BdSP17] J.-B. Bru and W. de Siquiera Pedra, *Lieb-Robinson Bounds for Multi-Commutators and Applications to Response Theory*, Springer Briefs in Mathematical Physics, vol. 13, Springer Nature, 2017.
- [Buc17] D. Buchholz, The resolvent algebra for oscillating lattice systems: Dynamics, ground and equilibrium states, *Commun. Math. Phys.* **353**, 691–716 (2017).
- [BG08] D. Buchholz and H. Grundling, The resolvent algebra: A new approach to canonical quantum systems, *J. Funct. Anal.* **254**, 2725–2779 (2008).
- [CMPGS13] J. Cirac, S. Michalakis, D. Perez-Garcia and N. Schuch, Robustness in projected entangled pair states, *Phys. Rev. B* **88**, 115108 (2013).
- [DRS19] W. De Roeck and M. Salmhofer, Persistence of exponential decay and spectral gaps for interacting fermions, *Commun. Math. Phys.* **365**, 773–796 (2019).
- [Dys52] F. Dyson, Divergence of perturbation theory in quantum electrodynamics, *Phys. Rev.* **85**, 631–632 (1952).
- [EESY04] A. Elgart, L. Erdős, B. Schlein and H.-T. Yau, Nonlinear Hartree equation as the mean field limit of weakly coupled fermions, *J. Math. Pures Appl.* **83**, 1241–1273 (2004).
- [Fol99] G. Folland, *Real Analysis: Modern Techniques and Their Applications*, John Wiley & Sons, 1999.
- [FMU03] J. Fröhlich, M. Merkli and D. Ueltschi, Dissipative transport: Thermal contacts and tunnelling junctions, *Ann. Henri Poincaré* **4**, 897–945 (2003).
- [FP18] J. Fröhlich and A. Pizzo, Lie-Schwinger block-diagonalization and gapped quantum chains (2018), [arXiv:1812.02457](#).
- [GR07] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, seventh ed., Elsevier/Academic Press, Amsterdam, 2007, translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger.
- [GR17] H. Grundling and G. Rudolph, Dynamics for QCD on an infinite lattice, *Commun. Math. Phys.* **349**, 1163–1202 (2017).
- [Has19] M. B. Hastings, The stability of free Fermi Hamiltonians, *J. Math. Phys.* **60**, 042201 (2019), [arXiv:1706.02270](#).
- [HK06] M. B. Hastings and T. Koma, Spectral gap and exponential decay of correlations, *Commun. Math. Phys.* **265**, 781–804 (2006).
- [HM15] M. B. Hastings and S. Michalakis, Quantization of hall conductance for interacting electrons on a torus, *Commun. Math. Phys.* **334**, 433–471 (2015).
- [Kli10] I. Klich, On the stability of topological phases on a lattice, *Ann. Phys.* **325**, 2120–2131 (2010).
- [LS15] M. Lewin and J. Sabin, The Hartree equation for infinitely many particles. I. Well-posedness theory, *Commun. Math. Phys.* **334**, 117–170 (2015).
- [LR72] E. Lieb and D. Robinson, The finite group velocity of quantum spin systems, *Commun. Math. Phys.* **28**, 251–257 (1972).
- [MZ13] S. Michalakis and J. Zwolak, Stability of frustration-free Hamiltonians, *Commun. Math. Phys.* **322**, 277–302 (2013).
- [NOS06] B. Nachtergaele, Y. Ogata and R. Sims, Propagation of correlations in quantum lattice systems, *J. Stat. Phys.* **124**, 1–13 (2006).
- [NRSS09] B. Nachtergaele, H. Raz, B. Schlein and R. Sims, Lieb-Robinson bounds for harmonic and anharmonic lattice systems, *Commun. Math. Phys.* **286**, 1073–1098 (2009).

- [NSSSZ10] B. Nachtergaele, B. Schlein, R. Sims, S. Starr and V. Zagrebnov, On the existence of the dynamics for anharmonic quantum oscillator systems, *Rev. Math. Phys.* **22**, 207–231 (2010).
- [NSY18] B. Nachtergaele, R. Sims and A. Young, Lieb-Robinson bounds, the spectral flow, and stability for lattice fermion systems, in *Mathematical Results in Quantum Physics*, Contemporary Mathematics, vol. 717, F. Bonetto, D. Borthwick, E. Harrell and M. Loss, eds., Amer. Math. Soc., 2018, pp. 93–115.
- [NSY19] B. Nachtergaele, R. Sims and A. Young, Quasi-locality bounds for quantum lattice systems. I. Lieb-Robinson bounds, quasi-local maps, and spectral flow automorphisms, *J. Math. Phys.* **60**, 061101 (2019).
- [NSY20] B. Nachtergaele, R. Sims and A. Young, Quasi-locality bounds for quantum lattice systems and perturbations of gapped ground states II. (2020), in preparation.
- [NT90] H. Narnhofer and W. Thirring, Quantum field theories with Galilei-invariant interactions, *Phys. Rev. Lett.* **64**, 1863–1866 (1990).
- [RS75] M. Reed and B. Simon, *Fourier Analysis, Self-Adjointness. Methods of Modern Mathematical Physics*, vol. 2, Academic Press, 1975.
- [RS80] M. Reed and B. Simon, *Functional Analysis. Methods of Modern Mathematical Physics*, revised and enlarged ed., vol. 1, Academic Press, 1980.
- [Rob68] D. W. Robinson, Statistical mechanics of quantum spin systems II, *Commun. Math. Phys.* **7**, 337–348 (1968).
- [Rue69] D. Ruelle, *Statistical Mechanics: Rigorous Results*, Benjamin, New York, 1969.
- [Sak91] S. Sakai, *Operator algebras in dynamical systems*, Encyclopedia of Mathematics and its Applications, vol. 41, Cambridge University Press, 1991, the theory of unbounded derivations in C^* -algebras.
- [SPGC11] N. Schuch, D. Perez-Garcia and I. Cirac, Classifying quantum phases using matrix product states and peps, *Phys. Rev. B* **84**, 165139 (2011).
- [Str68] R. F. Streater, On certain non-relativistic quantized fields, *Commun. Math. Phys.* **7**, 93–98 (1968).
- [SW70] R. F. Streater and I. F. Wilde, The time evolution of quantized fields with bounded quasi-local interaction density, *Commun. Math. Phys.* **17**, 21–32 (1970).
- [Tes09] G. Teschl, *Mathematical methods in quantum mechanics*, Graduate Studies in Mathematics, vol. 99, American Mathematical Society, Providence, RI, 2009, with applications to Schrödinger operators.
- [Ver11] A. F. Verbeure, *Many-Body Boson Systems*, Theoretical and Mathematical Physics, Springer-Verlag, 2011.
- [Wei80] J. Weidmann, *Linear Operators in Hilbert Spaces*, Graduate Texts in Mathematics, vol. 68, Springer, 1980.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, DAVIS, CA 95616, USA

E-mail address: mgebert@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS AND CENTER FOR QUANTUM MATHEMATICS AND PHYSICS, UNIVERSITY OF CALIFORNIA, DAVIS, DAVIS, CA 95616, USA

E-mail address: bxn@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, DAVIS, CA 95616, USA

E-mail address: jreschke@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUSCON, AZ 85721, USA

E-mail address: rsims@math.arizona.edu