On the accurate reproduction of strongly repulsive interatomic potentials

Susi Lehtola

Department of Chemistry, University of Helsinki,

P.O. Box 55 (A. I. Virtasen aukio 1), FI-00014 University of Helsinki, Finland*

Knowledge of the repulsive behavior of potential energy curves V(R) at $R \to 0$ is necessary for understanding and modeling irradiation processes of practical interest. V(R) is in principle straightforward to obtain from electronic structure calculations; however, commonly-used numerical approaches for electronic structure calculations break down in the strongly repulsive region due to the closeness of the nuclei. In the present work, we show by comparison to fully numerical reference values that a recently developed procedure [S. Lehtola, J. Chem. Phys. 151, 241102 (2019)] can be employed to enable accurate linear combination of atomic orbitals calculations of V(R) even at small R by a study of the seven nuclear reactions $\text{He}_2 \longrightarrow \text{Be}$, $\text{HeNe} \longrightarrow \text{Mg}$, $\text{Ne}_2 \longrightarrow \text{Ca}$, $\text{HeAr} \longrightarrow \text{Ca}$, $\text{MgAr} \longrightarrow \text{Zn}$, $\text{Ar}_2 \longrightarrow \text{Kr}$, and $\text{NeCa} \longrightarrow \text{Zn}$.

I. INTRODUCTION

The interaction of high-energy particles with matter is typically modeled using pairwise potentials (see *e.g.* chapter 6 of [1]), as the dominant interactions are determined by the highly repulsive nuclear Coulomb barriers that are pairwise terms; see *e.g.* [2] for a recent numerical demonstration for low-energy projectiles incident on copper surfaces. Most practical simulations employ the universal potential by Ziegler, Biersack and Littmark [3] (ZBL) which is based on Thomas–Fermi calculations of the repulsive barrier. However, Thomas–Fermi theory has significant shortcomings; for instance, it is well known not to bind any molecules, and a method lacking these shortcomings like Hartree–Fock (HF) or densityfunctional theory [4, 5] (DFT) would certainly be more attractive.

Ab initio calculations of the diatomic potential energy curve (PEC), denoted here as $V_{AB}(R)$, are, however, challenging at small internuclear distances R due to the closeness of the two nuclei. In contrast to chemistry at ambient conditions, even the innermost core electrons may be significantly affected by the interaction between the two atoms: for instance, in the $\mathrm{Ar}_2 \Longleftarrow \mathrm{Kr}$ nuclear reaction obtained as $R \rightarrow 0$, the two [Ne] $3s^23p^6$ electronic configurations of the argon atoms deform into the single $[Ne]3s^23p^64s^23d^{10}4p^6$ configuration of the krypton atom. An extremely flexible numerical approach must be used in order to describe such changes accurately, obviously disallowing the use of pseudopotential and frozencore approaches. Although some efforts for the *ab initio* description of the Coulomb barrier have been made in the literature (see e.g. [2], [6-17] and references therein), the problem of facile computation of $V_{AB}(R)$ for $R \to 0$ remains still unsolved in the general case.

All-electron calculations are typically undertaken within the linear combination of atomic orbitals (LCAO) approach. However, also the LCAO approach fails in this case, because the basis functions on the atoms A and B

quickly become linearly dependent when $R \rightarrow 0$. Moreover, large atomic basis sets should be used in order to allow the necessary flexibility for the core orbitals to deform in presence of the other nucleus and its electrons. But, the more functions are included in the calculations, the more linear dependencies are generated when the nuclei start coinciding, and the calculations become numerically unstable as the basis set becomes ill-behaved.

As always, fully numerical electronic structure calculations are one option, see [18] for a recent review. Here, the numerical basis set can always be chosen in such a way that linear dependencies do not arise even at small R. However, fully numerical approaches carry a much higher computational cost than that of LCAO calculations using *e.g.* Gaussian basis sets, and may also be harder to set up; see the discussion in [18, 19]. Moreover, fully numerical electronic structure programs are lessdeveloped than Gaussian-basis ones, because the huge number of basis functions in a fully numerical approach may *e.g.* make sophisticated convergence algorithms intractable [18], making it more difficult to carry out the wanted electronic structure calculations.

Despite the numerical problems encountered in standard LCAO approaches, it should be perfectly well possible to describe diatomic molecules using atomic basis sets even at small internuclear distances, because at small Rthe molecule looks like the compound atom that is especially easy to describe with atomic basis sets. This means that the problems in LCAO calculations should be circumventable by adopting a basis set that is adapted to the molecular geometry. (In contrast, significant distortions to the electronic structure of atoms and molecules can be observed *e.g.* in strong magnetic fields as discussed in [20] and references therein, in which case LCAO calculations become unreliable.)

Because the electronic structure at $R \rightarrow 0$ may be quite far from those for which typical basis sets have been optimized, one can customize the basis set for the system by hand as in [7]. (Alternatively, one could also optimize a new basis set from scratch for the system.) However, given that this would lead to a different basis set for every molecule and for every molecular geometry, a systematic study of the repulsive potentials of all the elements in

^{*} susi.lehtola@alumni.helsinki.fi

the periodic table would be faced with a gargantuan task for basis set generation. For instance, the PECs for all 4186 diatomic molecules from Z = 1 to Z = 92 were calculated in [17] at internuclear distances ranging from R = 0.002 Å to R = 1000 Å; this is only feasible with a fully automatic approach. (Convergence to the basis set limit was not checked in [17], and we will show later in the present work that the values are not converged.)

Issues with linear dependencies are encountered also in other applications of quantum chemistry. For instance, an accurate description of weakly bound anions may require the use of several shells of diffuse functions on each atom, making the molecular basis set ill-behaved due to linear dependencies [21]. An approach for curing overcompleteness issues in the study of weakly bound anions with LCAO basis sets by pivoted Cholesky decompositions was recently proposed in [22]. Cholesky decomposition algorithms have a long history in quantum chemistry, starting with with the decomposition of the two-electron integral tensor proposed over 40 years ago by Beebe and Linderberg [23] that has recently become popular due to efficient implementations afforded by modern computer architectures, see e.g. [24, 25]. The tractability of the Cholesky decomposition of the twoelectron integrals relies on the set of basis function products $\chi_i(\mathbf{r})\chi_i(\mathbf{r})$ being highly linearly dependent: due to these dependencies, the number of Cholesky vectors of the two-electron integrals tensor grows only linearly with system size. Also method-specific variants related to the two-electron integrals tensor decomposition have been suggested, see [26] for a review. The construction of localized orbitals by the Cholesky decomposition of the LCAO one-electron density matrix has also been proposed [27].

The mathematical closeness of the Cholesky decomposition of the two-electron integrals tensor, $(ij|kl) = \sum_{P} L^{P}_{(ij)} L^{P}_{(kl)}$ in chemists' notation, to resolution-of-the-identity methods [28], $(ij|kl) \approx \sum_{AB} (ij|A)(A|B)^{-1}(B|kl)$, led to a black-box procedure for the formation of auxiliary basis sets [29, 30], in which the pivot index of the Cholesky decomposition of the twoelectron integrals is used to determine the auxiliary functions; the resulting set of auxiliary functions is better behaved than the set of basis function products. Building on the work of [29, 30], we proposed curing significant linear dependencies in overcomplete basis sets by the extraction of a well-behaved subset of the basis functions using a pivoted Cholesky procedure on the overlap matrix $S_{\mu\nu} = \langle \mu | \nu \rangle$ to a predefined threshold τ [22]; this yields an optimal approximation for the original over-complete S [31]. Electronic structure calculations can be carried out in the basis set defined by the set of pivot functions without problems. Before [22], Cholesky decompositions of the overlap matrix appear to have only been used in the full, unpivoted form [32–34] that is not safe for illconditioned matrices S.

In the present work, we show that the partial Cholesky decomposition algorithm proposed in [22] for calculations with basis sets containing a large number of linearly dependent diffuse functions presents a solution to the present problem of the calculation of strongly repulsive interatomic potentials by allowing the use of standard atomic basis sets even at $R \rightarrow 0$, since the basis function degeneracies that would otherwise prevent reliable electronic structure calculations from taking place are cleaned away automatically.

As our aim is simply to prove that the basis set limit can be reached without problem even at tiny values of R, we have chosen to study a set of seven nuclear reactions involving only closed-shell atoms: $He_2 \implies Be_2$ $\operatorname{HeNe} \rightleftharpoons \operatorname{Mg}, \operatorname{Ne}_2 \rightleftharpoons \operatorname{Ca}, \operatorname{HeAr} \rightleftharpoons \operatorname{Ca}, \operatorname{MgAr} \rightleftharpoons \operatorname{Ca}$ Zn, $Ar_2 \rightleftharpoons Kr$, and NeCa \rightleftharpoons Zn. We show that the suggested Cholesky procedure reproduces fully numerical HF reference values for the reactions, while the values reported in [17] are not converged for small R. The present case is much more challenging than that of [22], as *e.g.* in He_2 , Ne_2 , and Ar_2 all basis functions of an atom become fully degenerate with the basis function of the other atom for $R \to 0$. Our calculations will be described in section II and our results reported in section III. The work is briefly summarized and discussed in section IV. Atomic units are used throughout the manuscript.

II. COMPUTATIONAL DETAILS

The PEC for atoms A and B is defined as

$$V_{AB}(R) = E_{\rm tot}^{A+B}(R) - E_{\rm el}^{A} - E_{\rm el}^{B}$$
 (1)

where $E_{\text{tot}}^{A+B}(R)$ is the total energy from the electronic structure calculation for the nuclei A and B separated by a distance of R, and E_{el}^A and E_{el}^B are the electronic energies of the non-interacting atoms, respectively. The total energy E_{tot}^{A+B} can be decomposed into a sum of the electronic energy $E_{\text{el}}^{A+B}(R)$ and the nuclear repulsion energy $E_{\text{nuc}}^{A+B}(R)$. Since the electronic energy of the compound atom (A+B) is finite, $E_{\text{tot}}^{A+B}(R)$ behaves asymptotically as $E_{\text{tot}}^{A+B}(R) \approx E_{\text{nuc}}^{A+B}(R) = Z_A Z_B R^{-1}$ for small R. Because $V_{AB}(R)$ thus diverges for small R, it is typical to report the PEC in terms of a screening function

$$\Phi_{AB}(R) = \frac{V_{AB}(R)}{E_{\text{nuc}}^{A+B}(R)} = \frac{RV_{AB}(R)}{Z_A Z_B}$$
(2)

as it is more easily manipulable, having the limits $\Phi_{AB}(0) = 1$ and $\Phi_{AB}(\infty) = 0$.

Although the procedure of [22] can be used with any type of atomic basis set (see [18] for a review thereof), Gaussian basis sets are employed in the present work. Furthermore, while the approach of [22] can also be applied to density functional or post-HF calculations, the HF level of theory is used in the present work as it has been found to be sufficient for the reproduction of repulsive potentials [10].

The ERKALE program [35, 36] is used for the Gaussianbasis calculations. The nuclei A and B are placed in the ERKALE calculations along the z axis at (0, 0, -R/2) and (0, 0, R/2), respectively, along with their atomic basis functions. Next, in order to be able to describe the compound atom (A + B) limit, basis functions for the compound atom are included in the calculation; placing the compound nucleus at the center of charge at $(0, 0, (Z_B - Z_A)R/[2(Z_B + Z_A)])$ leads to a vanishing dipole moment of the nuclear charge distribution, and hopefully a more accurate calculation. Once the basis functions for the compound nucleus have been added, the construction of the one-electron basis $\{|\mu\rangle\}$ is complete; however, by this stage the basis set is likely overcomplete.

Next, the overlap matrix $S_{\mu\nu} = \langle \mu | \nu \rangle$, its eigenvalues λ_i and its reciprocal condition number

$$r = \frac{\lambda_{\min}}{\lambda_{\max}} \tag{3}$$

are computed. If the basis set is found to be overcomplete, *i.e.*, r is found to be smaller than the machine epsilon, the Cholesky procedure of [22] is used to regularize the molecular basis set. The procedure uses a pivoted Cholesky decomposition to pick a subset of the basis functions $\{|\mu\rangle\}$ that spans all of the functions in the original basis set up to a predefined threshold. The resulting reduced-size basis is numerically well-conditioned, and poses no problems to electronic structure calculations which then proceed as usual. The procedure is implemented as a modification [22] to the canonical orthogonalization method [37]; a Cholesky threshold of 10^{-7} and a linear dependence threshold of 10^{-5} are used in the present work. As the basis set is normalized, $S_{\mu\mu} = 1$, the first function of the Cholesky procedure will be the first function in the basis set. However, since diffuse basis functions may be representable as superpositions of more localized basis functions while the converse is unlikely to be the case, the basis functions are reorganized from tight to diffuse before the Cholesky procedure. This procedure was found to yield more compact reduced basis sets in [22].

The screening function $\Phi(R)$ is computed with ERKALE on a logarithmic grid consisting of 121 points ranging from $R = 10^{-5}$ Å to R = 10 Å. The Gaussianbasis values are then compared to a set of fully numerical reference values obtained with the HELFEM program [19, 38, 39]. The superposition of atomic potentials (SAP) initial guess [40] is used in all ERKALE and HELFEM calculations in combination with local exchange potentials recently determined at the complete basis set limit [41]. The SAP guess correctly includes the significant Pauli repulsion between the electrons on the two nuclei at small R in contrast to its commonlyused alternatives discussed in [40], thus leading to faster convergence of the self-consistent field procedure.

Only singlet Σ wave functions are considered in the present work, in analogy to [10]. In the cases of He₂ \rightleftharpoons Be, HeNe \rightleftharpoons Mg, Ne₂ \rightleftharpoons Ca, and HeAr \rightleftharpoons Ca, the large-R and small-R wave functions have the same electronic configurations: two occupied σ orbitals for He₂ and Be, four σ and one π orbital for HeNe and Mg, and

six σ and two π orbitals for Ne₂, Ca, and HeAr; each σ and π orbital fitting two and four electrons, respectively [18]. For the heavier systems, MgAr \Longrightarrow Zn, Ar₂ \rightleftharpoons Kr, and NeCa \rightleftharpoons Zn, the electronic configurations are different at small R and at large R, and both states were calculated: nine σ and three π in MgAr and NeCa; seven σ , three π and one δ orbital in Zn; ten σ and four π orbitals in Ar₂; and eight σ , four π and one δ orbital in Kr; δ orbitals likewise fitting four electrons [18]. The values reported correspond to the lower state in each case; for instance, the Kr configuration is lower in Ar₂ for $R \leq$ 0.56 Å, the state crossing depending on the used basis set.

III. RESULTS

Very accurate LCAO calculations can be performed both at small R and at large R, as in the former case a single expansion center is sufficient, and as in the latter the basis functions on the two centers do not develop strong linear dependencies. For this reason, we start off in table I by comparing the values of the screening function $\Phi(R)$ at intermediate values of R for the decontracted double- to quadruple- ζ pc-n basis sets [42] (denoted as un-pc-1, un-pc-2, and un-pc-3, respectively) as well as for the universal Gaussian basis set [43] (UGBS) to fully numerical reference values.

Examination of the data in table I shows that good results are already obtained with the double- ζ un-pc-1 basis set, while the UGBS basis set appears to reproduce values that are in-between those of the triple- ζ unpc-2 and the quadruple- ζ un-pc-3 basis set at small R. This suggests that the screening function is insensitive to polarization functions at small R; however, some polarization effects are already described by the compound nucleus basis functions included at the center of charge. As the UGBS basis set is available for most of the periodic table and equivalent atomic basis sets can be easily generated, see [44], we choose the UGBS basis set for the rest of the work.

To confirm the finding of [10] that the screening function has a negligible dependence on the employed level of theory, we also report fully numerical reference values for Ar_2 calculated with HELFEM using the local density approximation (LDA), in which the local exchange functional [45, 46] is combined with the Vosko–Wilk–Nusair correlation functional [47] as in [10]. The differences of the HF and LDA screening functions are only seen in the third decimal, confirming that HF or DFT is suitable for the present purposes.

The screening functions for the seven nuclear reactions computed with the UGBS basis set are shown in figure 1 for He₂ \implies Be, figure 2 for HeNe \implies Mg, figure 3 for Ne₂ \implies Ca, figure 4 for HeAr \implies Ca, figure 5 for MgAr \implies Zn, figure 6 for Ar₂ \implies Kr, and figure 7 for NeCa \implies Zn. The curves are smooth and the agreement with fully numerical reference values is superb in



Figure 1. UGBS screening function for the $\text{He}_2 \iff$ Be reaction with fully numerical reference values (+).

all cases.

All of these reactions have also been studied in [17] with the LDA approach of [10] [48]. However, out of the seven reactions currently examined, [17] only only reports data for $\operatorname{Ar}_2 \rightleftharpoons \operatorname{Kr}$. A comparison to the UGBS results and fully numerical HF and LDA reference values is shown in figure 8. The data from [17] agree with the present values at large R, but discrepancies are visible for R < 0.1 Å. The UGBS data is agrees with the fully numerical HF and LDA reference data, indicating that an insufficient basis set was used in [17].

IV. SUMMARY AND DISCUSSION

We have shown by comparison to fully numerical Hartree–Fock reference values that accurate potential energy curves can be reproduced with linear combination of atomic orbitals (LCAO) calculations even in the strongly repulsive region at small internuclear distances—where even the core orbital basis functions become fully linearly dependent—by using a recently suggested procedure [22] to eliminate linear dependencies from the basis set. As LCAO calculations are faster and easier to run than fully numerical ones, the automated procedure of the present work enables the systematical calculation of screening functions along the lines of [17] but with guaranteed accuracy.

Although the present study has been limited to diatomic molecules, the underlying Cholesky decomposition has already been shown to work for polyatomic molecules [22], making the calculation of repulsive manybody potentials also possible.



Figure 2. UGBS screening function for the HeNe \implies Mg reaction with fully numerical reference values (+).



Figure 3. UGBS screening function for the Ne₂ \implies Ca reaction with fully numerical reference values (+).

The facile computation of the repulsive barrier afforded by the present method should make it easier to study various irradiation processes, in which the purely repulsive part of the potential plays a pivotal role. For instance, defect formation and migration in materials subjected to particle bombardment is determined purely by the repulsive part of the potential [15], and accounting for this kind of radiation damage is an important aspect in the design of radiation shielding materials of fusion reactors

		10^{-3}	$10^{-2.5}$	10^{-2}	$10^{-1.5}$	$10^{-1.25}$	10^{-1}	$10^{-0.75}$	$10^{-0.5}$	$10^{-0.25}$	10^{0}
He_2	HelfEM	0.99582	0.98678	0.95831	0.87089	0.77943	0.64265	0.47484	0.34264	0.21805	0.07205
	Δun -pc-1	0.00000	-0.00001	-0.00005	-0.00007	0.00027	0.00095	0.00068	0.00023	0.00162	0.00011
	Δ un-pc-2	0.00000	0.00000	0.00000	0.00004	0.00006	0.00008	0.00007	0.00005	0.00041	-0.00005
	Δ un-pc-3	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00001	0.00002	0.00002	0.00000
	ΔUGBS	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00001	0.00006	0.00021	0.00022
HeNe	HelfEM	0.99356	0.97966	0.93658	0.81686	0.71223	0.57716	0.40213	0.21037	0.09810	0.03762
	Δun -pc-1	-0.00001	-0.00003	-0.00007	0.00000	0.00013	-0.00007	-0.00004	0.00016	-0.00022	-0.00111
	Δ un-pc-2	0.00000	0.00000	0.00000	0.00006	0.00006	0.00009	0.00024	0.00037	0.00014	-0.00010
	Δ un-pc-3	0.00000	0.00000	0.00000	0.00000	0.00001	0.00004	0.00006	0.00003	0.00002	0.00000
	ΔUGBS	0.00000	0.00000	0.00000	0.00000	0.00001	0.00008	0.00031	0.00085	0.00120	0.00031
Ne_2	HelfEM	0.99207	0.97503	0.92326	0.78984	0.67932	0.53204	0.36532	0.20790	0.07321	0.02656
	Δ un-pc-1	0.00000	-0.00001	0.00001	-0.00001	0.00008	0.00053	0.00065	0.00029	-0.00007	-0.00035
	Δ un-pc-2	0.00000	0.00000	0.00000	0.00001	0.00006	0.00009	0.00007	0.00009	0.00007	-0.00001
	Δ un-pc-3	0.00000	0.00000	0.00000	0.00000	0.00001	0.00001	0.00002	0.00003	0.00001	0.00001
	ΔUGBS	0.00000	0.00000	0.00000	0.00000	0.00000	0.00001	0.00002	0.00012	0.00020	0.00011
HeAr	HELFEM	0.99228	0.97569	0.92539	0.79716	0.69117	0.54811	0.38703	0.24271	0.10196	0.04028
	∆un-pc-1	0.00000	0.00000	0.00001	0.00012	0.00029	0.00058	0.00060	0.00043	0.00032	0.00001
	Δun -pc-2	0.00000	0.00000	0.00001	0.00008	0.00008	0.00019	0.00028	0.00020	0.00009	0.00001
	Δun-pc-3	0.00000	0.00000	0.00000	0.00001	0.00004	0.00010	0.00011	0.00006	0.00002	0.00001
	$\Delta 0GBS$	0.00000	0.00000	0.00000	0.00001	0.00004	0.00016	0.00038	0.00043	0.00084	0.00083
MaAn	UDIFEM	0.00091	0.07114	0.01200	0 77106	0.65400	0 50202	0 22521	0 17096	0 07499	0 00000
MgAI	Aun ng 1	0.99081	0.97114	0.91299	0.77190	0.00409	0.00202	0.00021	0.17920	0.07422	0.02229
	$\Delta un-pc-1$	0.00000	0.00000	0.00000	0.00003	0.00000	0.00008	0.00024 0.00017	0.00091	0.00032	0.00018
	Aun-pc-2	0.00000	0.00000	0.00000	0.00001	0.00002	0.00004	0.00017	0.00000	0.00012	0.00000
	AUGBS	0.00000	0.00000	0.00000	0.00000	0.00000	0.00002	0.00000	0.00010	0.00000	0.00002
		0.00000	0.00000	0.00000	0.00000	0.00001	0.00000	0.00020	0.00000	0.00025	0.00020
Ara	Helfem	0.99011	0.96900	0.90749	0.76090	0.63761	0.48451	0.31334	0.17568	0.07255	0.02137
2	ΔLDA^a	0.00001	0.00003	0.00009	0.00012	0.00017	0.00028	0.00073	-0.00130	0.00174	no data
	Δun -pc-1	0.00000	0.00000	0.00000	0.00003	0.00005	0.00013	0.00089	0.00158	0.00290	0.00155
	Δun -pc-2	0.00000	0.00000	0.00000	0.00001	0.00002	0.00009	0.00080	0.00117	0.00115	0.00066
	Δ un-pc-3	0.00000	0.00000	0.00000	0.00000	0.00001	0.00006	0.00055	0.00029	0.00010	0.00030
	ΔUGBS	0.00000	0.00000	0.00000	0.00000	0.00001	0.00009	0.00078	0.00141	0.00275	0.00078
NeCa	HelfEM	0.99082	0.97117	0.91310	0.77230	0.65438	0.50203	0.33509	0.17662	0.07093	0.01768
	Δ un-pc-1	0.00000	0.00000	0.00002	0.00005	0.00012	0.00032	0.00057	0.00061	0.00023	0.00008
	Δ un-pc-2	0.00000	0.00000	0.00000	0.00002	0.00005	0.00007	0.00016	0.00026	0.00011	0.00004
	Δ un-pc-3	0.00000	0.00000	0.00000	0.00000	0.00001	0.00002	0.00006	0.00007	0.00002	0.00001
	ΔUGBS	0.00000	0.00000	0.00000	0.00000	0.00001	0.00003	0.00021	0.00095	0.00047	0.00049

Table I. Values of screening function $\Phi(R)$ computed at various points R (value in Å given on the first row) with the fully numerical HELFEM program. The Gaussian-basis-set truncation errors $\Delta \text{basis} = \Phi^{\text{basis}}(R) - \Phi^{\text{reference}}(R)$ of the un-pc-n and UGBS basis sets are also shown; these calculations were done with ERKALE. The data for Ar₂ also includes the differences between the LDA and HF screening functions' reference values $\Delta \text{LDA} = \Phi^{\text{LDA}}(R) - \Phi^{\text{HF}}(R)$, both of which have been computed with HELFEM.



Figure 4. UGBS screening function for the HeAr \iff Ca reaction with fully numerical reference values (+).



Figure 5. UGBS screening function for the MgAr \implies Zn reaction with fully numerical reference values (+).

[49, 50].

The present study has been limited to non-relativistic calculations on light, closed-shell atoms. As relativistic effects increase rapidly in Z [51, 52], they are more important at the compound nucleus limit $R \to 0$ than at large R. Note also that in contrast to usual applications to chemistry, the screening function merits from no systematic error cancellation from the subtraction of atomic energies. The present procedure can, however,



Figure 6. UGBS screening function for the ArAr \iff Kr reaction with fully numerical reference values (+).



Figure 7. UGBS screening function for the NeCa \implies Zn reaction with fully numerical reference values (+).

be straightforwardly extended to relativistic methods as well, making it possible to model the relativistic effects. Open-shell atoms as well as relativistic effects will be visited in future work.





Figure 8. Comparison of the ArAr \iff Kr UGBS/Hartree-Fock screening function (black line) against LDA data from [17] (red line). Fully numerical Hartree–Fock (black +) as well as LDA (red \times) reference values are also shown.

[3] J. F. Ziegler, U. Littmark, and J. P. Biersack, The stopping and range of ions in solids (Pergamon, New York,

and

W. Kohn and L. J. Sham, Phys. Rev. 140, A1133 (1965).

[6] N. H. Sabelli, M. Kantor, R. Benedek, and T. L. Gilbert,

[8] J. Keinonen, A. Kuronen, P. Tikkanen, H. G.

Börner, J. Jolie, S. Ulbig, E. G. Kessler, R. M.

Kuronen,

and

Nucl. Instrum. Methods Phys. Res., Sect. B 88, 382 (1994).

Nucl. Instrum. Methods Phys. Res., Sect. B 132, 45 (1997).

W.

and T. L. Gilbert,

and A. P. Seitsonen,

and D. Sundholm,

Κ.

Ρ.

E.

Phys. Rev. B 70, 035106 (2004),

Α.

and

617.

1985).

Hohenberg

Phys. Rev. 136, B864 (1964).

J. Chem. Phys. 68, 2767 (1978).

[7] N. H. Sabelli, R. Benedek,

Nieminen, M. J. Puska,

[10] K. Nordlund, N. Runeberg,

arXiv:0401265 [cond-mat].

Keinonen,

М.

M.

Phys. Rev. Lett. 67, 3692 (1991).

Nieminen,

Α.

Pruneda

Phys. Rev. A 20, 677 (1979).

[4] P.

[9] J.

[11] J.

R.

cho,

- I thank Kai Nordlund and Dage Sundholm for discussions on repulsive potentials. This work has been supported by the Academy of Finland (Suomen Akatemia) through project number 311149. Computational resources provided by CSC - It Center for Science Ltd (Espoo, Finland) and the Finnish Grid and Cloud Infrastructure (persistent identifier urn:nbn:fi:research-infras-2016072533) are gratefully acknowledged.
- [1] P. Sigmund, Particle Penetration and Radiation Effects Volup16 2M. Perkennetkenski, Rtadiat. dt. Molects & Olives 167, 666 (2012). Ν. Zinoviev Springer Series in Solid-State Sciences, Vol. 179 (Springer [17] A. and Κ. Nordlund. Cham Heidelberg New York Dordrecht London, 2014) p. Nucl. Instrum. Methods Phys. Res., Sect. B 406, 511 (2017).

Kohn,

- [18] S. Lehtola, Int. J. Quantum Chem. **119**, e25968 (2019),
- [2] M. A. Karolewski, Nucl. Instrum. Methods Phys. Res., Sect. B 256;X3541(02007)431.
 - [19] S. Lehtola, Int. J. Quantum Chem. 119, e25944 (2019), arXiv:1810.11653.
 - [20] S. Lehtola, M. Dimitrova, and D. Sundholm, Mol. Phys. doi: 10.1080/00268976.2019.1597989 (2019), arXiv:1812.06274.
 - [21] J. M. Herbert, in Rev. Comput. Chem., Vol. 28 (John Wiley & Sons, Inc., 2015) pp. 391–517.
 - [22] S. Lehtola, J. Chem. Phys. 151, 241102 (2019), arXiv:1911.10372.
 - [23] N. Η. F. Beebe J. Linderberg, and Int. J. Quant. Chem. 12, 683 (1977).
 - [24] S. D. Folkestad, E. F. Kjønstad, and H. Koch, J. Chem. Phys. 150, 194112 (2019), arXiv:1811.12890.
 - [25] X. Feng, E. Epifanovsky, J. Gauss, and A. I. Krylov, J. Chem. Phys. 151, 014110 (2019).
 - [26] F. Aquilante, L. Boman, J. Boström, H. Koch, R. Lindh, A. S. de Merás, and T. B. Pedersen, in Linear-Scaling Tech. Comput. Chem. Phys., Vol. 13 (Springer Netherlands, Dordrecht, 2011) pp. 301–343.
 - [27] F. Aquilante, T. B. Pedersen, A. Sánchez de Merás, and H. Koch, J. Chem. Phys. 125, 174101 (2006).
 - [28] O. Vahtras, J. Almlöf, and M. W. Feyereisen,
- [12] V. Kuzmin, Nucl. Instrum. Methods Phys. Res., Sect. B 249, 13 (2006). Phys. Lett. 213, 514 (1993).

Arta-

Nordlund,

Seitsonen.

- [14] V. Kuzmin, Surf. Coatings Technol. 201, 8388 (2007).
- [15] N. Juslin Κ. Nordlund, and J. Nucl. Mater. 382, 143 (2008).
- [13] M. A. Karolewski, Nucl. Instrum. Methods Phys. Res., Sect. [29]4B, 43 (2016)te, R. Lindh, and T. B. Pedersen, J. Chem. Phys. 127, 114107 (2007).
 - [30] F. Aquilante, L. Gagliardi, T. B. Pedersen, and R. Lindh, J. Chem. Phys. 130, 154107 (2009).

- [31] H. Harbrecht, M. Peters, and R. Schneider, Appl. Numer. Math. 62, 428 (2012).
- [32] J. Millam Ε. М. and G. Scuseria, J. Chem. Phys. 106, 5569 (1997).
- [33] M. Challacombe, J. Chem. Phys. 110, 2332 (1999).
- [34] Y. Shao, C. Saravanan, M. Head-Gordon, and C. A. White, J. Chem. Phys. 118, 6144 (2003).
- [35] J. Lehtola, M. Hakala, A. Sakko, and K. Hämäläinen, J. Comput. Chem. 33, 1572 (2012).
- [36] S. Lehtola, "ERKALE HF/DFT from Hel," (2018).
- [37] P.-O. Löwdin, Adv. Phys. 5, 1 (1956).
- [38] S. Lehtola, Int. J. Quantum Chem. **119**, e25945 (2019), arXiv:1810.11651.
- [39] S. Lehtola, "HelFEM Finite element methods for electronic studyetuPsyckköulations Resmallysystemen". 63, 45 (2012). (2018).
- [40] S. Lehtola, J. Chem. Theory Comput. 15, 1593 (2019), arXiv:1810.11659.
- [41] S. Lehtola, Phys. Rev. A 101, 012516 (2020), arXiv:1908.02528.

- [42] F. Jensen, J. Chem. Phys. 115, 9113 (2001).
- [43] E. V. R. de Castro and F. E. Jorge, J. Chem. Phys. 108, 5225 (1998).
- [44] S. Lehtola, (2020), arXiv:2001.04224.
- [45] F. Bloch, Zeitschrift für Phys. 57, 545 (1929).
- [46] P. A. M. Dirac, Math. Proc. Cambridge Philos. Soc. 26, 376 (1930). [47] S. H. Vosko, L. Wilk, and M. Nusair, Can. J. Phys. 58, 1200 (1980).
- [48] Note1, Kai Nordlund, private communication, 2020.
- [49] C. S. Becquart, M. F. Barthe, and A. De Backer, Phys. Scr. T145, 014048 (2011).
- [50] A. E. Sand, K. Nordlund, and S. L. Dudarev, J. Nucl. Mater. 455, 207 (2014).
- - [52] P. Pyykkö, Chem. Rev. **112**, 371 (2012).