

# Brouwer's fan theorem

Josef Berger

December 28, 2021

## Abstract

Brouwer's fan theorem states that every bar is a uniform bar. We give an overview of the status of this axiom in Bishop's constructive mathematics. In particular, we describe the relationship between the fan theorem, the weak König lemma, and the uniform continuity theorem.

*Keywords:* Brouwer's fan theorem, the weak König lemma, the uniform continuity theorem

2010 Mathematics Subject Classification: 03F60, 03A99, 54E45

## 1 Introduction

We analyse the relationship between Brouwer's fan theorem for detachable bars, the weak König lemma, and the uniform continuity theorem:

FAN Every detachable bar is a uniform bar.

WKL Every infinite tree has a path.

UC Every pointwise continuous function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is uniformly continuous.

Extending the well-known basic picture

$$\text{WKL} \Rightarrow \text{UC} \Rightarrow \text{FAN},$$

we prove the implications of the following diagram:

$$\begin{array}{ccccc}
\text{WKL} & \Rightarrow & \text{FAN} \wedge \text{DEFU} & \Leftrightarrow & \text{c-FAN} \Rightarrow \text{FAN} \\
& & \Downarrow & & \Downarrow \\
& & \text{FAN} \wedge \text{DECO} & \Leftrightarrow & \text{UC}
\end{array}$$

The axiom c-FAN was introduced in [2] as a version of the fan theorem which is equivalent to UC. The axioms DEFU and DECO are auxiliary statements clarifying the gap between FAN and c-FAN, UC respectively.

Mosts results mentioned in this article have been published before, see

- [9, 10] for results about WKL and FAN,
- [4, 5] for results about unique existence, and
- [1, 2, 8] for results about UC.

## 2 Notation

Let  $\mathbb{N} = \{0, 1, 2, \dots\}$  denote the set of natural numbers. We use the variables  $k, l, m, n, N, i, j$  for natural numbers. Let  $\{0, 1\}^*$  denote the set of all finite binary sequences, typically denoted by variables like  $u, v, w$ . The symbol  $\emptyset$  denotes the empty sequence. Note that  $\emptyset \in \{0, 1\}^*$ . Let  $\{0, 1\}^{\mathbb{N}}$  denote the set of all infinite binary sequences. We use the variables  $\alpha, \beta, \gamma$  for elements of  $\{0, 1\}^{\mathbb{N}}$ . The *length* of  $u$  is denoted by  $|u|$ . We define

$$\mathcal{N} = \{u \mid |u| > 0 \wedge \forall i < |u| (u_i = 0)\}$$

and

$$\mathcal{E} = \{u \mid |u| > 0 \wedge \forall i < |u| (u_i = 1)\}.$$

Let  $\{0, 1\}^n = \{u \mid |u| = n\}$  denote the set of all binary sequences of lenght  $n$ . The *concatenation* of  $u$  and  $v$  is denoted by  $u * v$  and defined by

$$(u_0, \dots, u_{n-1}) * (v_0, \dots, v_{m-1}) = (u_0, \dots, u_{n-1}, v_0, \dots, v_{m-1}).$$

The *concatenation* of  $u$  and  $\alpha$  is denoted by  $u * \alpha$  and defined by

$$(u_0, \dots, u_{n-1}) * \alpha = (u_0, \dots, u_{n-1}, \alpha_0, \alpha_1, \alpha_2, \dots).$$

If  $n \leq |w|$ , we denote the *restriction* of  $w$  to the first  $n$  elements by  $\overline{w}n$ . Note that  $\overline{w}0 = \emptyset$ . We denote the *restriction* of  $\alpha$  to the first  $n$  elements by  $\overline{\alpha}n$ . We write  $u < v$  for

$$|u| = |v| \wedge \exists k < |u| (\overline{u}k = \overline{v}k \wedge u_k = 0 \wedge v_k = 1).$$

Fix a subset  $A$  of  $\{0, 1\}^*$ . We define

- $A^\circ = \{u \mid \forall w (u * w \in A)\}$ ,
- $\overline{A} = \{u * w \mid u \in A \wedge w \in \{0, 1\}^*\}$ , and
- for any  $u$ ,  $A_u = \{w \mid u * w \in A\}$ .

A *path* of  $A$  is a sequence  $\alpha$  such that  $\forall n (\overline{\alpha}n \in A)$ . The set  $A$  has *at most one path* if

$$\forall \alpha, \beta (\exists n (\alpha_n \neq \beta_n) \Rightarrow \exists n (\overline{\alpha}n \notin A \vee \overline{\beta}n \notin A)).$$

A *longest path* of  $A$  is a sequence  $\alpha$  such that  $\forall u \in A (\overline{\alpha}|u| \in A)$ . Note that each path of  $A$  is a longest path of  $A$  but not necessarily vice versa.

The set  $A$  is said to be

- *detachable* (from  $\{0, 1\}^*$ ) if

$$\forall u (u \in A \vee u \notin A);$$

- a *c-set* if there exists a detachable set  $D \subseteq \{0, 1\}^*$  such that  $A = D^\circ$ ;
- *closed under extension* if

$$\forall u \in A \forall w (u * w \in A);$$

- *closed under restriction* if

$$\forall u, w (u * w \in A \Rightarrow u \in A);$$

- a *tree* if it is detachable and closed under restriction;
- *infinite* if

$$\forall n \exists u (|u| = n \wedge u \in A);$$

- a *bar* if

$$\forall \alpha \exists n (\overline{\alpha}n \in A);$$

- a *uniform bar* if

$$\exists N \forall \alpha \exists n \leq N (\overline{\alpha}n \in A);$$

- *convex* if

$$\forall u, w \in A \forall v (u < v < w \Rightarrow v \in A);$$

- *co-convex* if  $\{u \mid u \notin A\}$  is convex.

### 3 The weak König lemma

The *weak König lemma* is the following axiom:

WKL Every infinite tree has a path.

**Lemma 1.** *Let  $T$  be a tree. The following are equivalent:*

- (a)  *$T$  is an infinite tree.*
- (b) *Every longest path of  $T$  is a path of  $T$ .*

The notion of longest path is commonly used in graph theory. In the context of constructive mathematics, it was introduced in [10], along with the *longest path lemma*:

LPL Every tree has a longest path.

When working with trees and longest paths, it is convenient to have at hand an algorithm which transforms an arbitrary tree  $T$  into an infinite tree  $T'$ , where  $T'$  is as closely related to  $T$  as possible. Given a tree  $T$ , we define subsets  $L_T$  and  $T'$  of  $\{0, 1\}^*$  by

$$u \in L_T \Leftrightarrow (u \in T \wedge \forall w ((|u| < |w| \vee u < w) \Rightarrow w \notin T) \vee (\emptyset \notin T \wedge u = \emptyset))$$

and

$$u \in T' \Leftrightarrow u \in T \cup \{\emptyset\} \vee \exists v \in L_T \exists w \in \mathcal{N} (u = v * w).$$

Note that

- if  $T = \emptyset$  then  $L_T = \{\emptyset\}$  and  $T' = \mathcal{N} \cup \{\emptyset\}$ ;
- if  $T$  is infinite, then  $L_T = \emptyset$  and  $T' = T$ ;
- if  $\emptyset \in T$  and the set

$$\{|u| \mid u \in T\}$$

is bounded, then  $L_T = \{u\}$ , where  $u$  is an element of  $T$  of maximal length.

**Proposition 1.** *For every tree  $T$  there is an infinite tree  $T'$  such that*

- (a)  $T \subseteq T'$ ;
- (b) if  $T$  is infinite, then  $T = T'$ ;
- (c) if  $T$  is convex, then  $T'$  is convex as well;
- (d) if  $T$  is finite, then  $T'$  has at most one path;
- (e) every path of  $T'$  is a longest path of  $T$ ;
- (f) if  $T$  has at most one path, then  $T'$  has at most one path.

The lesser limited principle of omniscience is the following axiom:

LLPO For all  $\alpha$ ,

$$\forall n, m (\alpha_n = \alpha_m = 1 \Rightarrow n = m) \Rightarrow \forall n (\alpha_{2n} = 0) \vee \forall n (\alpha_{2n+1} = 0).$$

**Proposition 2.** LLPO  $\Rightarrow$  WKL

*Proof.* Let  $T$  be an infinite tree. Let  $\zeta_T(u, m)$  be defined by

$$\zeta_T(u, m) \Leftrightarrow \exists w \in \{0, 1\}^m (u * w \in T).$$

Moreover, set

$$Z_T = \{u \mid \forall m \zeta_T(u, m)\}.$$

Define  $\beta$  by

$$\beta_{2n} = 0 \Leftrightarrow \zeta_T((0), n)$$

and

$$\beta_{2n+1} = 0 \Leftrightarrow \zeta_T((1), n)$$

and define  $\alpha$  by

$$\alpha_n = 1 \Leftrightarrow \beta_n = 1 \wedge \forall m < n (\beta_m = 0).$$

By LLPO, either

$$\forall n (\alpha_{2n} = 0) \tag{1}$$

or else

$$\forall n (\alpha_{2n+1} = 0). \tag{2}$$

If (1) holds, we can conclude that

$$\forall n (\beta_{2n} = 0),$$

which implies that  $(0) \in Z_T$ .

Analogously, we can show that (2) implies that  $(1) \in Z_T$ .

Iterating this procedure yields a path  $\gamma$  of  $Z_T$ . Since  $Z_T \subseteq T$ , the sequence  $\gamma$  is also a path of  $T$ .

□

**Proposition 3.** *The following are equivalent:*

- (a) WKL
- (b) LPL
- (c) *Every convex tree has a longest path.*
- (d) *Every infinite convex tree has a path.*
- (e) LLPO

*Proof.* “(a)  $\Rightarrow$  (b)” This follows from Proposition 1.

“(c)  $\Rightarrow$  (d)” Let  $T$  be an infinite convex tree. By (c), the tree  $T$  has a longest path  $\alpha$ . By Lemma 1, the sequence  $\alpha$  is a path of  $T$ .

“(d)  $\Rightarrow$  (e)” Fix a sequence  $\alpha$  with

$$\forall n, m (\alpha_n = \alpha_m = 1 \Rightarrow n = m).$$

We define an infinite convex tree  $S$  by

$$S = \{\emptyset, (0), (1)\} \cup S_0 \cup S_1,$$

where

$$S_0 = \{(0) * w \mid w \in \mathcal{E} \wedge \forall i \leq |w| (\alpha_{2i} = 0)\},$$

$$S_1 = \{(1) * w \mid w \in \mathcal{N} \wedge \forall i \leq |w| (\alpha_{2i+1} = 0)\}.$$

Let  $\beta$  be a path of  $S$ . If  $\beta_1 = 0$ , then  $\forall i (\alpha_{2i} = 0)$ . If  $\beta_1 = 1$ , then  $\forall i (\alpha_{2i+1} = 0)$ .

“(e)  $\Rightarrow$  (a)” This is Proposition 2.

□

Proposition 2 and Proposition 3 (without the parts about convexity) were proved in [9] and [10].

## 4 The fan theorem

*Brouwer’s fan theorem* is the following axiom:

FAN Every detachable bar is a uniform bar.

**Lemma 2.** *If  $B \subseteq \{0, 1\}^*$  is closed under extension, the following are equivalent:*

(a)  $B$  is a uniform bar.

(b)  $\exists n (\{0, 1\}^n \subseteq B)$

The next lemma enables us to switch freely between a set  $B$  itself and  $\overline{B}$ , when discussing bar-related properties. This technique was introduced in [10].

**Lemma 3.** *Let  $B$  be a subset  $\{0, 1\}^*$ . Then*

(a)  $\overline{B}$  is closed under extension,

(b)  $B \subseteq \overline{B}$ ,

(c)  $B$  is detachable  $\Rightarrow \overline{B}$  is detachable,

(d)  $B$  is a bar  $\Rightarrow \overline{B}$  is a bar,

(e)  $\overline{B}$  is a uniform bar  $\Rightarrow B$  is a uniform bar.

**Proposition 4.** *The following are equivalent:*

(a) WKL!

(b) Each tree with at most one path has a longest path.

(c) Each detachable bar which is closed under extension is a uniform bar.

(d) FAN

*Proof.* “(a)  $\Rightarrow$  (b)” This follows from Proposition 1.

“(b)  $\Rightarrow$  (c)” Let  $B$  be a detachable bar which is closed under extension. Note that the set

$$T = \{u \mid u \notin B\}$$

is a tree. Since  $B$  is a bar, the tree  $T$  has at most one path. Let  $\alpha$  be a longest path of  $T$ . There exists  $n$  such that  $\overline{\alpha}n \in B$ . This implies that  $T \cap \{0, 1\}^n = \emptyset$  and therefore  $\{0, 1\}^n \subseteq B$ . Thus, by Lemma 2,  $B$  is a uniform bar.

“(c)  $\Rightarrow$  (d)” This follows from Lemma 3.

“(d)  $\Rightarrow$  (a)” Fix an infinite tree  $T$  with at most one path. Define a set  $B$  by

$$u \in B :\Leftrightarrow (0) * u \notin T \vee \forall v \in \{0, 1\}^{|u|} ((1) * v \notin T).$$

We show that the  $B$  is a bar. Fix a sequence  $\alpha$ . We define  $B_\alpha \subseteq \{0, 1\}^*$  by

$$v \in B_\alpha :\Leftrightarrow ((0) * \overline{\alpha}|v| \notin T \vee (1) * v \notin T).$$

Note that  $B_\alpha$  is closed under extension. Since  $T$  has a most one path,  $B_\alpha$  is a bar and therefore a uniform bar. Thus, there exists  $m$  such that  $\{0, 1\}^m \subseteq B_\alpha$ . Thus,  $\overline{\alpha}m \in B$ . This concludes the proof that  $B$  is a bar.

Since  $B$  is a uniform bar and closed under extension, there exists an  $n$  such that  $\{0, 1\}^n \subseteq B$ . We show that either

(i)  $\forall u \in \{0, 1\}^n ((0) * u \notin T)$  or else

(ii)  $\forall u \in \{0, 1\}^n ((1) * u \notin T)$ .

If (i) fails to hold, there exists  $u \in \{0, 1\}^n$  such that  $(0) * u \in T$ . Since  $u \in B$ , this implies (ii).

Let  $Z_T$  be defined as in the proof of Proposition 2. If (i) holds, then  $(1) \in Z_T$ . If (ii) holds, then  $(0) \in Z_T$ . Iterating this construction yields a path  $\gamma$  of  $Z_T$  and therefore of  $T$ .

□

The equivalence of FAN and WKL! was proved in [5].

**Corollary 1.**  $\text{WKL} \Rightarrow \text{FAN}$

Corollary 1 was originally published in [10].

**Lemma 4.** *If  $A \subseteq \{0, 1\}^*$  is co-convex, then  $\overline{A}$  is also co-convex.*

*Proof.* Fix  $u, v, w$  such that  $u < v < w$  and  $u, w \notin \overline{A}$ . We have to show that  $v \notin \overline{A}$ . To this end, fix  $m \leq |v|$ . Then  $\overline{u}m \notin A$  and  $\overline{w}m \notin A$ . The co-convexity of  $A$  yields  $\overline{v}m \notin A$ . Since  $m \leq |v|$  was chosen arbitrarily, this implies that  $v \notin \overline{A}$ .

□

**Proposition 5.**

- (a) *Every infinite convex tree with at most one path has a path.*
- (b) *Every convex tree with at most one path has a longest path.*
- (c) *Every detachable co-convex bar is a uniform bar.*

*Proof.* (a) Let  $T$  be an infinite convex tree with at most one path. Set

$$\alpha = (0, 1, 1, 1, \dots)$$

and

$$\beta = (1, 0, 0, 0, \dots).$$

Since  $T$  has at most one path, there exists  $n$  such that

$$\overline{\alpha}(n+1) \notin T \vee \overline{\beta}(n+1) \notin T.$$

Since  $T$  is convex, we can conclude that either

(i)  $\forall u \in \{0, 1\}^n ((0) * u \notin T)$  or else

(ii)  $\forall u \in \{0, 1\}^n ((1) * u \notin T)$ .

In the first case, choose  $\alpha_0 = 1$  and in the second case, choose  $\alpha_0 = 0$ . Iterating this construction leads to a path  $\alpha$  of  $T$ .

(b) follows from (a) and Proposition 1.

(c) Let  $B$  be a detachable, co-convex bar. By Lemma 4, we can assume that  $B$  is closed under extension. The set

$$T = \{u \mid u \notin B\}$$

is a convex tree. Since  $B$  is a bar, the tree  $T$  has at most one path. Let  $\alpha$  be a longest path of  $T$ . There exists  $n$  such that  $\bar{\alpha}n \in B$ . Then  $T \cap \{0, 1\}^n = \emptyset$ . Thus,  $\{0, 1\}^n \subseteq B$ . This implies that  $B$  is a uniform bar.  $\square$

The notion of co-convex bars was introduced in [7]. In [12], the fan theorem for co-convex bars—every detachable co-convex bar is a uniform bar—is used for a case study in program extraction from proofs. For a treatment of convex trees in constructive reverse mathematics see [6].

## 5 The uniform continuity theorem

Fix a function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  and  $u \in \{0, 1\}^*$ . We define

$$F(u) = F(u * \mathbf{0}),$$

where  $\mathbf{0} = (0, 0, 0, \dots)$ . Moreover, we define a new function  $F_u : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  by

$$F_u(\alpha) = F(u * \alpha).$$

For  $n \in \mathbb{N}$ , we define

$$F \equiv n \Leftrightarrow \forall \alpha (F(\alpha) = n).$$

A function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is

- *pointwise continuous* if

$$\forall \alpha \exists n \forall \beta (\bar{\alpha}n = \bar{\beta}n \Rightarrow F(\alpha) = F(\beta));$$

- *uniformly continuous* if

$$\exists n \forall \alpha, \beta (\overline{\alpha}n = \overline{\beta}n \Rightarrow F(\alpha) = F(\beta));$$

- *bounded* if

$$\exists N \forall \alpha (F(\alpha) \leq N);$$

- *constant* if

$$\exists n (F \equiv n).$$

**Lemma 5.** *If  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is uniformly continuous,*

$$F \equiv 0 \vee \neg (F \equiv 0).$$

**Lemma 6.** *Fix  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$ .*

(a)  *$F$  is pointwise continuous  $\Leftrightarrow \forall \alpha \exists n (F_{\overline{\alpha}n} \text{ is constant})$*

(b)  *$F$  is uniformly continuous  $\Leftrightarrow \exists N \forall u \in \{0, 1\}^N (F_u \text{ is constant})$*

**Lemma 7.** *Suppose that  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is pointwise continuous. Then the set*

$$\{u \mid F(u) \leq |u|\}$$

*is a bar.*

*Proof.* Fix  $\alpha$ . There exists  $N$  such that

$$\forall \beta (\overline{\alpha}N = \overline{\beta}N \Rightarrow F(\alpha) = F(\beta)).$$

We may assume that  $F(\alpha) \leq N$ . Note that

$$F(\overline{\alpha}N) = F(\alpha) \leq N.$$

This implies that

$$\overline{\alpha}N \in \{u \mid F(u) \leq |u|\}.$$

□

**Lemma 8.** *Every uniformly continuous function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is bounded.*

We now consider the *uniform continuity theorem*:

UC Every pointwise continuous function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  is uniformly continuous.

**Proposition 6.** UC  $\Rightarrow$  FAN

*Proof.* Let  $B$  be a detachable bar. The function

$$F(\alpha) = \min \{n \mid \bar{\alpha}n \in B\} \quad (3)$$

is pointwise continuous. In view of UC, the function  $F$  is uniformly continuous and therefore, by Lemma 8, bounded. Thus,  $B$  is a uniform bar.  $\square$

Fix functions  $F, M : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$ . The function  $M$  is a *modulus* for  $F$  if

- (a)  $M$  is pointwise continuous;
- (b)  $\forall \alpha, \beta, (\bar{\alpha}M(\alpha) = \bar{\beta}M(\alpha) \Rightarrow F(\alpha) = F(\beta))$ .

MUC Fix  $F, M : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$ . Assume that  $M$  is a pointwise continuous modulus of  $F$ . Then  $F$  is uniformly continuous.

**Proposition 7.** MUC  $\Leftrightarrow$  FAN

*Proof.* “ $\Rightarrow$ ” Let  $B$  be a detachable bar. Then the function  $F$ , given by (3), is a pointwise continuous modulus of itself, which implies that  $F$  is uniformly continuous and therefore bounded. Thus,  $B$  is a uniform bar.

“ $\Leftarrow$ ” Fix  $F, M : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  and assume that  $M$  is a pointwise continuous modulus of  $F$ . By Lemma 7, the set

$$B = \{u \mid M(u) \leq |u|\}$$

is a bar and therefore, in view of FAN, a uniform bar. Thus, there exists  $N$  such that

$$\forall \alpha \exists n \leq N (M(\bar{\alpha}n) \leq n).$$

This implies

$$\forall \alpha (F_{\bar{\alpha}N} \text{ is constant}).$$

By Lemma 6, we can conclude that  $F$  is uniformly continuous.  $\square$

The axiom MUC and Proposition 7 are taken from [1]. For more about uniform continuity and FAN see [8].

## 6 The fan theorem for c-sets

In the previous section, we have matched uniform continuity with the fan theorem by replacing UC with its weaker version MUC. Another option is to strengthen FAN by relaxing the requirement that the bar be detachable. This leads to the axiom c-FAN, which was introduced in [2].

**Lemma 9.** *For all subsets  $A$  of  $\{0, 1\}^*$  and all  $u$  we have*

$$(a) \ u \in A^\circ \Leftrightarrow A_u = \{0, 1\}^* \text{ and}$$

$$(b) \ (A_u)^\circ = (A^\circ)_u.$$

*Proof.* In order to prove (b), fix  $u, v$  and  $w$  and note that

$$\begin{aligned} w \in (A_u)^\circ &\Leftrightarrow \forall v (w * v \in A_u) \Leftrightarrow \\ \forall v (u * (w * v) \in A) &\Leftrightarrow \forall v ((u * w) * v \in A) \Leftrightarrow \\ u * w \in A^\circ &\Leftrightarrow w \in (A^\circ)_u. \end{aligned}$$

□

c-FAN Every bar which is a c-set is a uniform bar.

We introduce the auxiliary axioms DEFU (**d**ecidability whether certain bars are **full**) and DECO (**d**ecidability whether certain functions are **c**onstant).

DEFU Let  $D$  be a detachable subset of  $\{0, 1\}^*$  such that  $D^\circ$  is a bar. Then,

$$\exists u (u \notin D) \vee \neg \exists u (u \notin D).$$

DECO Let  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$  be pointwise continuous. Then,

$$\exists \alpha, \beta (F(\alpha) \neq F(\beta)) \vee \neg \exists \alpha, \beta (F(\alpha) \neq F(\beta)).$$

**Lemma 10.**  $\text{DEFU} \Leftrightarrow \text{DECO}$

*Proof.* “ $\Rightarrow$ ” Fix a pointwise continuous function  $F : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{N}$ . Set

$$D = \{u \mid F(u) = F(u * (1))\}.$$

By Lemma 6,  $D^\circ$  is a bar. Note that

$$\exists u (u \notin D) \Leftrightarrow \exists \alpha, \beta (F(\alpha) \neq F(\beta)).$$

“ $\Leftarrow$ ” Fix  $D \subseteq \{0, 1\}^*$  and suppose that  $D^\circ$  is a bar. Define

$$F(\alpha) = \min \{n \mid \forall m \geq n (\bar{\alpha}m \in D)\}$$

and note that

$$\exists u (u \notin D) \Leftrightarrow \exists \alpha (F(\alpha) > 0). \quad (4)$$

In presence of DECO, the right hand side of (4) is decidable, which implies that the left hand side of (4) is decidable.  $\square$

The diagram in the next proposition sheds some light on the relationship between bars and functions. Its core part is the equivalence between c-FAN and UC.

**Proposition 8.**

$$\begin{array}{ccc} \text{WKL} & \Rightarrow & \text{FAN} \wedge \text{DEFU} \Leftrightarrow \text{c-FAN} \Rightarrow \text{FAN} \\ & \Updownarrow & \Updownarrow \\ & \text{FAN} \wedge \text{DECO} \Leftrightarrow & \text{UC} \end{array}$$

*Proof.* “WKL  $\Rightarrow$  DEFU” Let  $D$  be a detachable subset  $D$  of  $\{0, 1\}^*$  such that  $D^\circ$  is a bar. Define an infinite tree  $T$  by

$$u \in T \Leftrightarrow \forall w (|w| \leq |u| \wedge w \notin D \Rightarrow \exists k \leq |w| (\bar{u}k \notin D)).$$

For each  $u \in T$  of length  $n$  we have

$$\{\bar{u}k \mid k \leq n\} \subseteq D \Leftrightarrow \{w \mid |w| \leq n\} \subseteq D.$$

Let  $\alpha$  be a path of  $T$ . The following are equivalent:

- (i)  $\exists u (u \notin D)$

(ii)  $\exists n (\bar{\alpha}n \notin D)$

There exists  $N$  such that  $\bar{\alpha}N \in D^\circ$ . Thus, (ii) is decidable, which yields the decidability of (i).

“FAN  $\wedge$  DEFU  $\Rightarrow$  c-FAN” Fix a detachable subset  $D$  of  $\{0, 1\}^*$  such that  $D^\circ$  is a bar. We show that  $D^\circ$  is detachable. To this end, fix some  $u$ . Note that  $(D^\circ)_u$  is a bar and therefore  $(D_u)^\circ$  is a bar. We can conclude that

$$D_u = \{0, 1\}^* \vee \exists w (w \notin D_u),$$

which implies

$$u \in D^\circ \vee u \notin D^\circ.$$

“FAN  $\wedge$  DECO  $\Rightarrow$  UC” Let  $F : \{0, 1\}^\mathbb{N} \rightarrow \mathbb{N}$  be pointwise continuous. Set

$$B = \{u \mid F_u \text{ is constant}\}.$$

In view of DECO, the set  $B$  is detachable. Since  $F$  is pointwise continuous,  $B$  is a bar. Thus,  $B$  is a uniform bar, which implies the uniform continuity of  $F$ .

“UC  $\Rightarrow$  FAN  $\wedge$  DECO” The implication UC  $\Rightarrow$  FAN holds by Proposition 6. The implication UC  $\Rightarrow$  DECO follows from Lemma 6.

“c-FAN  $\Rightarrow$  FAN” Let  $B$  be a detachable bar which is closed under extension. Then  $B = B^\circ$ , which implies that  $B$  is a c-set and therefore, in presence of c-FAN, a uniform bar.

□

Another equivalent of c-FAN is the so-called anti-Specker theorem; see [3]. For more on the fan theorem, see [11] and [13].

## References

- [1] Josef Berger. The Fan Theorem and Uniform Continuity. *New Computational Paradigms. CiE 2005. Lecture Notes in Computer Science*, 3526, 2005.

- [2] Josef Berger. The logical strength of the uniform continuity theorem. *Logical Approaches to Computational Barriers. CiE 2006. Lecture Notes in Computer Science*, 3988, 2006.
- [3] Josef Berger and Douglas Bridges. A fan-theoretic equivalent of the antithesis of Specker’s theorem. *Indagationes Mathematicae*, 18, 2007.
- [4] Josef Berger, Douglas Bridges, and Peter Schuster. The fan theorem and unique existence of maxima. *Journal of Symbolic Logic*, 71, 2006.
- [5] Josef Berger and Hajime Ishihara. Brouwer’s fan theorem and unique existence in constructive analysis. *Mathematical Logic Quarterly*, 51, 2005.
- [6] Josef Berger, Hajime Ishihara, Takayuki Kihara, and Takako Nemoto. The binary expansion and the intermediate value theorem in constructive reverse mathematics. *Archive for Mathematical Logic*, 58(1-2):203–217, 2019.
- [7] Josef Berger and Gregor Svindland. Brouwer’s fan theorem and convexity. *J. Symbolic Logic*, 83(4):1363–1375, 2018.
- [8] Douglas Bridges and Fred Richman. *Varieties of Constructive Mathematics*. Cambridge Univ. Press, 1987.
- [9] Hajime Ishihara. An omniscience principle, the König lemma and the Hahn-Banach theorem. *Z. Math. Logik Grundlagen Math.*, 36(2):237–240, 1990.
- [10] Hajime Ishihara. Weak König’s Lemma Implies Brouwer’s Fan Theorem: A Direct Proof. *Notre Dame Journal of Formal Logic*, 47(2):249–252, 2006.
- [11] Robert S. Lubarsky and Hannes Diener. Separating the fan theorem and its weakenings. *The Journal of Symbolic Logic*, 79(3):792–813, 2014.
- [12] Helmut Schwichtenberg. Program extraction from proofs: The fan theorem for uniformly coconvex bars. In *Mathesis Universalis, Computability and Proof*, pages 333–341. Springer International Publishing, 2019.

- [13] Wim Veldman. Brouwer's Fan Theorem as an axiom and as a contrast to Kleene's alternative. *Archive for Mathematical Logic*, 53(5-6):621–693, 2014.