Optimal Control for a Nonlinear Stochastic Parabolic Model of Language Competition

^{*a*} Sakine Esmaili^{*} ^{*a*} M. R. Eslahchi

^a Department of Applied Mathematics, Faculty of Mathematical Sciences, Tarbiat Modares University, P.O. Box 14115-134, Tehran, Iran

February 1, 2022

Abstract

In this investigation, an optimal control problem for a stochastic mathematical model of language competition is studied. We have considered the stochastic model of language competition by adding the stochastic terms to the deterministic model to take into account the random perturbations and uncertainties caused by the environment to have more reliable model. The model has formulated the population densities of the speakers of two languages, which are competing against each other to be saved from destruction, attract more speakers and so on, using two nonlinear stochastic parabolic equations. Four factors including the status of the languages and the growth rates of the populations are considered as the control variables (which can be controlled by the speakers of the populations or policy makers who make decisions for the populations for particular purposes) to control the evolution of the population densities. Then, the optimal control problem for the stochastic model of language competition is studied. Employing the tangent-normal cone techniques, the Ekeland variational principle and other theorems proved throughout the paper, we have shown there exists unique stochastic optimal control. We have also presented the exact form of the optimal control in terms of stochastic adjoint states.

Keywords: Optimal control, Nonlinear stochastic parabolic equation, Existence and uniqueness of stochastic optimal control, Language competition.

MSC: 49K20, 49J20, 35R60.

1 Introduction

There are different languages spoken by different populations, which are affected by a variety of factors consisting of growth rates of the populations, status of the languages and movement of the speakers, and may be in danger of extinction. It seems that there is a competition

^{*} E-mail addresses: sakine.esmaili@modares.ac.ir (Sakine Esmaili), eslahchi@modares.ac.ir (M. R. Eslahchi).

between the languages to attract more speakers, increase their population sizes and be saved from extinction, especially when different languages are spoken in the same area. For this reason, it is of great importance to study the evolution of endangered languages to be able to save them from extinction. Hence, the mathematical modelling of language competition is applied to study the evolution of speaker populations of languages [1-8], which leads us to predict the influence of different factors on the future of them and detect the efficient factors in saving at-risk languages. For instance, since India, with a 125 million English speakers, has the second largest number of English speakers in the world, the model of interaction between Monolingual Hindi speakers, Hindi/English Bilinguals and Hinglish speakers is studied in [1]. In another paper [2], the authors have investigated a society with two official languages consisting of A, spoken by every individual in the society, and B, spoken by a bilingual minority. In [3], time evolution of the density of speakers of an at-risk language, Aromanian, which is spoken by a bilingual community in North-Western Greece, is studied. In investigation [4], the language competition with game theoretic approach is analyzed when one language is known by all individuals and the other one is spoken by a minority. The author of [5], has also used game theory to contribute to the economics of language and offered some explanations of how past languages like Latin or Sanskrit can develop into a standard for literary production. In another investigation [6], the authors studied a mathematical model of language competition consisting of two coupled nonlinear parabolic equations. Each parabolic equation presents the population density of the speakers of one language, which is competing against another one to be saved from destruction. In this model, the effects of some factors including growth rates of the populations and status of the languages on the evolution of the population densities are modelled. Since the language with lower status or the population with lower growth rate may be destroyed, one may think that by changing these factors, we can save the languages. But in reality, every change can have negative effect on something else and can be so costly. Therefore, considering optimal control for the model is of great importance and helps us to adopt accurate policies for saving at-risk languages. Some other examples of competition between populations can be seen in [9, 10, 11].

Owing to the fact that all systems are affected by many uncertainties caused by the environment, considering the models in which the stochastic terms are included, enables us to deal with more trustable models. In this paper, we have studied a stochastic optimal bounded control strategy for a nonlinear stochastic parabolic model of language competition with a quadratic cost function. We have considered the mathematical model of language competition studied in [6] but we have added stochastic terms to the model to consider uncertainties and random perturbations and study more reliable model. In order to control the population densities of the speakers of two languages, our control strategy is considering some factors in the nonlinear stochastic parabolic model including the status of the languages and the growth rates of the populations to play role as the control variables, which have direct effects on the languages to attract more speakers. Since changing the control variables unthinkingly in order to save an at-risk language can be very costly and endanger another one, we have considered an optimal control problem for the model to minimize a quadratic cost function in which the cost of changing the values of status and growth rates are considered and our goals for population sizes are included. Designing this cost function helps us to make accurate and efficient decisions. For deriving the bounded optimal control variables, we have used some stochastic auxiliary equations (so-called adjoint equations). Employing the stochastic adjoint equations, some variational techniques and the tangent-normal cone techniques, the necessary conditions for the stochastic optimal control variables (which are stochastic because they are obtained in terms of the stochastic adjoint equations) are presented. After that, applying the Ekeland variational principle and adjoint equations, we have made a sequence converging to the exact optimal bounded control variables. Using the constructed sequence, the existence and uniqueness of the stochastic optimal control are proved. Since the control variables are assumed to be bounded, the obtained optimal control variables are continuous piecewise-defined functions of the stochastic adjoint states.

The organization of paper is as follows. In Section 2, the model and the optimal control problem are presented. The adjoint equations are introduced in Subsection 2.1. The existence and uniqueness of optimal control are studied in Subsection 2.2. Concluding remarks are offered in Section 3. Some lemmas and mathematical concepts, which are used throughout the paper are included in Appendix.

2 Model and Optimal Control Problem

In this section, first we introduce the stochastic mathematical model of language competition including two nonlinear stochastic parabolic equations describing the evolution of the population densities of the speakers of two languages. The stochastic model is obtained by adding stochastic terms to the model studied in [6] to have more reliable model by considering the random perturbations and uncertainties.

$$df_{1} = \nabla (D_{1}\nabla f_{1})dt - \nabla (F_{1}f_{1})dt + R_{1}(f_{1}, f_{2}, \beta_{1})dt + R(f_{1}, f_{2}, s_{1}, s_{2})dt + f_{1}\sum_{i=0}^{n}h_{i,1}(x, y)dB_{i},$$
(1)
$$(x, y, t) \in \Omega \times (0, \infty),$$

$$f_1(x, y, t) = f_1^b(x, y, t), \ (x, y) \in \partial\Omega, \ t > 0, \ f_1(x, y, 0) = f_1^0(x, y), \ (x, y) \in \Omega,$$
(2)

$$df_2 = \nabla (D_2 \nabla f_2) dt - \nabla (F_2 f_2) dt + R_2(f_1, f_2, \beta_2) dt - R(f_1, f_2, s_1, s_2) dt + f_2 \sum_{i=0}^{n} h_{i,2}(x, y) dB_i,$$
(3)

$$(x, y, t) \in \Omega \times (0, \infty),$$

$$f_2(x, y, t) = f_2^b(x, y, t), \ (x, y) \in \partial\Omega, \ t > 0, \ f_2(x, y, 0) = f_2^0(x, y), \ (x, y) \in \Omega,$$
(4)

$$R_1(f_1, f_2, \beta_1) = \beta_1 f_1(1 - \frac{f_1 + f_2}{N}), \quad R_2(f_1, f_2, \beta_2) = \beta_2 f_2(1 - \frac{f_1 + f_2}{N}),$$
$$R(f_1, f_2, s_1, s_2) = k(s_1 f_1^{\alpha} f_2 - s_2 f_1 f_2^{\alpha}),$$

where $B(t) = (B_1(t), \dots, B_n(t))$ is an *n*-dimensional Brownian motion on $(\Omega, \mathfrak{F}, \mathbb{P})$ with associated filtration $\{\mathfrak{F}(t)\}_{t>0}$, $f_i(x, y, t)$ describes the population density of speakers of language i, α is a positive constant, $s_i(t)$ represents the status of language i at time t and the logistic terms with carrying capacity N and Malthus rates β_1 and β_2 take into account the growth of the populations. The population movement is described by the advection terms with the external force fields $F_1(x, y) = (F_1^x(x, y), F_1^y(x, y))$ and $F_2(x, y) = (F_2^x(x, y), F_2^y(x, y))$ and by the diffusion terms with the diffusion coefficients D_1 and D_2 .

In this model, we have assumed that the following properties are satisfied:

A. $\Omega \subset \mathbb{R}^2$ is an open, bounded set with boundary $\partial \Omega \in C^2$.

B. There exist C^2 -smooth functions $u_1(x, y, t)$ and $u_2(x, y, t)$ such that $u_i(x, y, t) = f_i^b(x, y, t)$ (i = 1, 2) on $\partial\Omega$ and $u_i(x, y, 0) = f_i^0(x, y)$ (i = 1, 2) on Ω .

C. F_1 and F_2 are C^1 -smooth functions, $h_{i,j}$ $(i = 0, \dots, n, j = 1, 2)$ are C^2 -smooth functions and $h_{i,j}(x, y) = 0$ on $\partial\Omega$.

In the following theorem the existence and uniqueness of the solution of (1)-(4) are presented.

Theorem 1. Let assumptions $\mathbf{A}-\mathbf{C}$ be satisfied, then the problem (1)–(4) for almost every $\omega \in \mathbf{\Omega}$ has a unique solution. For every T > 0, f_1 and f_2 are $\mathfrak{F}(t)$ -adapted and for almost every $\omega \in \mathbf{\Omega}$, we also have

$$f_1, f_2 \in C\left(\Omega \times [0, T]\right)$$

Proof Using Lemma 3 (see Appendix) and fixed point theorems, we can prove this theorem. \Box Now, we consider the optimal control for the presented model of language competition. In this language competition optimal control problem (LCOCP), we purpose to control the population densities of the speakers by considering some time dependant control variables β_1 , β_2 , s_1 , and s_2 , which control the growth of the populations and the status of the languages. Then, we minimize

$$J(\beta_1, \beta_2, s_1, s_2) := \int_{\Omega} \int_0^T ((\beta_1 - b_1)^2 + (\beta_2 - b_2)^2 + \lambda_1 (f_1 - r_1^*)^2 + \lambda_2 (f_2 - r_2^*)^2) dt dx dy + \int_{\Omega} \int_0^T (s_1 - r_3^*)^2 + (s_2 - r_4^*)^2 dt dx dy,$$

$$(5)$$

$$(\beta_1, \beta_2, s_1, s_2) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1,$$

where $r_1^*, r_2^*, r_3^*, r_4^*, b_1, b_2 \in C^{\alpha, \alpha/2}(\overline{\Omega} \times [0, T])$ for some $0 < \alpha < 1$, also

$$\begin{aligned} \mathcal{C}_{ad} &:= \left\{ v \in C(\overline{\Omega} \times [0,T]) \ a.s.: \ v \text{ is } \mathfrak{F}(t) - \text{adapted}, \ l_{1}^{*}(x,y,t) \leq v(x,y,t) \leq l_{1}^{**}(x,y,t) \ a.s. \right\}, \\ (6) \\ \mathcal{W}_{ad} &:= \left\{ v \in C(\overline{\Omega} \times [0,T]) \ a.s.: \ v \text{ is } \mathfrak{F}(t) - \text{adapted}, \ l_{2}^{*}(x,y,t) \leq v(x,y,t) \leq l_{2}^{**}(x,y,t) \ a.s. \right\}, \\ (7) \\ \mathcal{C}_{ad}^{1} &:= \left\{ v \in C(\overline{\Omega} \times [0,T]) \ a.s.: \ v \text{ is } \mathfrak{F}(t) - \text{adapted}, \ l_{3}^{*}(x,y,t) \leq v(x,y,t) \leq l_{3}^{**}(x,y,t) \ a.s. \right\}, \\ (8) \\ \mathcal{W}_{ad}^{1} &:= \left\{ v \in C(\overline{\Omega} \times [0,T]) \ a.s.: \ v \text{ is } \mathfrak{F}(t) - \text{adapted}, \ l_{4}^{*}(x,y,t) \leq v(x,y,t) \leq l_{4}^{**}(x,y,t) \ a.s. \right\}, \\ (9) \\ l_{i}^{*}, l_{i}^{**} \in C^{\alpha, \alpha/2}(\overline{\Omega} \times [0,T]), \ i = 1, 2, 3, 4, \end{aligned}$$

such that the equations (1)-(4) are satisfied.

Adjoint Equations 2.1

Before presenting our main results, we introduce the following stochastic adjoint equations, which are instrumental in presenting the necessary conditions and proving the existence and uniqueness of stochastic optimal control variables. The optimal control variables are stochastic because they will obtained in terms of the stochastic adjoint states.

$$dz_{f_1} = -\nabla (D_1 \nabla z_{f_1}) dt + H_1^* \underbrace{(z_{f_1}, z_{f_2}, f_1, f_2, \beta_1, \beta_2, s_1, s_2)}_{Z^*} dt - z_{f_1} \sum_{i=0}^n h_{i,1}(x, y) dB_i,$$
(10)

$$z_{f_1}(x, y, t) = 0, \ (x, y) \in \partial\Omega, \ 0 < t < T, \ z_{f_1}(x, y, T) = 0, \ (x, y) \in \Omega,$$
(11)

$$dz_{f_2} = -\nabla (D_2 \nabla z_{f_2}) dt + H_2^* (Z^*) dt - z_{f_2} \sum_{i=0}^n h_{i,2}(x, y) dB_i, \ (x, y, t) \in \Omega \times (0, T),$$
(12)

$$z_{f_2}(x, y, t) = 0, \ (x, y) \in \partial\Omega, \ 0 < t < T, \ z_{f_2}(x, y, T) = 0, \ (x, y) \in \Omega,$$
(13)

$$dz_{g_1} = G_1^* (Z^*) dt, \ dz_{g_2} = G_2^* (Z^*) dt, \ (x, y) \in \Omega, \ 0 < t < T,$$
(14)

$$dz_{g_3} = G_3^* \left(Z^* \right) dt, \ dz_{g_4} = G_4^* \left(Z^* \right) dt, \ \left(x, y \right) \in \Omega, \ 0 < t < T,$$

$$(15)$$

$$z_{g_i}(x, y, T) = 0, \ (x, y) \in \Omega, \ i = 1, 2, 3, 4,$$
(16)

where

$$H_{1}^{*ne} = -\frac{\partial \left(R_{1}(f_{1}, f_{2}, \beta_{1}) + R(f_{1}, f_{2}, s_{1}, s_{2})\right)}{\partial f_{1}} z_{f_{1}} - \frac{\partial \left(R_{2}(f_{1}, f_{2}, \beta_{2}) - R(f_{1}, f_{2}, s_{1}, s_{2})\right)}{\partial f_{1}} z_{f_{2}} - \nabla (z_{f_{1}}) \cdot F_{1} + \lambda_{1}(f_{1} - r_{1}^{*}) + z_{f_{1}} \sum_{i=0}^{n} (h_{i,1}(x, y))^{2}, \\H_{2}^{*}(Z^{*}) = -\frac{\partial \left(R_{1}(f_{1}, f_{2}, \beta_{1}) + R(f_{1}, f_{2}, s_{1}, s_{2})\right)}{\partial f_{2}} z_{f_{1}} - \frac{\partial \left(R_{2}(f_{1}, f_{2}, \beta_{2}) - R(f_{1}, f_{2}, s_{1}, s_{2})\right)}{\partial f_{2}} z_{f_{2}} - \nabla (z_{f_{2}}) \cdot F_{2} + \lambda_{2}(f_{2} - r_{2}^{*}) + z_{f_{2}} \sum_{i=0}^{n} (h_{i,2}(x, y))^{2}, \\G_{1}^{*}(Z^{*}) = -\frac{\partial R_{1}(f_{1}, f_{2}, \beta_{1})}{\partial \beta_{1}} z_{f_{1}}, \qquad (17)$$

/

$$G_2^*(Z^*) = -\frac{\partial R_2(f_1, f_2, \beta_2)}{\partial \beta_2} z_{f_2},$$
(18)

$$G_3^*(Z^*) = -\frac{\partial R(f_1, f_2, s_1, s_2)}{\partial s_1} z_{f_1} + \frac{\partial R(f_1, f_2, s_1, s_2)}{\partial s_1} z_{f_2},$$
(19)

$$G_4^*(Z^*) = -\frac{\partial R(f_1, f_2, s_1, s_2)}{\partial s_2} z_{f_1} + \frac{\partial R(f_1, f_2, s_1, s_2)}{\partial s_2} z_{f_2}.$$
 (20)

In the following theorem, the existence and uniqueness of the solution of the system of stochastic adjoint equations are presented.

Theorem 2. Let assumptions $\mathbf{A}-\mathbf{C}$ be satisfied, then for almost every $\omega \in \Omega$, the adjoint system (10)–(16) has a unique solution $(z_{f_1}, z_{f_2}, z_{g_1}, z_{g_2}, z_{g_3}, z_{g_4})$ and for every T > 0, we have $z_{f_1}, z_{f_2}, z_{g_1}, z_{g_2}, z_{g_3}$ and z_{g_4} are $\mathfrak{F}(t)$ -adapted and

$$z_{f_1}, z_{f_2}, z_{g_1}, z_{g_2}, z_{g_3}, z_{g_4} \in C(\Omega \times [0, T]).$$
 a.s.

Proof Employing Lemma 3 (see Appendix) and Theorem 1, we obtain the result.

2.2 Existence and Uniqueness of Optimal Control

In this subsection, we present the necessary conditions for optimal control, which give us the explicit forms of stochastic optimal control variables in terms of the stochastic adjoint states, also we prove the existence and uniqueness of optimal control variables.

In the next theorem, we introduce the necessary conditions for optimal control variables of LCOCP.

Theorem 3. (Necessary Conditions) Let $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$ be the optimal control of LCOCP, then for almost every $\omega \in \Omega$, we have

$$\beta_1^* = \mathcal{F}\Big(b_1 - G_1^*(Z^*), l_1^*, l_1^{**}\Big), \ \beta_2^* = \mathcal{F}\Big(b_2 - G_2^*(Z^*), l_2^*, l_2^{**}\Big),$$
(21)

$$s_1^* = \mathcal{F}\left(r_3^* - G_3^*\left(Z^*\right), l_3^*, l_3^{**}\right), \ s_2^* = \mathcal{F}\left(r_4^* - G_4^*\left(Z^*\right), l_4^*, l_4^{**}\right),$$
(22)

where $(z_{f_1}, z_{f_2}, f_1, f_2)$ is the exact solution of (1)-(4) and (10)-(13) corresponding to $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$, $\beta_1^*, \beta_2^*, s_1^*$ and s_2^* are $\mathfrak{F}(t)$ -adapted, l_i^* and l_i^{**} (i = 1, 2, 3, 4) are defined in (6)-(9), and

$$\mathcal{F}(x, u, v) = \begin{cases} u, & x < u, \\ x, & u \le x \le v, \\ v, & v < x. \end{cases}$$
(23)

Proof Let (f_1^*, f_2^*) and $(f_1^{\epsilon}, f_2^{\epsilon})$ be the solutions of (1)–(4) corresponding to the controls

$$(\beta_1^*, \beta_2^*, s_1^*, s_2^*) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1,$$

and

$$(\beta_1^{\epsilon}, \beta_2^{\epsilon}, s_1^{\epsilon}, s_2^{\epsilon}) = (\beta_1^* + \epsilon \beta_1^0, \beta_2^* + \epsilon \beta_2^0, s_1^* + \epsilon s_1^0, s_2^* + \epsilon s_2^0) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1,$$

respectively, where C_{ad} , W_{ad} , C_{ad}^1 and W_{ad}^1 are defined in (6)–(9) and ϵ is a positive constant. Also, assume that

$$J(\beta_1^*, \beta_2^*, s_1^*, s_2^*) = \min\{J(\beta_1, \beta_2, s_1, s_2) : (\beta_1, \beta_2, s_1, s_2) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1\},$$
$$f_{1,1}^{\epsilon} = \frac{f_1^{\epsilon} - f_1^*}{\epsilon}, \ f_{2,1}^{\epsilon} = \frac{f_2^{\epsilon} - f_2^*}{\epsilon}.$$

Therefore, one can derive that

$$\frac{J(\beta_1^\epsilon,\beta_2^\epsilon,s_1^\epsilon,s_2^\epsilon)-J(\beta_1^*,\beta_2^*,s_1^*,s_2^*)}{\epsilon} =$$

$$\int_{\Omega} \int_{0}^{T} ((\beta_{1}^{*} + \beta_{1}^{\epsilon} - 2b_{1})\beta_{1}^{0} + (\beta_{2}^{*} + \beta_{2}^{\epsilon} - 2b_{2})\beta_{2}^{0} + \lambda_{1}(f_{1}^{*} + f_{1}^{\epsilon} - 2r_{1}^{*})f_{1,1}^{\epsilon} + \lambda_{2}(f_{2}^{*} + f_{2}^{\epsilon} - 2r_{2}^{*})f_{2,1}^{\epsilon})dtdxdy + \int_{\Omega} \int_{0}^{T} (s_{1}^{*} + s_{1}^{\epsilon} - 2r_{3}^{*})s_{1}^{0} + (s_{2}^{*} + s_{2}^{\epsilon} - 2r_{4}^{*})s_{2}^{0}dtdxdy \ge 0, \quad a.s.$$

$$(24)$$

and

$$df_{1,1}^{\epsilon} = \nabla .(D_1 \nabla f_{1,1}^{\epsilon})dt + E_1^{\epsilon}(\underbrace{f_1^*, f_2^*, \beta_1^*, \beta_2^*, s_1^*, s_2^*, f_1^{\epsilon}, f_2^{\epsilon}, \beta_1^{\epsilon}, \beta_2^{\epsilon}, s_1^{\epsilon}, s_2^{\epsilon}}_{Z^{*,\epsilon}})dt + f_{1,1}^{\epsilon}\sum_{i=0}^n h_{i,1}dB_i, \quad (25)$$

$$(x, y, t) \in \Omega \times (0, \infty),$$

$$f_{1,1}^{\epsilon}(x,y,t) = 0, \ (x,y) \in \partial\Omega, \ t > 0, \ f_{1,1}^{\epsilon}(x,y,0) = 0, \ (x,y) \in \Omega,$$
(26)

$$df_{2,1}^{\epsilon} = \nabla (D_2 \nabla f_{2,1}^{\epsilon}) dt + E_2^{\epsilon} (Z^{*,\epsilon}) dt + f_{2,1}^{\epsilon} \sum_{i=0}^n h_{i,2} dB_i, \ (x, y, t) \in \Omega \times (0, \infty),$$
(27)

$$f_{2,1}^{\epsilon}(x,y,t) = 0, \ (x,y) \in \partial\Omega, \ t > 0, \ f_{2,1}^{\epsilon}(x,y,0) = 0, \ (x,y) \in \Omega,$$
(28)

where

$$E_1^{\epsilon}(Z^{*,\epsilon}) = -\nabla \cdot (F_1 f_{1,1}^{\epsilon}) + \frac{R_1(f_1^{\epsilon}, f_2^{\epsilon}, \beta_1^{\epsilon}) + R(f_1^{\epsilon}, f_2^{\epsilon}, s_1^{\epsilon}, s_2^{\epsilon}) - \left(R_1(f_1^{*}, f_2^{*}, \beta_1^{*}) + R(f_1^{*}, f_2^{*}, s_1^{*}, s_2^{*})\right)}{\epsilon},$$

$$E_{2}^{\epsilon}(Z^{*,\epsilon}) = -\nabla \cdot (F_{2}f_{2,1}^{\epsilon}) + \frac{R_{2}(f_{1}^{\epsilon}, f_{2}^{\epsilon}, \beta_{2}^{\epsilon}) - R(f_{1}^{\epsilon}, f_{2}^{\epsilon}, s_{1}^{\epsilon}, s_{2}^{\epsilon}) - \left(R_{2}(f_{1}^{*}, f_{2}^{*}, \beta_{2}^{*}) - R(f_{1}^{*}, f_{2}^{*}, s_{1}^{*}, s_{2}^{*})\right)}{\epsilon}.$$

Therefore using (10)–(16) and Theorem 9 (see Appendix), one can deduce that

$$\int_{0}^{T} \int_{\Omega} d(f_{1,1}^{\epsilon} z_{f_{1}}) + d(f_{2,1}^{\epsilon} z_{f_{2}}) + \int_{0}^{T} \int_{\Omega} \beta_{1}^{0} dz_{g_{1}} + \beta_{2}^{0} dz_{g_{2}} + s_{1}^{0} dz_{g_{3}} + s_{2}^{0} dz_{g_{4}} dx dy = \int_{0}^{T} \int_{\Omega} \left(f_{1,1}^{\epsilon} dz_{f_{1}} + z_{f_{1}} df_{1,1}^{\epsilon} + dz_{f_{1}} df_{1,1}^{\epsilon} + f_{2,1}^{\epsilon} dz_{f_{2}} + z_{f_{2}} df_{2,1}^{\epsilon} + dz_{f_{2}} df_{2,1}^{\epsilon} \right) +$$

$$\int_0^T \int_\Omega \left(\beta_1^0 dz_{g_1} + \beta_2^0 dz_{g_2} + s_1^0 dz_{g_3} + s_2^0 dz_{g_4} \right) dx dy = \int_0^T \int_\Omega \left(\lambda_1 (f_1^* - r_1^*) f_{1,1}^\epsilon + \lambda_2 (f_2^* - r_2^*) f_{2,1}^\epsilon \right) dx dy dt + \varpi(\epsilon, T),$$

where $(z_{f_1}, z_{f_2}, z_{g_1}, z_{g_2}, z_{g_3}, z_{g_4})$ is the exact solution of the adjoint system (10)–(16) corresponding to $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$ and $\lim_{\epsilon \to 0} \overline{\omega}(\epsilon, T) = 0$, thus we have

$$\int_0^T \int_\Omega \left(\lambda_1 (f_1^* - r_1^*) f_{1,1}^\epsilon + \lambda_2 (f_2^* - r_2^*) f_{2,1}^\epsilon - \beta_1^0 \frac{\partial z_{g_1}}{\partial t} - \beta_2^0 \frac{\partial z_{g_2}}{\partial t} - s_1^0 \frac{\partial z_{g_3}}{\partial t} - s_2^0 \frac{\partial z_{g_4}}{\partial t} \right) dx dy dt = -\varpi(\epsilon, T)$$

$$\tag{29}$$

Hence, using (24) and (29), we arrive at

$$\int_{\Omega} \int_{0}^{T} (\beta_{1}^{*} - b_{1} + \frac{\partial z_{g_{1}}}{\partial t}) \beta_{1}^{0} + (\beta_{2}^{*} - b_{2} + \frac{\partial z_{g_{2}}}{\partial t}) \beta_{2}^{0} dt dx dy + \\ \int_{\Omega} \int_{0}^{T} (s_{1}^{*} - r_{3}^{*} + \frac{\partial z_{g_{3}}}{\partial t}) s_{1}^{0} + (s_{2}^{*} - r_{4}^{*} + \frac{\partial z_{g_{4}}}{\partial t}) s_{2}^{0} dt dx dy \geq 0, \quad a.s.$$
(30)

Therefore using tangent-normal cone techniques [19] and (6)–(9), $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$ is as follows

$$\beta_1^* = \mathcal{F}\left(b_1 - \frac{\partial z_{g_1}}{\partial t}, l_1^*, l_1^{**}\right), \ \beta_2^* = \mathcal{F}\left(b_2 - \frac{\partial z_{g_2}}{\partial t}, l_2^*, l_2^{**}\right), \ a.s.$$
(31)

$$s_{1}^{*} = \mathcal{F}\left(r_{3}^{*} - \frac{\partial z_{g_{3}}}{\partial t}, l_{3}^{*}, l_{3}^{**}\right), \ s_{2}^{*} = \mathcal{F}\left(r_{4}^{*} - \frac{\partial z_{g_{4}}}{\partial t}, l_{4}^{*}, l_{4}^{**}\right). \ a.s.$$
(32)

Also from (31)–(32), we can conclude that β_1^* , β_2^* , s_1^* and s_2^* are $\mathfrak{F}(t)$ -adapted. \Box The following theorem gives the existence and uniqueness of the optimal control variables for LCOCP.

Theorem 4. (Existence and Uniqueness) *LCOCP*, for almost every $\omega \in \Omega$, has a unique optimal control, which satisfies (21)–(22).

Proof Using Theorem 9 (see Appendix), we conclude that for each

$$(\beta_1, \beta_2, s_1, s_2) \in \mathcal{C}_{ad}^* \times \mathcal{W}_{ad}^* \times \mathcal{C}_{ad}^{1*} \times \mathcal{W}_{ad}^{1*} \setminus C_4(\underbrace{\Omega \times [0, T]}_{\overline{Q_T}}),$$

where

$$C_4(\overline{Q_T}) = C(\overline{Q_T}) \times C(\overline{Q_T}) \times C(\overline{Q_T}) \times C(\overline{Q_T}),$$

$$\mathcal{C}^*_{ad} := \left\{ v \in L^{\infty}(\overline{Q_T}) \quad a.s. : l_1^*(x, y, t) \leq v(x, y, t) \leq l_1^{**}(x, y, t), \quad a.e. \quad a.s. \right\},$$

$$\mathcal{W}^*_{ad} := \left\{ v \in L^{\infty}(\overline{Q_T}) \quad a.s. : l_2^*(x, y, t) \leq v(x, y, t) \leq l_2^{**}(x, y, t), \quad a.e. \quad a.s. \right\},$$

$$\mathcal{C}^{1*}_{ad} := \left\{ v \in L^{\infty}(\overline{Q_T}) \quad a.s. : l_3^*(x, y, t) \leq v(x, y, t) \leq l_3^{**}(x, y, t), \quad a.e. \quad a.s. \right\},$$

$$\mathcal{W}_{ad}^{1*} := \Big\{ v \in L^{\infty}(\overline{Q_T}) \ a.s. : l_4^*(x, y, t) \le v(x, y, t) \le l_4^{**}(x, y, t), \ a.e. \ a.s. \Big\},$$

there exists a unique value $J^*(\beta_1, \beta_2, s_1, s_2)$ such that for every sequence $\left\{ (\beta_1^n, \beta_2^n, s_1^n, s_2^n) \right\}_{n=0}^{\infty}$ in $\mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1$ converging to $(\beta_1, \beta_2, s_1, s_2)$ in

$$L_4^2(\overline{Q_T}) := L^2(\overline{Q_T}) \times L^2(\overline{Q_T}) \times L^2(\overline{Q_T}) \times L^2(\overline{Q_T}),$$

we have $J(\beta_1^n, \beta_2^n, s_1^n, s_2^n) \to J^*(\beta_1, \beta_2, s_1, s_2)$. Using the Lebesgue Dominated Convergence Theorem [20], we deduce that the function

$$\mathcal{J}(\beta_{1},\beta_{2},s_{1},s_{2}) = \begin{cases} J(\beta_{1},\beta_{2},s_{1},s_{2}), & (\beta_{1},\beta_{2},s_{1},s_{2}) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^{1} \times \mathcal{W}_{ad}^{1}, \\ J^{*}(\beta_{1},\beta_{2},s_{1},s_{2}), & (\beta_{1},\beta_{2},s_{1},s_{2}) \in \mathcal{C}_{ad}^{*} \times \mathcal{W}_{ad}^{*} \times \mathcal{C}_{ad}^{1*} \times \mathcal{W}_{ad}^{1*} \setminus C_{4}(\overline{Q_{T}}), \\ +\infty, & (\beta_{1},\beta_{2},s_{1},s_{2}) \in (\mathcal{C}_{ad}^{*} \times \mathcal{W}_{ad}^{*} \times \mathcal{C}_{ad}^{1*} \times \mathcal{W}_{ad}^{1*})^{c} \cap L_{4}^{1}(\overline{Q_{T}}), \end{cases}$$
(33)

is lower semicontinuous with respect to $(\beta_1, \beta_2, s_1, s_2)$ in $L^1_4(\overline{Q_T})$. Thus, from Theorem 8 (see Appendix), we conclude that for each positive ϵ , there exists $(\beta_1^{\epsilon}, \beta_2^{\epsilon}, s_1^{\epsilon}, s_2^{\epsilon})$ such that

$$\mathcal{J}(\beta_{1}^{\epsilon}, \beta_{2}^{\epsilon}, s_{1}^{\epsilon}, s_{2}^{\epsilon}) < \mathcal{J}(\beta_{1}, \beta_{2}, s_{1}, s_{2}) + \epsilon^{\frac{1}{2}} \Big(\|\beta_{1} - \beta_{1}^{\epsilon}\|_{L^{1}(\overline{Q_{T}})} + \|\beta_{2} - \beta_{2}^{\epsilon}\|_{L^{1}(\overline{Q_{T}})} + \|s_{1} - s_{1}^{\epsilon}\|_{L^{1}(\overline{Q_{T}})} + \|s_{2} - s_{2}^{\epsilon}\|_{L^{1}(\overline{Q_{T}})} \Big), \qquad (34)$$
$$\forall (\beta_{1}, \beta_{2}, s_{1}, s_{2}) \neq (\beta_{1}^{\epsilon}, \beta_{2}^{\epsilon}, s_{1}^{\epsilon}, s_{2}^{\epsilon}).$$

In addition, each sequence $\left\{ (\beta_1^n, \beta_2^n, s_1^n, s_2^n) \right\}_{n=0}^{\infty}$ in $\mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1$ converging to $(\beta_1^{\epsilon}, \beta_2^{\epsilon}, s_1^{\epsilon}, s_2^{\epsilon})$ in $L_4^2(\overline{Q_T})$ is a Cauchy sequence in $L_4^2(\overline{Q_T})$. If we assume that for each $n \in \mathbb{N}_0$, (f_1^n, f_2^n) is the solution of (1)–(4) corresponding to $(\beta_1^n, \beta_2^n, s_1^n, s_2^n)$, according to Theorem 9, the sequence $\left\{ (f_1^n, f_2^n) \right\}_{n=0}^{\infty}$ is a Cauchy sequence in $L_2^2(\overline{Q_T})$. So, using the Lebesgue Dominated Convergence Theorem [20], we derive that for each p > 5, there exist subsequences $\left\{ (\beta_1^{n_k}, \beta_2^{n_k}, s_1^{n_k}, s_2^{n_k}) \right\}_{k=0}^{\infty}$ and $\left\{ (f_1^{n_k}, f_2^{n_k}) \right\}_{k=0}^{\infty}$, which are Cauchy in

$$L_4^p(\overline{Q_T}) := L^p(\overline{Q_T}) \times L^p(\overline{Q_T}) \times L^p(\overline{Q_T}) \times L^p(\overline{Q_T}),$$

and

$$L_2^p(\overline{Q_T}) := L^p(\overline{Q_T}) \times L^p(\overline{Q_T}),$$

respectively. Thus, using Lemma 2 (see Appendix), it is easy to see that the sequences

$$\left\{ \left(\mathfrak{f}_{1}^{n_{k}}, \mathfrak{f}_{2}^{n_{k}} \right) \right\}_{k=0}^{\infty} = \left\{ \left(e^{-\sum_{i=0}^{n} h_{i,1}B_{i}} f_{1}^{n_{k}}, e^{-\sum_{i=0}^{n} h_{i,2}B_{i}} f_{2}^{n_{k}} \right) \right\}_{k=0}^{\infty},$$

and

$$\left\{ (\mathfrak{z}_{f_1^{n_k}},\mathfrak{z}_{f_2^{n_k}}) \right\}_{k=0}^{\infty} = \left\{ (e^{\sum_{i=0}^n h_{i,1}B_i} z_{f_1^{n_k}}, e^{\sum_{i=0}^n h_{i,2}B_i} z_{f_2^{n_k}}) \right\}_{k=0}^{\infty},$$

are Cauchy in

$$W_p^{2,1,2}(\overline{Q_T}) := W_p^{2,1}(\overline{Q_T}) \times W_p^{2,1}(\overline{Q_T}),$$

where $(z_{f_1^{n_k}}, z_{f_2^{n_k}})$ is the solution of (10)–(13) corresponding to $(\beta_1^{n_k}, \beta_2^{n_k}, s_1^{n_k}, s_2^{n_k})$. Therefore, we conclude that there exist

$$A = (f_1^{\epsilon}, f_2^{\epsilon}, z_{f_1}^{\epsilon}, z_{f_2}^{\epsilon}, z_{g_1}^{\epsilon}, z_{g_2}^{\epsilon}, z_{g_3}^{\epsilon}, z_{g_4}^{\epsilon}),$$

and sequences

$$\left\{\underbrace{(f_1^m, f_2^m, z_{f_1^m}, z_{f_2^m}, z_{g_1^m}, z_{g_2^m}, z_{g_3^m}, z_{g_4^m})}_{A_m}\right\}_{m=0}^{\infty},$$

and

$$\left\{\underbrace{(\beta_1^m, \beta_2^m, s_1^m, s_2^m)}_{B_m}\right\}_{m=0}^{\infty}$$

such that

$$\lim_{m \to \infty} A_m^* = A^*, \text{ in } W_p^{2,1,4}(\overline{Q_T}) \times L_4^p(\overline{Q_T}), \quad \lim_{m \to \infty} B_m = (\beta_1^\epsilon, \beta_2^\epsilon, s_1^\epsilon, s_2^\epsilon), \text{ in } L_4^p(\overline{Q_T}),$$

where

$$A_m^* = (e^{-\sum_{i=0}^n h_{i,1}B_i} f_1^m, e^{-\sum_{i=0}^n h_{i,2}B_i} f_2^m, e^{\sum_{i=0}^n h_{i,1}B_i} z_{f_1^m}, e^{\sum_{i=0}^n h_{i,2}B_i} z_{f_2^m}, z_{g_1^m}, z_{g_2^m}, z_{g_3^m}, z_{g_4^m}),$$

and

$$A^* = (e^{-\sum_{i=0}^n h_{i,1}B_i} f_1^{\epsilon}, e^{-\sum_{i=0}^n h_{i,2}B_i} f_2^{\epsilon}, e^{\sum_{i=0}^n h_{i,1}B_i} z_{f_1}^{\epsilon}, e^{\sum_{i=0}^n h_{i,2}B_i} z_{f_2}^{\epsilon}, z_{g_1}^{\epsilon}, z_{g_2}^{\epsilon}, z_{g_3}^{\epsilon}, z_{g_4}^{\epsilon})$$

Using the proof of Theorem 3 and tangent-normal cone techniques [19], for ϵ small enough we have

$$|\beta_1^{\epsilon} - \mathcal{F}\left(b_1 - \frac{\partial z_{g_1}^{\epsilon}}{\partial t}, l_1^*, l_1^{**}\right)| \le \epsilon^{\frac{1}{2}}, \ |\beta_2^{\epsilon} - \mathcal{F}\left(b_2 - \frac{\partial z_{g_2}^{\epsilon}}{\partial t}, l_2^*, l_2^{**}\right)| \le \epsilon^{\frac{1}{2}}, \ a.e. \ (x, y, t) \in \Omega \times [0, T], \ a.s.$$
(35)

$$|s_1^{\epsilon} - \mathcal{F}\left(r_3^* - \frac{\partial z_{g_3}^{\epsilon}}{\partial t}, l_3^*, l_3^{**}\right)| \le \epsilon^{\frac{1}{2}}, \ |s_2^{\epsilon} - \mathcal{F}\left(r_4^* - \frac{\partial z_{g_4}^{\epsilon}}{\partial t}, l_4^*, l_4^{**}\right)| \le \epsilon^{\frac{1}{2}}, \ a.e. \ (x, y, t) \in \Omega \times [0, T]. \ a.s.$$

$$(36)$$

From Theorem 10 (see Appendix) and the Gronwall inequality, it is clear that the function

$$\mathcal{H}: \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^{1} \times \mathcal{W}_{ad}^{1} \to \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^{1} \times \mathcal{W}_{ad}^{1},$$

$$\mathcal{H}(\beta_{1}, \beta_{2}, s_{1}, s_{2}) =$$

$$\left(\mathcal{F}\left(b_{1} - \frac{\partial z_{g_{1}}}{\partial t}, l_{1}^{*}, l_{1}^{**}\right), \mathcal{F}\left(b_{2} - \frac{\partial z_{g_{2}}}{\partial t}, l_{2}^{*}, l_{2}^{**}\right), \mathcal{F}\left(r_{3}^{*} - \frac{\partial z_{g_{3}}}{\partial t}, l_{3}^{*}, l_{3}^{**}\right), \mathcal{F}\left(r_{4}^{*} - \frac{\partial z_{g_{4}}}{\partial t}, l_{4}^{*}, l_{4}^{**}\right) \right),$$

where $z_{g_1}, z_{g_2}, z_{g_3}$ and z_{g_4} are adjoint states corresponding to $(\beta_1, \beta_2, s_1, s_2)$, for almost every $\omega \in \mathbf{\Omega}$ has a unique fixed point $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$. Also using (34) and Theorems 9 and 10, it is easy to show that the sequence $\{(\beta_1^\epsilon, \beta_2^\epsilon, s_1^\epsilon, s_2^\epsilon)\}$ has a subsequence, which converges to $(\beta_1^*, \beta_2^*, s_1^*, s_2^*)$ in $L_4^2(\overline{Q_T})$, as ϵ converges to zero. Therefore, one can arrive at

$$\mathcal{J}(\beta_1^*, \beta_2^*, s_1^*, s_2^*) \leq \mathcal{J}(\beta_1, \beta_2, s_1, s_2), \quad \forall (\beta_1, \beta_2, s_1, s_2) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1, \quad a.s.$$

which results in

$$J(\beta_1^*, \beta_2^*, s_1^*, s_2^*) \le J(\beta_1, \beta_2, s_1, s_2), \quad \forall (\beta_1, \beta_2, s_1, s_2) \in \mathcal{C}_{ad} \times \mathcal{W}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad}^1. \quad a.s.$$

3 Conclusions

The model considered in this paper is a system of two stochastic nonlinear parabolic equations modelling the competition between two languages spoken in the same area. Since the evolution of the population of the speakers of each language depends on several factors including the growth rate of the population and the status of the language, the population with lower growth rate or status in compare with another one may be in danger of extinction. Clearly, by controlling the mentioned factors we can control the evolution of the populations. But, it is worth mentioning that changing these factors can be very costly and may have dangerous side effects on the population densities. These reasons encourage us to study the optimal control for the model of language competition to adopt more accurate and efficient policies in order to control the evolution of the populations. For the reader's convenience, we briefly highlight our contributions as follows:

- We have added the stochastic terms to the model of competition between the languages to deal with a trustable model, which enables us to adopt more accurate policies. It is proved that the stochastic model has a unique solution.
- We have considered four factors consisting of the growth rates of the populations and the status of the languages as the control variables. Then, by considering the cost function (5), we have studied the optimal control problem for the model.
- The stochastic adjoint equations (10)–(16) are obtained, which help us to present the explicit forms of control variables and prove the existence and uniqueness of optimal control. It is shown the system of adjoint equations has a unique solution.
- By presenting some properties for the solutions of the problem (1)–(4) and the adjoint system (10)–(16) corresponding to different control variables in their admissible control sets (see Theorems 9 and 10 in Appendix) and employing normal cone techniques and the adjoint equations, we have derived the necessary conditions for the stochastic optimal control variables.
- The Ekeland variational principle (Theorem 8 in Appendix) is an effective tool in proving the existence and uniqueness of stochastic optimal control variables. This principle together with the properties of the solutions of the problem (1)–(4) and the adjoint system (10)–(16) (see Theorems 9 and 10 in Appendix) and some useful results enable us to prove that there exists unique optimal control.
- It is worthy of attention that we have presented the explicit forms of the stochastic optimal control variables in Theorem 3.

Acknowledgments

The authors are very grateful to reviewers for carefully reading the paper and for their valuable comments and suggestions which improved the original submission of this paper.

Appendix

We provide here some essential mathematical concepts, lemmas and theorems which have been used in the mathematical analysis throughout the paper.

Definition 1. [13] Let B(t) be 1-dimensional Brownian motion and $\mathfrak{F}(t)$ be an associated filtration. An Ito process is a stochastic process of the form

$$X(t) = X(0) + \int_0^t u(s)ds + \int_0^t v(s)dB(s),$$

where X(0) is nonrandom and u(t) and v(t) are adapted stochastic processes and

$$\mathbb{P}\Big[\int_0^t (v(s,\omega))^2 ds < \infty \text{ for all } t \ge 0\Big] = 1, \quad \mathbb{P}\Big[\int_0^t |u(s,\omega)| ds < \infty \text{ for all } t \ge 0\Big] = 1.$$

Theorem 5. [13] Let f(t, x) be a function with continuous partial derivative f_t , f_x and f_{xx} . Also assume that X(t) be an Ito process as described in Definition 1. Then, for every T > 0, we have

$$f(T, X(T)) = f(0, X(0)) + \int_0^T f_t(t, X(t))dt + \int_0^T f_x(t, X(t))u(t)dt + \int_0^T f_x(t, X(t))v(t)dB(t) + \frac{1}{2}\int_0^T f_{xx}(t, X(t))v^2(t)dt.$$

Theorem 6. [12] Let $f(t, x) = (f_1(t, x), \dots, f_n(t, x))$ be a C^2 -smooth map from $[0, \infty) \times \mathbb{R}^n$ into \mathbb{R}^m and dX(t) = u(t)dt + v(t)dB(t) be an n-dimensional Ito process, which is of the following form

$$\begin{cases} dX_1 = u_1 dt + v_{11} dB_1 + \dots + v_{1m} dB_m, \\ \vdots \\ dX_n = u_n dt + v_{n1} dB_1 + \dots + v_{nm} dB_m, \end{cases}$$
(37)

where $B = (B_1, \dots, B_m)$ is an m-dimensional Brownian motion. Then, the process

$$Y(t) = f(t, X(t)),$$

is an Ito process and

$$dY_k(t) = \frac{\partial f_k}{\partial t}(t, X(t))dt + \sum_{i=0}^n \frac{\partial f_k}{\partial x_i}(t, X(t))dX_i(t) + \frac{1}{2}\sum_{i=0}^n \sum_{j=0}^n \frac{\partial^2 f_k}{\partial x_i \partial x_j}(t, X(t))dX_i(t)dX_j(t),$$

where $dB_i dB_j = \delta_{ij} dt$ and $dt dt = dB_i dt = dt dB_i = 0$.

Lemma 1. [13] Let $X_1(t)$ and $X_2(t)$ be Ito processes. Then

$$d(X_1(t)X_2(t)) = X_1(t)dX_2(t) + X_2(t)dX_1(t) + dX_1(t)dX_2(t).$$

Theorem 7. Let (X, Σ) be a measurable space and $(\mathbb{R}, \mathcal{B})$ be the Borel measurable space. Also, assume that $f: X \to \mathbb{R}$ and $g: X \to \mathbb{R}$ are measurable functions and $h: \mathbb{R} \longrightarrow \mathbb{R}$ is a continuous function, then

fg is measurable.

h is measurable.

hof is measurable if f is finite.

If u = f a.e. on X, then u is a measurable function.

Definition 2. [14, 17] Let $0 < \alpha < 1$ and $\Omega \subset \mathbb{R}^n$ be bounded. Then $f \in C^{\alpha, \frac{\alpha}{2}}(\overline{\Omega} \times [0, T])$ if there exists a positive constant C such that

$$|f(x_1,t_1) - f(x_2,t_2)| \le C \Big(|x_1 - x_2|^2 + |t_1 - t_2| \Big)^{\frac{\alpha}{2}}, \ \forall x_1, x_2 \in \overline{\Omega}, \ \forall t_1, t_2 \in [0,T].$$

Furthermore, for any nonnegative integer k

$$C^{2k+\alpha,k+\frac{\alpha}{2}}(\overline{\Omega}\times[0,T]):=\{f\in C^{\alpha,\frac{\alpha}{2}}(\overline{\Omega}\times[0,T]):\partial_x^\beta\partial_t^i f\in C^{\alpha,\frac{\alpha}{2}}(\overline{\Omega}\times[0,T]), \quad |\beta|+2i\leq 2k\}.$$

Theorem 8. [18] Let X be a complete metric space and let $f : X \to (-\infty, +\infty]$ be lower semicontinuous and bounded from below and $\neq +\infty$. Let $\epsilon > 0$ and $x_{\epsilon} \in X$ be such that $f(x_{\epsilon}) \leq \inf \{f(x) : x \in X\} + \epsilon$. Then, there exists $y_{\epsilon} \in X$ such that

$$f(y_{\epsilon}) \le f(x_{\epsilon}), \quad d(x_{\epsilon}, y_{\epsilon}) \le \epsilon^{\frac{1}{2}},$$
$$f(y_{\epsilon}) < f(x) + \epsilon^{\frac{1}{2}} d(y_{\epsilon}, x), \quad \forall x \ne y_{\epsilon}.$$

Definition 3. Let $Q_T = \Omega \times (0, T)$, then we define

$$W_p^{2,1}(Q_T) := \left\{ u \in L^p(Q_T) : \partial_x^\alpha \partial_t^k u \in L^p(Q_T), \ |\alpha| + 2k \le 2 \right\},$$

with the norm $||u||_{W_p^{2,1}(Q_T)} := \sum_{|\alpha|+2k \le 2} ||\partial_x^{\alpha} \partial_t^k u||_{L^p(Q_T)}.$

Lemma 2. Let p > 5, $\Omega \subset \mathbb{R}^n$ be bounded with boundary $\partial \Omega \in C^2$ and

$$\frac{\partial c}{\partial t} - \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij}c + \sum_{i=1}^{n} b_i(x,t) D_i c + d(x,t) c = g(x,t), \quad (x,t) \in Q_T = \Omega \times (0,T), \quad (38)$$
$$c(x,t) = \psi(x,t), \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}),$$

where $\psi \in W_p^{2,1}(Q_T)$, $a_{ij}, b_i, d \in C(\overline{\Omega} \times [0,T])$, $a_{ij} = a_{ji}$ and for constants $0 < C_1 \leq C_2$

$$C_1|\lambda|^2 \le \sum_{i=1}^n \sum_{j=1}^n a_{ij}(x,t)\lambda_i\lambda_j \le C_2|\lambda|^2, \quad \forall \lambda \in \mathbb{R}^n, (x,t) \in Q_T.$$

Also assume that $g \in L^p(Q_T)$, then (38) has a unique solution $c \in W_p^{2,1}(Q_T)$ such that

$$||c||_{W_p^{2,1}(Q_T)} \le \mu \left(||c||_{L^p(Q_T)} + ||\psi||_{W_p^{2,1}(Q_T)} + ||g||_{L^p(Q_T)} \right), \tag{39}$$

where μ depends on T, p, Ω , $\|a_{ij}\|_{L^{\infty}(Q_T)}$, $\|b_i\|_{L^{\infty}(Q_T)}$ and $\|d\|_{L^{\infty}(Q_T)}$. Moreover, if $g \in C(\overline{Q_T})$, then

$$||c||_{L^{\infty}(Q_T)} \le e^{\mu_0 T} \Big(\|\psi\|_{L^{\infty}(\partial Q_T \setminus (\Omega \times \{t=T\})} + T||g||_{L^{\infty}(Q_T)} \Big), \tag{40}$$

where $\mu_0 = 0$ if $d \ge 0$ and $\mu_0 = -\frac{\inf}{Q_T} d$ otherwise.

Proof See [14, 15, 16].

Lemma 3. Let p > 5, $\Omega \subset \mathbb{R}^n$ be bounded with boundary $\partial \Omega \in C^2$ and

$$dc - \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij} c dt + \sum_{i=1}^{n} b_i(x,t) D_i c dt + h(x,t) c dt = g(x,t) dt + c \sum_{k=0}^{m} h_k(x) dB_k, \quad (41)$$
$$(x,t) \in Q_T = \Omega \times (0,T),$$
$$c(x,t) = \psi(x,t), \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}),$$

 $\mathfrak{M} = \max\{\|a_{ij}\|_{L^{\infty}(Q_T)}, \|b_i\|_{L^{\infty}(Q_T)}, \|h\|_{L^{\infty}(Q_T)}, \|D_ih_k\|_{L^{\infty}(Q_T)}, \|D_{ij}h_k\|_{L^{\infty}(Q_T)}\},\$

where $\psi \in W_p^{2,1}(Q_T)$, $a_{ij}, b_i, h, D_{i,j}h_k \in C(\overline{\Omega} \times [0,T])$, $a_{ij} = a_{ji}$, and $h_k(x) = 0$ on $\partial\Omega$, and for constants $0 < C_1 \leq C_2$

$$C_1|\lambda|^2 \le \sum_{i=1}^n \sum_{j=1}^n a_{ij}(x,t)\lambda_i\lambda_j \le C_2|\lambda|^2, \quad \forall \lambda \in \mathbb{R}^n, (x,t) \in Q_T.$$

$$(42)$$

Also assume that $g \in L^p(Q_T)$, then for almost every $\omega \in \Omega$, the problem (41) has a unique solution $c(x,t) \in C(Q_T)$ such that $c(x,t) = e^{\sum_{k=0}^{m} h_k(x)B_k}C(x,t)$ and

$$||C||_{W_{p}^{2,1}(Q_{T})} \leq \mu(\omega) \left(||C||_{L^{p}(Q_{T})} + ||\psi||_{W_{p}^{2,1}(Q_{T})} + ||e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}g||_{L^{p}(Q_{T})} \right), \quad (43)$$

where $\mu(\omega)$ depends on T, p, Ω , and \mathfrak{M} . Moreover, if $g \in C(\overline{Q_T})$, then c is $\mathfrak{F}(t)$ -adapted and for almost every $\omega \in \Omega$, we have

$$||c||_{L^{\infty}(Q_T)} \le \mu'_0(\omega) \Big(\|\psi\|_{L^{\infty}(\partial Q_T \setminus (\Omega \times \{t=T\})} + T||g||_{L^{\infty}(Q_T)} \Big), \tag{44}$$

where $\mu'_0(\omega)$ depends on T, p, Ω , and \mathfrak{M} . Also if $\psi = 0$, then

$$\int_{\Omega} (c(x,t))^2 dx \le e^{\lambda_2^*(\omega)t} \int_0^t \int_{\Omega} (g(x,s))^2 dx ds,$$
(45)

where $\lambda_2^*(\omega)$ depends on T, p, Ω , and \mathfrak{M} .

Proof Let C be the solution of

$$dC - \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij}Cdt + \sum_{i=1}^{n} \left(\underbrace{b_i(x,t) - 2\sum_{j=1}^{n} a_{ij} \sum_{k=0}^{m} B_k D_j h_k(x)}_{b'_i} \right) D_iCdt + \psi_1Cdt = \underbrace{e^{-\sum_{k=0}^{m} h_k(x) B_k} g(x,t) dt, \quad (x,t) \in Q_T, \quad (46)}_{C(x,t) = \psi(x,t), \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}), \quad (46)$$

where

$$\psi_1 = h(x,t) + \sum_{i=1}^n b_i(x,t) \sum_{k=0}^m B_k D_i h_k(x) + \frac{1}{2} \sum_{k=0}^m (h_k(x))^2 - e^{-\sum_{k=0}^m h_k(x)B_k} \sum_{i=1}^n \sum_{j=1}^n a_{ij}(x,t) D_{ij}(e^{\sum_{k=0}^m h_k(x)B_k}).$$

Now, we define A_n and $\mathfrak{F}_n(t)$ as follows

$$A_n = \{ \omega \in \mathbf{\Omega} : |B_k(s)| < n+1, \ 0 \le s \le t, \ k = 0, 1, \cdots, m \} \cap \mathbb{A}$$

and

$$\mathfrak{F}_n(t) = \{A \cap A_n : A \in \mathfrak{F}(t)\},\$$

where

$$\mathbb{A} = \{ \omega \in \mathbf{\Omega} : B_K \text{ is a countinuous function with respect to } t \}$$

Clearly $B_k(t)\Big|_{A_n}$ is a measurable function with respect to $(A_n, \mathfrak{F}_n(t))$. Therefore, using Lemma 2 for almost every $\omega \in \mathbf{\Omega}$, the problem (46) has a unique solution such that $C(x,t) \in W_p^{2,1}(Q_T)$ and there exists positive μ_n such that

$$||C||_{W_p^{2,1}(Q_T)} \le \mu_n \left(||C||_{L^p(Q_T)} + ||\psi||_{W_p^{2,1}(Q_T)} + ||e^{-\sum_{k=0}^m h_k(x)B_k}g||_{L^p(Q_T)} \right), \quad \forall \omega \in A_n, \quad (47)$$

where μ_n depends on T, p, Ω , $\|a_{ij}\|_{L^{\infty}(Q_T)}$, $\|b'_i\|_{L^{\infty}(Q_T)}$ and $\|\psi_1\|_{L^{\infty}(Q_T)}$. Moreover, if $g \in C(\overline{Q_T})$, then there exists positive μ_0^n such that

$$||C||_{L^{\infty}(Q_T)} \le e^{\mu_0^n T} \Big(\|\psi\|_{L^{\infty}(\partial Q_T \setminus (\Omega \times \{t=T\})} + T||g||_{L^{\infty}(Q_T)} \Big), \quad \forall \omega \in A_n.$$

$$\tag{48}$$

It is also clear that the solution of (46), C, is a function of B_k $(k = 0, \dots, m)$. From (46) and (48), $C\Big|_{A_n} = \mathfrak{G}(B_0\Big|_{A_n}, \dots, B_m\Big|_{A_n})$ is a continuous function with respect to $B_k\Big|_{A_n}$ $(k = 0, \dots, m)$. Therefore, from Theorem 7, $C\Big|_{A_n}$ is $\mathfrak{F}_n(t)$ -adapted. Thus $C\Big|_{A_n}$ is $\mathfrak{F}(t)$ -adapted. Also, C the solution of (46) can be defined (almost surely) as follows

$$C = C \Big|_{A_n}, \ \omega \in A_n, \ n \in \mathbb{N}_0.$$

Therefore, for every borel set B, $C^{-1}(B) = \bigcup_{n=0}^{\infty} C \Big|_{A_n}^{-1}(B) \in \mathfrak{F}(t)$. So, C is $\mathfrak{F}(t)$ -adapted. On the other hand, we have

$$d(C^{2}) = 2CdC = 2C\sum_{i=1}^{n}\sum_{j=1}^{n}a_{ij}(x,t)D_{ij}Cdt - 2C\sum_{i=1}^{n}b'_{i}D_{i}Cdt - 2\psi_{1}C^{2}dt + 2e^{-\sum_{k=0}^{m}h_{k}(x)B_{k}}g(x,t)Cdt,$$
$$(x,t) \in Q_{T},$$
$$C(x,t) = \psi(x,t), \quad (x,t) \in \partial Q_{T} \setminus (\Omega \times \{t=T\}).$$

Therefore if $\psi = 0$, then

$$\int_{\Omega} d(C^{2}) dx = \int_{\Omega} \left(2C \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij}C - 2C \sum_{i=1}^{n} b'_{i}(x,t) D_{i}C - 2\psi_{1}(x,t)C^{2} + 2e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}} g(x,t)C \right) dt dx = \int_{\Omega} \left(-2 \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{i}C D_{j}C - 2C \sum_{i=1}^{n} b'_{i}(x,t) D_{i}C - 2\psi_{1}(x,t)C^{2} + 2e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}} g(x,t)C \right) dt dx$$

Hence, from (42), we conclude that if $\psi = 0$, then

$$\int_{\Omega} (C(x,t))^2 dx = \int_{0}^{t} \int_{\Omega} \left(-2\sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,s) D_i C D_j C - 2C \sum_{i=1}^{n} b'_i(x,s) D_i C - 2\psi_1(x,s) (C)^2 + 2e^{-\sum_{k=0}^{m} h_k(x) B_k} g(x,s) C \right) ds dx \le \int_{0}^{t} \int_{\Omega} \lambda^*(\omega) C^2 dx ds + \int_{0}^{t} \int_{\Omega} e^{-2\sum_{k=0}^{m} h_k(x) B_k} (g(x,s))^2 dx ds, \quad a.s.$$

where $\lambda^*(\omega)$ depends on T, p, Ω and \mathfrak{M} . Employing Gronwall inequality, we deduce that

$$\int_{\Omega} (C(x,t))^2 dx \le e^{\lambda_1^*(\omega)t} \int_0^t \int_{\Omega} (g(x,s))^2 dx ds, \quad a.s.$$
(49)

where $\lambda_1^*(\omega)$ depends on T, p, Ω , and \mathfrak{M} .

Since C is $\mathfrak{F}(t)$ -adapted, c is $\mathfrak{F}(t)$ -adapted. Thus, using the Ito product rule for $c = e^{\sum_{k=0}^{m} h_k(x)B_k}C$, we have

$$d(\underbrace{e^{\sum_{k=0}^{m} h_{k}(x)B_{k}}C}_{c}) = d(e^{\sum_{k=0}^{m} h_{k}(x)B_{k}})C + e^{\sum_{k=0}^{m} h_{k}(x)B_{k}}dC =$$

$$Ce^{\sum_{k=0}^{m} h_{k}(x)B_{k}}\sum_{k=0}^{m} h_{k}(x)dB_{k} + \frac{C}{2}e^{\sum_{k=0}^{m} h_{k}(x)B_{k}}\sum_{k=0}^{m} (h_{k}(x))^{2}dt + e^{\sum_{k=0}^{m} h_{k}(x)B_{k}}dC.$$
(50)

Thus, from (46) and (50), we can conclude that

$$dc - \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij}cdt + \sum_{i=1}^{n} b_i(x,t) D_icdt + h(x,t)cdt = g(x,t)dt + c \sum_{k=0}^{m} h_k dB_k, \quad (x,t) \in Q_T,$$

$$(51)$$

$$c(x,t) = \psi(x,t), \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}).$$

Hence, from (48) and (51), we conclude that for almost every $\omega \in \Omega$, (41) has a continuous solution c. Moreover, if $g \in C(\overline{Q_T})$, then

$$||c||_{L^{\infty}(Q_T)} \le \mu'_0(\omega) \Big(\|\psi\|_{L^{\infty}(\partial Q_T \setminus (\Omega \times \{t=T\})} + T||g||_{L^{\infty}(Q_T)} \Big), \quad a.s.$$

$$(52)$$

where $\mu'_0(\omega)$ depends on T, p, Ω , and \mathfrak{M} . Also if $\psi = 0$, then from (49) we have

$$\int_{\Omega} (c(x,t))^2 dx \le e^{\lambda_2^*(\omega)t} \int_0^t \int_{\Omega} (g(x,s))^2 dx ds, \quad a.s.$$
(53)

where $\lambda_2^*(\omega)$ depends on T, p, Ω and \mathfrak{M} . Now, we want to show that the problem (41) has a unique solution. Let C^1 and C^2 be the solutions of (41). Thus, $U = C^1 - C^2$ is the solution of

$$dU - \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij} U dt + \sum_{i=1}^{n} b_i(x,t) D_i U dt + h(x,t) U dt = U \sum_{k=0}^{m} h_k dB_k, \quad (x,t) \in Q_T, \quad (54)$$
$$U(x,t) = 0, \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}).$$

On the other hand, using the Ito product rule for $U_1 = e^{-\sum_{k=0}^m h_k(x)B_k}U$, one can conclude that

$$dU_{1} = e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}dU + Ude^{-\sum_{k=0}^{m} h_{k}(x)B_{k}} + de^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}dU =$$

$$e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}dU - Ue^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}\sum_{k=0}^{m} h_{k}(x)dB_{k} + \frac{U}{2}e^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}\sum_{k=0}^{m} (h_{k}(x))^{2}dt =$$

$$Ue^{-\sum_{k=0}^{m} h_{k}(x)B_{k}}dU - U_{1}\sum_{k=0}^{m} h_{k}(x)dB_{k} - \frac{U_{1}}{2}\sum_{k=0}^{m} (h_{k}(x))^{2}dt.$$

Therefore, $U_1 = e^{-\sum_{k=0}^m h_k(x)B_k}U$ is the solution of

$$dU_1 - \sum_{i=1}^n \sum_{j=1}^n a_{ij}(x,t) D_{ij} U_1 dt + \sum_{i=1}^n \left(b_i(x,t) - 2 \sum_{j=1}^n a_{ij} \sum_{k=0}^m B_k D_j h_k(x) \right) D_i U_1 dt + \left(h\left(x,t\right) + \sum_{i=1}^n b_i(x,t) \sum_{k=0}^m B_k D_i h_k(x) + \frac{1}{2} \sum_{k=0}^m (h_k(x))^2 - \frac{1}{2} \sum_{k=0}^$$

$$e^{-\sum_{k=0}^{m} h_k(x)B_k} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}(x,t) D_{ij}(e^{\sum_{k=0}^{m} h_k(x)B_k}) U_1 dt = 0,$$

$$U_1(x,t) = 0, \quad (x,t) \in \partial Q_T \setminus (\Omega \times \{t=T\}).$$

Finally, using Lemma 2, one can deduce that $U_1 = 0$ for almost every $\omega \in \Omega$, which results in the uniqueness of the solution of the problem (41) for almost every $\omega \in \Omega$.

Theorem 9. Let p > 5 and (f_1^{-1}, f_2^{-1}) and (f_1^{-2}, f_2^{-2}) be the exact solutions of the problem (1)–(4) corresponding to $(\beta_1^1, s_1^1, \beta_2^1, s_2^1) \in \mathcal{C}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad} \times \mathcal{W}_{ad}^1$ and $(\beta_1^2, s_1^2, \beta_2^2, s_2^2) \in \mathcal{C}_{ad} \times \mathcal{C}_{ad}^1 \times \mathcal{W}_{ad} \times \mathcal{W}_{ad}^1$, respectively. Then, for almost every $\omega \in \Omega$, there exist positive $\iota_0(\omega)$ and $\iota(\omega)$ such that

$$\|(f_1^{-1}, f_2^{-1}) - (f_1^{-2}, f_2^{-2})\|_{L^{\infty}(\Omega)}^{t} \leq \iota_0(\omega) \int_0^t \|(\beta_1^{-1}, \beta_2^{-1}, s_1^{-1}, s_2^{-1}) - (\beta_1^{-2}, \beta_2^{-2}, s_1^{-2}, s_2^{-2})\|_{L^{\infty}(\Omega)}^p ds,$$

and

$$\|(f_1^{1}, f_2^{1}) - (f_1^{2}, f_2^{2})\|_{L^2(Q_t)}^2 \le$$

$$\iota(\omega) \int_0^t \|(\beta_1^1, \beta_2^1, s_1^1, s_2^1) - (\beta_1^2, \beta_2^2, s_1^2, s_2^2)\|_{L^2(Q_s)}^2 ds, \quad Q_t = \Omega \times (0, t),$$

where $\iota_0(\omega)$ and $\iota(\omega)$ depend on $p, \Omega, \|D_i h_{k,l}\|_{L^{\infty}(Q_T)}, \|D_{ij} h_{k,l}\|_{L^{\infty}(Q_T)}$ and $\|h_{k,l}\|_{L^{\infty}(Q_T)}$ $(k = 0, 1, \dots, n, l = 1, 2).$

Proof From the problem (1)–(4), for $\dot{f}_1 = f_1^1 - f_1^2$, we have

$$df_{1} = \nabla .(D_{1}\nabla f_{1})dt - \nabla .(F_{1}f_{1})dt + \left(\underbrace{R_{1}(f_{1}^{1}, f_{2}^{1}, \beta_{1}^{1}) + R(f_{1}^{1}, f_{2}^{1}, s_{1}^{1}, s_{2}^{1}) - R_{1}(f_{1}^{2}, f_{2}^{2}, \beta_{1}^{2}) - R(f_{1}^{2}, f_{2}^{2}, s_{1}^{2}, s_{2}^{2})}_{g_{1}^{*}}\right)dt + \frac{\dot{f}_{1}\sum_{k=0}^{n} h_{k,1}(x, y)dB_{k}, \text{ on } \Omega, t > 0,}{\dot{f}_{1}(x, y, t) = 0, \text{ on } \partial\Omega, t > 0, \dot{f}_{1}(x, y, 0) = 0, \text{ on } \Omega.}$$

$$(55)$$

Using (43), one can deduce that $\dot{f}_1(x, y, t) = e^{\sum_{k=0}^n h_{k,1}(x,y)B_k}\dot{F}$ and

$$||\dot{F}||_{W_{p}^{2,1}(Q_{t})} \leq \mu^{*}(\omega) \left(||\dot{F}||_{L^{p}(Q_{t})} + ||g_{1}^{*}||_{L^{p}(Q_{t})} \right),$$
(56)

where $\mu^*(\omega)$ depends on $\|D_i h_{k,1}\|_{L^{\infty}(Q_T)}$, $\|D_{ij} h_{k,1}\|_{L^{\infty}(Q_T)}$ and $\|h_{k,1}\|_{L^{\infty}(Q_T)}$. Using t-Anisotropic Embedding Theorem [14], one can conclude that

$$(\dot{f}_1(x,y,t))^p \le \mu_1^*(\omega) \left(||\dot{f}_1(x,y,s)||_{L^p(Q_t)}^p + ||g_1^*(x,y,s)||_{L^p(Q_t)}^p \right). \quad a.s.$$
(57)

Similarly, it can be shown that

$$(\dot{f}_2(x,y,t))^p \le \mu_2^*(\omega) \left(||\dot{f}_2(x,y,s)||_{L^p(Q_t)}^p + ||g_2^*(x,y,s)||_{L^p(Q_t)}^p \right), \quad a.s.$$
(58)

where

$$\dot{f}_2 = f_2^1 - f_2^2,$$

and

$$g_2^* = R_2(f_1^1, f_2^1, \beta_2^1) - R(f_1^1, f_2^1, s_1^1, s_2^1) - R_2(f_1^2, f_2^2, \beta_2^2) + R(f_1^2, f_2^2, s_1^2, s_2^2).$$

From (57) and (58), we have

$$\begin{split} \|\dot{f}_{1}(x,y,t)\|_{L^{\infty}(\Omega)}^{p} + \|\dot{f}_{2}(x,y,t)\|_{L^{\infty}(\Omega)}^{p} \leq \\ \mu_{3}^{*}(\omega) \Big(\int_{0}^{t} \|\dot{f}_{1}(x,y,s)\|_{L^{\infty}(\Omega)}^{p} + \|\dot{f}_{2}(x,y,s)\|_{L^{\infty}(\Omega)}^{p} ds + \\ \int_{0}^{t} \|(\beta_{1}^{1},\beta_{2}^{1},s_{1}^{1},s_{2}^{1}) - (\beta_{1}^{2},\beta_{2}^{2},s_{1}^{2},s_{2}^{2})\|_{L^{\infty}(\Omega)}^{p} ds \Big). \ a.s. \end{split}$$

Thus, applying the Gronwall inequality results in

$$\|\dot{f}_{1}(x,y,t)\|_{L^{\infty}(\Omega)} + \|\dot{f}_{2}(x,y,t)\|_{L^{\infty}(\Omega)} \leq \iota_{0}^{*}(\omega) \left(\int_{0}^{t} \|(\beta_{1}^{1},\beta_{2}^{1},s_{1}^{1},s_{2}^{1}) - (\beta_{1}^{2},\beta_{2}^{2},s_{1}^{2},s_{2}^{2})\|_{L^{\infty}(\Omega)}^{p} ds\right)^{\frac{1}{p}} \cdot a.s.$$

Also, from (45), (55) and the Gronwall inequality, one can deduce that

$$\|(f_1^{1}, f_2^{1}) - (f_1^{2}, f_2^{2})\|_{L^2(Q_t)}^2 \le \iota(\omega) \int_0^t \|(\beta_1^{1}, \beta_2^{1}, s_1^{1}, s_2^{1}) - (\beta_1^{2}, \beta_2^{2}, s_1^{2}, s_2^{2})\|_{L^2(Q_s)}^2 ds. \quad a.s.$$

Theorem 10. Let p > 5 and $(z_{f_1}^1, z_{f_2}^1, z_{g_1}^1, z_{g_2}^1, z_{g_3}^1, z_{g_4}^1)$ and $(z_{f_1}^2, z_{f_2}^2, z_{g_1}^2, z_{g_2}^2, z_{g_3}^2, z_{g_4}^2)$ be the exact solutions of the stochastic adjoint system (10)–(16) corresponding to $(\beta_1^1, s_1^1, \beta_2^1, s_2^1) \in C_{ad} \times C_{ad}^1 \times W_{ad} \times W_{ad}^1$ and $(\beta_1^2, s_1^2, \beta_2^2, s_2^2) \in C_{ad} \times C_{ad}^1 \times W_{ad} \times W_{ad}^1$, respectively. Then, for almost every $\omega \in \Omega$, there exists positive $\kappa^*(\omega)$ such that

$$\begin{split} \|(z_{f_1}^1, z_{f_2}^1, z_{g_1}^1, z_{g_2}^1, z_{g_3}^1, z_{g_4}^1) - (z_{f_1}^2, z_{f_2}^2, z_{g_1}^2, z_{g_2}^2, z_{g_3}^2, z_{g_4}^2)\|_{L^{\infty}(\Omega)}^p &\leq \\ \kappa^*(\omega) \Big(\int_t^T \int_0^s \|(\beta_1^1, \beta_2^1, s_1^1, s_2^1) - (\beta_1^2, \beta_2^2, s_1^2, s_2^2)\|_{L^{\infty}(\Omega)}^p dl ds + \\ \int_t^T \|(\beta_1^1, \beta_2^1, s_1^1, s_2^1) - (\beta_1^2, \beta_2^2, s_1^2, s_2^2)\|_{L^{\infty}(\Omega)}^p ds \Big), \end{split}$$

where $\kappa^*(\omega)$ depends on p, Ω , $\|D_i h_{k,l}\|_{L^{\infty}(Q_T)}$, $\|D_{ij} h_{k,l}\|_{L^{\infty}(Q_T)}$ and $\|h_{k,l}\|_{L^{\infty}(Q_T)}$ $(k = 0, 1, \dots, n, l = 1, 2)$.

Proof Similar to the proof of Theorem 9, we can get the result.

References

- Parshad RD, Bhowmick S, Chand V, Kumari N, Sinha N. What is India speaking? Exploring the "Hinglish" invasion. Physica A. 2016;449: 375–389.
- [2] Iriberri N, Uriarte J. Minority language and the stability of bilingual equilibria. Ration Soc. 2012;24:442–462.
- [3] Bakalis E, Galani A. Modeling language evolution: Aromanian, an endangered language in Greece. Physica A. 2012;391:4963–4969.
- [4] Patriarca M, Castelló X, Uriarte JR, Eguíluz VM, San Miguel M. Modeling two-language competition dynamics. Adv Complex Syst. 2012;15:1–24.
- [5] Wiese H. Language competition: an economic theory of language learning and production. Int J Sociol Lang. 2015;2015: 295–329.
- [6] Patriarca M, Heinsalu E. Influence of geography on language competition. Physica A. 2009;388:174–186.
- [7] Minett JW, Wang WS-Y. Modelling endangered languages: The effects of bilingualism and social structure. Lingua. 2008;118:19–45.
- [8] Gong T, Shuai L, Zhang M. Modelling language evolution: Examples and predictions. Phys Life Rev. 2014;11:280–302.
- [9] Tse YC, Chan CKP, Luk MH, Kwong NH, Leung PT, Binder R, Schumacher S. A population-competition model for analyzing transverse optical patterns including optical control and structural anisotropy. New J Phys. 2015;17:083054.
- [10] Hou R, Wu J-w, Du HS. Customer social network affects marketing strategy: A simulation analysis based on competitive diffusion model. Physica A. 2017;469:644–653.
- [11] Parta P, Vassallo CN, Wall D, Igoshin OA. Mechanism of kin-discriminatory demarcation line formation between colonies of swarming bacteria. Biophys J. 2017;113:2477–2486.
- [12] Øksendal B. Stochastic Differential Equations: An Introduction with Applications. 2th ed., Berlin: Springer; 1985.
- [13] Shreve SE. Stochastic Calculus for Finance II: Continuous-Time Models. New York: Springer; 2004.
- [14] Wu Z, Yin J, Wang C. Elliptic and Parabolic Equations. Singapore: World Scientific; 2006.
- [15] Ladyzenskaja OA, Solonnikov VA, Ural'ceva NN. Linear and Quasi-Linear Equations of Parabolic Type. Providence: American Mathematical Society; 1968.

- [16] Esmaili S, Eslahchi MR. Optimal control for a parabolic–hyperbolic free boundary problem modeling the growth of tumor with drug application. J Optim Theory Appl. 2017;173:1013– 1041.
- [17] Brezis H. Functional Analysis, Sobolev Spaces and Partial Differential Equations. New York: Springer; 2011.
- [18] Barbu V. Mathematical Methods in Optimization of Differential Systems. Dordrecht: Kluwer Academic Publishers; 1994.
- [19] Barbu V, Iannelli M. Optimal control of population dynamics. J Optim Theory Appl. 1999;102:1–14.
- [20] Kurtz DS, Swartz CW. Theories of Integration: the Integrals of Riemann, Lebesgue, Henstock-Kurzweil, and Mcshane. Singapore: World Scientific; 2004.