DYNAMICAL SIMULATIONS OF THE FIRST GLOBULAR CLUSTERS

RAYMOND G. CARLBERG

Department of Astronomy & Astrophysics, University of Toronto, Toronto, ON M5S 3H4, Canada

Draft version January 24, 2022

ABSTRACT

A Milky Way-like halo is simulated with tidally limited star clusters inserted in the dark matter halos present at high redshift. An n-body code augmented with velocity relaxation in the star clusters evolves the system. The stripped stars and remnant clusters that began in the lower mass sub-halos have a distribution somewhat more extended than the dark matter halo, with a mean galactic radius of about 60 kpc inside 150 kpc, whereas the distribution of all stripped stars and clusters is more centrally concentrated than the dark matter. The clusters from low mass sub-halos, those with a peak circular velocity of $12 - 18 \text{ km s}^{-1}$, also produce most of the population's thin stellar streams. Together these suggest a search strategy for extremely metal poor cluster stars and their remnants. The dependence of the stellar population distribution on sub-halo mass is not seen in simulations that start clusters at lower redshift. The half mass radii of the clusters are set by the tidal fields of the initial sub-halo, "dwarf galaxy" location, which causes the average half mass radius to decrease with increasing redshift of formation. Starting clusters at greater than redshift 4 leads to cluster half mass radii approximately as seen in the Milky Way.

Subject headings: Milky Way dynamics; Milky Way dark matter halo

1. INTRODUCTION

The first globular clusters can be defined as those that formed above redshift 6, approximately the first billion years of the universe, the age over which reionization was largely completed (Planck Collaboration et al. XLVII 2016). Globular clusters may have played an important role in providing the ionizing photons for reionization (Ricotti 2002; Boylan-Kolchin 2018). Even if globular clusters are sub-dominant ionizing sources they are expected to be valuable tracers of star formation and structure at high redshift during their bright phase when O and B stars are present, being within the observational capabilities of the James Webb Space Telescope (Carlberg 2002; Renzini 2017; Boylan-Kolchin 2018). Confident identification of the remnants of the first clusters in current epoch galaxies, including the Milky Way, will enable detailed studies of their chemical makeup at low metallicity, which provides clues to the nature of even more metal poor stars, and to provide insights into the cluster's dynamical evolution and mass loss history.

Globular clusters with metallicities below $[Fe/H] \lesssim -1$ have halo kinematics and there is considerable evidence that these are predominantly accreted onto the Milky Way as part of the hierarchical buildup of the galaxy (Searle & Zinn 1978; Brodie, & Strader 2006). Milky Way globular clusters show an age-metallicity relation (VandenBerg et al. 2013) for $[Fe/H] \ge -1.7$, but clusters with $[Fe/H] \leq -1.7$ are consistent with a common age that would place their formation in the z > 6 regime. In the Milky Way approximately 1/3 of the halo clusters have [Fe/H] < -1.7 and are within one Giga-year of the Hubble age, although the age dating uncertainties do not guarantee that they were formed at or above redshift 6. Observational confirmation of the numbers of the high redshift population is an important goal. Nevertheless, the globular clusters that formed above redshift 6 likely

are a population with significant numbers.

The oldest of the first globular clusters should have very low metal abundances. The metallicity distribution function of field halo stars is approximately in accord with a simple closed box model (Hartwick 1976), for [Fe/H] ≥ -3.5 , below which the numbers begin to fall below the relation (Da Costa et al. 2019; Youakim et al. 2020). On the other hand, globular clusters appear to exhibit a metallicity floor at about [Fe/H] $\simeq -2.5$ (Beasley et al. 2019). There is some theoretical support for [Fe/H] $\simeq -2.5$ being the minimum to produce sufficient cooling to allow a globular cluster to form (Kruijssen 2019), but it is dependent on abundances of individual elements at high redshift. A floor could be at a yet lower metallicity (Hartwick 2018; Yoon et al. 2019).

The apparent metallicity floor for the globular clusters may indicate that extremely metal poor globular clusters did not form, which would limit the usefulness of globular clusters as a tracer population of general star formation at high redshift. On the other hand, extremely metal poor globular clusters could have formed, then dissolved in the tidal field of the galaxy and the progenitor systems that hosted them. Moreover, there remains the intriguing possibility that there are a few of these rare objects waiting to be found. Encouragement comes from high resolution spectra of two stars in inner halo Sylgr stream (Ibata et al. 2019) which have essentially identical abundance patterns at a metallicity of [Fe/H] = -2.9(Roederer, & Gnedin 2019). Since thin stellar streams generally arise from a globular cluster progenitor, further confirmation of the Sylgr metallicity will push the globular cluster metallicity floor down and increase the prospects for a successful hunt for extremely metal poor, $[Fe/H] \lesssim -3$, globular clusters and additional extremely metal poor stellar streams.

The theoretical study of the formation of globular clusters has some basic physical considerations of sufficient gas cooling to allow stars to form (Peebles, & Dicke 1968;

carlberg@astro.utoronto.ca

Gunn 1980). The study of globular cluster formation in a cosmological context is coming within the capabilities of high resolution numerical simulations, although there is not yet good agreement on details. High redshift clusters likely form from the gas present in central regions of the relatively low mass dark matter halos that are present at that time. There could be a single process, or, a two-stage globular cluster formation history (Katz, & Ricotti 2013). Empirical study at low redshift finds that stars form in a wide range of group sizes and densities (Lada & Lada 2003). The globular clusters that survive are likely to be the dense, massive cluster end of general star formation that produces young clusters of a wide range of densities and masses, (Bromm, & Clarke 2002; Leaman et al. 2013; Phipps et al. 2019). Tidal fields disperse most of the star clusters formed, leaving those dense enough to survive in their tidal field environment (Fall & Zhang 2001; Chandar et al. 2017). An alternate model emphasizes the creation of "naked" of dark matter globular clusters in the collision of gas-rich dark matter sub-halos which can produce the very high gas densities required for a globular cluster (Trenti et al. 2015; Lahén et al. 2019; Madau et al. 2020). A key issue is the dominant physics that regulates the characteristic size of halo globular clusters. Tidal fields present during formation and afterwards will always set an upper limit, but gas and radiation processes could produce smaller clusters.

The evolution of globular clusters within a cosmological distribution of dark matter is largely a dynamical problem. The dynamical phase begins after a complex and poorly understood phase of rapid star formation and vigorous stellar mass loss and gas expulsion in the first 100 Myr of their lifetime (Calura et al. 2019; Howard et al. 2019). The ubiquitous presence of distinct populations of chemically enriched second generation stars within globular clusters (summarized in Renzini et al. (2015) points to the complexity of the process but also suggests a cosmologically short timescale during which gas processes continued (D'Antona et al. 2016). The simulations here start when the clusters effectively become stellar dynamical objects, which is sufficiently close in time to their star formation phase that they are near their site of formation.

One of the goals of this study is to provide some guidance on the likelihood of survival of very high redshift globular clusters and where the remnant clusters or tidal star streams are likely to be found. In a hierarchical assembly picture, the oldest, hence most metal poor, stars in the galactic halo should be centrally concentrated (Tumlinson 2010), which is then likely true for globular clusters as a tracer population, although globular clusters that orbit in the central few kilo-parsecs of the galaxy cannot survive long (Gnedin, & Ostriker 1999; Gnedin et al. 2014). Another goal is to follow the mass-radius relation of the clusters from their time of formation in the low mass dark halos present at high redshift to their current epoch orbits in the dark matter halo of a Milky Way-like galaxy.

In this paper dynamically realistic globular clusters are started in a cosmologically realistic (cosmic web-like) distribution of dark matter present at redshift 8. The dark matter and star clusters evolve under their gravitational forces to assemble into a Milky Way-like galactic halo, with most of the mass accretion before redshift one. The analysis is supplemented with simulations started at redshift 4.6 and 3.2. The simulations are useful to answer questions about the rate of mass loss from the clusters subject to the strong tidal fields present, and, the galactic distribution of the clusters, their half mass radii, and their tidally removed stars. The next section briefly discusses the numerical methods. Section §3 discusses the time evolution of the simulations, §4 the galactic distribution of clusters and stripped stars, comparing those from low and higher mass sub-halos. In §5 the relationship between the final cluster sizes and the starting conditions is explored. The Discussion, §6, proposes connections between these dynamical results and the evolution and distribution of metal poor globular clusters.

2. COSMOLOGICAL STAR CLUSTER SIMULATIONS

The range of about 10^5 in dynamical length scale from star clusters, with a half mass scale of 5-10 parsecs, to a galactic scale dark matter halo a virial radius of 200 kilo-parsecs is a numerical challenge. The stellar dynamics of stars clusters is fundamentally collisional (Spitzer 1987) in which gravitational interactions between individual stars, binaries and other multiples allows some stars to increase their orbital energy within the cluster at the expense of others. External tidal fields heat and pull stars away from the clusters. To tackle these problems a modified version of the standard cosmological n-body code, Gadget3 (Springel 2005) is employed without any hydrodynamics. The dark matter particles have a mass of $4 \times 10^4 \,\mathrm{M_{\odot}}$ with a softening of 200 pc. Star particle clusters are added into the simulation. The star particles are made as close to stellar masses as practical, $5 M_{\odot}$, with a softening of 2 pc. in total there are about 10^8 particles in the simulation, about 60% of them dark matter.

A Monte Carlo heating of the star particles in the clusters is added to Gadget3, in the form of random velocity increments of sizes determined by the half mass velocity relaxation rate (Spitzer 1987; Binney & Tremaine 2008). The heating is calibrated using NBODY6 runs (Aarseth 1999; Carlberg 2018, 2020). Calibration runs of large N clusters remain a challenge. However, as cluster masses increase tidal heating dominates over internal heating, so the outcome of massive cluster evolution is expected not to depend sensitively on the star heating. Additional calibration and increasing the total number of particles to further refine and check the results will be valuable.

To create a Milky Way-like final system, the simulations are conveniently initiated using the Via Lactea II halo catalogs (Madau et al. 2008) which are reconstituted into particle halos using the Hernquist (1990) profile. The halos are traced back from the virialized halos at the final state of the simulation so need to be boosted with extended halos that allows for the mass not associated with virialized sub-halos at redshift 8, for which the average extended mass halo boosts the total mass a factor of 7-8. The extended halos are about the same mass as the virialized core at redshift 4.6, and even less at redshift 3.2. The approach of reconstituting halos offers some advantages and creates a realistic dark matter distribution, but is being pressed to its limits at high redshift.

The star clusters are created using a King model equi-




FIG. 1.— The evolving spatial distribution of globular cluster stars in a redshift 8 simulation. Stars initially in clusters started in sub-halos below a virialized mass $2.5 \times 10^8 M_{\odot}$ are shown in the left panels and above $25 \times 10^8 M_{\odot}$, one decade in mass higher in the right panels, all in a ± 100 kpc box. Times shown are 1.7, 3.7, 5.7, 7.7, 9.7, 11.7 and 13.7 (units of 1.022 Gyr) from the top. Both sets of stars are from the same simulation.

librium (King 1966) with a dimensionless central potential, $W_0 = 7$. The outer radius of a cluster of mass M_c is scaled to the simple tidal radius, $r_t = [M_c/(3M(<$ $[r]^{1/3}r$, where M(< r) is the mass inside sub-halo at a cluster's initial orbital radius r. The clusters are usually set up to initially exactly fill this tidal radius. Tests show that tidally under-filling initial clusters quickly expand to the local tidal radius, whatever it is. For the chosen value of W_0 the ratio of cluster half mass to tidal radius is then 0.22. Clusters are inserted into the sub-halo in a disk-like distribution. The initial radii are drawn from an exponential disk distribution with a scale radius that is 20% of the radius of the peak of the rotation curve for the sub-halo, roughly what is expected based on the average angular momentum of the sub-halo. The number of clusters is based on a ratio of cluster mass to virial mass of 10^{-4} , which is about 1.3×10^{-5} allowing for the extended mass halo (Hudson et al. 2014). There are typically one or two clusters per sub-halo. The clusters are started with a system velocity of the local circular velocity plus 5 km s⁻¹ of random velocity.

3. EVOLUTION OF THE CLUSTERS

The simulations provide representative orbital histories of high redshift globular clusters that are accreted onto the Milky Way-like halo. Stars are pulled away from the clusters by the tidal fields in their initial sub-halos, and later the main halo of the galaxy. The orbits of the tidal stream stars are accurately followed. The distribution of those tidal stars is one of the most important outcomes of these simulations to guide searches for extremely metal poor clusters.

Figure 1 shows the evolution of the spatial distribution of the clusters (unresolved in the plot) and stars that tidal fields pull away from clusters. The left panel shows the stars from clusters originally in low mass sub-halos, those with a virialized halo of less than $2.5 \times 10^8 \,\mathrm{M_{\odot}}$, total extended mass about $2 \times 10^9 \,\mathrm{M_{\odot}}$ or less. The right panel shows the stars from clusters started in sub-halos at least ten times more massive, $2.5 \times 10^9 M_{\odot}$ and a total extended mass of about $2 \times 10^{10} M_{\odot}$ and greater. The stars from clusters started in more massive sub-halos are much more concentrated to the center of the dark halo in a nearly smooth density distribution, although one or two short streams appear near the center at late times. The clusters started in the low mass sub-halos produce a rich set of star streams, which are visible from the inner few kpc to 100 kpc. Because tidal fields pull stars away from a cluster near pericenter, at the inner turnaround point the stream width is comparable to the local tidal radius of the progenitor cluster, say 0.1 kpc. As the stream travels out to its outer turnaround radius the width expands, often to a few kpc.

The time dependence of the fractional mass remaining in the clusters that are located within 150 kpc of the center of the halo at the end of the simulation is shown in Figure 2. On the average, the clusters started at high redshift lose a bit more than half of their mass of the course of the simulation. Most of a cluster's mass is lost when it is in the strongest tidal fields at the beginning of the simulation. The red lines are for the most massive clusters, which suffer the greatest mass loss. The problem of the massive clusters for high redshift of formation




FIG. 2.— The fractional mass remaining as a function of time for clusters within 150 kpc of the main halo at the end of the simulation. The colors scale gives the initial cluster mass, M_c , in log $10(M_c/M_{\odot})$ of the cluster, ranging from turquoise (4.5-4.95), blue (4.95-5.4), green (5.4-5.85) and red (5.85-6.3). The black dotted line is the average.

is an interesting one that will be considered in a future paper.

The tidal tensor is calculated as the second derivatives of the potential with respect to the coordinates, using an offset from the cluster center of 50 pc. The largest absolute eigenvalue of the tidal matrix is the plotted quantity. Ongoing merging of the sub-halos into the growing main halo causes the mean tidal fields to drop with time, Figure 3. The tidal field in a dark halo with circular velocity V_c scales with V_c^2/r^2 at radius r. The star clusters begin in dark halos with circular velocities of some 20 km s⁻¹ which merge to create a galactic halo with a circular velocity the increase in orbital radius as a result of halo buildup.

The tidal field increases with distance from the center of a star cluster, driving an approximately linear increase in impulsive velocity change in the outskirts of the cluster, or a quadratic increase in kinetic energy with size. Although the clusters are set up to have about the same mean local density, $r \propto M^{0.33}$ the variation in location within a sub-halo and that massive clusters are more likely to be found in more massive sub-halos leads to an outcome where the population of clusters has a size mass relation, $r \propto M^{0.6}$. High mass clusters are then relatively more affected by tidal heating.

The rate of mass loss drops very quickly with time in Figure 2. Mass loss is largely the result of tidal heating, which drop quickly as the substructure merges together to reduce the average density, hence tidal field strength leads to a fairly rapid decline in the mass loss rate. A simulation started at redshift 4.56 with the same setup principles has a much lower mass loss rate as shown in Figure 4. The small dense halos present in the redshift 8 start are a challenge to the numerical resolution, so there is also likely to be some two-body relaxation between the heavy dark matter particles and the light star particles which requires larger N simulations.

It is observationally well established that globular clusters lose mass to tidal fields, perhaps as best illustrated with the Pal 5 cluster (Grillmair & Dionatos 2006) al-




FIG. 3.— The largest eigenvalue of the tidal matrix as a function of time for the clusters in the high mass sub-halos (top panel) and the low mass sub-halos (bottom panel).




FIG. 4.— The fractional mass remaining in the clusters as a function of time for a redshift 4.56 start. The colors are the same as in Figure 2.

though there is an ever growing list of globular clusters with clearly detected tidal tails. The chemically unique second generation stars are used to estimate the fraction of the stellar halo that originated within globular clusters (Martell et al. 2011) with current estimates being about




FIG. 5.— The radial density distribution of stars lost from clusters in sub-halos with mass below $0.025\times10^{10}~M_{\odot}$ (blue solid line) and the higher sub-halos mass above $0.25\times10^{10}~M_{\odot}$ (blue dashed line) at the end of the simulation. The dark matter density profile is shown as the black solid line.

10% (Koch et al. 2019). Their results imply that clusters as a population have lost some 50+% of their stellar mass and about half of the initially present clusters have dissolved. The empirical estimates contain a number of parameters with uncertain values, however, on the whole the substantial presence of stars originating in globular clusters favors a redshift higher than 5 to give the strong tides required for the observationally inferred halo fraction of globular cluster stars.

4. THE DISTRIBUTION OF CLUSTER REMNANTS

The number of unbound cluster stars per unit volume as a function of radial distance in the halo is shown in Figure 5. The lines show the densities of the stars from clusters in the lowest mass sub-halos, those below $0.025 \times 10^{10} \,\mathrm{M_{\odot}}$ (solid line) and the highest mass sub-halos, those above $0.25 \times 10^{10} \,\mathrm{M_{\odot}}$.

The much more concentrated density distribution that develops for the star clusters that begin in the more massive sub-halos, see Figure 1, is not present in the early time distribution, although the clusters are stronger tidal fields on the average, see Figure 3. Figure 6 shows the radial distribution of the same clusters as in Figure 5, but at time 1.45, the model universe has an age of 1.41 Gyr and the simulation is 0.73 Gyr old. The differences in radial distribution of the two sub-populations that develop over the course of the simulation is likely due to dynamical friction during infall being a larger effect for high mass sub-halo infall than for low mass sub-halos.

The segregation of clusters formed in low and high mass sub-halos is not prominent for simulations started at lower redshifts. Density profiles of dark matter, stars lost from clusters in sub-halos below $0.035 \times 10^{10} \,\mathrm{M_{\odot}}$ and stars from clusters in sub-halos with mass above $0.35 \times 10^{10} \,\mathrm{M_{\odot}}$ are shown in Figure 7 for a simulation started at redshift 4.56. Although the mass ranges are comparable to the higher redshift start, the density profiles of the clusters in different mass sub-halos do not show a dependence on sub-halo mass. The overall density distribution of this lower redshift infall population is more extended than the dark matter. A redshift 3 start




FIG. 6.— The radial density distribution of the stars lost from clusters in low (solid blue) and high (dashed blue) mass sub-halos near the beginning of the simulation, 0.73 Gyrs after the start. There is no initial difference in density distribution of the star clusters with sub-halo mass.




FIG. 7.— Similar to Figure 5 but for a redshift 4.56 start. The radial density distribution of stars removed from clusters in subhalos with mass below $0.035\times 10^{10}\,M_\odot$ (blue solid line) and the higher sub-halos mass above $0.35\times 10^{10}\,M_\odot$ (blue dashed line) at the end of the simulation. The dark matter density profile is shown as the black solid line.

has an even less centrally concentrated distribution of remnant clusters and stellar debris.

The endpoint cluster masses with their distances from the center in the galactic halo are shown in Figure 8. The numbers of clusters are based on starting with a ratio of cluster mass to halo mass of 1.2×10^{-5} , which may be appropriate for the entire halo cluster population, of which redshift 8 clusters would be some currently unknown fraction. The red points in Figure 8 are the 3 surviving clusters that began in the six most massive sub-halos. They are at such small galactic radii that they would certainly be merged into the baryonic disk and bulge. The blue dots in Figure 8 are the clusters started in the 58 lowest mass sub-halos that were seeded with globular clusters. These clusters lose significant mass when within 20 kpc of the center, but are largely a more distant population. The green dots are for

FIG. 8.— Current radial locations of clusters started in sub-halos of mass less than $2.5\times10^8\,M_\odot$ (blue) greater than $10\times10^8\,M_\odot$ (green) and greater than $25\times10^8\,M_\odot$ (red). Two of the five most massive clusters have completely evaporated, and there are four other clusters inside 10 kpc that have lost so much mass that they fall below the plotted mass range. The inset shows the distribution of clusters in the Harris (1996) catalog, assuming a mass-to-light of 2. The colors of the inset indicate the metallicity range $-1 \geq$ [Fe/H] > -2 (red) [Fe/H] ≤ -2 (blue).

10

R_{GC} [kpc]

100

a cluster population started in the 31 heaviest halos and are an intermediate population of lower mass and smaller galactic radii. The blue dots in the Figure 8 inset are the [Fe/H] < -2 clusters in the Harris (1996) catalog and appear to have a distribution closest to the clusters started in the low mass sub-halos. The simulated clusters do not have final time clusters with masses quite as large as the most massive in the Milky Way.

The clusters that begin in low-mass sub-halos have an average distance from the center of the galaxy of approximately 60 kpc, for those within 150 kpc, and the median radius is approximately 40 kpc. Therefore, in contrast to the general distribution of stars formed at high redshift which are concentrated to the center of the final dark matter halo, the stars formed in the lower mass subhalos present at high redshift, and sufficiently massive to be able to support star formation, the stars and clusters are distributed in a slightly shallower radial distribution than the dark matter.

5. CLUSTER HALF MASS RADIUS DEPENDENCE ON STARTING REDSHIFT

The simulated redshift 8 clusters substantially overlap in half-mass radii, Figure 9, with those measured in the Milky Way, shown as the inset in Figure 9. The Harris (1996) projected half-light radii have been multiplied by 1.33 to get to a 3D value (Baumgardt et al. 2010) under the mass follows light assumption. After setting the [Fe/H] ≥ -1 Milky Way disk clusters aside, the distribution of very metal poor clusters, [Fe/H] ≤ -2 is broadly similar to the simulated clusters, although somewhat larger. The Milky Way contains a few clusters more massive than in the simulation and are probably not adequately modeled in these simulations.

The tidal field sets an upper limit to the size of clusters. In principle gas cooling and star formation processes could produce significantly smaller clusters, although as clusters become denser the internal star-star




FIG. 9.— Cluster half mass radii as a function of their galactocentric distance for the clusters at the end of the redshift 8 simulation. The Milky Way $[Fe/H] \leq -1$ clusters from the Harris (1996) catalog are shown in the inset, with the colors indicating metallicity, as in Figure 8.




FIG. 10.— Cluster half mass radii vs galactic radii for clusters started at redshift 8 (red), 4.6 (green) and 3.2 (blue). The highest redshift clusters have the smallest half mass radius. The redshift 3.2 clusters only have a small fraction with sizes comparable to Milky Way halo clusters.

relaxation will increase. It is an interesting possibility that tidal fields, largely from the dark matter distribution in which they form, play the dominant role in determining the size of the clusters formed at high redshift which are later incorporated into the galactic halo. Figures 10 and 11 plot respectively, r_h , the half mass radius, and, r_h/r_t , the ratio of the half mass radius to the tidal radius, against the final time galactic radius for simulations started at redshift 8, 4.6 and 3.2. The redshift 4.6simulation was run again starting with clusters 2.5 times smaller, but the clusters quickly expand to fill their tidal radii in the sub-halo. Consequently the results are not significantly dependent how under-filling of the tidal radius clusters are at the outset. The half mass radius has a weak rise with galactic radius, $r_h \propto R_{GC}^{0.26\pm0.2}$. The ratio r_h/r_t in the final galactic halo is a strong function of galactic radius, as a result of the dependence of tidal radius of galactic radius. Most of the clusters have a half

10⁶

Current Mass [M₍₂₂₈₄₎] 10⁵

1000




FIG. 11.— The ratio of individual cluster half mass radius to its current location tidal radius as a function of galactic radius for clusters started at redshift 8 (red), 4.6 (green) and 3.2 (blue).




FIG. 12.— The half mass radius-mass relation for the clusters in the redshift 8 simulation at times 0.7 Gyr (red), 7.5 Gyr (green) and 13.4 Gyr (blue). The relations varies little with time for clusters that remain bound, although the clusters lose considerable mass with time.

mass relaxation time longer than an orbital time so the clusters do not change much in half mass size as they orbit, whereas the local tidal radius varies around the orbit, leading to the strong dependence of the ratio on the galactic radius. The half mass radii of the clusters at the beginning, halfway point, and end of the redshift 8 simulation are shown in Figure 12 which shows that the cluster half mass radii are largely determined by the tidal fields present at the starting location of the clusters. Although clusters lose considerable mass with time, Figure 2, much of it occurs at early times when the tidal fields are similar to those at the start, Figure 3. These simulations apply only to halo clusters where the strong tidal fields of a baryonic galaxy can be ignored.

Figures 10 and 11 also show that at any given galactic radius the clusters are ordered with decreasing r_h , and the ratio r_h/r_t , with increasing redshift at which they were started. For a more realistic simulation with a continuous cluster formation history the values of r_h would be smoothly distributed and plausibly would resemble the same relation in the Milky Way (Baumgardt et al. 2010). The width of the r_h relation for a single starting redshift is likely due to the range of tidal forces at the spread of the initial radial locations of the clusters in their starting sub-halos, see Figure 3.

It is likely that the starting redshift of formation and the initial orbital radius at which the clusters are initially placed are somewhat degenerate quantities in determining final time r_h values for individual clusters. The redshift 3.2 starting conditions leads to a substantial fraction of the population have half mass radii larger than are seen in the Milky Way globular cluster population. The simulation results here indicate that halo clusters must have formed largely above redshift 4, for which there is some observational support (Katz, & Ricotti 2013).

Had the clusters been started closer to center of the initial sub-halos the stronger tidal fields (assuming a cusped density profile) would lead to somewhat smaller tidal radii, hence, r_h values. However, for a central $\rho(r) \propto r^{-1}$, $M(< r) \propto r^2$, the local tidal radius for a fixed mass cluster varies slowly with radial location, $r_t \propto r^{1/3}$. That is, a factor of two decrease in tidal radius requires a factor of 8 decrease in orbital radius in the sub-halo, which would move the clusters inside 100 pc. Even at a 1 kpc starting radius the clusters will sink due to dynamical friction (Oh et al. 2000; Cole et al. 2012) which happens naturally in these simulations, as visible in a few clusters where the tidal fields increase with time in Figure 3. Since most sub-halos have only a single globular cluster at the outset, friction drawing them to the center is not necessarily a problem, and sub-cluster merging is working to place clusters in larger orbits in merged halos where dynamical friction is reduced. Further simulations are required to clarify these issues.

The most massive star cluster allowed in the initial conditions was $2 \times 10^6 \,\mathrm{M_{\odot}}$ and the most massive clusters have a high relative mass loss rate than lower mass clusters, see Figure 2. It would be straightforward to include more massive clusters in the simulation. Obtaining a cluster like M15/NGC7078 with a mass of nearly $10^6 \,\mathrm{M_{\odot}}$ and a half mass radius of 5 pc (Baumgardt et al. 2010), requires yet stronger tidal fields or special formation conditions.

6. DISCUSSION

The dynamical simulations here find that clusters formed at redshift 8 in higher mass dark matter halos are brought into the center of a Milky Way-like halo and either destroyed at high redshift by the dark halo, or, later as the dense baryonic galaxy develops. On the other hand, the accreted lower mass dark halos put their globular clusters into large radius orbits, an average galactocentric distance of about 60 kpc inside 150 kpc. Most of the stars lost to early time tidal fields are eventually spread out over the halo, however late time tidal streams are thin and are detectable at the current epoch. The Sylgr stream may be an example (Ibata et al. 2019; Roederer, & Gnedin 2019).

The relation between initial dark matter sub-halo mass and the orbits of clusters found at redshift 8 is not seen at redshift 4.6. The differing radial distribution is likely the result of dynamical friction as sub-halos of varying mass fall into the main halo. The time scale for dynamical friction is derived in Binney & Tremaine (2008), Equation (8.13),

$$t_{df} = \frac{1.17}{\log \Lambda} \frac{M(r)}{M_{sat}} t_{cross},\tag{1}$$

where t_{cross} is the crossing time, defined as the travel time for one radian of orbit, $\log \Lambda$ is the Coulomb logarithm, and M(r) is the host halo mass inside radius r(assumed to have $M(r) \propto r$), and M_{sat} is the sub-halo mass, approximated over an orbit as a constant mass satellite. The mass of the dominant halo declines drops quickly with increasing redshift. In the simulations here the most massive halo at redshift 8 is $0.94 \times 10^{10} \,\mathrm{M_{\odot}}$, whereas at redshift 4.6 the most massive is $7.3 \times 10^{10} \,\mathrm{M_{\odot}}$, 7.8 times more massive. In addition the dynamical time in the halo at high redshift is shorter because the universe is denser by a factor of $(1 + z)^3$. Therefore, the same mass sub-halo falling into the dominant halo at redshift 8 has about a factor of 15 times more dynamical friction per unit time than at redshift 5.6. At lower redshift dynamical friction time for the sub-halos becomes longer than the dynamical time for both high and low mass sub-halos so that all sub-halos merge into the main halo with relatively little loss of orbital energy.

Observations show that lower luminosity galaxies host globular clusters that tend to have lower metallicities than the field stars (Lamers et al. 2017) with the Fornax dwarf being a notable example (Larsen et al. 2012). Lower luminosity galaxies also have a higher number of globular clusters per unit galaxy luminosity (Georgiev et al. 2010). Although the relation between galaxy luminosity and dark halo mass remains an active research topic the strong correlation is well established in local group galaxies (Brook et al. 2014) and much of the relation must have been put in place at high redshift.

Low metallicity stars and globular clusters appear to have numbers as a function of metal abundance, Z, in quite good agreement with the simple one zone halo chemical evolution model. The model has a single parameter, the effective enrichment yield, Y, of a generation of stars, and then predicts cumulative numbers $N(< Z) \propto 1 - \exp(-Z/Y)$ with a normalization that depends on when star formation ceases, or, is matched to an observational data set (Hartwick 1976). Depending on the details of calibrating to Milky Way halo numbers, if globular clusters trace the field star metallicity distribution, then 3-10 clusters are expected below the apparent floor of [Fe/H] < -2.5 if the clusters continue to be in proportion to the numbers of field stars with metallicity (Youakim et al. 2020).

The presence of unbound, chemically unique second generation globular cluster stars in the stellar halo of the Milky Way suggests that clusters as a population have lost 50+% of their stellar mass over their lifetime (Koch et al. 2019). In the tidally limited cluster framework advocated in this paper a large mass loss for halo clusters requires that they were formed at greater than redshift 4, Figures 2 and 4, when tides were sufficiently strong to drive substantial mass loss.

Some of the very rare extremely metal poor stars are in a thin distribution aligned with the Milky Way disk (Sestito et al. 2019). These stars could be the remnant of a very early gas disk of the Milky Way, or, the result of late infall of dwarf galaxy sufficiently massive and dense to be pulled into the plane of the current disk (Toth, & Ostriker 1992; Walker et al. 1996; Huang, & Carlberg 1997; Velazquez, & White 1999). The distribution of extremely metal poor stars will be affected by the buildup of the baryonic galaxy, which is another motivation to search for such stars at distances in the halo that are not significantly affected by the baryonic galaxy.

Observations at high redshift will eventually be able to measure the circular velocities of the dark matter halos containing globular clusters. The circular velocities of the sub-halos are observational quantities that are straightforward to compare to the same measurements in the simulations. The low mass sub-halos have a mean peak circular velocity of $12 - 18 \text{ km s}^{-1}$, comparable to the Fornax dwarf (Walker et al. 2006), although the clusters here are much denser. The high mass sub-halos have $26 - 55 \text{ km s}^{-1}$. The clusters are inserted about 20-30% of this characteristic radius, so have a initial velocity typically 30-50% of the peak circular velocity of a halo.

7. CONCLUSIONS

The dynamical simulations of tidally limited star clusters reported here find that clusters that formed in the sub-halos present at redshift 8 have orbits in the resulting Milky Way-like halo that depend on the mass of the dark halo in which they formed. The clusters that are formed in the more massive halos are deposited deep in the potential of the forming galactic halo, where the strong tides cause a lot of mass loss and leave the clusters at such small orbital radii that a baryonic bulge and disk would likely completely disperse the clusters. On the other hand, the clusters formed in the lower mass halos are spread out over the galaxy in a distribution which is similar or somewhat more extended than the dark matter density. The peak circular velocity of the lower mass halos is 12 - 18 km s⁻¹. The same starting procedure at redshift 4.6 leads to negligible differences between the radial distribution of stars from the clusters in low and higher mass sub-halos.

The sizes of the simulated clusters are limited by by their tidal radii, which are dependent on the tidal field present at the starting time and location. The clusters formed at higher redshifts will, on the average, be in stronger tidal fields and will be the relatively smaller clusters. The simulations here find that tidally clusters need to have been formed at $z \gtrsim 4$ to be a reasonable match to the Milky Way distribution of sizes. Together these results indicate that a large fraction of the halo clusters formed at z > 6.

The simulations suggest that many of the first globular clusters to form, which will be the most metal poor, will be fairly evenly spread over the entire galactic halos. Those that are dissolved could be indirectly found through the ongoing search for stellar streams. As shown in the bottom left panel of Figure 1, stellar streams from the high redshift clusters are visible in the inner galaxy and out to 100 kpc.

This research was supported by NSERC of Canada. Computations were performed on the niagara supercomputer at the SciNet HPC Consortium. SciNet is funded by: the Canada Foundation for Innovation; the Government of Ontario; Ontario Research Fund - Research Ex-

cellence; and the University of Toronto.

REFERENCES

- Aarseth, S. J. 1999, PASP, 111, 1333
- Baumgardt, H., Parmentier, G., Gieles, M., et al. 2010, MNRAS, 401, 1832
- Beasley, M. A., Leaman, R., Gallart, C., et al. 2019, MNRAS, 487, 1986

- 487, 1986 Binney, J., & Tremaine, S. 2008, Galactic Dynamics: Second Edition, Princeton University Press Boylan-Kolchin, M. 2018, MNRAS, 479, 332 Brodie, J. P., & Strader, J. 2006, ARA&A, 44, 193 Bromm, V., & Clarke, C. J. 2002, ApJ, 566, L1 Brook, C. B., Di Cintio, A., Knebe, A., et al. 2014, ApJ, 784, L14 Calura, F., D'Ercole, A., Vesperini, E., et al. 2019, MNRAS, 489, 3269

- Carlberg, R. G. 2002, ApJ, 573, 60 Carlberg, R. G. 2018, ApJ, 861, 69 Carlberg, R. G. 2020, ApJ, 889, 107 Chandar, R., Fall, S. M., Whitmore, B. C., et al. 2017, ApJ, 849, 128
- Cole, D. R., Dehnen, W., Read, J. I., et al. 2012, MNRAS, 426, 601
- D'Antona, F., Vesperini, E., D'Ercole, A., et al. 2016, MNRAS, 458, 2122

- 458, 2122
 Da Costa, G. S., Bessell, M. S., Mackey, A. D., et al. 2019, MNRAS, 489, 5900
 Fall, S. M., & Zhang, Q. 2001, ApJ, 561, 751
 Georgiev, I. Y., Puzia, T. H., Goudfrooij, P., et al. 2010, MNRAS, 406, 1967
 Gnedin, O. Y., & Ostriker, J. P. 1999, ApJ, 513, 626
 Gnedin, O. Y., & Ostriker, J. P., & Tremaine, S. 2014, ApJ, 785, 71
 Grillmair, C. J., & Dionatos, O. 2006, ApJ, 641, L37
 Gunn, J. E. 1980, Globular Clusters, ed. D. Hanes & B. Madore (Cambridge: Cambridge Univ. Press). p 301

- Gunn, J. E. 1980, Globular Clusters, ed. D. Hanes & B. Madore (Cambridge: Cambridge Univ. Press), p 301
 Harris, W. E. 1996, AJ, 112, 1487
 Hartwick, F. D. A. 1976, ApJ, 209, 418
 Hartwick, F. D. A. 2018, Research Notes of the American Astronomical Society, 2, 204
 Hernquist, L. 1990, ApJ, 356, 359
 Howard, C. S., Pudritz, R. E., Sills, A., et al. 2019, MNRAS, 486, 1146 1146
- Huang, S., & Carlberg, R. G. 1997, ApJ, 480, 503
 Hudson, M. J., Harris, G. L., & Harris, W. E. 2014, ApJ, 787, L5
 Ibata, R. A., Malhan, K., & Martin, N. F. 2019, ApJ, 872, 152
 Katz, H., & Ricotti, M. 2013, MNRAS, 432, 3250
 Koch, A., Grebel, E. K., & Martell, S. L. 2019, A&A, 625, A75

- King, I. R. 1966, AJ, 71, 64Kruijssen, J. M. D. 2019, MNRAS, 486, L20
- Kruijssen, J. M. D. 2019, MNRAS, 486, L20
 Lada, C. J., & Lada, E. A. 2003, ARA&A, 41, 57
 Lahén, N., Naab, T., Johansson, P. H., et al. 2019, ApJ, 879, L18
 Lamers, H. J. G. L. M., Kruijssen, J. M. D., Bastian, N., et al. 2017, A&A, 606, A85
 Larsen, S. S., Strader, J., & Brodie, J. P. 2012, A&A, 544, L14
 Leaman, R., VandenBerg, D. A., & Mendel, J. T. 2013, MNRAS, 436, 122
 Madau, P., Diemand, J., & Kuhlen, M. 2008, ApJ, 679, 1260-1271
 Madau, P., Lupi, A., Diemand, J., et al. 2020, ApJ, 890, 18
 Martell, S. L., Smolinski, J. P., Beers, T. C., et al. 2011, A&A, 534, A136

- 534, A136 Oh, K. S., Lin, D. N. C., & Richer, H. B. 2000, ApJ, 531, 727 Peebles, P. J. E., & Dicke, R. H. 1968, ApJ, 154, 891

- Planck Collaboration, Adam, R., Aghanim, N., et al. 2016, A&A, 596, A108
- Phipps, F., Khochfar, S., Varri, A. L., et al. 2019, arXiv e-prints, arXiv:1910.09924
- Renzini, A., D'Antona, F., Cassisi, S., et al. 2015, MNRAS, 454, 4197
- Renzini, A. 2017, MNRAS, 469, L63

- Renzini, A. 2017, MNRAS, 469, L63
 Ricotti, M. 2002, MNRAS, 336, L33
 Roederer, I. U., & Gnedin, O. Y. 2019, ApJ, 883, 84
 Searle, L., & Zinn, R. 1978, ApJ, 225, 357
 Sestito, F., Martin, N. F., Starkenburg, E., et al. 2019, arXiv e-prints, arXiv:1911.08491
 Spitzer, L. 1987, Dynamical evolution of globular clusters, Princeton, NJ, Princeton University Press, 1987, 191 p.
 Springel, V. 2005, MNRAS, 364, 1105
 Toth, G., & Ostriker, J. P. 1992, ApJ, 389, 5
 Trenti, M., Padoan, P., & Jimenez, R. 2015, ApJ, 808, L35
 Tumlinson, J. 2010, ApJ, 708, 1398
 VandenBerg, D. A., Brogaard, K., Leaman, R., et al. 2013, ApJ, 775, 134 775, 134

- Velazquez, H., & White, S. D. M. 1999, MNRAS, 304, 254
 Walker, I. R., Mihos, J. C., & Hernquist, L. 1996, ApJ, 460, 121
 Walker, M. G., Mateo, M., Olszewski, E. W., et al. 2006, AJ, 131, 2114
- Yoon, J., Beers, T. C., Tian, D., et al. 2019, ApJ, 878, 97
- Youakim, K., Starkenburg, E., Martin, N. F., et al. 2020, MNRAS, 492, 4986