Modelling the strongest silicate emission features of local type 1 AGN

M. Martínez-Paredes, ¹ O. González-Martín, ² D. Esparza-Arredondo, ² M. Kim, ¹ A. Alonso-Herrero, ³ Y. Krongold, ⁴ T. Hoang, ¹ C. Ramos Almeida, ^{5,6} I. Aretxaga, ⁷ D. Dultzin, ⁴ and J. Hodgson ¹

¹Korea Astronomy and Space Science Institute 776, Daedeokdae-ro, Yuseong-gu, Daejeon, Republic of Korea (34055)
 ²Instituto de Radioastronomía y Astrofísica UNAM Apartado Postal 3-72 (Xangari), 58089 Morelia, Michoacán, Mexico
 ³Centro de Astrobiología, CSIC-INTA, ESAC Campus, E-28692 Villanueva de la Caada, Madrid, Spain
 ⁴Instituto de Astronomía UNAM, México, CDMX., C.P. 04510
 ⁵Instituto de Astrofísica de Canarias (IAC),E-38205 La Laguna, Tenerife, Spain

⁶Departamento de Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain

⁷Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Luis Enrrique Erro 1, Sta. Ma. Tonantzintla, Puebla, Mexico

(Received September ??, 2019; Revised November ??, 2019; Accepted December ??, 2019)

Submitted to ApJ

ABSTRACT

We measure the 10 and $18\mu \text{m}$ silicate features in a sample of 67 local (z < 0.1) type 1 active galactic nuclei (AGN) with available Spitzer spectra dominated by non-stellar processes. We find that the $10\mu \text{m}$ silicate feature peaks at $10.3^{+0.7}_{-0.9}\mu \text{m}$ with a strength ($\text{Si}_p = \ln f_p(\text{spectrum})/f_p(\text{continuum})$) of $0.11^{+0.15}_{-0.36}$, while the $18\mu \text{m}$ one peaks at $17.3^{+0.4}_{-0.7}\mu \text{m}$ with a strength of $0.14^{+0.06}_{-0.06}$. We select from this sample sources with the strongest $10\mu \text{m}$ silicate strength ($\sigma_{Si_{10}\mu m} > 0.28$, 10 objects). We carry out a detailed modeling of the IRS/Spitzer spectra by comparing several models that assume different geometries and dust composition: a smooth torus model, two clumpy torus models, a two-phase medium torus model, and a disk+outflow clumpy model. We find that the silicate features are well modeled by the clumpy model of Nenkova et al. 2008, and among all models those including outflows and complex dust composition are the best (Hoenig et al. 2017). We note that even in AGN-dominated galaxies it is usually necessary to add stellar contributions to reproduce the emission at the shortest wavelengths.

Keywords: (galaxies:) active: general galaxies: infrared: galaxies

1. INTRODUCTION

A dusty torus surrounds the central engine of active galactic nuclei (AGN) on a scale of few pc (e.g., Krolik & Begelman 1988; Antonucci 1993; Robson et al. 1995; Peterson 1997). It shines at infrared (IR) wavelengths between 1 to 1000 μ m peaking at around 20 μ m (e.g., Sanders et al. 1989; Elvis et al. 1994). This emission is the result of the IR re-radiation of optical-UV light generated around the central black hole (BH) that has been absorbed by the dusty torus (Neugebauer et al. 1979). The main observational components that describe it are the slope of the spectral energy distribution (SED) between 1-8 μ m, and the strength of the silicate features around 10μ m and 18μ m, that are produced within the warm dust of the torus that the AGN directly illuminates. The silicate features have been observed mostly in emission with the mid-infrared (MIR) infrared spectrometer (IRS) Spitzer in type 1 AGN (Siebenmorgen et al. 2005; Hatziminaoglou et al. 2015). In these AGN the 10μ m silicate feature is broader and peaks at much longer wavelengths ($\sim 10.0-11.5 \mu$ m) than the "normal" silicate emission feature of the Galactic inter-stellar medium (ISM) (e.g., Hao et al. 2005; Siebenmorgen et al. 2005; Sturm et al. 2005; Li et al. 2008). This suggests different silicate compositions, such as a different proportion and/or size of grains (e.g., Shi et al. 2006; Li et al. 2008).

Martínez-Paredes et al. (2017) used the starburst-subtracted IRS/Spitzer spectra of 20 quasi stellar objects (QSOs) between ~ 7.5 to 15 μ m plus the unresolved near-infrared (NIR) emission to constrain the parameters of the dusty torus using the CLUMPY models of Nenkova et al. (2008a,b). They noted that in most cases the spectral range around 8μ m is underestimated by the CLUMPY models and that trying to fit this part of the spectrum resulted in a bad fit of the silicate feature at 10μ m. Furthermore, in most cases the peak of the 10μ m silicate feature was shifted from the model location.

Considering that the shape and peak of the silicate features are strongly correlated with the properties of the dust (e.g., Draine et al. 2007; Sirocky et al. 2008), in this work we use five of the most popular torus models to investigate how well they reproduce the strongest silicate emission features observed in type 1 AGN. Recently, González-Martín et al. (2019a) compared these models and found that they can be distinguished according to the continuum slopes and silicate strengths, and in a second work (González-Martín et al. 2019b) they used MIR spectroscopy data of type 1 and type 2 AGNs in the Swift/BAT survey to investigate how these models reproduce their spectral continuum. We focus our study on type 1 AGN with strong silicate emission features because: 1) for type 2 AGN, the inclusion of a fraction ($\sim 50\%$) of silicate grains, either based on Draine (2003) or Ossenkopf et al. (1992) are enough to reproduce the observed silicate features in absorption (see e.g., Ramos Almeida et al. 2011; Alonso-Herrero et al. 2011; Martínez-Paredes et al. 2015); and 2) the silicate features are well above the continuum in the low resolution *Spitzer* spectra, which support the interpretation that the silicate features are due to AGN activity.

In this paper we aim at exploring how accurately smooth, clumpy, two-phase, and outflow torus models reproduce the IRS/Spitzer spectra of these objects, and analyzing the results in terms of physical differences like the geometry and dust composition assumed. The first model that we consider is the smooth torus model of (Fritz et al. 2006, Fritz06 hereafter), which assumes a continuous distribution of dust composed of graphite and silicate grains in almost equal percentages. The second model is the CLUMPY torus model of (Nenkova et al. 2008a,b, Nenkova08), which assumes a standard ISM dust composition of 53% silicates and 47% graphite with sizes between 0.025 and 0.25 μ m. The third torus model is clumpy (Hönig & Kishimoto 2010b, Hoenig10) and assumes three dust components that include the standard ISM composition, the standard ISM composition with larger grains (0.1 - 1.0 μ m), and a larger fraction of graphite grains (30% silicates and 70% graphite, with sizes between 0.05 and 0.25 μ m). The fourth model (Hönig & Kishimoto 2017, Hoenig17) has two components: a dusty clumpy disk that takes into account the emission from the hotter dust close to the central engine; and a hollow cone formed by a wind of clumpy dusty clouds elongated toward the polar direction. This model assumes that the wind is originated close to the sublimation zone of the dusty disk. Because of that, the dust in the wind has similar properties to the dust in the sublimation zone.

The paper is organized as follows: in Section 2 we present our sample selection and data; in Section 3 we describe the models; in Section 4 we present the analysis; and in Section 5 the discussion. Our summary and conclusions are presented in Section 6. We adopt the following cosmology: $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.30 \text{ and } \Omega_{\Lambda} = 0.70$.

2. THE SAMPLE AND DATA

2.1. The silicate dominated local AGN

We use the latest version of the AGN catalog of Véron-Cetty & Véron (2010) and the sample of low luminosity type 1 AGN in Mason et al. (2012) to select AGN that have redshifts z < 0.1 and available 2D low-resolution IRS/Spitzer spectra in the CASSIS database (v6., Lebouteiller et al. 2011). We select those objects with spectra extracted as point-sources that cover the $5-35\mu$ m spectral range. The first criterion allows us to probe the MIR emission from the central region of the AGN, and the second one allows us to study both the 10 and 18μ m silicate features. As an exception, we include NGC 3998, a strong silicate emitter (e.g., Mason et al. 2012), which has a low-resolution spectrum in the ~ 7.5 -14.5 μ m range and a high-resolution spectrum in the ~ 14 -35 μ m range. In order to ensure that the emission in the IRS/Spitzer spectra is mostly dominated by dust heated by the AGN, we decompose them into their stellar, ISM and AGN emission (dust heated by AGN) components at $\sim 5-15\,\mu$ m, using the spectral decomposition tool Deblendirs (Hernán-Caballero et al. 2015). For more details on the spectral decomposition we refer the reader to Appendix A. We select only those objects (67) for which the spectral contribution due to AGN is > 80% (see Figure 1). Hereafter, we refer to the spectra of these objects as the AGN-dominated IRS/Spitzer spectra. Note that we refer here to the original spectra, not to the component obtained from the decomposition.




Figure 1. Properties of the sample. **Left:** distribution of redshifts. **Right:** distribution of AGN contributions to the IRS/Spitzer spectra. The black line represents the full sample, and the blue line is a subsample defined in section 2.1.

We obtained the reduced low resolution ($R \sim 60-127$) IRS/Spitzer spectra from the CASSIS database (v6., Lebouteiller et al. 2011). The spectra include the SL1 ($\lambda \sim 7.4-14.5\,\mu\mathrm{m}$) and SL2 ($\lambda \sim 5.2-7.7\,\mu\mathrm{m}$) modules with a slit-width of 3.6 arcsec, and the LL1 ($\lambda \sim 19.9-39.9\mu\mathrm{m}$) and LL2 ($\lambda \sim 13.9-21.3\,\mu\mathrm{m}$) modules with a slit-width of 10.5 arcsec (Werner et al. 2004; Houck et al. 2004). We use the final stitched spectra between $5-35\,\mu\mathrm{m}$. The different module spectra were stitched by scaling the LL and SL1 flux modules to the shortest module SL2 flux.

2.3. Measuring the silicate emission features

To measure the silicate emission features we start by interpolating a local continuum between both sides of the emission line complexes. We inspect each spectrum visually and choose three bands in which to measure the continuum. Band 1 is located at the blue extreme of the $10\mu \rm m$ silicate feature, while band 3 is located at the red extreme of the $18\mu \rm m$ silicate feature, and band 2 is located between the two silicates features. Band 1, 2, and 3 are located within the 7.8-8.5, 13.5-14.0 and $20.0-21.0\,\mu \rm m$ ranges.

In order to better define uncertainties in the continuum definition, which might be affected by the presence of Polycyclic Aromatic Hydrocarbon (PAH) molecular emission lines around 7μ m, and in some cases by the artificial "teardrop" feature around 14μ m, we trace fiducial mean values of each continuum band by bootstrapping on the measured fluxes (hatched pink regions in Figure 2). We generate 100 continuum values, considering the uncertainties, between band 1 and 2, and between band 2 and 3. We randomly associate shorter and longer wavelength mean continuum values to generate linear continua below the silicate features. We note that fitting a spline continuum gives similar results. The dark blue solid lines in Figure 2 are bootstrapped local continua derived from the continuum bands and they define the regions where we measure the strength of the silicate features.

Considering that the peak of the silicate features vary from object to object, we choose to measure the strength of the features at the wavelengths where they peak. However, for simplicity we still call them the 10 and 18μ m silicate features. The silicate strength is, hence, defined as the silicate peak relative to the continuum, at the wavelength where the silicate feature peaks (see e.g., Hao et al. 2007), according to the equation:

$$Si_p = \ln \frac{f_p \text{ (spectrum)}}{f_p \text{ (continuum)}}.$$
 (1)

In Table 1 we list the mean and the 68 percent intervals for the objects with the largest silicate feature strengths. In Appendix B (Table 4) we list the silicate emission strengths for the full sample. On average, we find that type

¹ An excess of emission present in the 2D SL1 spectrum of some objects.




Figure 2. IRS/Spitzer spectrum of PG 2214+139 (light blue solid line). The red line is the local continuum that follows the broad features of the IRS/Spitzer spectrum. The blue solid lines are the bootstrapped local continua and the vertical pink dashed bars are the bands used to fit the continua around the features. The vertical black dashed-lines indicate the wavelength where the silicate strength is measured. The vertical grey dashed-lines mark other emission lines.

Table 1. Basic properties of objects in the Si-s sample and their silicate feature strengths measured from the IRS/Spitzer spectra. Columns 1 and 2 list the name and type of activity, column 3 gives the redshift, and columns 4 and 5 list the logarithm of the intrinsic hard X-ray luminosity and the corresponding reference. Columns 6 and 7 list the wavelength where the 10μ m silicate emission feature peaks, and the 10μ m silicate strength, and columns 8 and 9 are are the wavelength and strength of the 18μ m silicate emission feature.

| Name | Activity ^a | \mathbf{z} | $log L_{X(2-10 \text{keV})}$ | Ref. | λ_p | $\mathrm{Si}_{10\mu\mathrm{m}}$ | λ_p | $\mathrm{Si}_{18\mu\mathrm{m}}$ |
|--------------|-----------------------|--------------|------------------------------|------|--------------------|---------------------------------|--------------------|---------------------------------|
| | | | ${\rm erg~s^{-1}}$ | | $(\mu \mathrm{m})$ | | $(\mu \mathrm{m})$ | |
| NGC7213 | LINER | 0.0058 | 42.2 | 6 | 10.7 ± 0.1 | 0.52 ± 0.05 | 17.3 ± 0.1 | 0.23 ± 0.04 |
| PG2304+042 | Sy1.2 | 0.0420 | 43.4^{*} | 9 | 10.3 ± 0.1 | 0.30 ± 0.06 | 17.3 ± 0.1 | 0.33 ± 0.08 |
| PKS0518-45 | LINER | 0.0420 | 44.9 | 8 | 10.4 ± 0.2 | 0.31 ± 0.05 | 17.6 ± 0.1 | 0.18 ± 0.04 |
| PG0844 + 349 | Sy1/QSO | 0.0640 | 43.7 | 1 | 10.3 ± 0.1 | 0.33 ± 0.06 | 16.7 ± 0.1 | 0.18 ± 0.04 |
| PG1351+640 | Sy1.5/QSO | 0.0882 | 43.1 | 6 | 9.8 ± 0.1 | 0.52 ± 0.03 | 18.4 ± 0.1 | 0.14 ± 0.03 |
| PG2214+139 | Sy1.0/QSO | 0.0658 | 43.8 | 1 | 10.2 ± 0.1 | 0.29 ± 0.04 | 17.8 ± 0.2 | 0.14 ± 0.04 |
| PG0804+761 | Sy1/QSO | 0.1000 | 44.5 | 1 | 9.9 ± 0.1 | 0.36 ± 0.03 | 17.3 ± 0.1 | 0.11 ± 0.04 |
| OQ208 | Sy1.5 | 0.0766 | 40.8 | 3 | 10.2 ± 0.1 | 0.34 ± 0.05 | 16.5 ± 0.1 | 0.12 ± 0.04 |
| NGC4258 | $LINER^b$ | 0.0015 | 40.9 | 5 | 11.0 ± 0.1 | 0.29 ± 0.05 | 17.3 ± 0.1 | 0.17 ± 0.04 |
| NGC3998 | LINER | 0.0035 | 41.2 | 4 | 10.7 ± 0.1 | 0.36 ± 0.01 | $16.3\pm0.1^*$ | 0.23 ± 0.02 |

Note.-*This value needs to be used carefully, since we are using the high angular resolution spectrum to fix the band 3 and measure the 18μm silicate feature strength. References: NED and Véron-Cetty & Véron (2010), Mason et al. (2012).

*Estimated from (15-150)+keV X-ray luminosity assuming a spectral power law with an index α = 1.8. References for hard X-ray luminosity: 1: Zhou & Zhang (2010); 2: Sambruna et al. (2011); 3: Ueda et al. (2005); 4: Younes et al. (2011); 5: Cappi et al. (2006); 6: Bianchi et al. (2009); 7:Brightman & Nandra (2011); 8: Winter et al. (2012); 9: Tueller et al. (2010); 10: Cusumano et al. (2010).

1 AGN have a $10\mu \text{m}$ silicate strength $Si_{10.3\mu \text{m}} = 0.13^{+0.15}_{-0.37}$ that peaks at $10.3^{+0.7}_{-0.9}\mu \text{m}$, and a $18\mu \text{m}$ silicate strength $Si_{17.3\mu \text{m}} = 0.14^{+0.06}_{-0.06}$ that peaks at $17.3\mu \text{m}$.

In most cases, our measurements, which were performed using the AGN-dominated *Spitzer* spectra, produce similar results to previous works (Hao et al. 2005; Thompson et al. 2009; Sirocky et al. 2008; Mendoza-Castrejón et al. 2015)




Figure 3. Distribution of the 10μ m silicate strength feature (see Table 1). The solid and dashed black lines indicate the mean and its 1σ confidence interval.

| Model name | Geometry | Dust composition | Scattering and absorption | Grain si | $zes^*(\mu m)$ |
|-------------|-----------------------|----------------------------------|------------------------------------|--------------|----------------|
| | | | coefficients | Graphites | Silicates |
| Fritz06 | Smooth torus | Graphite and Silicates | graphite and silicates | 0.005-0.25 | 0.025-0.25 |
| | | | Laor & Draine (1993) | | |
| Nenkova08 | Clumpy torus | Standard ISM | graphites, Draine (2003) | (0.005-0 | .01)-0.25 |
| | | | silicates, Ossenkopf et al. (1992) | | |
| Hoenig10 | Clumpy torus | Standard ISM | graphites, Draine (2003) | (0.005-0 | .01)-0.25 |
| | | +standard ISM with large grains | silicates, Draine (2003) | | 0.1 - 1.0 |
| | | +graphite dominated dust | Ossenkopf et al. (1992) | | 0.05 - 0.25 |
| | | (70% graphite and 30% silicates) | | | |
| Hoenig17 | Disk+Outflow | Standard ISM | like Hoenig10 | (0.005-0 | .01)-0.25 |
| | | +standard ISM with large grains | | | |
| Stalevski16 | Two phase media torus | Graphite and Silicates | Laor & Draine (1993) | 0.005 - 0.25 | 0.005 - 0.25 |
| | | | Li & Draine (2001) | | |

Table 2. Dust properties of the models.

Note. The standard ISM is composed by 47% of graphite and 53% silicates. *Assuming MNR distribution (Mathis et al. 1977). The numbers inside the parenthesis indicate the minimal range of sizes for the smaller grains.

(see Tables 4). Additionally, we compare our measurements of the silicate strengths with the values obtained using DEBLENDIRS and find that they are similar within the uncertainties (see Figure 13 in Appendix A).

$2.4. \ Sample \ selection$

We build our final sample by selecting only those objects that show the strongest silicate emission features (see Table 1). From the original sample of 67 local type 1 AGNs we select those objects with the largest 10μ m silicate strength ($\sigma_{Si_{10\mu m}} > 0.28$, see Figure 3). The final sample is composed of 10 objects: six Seyfert (Sy) galaxies, and four low-ionization nuclear emission-line region (LINER) galaxies. Four of the Seyfert galaxies are also classified as PG QSOs. Note that we are not selecting the sample according to the type of AGN. We list the type only to give information about their basic properties.

Hereafter, we will refer to this sample as the type 1 AGN strong Silicate selected sample (Si-s). The sample spans a range of hard (2-10 keV) X-ray unobscured luminosities between $\sim 10^{41}$ erg s⁻¹ and $\sim 10^{45}$ erg s⁻¹.

3. DUSTY TORUS MODELS

There are two types of torus models widely used in the literature: smooth torus models and clumpy torus models. Smooth models assume a continuous distribution of the dust in the torus (Pier & Krolik 1992; Granato et al. 1997;




Figure 4. Cartoons of torus and disk+outflow models: a) smooth models (Fritz et al. 2006), b) clumpy models (Nenkova et al. 2008a,b; Hönig & Kishimoto 2010b) and, c) disk + outflow clumpy models (Hönig & Kishimoto 2017). For smooth models: viewing angle i (degrees) = [0-90], angular width Θ (degrees) = [20-60], parameter of the dust distribution $\gamma = [0-6]$, and the outer-to-inner radius ratio $Y = R_{\text{out}}/R_{\text{in}} = [10-150]$. For clumpy models: viewing angle i (degrees) = [0-90], angular width σ (degrees) = [15-70], number of clouds along the equatorial line $N_0 = [1-15]$ (for Nenkova08) and [2.5-10] (for Hoenig10), parameter of radial distribution ($\propto r^{-q}$, in Nenkova08) q = [0-2.5], and parameter of radial distribution ($\propto r^{a}$, in Hoenig10) a = [-2.5-0], the outer-to-inner radius ratio $Y = R_{\text{out}}/R_{\text{in}} = [5-100]$ (for Nenkova08), and $R_{\text{max}} = 170R_{\text{in}}$ (for Hoenig10). For disk+outflow clumpy models: viewing angle i (degrees) = [0-90], number of clouds along the equatorial line $N_0 = [2.5-10]$, half-opening angle of the outflow σ_{Θ} (degrees) = [30-45], and angular width of the disk (degrees) $\Theta_{\text{w}} = [7.0-15]$. For a more detailed comparison between the parameters of the models see González-Martín et al. (2019a,b).

Efstathiou & Rowan-Robinson 1995; Fritz et al. 2006), while clumpy models assume a distribution of dusty clouds or clumps in a toroidal structure (Dullemond & van Bemmel 2005; Nenkova et al. 2008a,b; Hönig & Kishimoto 2010b, 2017).

For simplicity, smooth models assumed a continuous distribution of the dust in the torus, so the temperature of the dust decreases monotonically with the distance from the central BH. According to this type of model only type 1 AGN produce $10\mu m$ silicate features in emission and type 2 AGN produce silicate features in absorption, implying an edge-on orientation with respect to the observer. In type 2 AGN, therefore, the outer region of the torus hides the emission produced by the hotter and warmer dust in the inner part of the torus. However, the silicate depth that this model predicts is larger than observed in type 2s, and it fails to explain silicate features in emission in type 2 QSOs (e.g., Sturm et al. 2006).

Conversely, clumpy models (e.g., Nenkova et al. 2008a,b) never produce very deep absorption silicate features, and they predict silicate features in emission in type 1 and type 2 AGN for a large combination of parameters (see e.g., Nikutta et al. 2009). A third type of model combines clumpy and smooth properties to produce a two-phase medium dusty torus model (Stalevski et al. 2012, 2016). This model assumes a distribution of dusty clumps with constant and high density, embedded in a smooth dusty component of low density.

It was motivated by observational evidence that suggests that dust around AGN seems to have a multiphase filamentary structure (Wada et al. 2009; Wada 2012).

A more recent model proposes a geometry composed by a compact and geometrically thin disk in the equatorial region plus an extended elongated polar structure of clumpy dust, that is cospatial with the outflow of the AGN (Hönig & Kishimoto 2017). This geometry is inferred from interferometric observations of nearby Seyfert galaxies, where the bulk of the MIR emission originates from the polar region rather than the equatorial plane (e.g., Raban et al. 2009; Hönig et al. 2012, 2013; Tristram et al. 2014; López-Gonzaga, & Jaffe 2016).

In this paper, we use the AGN-dominated IRS/Spitzer spectra of our Si-s sample of type 1 AGN to investigate which models are able to better reproduce the peak and shape of the strongest silicate emission features observed. In the following subsections we briefly describe the main geometrical and physical properties of the models and Table 2 summarizes the different geometries and dust compositions assumed for each model.

3.1. Smooth torus model of Fritz et al. (2006)

This model is represented by a flared disc delimited by the inner and outer torus radius (e.g., Efstathiou & Rowan-Robinson 1995). The inner radius (R_{\min}) is defined by the sublimation temperature of dust grains (1500 K) under the strong radiation produced by the central engine. The size of the torus is determined by the outer radius $(Y = R_{\text{out}}/R_{\text{in}}, \text{where } Y \text{ is a free parameter})$, and the angular width of the torus Θ (see the cartoon (a) in Figure 4). This model assumes that the dust is composed of graphite grains (53 per cent) with sizes from 0.005 to $0.25\mu\text{m}$, and silicate grains (47 per cent) with sizes from 0.025 to $0.25\mu\text{m}$. The grain sizes are distributed according to Mathis et al. (1977). For the two species they used the scattering and absorption coefficients given by Laor & Draine (1993). The dust is illuminated by an isotropic central point-like emitting source, which is represented by a broken power law of the form $\lambda L_{\lambda} \propto \lambda^{\alpha}$, with $\alpha = 1.2$ if $0.001 < \lambda < 0.03 \,\mu\text{m}$, $\alpha = 0$ if $0.03 < \lambda < 0.125 \,\mu\text{m}$, and $\alpha = 0.5$ if $0.125 < \lambda < 20.0 \,\mu\text{m}$. Other parameters of the model are the viewing angle i, the polar (γ) and radial (β) indices of the gas density distribution $\rho(r,\Theta) \propto r^{\beta} e^{-\gamma \times \cos(\Theta)}$ within the torus, and the optical depth $\tau_{9.7 \,\mu\text{m}}$. For a more complete description of this model see Fritz et al. (2006).

3.2. Clumpy torus models of Nenkova et al. (2008a,b)

This type of model assumes a central point-like emitting source surrounded by a toroidal distribution of clouds. The emission of the central source is characterized by a broken power law of the form $\lambda f_{\lambda} \propto \lambda^{1.2}$ for $\lambda \leq 0.01 \mu \text{m}$, $\propto \lambda^{-0.5}$ for $0.1 \mu \text{m} \leq \lambda \leq 1 \mu \text{m}$, $\propto \lambda^{-3}$ for $\lambda \geq 1 \mu \text{m}$, and a constant power-law between 0.01 and $0.1 \mu \text{m}$. Due to its clumpy nature, the central source can directly heat the dust in the inner region of the torus and the dust located at several sublimation radii from the central source. This model assumes standard Galactic spherical dust grains (standard ISM) composed of graphites (47%, Draine 2003) and silicates (53%, Ossenkopf et al. 1992), with a power-law distribution of grain sizes ($\propto a^{-3.5}$), where $a_{\min} = 0.05 - 0.01 \mu \text{m}$ and $a_{\max} = 0.25 \mu \text{m}$, respectively. This model has six free parameters: the viewing angle i, the number of clouds along the equatorial ray N_0 , the angular width σ , the radial extend $Y = R_{\text{outer}}/R_{\text{inner}}$, the index of the radial distribution of clouds q, and the optical depth per cloud τ_V (see cartoon (b) in Figure 4). For a more complete description of this model see Nenkova et al. (2008a,b).

3.3. Clumpy torus models of Hönig & Kishimoto (2010b)

In this case the toroidal distribution of dusty clumps surrounds a central point-like emitting source, that is described by a broken power-law of the form $\lambda f_{\lambda} \propto \lambda$ for $\lambda < 0.03 \mu \text{m}$, $\propto \lambda^{-3}$ for $\lambda > 3 \mu \text{m}$, constant for λ between $0.03 \mu \text{m}$ and $0.3 \mu \text{m}$, and $\propto \lambda^{-4/3}$ for λ between 0.3 and $3 \mu \text{m}$. They modeled the torus following a 3D Monte Carlo radiative transfer simulation. These kinds of simulations fail to properly sample optically thick surface regions with enough grid cells so that each cell is optically thin, which results in underestimating the emission temperature and resulting in a smaller number of model clouds with respect to Nenkova et al. (2008a,b). The dust is composed of three components: a standard ISM component, a standard ISM component with larger grains ($a_{\min} = 0.1 \mu \text{m}$ and $a_{\max} = 1.0 \mu \text{m}$), and intermediate to larger grains ($a_{\min} = 0.05 \mu \text{m}$ and $a_{\max} = 0.25 \mu \text{m}$) with 70% graphite and 30% silicates. The free

parameters that describe this model are the viewing angle i, the number of clouds along the equatorial ray N_0 , the half-opening angle Θ , the index of the radial distribution a, and the optical depth τ_V (see cartoon (b) in Figure 4). For a more complete description of this model see Hönig & Kishimoto (2010b).

3.4. Clumpy disk+outflow models of Hönig & Kishimoto (2017)

This type of model is based on the parametrization of the clumpy torus model of Hönig & Kishimoto (2010b), but instead of adding a blackbody component to take into account the NIR emission, they include a set of different sublimation temperatures for silicate and graphite dust that results in more emission from graphite located in the inner edge of the torus. In this way, when the dust is heated to temperatures > 1200 K smaller silicate grains are destroyed leaving only graphite grains that can be heated up to 1900 K. In the innermost radius only, grains with a size between 0.075 and $1\mu m$ survive.

The clouds are distributed according to a radial power-law $\propto r^a$, where a is the power law index and r the distance from the black hole in units of the sublimation radius $r_{\rm sub}$. They also add a polar outflow, modeled as a hollow cone that can be characterized by the radial distribution of dust clouds in the wind $a_{\rm w}$, the half-opening angle of the wind $(\Theta_{\rm w})$, and the angular width (σ_{Θ}) . Other parameters are the number of clouds along the equatorial ray N_0 , and the scale height in the vertical Gaussian distribution of clouds h in the disk, (see cartoon (c) in Figure 4). For a more complete description of this model see Hönig & Kishimoto (2017).

3.5. Two-phase medium dusty torus models of Stalevski et al. (2016)

This kind of models assumes a distribution of high density dusty clumps embedded in a low density smooth dusty component. This assumption produces both weaker silicate features and a pronounced NIR emission. They assume that the accretion disk in the nucleus radiates as a broken power-law of the form $\lambda L_{\lambda} \propto \lambda^{\alpha}$, where $\alpha = 1.2$ for a spectral range of $0.001 \leq \lambda \leq 0.01~\mu\text{m}$, $\alpha = 0$ for $0.01 < \lambda \leq 0.1~\mu\text{m}$, $\alpha = -0.5$ for $0.1 < \lambda \leq 5~\mu\text{m}$, and $\alpha = -3$ for $5 < \lambda \leq 50~\mu\text{m}$. The dust is distributed following a law that allows a density gradient along the radial (r) and polar (θ) directions, inside a flare disc defined by the inner $(R_{\rm in})$, outer radii $(R_{\rm out})$ and half opening angle. The inner radius is defined by the sublimation temperature of 1500 K for an average dust grain size of 0.05 μ m. They assumed a standard ISM dust composition with optical properties from Laor & Draine (1993) and, Li & Draine (2001). For a more complete description of this model see Stalevski et al. (2012, 2016).

4. ANALYSIS

4.1. Synthetic and observed silicate peak wavelengths and strengths

In this section we explore how well the dusty torus models reproduce the central wavelength and strength of both 10 and 18 μ m silicate features, and the NIR (α_{NIR}) and MIR (α_{MIR}) spectral indexes. In order to make a proper comparison the synthetic and observed central wavelength and strength of both the 10 and 18 μ m silicate features are measured following the same methodology described in Section 2.3, and fixing the bands to the sides of the silicate features between 7-7.5 μ m, 14-15 μ m, and 25 – 26 μ m. The synthetic and observed spectral indexes α_{NIR} and α_{MIR} are measured between 5.5-7.5 μ m, and between 7.5-14.0 μ m, respectively, according to the following definition $\alpha_{2,1} = -log(f_{\nu}(\lambda_2)/f_{\nu}(\lambda_1))/log(\lambda_2/\lambda_1)$, with $\lambda_2 > \lambda_1$ (see e.g., Buchanan et al. 2006).

In Figures 5 and 6 we plot the wavelength of the peak of the both 10 and $18\mu m$ silicate features and the strengths of both silicate features as predicted by the models, and as observed in the AGN-dominated IRS/Spitzer spectra of the Si-s sample. Additionally, we color-coded the objects according to their bolometric luminosity, which we estimate using the hard X-ray luminosity and the relation derived by Marconi et al. (2004) and Alexander, & Hickox (2012).

Since the models are probabilistic in nature, we compare the envelope of measurements performed on the models to the measurements of the Si-s AGN sample. While the Fritz 06, Stalev16 and Nenkova08 models envelope covers the wavelength space (Figure 5) where our Si-s AGN measurements lie, the Hoenig10 and Hoenig17 models leave miss one and two objects, respectively, with the lowest value central wavelength of the $18\mu m$ silicate feature. Curiously, these objects are of low bolometric luminosity (OQ 208 and NGC 4258).

For the silicate strengths (Figure 6) we note that the range of synthetic values sampled by the models mostly match the observations. However, we also note that Hoenig10 and Hoenig17 models show a narrower range of values respect to the other models. Additionally, the Hoenig17 model never predicts both silicates in absorption. Similar results were reported for a larger sample of AGN (González-Martín et al. 2019b). Finally, all the models tend to produce extremely prominent silicate emission features that have not been observed.




Figure 5. Central wavelength of the synthetic and observed peaks of the 10 and $18\mu m$ silicate features. Grey circles represent the values measured in the models, while the stars are the measurements obtained from the AGN-dominated IRS/Spitzer spectra of the Si-s sample. The colours of the stars indicate different range of bolometric luminosities (L_{bol}). White stars are for $logL_{bol} < 42$, light-green stars are for $43 \le logL_{bol} < 44$, blue stars for $44 \le logL_{bol} < 45$, green stars for $45 \le logL_{bol} < 46$, and black star for $logL_{bol} > 46$. Small stars indicate lower luminosities, while larger stars are for higher luminosities.

In Figure 7 we compare the synthetic and observed silicate strength of the $10\mu m$ silicate feature with the α_{NIR} and α_{MIR} spectral indexes. We note that the range of synthetic values sampled by the Fritz06, Hoenig10, and Hoenig17 models mostly match the observations, independently of the bolometric luminosity. However, the Nenkova08 and Stalev16 models miss several of the NIR and MIR spectral indexes observed. Note that NGC 3998 is excluded from plots in Figures 5, 6 and 7 because the short spectral range ($\sim 7.5-14.5\mu m$) cover by the low resolution IRS/Spitzer spectrum of this object.

4.2. Modeling

We apply the torus models using the computational spectral fitting tool XSPEC, which is part of the HEASOFT² software. These models were recently converted into the XSPEC format in order to fit data in a similar way (see section 2 in González-Martín et al. 2019a). We also use the set of synthetic stellar and empirical HII components (ISM) previously converted into XSPEC format (González-Martín et al. 2019a). The former corresponds to a stellar population of 10¹⁰ years and solar metallicity from the stellar spectral libraries of Bruzual & Charlot (2003), while the empirical HII components are average starburst templates from Smith et al. (2007).

² https://heasarc.gsfc.nasa.gov




Figure 6. Silicate strength of the 10 and $18\mu m$ silicate features. Symbols and colours are as in Figure 5.

We model the AGN-dominated IRS/Spitzer spectrum of each object in the Si-s sample, using each one of the four torus family models and the disk+outflow models. Additionally, we add a stellar synthetic and/or HII empirical component to investigate in which cases adding one or both components really improves the fitting. We also add foreground extinction to the torus models by using the ZDUST component (Pei 1992).

In detail, we start converting the IRS/Spitzer spectra into the XSPEC format, loading the data into XSPEC, removing the parts of the spectra dominated by emission lines, and modeling the spectra assuming the following component combinations:

- 1: AGN dust emission: torus models or disk+outflow model
- 2: AGN + stellar
- 3: AGN + HII
- 4: AGN + stellar + HII

For each step, we save the reduced $\chi^2_{\rm red}$, the parameters of the model with their uncertainties, and the emission contribution of each component to the total emission of the IRS/Spitzer spectrum between 5.5 to 30 μ m. In those cases where $\chi^2_{\rm red} > 2$ we reported the case as "non-modeled". We perform this procedure for each torus model, resulting in 20 spectral fits per object. Figure 8 shows, for each torus model, the number of spectral fits obtained, using one, two, or three components. Those cases in which none of the component combinations (1,2,3, or 4) are able to model the spectrum with a $\chi^2_{\rm red} < 2$ are called "non-modeled" spectral fits (see Table 5 in Appendix C). In the next section we investigate which model best fits the peak and strength of both silicate emission features in the Si-s sample.




Figure 7. Synthetic and observed silicate strength of the $10\mu m$ silicate feature, and the NIR α_{NIR} and MIR α_{MIR} , spectral indexes. Symbols and colours are as in Figure 5.

4.3. Model comparison

In most cases, we find that the same object can be modeled assuming one (AGN), two (AGN+stellar or AGN+HII), or three components (AGN+Stellar+HII) with a $\chi^2_{\rm red} \sim 1$. For each object and torus model we use the statistical f-test, which allows us to evaluate in which cases the addition of a new component improves the fit from a statistical point of view. We add a new component when the f-test probability is $< 10^{-4}$. The computational tool XSPEC includes the FTEST command line, which allows us to calculate the f-statistic and its probability, when new and old values of χ^2 and the degrees of freedom (dof) are provided. As an example, in Figure 9 we show the best fit of the IRS/Spitzer spectrum of PG 0844+349 for each torus model. For this particular object the Fritz06, Nenkova08, and Stalev16 models need an additional stellar component to fit the spectrum, while the Hoenig10 and Hoenig17 models need also the HII component. Additionally, for this object, we observe that clumpy and disk+outflow models produce the flattest residuals within the uncertainties and smaller $\chi^2_{\rm red}$ (see bottom panel in Figure 9 and Table 3).

In Figure 10 we plot the histogram of the fractional contribution of each spectral component. The AGN component dominates the emission in all cases, which is expected due to our selection of AGN-dominated IRS/Spitzer spectrum sources.

Fritz06 models are able to reproduce the IRS/Spitzer spectra in seven of 10 objects. The AGN component contributes more than 93%. The Nenkova08 models reproduce the IRS/Spitzer spectra of 10 objects, in most cases with an AGN contribution > 81%, except for NGC4258 for which the AGN component contributes 66%. For Hoenig10 models we find that four of the objects can be modeled with an AGN contribution > 80%, and three other (PG2214+139,




Figure 8. Fitting components for each torus model: from top to bottom the horizontal bars show the number of fits with one (AGN), two (AGN + stellar or AGN + HII), and/or three (AGN + stellar + HII) components for each dusty torus model. Additionally, for each torus model, the two last horizontal bars represent the total number of cases modeled and not modeled ($\chi^2_{\text{red}} > 2$), respectively.

PG0804+761, and NGC4258) with a contribution between 67 and 76 %. The Hoenig17 models are able to fit seven of the objects with an AGN contribution > 87%, and NGC 4258 with an AGN component that contributes > 62%. Finally, in the case of two-phase medium models of Stalevski et al. (2016) only five of the 10 objects can be reproduced with a contribution of the AGN > 82%.

Irrespective of the model used, NGC4258 always need a large ($\sim 20-38\%$) contribution from the HII component. It is possible that in this case the spectral decomposition tool DEBLENDIRS is not perfectly separating the different spectral contributions at MIR. We also note that in all cases the stellar component is necessary to take into account the emission in the bluer extreme of the spectrum, while the HII component is necessary to take into account the emission in the redder extreme of the spectrum. A similar result was also found by González-Martín et al. (2019a) for a large sample of AGN.

We obtain upper (or lower) limits of the resulting free parameters and covering factors. The covering factor is defined as $1 - P_{\rm esc}$, where $P_{\rm esc}$ is the probability that a photon emitted in the central engine is able to escape without being absorbed by the torus (in the case of the smooth torus and two-phase medium torus models) or by a dusty cloud in clumpy torus models (Nenkova08, Hoenig10). In the case of Hoenig17 models, the covering factor derived is the sum of the geometrical covering factors of the disk and the covering factor of the outflow (see González-Martín et al. 2019b).

To obtain well constrained parameters a detailed modeling that includes NIR and far-infrared (FIR) data is necessary (see e.g., Ramos Almeida et al. 2014). However, modeling simultaneously the NIR and MIR components of the IRS/Spitzer spectrum of type 1 AGNs has been a challenge (e.g., Mor et al. 2009; Martínez-Paredes et al. 2017; Hernán-Caballero et al. 2015). In this work we assume the NIR as a stellar component. Our purpose is to show if any of the proposed models are able to explain both the peak and shape of the strongest silicate emission features. For this purpose, we only need to check that the range of values of the covering factors obtained from modeling the IRS/Spitzer spectra with each model are within the range of values expected for type 1 AGNs (see e.g., Ramos Almeida et al. 2011;




Figure 9. Modeling and residuals of the IRS/Spitzer spectrum of PG 0844+349. From top to bottom we assume models Fritz06 ($\chi^2_{\rm red} \sim 1.20$), Nenkova08 ($\chi^2_{\rm red} \sim 0.32$), Hoenig10 ($\chi^2_{\rm red} \sim 0.20$), Hoenig17 ($\chi^2_{\rm red} \sim 0.13$), and Stalevski16 ($\chi^2_{\rm red} \sim 0.81$). The $\chi^2_{\rm red}$ values refer to fits with both torus models and other components. The last panel shows the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in erg s⁻¹cm⁻², and the red solid line is the fitted torus model. The orange line is the total SED that results when the stellar (green dot-dashed line) and/or the HII (blue dotted line) components are added to model the spectrum.




Figure 10. Fractional contributions of the AGN (brown histogram), stellar (green histogram) and HII (blue histogram) components for Fritz06, Nenkova08, Hoenig10, Hoenig17, and Stalevski16 models.

Alonso-Herrero et al. 2011; Ichikawa et al. 2015; Mateos et al. 2016; Martínez-Paredes et al. 2017; González-Martín et al. 2017).

We note that on average the smooth models produce a dusty torus with a small angular widths, with low and high viewing angles, although the angular width is poorly constrained. The dusty clumpy torus models of Nenkova08 produce both large and low viewing angles and a range of angular width from low (15 degrees) to high (70 degrees) values, and a number of clouds along the equatorial ray that are in general $\lesssim 7$ clouds, resulting in escape probabilities $\gtrsim 40\%$. The clumpy models of Hoenig10 produce values of the viewing angle that range from nearly 30 to 80 degrees, angular widths around 55 degrees, and number of clouds in the range 2.5 to 10.0. The viewing angles in disk+outflow models of Hoenig17 that we find is between 0 and 50 degrees and the angular widths between 30 and 45 degrees. Although in three cases we obtain lower limits of the angular width, indicating that this could be larger. The two-phase models of Stalev16 produce on average large viewing angles (~ 80), only in one case we obtain a viewing angle around 10 degrees, and the angular width is around 80 degrees, resulting in very obscured AGNs (see Tables 6,7,8, 9, and Figure 17 in Appendix C).

In general, Fritz06 and Hoenig17 models produce lower covering factors than Nenkova08, Hoenig10 and Stalevski16 models. Fritz06 and Hoenig17 models produce covering factors around 0.2, although they range from 0 to 1. The Nenkova08 model produces covering factors around 0.6 with a range from 0.3 to 1.0. These values are consistent with the range of values obtained by Martínez-Paredes et al. (2017) for a sample of PG QSOs using Nenkova08 models. Feltre et al. (2012) compare smooth and clumpy torus models of Nenkova08 and found that both torus model families produce similar MIR continuum shapes for different model parameters. In the Hoenig10, and Stalevski16 models the covering factors are large (around 0.8), probably due the fact that in these models the angular width of the torus tends to be larger, which leads to a more obscured AGN.

Table 3. Best fitting from smooth models. Column 1 lists the name of the object, column 2 the χ^2_{red} , column 3 the combination of components that best fit the IRS/Spitzer spectra, and columns 4, 5, 6 the percentage contribution of each component.

| | | | Comp | onent cont | tributions | | | | Comp | onent cont | ributions | |
|------------------|----------------|----------------------------|----------|------------|------------|-------------|-------------------------------------------------|-----------------------------------|----------|------------|-----------|--|
| Name | χ^2_{red} | Best combination | AGN | Stellar | HII | Name | χ^2_{red} | Best combination | AGN | Stellar | HII | |
| | | | (%) | (%) | (%) | | | | (%) | (%) | (%) | |
| S | mooth | torus models of Fritz | et al. (| 2006) | | D | Disk+outflow models of Hönig & Kishimoto (2017) | | | | | |
| PG2304+042 | 1.92 | Torus+Stellar | 93 | 7 | 0 | PG2304+042 | 0.57 | Disk+outflow+Stellar | 93 | 7 | 0 | |
| PG 0844 + 349 | 1.20 | Torus+Stellar | 93 | 7 | 0 | PKS0518-45 | 0.31 | Disk+outflow | 100 | 0 | 0 | |
| PG 1351+640 | 1.52 | Torus+Stellar | 97 | 3 | 0 | PG 0844+349 | 0.13 | $_{\rm Disk+outflow+Stellar+HII}$ | 87 | 5 | 8 | |
| PG 2214+139 | 1.52 | Torus+Stellar | 96 | 4 | 0 | PG 2214+139 | 0.20 | $_{\rm Disk+outflow+Stellar+HII}$ | 95 | 1 | 4 | |
| PG 0804+761 | 0.63 | Torus+Stellar | 94 | 6 | 0 | PG 0804+761 | 0.49 | Disk+outflow+HII | 95 | 0 | 5 | |
| NGC 3998 | 1.91 | Torus+Stellar | 100 | 0 | 0 | OQ 208 | 1.78 | Disk+outflow+Stellar | 98 | 2 | 0 | |
| | | | | | | NGC 4258 | 1.88 | Disk+outflow+HII | 62 | 0 | 38 | |
| | | | | | | NGC 3998 | 0.39 | Disk+outflow | 100 | 0 | 0 | |
| Clur | npy to | rus models of Nenkova | et al. (| (2008a,b) | | Two | -phase | media torus models of Stalev | ski et a | l. (2016) | | |
| NGC7213 | 1.11 | Torus+Stellar | 96 | 4 | 0 | PG2304+042 | 1.99 | Torus+Stellar | 93 | 7 | 0 | |
| PG2304+042 | 1.25 | Torus+Stellar | 92 | 8 | 0 | PKS0518-45 | 1.65 | Torus | 100 | 0 | 0 | |
| PKS0518-45 | 1.29 | Torus+Stellar | 97 | 3 | 0 | PG 0844+349 | 0.81 | Torus+Stellar | 92 | 8 | 0 | |
| PG 0844 + 349 | 0.32 | Torus+Stellar | 91 | 9 | 0 | PG 2214+139 | 0.91 | Torus+Stellar+HII | 82 | 10 | 8 | |
| $PG\ 1351+640$ | 1.92 | Torus+Stellar | 99 | 1 | 0 | PG 0804+761 | 1.11 | Torus+Stellar | 90 | 10 | 0 | |
| PG 2214+139 | 1.92 | Torus+Stellar | 87 | 13 | 0 | | | | | | | |
| PG 0804 + 761 | 1.60 | Torus+Stellar | 89 | 11 | 0 | | | | | | | |
| OQ 208 | 0.32 | $_{\rm Torus+Stellar+HII}$ | 91 | 3 | 6 | | | | | | | |
| NGC 4258 | 0.78 | $_{\rm Torus+Stellar+HII}$ | 66 | 4 | 30 | | | | | | | |
| NGC 3998 | 0.32 | Torus+Stellar | 81 | 19 | 0 | | | | | | | |
| Clump | y toru | s models of Hönig & I | Kishimo | to (2010b) |) | • | | | | | | |
| PG2304+042 | 1.18 | Torus | 100 | 0 | 0 | | | | | | | |
| PKS0518-45 | 0.46 | $_{\rm Torus+Stellar+HII}$ | 89 | 6 | 5 | | | | | | | |
| PG 0844 + 349 | 0.20 | $_{\rm Torus+Stellar+HII}$ | 80 | 10 | 10 | | | | | | | |
| $PG\ 2214{+}139$ | 0.71 | $_{\rm Torus+Stellar+HII}$ | 76 | 9 | 14 | | | | | | | |
| PG 0804 + 761 | 1.79 | $_{\rm Torus+Stellar+HII}$ | 76 | 9 | 15 | | | | | | | |
| NGC 4258 | 1.64 | $_{\rm Torus+Stellar+HII}$ | 67 | 3 | 29 | | | | | | | |
| NGC 3998 | 0.40 | Torus | 100 | 0 | 0 | | | | | | | |

4.4. Residuals

Using the spectral residuals for all modeled and non-modeled cases we calculate the average spectral residuals for each torus model. The vertical black solid lines in Figure 11 represents the mean wavelength where the silicate features peak in the IRS/Spitzer spectra, and the black dotted lines their 1σ intervals.

In order to discuss qualitatively the similarities and differences between the residuals of the models we divide the spectral range into three parts in Figure 11. These parts are the region bluewards of $10\mu\text{m}$, between $10\mu\text{m}$ and $18\mu\text{m}$, and redwards $18\mu\text{m}$. In part (a) we observe that on average, around $5\mu\text{m}$, Fritz06, Nenkova08, Hoenig10, Stalev16 models are the worst at reproducing the bluer extreme of the spectra within the uncertainties. However, at longer wavelengths the Fritz06 and Stalevski16 models are the least accurate at reproducing the shape of the spectra. The Hoenig17 models best reproduce the spectra at all wavelengths within the uncertainties. In the part (b) Nenkova08 show the flattest residual, while in the redder extreme the Fritz06 and Stalevski16 models show the largest residuals, while the Hoenig10 and Hoenig17 models shows flatter residuals. In part (c) all models show similar residuals, although Nenkova08 show the flattest residual within the uncertainties.

We also observe that the Fritz06 and Stalevski16 models underestimate the strength of the 10 and $18\mu m$ silicate features, while the Nenkova08, Hoenig10, and Hoenig17 models best reproduce the peak and the shape of both features. In general we note that Hoenig17 and Nenkova08 models show flatter residuals, resulting in the models that best reproduces the shape and the peak of the strong silicate features observed in these objects.

Some objects deserve particular attention. For instance, in Figure 19 we can see that for NGC 7213 we require the stellar component to fit the bluer extreme of the spectrum, but that both silicate peaks are still underestimated by the Nenkova08 models.

PG 1351+640 is modeled only by the smooth and clumpy models of Nenkova08. In both cases it is necessary to add the stellar component, which contributes 3% and 1% in the case of smooth (Fritz06) and clumpy models (Nenkova08 and Hoenig10), respectively. OQ208 is modeled only by the Nenkova08 and Hoenig17 models. In this case the residuals from the Nenkova08 model are flatter than the residuals from the Hoenig17 models, although in the case of the Nenkova08 model it is necessary to include the HII component with a contribution of 6%, in addition to




Figure 11. Average residuals (data/model) for all modeling combination including the modeled and non-modeled cases. The red points represent the average residuals of objects fitted with the smooth models. The purple and blue points represent the average residuals obtained from those objects fitted with the clumpy models of Nenkova08 and Hoenig10, respectively. The green points represent the average residuals obtained from the objects fitted using the disk+outflow (Hoenig17) models, and the magenta points are the residuals obtained from using the two-phase dusty torus model of Stalevski16. The vertical grey solid line indicates the mean wavelength where the 10 and 18μ m silicate features peak. The grey dashed lines are the 1σ confidence intervals.

the stellar component (3%) in order to fit the redder extreme of the spectrum. In contrast, the Hoenig17 model is able to reproduce the entire spectral range requiring only a small contribution from the stellar component (2%). The remaining objects are modeled successfully by the Nenkova08, Hoenig10, and Hoenig17 models.

We divide the objects in the Si-s sample into three groups according to their bolometric luminosities and BH accretion rates. For each group and torus model we combine the residual obtained from fitting the AGN-dominated Spitzer spectrum with the components C1 (AGN), C2 (AGN+Stellar), C3 (AGN+HII), and C4 (AGN+Stellar+HII). In Figure 12 we plot the average residual for each group. We note that for the first group, all models are unable to produce a flat residual around the $10\mu m$ silicate feature. But, for the second group, which covers a larger range of bolometric luminosities and BH accretion rates, all the residuals become flatter. At the largest bolometric luminosites of the third group all models show the flattest residuals. These results show that all models fail in reproduce the central wavelength of the $10\mu m$ silicate feature in the objects with lower bolometric luminosities, as we see in Figure 5 in Section 4.1. Additionally, we note that the Hoenig17 models always produce the flattest residual around $5\mu m$ for low, intermediate, and high luminosities.

5. DISCUSSION

In the previous section we used four different torus models (smooth, clumpy, two-phase medium, and disk+outflow) to reproduce the strongest silicate features observed in type 1 AGNs. These features appear broader and shifted with respect to the silicate emission features observed in the standard ISM, suggesting a different dust composition (or geometry) of the torus or surrounding dust of AGNs. We discuss below the dust distribution (Section 5.1) and composition (Section 5.2) of the dust. We also discuss the deficiencies of the models to reproduce the strong silicate emission features in Section 5.3.




Figure 12. Average residuals for three ranges of bolometric luminosities and BH accretion rates. Colours are as in Figure 11.

5.1. Dust distribution

Based on our analysis the clumpy models of Nenkova08 ($\chi^2_{\rm red} \sim 1.03$), Hoenig10 ($\chi^2_{\rm red} \sim 0.89$), and Hoenig17 ($\chi^2_{\rm red} \sim 0.70$) produce a better fitting of the IRS/Spitzer spectra than the smooth ($\chi^2_{\rm red} \sim 1.36$) and two-phase medium ($\chi^2_{\rm red} \sim 1.20$) models. This result is in agreement with the previous evidence that the surrounding medium around the AGN should be clumpy (see Ramos Almeida, & Ricci 2017, and references therein). Indeed, Mendoza-Castrejón et al. (2015) found that isolated type 1 AGN have a clumpy dust distribution, while interacting type 1 AGN can have both clumpy or smooth dusty distributions. Additionally, we find that all models produces flatter residual in high luminosity AGN than in low luminosity AGN, probably due to all models better sample the central wavelength and both 10 and 18 μ m silicate features of high luminosity AGNs. However, a high luminosities Hoenig17 models produce the flattest residual along all the spectral range.

Mor et al. (2009) used IRS/Spitzer spectra of a sample of 26 nearby QSOs to constrain the clumpy models of Nenkova08. They argue that in order to model the entire spectral range between 5-30 μ m they need to add two more components, one that takes into account the emission produced by the dust in the narrow line region (NLR) (Schweitzer et al. 2008), and another one that takes into account the emission produced by dust close to the AGN, not directly related with the dusty torus (e.g., Minezaki, et al. 2004; Riffel et al. 2009; Kishimoto et al. 2007). However, Alonso-Herrero et al. (2011) showed, for a sample of local Seyferts, that when high angular resolution data at NIR and MIR is used, it is not necessary to add any additional components in order to constrain the clumpy models of Nenkova08, although for some type 1 Seyferts galaxies they found that the NIR emission is underpredicted by the fitted SED. In a previous work, we used the starburst-subtracted IRS/Spitzer spectra from $\sim 7.5 - 15\mu$ m plus NIR high angular resolution data from the Near Infrared Camera and Multi-Object Spectrometer (NICMOS) on HST for a sample of

20 nearby QSOs in order to constrain the clumpy torus models of Nenkova08. Martínez-Paredes et al. (2017) found that including the spectral range between $5-8\mu m$ resulted in a poor fitting of the $10\mu m$ silicate emission feature. In this work we find that the AGN-dominated *Spitzer* spectra of the Si-s type 1 AGN can be fitted by Nenkova08 and/or Hoenig10 models by adding a stellar component that takes into account the bluer spectral range, and in some cases the HII component, in order to improve the fitting in the redder spectral range. A similar result was also found for a large sample of AGNs (González-Martín et al. 2019a,b).

Hernán-Caballero et al. (2017) found, for a sample of 85 QSOs, that a superposition of two blackbodies between 1.7-8.4 μ m, with temperatures around 1000 K for the hot blackbody component, and 400 K for the warm blackbody component, can fit the *Spitzer* spectra between 0.1 and 10 μ m. They argue that an additional hotter component of dust is necessary to reproduce the excess emission at 1-2 μ m. On the other hand, Lyu et al. (2017) argues that the strong silicate emission features observed in dust-deficient PG QSOs can be explained assuming a reduced height scale of the warm dust, i.e., the dust between the sublimation zone and the outer region of cold dust, allowing for less interception of the radiation from the accretion disk with the inner dust, resulting in a decrease of the MIR continuum, but keeping the heating of the outer dust responsible for the silicate emission features invariant (see Figure 20 in Lyu et al. 2017).

5.2. Dust composition

In general, the Nenkova08 models produce a better fit than the Hoenig10 ones. They assume the same geometry, but the former uses a standard dust composition, while the latter includes standard ISM dust plus standard ISM with large grains, and also a composition of dust mostly dominated by graphite. The Hoenig10 models assume this dust composition in order to take into account the observational suggestion that the dust composition in AGN deviates from standard ISM dust (Suganuma et al. 2006; Kishimoto et al. 2007). One of the clear improvements in the Hoenig17 models is that they allow the existence of different dust compositions in different parts of the surrounding dusty structure (disk+outflow).

The chemical composition of the dust has been largely studied for the ISM (see Henning 2010, and references therein), and to a lesser extent in AGN (see Lyu, Hao & Li 2014, and references therein). However, determining the exact chemical composition of the dust has been challenging. Srinivasan et al. (2017) studied the dust composition of a large sample of PG QSOs with a redshift z < 0.5, that showed the 10μ m silicate feature in emission. They found that the dust is mostly composed of amorphous oxides and silicates, plus a small fraction in crystalline form. However, this small fraction is nearly four times larger than the last upper limit ($\lesssim 2.2$) reported by Kemper et al. (2005) for the ISM, and more similar to the upper limit ($\lesssim 5$) previously reported by Li & Draine (2001) for the ISM.

5.3. Deficiencies of the models

In general we find that neither the peak nor the shape of the silicate features of the AGN-dominated IRS/Spitzer spectra of the Si-s type 1 AGN are perfectly reproduced by the models. However, we note that for each object either the Hoenig17 or the Nenkova08 models produce the smallest $\chi^2_{\rm red}$ and flattest residuals. Hönig & Kishimoto (2017) point out that their dust composition explains both the observed small NIR reverberation mapping and interferometric sizes with dust sublimation physics. They argue that this combination of disk+outflow clumpy models is able to reproduce the $3-5\mu$ m bump in type 1 AGN, and preserve the MIR bump produced by the wind. Indeed, according to this model a standard ISM composition of the dust in the wind would be the responsible for the emission of the silicate features. García-González et al. (2017) found that the Hoenig17 (disk+outflow) models predict MIR slopes (between $8.1-12.5\mu$ m) and silicate strengths at 10 μ m that are in agreement with the values observed in type 1 AGN. Particularly, they noted that when clouds are more concentrated towards the inner region of the torus (see e.g., Hönig & Kishimoto 2010b; Ramos Almeida et al. 2011; Ichikawa et al. 2015; Martínez-Paredes et al. 2017), the MIR spectral indices are flatter and the silicate features are stronger than those observed in Seyferts and QSOs.

6. SUMMARY AND CONCLUSIONS

In order to investigate which model better reproduces the shape and peak of the strongest 10μ m silicate emission features observed in type 1 AGN, we measure the 10μ m silicate emission strength for a sample of local (z < 0.1) type 1 AGN, for which their IRS/Spitzer spectra is mostly dominated by the emission of non-stellar processes (> 80%).

We find that these objects show silicate features in emission. On average the 10μ m silicate feature has a strength of $0.13^{+0.15}_{-0.36}$ that peaks at 10.3μ m, and a 18μ m silicate strength of $0.14^{+0.06}_{-0.06}$ that peaks at 17.3μ m. We find that 10 objects are among the AGN with the largest 10μ m silicate strengths ($\sigma_{Si_{10\mu}} > 0.28$, Si-s sample), and that some of them have been previously classified as objects with prominent silicate features.

We use four different torus models, Fritz06 (Fritz et al. 2006), Nenkova08 (Nenkova et al. 2008a,b), Hoenig10 (Hönig & Kishimoto 2010b), Stalevski16 (Stalevski et al. 2016) and a disk+outflow (Hoenig17, Hönig & Kishimoto 2017) model to fit the IRS/Spitzer spectra of the Si-s sample, and investigate which model better reproduces the peak and shape of both silicate emission features. The models assume different dust distributions, namely, smooth, clumpy, and a two-phase medium, as well as different dust compositions. We find that in most cases it is necessary to add a stellar or HII component in order to improve the fit. In most cases we find that the contribution of these components is << 20%, in agreement with our selection requirement that the spectra be dominated by the emission of the AGN. The exceptions is NGC 4258 for which the spectral decomposition seems to underestimate the HII component.

We find that in general Fritz06 and Hoenig17 models produce lower covering factors than the Nenkova08, Hoenig10, and Stalevski17 models. The values are consistent with those reported in previous works for type 1 AGN. We find that the individual and average spectra are reproduced better with clumpy torus models than smooth models. Moreover, the Hoenig17 model shows the flattest residuals along all the spectral range between $\sim 5-35\mu m$, while the rest of models fail to reproduce the bluer extreme of the spectrum. However, on average none of the models are able to exactly reproduce the peak and shape of the silicate features.

In the near future the *Mid-Infrared Instrument* (MIRI) onboard of the *James Webb Space Telescope* will provide high angular resolution with higher sensitivity and spectral resolution observations which will allow an in-depth investigation of the dust properties in active galaxies. Additionally, new models that include a better description of the properties of the dust will be required.

ACKNOWLEDGEMENTS

MM-P acknowledges support by the KASI and UNAM-DGAPA postdoctoral fellowships. This work is partially supported by the KASI project 2019184100 and Conacyt project CB-2016-281948. O.G.-M. acknowledges support by the PAPIIT projects IA100516. A.A.-H. acknowledges support through grant PGC2018-094671-B-I00 (MCIU/AEI/FEDER,UE). AAHs work was done under project No. MDM-2017-0737 Unidad de Excelencia "María de Maeztu"- Centro de Astrobiología (INTA-CSIC). YK acknowledges support from grant DGAPA-PAPIIT 106518, and from program DGAPA-PASPA. T.H acknowledges the support from the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (2019R1A2C1087045). CRA acknowledges financial support from the Spanish Ministry of Science and Innovation (MICINN) through project PN AYA2013-47742- C4-2-P. CRA also acknowledges the Ramon y Cajal Program of the Spanish Ministry of Economy and Competitiveness. This work is based on observations obtained with the Spitzer Space Observatory, which is operated by JPL, Caltech, under NASA contract 1407. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by JPL, Caltech, under contract with the National Aeronautics and Space Administration. CASSIS is a product of the Infrared Science Center at Cornell University, supported by NASA and JPL.

REFERENCES

Antonucci, R. 1993, ARA&A, 31, 473

Alexander, D. M., & Hickox, R. C. 2012, New Astronomy Reviews, 56, 93

Alonso-Herrero, A., Ramos Almeida, C., Mason, R., et al. 2011, ApJ, 736, 82

Alonso-Herrero, A., Poulton, R., Roche, P. F., et al. 2016, MNRAS, 463, 2405

Bianchi, S., Bonilla, N. F., Guainazzi, M., Matt, G., & Ponti, G. 2009, A&A, 501, 915

Brightman, M., & Nandra, K. 2011, MNRAS, 413, 1206

Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000

Buchanan, C. L., Gallimore, J. F., O'Dea, C. P., et al. 2006, AJ, 132, 401

Cappi, M., Panessa, F., Bassani, L., et al. 2006, A&A, 446, 459 Cusumano, G., La Parola, V., Segreto, A., et al. 2010, Astronomy and Astrophysics, 510, A48

Draine, B. T. 2003, ApJ, 598, 1026

Draine, B. T., Dale, D. A., Bendo, G., et al. 2007, ApJ, 663, 866

Dullemond, C. P., & van Bemmel, I. M. 2005, A&A, 436, 47

Dudik, R. P., Weingartner, J. C., Satyapal, S., et al. 2007, ApJ, 664, 71

Efstathiou, A., & Rowan-Robinson, M. 1995, MNRAS, 273, 649

Elvis, M., Wilkes, B. J., McDowell, J. C., et al. 1994, ApJS, 95, 1

Feltre, A., Hatziminaoglou, E., Fritz, J., & Franceschini, A. 2012, MNRAS, 426, 120

- Fritz, J., Franceschini, A., & Hatziminaoglou, E. 2006, MNRAS, 366, 767
- García-González, J., Alonso-Herrero, A., Hönig, S. F., et al. 2017, MNRAS, 470, 2578
- Granato, G. L., Danese, L., & Franceschini, A. 1997, Mem. Soc. Astron. Italiana, 68, 39
- González-Martín, O., Rodríguez-Espinosa, J. M., Díaz-Santos, T., et al. 2013, A&A, 553, A35
- González-Martín, O., Masegosa, J., Hernán-Caballero, A., et al. 2017, ApJ, 841, 37
- González-Martín, O., Masegosa, J., García-Bernete, I., et al. 2019, arXiv e-prints, arXiv:1908.11381
- González-Martín, O., Masegosa, J., García-Bernete, I., et al. 2019, arXiv e-prints, arXiv:1908.11389
- Hatziminaoglou, E., Hernán-Caballero, A., Feltre, A., et al. 2015, ApJ, 803, 110
- Hao, L., Spoon, H. W. W., Sloan, G. C., et al. 2005, ApJL, 625, L75
- Hao, L., Weedman, D. W., Spoon, H. W. W., et al. 2007, ApJL, 655, L77
- Hernán-Caballero, A., Alonso-Herrero, A., Hatziminaoglou, E., et al. 2015, ApJ, 803, 109
- Henning, T. 2010, ARA&A, 48, 21
- Hernán-Caballero, A., Hatziminaoglou, E., Alonso-Herrero, A., et al. 2017, Frontiers in Astronomy and Space Sciences, 4, 30
- Hönig, S. F., Kishimoto, M., Gandhi, P., et al. 2010, A&A, 515, A23
- Hönig, S. F., & Kishimoto, M. 2010, A&A, 523, A27
- Hönig, S. F., Kishimoto, M., Antonucci, R., et al. 2012, The Astrophysical Journal, 755, 149
- Hönig, S. F., Kishimoto, M., Tristram, K. R. W., et al. 2013, The Astrophysical Journal, 771, 87
- Hönig, S. F., & Kishimoto, M. 2017, ApJL, 838, L20
- Houck, J. R., Roellig, T. L., Van Cleve, J., et al. 2004, Proc. SPIE, 5487, 62
- Ichikawa, K., Packham, C., Ramos Almeida, C., et al. 2015, ApJ, 803, 57
- Kemper, F., Vriend, W. J., & Tielens, A. G. G. M. 2005, ApJ, 633, 534
- Kishimoto, M., Hönig, S. F., Beckert, T., et al. 2007, A&A, 476, 713
- Krolik, J. H., & Begelman, M. C. 1988, ApJ, 329, 702
- Laor A., Draine B. T., 1993, ApJ, 402, 441
- Li A., Draine B. T., 2001, AAS, 199, 97.13
- Li, M. P., Shi, Q. J., & Li, A. 2008, MNRAS, 391, L49
- Lebouteiller, V., Barry, D. J., Spoon, H. W. W., et al., 2011, ApJS, 196, 8
- Lyu J., Hao L., Li A., 2014, ApJL, 792, L9
- Lyu, J., Rieke, G. H., & Shi, Y. 2017, ApJ, 835, 257

- López-Gonzaga, N., & Jaffe, W. 2016, Astronomy and Astrophysics, 591, A128
- Marconi, A., Risaliti, G., Gilli, R., et al. 2004, Monthly Notices of the Royal Astronomical Society, 351, 169
- Martínez-Paredes, M., Alonso-Herrero, A., Aretxaga, I., et al. 2015, MNRAS, 454, 3577
- Martínez-Paredes, M., Aretxaga, I., Alonso-Herrero, A., et al. 2017, MNRAS, 468, 2
- Mason, R. E., Lopez-Rodriguez, E., Packham, C., et al. 2012, AJ, 144, 11
- Mason, R. E., Ramos Almeida, C., Levenson, N. A., Nemmen, R., & Alonso-Herrero, A. 2013, ApJ, 777, 164
- Mateos, S., Carrera, F. J., Alonso-Herrero, A., et al. 2016, The Astrophysical Journal, 819, 166
- Mathis, J. S., Rumpl, W., & Nordsieck, K. H. 1977, ApJ, 217, 425
- Mendoza-Castrejón, S., Dultzin, D., Krongold, Y., et al. 2015, MNRAS, 447, 2437
- Minezaki T., et al., 2004, ApJL, 600, L35
- Mor, R., Netzer, H., & Elitzur, M. 2009, ApJ, 705, 298
- Nenkova, M., Ivezić, Ž., & Elitzur, M. 2002, ApJL, 570, L9
- Nenkova, M., Sirocky, M. M., Ivezić, Ž., & Elitzur, M. 2008, ApJ, 685, 147
- Nenkova, M., Sirocky, M. M., Nikutta, R., Ivezić, Ž., & Elitzur, M. 2008, ApJ, 685, 160
- Neugebauer, G., Oke, J. B., Becklin, E. E., & Matthews, K. 1979, ApJ, 230, 79
- Nikutta, R., Elitzur, M., & Lacy, M. 2009, ApJ, 707, 1550
 Ossenkopf, V., Henning, T., & Mathis, J. S. 1992, A&A,
 261, 567
- Pei, Y. C. 1992, ApJ, 395, 130
- Peterson, B. M. 1997, IAU Colloq. 159: Emission Lines in Active Galaxies: New Methods and Techniques, 113, 489
- Pier, E. A., & Krolik, J. H. 1992, ApJ, 401, 99
- Raban, D., Jaffe, W., Röttgering, H., et al. 2009, Monthly Notices of the Royal Astronomical Society, 394, 1325
- Ramos Almeida, C., Levenson, N. A., Alonso-Herrero, A., et al. 2011, ApJ, 731, 92
- Robson, I., Dunlop, J., Taylor, G., & Hughes, D. 1995, BAAS, 27, 1415
- Ramos Almeida, C., Alonso-Herrero, A., Esquej, P., et al. 2014, MNRAS, 445, 1130
- Ramos Almeida, C., & Ricci, C. 2017, Nature Astronomy, 1, 679
- Riffel, R. A., Storchi-Bergmann, T., & McGregor, P. J. 2009, ApJ, 698, 1767
- Rowan-Robinson, M. 1977, ApJ, 213, 635
- Sambruna, R. M., Tombesi, F., Reeves, J. N., et al. 2011, ApJ, 734, 105

- Sanders, D. B., Phinney, E. S., Neugebauer, G., Soifer,
 B. T., & Matthews, K. 1989, ApJ, 347, 29
- Shi, Y., Rieke, G. H., Hines, D. C., et al. 2006, ApJ, 653, 127
- Siebenmorgen, R., Haas, M., Krügel, E., & Schulz, B. 2005, A&A, 436, L5
- Sirocky, M. M., Levenson, N. A., Elitzur, M., Spoon, H. W. W., & Armus, L. 2008, ApJ, 678, 729
- Srinivasan, S., Kemper, F., Zhou, Y., et al. 2017, Planet. Space Sci., 149, 56
- Sturm, E., Schweitzer, M., Lutz, D., et al. 2005, ApJL, 629, L21
- Sturm, E., Hasinger, G., Lehmann, I., et al. 2006, ApJ, 642, 81
- Schinnerer, E., Eckart, A., & Tacconi, L. J. 1998, ApJ, 500, 147
- Schweitzer, M., Groves, B., Netzer, H., et al. 2008, ApJ, 679, 101-117
- Smith, J. D. T., Draine, B. T., Dale, D. A., et al. 2007, ApJ, 656, 770
- Stalevski, M., Fritz, J., Baes, M., et al. 2012, MNRAS, 420, 2756
- Stalevski, M., Ricci, C., Ueda, Y., et al. 2016, MNRAS, 458, 2288

- Suganuma, M., Yoshii, Y., Kobayashi, Y., et al. 2006, ApJ, 639, 46
- Thompson, G. D., Levenson, N. A., Uddin, S. A., et al. 2009, ApJ, 697, 182
- Tristram, K. R. W., Burtscher, L., Jaffe, W., et al. 2014, Astronomy and Astrophysics, 563, A82
- Tueller, J., Baumgartner, W. H., Markwardt, C. B., et al. 2010, VizieR Online Data Catalog, J/ApJS/186/378
- Ueda, Y., Ishisaki, Makishima, K., & Ohashi, T. 2005, ApJS, 161, 185
- Véron-Cetty, M.-P., & Véron, P. 2010, A&A, 518, A10
- Wada, K., Papadopoulos, P. P., & Spaans, M. 2009, ApJ, 702, 63
- Wada, K. 2012, ApJ, 758, 66
- Werner, M. W., Roellig, T. L., Low, F. J., et al. 2004, ApJS, 154, 1
- Winter, L. M., Veilleux, S., McKernan, B., & Kallman, T. R. 2012, ApJ, 745, 107
- Younes, G., Porquet, D., Sabra, B., & Reeves, J. N. 2011, A&A, 530, A149
- Zhou, X.-L., & Zhang, S.-N. 2010, ApJL, 713, L11




Figure 13. 10μ m silicate strength as measured using our own methodology (see Section 2.3) and deblendIRS. The grey dashed line represent the 1:1 comparison.

7. APPENDIX A: SPECTRAL DECOMPOSITION

In order to select those type 1 AGNs in which the IRS/Spitzer spectrum is mostly (> 80%) dominated by emission from dust heated by the AGN we use the spectral decomposition tool DEBLENDIRS from Hernán-Caballero et al. (2015).

DEBLENDIRS is a spectral decomposition tool that uses a set of starburst, stellar and AGN templates with IRS/Spitzer spectrum. This spectral decomposition assumes that the spectral shape of the AGN and its host galaxy are found in others sources where the emission from the AGN or host galaxy completely dominate the spectral emission. The spectral emission of the host galaxy is composed by the stellar emission (passive stellar population) and the emission from the interstellar medium (ISM), which is call PAH, from Polycyclic Aromatic Hydrocarbon, since the emission of these molecules is relate with the presence of young star forming regions (starburst, SB). The spectral decomposition is carry on trying every possible combination between stellar, PAH, and AGN, according to the following linear spectral combination:

$$f_{i,j,k}(\lambda) = af_i^{stellar}(\lambda) + bf_i^{PAH} + cf_k^{AGN}(\lambda),$$
(2)

where the i,j and k indices range all the stellar, PAH, and AGN templates, respectively. The a, b and c coefficients are obtain through the χ^2 minimization. In Figure 14 we show an example of the spectral decomposition. After find the combination of components that best reproduce the IRS/Spitzer spectrum DEBLENDIRS estimates the fractional contribution of each component to the integrated 5 - 15 μ m luminosity, the luminosity of the starburst at 12μ m, and the luminosity of the AGN at 12, and 6 μ m. Additionally, it gives for the AGN component the silicate strength measured at the wavelength where the silicate feature peaks, and the MIR spectral index measured between 8.1 and 12.5 μ m.




Figure 14. Upper. Spectral decomposition: the green dot-dashed, blue dashed, and red dotted lines represent the stellar, starburst, and AGN components respectively. The orange solid line is the sum of the three (stellar, starburst, and AGN) components. The black solid line is the resampled IRS/Spitzer spectrum with their errors (grey shadow). The horizontal solid black line around zero is the residual. Bottom. Probability distributions: rSTR, rPAH, and rAGN are the fractional contribution of the stellar, PAH, and AGN components. L_{12} SB fraction, L_{12} AGN fraction, and L_{6} AGN fraction are the monochromatic luminosity of the SB and AGN at 12 and 6 μ m respectively. S_{sil} and α AGN are the silicate strength and the spectral index of the AGN component.

8. APPENDIX B: SILICATE STRENGTH MEASUREMENTS

Here we report the silicate feature strengths measurements for the full sample.

Table 4. Silicate feature strengths measured from the IRS/Spitzer spectrum. Columns 1 lists the name. Columns 2 and 3 list the wavelength where the $10\mu m$ silicate emission feature peaks, and the $10\mu m$ silicate strength. Columns 4 and 5 are like columns 2 and 3 but for the $18\mu m$ silicate emission feature. Columns 6, 7, 8, and 9 are like columns 2, 3, 4, and 5 but for the values previously reported in the literature. Column 10 list the reference.

| Name | λ_p (μ m) | $\mathrm{Si}_{10\mu\mathrm{m}}$ | λ_p (μ m) | $\mathrm{Si}_{18\mu\mathrm{m}}$ | λ_p | $\mathrm{Si}_{10\mu\mathrm{m}}$ | λ_p | $\mathrm{Si}_{18\mu\mathrm{m}}$ | Ref |
|----------------------------------|----------------------------------|--------------------------------------|----------------------------------|------------------------------------|---------------|---------------------------------|----------------|---------------------------------|-----|
| NGC7213 | 10.7 ± 0.1 | 0.52 ± 0.05 | 17.3 ± 0.1 | 0.23 ± 0.04 | 10.1 | 0.6 | 17.9 | 0.16 | 1 |
| 3C321 | 9.8 ± 0.1 | -0.9 ± 0.1 | 18.1 ± 0.1 | -0.09 ± 0.05 | | | | | |
| 3C405 | 9.8 ± 0.1 | -0.73 ± 0.07 | 18.2 ± 0.1 | -0.06 ± 0.04 | | | | | |
| 3C33 | 9.7 ± 0.1 | -0.17 ± 0.04 | 17.4 ± 0.1 | 0.12 ± 0.04 | | | | | |
| ESO434-G40 | 9.9 ± 0.1 | -0.37 ± 0.03 | 17.8 ± 0.1 | 0.07 ± 0.02 | | | | | |
| IIIZw2 | 10.4 ± 0.8 | 0.04 ± 0.05 | 16.6 ± 0.1 | 0.11 ± 0.03 | | | | | |
| IRAS03450+0055 | 10.3 ± 0.1 | 0.15 ± 0.05 | 16.7 ± 0.1 | 0.08 ± 0.05 | | | | | |
| IRAS05218-1212 | 9.2 ± 0.1 | -0.06 ± 0.03 | 17.8 ± 0.1 | 0.13 ± 0.04 | | | | | |
| MCG-6-30-15 | 9.0 ± 0.1 | -0.13 ± 0.03 | 17.3 ± 0.1 | 0.07 ± 0.04 | 10.2 | 0.02 | 18.3 | 0.10 | 1 |
| MRK1218 | 9.2 ± 0.1 | -0.22 ± 0.1 | 16.5 ± 0.4 | 0.24 ± 0.07 | | | | | |
| Mrk10 | 9.4 ± 0.1 | -0.14 ± 0.05 | 17.4 ± 0.1 | 0.1 ± 0.06 | | | | | |
| Mrk110 | 10.3 ± 0.1 | 0.19 ± 0.03 | 18.0 ± 0.1 | 0.25 ± 0.03 | | | | | |
| Mrk176 | 11.6 ± 0.2 | 0.07 ± 0.03 | 17.3 ± 0.1 | 0.18 ± 0.04 | | | | | |
| Mrk231 | 9.8 ± 0.1 | -0.73 ± 0.02 | 17.3 ± 0.1 | -0.18 ± 0.02 | 9.8 | -0.62 | 17.9 | -0.23 | 1 |
| Mrk3 | 9.8 ± 0.1 | -0.45 ± 0.04 | 17.9 ± 0.1 | 0.19 ± 0.02 | | | | | |
| Mrk348 | 9.4 ± 0.1 | -0.29 ± 0.04 | 16.6 ± 0.1 | 0.18 ± 0.02 | | | | | |
| Mrk463E | 9.8 ± 0.1 | -0.45 ± 0.02 | 18.0 ± 0.1 | 0.05 ± 0.03 | 9.8 | -0.4 | 18.3 | 0.09 | 2 |
| Mrk50 | 10.8 ± 0.1 | 0.22 ± 0.08 | 17.9 ± 0.1 | 0.27 ± 0.26 | 10.8 | 0.24 | 18.3 | 0.37 | 3 |
| Mrk573 | 9.3 ± 0.1 | -0.14 ± 0.03 | 17.3 ± 0.1 17.3 ± 0.1 | 0.27 ± 0.20 0.11 ± 0.02 | 9.4 | -0.10 | 17.2 | 0.04 | 3 |
| Mrk734 | 10.7 ± 0.1 | 0.06 ± 0.04 | 18.0 ± 0.1 | 0.09 ± 0.04 | | | | | |
| Mrk915 | 9.4 ± 0.1 | -0.11 ± 0.07 | 17.3 ± 0.1 | 0.03 ± 0.04 0.18 ± 0.05 | 10.4 | 0.05 | 17.1 | 0.27 | 3 |
| NGC3081 | 9.4 ± 0.1 9.3 ± 0.1 | -0.11 ± 0.07 -0.26 ± 0.03 | 17.3 ± 0.1 17.3 ± 0.1 | 0.10 ± 0.05 0.12 ± 0.06 | | | | | |
| NGC7212 | 9.9 ± 0.1 | -0.26 ± 0.03 -0.46 ± 0.09 | 17.3 ± 0.1 17.3 ± 0.1 | 0.12 ± 0.00 0.18 ± 0.04 | | | | | |
| NGC7212 NGC788 | 9.9 ± 0.1 9.4 ± 0.1 | -0.40 ± 0.09 -0.25 ± 0.04 | 17.3 ± 0.1 16.7 ± 0.1 | 0.16 ± 0.04 0.14 ± 0.03 | 9.5 | -0.08 | 20.5 | 0.02 | 3 |
| PG1149-110 | 9.4 ± 0.1 11.6 ± 0.1 | -0.25 ± 0.04 0.06 ± 0.05 | 10.7 ± 0.1 17.3 ± 0.1 | | | | | | |
| | | | | 0.13 ± 0.03 | | | | | |
| PG1244+026 | 10.9 ± 0.1 | 0.14 ± 0.06 | 17.3 ± 0.1 | 0.06 ± 0.04 | | | | | |
| PKS2048-57 ΓΟΝ1542 | 9.9 ± 0.1 | -0.33 ± 0.02 | 15.3 ± 0.1 | 0.02 ± 0.02 | | | | | |
| UGC3601 | 10.9 ± 0.1 11.6 ± 0.1 | 0.08 ± 0.06 0.1 ± 0.06 | 17.8 ± 0.1 16.6 ± 0.1 | 0.12 ± 0.03 | | | | ••• | |
| | | | | 0.20 ± 0.08 | 10.7 | 0.17 | 17.2 | 0.16 | 3 |
| JM614 | 10.9 ± 0.1 | 0.12 ± 0.05 | 17.6 ± 0.1 | 0.19 ± 0.1 | 10.7 | 0.17 | | 0.16 | |
| 33-0754+394 | 10.1 ± 0.2 | 0.13 ± 0.09 | 17.7 ± 0.1 | 0.13 ± 0.17 | | | | ••• | |
| F9 | 10.7 ± 0.1 | 0.13 ± 0.02 | 17.4 ± 0.1 | 0.13 ± 0.03 | | | | ••• | |
| IZw136 | 9.2 ± 0.1 | -0.06 ± 0.03 | 17.9 ± 0.1 | 0.09 ± 0.04 | | | | | |
| Zw1 | 9.7 ± 0.1 | 0.22 ± 0.03 | 17.8 ± 0.1 | 0.08 ± 0.03 | | | | | |
| PG0007+106/Mrk1501 | 11.6 ± 0.1 | 0.04 ± 0.05 | 16.3 ± 0.3 | 0.10 ± 0.03 | | | | | |
| PG0923+129/Mrk705 ^a | 9.2 ± 0.1 | -0.09 ± 0.03 | 17.3 ± 0.1 | 0.16 ± 0.03 | | | | | |
| PG1211+143 | 10.7 ± 0.1 | 0.23 ± 0.03 | 17.6 ± 0.1 | 0.15 ± 0.04 | | ••• | | | |
| PG1229+204/Mrk771 | 9.2 ± 0.1 | -0.15 ± 0.08 | 19.1 ± 0.2 | 0.07 ± 0.03 | | ••• | | | |
| PG1411+442 | 10.8 ± 0.1 | 0.17 ± 0.04 | 18.4 ± 0.2 | 0.1 ± 0.03 | | | | | |
| PG1426+015/Mrk1383 | 10.7 ± 0.1 | 0.17 ± 0.04 | 17.4 ± 0.1 | 0.15 ± 0.03 | | | | | |
| G1440+356/Mrk478 | 9.3 ± 0.1 | -0.09 ± 0.05 | 17.3 ± 0.1 | 0.07 ± 0.02 | | | | ••• | |
| PG1448+273 | 11.6 ± 0.1 | 0.08 ± 0.04 | 17.7 ± 0.1 | 0.14 ± 0.03 | | | | ••• | |
| PG1501+106/Mrk841 | 9.4 ± 0.1 | -0.09 ± 0.03 | 18.2 ± 0.1 | 0.14 ± 0.03 | | | | | |
| PG1534+580/Mrk290 | 10.7 ± 0.1 | 0.11 ± 0.03 | 17.9 ± 0.1 | 0.15 ± 0.03 | | | | | |
| ESO198-G24 | 9.9 ± 0.1 | 0.23 ± 0.05 | 16.3 ± 0.1 | 0.25 ± 0.06 | | | | | |
| ESO548-G81 | 10.7 ± 0.1 | 0.21 ± 0.04 | 16.8 ± 0.1 | 0.22 ± 0.04 | 10.7 | 0.31 | 18.0 | 0.17 | 3 |
| NGC2110 | 10.7 ± 0.1 | " 0.22 ± 0.03 | 16.5 ± 0.1 | 0.16 ± 0.02 | | | | | |
| Mrk486/PG1535+547 | 10.9 ± 0.1 | 0.17 ± 0.05 | 17.4 ± 0.1 | 0.10 ± 0.06 | | | | | |
| $C120^{b}$ | 10.7 ± 0.1 | 0.19 ± 0.03 | 17.5 ± 0.1 | 0.13 ± 0.04 | 10.2 | 0.26 | 17.8 | 0.15 | 1 |
| NGC1275 | 10.6 ± 0.3 | 0.15 ± 0.02 | 17.9 ± 0.1 | 0.09 ± 0.03 | | | | | |
| NGC7603 | 11.0 ± 0.1 | 0.21 ± 0.02 | 16.5 ± 0.1 | 0.17 ± 0.03 | 10.2 | 0.12 | 17.7 | 0.13 | 1 |
| $^{\circ}$ $^{\circ}$ $^{\circ}$ | | | | | 11.7 | 0.16 | 17.1 | 0.13 | 3 |
| /IRK1018 | 10.2 ± 0.1 | 0.23 ± 0.06 | 17.7 ± 0.1 | 0.16 ± 0.05 | | | | | |
| C382 | 10.7 ± 0.1 | 0.24 ± 0.05 | 17.5 ± 0.1 | 0.20 ± 0.03 | | | | | |
| PG1404+226 | 10.0 ± 0.1 | 0.24 ± 0.06 | 16.6 ± 0.1 | 0.2 ± 0.08 | | | | | |
| $^{\circ}$ G2209+184 a | 11.0 ± 0.1 | 0.25 ± 0.06 | 16.4 ± 0.1 | 0.21 ± 0.10 | | | | | |
| UGC12282 | Sy1.9 | 0.0169 | 42.8* | 10 | 9.2 ± 0.1 | -0.31 ± 0.06 | 17.3 ± 0.1 | 0.15 ± 0.07 | |

Note.- Ref.: (1): Thompson et al. (2009), (2): Sirocky et al. (2008), (3): Mendoza-Castrejón et al. (2015). ^aThe band 1 range from $8.0-8.5\,\mu\text{m}$. ^bThe band 1 range from $8.0-8.8\,\mu\text{m}$. ^cThe band 1 and 2, range from $8.0-8.8\,\mu\text{m}$, and $14.0-14.5\,\mu\text{m}$, respectively.




Figure 15. IRS/Spitzer spectrum (light blue solid line). Upper panel: NGC3998 (left), NGC4258 (right). Middle panel: NGC7213 (left), OQ 208 (right). Bottom panel: PG0804+761 (left), PG0844+349 (right). The red line is the local continuum that follows the broad features of the IRS/Spitzer spectrum. The blue solid lines are the bootstrapped local continua and the vertical pink dashed bars are the bands used to fit the continua around the features. The vertical black dashed-lines indicate the wavelength where the silicate strength is measured. The vertical grey dashed-lines mark other emission lines. For NGC3998 the spectrum between 14 and 35μ m is the high angular resolution spectrum from Spitzer.




Figure 16. IRS/Spitzer spectrum. Upper panel: PG1351+640 (left), PG2214+139 (right). Bottom panel: PG2304+042 (left), PKS0518-45 (right). Lines as in Figure 15.

9. APPENDIX C: TABLES OF PARAMETERS

Here we report the χ^2 and degree of freedom for each object in the Si-s sample fitted with the model combination: AGN, AGN+Stellar, AGN+HII, and AGN+Stellar+HII. We also report the set of parameters obtained from the best fit, and the plots from modeling the non-stellar IRS/Spitzer spectrum of all objects in the Si-s sample.

Table 5. All fitting from smooth models of Fritz et al. (2006), clumpy models of Nenkova et al. (2008a,b) and Hönig & Kishimoto (2010b), and disk+outflow models of Hönig & Kishimoto (2017). Column 1 lists the name and models. Column 2, 3, 4 and 5 the χ^2 and dof of each model combination: C1 indicate AGN model (torus or disk+outflow), C2 AGN+Stellar, C3 AGN+HII, and C4 AGN+Stellar+HII. Column 6 lists the combination of components that best fit the IRS/Spitzer spectrum.

| | | χ^2 | dof dof | | | χ^2/dof | | | | | |
|--------------|-----------|-----------|-----------|-----------|------|-----------------------|-----------|-----------|-----------|-----------|------|
| Name | C1 | C2 | C3 | C4 | Best | Name | C1 | C2 | C3 | C4 | Best |
| NGC7213 | | | | | | PG2214+139 | | | | | |
| Fritz+06 | NM | NM | NM | NM | | Fritz06 | NM | 187.3/123 | NM | 187.3/122 | C2 |
| Nenkova08 | 169.6/114 | 125.1/113 | 169.6/113 | 125.0/112 | C2 | Nenkova+08 | NM | 236.4/123 | NM | 236.4/122 | C2 |
| Hoenig10 | NM | NM | NM | NM | | Hoenig+10 | NM | NM | NM | 84.6/120 | C4 |
| Hoenig17 | NM | NM | NM | NM | | Hoenig+17 | NM | NM | NM | 22.9/117 | C4 |
| Stalevski16 | NM | NM | NM | NM | | Stalevski16 | NM | 181.6/120 | NM | 107.9/119 | C4 |
| PG2304+042 | | | | | | PG0804+761 | | | | | |
| Fritz+06 | NM | 224.7/117 | NM | 224.7/116 | C2 | Fritz+06 | 124.0/125 | 77.9/124 | 124.0/124 | 77.9/123 | C2 |
| Nenkova+08 | 184.9/118 | 146.6/117 | 185.0/117 | 146.5/116 | C2 | Nenkova+08 | NF | 198.8/124 | NF | 190.2/123 | C2 |
| Hoenig+10 | 140/119 | 82.5/68 | NM | 75.9/117 | C1 | Hoenig+10 | NM | NM | NM | 222.5/124 | C4 |
| Hoenig+17 | 77.6/116 | 65.7/115 | 76.9/115 | 65.7/114 | C2 | Hoenig+17 | 89.0/123 | 89.0/122 | 59.6/122 | 59.6/121 | C3 |
| Stalevski16 | NM | 232.2/117 | NM | NM | C2 | Stalevski16 | NM | 137.5/124 | NM | 136.0/123 | C2 |
| PKS0518-45 | | | | | | OQ208 | | | | | |
| Fritz+06 | NM | NM | NM | NM | | Fritz+06 | NM | NM | NM | NM | |
| Nenkova+08 | 178/9118 | 150.8/117 | 178.9/117 | 151.0/116 | C2 | Nenkova+08 | 152.4/124 | 45.7/123 | 184.4/123 | 39.0/122 | C4 |
| Hoenig+10 | 87.4/119 | 59.7/118 | 87.4/118 | 54.0/117 | C4 | Hoenig+10 | NM | NM | NM | NM | |
| Hoenig+17 | 36.1/116 | 36.4/115 | 35.7/115 | 35.5/114 | C1 | Hoenig+17 | NM | 215.5/121 | NM | 211.1/120 | C2 |
| Stalevski16 | 194.8/118 | 191.1/117 | 196.4/117 | 191.1/116 | C1 | Stalevski16 | NM | NM | NM | NM | |
| PG0844 + 349 | | | | | | NGC4258 | | | | | |
| Fritz+06 | 241.4/121 | 144.2/120 | NM | 144.2/119 | C2 | Fritz+06 | NM | NM | NM | NM | |
| Nenkova+08 | 162.5/121 | 37.9/120 | 162.5/120 | 37.4/119 | C2 | Nenkova+08 | NM | 122.4/110 | 82.8/110 | 85.0/109 | C4 |
| Hoenig+10 | NM | 57.7/121 | NM | 24.2/120 | C4 | Hoenig+10 | NM | NM | NM | 180.1/110 | C4 |
| Hoenig+17 | 29.2/119 | 27.1/118 | 20.3/118 | 14.9/117 | C4 | Hoenig+17 | NM | NM | 202.5/108 | 202.8/107 | C3 |
| Stalevski16 | NM | 97.2/120 | NM | 97.1/119 | C2 | Stalevski16 | NM | NM | NM | NM | |
| PG1351 + 640 | | | | | | NGC3998 | | | | | |
| Fritz+06 | NM | 187.3/123 | NM | 187.3/122 | C2 | Fritz+06 | NM | 137.8/72 | 137.8/72 | 137.8/71 | C2 |
| Nenkova+08 | NM | 236.4/123 | NM | 236.4/122 | C2 | Nenkova+08 | 30.9/73 | 23.3/72 | 27.1/72 | 21.8/71 | C2 |
| Hoenig+10 | NM | NM | NM | NM | | Hoenig+10 | 29.5/74 | 27.2/73 | 28.8/73 | 28.7/72 | C1 |
| Hoenig+17 | NM | NM | NM | NM | | Hoenig+17 | 27.9/71 | 25.3/70 | 24.6/70 | 23.0/69 | C1 |
| Stalevski16 | NM | NM | NM | NM | | Stalevski16 | NM | NM | NM | NM | |

Note.-NM = Non-modeled, indicate that the spectrum is not fitted by this component or combination of components.

Table 6. Derived parameters from smooth models of Fritz et al. (2006). Column 1 lists the name of the object modeled. Columns from 2 to 7 list the torus parameters that best reproduce the AGN-dominated IRS/Spitzer spectrum: the viewing angle i, the torus angular width σ deg, polar index γ and radial index β of the gas density distribution $\rho(r,\Theta) \propto r^{\beta} e^{-\gamma \times cos(\Theta)}$ within the torus, the radial extend $Y = R_{outer}/R_{inner}$, and the optical depth $\tau_{9.7 \, \mu m}$. Column 7 lists the covering factor which is derived using the i, γ , and τ (see equation in González-Martín et al. 2019b)

| | Parameters | | | | | | | | | | | |
|------------------|-------------------|---------------------|---------------------|---------------------|-------------------|--------------------|-----------------|--|--|--|--|--|
| Name | i (min;max) | $\Theta(\min;\max)$ | $\gamma(\min;\max)$ | $\beta(\min;\max)$ | Y (min;max) | τ (min;max) | Covering Factor | | | | | |
| | [0 - 90] deg | [20 - 60] | [0 - 6] | [-1.0 - 0.0] | [10 - 150] | [0.3 - 10] | [0 - 1] | | | | | |
| PG2304+042 | 0.0* | 20.0* | 6.0* | -0.01* | 10.0* | 9.8 (9.1;10.0) | 0.2* | | | | | |
| PG 0844+349 | 12.9 (11.8;14.9) | 20.0* | 6.0^{*} | -0.25 (-0.26;-0.20) | 11.8 (11.6;12.0) | $6.0\ (5.9;\ 6.2)$ | 0.1* | | | | | |
| PG 1351+640 | > 84.5 | < 21.0 | 6.0^{*} | -0.01* | 35.9 (35.6;36.1) | 7.4 (6.9; 7.6) | 0.1* | | | | | |
| PG 2214+139 | 70.0 (66.3; 71.6) | < 22.1 | 0.0^{*} | -0.5 (-0.6;-0.4) | 10.4 (10.2; 10.7) | 1.0 (0.9; 1.1) | 0.6* | | | | | |
| $PG\ 0804{+}761$ | 44.3 (34.1; 51.9) | < 24 | > 5.7 | -0.7 (-0.8;-0.7) | 10.0* | $3.0\ (2.7;\ 3.2)$ | > 0.1 | | | | | |
| NGC 3998 | 0.0* | < 21 | > 5.9 | -0.01* | > 142 | $2.1\ (2.1;\ 2.2)$ | 0.1* | | | | | |

Note.-*Parameter unrestricted.

Table 7. Derived parameters from clumpy models of Nenkova et al. (2008a,b). Column 1 lists the name of the object modeled. Columns from 2 to 7 list the torus parameters that best reproduce the AGN-dominated IRS/Spitzer spectrum: the viewing angle i, the number of clouds along the equatorial ray N_0 , the angular width σ deg, the radial extend Y, the index of the radial distribution of clouds q, and the optical depth τ_V . Column 7 lists the covering factor which is derived using the i, N_0 , and Θ (see equation in González-Martín et al. 2019b)

| | | | Par | ameters | | | |
|------------|-------------------|-------------------|--------------------|---------------------|--------------------|----------------------------|-----------------|
| Name | i (min;max) | N_0 (min,max) | σ (min,max) | Y (min;max) | q (min,max) | $\tau_V \text{ (min,max)}$ | Covering Factor |
| | [0 - 90] deg | [1 - 15] | $[15 - 70] \deg$ | [5 - 100] | [0.0 - 2.5] | [5 - 300] | [0 - 1] |
| NGC7213 | 0.01 (0.00;1.47) | 7.0 (6.1;7.3) | < 15.0 | 10.2 (10.0; 10.5) | < 0.01 | 55.1 (51.6; 59.5) | < 0.4 |
| PG2304+042 | 76.3 (74.0;77.4) | 13.1 (10.5; 13.6) | 17.5 (15.0; 21.7) | 10.0 (9.8;10.1) | 0.9 (0.6; 1.0) | 10.0 (10.0;10.3) | 0.5 (04;0.6) |
| PKS0518-45 | 70.8 (69.0; 75.0) | 12.8 (12.1; 14.1) | 23.6 (20.7; 29.8) | 10.0 (9.9; 10.0) | $0.1\ (0.0;\ 0.5)$ | 10.0 (10.0 10.6) | 0.6 (0.6;0.8) |
| PG0844+349 | 78.3 (66.4; 90.0) | 6.7 (4.2; 10.9) | 20.3 (15.0; 41.0) | 11.7 (10.5; 14.7) | 0.5 (0.0; 1.2) | 13.9 (10.0; 22.3) | 0.5 (0.3;0.9) |
| PG1351+640 | 0.0* | 7.6 (7.3; 10.4) | 45.0 (44.3; 45.3) | 30.3 (29.9; 31.0) | < 0.01 | 15.5 (15.1;15.9) | 0.9* |
| PG2214+139 | > 88.6 | 2.7(2.1; 3.1) | 24.7 (15.1; 39.7) | > 91 | > 2.4 | 36.8 (33.2;39.3) | > 0.5 |
| PG0804+761 | > 76.9 | 1.0* | 54.5 (27.0; 70.0) | 100.0 (79.2; 100.0) | 2.3* | 87.8 (65.9; 100.9) | > 0.5 |
| OQ208 | 25.5 (0.0; 74.0) | 1.3 (1.2; 3.4) | 64.8 (24.4; 70.0) | 11.7 (11.3; 13.5) | 0.01 (0.00; 0.81) | > 265.6 | 0.6 (0.3;0.9) |
| NGC4258 | 81.2 (72.8; 86.0) | 1.5 (1.1; 1.8) | < 20.4 | 7.8 (7.2; 8.6) | 0.01 (0.00; 0.05) | > 254.9 | < 0.3 |
| NGC3998 | > 87.5 | 1.7 (1.6; 1.8) | < 36.4 | 100.0 (96.8; 100.0) | 0.01 (0.00; 0.02) | 20.1 (16.5; 24.3) | > 0.5 |

Note.-*Parameter unrestricted.

Table 8. Derived parameters from clumpy models of Hönig & Kishimoto (2010b). Column 1 lists the name of the object modeled. Columns from 2 to 6 list the torus parameters that best reproduce the AGN-dominated IRS/Spitzer spectrum: the viewing angle i, the number of clouds along the equatorial ray N_0 , the angular width Θ , the index of the radial distribution of clouds a, and the optical depth per cloud τ_V . Column 7 lists the covering factor which is derived using the i, N_0 , and Θ (see equation in González-Martín et al. 2019b)

| | | Parameters | | | | | | | | | | |
|--------------|-------------------|--------------------|--------------------|----------------------|--------------------------|-------------------|--|--|--|--|--|--|
| Name | i (min;max) | $N_0(\min,\max)$ | θ (min,max) | a (min,max) | $\tau_V(\text{min;max})$ | Covering Factor | | | | | | |
| | [0 - 90] | [2.5 - 10.0] | [50 - 60] | [-2.0 - 0.0] | [30 - 80] | [0 - 1] | | | | | | |
| PG2304+042 | 75.4 (49.6;87.6) | < 3.2 | > 55.0 | -0.03 (-0.06; -0.01) | < 32.7 | > 0.6 | | | | | | |
| PKS0518-45 | 51.8 (38.5; 55.8) | 5.7(3.1;6.9) | 58.5 (47.7; 60.0) | -0.01 (-0.06; -0.01) | 45.9 (41.2; 63.8) | 0.9 (0.8-1.0) | | | | | | |
| PG0844 + 349 | 44.9 (0.0; 53.3) | $3.1\ (2.6;\ 4.2)$ | 55.1 (46.2; 60.0) | -0.08 (-0.17; -0.03) | > 73.9 | 0.8 (0.7-0.9) | | | | | | |
| PG2214+139 | 30.0 (25.4;33.3) | 8.9 (8.0; 9.8) | 56.7 (54.1; 59.5) | -1.1 (-1.2; -1.1) | > 78.9 | 0.97 (0.95, 0.98) | | | | | | |
| PG0804+761 | 29.9 (25.2; 30.4) | 6.9 (6.4; 7.4) | > 59.7 | -0.7* | 80.0 (79.3; 80.0) | > 1.0 | | | | | | |
| NGC4258 | 31.0 (30.0; 32.2) | > 9.7 | > 59.8 | -0.01* | > 79.6 | 1.0 | | | | | | |
| NGC3998 | 75.1 (59.1; 90.0) | 2.9(2.5; 3.6) | 54.7 (32.7; 60.0) | -0.3 (-0.5; -0.2) | < 43.8 | 0.8 (0.6;0.9) | | | | | | |

Note.- *Parameter unrestricted.

Table 9. Derived parameters from disk+outflow models of Hönig & Kishimoto (2017). Column 1 lists the name of the object modeled. Columns from 2 to 9 list the disk and outflow parameters that best reproduce the AGN-dominated IRS/Spitzer spectrum: Disk: index of the radial distribution of clouds a, number of clouds along the equatorial ray N_0 , the scale height in the vertical Gaussian distribution of clouds h. Wind: index of the radial distribution of clouds in the wind a_w ; the half-opening angle of the wind Θ_w , and its angular width σ_{Θ} . The viewing angle i and the ratio between the number of clouds along the cone and N_0 , f_{wd} . Column 7 lists the covering factor which is derived using the i, N_0 , Θ_w and, σ_{Θ} (see equation in González-Martín et al. 2019b)

| | | | | Par | rameters | | | | |
|--------------|-------------------|------------------|-------------------|-----------------------|------------------------------|-------------------|-------------------|-------------------------|-----------------|
| Name | i(min,max) | $N_0(\min,\max)$ | $a(\min,\max)$ | $\Theta_w(\min,\max)$ | $\sigma_{\theta}(\min,\max)$ | $a_w(\min,\max)$ | $h(\min,\max)$ | f_{wd} (min,max) | Covering Factor |
| | [0 - 90] | [5 - 10] | [-3.00.5] | [7 - 15] deg | $[30 - 45] \deg$ | [-2.50.5] | [0.1 - 0.5] | [0.15 - 0.75] | [0 - 1] |
| PG2304+042 | 45.0 (44.1; 45.9) | < 5.3 | -2.5 (-2.7; -2.5) | 10.4 (9.2; 11.2) | > 44.6 | -0.5* | > 0.4 | 0.30 (0.30; 0.33) | > 0.3 |
| PKS0518-45 | 29.6 (21.2; 33.9) | < 6.1 | -2.2 (-2.4; -2.0) | > 9.0 | 38.8 (33.8; 43.0) | -0.5 (-0.9; -0.5) | 0.12 (0.10; 0.16) | > 0.41 | > 0.2 |
| PG0844 + 349 | 18.6 (0.1; 26.9) | > 6.7 | -2.0 (-2.3; -1.6) | > 8.5 | 34.8 (33.5; 42.7) | -0.5 (-0.8; -0.5) | 0.11 (0.10; 0.19) | $0.73 \ (0.42; \ 0.75)$ | > 0.2 |
| PG2214+139 | 15.6 (0.0; 16.8) | > 6.6 | -3.0 (-3.0; -2.8) | 8.5 (7.0; 13.5) | > 35.5 | -0.5 (-0.9; -0.5) | 0.4 (0.3; 0.5) | > 0.40 | > 0.4 |
| PG0804+761 | 0.0* | > 9.7 | -2.4 (-2.4; -1.8) | 10.3 (9.8; 10.8) | > 44.3 | -0.5* | 0.18 (0.17; 0.20) | > 0.73 | > 0.2 |
| OQ208 | 0.0* | 10.0^{*} | -0.5* | 9.9 (9.7; 10.1) | > 44.9 | -0.5 (-0.5; -0.5) | 0.5^{*} | $0.75 \ (0.74; \ 0.75)$ | > 0.4 |
| NGC4258 | 30.0 (25.6; 30.4) | > 9.7 | -2.1 (-2.1; -2.0) | 7.1 (7.0; 7.5) | > 44.5 | -0.5 (-0.5; -0.5) | 0.1^{*} | > 0.74 | > 0.1 |
| NGC3998 | < 26.31 | < 6.6 | -0.5 (-1.1; -0.5) | 12.7^{*} | 33.4 (31.0; 45.0) | -2.5 (-2.5; -1.5) | 0.1^* | > 0.27 | 0.2* |

Note.-*Parameter unrestricted.

Table 10. Derived parameters from two-phase media models of Stalevski et al. (2016). Column 1 lists the name of the object modeled. Column 2 lists the viewing angle i, column 2 lists the angular width σ of the torus. Column 3 and 4 list the indices of the radial (p) and angular distribution (q) of the clouds. Column 5 gives the ratio between the outer and inner radius. Column 6 gives the optical depth $\tau_{9.7}$. Columns 7 list the covering factor.

| | Parameters | | | | | | | | | | | |
|------------|-------------------|---------------------|----------------------|----------------------|-------------------|------------------------------|-----------------|--|--|--|--|--|
| Name | $i(\min,\max)$ | $\sigma(\min,\max)$ | $p(\min,\max)$ | $q(\min,\max)$ | $Y(\min,\max)$ | $\tau_{9.7}(\text{min,max})$ | Covering Factor | | | | | |
| | [0-90] | [10-80] | [0-1.5] | [0-1.5] | [10-30] | [3-11] | [0-1] | | | | | |
| PG2304+042 | > 83.5 | 80.0 (60.0;79.1) | 0.01 (0.0;0.02) | 1.0 (1.0; 1.1) | 10.0* | 3.0 (0.1; 3.1) | > 0.7 | | | | | |
| PKS0518-45 | > 66.5 | 80.0 (60.0;79.0) | $0.01\ (0.0;\ 0.01)$ | $0.7 \ (0.6; \ 0.7)$ | 10.0* | 3.0 (0.1; 3.1) | > 0.8 | | | | | |
| PG0844+349 | > 80.5 | 80.0 (60.0;76.2) | $0.01\ (0.0;\ 0.01)$ | 1.5^{*} | 12.3 (11.9; 12.8) | 3.1 (0.1; 3.3) | > 0.6 | | | | | |
| PG2214+139 | 63.0 (58.3; 64.9) | 70.0 (67.8; 74.4) | > 1.5 | 1.5^{*} | < 10.1 | 3.8 (3.6; 4.2) | 0.7* | | | | | |
| PG0804+761 | 10.0 (0.0; 14.1) | 42.3 (41.6; 44.1) | > 1.5 | 1.5* | < 10.2 | 10.8* | 1.0* | | | | | |

Note.-*Parameter unrestricted.




Figure 17. The viewing angle Vs the angular width derived from the Si-s sample for the smooth dusty torus of Fritz06, the clumpy torus models of Nenkova08 and Hoenig10, the disk+outflow torus model of Hoenig10, and the two-phase media models of stalev16.




Figure 18. Modeling and residuals of the IRS/Spitzer spectrum of PG 2304+042. From top to bottom is the spectrum fitted assuming Fritz06 ($\chi^2_{red} \sim 1.20$), Nekova08 ($\chi^2_{red} \sim 0.32$), Hoenig10 ($\chi^2_{red} \sim 0.20$), Hoenig17 ($\chi^2_{red} \sim 0.13$), and Stalevski16 ($\chi^2_{red} \sim 0.81$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 19. Modeling and residuals of the IRS/Spitzer spectrum of NGC 7213. From top to bottom is the spectrum fitted assuming Nekova08 ($\chi^2_{red} \sim 1.11$). The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in erg s⁻¹cm⁻², and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 20. Modeling and residuals of the IRS/Spitzer spectrum of PKS 0518-45. From top to bottom is the spectrum fitted assuming Nekova08 ($\chi^2_{red} \sim 1.29$), Hoenig10 ($\chi^2_{red} \sim 0.46$), Hoenig17 ($\chi^2_{red} \sim 0.31$), and Stalevski16 ($\chi^2_{red} \sim 1.65$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 21. Modeling and residuals of the IRS/Spitzer spectrum of PG 1351+640. From top to bottom is the spectrum fitted assuming Fritz06 ($\chi^2_{red} \sim 1.52$), Nekova08 ($\chi^2_{red} \sim 1.92$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 22. Modeling and residuals of the IRS/Spitzer spectrum of PG 2214+139. From top to bottom is the spectrum fitted assuming Fritz06 ($\chi^2_{red} \sim 1.52$), Nekova08 ($\chi^2_{red} \sim 1.92$), Hoenig10 ($\chi^2_{red} \sim 0.71$), Hoenig17 ($\chi^2_{red} \sim 0.20$), and Stalevski16 ($\chi^2_{red} \sim 0.91$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 23. Modeling and residuals of the IRS/Spitzer spectrum of PG 0804+761. From top to bottom is the spectrum fitted assuming Fritz06 ($\chi^2_{red} \sim 0.63$), Nekova08 ($\chi^2_{red} \sim 1.60$), Hoenig10 ($\chi^2_{red} \sim 1.79$), Hoenig17 ($\chi^2_{red} \sim 0.49$), and Stalevski16 ($\chi^2_{red} \sim 1.11$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 24. Modeling and residuals of the IRS/Spitzer spectrum of NGC4258. From top to bottom is the spectrum fitted assuming Nekova08 ($\chi^2_{red} \sim 0.78$), Hoenig10 ($\chi^2_{red} \sim 1.64$), and Hoenig17 ($\chi^2_{red} \sim 1.88$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 25. Modeling and residuals of the IRS/Spitzer spectrum of OQ208. From top to bottom is the spectrum fitted assuming Nekova08 ($\chi^2_{red} \sim 0.32$), and Hoenig17 ($\chi^2_{red} \sim 1.78$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.




Figure 26. Modeling and residuals of the IRS/Spitzer spectrum of NGC3998. From top to bottom is the spectrum fitted assuming Fritz06 ($\chi^2_{red} \sim 1.91$), Nekova08 ($\chi^2_{red} \sim 0.32$), Hoenig10 ($\chi^2_{red} \sim 0.40$), and Hoenig17 ($\chi^2_{red} \sim 0.39$) model. The last panel show the residuals defined as the ratio between the data and model. In all panels the black points are the IRS/Spitzer spectrum and its error in (erg s⁻¹cm⁻²), and the red solid line is the fitted torus model. The orange line is the total SED that results when more than one component, the stellar (green dot-dashed line) and/or HII (blue dotted line), is added to model the spectrum.