# Asymptotic behavior at infinity of solutions of Lagrangian mean curvature equations \*

Jiguang Bao<sup>†</sup>, Zixiao Liu<sup>‡</sup>

School of Mathematical Sciences, Beijing Normal University Laboratory of Mathematics and Complex Systems, Ministry of Education Beijing 100875, China

February 17, 2022

#### Abstract

We studied the asymptotic behavior of solutions with quadratic growth condition of a class of Lagrangian mean curvature equations  $F_{\tau}(\lambda(D^2u)) = f(x)$  in exterior domain, where fsatisfies a given asymptotic behavior at infinity. When f(x) is a constant near infinity, it is not necessary to demand the quadratic growth condition anymore. These results are a kind of exterior Liouville theorem, and can also be regarded as an extension of theorems of Pogorelov [33], Flanders [16] and Yuan [42, 43].

Mathematics Subject Classification 2000: 35J60; 35B40

Keywords: Special Lagrangian equation, Entire solutions, Asymptotic behaviors.

#### Contents

2Constant Right Hand Side Situation73Convergence of Hessian at infinity134Analysis of Linearized Equation205Proof of Theorem 1.5246Proof of Theorem 1.6287Interior Estimates36	1	Introduction	2
3Convergence of Hessian at infinity134Analysis of Linearized Equation205Proof of Theorem 1.5246Proof of Theorem 1.6287Interior Estimates36	2	Constant Right Hand Side Situation	7
4Analysis of Linearized Equation205Proof of Theorem 1.5246Proof of Theorem 1.6287Interior Estimates36	3	Convergence of Hessian at infinity	13
5Proof of Theorem 1.5246Proof of Theorem 1.6287Interior Estimates36	4	Analysis of Linearized Equation	20
6Proof of Theorem 1.6287Interior Estimates36	5	Proof of Theorem 1.5	24
7 Interior Estimates 36	6	Proof of Theorem 1.6	28
	7	Interior Estimates	36

\*Supported in part by National Natural Science Foundation of China (11871102 and 11631002). \*E-mail address: jgbao@bnu.edu.cn

<sup>&</sup>lt;sup>‡</sup>Corresponding author, E-mail address: liuzixiao@mail.bnu.edu.cn

#### 8 Perturbed Results for $\tau = \frac{\pi}{4}$

# **1** Introduction

In 2010, M.Warren [38] first studied the minimal/maximal Lagrangian graph in  $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$ , where

$$g_{\tau} = \sin \tau \delta_0 + \cos \tau g_0, \quad \tau \in \left[0, \frac{\pi}{2}\right],$$

is the linearly combined metric of standard Euclidean metric

$$\delta_0 = \sum_{i=1}^n dx_i \otimes dx_i + \sum_{j=1}^n dy_j \otimes dy_j,$$

and the pseudo-Euclidean metric

$$g_0 = \sum_{i=1}^n dx_i \otimes dy_i + \sum_{j=1}^n dy_j \otimes dx_j.$$

He showed that if  $u \in C^2(\Omega), \ \Omega \subset \mathbb{R}^n$  is a solution of

$$F_{\tau}\left(\lambda\left(D^{2}u\right)\right) = C_{0}, \quad x \in \Omega, \tag{1.1}$$

then the volume of (x, Du(x)) is a maximal (for  $\tau \in (0, \frac{\pi}{4})$ ) /minimal (for  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$ ) among all homologous,  $C^1$ , space-like *n*-surfaces in  $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$ , where  $C_0$  is a constant,  $\lambda(D^2u) = (\lambda_1, \lambda_2, \cdots, \lambda_n)$  are *n* eigenvalues of the Hessian matrix  $D^2u$  and

$$F_{\tau}(\lambda) := \begin{cases} \frac{1}{n} \sum_{i=1}^{n} \ln \lambda_{i}, & \tau = 0, \\ \frac{\sqrt{a^{2} + 1}}{2b} \sum_{i=1}^{n} \ln \frac{\lambda_{i} + a - b}{\lambda_{i} + a + b}, & 0 < \tau < \frac{\pi}{4}, \\ -\sqrt{2} \sum_{i=1}^{n} \frac{1}{1 + \lambda_{i}}, & \tau = \frac{\pi}{4}, \\ \frac{\sqrt{a^{2} + 1}}{b} \sum_{i=1}^{n} \arctan \frac{\lambda_{i} + a - b}{\lambda_{i} + a + b}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \sum_{i=1}^{n} \arctan \lambda_{i}, & \tau = \frac{\pi}{2}, \end{cases}$$

 $a = \cot \tau, b = \sqrt{|\cot^2 \tau - 1|}.$ 

If  $\tau = 0$ , then (1.1) becomes the famous Monge-Ampère equation

$$\det D^2 u = e^{nC_0}.$$

If  $\tau = \frac{\pi}{2}$ , then (1.1) becomes the special Lagrangian equation

$$\sum_{i=1}^{n} \arctan \lambda_i \left( D^2 u \right) = C_0.$$

2

In the same paper [38], M.Warren used change of variable and restated the Bernstein-type results of Jörgens [23]-Calabi [10]-Pogorelov [33], Flanders [16], and Yuan [42, 43] to give the following statement.

**Theorem 1.1.** u is a quadratic polynomial if  $u \in C^2(\mathbb{R}^n)$  is a solution of (1.1) respectively, in the following cases, with

- 1.  $D^2u > (-a+b)I$  for  $\tau \in (0, \frac{\pi}{4})$ ;
- 2.  $D^2 u > 0$  for  $\tau = \frac{\pi}{4}$ ;
- 3. either

$$D^2 u \ge -aI,\tag{1.2}$$

or

$$D^2 u > -(a+b)I, \quad \left|\frac{bC_0}{\sqrt{a^2+1}} + \frac{n\pi}{4}\right| > \frac{n-2}{2}\pi,$$
 (1.3)

for  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$ .

The aim of this paper is to study the asymptotic behavior of classical solutions of equation (1.1) in exterior domain, including the stability with respect to the right hand side term. There are plenty of marvelous work on these type of questions. The classical theorem by Jörgens [23], Calabi [10] and Pogorelov [33] states that any convex classical solution of det  $D^2u = 1$  on  $\mathbb{R}^n$  must be a quadratic polynomial. See Cheng and Yau [12], Caffarelli [8] and Jost and Xin [24] for different proofs and extensions. For Monge-Ampère equations with constant right hand side term in exterior domain, there are exterior Jörgens-Calabi-Pogorelov type results [15] (for n = 2) and [5], which state that all convex solutions are asymptotic to quadratic polynomials (for n = 2 we need additional log-term) near infinity. Pioneered by Gutièrrez and Huang [19], Bernstein-type result of parabolic convex entire solution also holds for parabolic Monge-Ampère type equation  $-u_t \det D^2 u = 1$ . Many other type of equations including parabolic version with constant right hand side term in exterior domain is also studied through different strategies, see for instance [2, 40, 42, 43, 36, 45, 41, 30]. Also, these type of equations with perturbed constant right hand side or periodic right hand side are also studied a lot, see for instance [3, 44, 4, 34].

First, we consider the equation with constant right hand side in exterior domain

$$F_{\tau}(\lambda(D^2 u)) = C_0, \quad \text{in} \quad \mathbb{R}^n \setminus \overline{B_1}.$$
(1.4)

We prove the following results that provide an asymptotic behavior at infinity. The idea is to use Legendre transform and apply the strategies in [26].

From now on, we let  $n \ge 3$ , I stand for the unit  $n \times n$  matrix and let Sym(n) denote the set of symmetric  $n \times n$  constant matrix. For any  $C, D \in \mathbb{R}, \tau \in [0, \frac{\pi}{2}]$ , let

$$\mathcal{A}_{\tau}(C,D) := \{ A \in \operatorname{Sym}(n) : F_{\tau}(\lambda(A)) = C, \text{ and } A \ge DI \}.$$

To shorten notation, if  $\tau \in [0, \frac{\pi}{2}]$  is fixed, we write  $\mathcal{A}(C, D)$  instead of  $\mathcal{A}_{\tau}(C, D)$ .

**Theorem 1.2.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) for some constant  $C_0$  with  $\tau \in (0, \frac{\pi}{4})$  and satisfy

$$D^2 u > (-a+b)I, \quad in \quad \mathbb{R}^n \setminus \overline{B_1}.$$
 (1.5)

Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(C_0, -a+b)$  such that

$$\limsup_{|x|\to\infty} |x|^{n-2+k} \left| D^k \left( u(x) - \left( \frac{1}{2} x' A x + \beta \cdot x + \gamma \right) \right) \right| < \infty, \quad k = 0, 1, 2.$$

$$(1.6)$$

**Theorem 1.3.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) for some constant  $C_0$  with  $\tau = \frac{\pi}{4}$  and satisfy

$$D^2 u > -I, \quad in \quad \mathbb{R}^n \setminus \overline{B_1},$$
 (1.7)

Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(C_0, -1)$  such that (1.6) holds.

**Theorem 1.4.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) for some constant  $C_0$  with  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$  and satisfy

$$D^2 u \ge -(a - \varepsilon_0)I,\tag{1.8}$$

for some  $\varepsilon_0 > 0$  or (1.3). Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(C_0, -a + \varepsilon_0)$  or  $A \in \mathcal{A}(C_0, -(a+b))$  respectively such that (1.6) holds.

By comparison principle as in [5], the global Theorem 1.1 can be obtained by these exterior behavior results i.e. Theorems 1.2, 1.3 and 1.4, apart from the situation of  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$  with only (1.2) instead of (1.8).

Second, we also consider equation with right hand side of a perturbation of suitable constant as the following

$$F_{\tau}\left(\lambda(D^2 u)\right) = \frac{\sqrt{a^2 + 1}}{2b} \sum_{i=1}^n \ln \frac{\lambda_i + a - b}{\lambda_i + a + b} = f(x), \quad \text{in} \quad \mathbb{R}^n, \tag{1.9}$$

for  $\tau \in (0, \frac{\pi}{4})$ . Especially in this situation, a > b always holds.

The main results in this part include

**Theorem 1.5.** Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.9) and satisfy

$$(-a+b)I < D^2 u \le MI, \tag{1.10}$$

for some constant M. Assume that  $f \in C^2(\mathbb{R}^n)$ , and

$$\limsup_{|x|\to\infty} |x|^{n+\varepsilon} |D^k(f(x) - f(\infty))| < \infty, \quad k = 0, 1, 2,$$
(1.11)

for some  $\varepsilon > 0$ . Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(f(\infty), -a+b)$  such that (1.6) holds.

**Theorem 1.6.** Let  $u \in C^{m+2}(\mathbb{R}^n)$  be a classical solution of (1.9) and satisfy (1.10). Assume that  $f \in C^m(\mathbb{R}^n)$ , and

$$\lim_{|x|\to\infty} \sup_{|x|^{\zeta+k}} \left| D^k(f(x) - f(\infty)) \right| < \infty, \quad \forall k = 0, 1, \cdots, m,$$
(1.12)

for some  $\zeta > 2, m \ge 3$ . Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(f(\infty), -a+b)$  such that

$$\lim_{|x|\to\infty} \sup |x|^{\min\{\zeta,n\}-2+k} \left| D^k \left( u(x) - \left(\frac{1}{2}x'Ax + \beta \cdot x + \gamma\right) \right) \right| < \infty, \quad \forall k = 0, 1, \cdots, m+1.$$
(1.13)

**Remark 1.7.** We can see that these two conditions on f in Theorems 1.5 and 1.6 doesn't include each other. The condition (1.11) holds true for more "un-regular" f than condition (1.12), which is originally proposed by Bao-Li-Zhang [3]. Condition (1.11) only demands the behavior of up to 2-order derivative of f, while condition (1.12) demands the behavior of at least 3-order derivative of f. Generally speaking, for  $n \ge 5$ , the condition (1.11) in Theorem 1.5 demands a higher convergence speed than condition (1.12) in Theorem 1.6, but the demand on regularity is the other way around.

On the one side, we take  $f(x) = |x|^{-(2+\varepsilon)} - 1$  for  $|x| \ge 1$ ,  $1 > \varepsilon > 0$ . Then it satisfies the condition (1.12) in Theorem 1.6 by picking  $\zeta := 2 + \varepsilon > 2$  but doesn't satisfies the condition (1.11) in Theorem 1.5.

On the other side, we take  $f(x) := e^{-m|x|} \sin(e^{|x|}) - 1$ ,  $|x| \ge 1$ ,  $m \ge 3$ . Then it satisfies the condition (1.11) in Theorem 1.5 because it has exponential decay up to (m-1)-order derivatives. But it doesn't satisfies the condition (1.12) in Theorem 1.6 because its m-order derivative doesn't admits a limit at infinity.

Next, we can weaken the assumption (1.10) in Theorems 1.5 and 1.6 into the linear growth of gradient

$$|Du(x)| \le C(1+|x|), \quad \forall x \in \mathbb{R}^n, \tag{1.14}$$

for some constant C. And further more, this linear growth of gradient condition can be weakened into the quadratic growth condition of u itself

$$|u(x)| \le C(1+|x|^2), \quad \forall x \in \mathbb{R}^n,$$
(1.15)

for some constant C. Since in the following Theorems 1.8, 1.9 and Remark 1.10, we always assume a lower bound of Hessian to make the equation elliptic, we can weaken the additional condition (1.15) into

$$u(x) \le C(1+|x|^2), \quad \forall x \in \mathbb{R}^n,$$
(1.16)

for some constant C.

To be more precise, we have the following stronger theorems than Theorems 1.5 and 1.6.

**Theorem 1.8** (Linear Growth of Gradient). Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.9) with  $D^2u > (-a+b)I$  and satisfy (1.14). Suppose that f satisfies the assumptions as in Theorem 1.5 (resp. Theorem 1.6), then we have the same results as in Theorem 1.5 (resp. Theorem 1.6).

**Theorem 1.9** (Quadratic Growth of Solution). Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.9) with  $D^2u > (-a+b)I$  and satisfy (1.16). Suppose that f satisfies the assumptions as in Theorem 1.5 (resp. Theorem 1.6), then we have the same results as in Theorem 1.5 (resp. Theorem 1.6).

As in the famous paper of Bao-Chen-Guan-Ji [2], such a quadratic growth condition is common and necessary for general k-Hessian equations and Hessian quotient equations. See for instance [30, 11, 39]. And as for the classical special Lagrangian equation, the linear growth condition of gradient is also studied by M.Warren and Y.Yuan in [40]. Similar conditions are also considered in different paper, see for instance [27, 14, 25, 35, 31]. **Remark 1.10.** By using extension theorems, all the results in Theorems 1.5, 1.6, 1.8 and 1.9 still hold when the equation (1.9) only holds in exterior domain i.e.

$$F_{\tau}\left(\lambda(D^{2}u)\right) = \frac{\sqrt{a^{2}+1}}{2b} \sum_{i=1}^{n} \ln \frac{\lambda_{i}+a-b}{\lambda_{i}+a+b} = f(x), \quad in \quad \mathbb{R}^{n} \setminus \overline{\Omega},$$
(1.17)

where  $\Omega$  is a bounded domain in  $\mathbb{R}^n$ .

Apart from the asymptotic behavior of solutions of (1.9) we stated earlier, the method we adapt here also works for Monge-Ampère equation (i.e.  $\tau = 0$  situation) without demanding the boundedness of Hessian. The Theorem 1.11 below is proved by Bao-Li-Zhang [3] and we obtain a new result in Theorem 1.12 under a different setting.

**Theorem 1.11.** Let  $u \in C^0(\mathbb{R}^n)$  be a convex viscosity solution of

$$\det D^2 u = f(x), \quad in \quad \mathbb{R}^n, \tag{1.18}$$

where  $f \in C^0(\mathbb{R}^n)$ ,  $D^m f$   $(m \ge 3)$  exist outside a compact subset of  $\mathbb{R}^n$  and (1.12) holds for some given  $f(\infty) \in (0, +\infty)$ . Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}_0(\frac{1}{n}\exp(f(\infty)), 0)$ such that (1.13) holds.

**Theorem 1.12.** Let  $u \in C^0(\mathbb{R}^n)$  be a convex viscosity solution of (1.18), where  $f \in C^0(\mathbb{R}^n)$ ,  $D^2 f$  exist outside a compact subset of  $\mathbb{R}^n$  and there exists a  $\varepsilon > 0$  such that (1.11) holds for some given  $f(\infty) \in (0, +\infty)$ . Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}_0(\frac{1}{n} \exp(f(\infty)), 0)$  such that (1.6) holds.

**Remark 1.13.** The difference between the conditions on f in Theorems 1.11 and 1.12 is the same as in Remark 1.7. As in Remark 1.10, all the results in Theorems 1.11 and 1.12 still hold when the equation (1.18) only happens on exterior domain i.e.

$$\det D^2 u = f(x), \quad in \quad \mathbb{R}^n \setminus \overline{\Omega}, \tag{1.19}$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^n$ .

For perturbed right hand side case of  $\tau = \frac{\pi}{4}$ , since the equation after Legendre transform becomes Laplacian operator, the proof is even simpler than the proof of Theorems 1.5 and 1.6. To be more precise, we obtain the following results.

**Theorem 1.14.** Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of the following equation with  $\tau = \frac{\pi}{4}$ ,

$$F_{\tau}(\lambda(D^2 u)) = -\sqrt{2} \sum_{i=1}^{n} \frac{1}{1 + \lambda_i(D^2 u)} = f(x), \quad \text{in} \quad \mathbb{R}^n, \tag{1.20}$$

and satisfy

$$-I < D^2 u \le MI, \tag{1.21}$$

for some constant M. Assume that  $f \in C^2(\mathbb{R}^n)$ , and satisfy (1.11) for some  $\varepsilon > 0$ . Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(f(\infty), -1)$  such that (1.6) holds.

**Theorem 1.15.** Let  $u \in C^{m+2}(\mathbb{R}^n)$  be a classical solution of (1.20) with  $\tau = \frac{\pi}{4}$  and satisfy (1.21) for some constant M. Assume that  $f \in C^m(\mathbb{R}^n)$  and satisfies (1.12) for some  $\zeta > 2$ ,  $m \ge 3$ . Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(f(\infty), -1)$  such that (1.13) holds.

**Theorem 1.16.** Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.20) with  $D^2u > -I$  and satisfy (1.14). Suppose that f satisfies the assumptions as in Theorem 1.14 (resp. Theorem 1.15), then we have the same results as in Theorem 1.14 (resp. Theorem 1.15).

**Theorem 1.17.** Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.20) with  $D^2u > -I$  and satisfy (1.16). Suppose that f satisfies the assumptions as in Theorem 1.14 (resp. Theorem 1.15), then we have the same results as in Theorem 1.14 (resp. Theorem 1.15).

**Remark 1.18.** As in Remarks 1.7 and 1.10, similar discussion work for  $\tau = \frac{\pi}{4}$  as well. Whether similar result still holds for  $\frac{\pi}{4} < \tau \leq \frac{\pi}{2}$  is still unclear.

This paper is organized in the following order. In section 2, we prove the asymptotic behavior of solutions of equations (1.4) with constant right hand side. Then, we move on to study the equation with perturbed right hand side. In section 3 we prove that under suitable conditions, the Hessian  $D^2u$  converge to suitable constant matrix  $A \in \text{Sym}(n)$  with  $|x|^{-\alpha}$  convergence rate for some  $\alpha > 0$  (Theorem 3.5). In section 4 we prepare some necessary analysis of linearized elliptic equations that will be used later on. In sections 5 and 6 we capture the linear and constant part of the solution at infinity, due to the difference of tools we apply, the conditions on RHS term f(x)are slightly different from [3]. Finally in section 7 we prove that under suitable conditions of f, the bounded Hessian condition follows from interior Hessian estimate and linear growth condition of gradient, which follows from the line of [29]. Also, as a corollary of Y.Y.Li's paper [28], we notice that the linear growth condition of gradient follows from interior gradient estimate and quadratic growth condition of solution itself. Eventually in section 8, we provide the results for  $\tau = \frac{\pi}{4}$ .

### 2 Constant Right Hand Side Situation

In this section, we give an rigorous proof of Theorems 1.2, 1.3 and 1.4 based on Theorem 2.1 in [26]. For reading simplicity, we repeat the statement of this theorem.

Theorem 2.1 (Exterior Liouville Theorem ). Let u be a smooth solution of

$$F\left(D^2u\right) = 0,$$

in the exterior domain  $\mathbb{R}^n \setminus \overline{B_1}$ , F is smooth, uniformly elliptic and the level set  $\{M | F(M) = 0\}$  is concave. Suppose for some constant K,

$$\left\| D^2 u \right\|_{L^{\infty}\left(\mathbb{R}^n \setminus \overline{B}_1\right)} \le K.$$

Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \{A \in Sym(n) : F(A) = 0, ||A|| \le K\}$  such that (1.6) holds.

By the important characterise results on ellipticity and concavity by Caffarelli-Nirenberg-Spruck [6],  $F(D^2u) = f(\lambda(D^2u))$  is uniformly elliptic and concave if

$$\lambda \leq \frac{\partial f}{\partial \lambda_i} \leq \Lambda, \quad \forall i = 1, 2, \cdots, n$$

for some constants  $0 < \lambda \leq \Lambda < \infty$  and f is a concave function.

Now we move on to study our targets. The section is organized in the following order. We will discuss the situation of  $0 < \tau < \frac{\pi}{4}$  situation in the first part,  $\tau = \frac{\pi}{4}$  situation in the second part and  $\frac{\pi}{4} < \tau < \frac{\pi}{2}$  in the third part.

#### **2.1** $0 < \tau < \frac{\pi}{4}$ Situation

In this part, we focus on equation (1.4) with  $0 < \tau < \frac{\pi}{4}$ . We introduce the following Legendre transform and apply Theorem 2.1 twice to prove the asymptotic behavior.

Let  $u \in C^2(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) and satisfy (1.5). We can extend the function u sufficiently smooth to  $\mathbb{R}^n$  such that  $D^2u > -(a+b)I$  in  $\mathbb{R}^n$ . Let

$$\overline{u}(x) := u(x) + \frac{a+b}{2}|x|^2,$$

then by (1.5) we have

$$D^2 \overline{u} = D^2 u + (a+b)I > 2bI$$
, in  $\mathbb{R}^n$ 

Let  $(\tilde{x}, \tilde{v})$  be the Legendre transform of  $(x, \bar{u})$  i.e.

$$\left\{ \begin{array}{l} \widetilde{x} := D\overline{u}(x) = Du(x) + (a+b)x \\ D_{\widetilde{x}}v(\widetilde{x}) := x \end{array} \right.$$

and we have

$$D_{\tilde{x}}^2 v(\tilde{x}) = \left(D^2 \bar{u}(x)\right)^{-1} = \left(D^2 u(x) + (a+b)I\right)^{-1} < \frac{1}{2b}I$$

This transform is exactly the classical Legendre transform, hence it is proved rigourously that such a scaler function  $v(\tilde{x})$  exists. Then we take

$$\widetilde{u}(\widetilde{x}) := \frac{1}{2} |\widetilde{x}|^2 - 2b \cdot v(\widetilde{x}), \qquad (2.1)$$

which satisfies

$$\widetilde{\lambda}_i \left( D^2 \widetilde{u} \right) = 1 - 2b \cdot \frac{1}{\lambda_i + a + b} = \frac{\lambda_i + a - b}{\lambda_i + a + b} \in (0, 1).$$
(2.2)

Then we see that  $\widetilde{u}(\widetilde{x})$  satisfies the following Monge-Ampère type equation

$$G(\widetilde{\lambda}(D^{2}\widetilde{u})) := \sum_{i=1}^{n} \ln \widetilde{\lambda}_{i} = \frac{2b}{\sqrt{a^{2}+1}} C_{0}, \text{ in } \mathbb{R}^{n} \setminus \overline{\Omega},$$
(2.3)

for some bounded set  $\Omega = D\overline{u}(B_1) = Du(B_1) + (a+b)B_1$ .

Moreover, we see that for any  $x \in \mathbb{R}^n, \ \widetilde{x} = D\bar{u}(x)$ ,

$$|\tilde{x} - 0| = |Du(x) - Du(0) + (a+b)x| > 2b|x|.$$

Hence triangle inequality gives us

$$|\widetilde{x}| \ge -|\widetilde{0}| + |\widetilde{x} - \widetilde{0}| > -|\widetilde{0}| + 2b|x|.$$

$$(2.4)$$

Especially we have

$$\lim_{x|\to\infty} |\widetilde{x}| = \infty$$

Now we are ready to prove Theorem 1.2.

*Proof of Theorem 1.2.* First we prove that there exists a positive lower bound such that  $\lambda_i$  is uniformly strictly positive.

It follows from equation (1.4) and the fact that  $\ln \frac{\lambda_i + a - b}{\lambda_i + a + b} \leq 0$ ,  $\forall i = 1, 2, \dots, n$ , we have

$$\ln \widetilde{\lambda_i} = \ln \frac{\lambda_i + a - b}{\lambda_i + a + b} \ge \frac{2b}{\sqrt{a^2 + 1}} C_0, \quad \forall i = 1, 2, \cdots, n$$

Thus it follows immediately that  $\widetilde{\lambda_i} \ge e^{\frac{2b}{\sqrt{a^2+1}}C_0} > 0$  and by  $C_0 < 0$ ,

$$\lambda_i \ge \frac{2b}{1 - e^{\frac{2b}{\sqrt{a^2 + 1}}C_0}} - a - b = -a + b + \frac{e^{\frac{2b}{\sqrt{a^2 + 1}}C_0}}{1 - e^{\frac{2b}{\sqrt{a^2 + 1}}C_0}}2b =: -a + b + \delta.$$
(2.5)

Note that for  $\tilde{\lambda}_i \in [e^{\frac{2b}{\sqrt{a^2+1}}C_0}, 1]$ , taking derivatives and we see  $\frac{\partial G}{\partial \tilde{\lambda}_i}(\tilde{\lambda})$  has positive lower and upper bound,  $\frac{\partial^2 G}{\partial \tilde{\lambda}_i^2}(\tilde{\lambda})$  has negative upper bound. Hence the equation (2.3) is uniformly elliptic and concave.

Applying Theorem 2.1 we learn that there exists some

$$\widetilde{A} \in \{\widetilde{A} \in \operatorname{Sym}(n) : G(\lambda(\widetilde{A})) = \frac{2b}{\sqrt{a^2 + 1}} C_0, \ \lambda_i(\widetilde{A}) \in [e^{\frac{2b}{\sqrt{a^2 + 1}}C_0}, 1]\},$$

such that

$$\lim_{|\widetilde{x}| \to +\infty} D^2 \widetilde{u}(\widetilde{x}) = \widetilde{A}.$$
(2.6)

From the relationship of Legendre transform, we have

$$D^{2}\tilde{u}(\tilde{x}) = I - 2b(D^{2}u(x) + (a+b)I)^{-1}$$
, and  $\tilde{x} = Du(x) + (a+b)x$ .

Now we show that all the eigenvalues  $\lambda_i(\widetilde{A})$  are strictly less than 1, which tells us  $I - \widetilde{A}$  is an invertible matrix. Argue by contradiction, by rotating the  $\widetilde{x}$  -space to make  $\widetilde{A}$  diagonal, we may assume that  $\widetilde{A}_{11} = 1$ . Then by the asymptotic behavior of  $D\widetilde{u}$  and hence Dv (take k = 1 in formula (1.6) of Theorem 2.1), we have the following asymptote for some  $\widetilde{\beta}_1 \in \mathbb{R}$ 

$$D_{\widetilde{x}_1}v = \widetilde{\beta}_1 + O(|\widetilde{x}|^{1-n}),$$

as  $|\tilde{x}| \to \infty$ . Thus we infer from the definition of Legendre transform (2.1) and (2.4) that

$$x_1 = D_{\widetilde{x}_1} v(\widetilde{x}) = \widetilde{\beta}_1 + O\left(|\widetilde{x}|^{1-n}\right), \qquad (2.7)$$

as  $|\tilde{x}| \to \infty$ . which means that  $\mathbb{R}^n \setminus \overline{B_1}$  is bounded in the  $x_1$  -direction, hence this becomes a contradiction. Thus  $\lambda_i(\tilde{A}) < 1$  strictly for every  $i = 1, 2, \dots, n$ . Hereinafter we will state similar argument as "strip argument" for short.

Together with the fact (2.4) and (2.6), we see that

$$\lim_{|x| \to +\infty} D^2 u(x) = \frac{1}{2b} (I - \widetilde{A})^{-1} - (a+b)I,$$

which is a bounded matrix. This limit together with the regularity assumption of extended function  $u \in C^4(\mathbb{R}^n)$  tells us

$$D^2 u(x) \le M, \quad \forall x \in \mathbb{R}^n$$

Applying formula (2.5), we see that for  $\lambda_i \in [-a + b + \delta, M]$ ,

$$\frac{\partial F_{\tau}}{\partial \lambda_i}(\lambda) = \frac{\lambda_i + a + b}{\lambda_i + a - b} \cdot \frac{2b}{\left(\lambda_i + a + b\right)^2} = \frac{2b}{\left(\lambda_i + a\right)^2 - b^2} \in [\frac{2b}{\left(M + a\right)^2 - b^2}, \frac{2b}{\left(b + \delta\right)^2 - b^2}],$$

and

$$\frac{\partial^2 F_{\tau}}{\partial \lambda_i^2}(\lambda) = -\frac{4b\left(\lambda_i + a\right)}{\left[\left(\lambda_i + a\right)^2 - b^2\right]^2} < 0.$$

Thus  $F_{\tau}$  is uniformly elliptic and concave, then Theorem 2.1 gives us the result.

## 2.2 $\tau = \frac{\pi}{4}$ Situation

Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) with  $\tau = \frac{\pi}{4}$  and satisfy (1.7). Let

$$\overline{U}(x) := u(x) + \frac{1}{2}|x|^2.$$

We can extend the function u sufficiently smooth to  $\mathbb{R}^n$  such that  $D^2 u > -I$  in  $\mathbb{R}^n$  and hence  $D^2 \overline{U} > 0$  in  $\mathbb{R}^n$ .

Let  $(\tilde{x}, V)$  be the Legendre transform of  $(x, \overline{U})$  i.e.

$$\begin{cases} \widetilde{x} := D\overline{U}(x) = Du(x) + x \\ D_{\widetilde{x}}V(\widetilde{x}) := x \end{cases},$$
(2.8)

and we have

$$D_{\widetilde{x}}^2 V(\widetilde{x}) = (D^2 \overline{U}(x))^{-1}.$$

Then we see that  $V(\tilde{x})$  satisfies:

$$-\Delta_{\widetilde{x}}V = \frac{\sqrt{2}C_0}{2}, \quad \text{in} \quad \mathbb{R}^n \setminus \overline{\Omega}, \tag{2.9}$$

where  $\Omega = D\overline{U}(B_1) = Du(B_1) + B_1$ .

*Proof of Theorem 1.3.* By the Legendre transform (2.8), the original equation is changed into formula (2.9). Since  $D^2 u > -I$ , hence from the equation (1.4) we obtain that for all  $i = 1, 2, \dots, n$ 

$$\frac{\sqrt{2}}{\lambda_i(D^2u)+1} = \sum_{j \neq i} \frac{-\sqrt{2}}{\lambda_j(D^2u)+1} - C_0 < -C_0.$$

Hence

$$D_{\tilde{x}}^2 V(\tilde{x}) = (D^2 u(x) + I)^{-1} \le \frac{-C_0}{\sqrt{2}} I.$$
(2.10)

Thus  $D_{\widetilde{x}}^2 V(\widetilde{x})$  is positive and bounded, applying Theorem 2.1 and we learned that the limit of  $D_{\widetilde{x}}^2 V(\widetilde{x})$  exists as  $|\widetilde{x}| \to \infty$ .

As the argument in the proof of Theorem 1.2, strip argument (2.7) tells us the limit of  $D^2u(x)$ as  $|x| \to \infty$  also exists, hence  $D^2u$  is bounded from above. Hence  $F_{\tau}$  is uniformly elliptic and concave with respect to the set of solutions and Theorem 2.1 gives us the result.

# **2.3** $\frac{\pi}{4} < \tau < \frac{\pi}{2}$ Situation

In this part, we focus on equation (1.4) with  $\frac{\pi}{4} < \tau < \frac{\pi}{2}$ . We prove Theorem 1.4 by using the following important identity formula, which is proved by taking derivatives. (See for example [38, 20]). When  $\lambda_i > -a - b$ ,  $\forall i = 1, 2, \dots, n$ , then

$$\sum_{i=1}^{n} \arctan \frac{\lambda_i + a - b}{\lambda_i + a + b} = \sum_{i=1}^{n} \arctan \left(\frac{\lambda_i + a}{b}\right) - \frac{n\pi}{4}.$$
(2.11)

First, we introduce the following result on asymptotic behavior of classical solutions of special Lagrangian equation (1.1) with  $\tau = \frac{\pi}{2}$ . See Theorem 1.1 of [26].

**Theorem 2.2.** Let u be a smooth solution of (1.4) with  $\tau = \frac{\pi}{2}$ , and we assume that  $|C_0| > (n-2)\pi/2$ . Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}_{\tau}(C_0, -\infty)$  such that (1.6) holds.

Applying Theorem 2.2 and the identity formula (2.11), we can immediately obtain the second part of Theorem 1.4.

**Proposition 2.3.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) with  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$  and satisfy condition (1.3). Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(C_0, -(a+b))$  such that (1.6) holds.

Proof. Setting

$$v(x) = \frac{u(x)}{b} + \frac{a}{2b}|x|^2,$$
(2.12)

since  $u \in C^4$  satisfies equation (1.4) and condition (1.3), by (2.11) we obtain

$$\sum_{i=1}^{n} \arctan \frac{\lambda_i(D^2 u) + a}{b} = \frac{b}{\sqrt{a^2 + 1}} (C_0 + \frac{n}{4} \frac{\sqrt{a^2 + 1}}{b} \pi),$$

i.e.

$$\sum_{i=1}^{n} \arctan \lambda_i (D^2 v) = \frac{b}{\sqrt{a^2 + 1}} C_0 + \frac{n}{4} \pi.$$
(2.13)

Condition (1.3) together with Theorem 2.2 tells us there exist  $\overline{\gamma} \in \mathbb{R}, \ \overline{\beta} \in \mathbb{R}^n$ , and  $\overline{A} \in \mathcal{A}_{\frac{\pi}{2}}(\frac{b}{\sqrt{a^2+1}}C_0 + \frac{n}{4}\pi, -\infty)$  such that (1.6) holds.

Hence the result follows from the definition of v(x).

Moreover, if we assume that  $D^2 u \ge -(a - \varepsilon_0)I$  for some  $\varepsilon_0 > 0$ , then we can similarly obtain a corresponding asymptotic behavior result by Theorem 2.1 here. The strategy here doesn't work for  $D^2 u \ge -aI$  situation.

**Theorem 2.4.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) with  $\tau = \frac{\pi}{2}$  and satisfy  $D^2 u \geq \varepsilon_0 I$ . Then there exist  $\gamma \in \mathbb{R}, \beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}_{\frac{\pi}{2}}(C_0, \varepsilon_0)$  such that (1.6) holds.

*Proof.* Extend u sufficiently smooth to  $\mathbb{R}^n$  such that  $D^2 u > \frac{\varepsilon_0}{2}I$  in  $\mathbb{R}^n$ . Let  $(\tilde{x}, \tilde{u})$  be the Legendre transform of (x, u) i.e.

$$\begin{cases} \widetilde{x} := Du(x) \\ D_{\widetilde{x}}\widetilde{u}(\widetilde{x}) := x \end{cases},$$

and we have

$$D_{\bar{x}}^2 \widetilde{u}(\tilde{x}) = \left( D^2 u(x) \right)^{-1}.$$

Hence

$$\widetilde{\lambda}_i(D^2\widetilde{u}) = \frac{1}{\lambda_i} \in (0, \frac{2}{\varepsilon_0}).$$

Also, due to  $D^2 u > \frac{\varepsilon_0}{2} I$  in  $\mathbb{R}^n$ , we also have the following relationship for all  $x^1, x^2 \in \mathbb{R}^n$ 

$$\begin{aligned} |\widetilde{x}^2 - \widetilde{x}^1|^2 &= |Du(x^2) - Du(x^1)|^2 \\ &\geq \frac{\varepsilon_0^2}{4} |x^2 - x^1|^2. \end{aligned}$$
(2.14)

Thus  $\widetilde{u}$  satisfies

$$G(\widetilde{\lambda}_i) := \sum_{i=1}^n \arctan\left(\frac{1}{\widetilde{\lambda}_i}\right) = C_0, \quad \text{in} \quad \mathbb{R}^n \setminus \overline{\Omega},$$

where  $\Omega = Du(B_1)$  is a bounded domain. By taking derivatives, we see that

$$\begin{split} \frac{\partial}{\partial \widetilde{\lambda}_i} G(\widetilde{\lambda}_i) &= \frac{1}{1 + (\frac{1}{\widetilde{\lambda}_i})^2} \cdot (-\frac{1}{\widetilde{\lambda}_i^2}) = -\frac{1}{1 + \widetilde{\lambda}_i^2} \in (-1, -\frac{\varepsilon_0^2}{4 + \varepsilon_0^2}),\\ &\frac{\partial^2}{\partial \widetilde{\lambda}_i^2} \left( G(\widetilde{\lambda}_i) \right) = \frac{2\widetilde{\lambda}_i}{\left(1 + \widetilde{\lambda}_i^2\right)^2} > 0. \end{split}$$

Hence the equation satisfied by  $\tilde{u}$  is uniformly elliptic and convex, applying Theorem 2.1 we see that there exists a  $\tilde{A} \in \text{Sym}(n)$  satisfying  $\sum_{i=1}^{n} \arctan\left(\frac{1}{\lambda_i(\tilde{A})}\right) = C_0$  such that

$$\lim_{|\widetilde{x}| \to +\infty} D^2 \widetilde{u}(\widetilde{x}) = \widetilde{A}.$$

By strip argument as (2.7), we see that  $\widetilde{A}$  is invertible. Hence from the definition of Legendre transform and (2.14) tells us  $|\widetilde{x}| \to \infty$ , as  $|x| \to \infty$ , we see that the limit of  $D^2u(x)$  as  $|x| \to \infty$  exists as well. Hence  $D^2u$  is bounded on  $\mathbb{R}^n$ . Now we verify that the equation (1.4) with  $\tau = \frac{\pi}{2}$  under these conditions is also uniformly elliptic and concave. By taking derivatives, we have

$$\frac{\partial}{\partial \lambda_i} \left( \sum_{i=1}^n \arctan \lambda_i \right) = \frac{1}{1 + \lambda_i^2},$$
$$\frac{\partial^2}{\partial \lambda_i^2} \left( \sum_{i=1}^n \arctan \lambda_i \right) = -\frac{2\lambda_i}{(1 + \lambda_i^2)^2} < 0.$$

Due to  $D^2u$  is bounded and non-negative, hence the operator is uniformly elliptic and concave.

Apply Theorem 2.1 and the result follows immediately.

As in the proof of Proposition 2.3, we can similarly obtain the following result by using identity formula (2.11). **Proposition 2.5.** Let  $u \in C^4(\mathbb{R}^n \setminus \overline{B_1})$  be a classical solution of (1.4) with  $\tau \in (\frac{\pi}{4}, \frac{\pi}{2})$ , satisfying condition (1.8). Then there exist  $\gamma \in \mathbb{R}$ ,  $\beta \in \mathbb{R}^n$  and  $A \in \mathcal{A}(C_0, -a + \varepsilon_0)$  such that (1.6) holds.

*Proof.* Let v be the function defined as in (2.12), then formula (2.11) tells us v satisfies equation (2.13). Since  $D^2 u \ge -(a - \varepsilon_0)I$ , we have  $D^2 v \ge \varepsilon_0 I$ . 

Apply Theorem 2.4, the result follows immediately.

Theorem 1.4 is the combination of Proposition 2.3 and Proposition 2.5.

#### **Convergence of Hessian at infinity** 3

In this section, we study the asymptotic behavior at infinity of Hessian matrix of classical solution of (1.9) with a fixed  $\tau \in (0, \frac{\pi}{4})$ .

Some basic and important results on Monge-Ampère equation (1.18) are important in our proof. The following lemma can be found in the proof of Theorem 1.2 in [3], which is based on the argument by Caffarelli-Li [5]. In the following results on Monge-Ampère equation (1.18) i.e. Lemma 3.1, Corollary 3.2, Theorem 3.3 and Remark 3.4, we assume  $f(\infty) = 1$  for simplicity.

**Lemma 3.1.** Let  $u \in C^0(\mathbb{R}^n)$  be a convex viscosity solution of (1.18) with  $u(0) = \min_{\mathbb{R}^n} u = 0$ , where  $0 < f \in C^0(\mathbb{R}^n)$  and

$$f^{\frac{1}{n}} - 1 \in L^n(\mathbb{R}^n)$$

Then there exists a linear transform T satisfying det T = 1 such that  $v := u \circ T$  satisfies

$$\left|v - \frac{1}{2}|x|^2\right| \le C_1 |x|^{2-\varepsilon}, \quad \forall |x| \ge R_0.$$
(3.1)

for some  $C_1 > 0$ ,  $\varepsilon > 0$  and  $R_0 \gg 1$ .

**Corollary 3.2.** Let u satisfy the same conditions as in Lemma 3.1,  $0 < f \in C^0(\mathbb{R}^n)$  and  $\exists \beta > 1$ such that

$$\limsup_{|x| \to \infty} |x|^{\beta} |f(x) - 1| < \infty$$

Then the same result in Lemma 3.1 holds.

*Proof.* Note that for a, b near 1 (bounded and away from origin), we have the following inequality

$$a^{\frac{1}{n}} - b^{\frac{1}{n}} \le C \cdot (a - b),$$

due to the derivative of  $x^{\frac{1}{n}}$  at x = 1 is  $\frac{1}{n} < 1$ . Hence we have

$$\left(\int_{\mathbb{R}^n \setminus B_1} \left| f(z)^{\frac{1}{n}} - 1 \right|^n dz \right)^{\frac{1}{n}} \le C \cdot \left( \int_{\mathbb{R}^n \setminus B_1} \left| f(z) - 1 \right|^n dz \right)^{\frac{1}{n}},$$

where  $C = C(\inf f, \sup f, n)$ . Hence we have

$$LHS \le C \cdot \left( \int_{\mathbb{R}^n \setminus B_1} |\frac{1}{|x|^{\beta}}|^n dz \right)^{\frac{1}{n}} < \infty \quad (\beta > 1).$$

Then the conditions of Lemma 3.1 are met and the results holds.

As a consequence, we have the following result on Monge-Ampère equation. The proof is similar to the one in Bao-Li-Zhang [3] and in Caffarelli-Li [5]. Since there are some differences from their proof, we provide the details here for reading simplicity.

**Theorem 3.3.** Let  $u \in C^0(\mathbb{R}^n)$  be a convex viscosity solution of (1.18),  $f \in C^{\alpha}(\mathbb{R}^n)$ ,  $\alpha \in (0, 1)$ , and satisfy

$$\limsup_{|x| \to \infty} |x|^{\beta} |f(x) - 1| < \infty, \tag{3.2}$$

$$[f]_{C^{\alpha}(B_{\frac{3}{2}R} \setminus B_{\frac{R}{2}})} \cdot R^{\alpha + \gamma} \le C, \quad \forall R > 1,$$

$$(3.3)$$

for some  $\beta > 1$ ,  $\gamma > 0$ . Then there exist  $A \in \mathcal{A}_0(0,0)$ ,  $C(n, R_0, \alpha, \beta, \gamma, c_0)$ , such that

$$|D^2 u(x) - A| \le \frac{C}{|x|^{\min\{\beta,\varepsilon,\gamma\}}}, \quad \forall |x| \ge R_1,$$
(3.4)

and

$$||D^2u||_{C^{\alpha}(\mathbb{R}^n)} \le C, \tag{3.5}$$

where  $\varepsilon$ , and  $R_0$  are positive constants come from Corollary 3.2 (or say, Lemma 3.1) and  $R_1 > R_0$  depends only on  $n, R_0, \varepsilon, \beta, c_0$ .

*Proof.* By Corollary 3.2, we see that there exist a linear transform T and  $\varepsilon > 0$ ,  $C_1 > 0$ ,  $R_0 \gg 1$  such that  $v := u \circ T$  satisfies (3.1)

Now in order to obtain pointwise decay speed estimate at infinity, we focus on sufficiently far ball, scale back to unit size, and then apply interior regular estimate. Set

$$w(x) := v(x) - \frac{1}{2}|x|^2.$$

For  $|x| = R > 2R_0$ , let

$$v_R(y) = \left(\frac{4}{R}\right)^2 v\left(x + \frac{R}{4}y\right), \text{ and } w_R(y) = \left(\frac{4}{R}\right)^2 w\left(x + \frac{R}{4}y\right) \quad |y| \le 2.$$

By (3.1), we have

$$||v_R||_{L^{\infty}(B_2)} \le C, ||w_R||_{L^{\infty}(B_2)} \le CR^{-\varepsilon},$$

for some constants C uniform to  $R \ge R_0$  and

$$v_R(y) - \left(\frac{1}{2}|y|^2 + \frac{4}{R}x \cdot y + \frac{8}{R^2}|x|^2\right) = O\left(R^{-\varepsilon}\right), \quad \forall y \in B_2.$$

In order to apply interior estimate, we set

$$\overline{v}_R(y) := v_R(y) - \frac{4}{R}x \cdot y - \frac{8}{R^2}|x|^2.$$

If  $R > R_1$  with  $R_1$  sufficiently large, the set

$$\Omega_R := \{ y \in B_2; \quad \overline{v}_R(y) < 1 \}$$

is between  $B_{1,2}$  and  $B_{1,8}$ . The equation satisfied by  $\overline{v}_R$  is

$$\det\left(D^2\overline{v}_R(y)\right) = f\left(x + \frac{R}{4}y\right) =: f_R(y), \quad \text{in } B_2.$$

By definition of  $f_R$ , the assumptions (3.2) and (3.3), we have

$$||f_R - 1||_{C^0(\overline{B_2})} \le CR^{-\beta}, \quad \forall R > 1.$$

$$[f_R]_{C^{\alpha}(\overline{B_2})} = \max_{\substack{y_1, y_2 \in B_2 \\ y_1, y_2 \in B_2 \\ z_1, z_2 \in B_{\frac{3}{2}R}(x) \setminus B_{\frac{R}{2}}(x)}} \frac{|f(x + \frac{R}{4}y_1) - f(x + \frac{R}{4}y_2)|}{|y_1 - y_2|^{\alpha}} \\ = \max_{\substack{z_1, z_2 \in B_{\frac{3}{2}R}(x) \setminus B_{\frac{R}{2}}(x)}} \frac{|f(z_1) - f(z_2)|}{|z_1 - z_2|^{\alpha}} \cdot (\frac{R}{4})^{\alpha} \\ \leq CR^{-\gamma}.$$

Combine these two parts we have

$$||f_R - 1||_{C^{\alpha}(\overline{B_2})} \le CR^{-\min\{\beta,\gamma\}}.$$

Applying the interior estimate by Caffarelli [8], Jian and Wang [22] on  $\Omega_{1,v}$ , we have

$$\|D^2 v_R\|_{C^{\alpha}(B_{1,1})} = \|D^2 \overline{v}_R\|_{C^{\alpha}(B_{1,1})} \le C,$$
(3.6)

and hence

$$\frac{I}{C} \le D^2 v_R \le CI \quad \text{on } B_{1.1},\tag{3.7}$$

for some C independent of R. This tells us the result (3.5). More explicitly, by the definition of  $v_R$ , we see that  $D^2v_R(y) = D^2v(x + \frac{R}{4}y)$  are bounded uniform to  $x \in B_{R_1}^c$ , hence

$$||D^{2}v||_{C^{0}(B_{R_{1}}^{c})} \leq \sup_{x \in B_{R_{1}}^{c}} ||D^{2}v||_{C^{0}(B_{\frac{|x|}{4}}(x))} \leq C.$$

Since  $f \in C^{\alpha}(B_{2R_1})$ , the interior  $C^{2,\alpha}$  estimate by Caffarelli [8], Jian and Wang [22] tells us

$$||D^2v||_{C^{\alpha}(B_{2R_1})} \le C.$$
(3.8)

This provides us the boundedness of Hessian matrix  $D^2v$  and hence so is  $D^2u$ . Similarly, for Hölder semi-norm, by definition we have

$$\begin{split} [D^2 v]_{C^{\alpha}(\mathbb{R}^n)} &= \sup_{\substack{x_1, x_2 \in \mathbb{R}^n}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}} \\ &\leq \max \left\{ \sup_{\substack{x_1, x_2 \in B_{2R_1} \\ x_1, x_2 \in B_{2R_1}}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \lim_{x_1 \in B_{R_1}^c} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}}, \lim_{x_1 \in B_{R_1}^c} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}} \right\}.$$

The first term bounded due to  $C^{2,\alpha}$  estimate in (3.8). For the second term, note that

$$\sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}} \leq \sup_{x \in B_{R_1}^c} [D^2 v]_{C^{\alpha}(B_{\frac{1}{4}|x|}(x))} \\
= \sup_{x \in B_{R_1}^c} \sup_{y_1, y_2 \in B_{\frac{1}{4}|x|}(x)} \frac{|D^2 v(y_1) - D^2 v(y_2)|}{|y_1 - y_2|^{\alpha}} \\
= \sup_{x \in B_{R_1}^c} \sup_{z_1, z_2 \in B_1} \sup_{\frac{|D^2 v(x + \frac{|x|}{4}z_1) - D^2 v(x + \frac{|x|}{4}z_2)|}{(\frac{|x|}{4})^{\alpha} \cdot |z_1 - z_2|^{\alpha}} \\
\leq \sup_{\substack{x \in B_{R_1}^c \\ R = |x|}} [D^2 v_R]_{C^{\alpha}(B_1)} \cdot (\frac{4}{R})^{\alpha} \\
\leq C,$$

for some constant C from (3.6). For the third term, due to Hessian matrix  $D^2v$  is proved to be bounded, hence

$$\sup_{\substack{x_1 \in B_{R_1}^c \\ x_2 \in B_{\frac{1}{4}|x_1|}(x_1)^c}} \frac{|D^2 v(x_1) - D^2 v(x_2)|}{|x_1 - x_2|^{\alpha}} \le (\frac{4}{3R_1})^{\alpha} \cdot 2||D^2 v||_{C^0(\mathbb{R}^n)} \le C.$$

Combine these three parts and since the linear transform T from Lemma 3.1 doesn't degenerate, we obtain (3.5) immediately.

Using Newton-Leibnitz formula between det  $D^2 \overline{v}_R(y) = f_{1,R}(y)$  and det I = 1 gives

$$\widetilde{a}_{ij}\partial_{ij}w_R = f_{1,R}(y) - 1 = O\left(R^{-\min\{\beta,\gamma\}}\right),\,$$

where  $\widetilde{a}_{ij}(y) = \int_0^1 \operatorname{cof}_{ij} \left( I + tD^2 w_R(y) \right) dt$ . By (3.6), (3.7) and using Landau-Kolmogorov interpolation inequality, since  $v_R$  is bounded on  $B_2$ , there exists some constant C independent of R such that  $||v_R||_{C^{2,\alpha}(\bar{B}_1)} \leq C$ ., the  $C^{\alpha}$  norm of  $D^2 w_R$  is also bounded by some constant C independent of R, hence

$$\frac{I}{C} \leq \widetilde{a}_{ij} \leq CI, \text{ on } B_{1.1}, \quad \|\widetilde{a}_{ij}\|_{C^{\alpha}\left(\overline{B}_{1.1}\right)} \leq C.$$

Thus Schauder's estimate (note that Theorem 6.2 of [18] demands the coefficients have bounded Hölder norm, then the frozen coefficient method can be applied) gives us

$$||w_R||_{C^{2,\alpha}(B_1)} \le C\left(||w_R||_{L^{\infty}(\overline{B}_{1,1})} + ||f_{1,R} - 1||_{C^{\alpha}(\overline{B}_1)}\right) \le CR^{-\min\{\varepsilon,\beta,\gamma\}}.$$

Scale back to the original  $B_{\frac{|x|}{4}}(x)$  ball, this is exactly our result (3.4).

**Remark 3.4.** The condition (3.3) can be replaced by the following

$$\exists \gamma > 0 \text{ such that } \limsup_{|x| \to \infty} |x|^{1+\gamma} |Df(x)| < \infty.$$
(3.9)

Condition (3.9) demands f has at least one order derivative while condition (3.3) only demands the  $C^{\alpha}$ -semi norm of f has a vanishing speed at infinity.

For example, similar to the example in Remark 1.7, we take  $f(x) := e^{-|x|} \sin(e^{|x|}) + 1$ . On the one hand, Df(x) doesn't admit a limit at infinity, hence f doesn't satisfies condition (3.9).

On the other hand, we calculate its  $C^{\alpha}$  Hölder semi-norm directly

$$\begin{split} \left[f\right]_{C^{\alpha}\left(B_{\frac{3}{2}R}\setminus B_{\frac{R}{2}}\right)} &= \sup_{z_{1},z_{2}\in B_{\frac{3R}{2}}\setminus B_{\frac{R}{2}}} \frac{|f\left(z_{1}\right)-f\left(z_{2}\right)|}{|z_{1}-z_{2}|^{\alpha}} \\ &\leq \sup_{z_{1},z_{2}\in B_{\frac{3R}{2}}\setminus B_{\frac{R}{2}}} \left(e^{-|z_{2}|}\frac{|\sin(e^{|z_{1}|})-\sin(e^{|z_{2}|})|}{|z_{1}-z_{2}|^{\alpha}}+\sin(e^{|z_{1}|})\frac{|e^{-|z_{1}|}-e^{-|z_{2}|}|}{|z_{1}-z_{2}|^{\alpha}}\right) \\ &\leq C \sup_{z_{1},z_{2}\in B_{\frac{3R}{2}}\setminus B_{\frac{R}{2}}} \left(e^{-R}\cdot\frac{\max_{z\in B_{\frac{3R}{2}}\setminus B_{\frac{R}{2}}}|\cos(e^{|z|})|\cdot|z_{1}-z_{2}|}{|z_{1}-z_{2}|^{\alpha}}\right) \\ &+ 1\cdot\frac{\sup_{z\in B_{\frac{3R}{2}}\setminus B_{\frac{R}{2}}}|e^{-|z|}|\cdot|z_{1}-z_{2}|}{|z_{1}-z_{2}|^{\alpha}}}{|z_{1}-z_{2}|^{\alpha}}\right) \\ &\leq Ce^{-R}\cdot R^{1-\alpha}, \end{split}$$

for some constant C independent of R. Hence f satisfies condition (3.3) for all  $\alpha \in (0, 1), \gamma > 0$ .

The study of equation (1.9) is transformed into Monge-Ampère type by taking Legendre transform as formula (2.1). We can easily verify as in Section 2 that  $\tilde{u}(\tilde{x})$  satisfies the following Monge-Ampère type equation

$$\sum_{i=1}^{n} \ln \widetilde{\lambda}_{i} = \frac{2b}{\sqrt{a^{2}+1}} f(\frac{1}{2b}(\widetilde{x} - D\widetilde{u}(\widetilde{x}))).$$

This equation is equivalent (under the condition of  $\widetilde{\lambda_i} > 0$ ) to

$$\det D^2 \widetilde{u} = \exp\left\{\frac{2b}{\sqrt{a^2 + 1}} f\left(\frac{1}{2b}(\widetilde{x} - D\widetilde{u}(\widetilde{x}))\right)\right\} =: g(\widetilde{x}).$$
(3.10)

**Theorem 3.5.** Let  $u \in C^4(\mathbb{R}^n)$  be a classical solution of (1.9) and satisfy (1.10). Also, we assume that f satisfies condition (3.2), (3.9) for some  $f(\infty) < 0$ . Then there exist  $A \in \mathcal{A}(f(\infty), -a + b), \alpha > 0, C(n, f, \beta, \gamma)$ , and  $R_2(n, f, \beta, \gamma)$  such that

$$\left|D^{2}u(x) - A\right| \le \frac{C}{|x|^{\alpha}}, \quad \forall |x| \ge R_{2}.$$
(3.11)

*Proof.* By condition (1.10),  $-a+b < \lambda_i \leq M$  for  $i = 1, 2, \dots, n$ , hence for some  $\delta = \delta(M) > 0$ ,

$$\widetilde{\lambda}_i = 1 - \frac{2b}{\lambda_i + a + b} \quad \text{satisfies} \quad 0 < \widetilde{\lambda}_i < 1 - \delta.$$
(3.12)

Also note that  $f \in C^0(\mathbb{R}^n)$  has a finite lower bound. Due to  $\ln \lambda_i < 0$  holds for all  $i = 1, 2, \dots, n$ . Hence similar to the strategy used in [37], we naturally have a lower bound such that for all  $i = 1, 2, \dots, n$ ,

$$\ln \widetilde{\lambda_i} > \frac{2b}{\sqrt{a^2 + 1}} \inf_{\mathbb{R}^n} f > -\infty, \quad \text{i.e.} \quad \widetilde{\lambda_i} > \exp\{\frac{2b}{\sqrt{a^2 + 1}} \inf_{\mathbb{R}^n} f\} > 0.$$

Combine these two results, we have

$$0 < \delta < \widetilde{\lambda_i} < 1 - \delta,$$

for some  $\delta = \delta(a, b, \inf_{\mathbb{R}^n} f, M)$ . Furthermore, we have a reversed direction of formula (2.4). By the definition of Legendre transform (2.1), we have

$$2b|x| = |\widetilde{x} - D\widetilde{u}(\widetilde{x})| \ge |\widetilde{x}| - |D\widetilde{u}(\widetilde{x})| \ge |\widetilde{x}| - (1 - \delta)|\widetilde{x}| - |D\widetilde{u}(0)|,$$

i.e.

$$|x| \ge \frac{\delta}{2b} |\widetilde{x}| - \frac{1}{2b} |D\widetilde{u}(0)|. \tag{3.13}$$

Combine formula (2.4) and (3.13), we see that there exists some constant  $C_0 = C_0(\inf_{\mathbb{R}^n} f, a, b, M)$ such that

$$\frac{1}{C_0}|x| \le |\tilde{x}| \le C_0|x|, \quad \text{for} \quad |x| \gg 1.$$
(3.14)

This linear growth equivalence result (3.14) is the major reason that we demand the boundedness of Hessian.

We obtain the limit of Hessian for equation (3.10) by Theorem 3.3 and Remark 3.4 first. Thus we need to verify the asymptotic behavior of  $q(\tilde{x})$ .

Step 1 Verify  $q(\tilde{x})$  satisfies condition (3.2)

By the equivalence (3.14),

$$\lim_{\widetilde{x}\to\infty}g(\widetilde{x}) = \exp\{\frac{2b}{\sqrt{a^2+1}}f(\infty)\} =: g(\infty) \in (0,1).$$

Hence we have

$$|\widetilde{x}|^{\beta}|g(\widetilde{x}) - g(\infty)| = e^{\frac{2b}{\sqrt{a^2 + 1}}} \frac{e^{f(\infty)}|\widetilde{x}|^{\beta}}{\left|\frac{\widetilde{x} - D\widetilde{u}(\widetilde{x})}{2b}\right|^{\beta}} \cdot \left|\frac{\widetilde{x} - D\widetilde{u}(\widetilde{x})}{2b}\right|^{\beta} \cdot \left|e^{f(\frac{\widetilde{x} - D\widetilde{u}(\widetilde{x})}{2b}) - f(\infty)} - 1\right|.$$

Note that f is a bounded function and admits a limit  $f(\infty)$  at infinity, together with formula (3.14) tell us  $f(\frac{\tilde{x}-D\tilde{u}(\tilde{x})}{2b}) - f(\infty)$  is less than 1 for sufficiently large  $|\tilde{x}|$ . Due to  $|e^t - 1| \le e|t|$  as long as  $|t| \le 1$ , by  $\tilde{x} - D\tilde{u}(\tilde{x}) = 2bx$  and equivalence (3.14), we

have

$$\limsup_{|\widetilde{x}| \to \infty} |\widetilde{x}|^{\beta} |g(\widetilde{x}) - g(\infty)| \le C_0^{\beta} e^{\frac{2b}{\sqrt{a^2 + 1}} + f(\infty) + 1} \limsup_{|x| \to \infty} |x|^{\beta} |f(x) - f(\infty)| < \infty.$$

Step 2 Verify  $g(\tilde{x})$  satisfies condition (3.9) By taking derivative once, we have

$$Dg(\widetilde{x}) = \exp\{\frac{2b}{\sqrt{a^2+1}}f\} \cdot Df(\frac{1}{2b}(\widetilde{x} - D\widetilde{u}(\widetilde{x}))) \cdot \frac{1}{2b}[I - D^2\widetilde{u}].$$

Due to  $D^2 \widetilde{u}$  and  $\exp\{\frac{2b}{\sqrt{a^2+1}}f\}$  are bounded (for sufficiently large  $\widetilde{x}$ ), hence we only need to consider the rest part of  $Dg(\tilde{x})$ .

By equivalence (3.14) we obtain

$$\left|\widetilde{x}\right|^{\gamma+1} \left| Df(\frac{1}{2b}(\widetilde{x} - D\widetilde{u}(\widetilde{x}))) \right| \le C_0^{\gamma+1} |x|^{\gamma+1} \cdot |Df(x)|.$$

Take the limit  $|\tilde{x}| \to \infty$  and condition (3.9) gives us

$$\limsup_{|\widetilde{x}|\to\infty} |\widetilde{x}|^{\gamma+1} \left| |Df(\frac{1}{2b}(\widetilde{x} - D\widetilde{u}(\widetilde{x}))) \right| \le C_0^{\gamma+1} \limsup_{|x|\to\infty} |x|^{\gamma+1} \cdot |Df(x)| < \infty.$$

By Theorem 3.3 and Remark 3.4, we have

$$\left| D^2 \widetilde{u}(\widetilde{x}) - \widetilde{A} \right| \le \frac{C}{|\widetilde{x}|^{\min\{\beta,\varepsilon,\gamma\}}}, \quad \forall |\widetilde{x}| \ge R_1$$

for some  $\widetilde{A} \in \widetilde{\mathcal{A}} := \left\{ \widetilde{A} \in \operatorname{Sym}(n) : \det \widetilde{A} = g(\infty), \ \widetilde{A} > 0 \right\}$  and  $\varepsilon > 0, C(n, f, \beta, \gamma), R_1(n, f, \beta, \gamma)$ . Now we prove that from the definition of Legendre transform (2.1) and the equivalence (3.14),

there exists an  $A \in \mathcal{A}(f(\infty), -a+b)$  such that (3.11) holds with  $\alpha = \min\{\beta, \varepsilon, \gamma\} > 0$ .

By strip argument as in the proof of Theorem 1.2,  $D^2 \tilde{u}$ , A are away from the origin and identity matrix (by a positive distance  $\delta$ ). Take

$$A := \left(\frac{1}{2b}\left(I - \widetilde{A}\right)\right)^{-1} - (a+b)I,$$

which satisfies  $F_{\tau}(A) = f(\infty)$ , we obtain

$$|D^{2}u - A| = 2b \left| (I - D^{2}\widetilde{u}(\widetilde{x}))^{-1} - (I - \widetilde{A})^{-1} \right|$$
  
$$\leq C_{\delta} |D^{2}\widetilde{u}(\widetilde{x}) - \widetilde{A}|$$
  
$$\leq \frac{C}{|\widetilde{x}|^{\min\{\beta,\varepsilon,\gamma\}}}.$$

By the equivalence (3.14), for the same  $\alpha > 0$  as above, we have formula (3.11).

Moreover, we have not only the limit of Hessian at infinity, but also the  $C^{\alpha}$  bound of Hessian.

**Theorem 3.6.** Under the conditions as in Theorem 3.5, there exists  $C = C(n, f, \beta, \gamma, \alpha, a, b, M)$  such that

$$||D^2u||_{C^{\alpha}(\mathbb{R}^n)} \le C.$$

*Proof.* Again, we apply Legendre transform (2.1) as in Theorem 3.5 to obtain (3.10). By Theorem 3.3, there exists some constant C relying on  $\tilde{u}$  such that

$$||D_{\widetilde{x}}^2\widetilde{u}||_{C^{\alpha}(\mathbb{R}^n)} \le C.$$

Now we transform back this result to  $D^2u$ . From formula (2.1) it follows that

$$D^{2}u(x) = \left(\frac{1}{2b}\left(I - D_{\widetilde{x}}^{2}\widetilde{u}(\widetilde{x})\right)\right)^{-1} - (a+b)I$$

and hence for any  $x, y \in \mathbb{R}^n$ ,

$$D^{2}u(x) - D^{2}u(y) = 2b \left| \left( I - D_{\widetilde{x}}^{2}\widetilde{u}(\widetilde{x}) \right)^{-1} - \left( I - D_{\widetilde{x}}^{2}\widetilde{u}(\widetilde{y}) \right)^{-1} \right|.$$

As in the argument in Theorem 3.5, there exists some  $0 < \delta$  such that

$$0 < \delta I \le D_{\widetilde{x}}^2 \widetilde{u}(\widetilde{x}) \le (1 - \delta)I$$

Thus it follows that  $\exists C_1 = C_1(n, \delta) > 0$ ,  $C_2 = C_2(n, \delta) > 0$  such that

$$2b \cdot C_1(\delta) \cdot \left| D_{\widetilde{x}}^2 \widetilde{u}(\widetilde{x}) - D_{\widetilde{x}}^2 \widetilde{u}(\widetilde{y}) \right| \ge \left| D^2 u(x) - D^2 u(y) \right| \ge 2b \cdot C_2(\delta) \cdot \left| D_{\widetilde{x}}^2 \widetilde{u}(\widetilde{x}) - D_{\widetilde{x}}^2 \widetilde{u}(\widetilde{y}) \right|$$
(3.15)

Combine formula (3.15) and the equivalence (3.14), we see that  $D^2u$  has bounded  $C^{\alpha}$  seminorm if and only if  $D^2\tilde{u}$  has bounded  $C^{\alpha}$  seminorm.

Due to the important normalization lemma of John-Cordoba and Gallegos (see [13]) cannot be applied to classical special Lagrangian equation without changing the operator, the level set method developed in [5] cannot be applied easily to  $\tau > \frac{\pi}{4}$  situation.

### 4 Analysis of Linearized Equation

Now that we have obtained the limit of Hessian at infinity and it converge with a Hölder decay speed, it is time to capture the linear and constant part of the solution. In order to do this, we follow the line of Li-Li-Yuan [26] and analyze the linearized equation of (1.9).

By the characterise result of ellipticity and concavity structure of  $F(\lambda(D^2u))$  type equation in [6], equation (1.9) is uniformly elliptic and concave. For any direction  $e \in \partial B_1$ , we apply  $\partial_e$ ,  $\partial_{ee}$  to the equation  $F_{\tau}(\lambda) = f(x)$ , then we have,

$$D_{M_{ij}}F_{\tau}\left(D^{2}u\right)D_{ij}\left(u_{e}\right) = f_{e}(x), \quad \text{and} \quad D_{M_{ij}}F_{\tau}\left(D^{2}u\right)D_{ij}\left(u_{ee}\right) \ge f_{ee}(x), \tag{4.1}$$

where  $D_{M_{ij}}F_{\tau}(D^2u)$  stands for the value of partial derivative of  $F_{\tau}(M)$  with respect to the *i*-line, *j*-column position of M at  $D^2u(x)$ .

The major difference between the linearized equation we obtained here and the one in Li-Li-Yuan [26] is that our equation is non-homogeneous. We take advantage of the linearity and use Green's function to transform the non-homogeneous equation into homogeneous equation. Then by studying the decay speed of solution of non-homogeneous equation and applying the theory in [26] we obtain corresponding results as in Theorem 4.3 and Corollary 4.4.

We consider the following Dirichlet Problem

$$\begin{cases} Lw := a_{ij}(x)D_{ij}w(x) = f(x), & \text{in } \mathbb{R}^n, \\ \lim_{|x| \to \infty} w(x) = 0, & \text{as } |x| \to \infty, \end{cases}$$
(4.2)

where the coefficients are  $C^2$  (which is provided by  $u \in C^4$ ), satisfy

$$||a_{ij}||_{C^{\alpha}(\mathbb{R}^n)} \le M,\tag{4.3}$$

for some  $\alpha > 0, M < \infty$ , strictly elliptic for some  $\gamma > 0$ ,

$$a_{ij}(x)\xi_i\xi_j \ge \gamma|\xi|^2, \quad \text{for all} \quad x,\xi \in \mathbb{R}^n,$$

$$(4.4)$$

and for some symmetric matrix  $a_{ij}(\infty)$ ,  $\varepsilon_0 > 0$ ,  $C < \infty$ ,

$$|a_{ij}(x) - a_{ij}(\infty)| \le C|x|^{-\varepsilon_0}.$$
(4.5)

By using the criterion in [32] together with the Theorem 2.2 of [26], we see that under these conditions, the Green's function of operator L is equivalent to the Green's function of Laplacian. More precisely, let  $G_L(x, y)$  be the Green's function centered at y, there exists constant C such that

$$C^{-1}|x-y|^{2-n} \leq G_L(x,y) \leq C|x-y|^{2-n}, \quad \forall x \neq y,$$
  
$$|\partial_{x_i}G_L(x,y)| \leq C|x-y|^{1-n}, \quad i = 1, \cdots, n, \quad \forall x \neq y,$$
  
$$\left|\partial_{x_i}\partial_{x_j}G_L(x,y)\right| \leq C|x-y|^{-n}, \quad i,j = 1, \cdots, n, \quad \forall x \neq y.$$
(4.6)

Now we study the existence result of Dirichlet problem (4.2) and study its asymptotic behavior at infinity. For the weak solution u of a linear elliptic equation  $a_{ij}(x)D_{ij}u(x) = f(x)$  in  $\mathbb{R}^n$ hereinafter, we mean that for any  $\varphi \in C_0^{\infty}(\mathbb{R}^n)$ ,  $u \in W_{loc}^{1,1}(\mathbb{R}^n)$  satisfies

$$\int_{\mathbb{R}^n} D_i(a_{ij}(x)\varphi(x))D_ju(x) + f(x)\varphi(x)\mathrm{d}x = 0.$$

For the distribution solution u of a linear elliptic equation  $a_{ij}(x)D_{ij}u(x) = f(x)$  in  $\mathbb{R}^n$  hereinafter, we mean that for any  $\varphi \in \mathcal{S}(\mathbb{R}^n)$ ,

$$\langle a_{ij}(x)D_{ij}u(x),\varphi\rangle = \langle f,\varphi\rangle$$

where

$$\langle a_{ij}(x)D_{ij}u(x),\varphi\rangle := \int_{\mathbb{R}^n} u(x)D_{ij}(a_{ij}(x)\varphi(x))\mathrm{d}x, \text{ and } \langle f,\varphi\rangle := \int_{\mathbb{R}^n} f(x)\varphi(x)\mathrm{d}x.$$

We can easily see from  $C_0^{\infty}(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n)$  that if u is a  $W_{loc}^{1,1}$  distribution solution, then it is also a weak solution.

**Lemma 4.1.** Assume in addition that  $f \in C^0(\mathbb{R}^n)$  satisfies for some  $k \ge 2, 0 < \varepsilon \ll 1$ ,

$$\limsup_{|x| \to +\infty} |x|^{k+\varepsilon} |f(x)| < \infty.$$
(4.7)

Then there exists a weak solution  $u \in C^1(\mathbb{R}^n)$  to the Dirichlet problem (4.2) and satisfies

$$|u(x)| \le C|x|^{2-k-\varepsilon},$$

for some constant C.

*Proof.* By the definition of Green's function, the equivalence result and potential theory (or Calderón-Zygmund inequality), the following convolution

$$G_L*f(x):=\int_{\mathbb{R}^n}G_L(x,y)f(y)\mathrm{d} y$$

belongs to  $W^{2,p}(\mathbb{R}^n)$  for sufficiently large  $p > \frac{n}{k}$  and is a distribution solution of L[w](x) = f(x)(See for example [1, 46]). Hence by the embedding theory, it is also a weak solution. Now we only need to verify that it vanishes with the desired speed at infinity. Following the line of [3] (Lemma 2.2, formula (2.21)), we find out that as long as f(y) satisfies (4.7), we have

$$|G_L * f(x)| \le G_L * [C|x|^{-k-\varepsilon}] \to 0$$
, as  $|x| \to \infty$ .

In fact, this is proved in a standard way by separating the integral domain into the following three part

$$E_{1} := \{ y \in \mathbb{R}^{n}, |y| \le |x|/2 \}, E_{2} := \{ y \in \mathbb{R}^{n}, |y-x| \le |x|/2 \}, E_{3} := \mathbb{R}^{n} \setminus (E_{1} \cup E_{2}).$$

Thus

$$|G_L * f(x)| \le C \int_{E_1 \cup E_2 \cup E_3} \frac{1}{|x - y|^{n-2}} \cdot \frac{1}{|y|^{k+\varepsilon}} \mathrm{d}y$$

By our choice of  $E_i$ , it follows immediately that

$$\begin{split} &\int_{E_1} \frac{1}{|x-y|^{n-2} \cdot |y|^{k+\varepsilon}} \mathrm{d} y \leq \int_{B_{\frac{|x|}{2}}} \frac{1}{|y|^{k+\varepsilon}} \mathrm{d} y \cdot \frac{1}{|\frac{|x|}{2}|^{n-2}}, \\ &= C_{n,k,\varepsilon} \cdot |x|^{n-k-\varepsilon} \cdot \frac{1}{|x|^{n-2}} = O(|x|^{2-k-\varepsilon}), \quad \text{as} \quad |x| \to \infty. \end{split}$$

Similarly, note that in  $E_2$  case,  $|y - x| \le \frac{|x|}{2} \le |y|$  we have

$$\int_{E_2} \frac{1}{|x-y|^{n-2} \cdot |y|^{k+\varepsilon}} \mathrm{d}y \le C_k \int_{|x-y| \le \frac{|x|}{2}} \frac{1}{|x-y|^{n-2+\varepsilon}} \mathrm{d}y \cdot \frac{1}{|x|^k},$$
$$\le \int_0^{\frac{|x|}{2}} \frac{1}{r^{\varepsilon-1}} \mathrm{d}r \cdot \frac{1}{|x|^k} = O(|x|^{-k-\varepsilon}), \quad \text{as} \quad |x| \to \infty.$$

Now we separate  $E_3$  into two parts

$$E_3^+ := \{ y \in E_3 : |x - y| \ge |y| \}, \ E_3^- := E_3 \setminus E_3^+.$$

Then

$$\int_{E_3^+} \frac{1}{|x-y|^{n-2} \cdot |y|^{k+\varepsilon}} \mathrm{d} y \leq \int_{|y| \geq \frac{|x|}{2}} \frac{1}{|y|^{n+k+\varepsilon}} \mathrm{d} y = O(|x|^{-k-\varepsilon}), \quad \text{as} \quad |x| \to \infty,$$

and

$$\int_{E_3^-} \frac{1}{|x-y|^{n-2} \cdot |y|^{k+\varepsilon}} \mathrm{d} y \leq \int_{|y-x| \geq \frac{|x|}{2}} \frac{1}{|y-x|^{n+k+\varepsilon}} \mathrm{d} y = O(|x|^{-k-\varepsilon}), \quad \text{as} \quad |x| \to \infty.$$

Hence we immediately obtain

$$|w(x)| = |G_L * f(x)| \le C|x|^{-k-\varepsilon}$$
, as  $|x| \to \infty$ .

Thus  $w(x) := G_L * f(x)$  solves the Dirichlet problem (4.2) with  $O(|x|^{2-k-\varepsilon})$  order vanishing at infinity.

Combine Lemma 4.1 with the exterior Liouville theorem for homogeneous equation proved by Li-Li-Yuan [26], we have the exterior asymptotic behavior theory for a class of non-homogeneous equations. For reading simplicity, we recall the Theorem 2.2 of [26] as the following.

**Theorem 4.2.** Let v be a positive classical solution of  $a_{ij}(x)D_{ij}v(x) = 0$  in  $\mathbb{R}^n \setminus \overline{B_1}$ , then there exists a constant  $v_{\infty} \ge 0$  such that

$$v(x) = v_{\infty} + o\left(|x|^{2-n+\delta}\right)$$
 as  $|x| \to \infty$ , for all  $\delta > 0$ .

Moreover, if  $a_{ij}(x)$  satisfies condition (4.5), then

$$v(x) = v_{\infty} + O\left(|x|^{2-n}\right) \quad as \quad |x| \to \infty.$$

$$(4.8)$$

Then the equivalence of Green's function gives us the following corresponding results for non-homogeneous situation.

**Theorem 4.3.** Let v be a positive classical solution of  $a_{ij}(x)D_{ij}v(x) = f(x)$  in  $\mathbb{R}^n \setminus \overline{B_1}$ , the coefficients satisfy (4.5) and  $f \in C^0(\mathbb{R}^n)$  satisfy (4.7) with k = n. Then there exists a constant  $v_{\infty} \geq 0$  such that

$$v(x) = v_{\infty} + O\left(|x|^{2-n}\right) \quad as \quad |x| \to \infty.$$

$$(4.9)$$

*Proof.* Taking auxiliary function w(x) as the solution we constructed in Lemma 4.1 satisfying system (4.2). Then under the decay speed (4.7) of f at infinity with k = n, we have that  $w = O(|x|^{2-n})$  as  $|x| \to \infty$ .

By the linearity of operator, we learned that  $\tilde{v} := v - w$  is a  $C^1$  weak solution of  $a_{ij}(x)D_{ij}\tilde{v}(x) = 0$  in exterior domain. Since the coefficients are uniformly elliptic and has bounded  $C^{\alpha}$  semi-norm, interior Schauder estimate tells us  $\tilde{v}$  is a classical solution. Also, we learned that the auxiliary function w is bounded, hence  $v \ge 0$  implies that there exists a constant lower bound to  $\tilde{v}$ .

Applying the exteirior Liouville theorem for homogeneous linear elliptic equation i.e. Theorem 4.2, we have

$$\widetilde{v}(x) = \widetilde{v}_{\infty} + O\left(|x|^{2-n}\right) \quad \text{as} \quad |x| \to \infty,$$

for some constant  $\tilde{v}_{\infty}$ . Due to  $v = \tilde{v} + w$  is non-negative, the result follows immediately.  $\Box$ 

**Corollary 4.4.** Let v be a classical solution of  $a_{ij}(x)D_{ij}v(x) = f(x)$  in  $\mathbb{R}^n \setminus \overline{B_1}$  Suppose that

$$|Dv(x)| = O(|x|^{-1}), \quad as \quad |x| \to +\infty.$$

Also, we demand that  $f \in C^0(\mathbb{R}^n)$  satisfies (4.7) with k = n. Then there exists a constant  $v_{\infty} \ge 0$  such that (4.9) holds.

*Proof.* In virtue of Theorem 4.3, we only need to prove that v is bounded from at least on one side. Again, by the equivalence of Green's function, we transform this problem into homogeneous situation and apply Corollary 2.1 of [26].Let w be the auxiliary function as in Lemma 4.1 satisfying (4.2), then

$$w(x) = \int_{\mathbb{R}^n} G_L(x, y) f(y) \mathrm{d}y, \quad \text{and} \quad \partial_i w(x) = \int_{\mathbb{R}^n} \partial_{x_i} G_L(x, y) f(y) \mathrm{d}y.$$

Applying similar analysis as in Lemma 4.1, it follows that

$$Dw(x) = O(|x|^{1-n}), \text{ as } |x| \to +\infty.$$

By triangle inequality, we set u := v - w and obtain

$$|Du(x)| \le |Dv(x)| + |Dw(x)| = O(|x|^{-1}), \text{ as } |x| \to +\infty.$$

Now we prove that u is bounded from one side, then the asymptotic behavior of w tells us v is also bounded from one side.

Argue by contradiction, if u were unbounded on both sides, there would exist a sequence  $\{x_k\}_{k=1}^{\infty}$ , such that  $1 < |x_k| < |x_{k+1}| \to +\infty$  and  $v(x_k) = 0$  for all  $k \in \mathbb{Z}^+$ . Then, it follows from  $|Dv(x)| \le C/|x|$  (for all  $x \in \mathbb{R}^n \setminus \overline{B_1}$ ) that, for any  $k \in \mathbb{Z}^+$  and any  $x \in \partial B_{|x_k|}$ , we have

$$|v(x)| \le \frac{C}{|x_k|} \cdot 2\pi |x_k| = 2C\pi.$$

By the maximum principle, we conclude that  $|v(x)| \leq 2C\pi$  on  $\overline{B}_{|x_{k+1}|} \setminus B_{|x_k|}$  for all  $k \in \mathbb{Z}^+$ . Therefore,  $|v(x)| \leq 2C\pi$  on  $\mathbb{R}^n \setminus \overline{B_{|x_1|}}$ , contradicts the unboundedness assumption.

Translating v by a constant doesn't affect the equation, so we can apply Theorem 4.3 and obtain the asymptotic results.

#### 5 Proof of Theorem 1.5

In this section, we provide the proof of Theorem 1.5 following the line of Li-Li-Yuan [26], some barrier functions are necessary to enhance the convergence speed from (3.11).

**Lemma 5.1**  $(|x|^{-\frac{1}{2}}$  as a Barrier). If a smooth function u satisfies the differential inequality

$$a_{ij}(x)D_{ij}u \ge g(x), \quad in \quad \mathbb{R}^n \setminus \overline{B_1},$$
(5.1)

and  $u \to 0$ , as  $|x| \to \infty$ , where the coefficients are uniformly elliptic and satisfy  $a_{ij}(x) \to a_{ij}(\infty)$ , as  $|x| \to \infty$ , for some positive symmetric matrix  $a_{ij}(\infty)$ . Suppose that

$$\limsup_{|x| \to \infty} |x|^{\frac{3}{2}} |g(x)| < \infty.$$

Then for some constant C,  $u(x) \leq C|x|^{-\frac{1}{2}}, \forall |x| \geq 2$ .

*Proof.* By suitable change of coordinate as in Lemma 6.1 of [18], we may assume without loss of generality that  $a_{ij}(\infty) = \delta_{ij}$ . Since the change of coordinate only relies on  $a_{ij}(\infty)$ , which is bounded w.r.t x variable, hence the result still holds true by allowing C relies on  $a_{ij}(\infty)$ .

To be more explicit, let P be a positive constant matrix such that  $\widetilde{a_{ij}}(\infty) = P^T a_{ij}(\infty) P$  is diagonal matrix whose diagonal elements are the eigenvalues  $\lambda_1, \dots, \lambda_n$  of  $a_{ij}(\infty)$ . Furthermore, by taking  $D = \text{diag}(\lambda_i^{-1/2} \delta_{ij})$ , we have

$$Q := PD, \ \widetilde{u}(x) := u(xQ), \ [\widetilde{a_{ij}}(x)] := D^T P^T[a_{ij}(xQ)] PD \to \delta_{ij} \text{ as } |x| \to \infty.$$
(5.2)

Since P, D, Q only relies on  $a_{ij}(\infty)$ , if the result holds for  $a_{ij}(\infty) = \delta_{ij}$ , which means  $\tilde{u}(x) \leq C|x|^{-\frac{1}{2}}$ , then it follows that

$$u(x) = \widetilde{u}(Q^{-1}x) \le C|Q^{-1}x|^{-\frac{1}{2}} \le C|x|^{-\frac{1}{2}}.$$

Now we prove the result for  $a_{ij}(\infty) = \delta_{ij}$ . By linearity and comparison principle, we only need to proof that  $|x|^{-\frac{1}{2}}$  forms a supersolution of this uniformly elliptic equation. By direct calculating, we have

$$a_{ij}(x)D_{ij}(|x|^{-\frac{1}{2}}) = (-\frac{1}{2})\sum_{i=1}^{n} a_{ii}(x)|x|^{-\frac{3}{2}} + (-\frac{3}{2}) \cdot (-\frac{1}{2})\sum_{i,j=1}^{n} a_{ij}(x)x_ix_j|x|^{-\frac{5}{2}}.$$

Due to  $a_{ij}(x) \to \delta_{ij}$  as  $|x| \to \infty$ ,

$$LHS = -\frac{2n-3}{4}|x|^{-\frac{3}{2}} + o(|x|^{-\frac{3}{2}}).$$

Hence as long as

$$\lim_{|x|\to\infty} |x|^k |g(x)| < \infty, \quad \text{for some} \quad k \ge \frac{3}{2},$$

we can pick a sufficiently large constant C > 0, R > 2 such that

$$a_{ij}(x)D_{ij}(|x|^{-\frac{1}{2}}) \le -\frac{|g(x)|}{C}, \quad \forall |x| \ge R.$$

Then we can pick a even larger C such that

$$C|x|^{-\frac{1}{2}} > u(x), \quad \forall x \in \partial B_R.$$

Comparison principle tells us  $u(x) \leq C|x|^{-\frac{1}{2}}$  for  $R \leq |x|$ . Due to u is a smooth function, hence it maintains bounded inside  $B_R \setminus B_2$ . The result follows immediately by picking sufficiently large C.

**Lemma 5.2**  $(|x|^{2-n} - |x|^{2-n-\varepsilon})$  as a Barrier ). If a smooth function u satisfies (5.1) and  $u \rightarrow 0$ , as  $|x| \rightarrow \infty$ , with coefficients uniformly elliptic and satisfy

$$|a_{ij}(x) - a_{ij}(\infty)| \le C|x|^{-\alpha}, \quad as \quad |x| \to \infty,$$

for some positive symmetric matrix  $a_{ij}(\infty)$ . If  $0 < \varepsilon < \alpha$  and

$$g(x) \ge 0$$
, or  $\limsup_{|x|\to\infty} |x|^n |g(x)| < \infty$ .

 $\label{eq:constant} \textit{Then for some constant } C, \, u(x) \leq C(|x|^{2-n}-|x|^{2-n-\varepsilon}), \; \forall |x| \geq 2.$ 

*Proof.* As the argument in Lemma 5.1, we may assume without loss of generality that  $a_{ij}(\infty) = \delta_{ij}$ . Otherwise we use change of coordinate as in (5.2), if the result holds for  $a_{ij}(\infty) = \delta_{ij}$ , then  $\forall |x| > 2$ , there exists constant C such that

$$\begin{aligned} u(x) &= \widetilde{u}(Q^{-1}x) &\leq C(|Q^{-1}x|^{2-n} - |Q^{-1}x|^{2-n-\varepsilon}) \\ &\leq C(|Q^{-1}|^{2-n}|x|^{2-n} - |Q^{-1}|^{2-n-\varepsilon}|x|^{2-n-\varepsilon}). \end{aligned}$$

If  $|Q^{-1}|^{2-n} \leq |Q^{-1}|^{2-n-\varepsilon}$ , then we immediately obtain

$$u(x) \le C|Q^{-1}|^{2-n-\varepsilon}(|x|^{2-n}-|x|^{2-n-\varepsilon}).$$

If  $|Q^{-1}|^{2-n} > |Q^{-1}|^{2-n-\varepsilon}$ , then for sufficiently large  $|x| \gg 1$  there exists some constant  $c_0$  such that

$$c_0|Q^{-1}|^{2-n}(|x|^{2-n}-|x|^{2-n-\varepsilon}) \ge (|Q^{-1}|^{2-n}-|Q^{-1}|^{2-n-\varepsilon})|x|^{2-n-\varepsilon},$$

hence we have

$$\begin{aligned} u(x) &\leq C |Q^{-1}|^{2-n} (|x|^{2-n} - |x|^{2-n-\varepsilon}) + (|Q^{-1}|^{2-n} - |Q^{-1}|^{2-n-\varepsilon}) |x|^{2-n-\varepsilon} \\ &\leq (C+c_0) |Q^{-1}|^{2-n} (|x|^{2-n} - |x|^{2-n-\varepsilon}) \end{aligned} , \ \forall |x| \gg 1. \end{aligned}$$

Hence we only need to prove the result for  $a_{ij}(\infty) = \delta_{ij}$ . Again, we verify  $|x|^{2-n} - |x|^{2-n-\varepsilon}$  is a barrier by direct calculation

$$a^{ij}(x)D_{ij}|x|^{2-n} = (2-n)\sum_{i=1}^{n} a_{ii}(x)|x|^{-n} + (-n)\cdot(2-n)\sum_{i,j=1}^{n} a_{ij}(x)x_ix_j|x|^{-n-2}.$$

By the condition of coefficients, we have

$$|a^{ii}(x) - 1| \le \frac{C}{|x|^{\alpha}}, \ |a^{ij}(x) - 0| \le \frac{C}{|x|^{\alpha}}, \ \forall i \ne j,$$

hence

$$\begin{aligned} a^{ij}(x)D_{ij}|x|^{2-n} &\leq n(2-n)|x|^{-n} - n(2-n)|x|^{-n} + |(2-n)|\sum_{i=1}^{n} \frac{C}{|x|^{\alpha}}|x|^{-n} \\ &+ |(-n)| \cdot |(2-n)|\sum_{i,j=1}^{n} \frac{C}{|x|^{\alpha}} x_{i}x_{j}|x|^{-n-2} \\ &\leq 0+n \cdot |(2-n)|C|x|^{-n-\alpha}. \end{aligned}$$

Similarly we calculate

$$a^{ij}D_{ij}|x|^{2-n-\varepsilon} = (2-n-\varepsilon)\sum_{i=1}^{n}a_{ii}(x)|x|^{-n-\varepsilon} + (-n-\varepsilon)\cdot(2-n-\varepsilon)\sum_{i,j=1}^{n}a_{ij}(x)x_ix_j|x|^{-n-2-\varepsilon},$$

and hence

$$a^{ij}D_{ij}|x|^{2-n-\varepsilon} \ge -\varepsilon(2-n-\varepsilon)|x|^{-n-\varepsilon} - C_n|x|^{-n-\alpha-\varepsilon}.$$

Combining these two result we have

$$a^{ij}(x)D_{ij}(|x|^{2-n} - |x|^{2-n-\varepsilon}) \le C \cdot n |(2-n)| \cdot |x|^{-n-\alpha} + \varepsilon (2-n-\varepsilon) \cdot |x|^{-n-\varepsilon} + C_n |x|^{-n-\alpha-\varepsilon}.$$

As long as  $\varepsilon < \alpha$ , the negative term  $\varepsilon(2 - n - \varepsilon)|x|^{-n-\varepsilon}$  takes the lead and makes this less than |g(x)| when |x| is sufficiently large due to the constants here are universal. Then comparison principle tells us the result.

*Proof of Theorem 1.5.* By Theorem 3.5, there exist  $A \in \mathcal{A}(f(\infty), -a+b)$ ,  $\alpha > 0$ ,  $R_2 \gg 1$  sufficiently large such that (3.11) holds.

Set  $v := u(x) - \frac{1}{2}x^T A x$  then we have

$$\left\{ \begin{array}{l} F_{\tau}(D^2v+A) = F_{\tau}(D^2u) = f(x), \\ F_{\tau}(A) = \lim_{|x| \to \infty} F_{\tau}(D^2u(x)) = f(\infty) \end{array} \right.$$

Hence

$$\widetilde{f}(x) := f(x) - f(\infty) = \int_0^1 D_{M_{ij}} F_\tau(tD^2v + A) \mathrm{d}t \cdot D_{ij}v =: \overline{a^{ij}}(x) D_{ij}v.$$

Also, from taking derivatives to any  $e \in \partial B_1$  direction, we consider the linearized equation as (4.1),  $v_e$  and  $v_{ee}$  satisfy

$$D_{M_{ij}}F_{\tau}(D^2v+A)D_{ij}(v_e) = f_e(x), \text{ and } D_{M_{ij}}F_{\tau}(D^2v+A)D_{ij}(v_{ee}) \ge f_{ee}(x).$$
 (5.3)

Let  $\widehat{a^{ij}}(x)$  denote the coefficients  $D_{M_{ij}}F_{\tau}(D^2v+A)$ .

Due to  $F_{\tau}$  is uniformly  $C^2$  in a bounded subdomain of Sym(n), hence for the same  $\alpha$  from Theorem 3.5, there exists some constant C such that

$$\left|\overline{a_{ij}}(x) - D_{M_{ij}}F_{\tau}(A)\right|, \left|\widehat{a_{ij}}(x) - D_{M_{ij}}F_{\tau}(A)\right| \le \frac{C}{|x|^{\alpha}}.$$

Thus Theorem 4.3 and Corollary 4.4 hold for the linear equations with coefficients  $\overline{a_{ij}}(x)$ ,  $\widehat{a_{ij}}(x)$ . From condition (1.11) and Lemma 5.2, we have the following finer convergence speed estimate  $v_{ee}(x) \leq C|x|^{2-n}$ . hence

$$\lambda_{\max}\left(D^2 v\right)(x) \le C|x|^{2-n},$$

and the ellipticity of equation (5.3) tells us

$$\lambda_{\min} \left( D^2 v \right)(x) \ge -C\lambda_{\max} \left( D^2 v \right)(x) - C|\tilde{f}(x)| \ge -C|x|^{2-n}.$$

Hence

$$\left|D^2 v(x)\right| \le C|x|^{2-n}.$$

Therefore

$$\left|\overline{a^{ij}}(x) - D_{M_{ij}}F_{\tau}(A)\right| \le C|x|^{2-n}, \quad \forall x \in B_1^c,$$

and

$$\left|\widehat{a^{ij}}(x) - D_{M_{ij}}F_{\tau}(A)\right| \le C|x|^{2-n}, \quad \forall x \in B_1^c.$$

Thus for any  $e \in \partial B_1$  we have

$$|Dv_e| \le \left| D^2 v(x) \right| \le C |x|^{2-n},$$

which provides us "bounded from one side" condition. Since  $F_{\tau}$  is uniformly  $C^2$  on the range of  $\lambda(D^2u)$ , hence Theorem 3.6 tells us the coefficients  $\overline{a^{ij}}$ ,  $\widehat{a^{ij}}$  has bounded  $C^{\alpha}$  norm. Also note that

the coefficients have a Hölder convergence speed and |Df| has  $O(|x|^{-(n+\varepsilon)})$  order decay. These conditions enable us to apply Corollary 4.4 to equation  $\widehat{a^{ij}}(x)D_{ij}(v_e) = f_e(x)$  and obtain

$$\exists \beta_e \in \mathbb{R} \text{ s.t. } v_e(x) = \beta_e + O\left(|x|^{2-n}\right), \quad as|x| \to \infty.$$
(5.4)

Picking  $e_i := (0, \dots, 0, \underbrace{1}_{\substack{\text{the } i-\text{th variable}}}, 0, \dots, 0)$  and we use  $\beta_i$  denote  $\beta_{e_i}$  from formula (5.4). Set  $\beta := (\beta_1, \beta_2, \dots, \beta_n) \in \mathbb{R}^n$  and

$$\overline{v}(x) := v(x) - \beta^T x = u(x) - \left(\frac{1}{2}x^T A x + \beta^T x\right).$$

Then formula (5.4) tells us

$$D\overline{v}(x)| = |(\partial_1 v - \beta_1, \cdots, \partial_n v - \beta_n)| = O(|x|^{2-n}).$$

Also note that it satisfies the linearized equation

$$\overline{a}^{ij}(x)D_{ij}\overline{v} = \overline{a}^{ij}(x)D_{ij}v = \widetilde{f}(x)$$

By the arguments above again and adapt Corollary 4.4 we have

$$\exists \gamma \ s.t. \ \overline{v}(x) = \gamma + O(|x|^{2-n}), \ as \ |x| \to \infty.$$

Set 
$$Q(x) = \frac{1}{2}x^T A x + \beta^T x + \gamma$$
 and the formula above tells us  
 $|u - Q| = |\overline{v} - \gamma| = O(|x|^{2-n}), \text{ as } |x| \to \infty.$ 

#### 6 Proof of Theorem 1.6

In this section, we provide an asymptotic behavior result that is parallel to the one by Bao-Li-Zhang [3]. From Theorem 3.5, we have proved that under suitable assumption of f, the Hessian matrix  $D^2u$  admits a limit A at infinity with  $O(|x|^{-\alpha})$  order.

Instead of using the original results in [3], we transform back using the definition of Legendre transform and follow the line of proving Lemma 2.1 in [3] to obtain estimates of up to (m + 1)-order derivative of solutions and Hölder estimate of  $D^{m+1}u(x)$  directly.

**Lemma 6.1.** Under the conditions of Theorem 1.6, suppose that there exists  $\varepsilon > 0$  such that

$$|D^2 u(x) - A| \le c_1 |x|^{-\varepsilon}, \quad |x| \ge R_0.$$

Assume without loss of generality that u(0) = 0, Du(0) = 0. Let

$$w(x) = u(x) - \frac{1}{2}x^T A x,$$

then there exist  $C(n, R_0, \varepsilon, f(\infty), c_1, \zeta) > 0$  and  $R_1(n, R_0, \varepsilon, f(\infty), c_1, \zeta) > R_0$  such that for any  $\alpha \in (0, 1)$ ,

$$\begin{aligned} \left| D^{k}w(y) \right| &\leq C|y|^{2-k-\varepsilon_{\zeta}}, \quad k = 0, \dots, m+1, \quad |y| > R_{1}, \\ \frac{\left| D^{m+1}w(y_{1}) - D^{m+1}w(y_{2}) \right|}{|y_{1} - y_{2}|^{\alpha}} &\leq C|y_{1}|^{1-m-\varepsilon_{\zeta}-\alpha}, \quad |y_{1}| > R_{1}, \ y_{2} \in B_{\frac{|y_{1}|}{2}}(y_{1}), \end{aligned}$$

where  $\varepsilon_{\zeta} := \min\{\varepsilon, \zeta\}.$ 

*Proof.* By direct calculate and our assumption u(0) = 0,  $Du(0) = \overrightarrow{0}$ , it follows that for some  $C(n, u|_{B_{R_0}}, c_1)$ 

$$|w(x)| \le C|x|^{2-\varepsilon}, \ \forall |x| \ge R_0.$$
(6.1)

For  $|x| = R > 2R_0$ , let

$$u_R(y) = \left(\frac{4}{R}\right)^2 u\left(x + \frac{R}{4}y\right), \text{ and } w_R(y) = \left(\frac{4}{R}\right)^2 w\left(x + \frac{R}{4}y\right), |y| \le 2.$$

Then the boundedness of  $D^2u$  together with (6.1) tell us

$$||u_R||_{L^{\infty}(B_2)} \le C, ||w_R||_{L^{\infty}(B_2)} \le CR^{-\varepsilon},$$

for some constant C uniform to  $R > 2R_0$ .

Now we attack the original equation by scaling back to unit size. It is easy to verify that  $u_R$  satisfies

$$F_{\tau}(\lambda(D^2 u_R(y))) = f_R(y) := f(x + \frac{R}{4}y), \text{ in } B_2.$$
 (6.2)

Take difference with  $F_{\tau}(\lambda(A)) = f(\infty)$  gives us

$$\widetilde{a_{ij}}(y)D_{ij}w_R = f_R(y) - f(\infty) = O(R^{-\zeta}),$$

where  $\widetilde{a_{ij}}(y) := \int_0^1 D_{M_{ij}} F_{\tau}(A + tD^2 w_R(y)) dt$  is uniformly elliptic and Theorem 3.6 tells us  $\widetilde{a_{ij}}(y)$  have bounded  $C^{\alpha}$  norm.

Apply the classical Schauder's estimate to obtain the  $C^{2,\alpha}$  regularity of  $w_R$ 

$$\|w_R\|_{C^{2,\alpha}(\overline{B_1})} \le C\left(\|w_R\|_{L^{\infty}(\overline{B_{1,1}})} + \|f_R - f(\infty)\|_{C^{\alpha}(\overline{B_1})}\right) \le CR^{-\varepsilon_{\zeta}}.$$

Now we take derivative with respect to  $e \in \mathbb{S}^{n-1}$  direction to equation (6.2) and obtain

$$D_{M_{ij}}F_{\tau}(D^2u_R(y))D_{ij}(\partial_e u_R(y)) = \partial_e f_R(y), \quad \text{in } B_2.$$
(6.3)

Since  $D_{M_{ij}}F_{\tau}(D^2u_R(y)), \partial_e u_R$  and  $\partial_e f_R$  are bounded in  $C^{\alpha}$  norm, we have

$$||u_R||_{C^{3,\alpha}(\overline{B_1})} \le C,$$

which implies

$$|D_{M_{ij}}F_{\tau}(D^2u_R(y))||_{C^{3,\alpha}(\overline{B_1})} \le C.$$

By the definition of  $w_R$ , we see that

$$D^{2}w_{R}(y) = D^{2}w(x + \frac{R}{4}y) = D^{2}u(x + \frac{R}{4}y) - A = D^{2}u_{R}(y) - A$$

and hence  $D_{ij}(\partial_e u_R(y)) = D_{ij}(\partial_e w_R(y))$ . Hence equation (6.3) can also be written into

$$D_{M_{ij}}F_{\tau}\left(D^2u_R(y)\right)D_{ij}\left(\partial_e w_R(y)\right) = \partial_e f_R(y), \quad \text{in } B_2.$$

We obtain by Schauder's estimate

$$||w_R||_{C^{3,\alpha}(\overline{B_{1/2}})} \le C\left(||w_R||_{L^{\infty}(\overline{B_{3/4}})} + |Df_R||_{C^{\alpha}(\overline{B_{3/4}})}\right) \le CR^{-\varepsilon_{\zeta}},$$

which provides us

$$||D^3 u_R||_{C^{\alpha}(\overline{B_{1/2}})} \le CR^{-\varepsilon_{\zeta}}$$

Estimates on higher order derivatives can be obtained by further differentiation of the equation and Schauder estimate. The result follows by induction immediately.  $\Box$ 

Next we prove a bootstrap lemma to improve the estimates in Lemma 6.1. Lemma 6.2 is originally proved for Monge-Ampère equation in [3] and we work with general uniformly elliptic equation with bounded derivative of F operator. Consider the uniformly elliptic equation

$$F(D^2u) = f(x), \quad \text{in} \quad \mathbb{R}^n \setminus B_{R_0}, \tag{6.4}$$

where F is smooth up to boundary of  $D := \overline{D^2 u(\mathbb{R}^n \setminus \overline{B_{R_0}})}$  i.e. the range of  $D^2 u$  and  $f(x) \in C^m(\mathbb{R}^n \setminus \overline{B_{R_0}})$  satisfies condition (1.12) for some  $\zeta > 2$ ,  $m \ge 3$ . We write  $D_{M_{ij}}F(M)$  as  $F_{M_{ij}}(M)$  for simplicity.

**Lemma 6.2.** Let u be a solution of equation (6.4) described above. Let v be a 2-order polynomial such that  $F(D^2v) = f(\infty)$  and let w := u - v. Suppose that for some  $0 < \varepsilon < \frac{1}{2}, \alpha \in (0, 1)$ , we have

$$\left| D^{\kappa} w(x) \right| \le C |x|^{2-\varepsilon-\kappa}, \quad |x| > 2R_1, \quad k = 0, \dots, m+1,$$

$$\frac{\left| D^{m+1} w\left(y_1\right) - D^{m+1} w\left(y_2\right) \right|}{\left|y_1 - y_2\right|^{\alpha}} \le C \left|y_1\right|^{1-m-\varepsilon-\alpha}, \quad |y_1| > 2R_1, y_2 \in B_{|y_1|/2}\left(y_1\right)$$

Then

$$\begin{aligned} \left| D^{k} w(x) \right| &\leq C |x|^{2-2\varepsilon-k}, \quad |x| > 2R_{1}, \quad k = 0, \dots, m+1, \\ \frac{\left| D^{m+1} w\left(y_{1}\right) - D^{m+1} w\left(y_{2}\right) \right|}{|y_{1} - y_{2}|^{\alpha}} &\leq C \left| y_{1} \right|^{1-m-2\varepsilon-\alpha}, \quad |y_{1}| > 2R_{1}, y_{2} \in B_{|y_{1}|/2}\left(y_{1}\right). \end{aligned}$$

*Proof.* Applying  $\partial_k$  to equation (6.4) and we obtain

$$\widehat{a_{ij}}D_{ij}(\partial_k u(x)) = \partial_k f(x), \tag{6.5}$$

where  $\widehat{a_{ij}}(x) := F_{M_{ij}}(D^2u(x)).$ 

Then this linearized equation is also uniformly elliptic with the coefficients satisfies (from the assumptions)

$$|\widehat{a_{ij}}(x) - F_{M_{ij}}(A)| \le ||DF||_{C^0(D)} \cdot \frac{C}{|x|^{\varepsilon}}, \quad \text{and} \quad |D\widehat{a_{ij}}(x)| \le \frac{C}{|x|^{1+\varepsilon}}, \quad |x| > R_1.$$

Also, for the  $\alpha$  in condition, we have <sup>2</sup>

$$\frac{|Da_{ij}(x_1) - Da_{ij}(x_2)|}{|x_1 - x_2|^{\alpha}} \le C||D^2F||_{C^0(D)} \cdot |x_1|^{-1-\varepsilon-\alpha}, \quad |x_1| > 2R_1, \quad x_2 \in B_{|x_1|/2}(x_1).$$

Apply  $\partial_l$  to the (6.5) and set  $h_1 = \partial_{kl} u$  we further obtain

$$F_{M_{ij},M_{qr}}(D^2u)D_{ijk}uD_{qrl}u + F_{M_{ij}}(D^2u)D_{ij}h_1 = \partial_{kl}f(x).$$

Due to the Hessian matrix converge to  $D^2v$  (positive constant matrix) at infinity with Hölder speed  $F_{M_{ij}}(D^2u) \rightarrow F_{M_{ij}}(D^2v)$  as  $|x| \rightarrow \infty$ . The extremal matrix makes the operator also uniformly elliptic, hence the Green's function of  $F_{M_{ij}}(D^2v)D_{ij}u$  exists and equivalence to the one of Laplacian. Then we can write

$$F_{M_{ij}}(D^2v)D_{ij}h_1 = f_2 := \partial_{kl}f - F_{M_{ij},M_{qr}}(D^2u)D_{ijk}uD_{qrl}u - (F_{M_{ij}}(D^2u) - F_{M_{ij}}(D^2v))D_{ij}h_1 - f_{M_{ij}}(D^2v)D_{ij}h_1 - f_{M$$

From the assumptions we stated, it follows that for the  $\alpha \in (0, 1)$  in the assumption,

$$|f_2(x)| \le C|x|^{-2-2\varepsilon}, \ \forall |x| \ge 2R_1,$$
$$\frac{|f_2(x_1) - f_2(x_2)|}{|x_1 - x_2|^{\alpha}} \le \frac{C}{|x_1|^{2+2\varepsilon+\alpha}}, \quad x_2 \in B_{|x_1|/2}(x_1), |x_1| \ge 2R_1.$$

Note that this step demanded 3-order derivative of f. This is essential for this method due to it provides the Hölder continuity of  $f_2$ , which is used in Schauder estimate.

Note that  $h_1 \to D_{kl}v$  for fixed  $k, l \in \{1, 2, \dots, n\}$ ,  $h_1$  satisfies the following "part" Dirichlet problem

$$\begin{cases} F_{M_{ij}}(D^2v)D_{ij}h_1 = f_2, & in \mathbb{R}^n \setminus B_{R_1}, \\ h_1 \to D_{kl}v, & as |x| \to \infty. \end{cases}$$

The main target of this bootstrap lemma is to obtain a finer estimate on the convergence speed of Hessian matrix. From  $O(|x|^{-\varepsilon})$  enhanced into  $O(|x|^{-2\varepsilon})$  as long as  $2\varepsilon < 1$ . In order to do this, we use a Green's function to change the equation into homogeneous situation.

Let

$$h_2(x) := \int_{\mathbb{R}^n \backslash B_{R_1}} G_{F_{M_{ij}}(D^2v)}(x,y) f_2(y) \mathrm{d}y,$$

where  $G_{{\cal F}_{{\cal M}_{ii}}(D^2v)}(x,y)$  is the distribution solution (Green's function) of

$$\left\{ \begin{array}{cc} F_{M_{ij}}(D^2v)D_{ij}u(y)=\delta_x & \text{ in } \mathbb{R}^n\\ u(y)\to 0 & \text{ as } |y|\to\infty \end{array} \right.$$

<sup>&</sup>lt;sup>2</sup>For simplicity, we use  $a_{ij}$  stands for this  $\widehat{a_{ij}}$ . Other linearize processes are similar to this.

The existence of such a Green's function and equivalence to fundamental solution is well-known, see [32, 21] for example. Then  $F_{M_{ij}}(D^2v)D_{ij}h_2 = f_2$  and satisfies  $h_2 \to 0$  as  $|x| \to \infty$ . Moreover, argue as in the proof of Lemma 4.1, separate  $\mathbb{R}^n$  into three pieces  $E_1, E_2, E_3$  and

Moreover, argue as in the proof of Lemma 4.1, separate  $\mathbb{R}^n$  into three pieces  $E_1, E_2, E_3$  and we obtain

$$|D^{j}h_{2}(x)| \le C|x|^{-2\varepsilon-j}, \quad |x| > 2R_{1}, \text{ for } j = 0, 1.$$

Argue as in the proof of Lemma 6.1, we also obtain

$$\frac{\left|D^{2}h_{2}\left(x_{1}\right)-D^{2}h_{2}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq \frac{C}{\left|x_{1}\right|^{2+2\varepsilon+\alpha}}, \quad x_{2} \in B_{\frac{\left|x_{1}\right|}{2}}\left(x_{1}\right), \left|x_{1}\right| > 2R_{1}.$$

Indeed, for each  $x_0 \in \mathbb{R}^n \setminus B_{2R_1}$ , let  $R = |x_0|$ , we set

$$h_{2,R}(y) = h_2\left(x_0 + \frac{R}{4}y\right), \quad f_{2,R}(y) = \frac{R^2}{16}f_2\left(x_0 + \frac{R}{4}y\right), \quad |y| \le 2.$$

Then we have

$$|h_{2,R}||_{C^0(\overline{B_1})} = ||h_2||_{C^0(\overline{B_{\lfloor x_0 \rfloor}(x_0)})} \le CR^{-2\varepsilon}, \text{ and } ||f_{2,R}||_{C^\alpha(\overline{B_1})} \le CR^{-2\varepsilon}.$$

Hence Schauder Estimate tells us

$$\|h_{2,R}\|_{C^{2,\alpha}(\overline{B_1})} \le C\left(\|h_{2,R}\|_{L^{\infty}(B_2)} + \|f_{2,R}\|_{C^{\alpha}(\overline{B_2})}\right) \le CR^{-2\varepsilon}.$$

Meaning that we have obtained the desired result for  $h_2$  i.e.

$$|D^{j}h_{2}(x)| \leq C|x|^{-2\varepsilon-j}, \quad j = 0, 1, 2, \quad |x| > 2R_{1},$$

and

$$\frac{\left|D^{2}h_{2}\left(x_{1}\right)-D^{2}h_{2}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq \frac{C}{\left|x_{1}\right|^{2}+2\varepsilon+\alpha}, \quad x_{2}\in B_{\frac{\left|x_{1}\right|}{2}}\left(x_{1}\right), \left|x_{1}\right|>2R_{1}.$$

Now we only need to study the difference between  $h_1 - D_{kl}v$  and  $h_2$ . By the linearity of Laplacian operator we have

$$\begin{cases} F_{M_{ij}}(D^2v)D_{ij}(h_1 - D_{kl}v - h_2) = 0, & in \ \mathbb{R}^n \setminus B_{2R_1}, \\ h_1 - D_{kl}v - h_2 \to 0, & as \ |x| \to \infty. \end{cases}$$

By taking  $|x|^{2-n}$  as a barrier function, it immediately follows that for some constant C

$$|h_1(x) - D_{kl}v - h_2(x)| \le C|x|^{2-n}, \quad |x| > 2R_1.$$

Due to  $|x|^{2-n}$  converge faster than  $|x|^{-2\varepsilon}$ , triangle inequality tells us

$$|h_1(x) - D_{kl}v| \le C|x|^{-2\varepsilon}, \quad |x| > 2R_1.$$

Then Newton-Leibnitz formula tells us

$$|D^{j}w(x)| \le C|x|^{2-j-2\varepsilon}, \quad |x| > 2R_{1}, \quad j = 0, 1, 2.$$

Higher regularity (when m is larger than 3), the result follows by taking more derivatives.

*Proof of Theorem 1.6.* The theorem follows exactly as proved in [3], which is based on Level Set Method and bootstrap argument we proved earlier. Since the proof is slightly different, we provide the details here again.

From the boundedness of Hessian matrix, we can verify that the conditions of Theorem 3.5 still holds. Meaning that there exist  $A \in A_3$ ,  $\alpha > 0$ ,  $R_2 \gg 1$  sufficiently large such that

$$\left|D^2u(x) - A\right| \le \frac{C}{|x|^{\alpha}}, \quad \forall |x| \ge R_2.$$

This enable us to apply Lemma 6.1 which gives us the initial point of doing iteration as in Lemma 6.2 to obtain a faster decay speed. We can always do this finite times till the condition of  $0 < \varepsilon < \frac{1}{2}$  in Lemma 6.2 fails to hold. As in Lemma 6.1, we denote  $w(x) = u(x) - \frac{1}{2}x^T Ax$ , where A is the limit of  $D^2u$  at infinity, which is provided by Theorems 3.5 and 3.6 in Section 3.

Let  $k_0$  be the positive integer such that  $2^{k_0}\varepsilon < 1$  and  $2^{k_0+1}\varepsilon > 1$  (we choose  $\varepsilon$  smaller if necessary to make both inequalities hold). Let  $\varepsilon_1 = 2^{k_0}\varepsilon$ , clearly we have  $1 < 2\varepsilon_1 < 2$ . Applying Lemma 6.1  $k_0$  times we have

$$|D^k w(x)| \le C|x|^{2-\varepsilon_1-k}, \quad k = 0, \dots, m+1, \quad |x| > 2R_1,$$

and

$$\frac{\left|D^{m+1}w(x_{1}) - D^{m+1}w(x_{2})\right|}{|x_{1} - x_{2}|^{\alpha}} \le C |x_{1}|^{1 - m - \varepsilon_{1} - \alpha}, |x_{1}| > 2R_{1}, x_{2} \in B_{|x_{1}|/2}(x_{1}).$$

Let  $h_1$  and  $f_2$  be the same as in Lemma 6.2. Then we have

$$|f_2(x)| \le C|x|^{1-m-2\varepsilon_1} + C|x|^{-2-\zeta} \quad |x| \ge 2R_1,$$

and

$$\frac{\left|f_{2}\left(x_{1}\right)-f_{2}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq \frac{C}{\left|x_{1}\right|^{m-1+2\varepsilon_{1}+\alpha}} + \frac{C}{\left|x\right|^{\zeta+2+\alpha}}, \quad \left|x_{1}\right| \geq 2R_{1}, x_{2} \in B_{\left|x_{1}\right|/2}\left(x_{1}\right).$$

Constructing  $h_2$  as in the proof of Lemma 6.2, we have

$$|D^{j}h_{2}(x)| \leq C|x|^{-2\varepsilon_{1}-j}, \quad j = 0, 1, 2, |x| > 2R_{1},$$

and

$$\frac{D^{2}h_{2}(x_{1}) - D^{2}h_{2}(x_{2})|}{|x_{1} - x_{2}|^{\alpha}} \leq \frac{C}{|x_{1}|^{2+2\varepsilon_{1}+\alpha}}, x_{2} \in B_{\frac{|x_{1}|}{2}}(x_{1}), |x_{1}| > 2R_{1}.$$

Similar to the proof of Lemma 6.2, it follows from the estimate above that

$$|h_1(x) - h_2(x)| \le C|x|^{2-n}, \quad |x| > 2R_1.$$

Since  $2\varepsilon_1 > 1$ ,

$$|h_1(x)| \le |h_2(x)| + C|x|^{2-n} \le C|x|^{-1}$$

By Theorem 4 of [17], for any  $i = 1, 2, \dots, n$ ,  $\partial_i w(x) \to c_i$  for some  $c_i \in \mathbb{R}$  as  $|x| \to \infty$ . Let  $\beta \in \mathbb{R}^n$  be the limit of Dw and  $w_1(x) = w(x) - \beta \cdot x$ . The equation satisfied by  $\partial_e w_1$  can be written as (for  $e \in \mathbb{S}^{n-1}$ )

$$a_{ij}\partial_{ij} \left(\partial_e w_1\right) = \partial_e f_v$$
, where  $a_{ij} = D_{M_{ij}}F_{\tau}(D^2 u) \to D_{M_{ij}}F_{\tau}(A)$ .

Write the equation into the perturbation of elliptic equation with constant coefficients, then from the estimate above we have

$$D_{M_{ij}}F_{\tau}(A)\left(\partial_{e}w_{1}\right) = f_{3} := \partial_{e}f_{v} - \left(D_{M_{ij}}F_{\tau}(D^{2}u) - D_{M_{ij}}F_{\tau}(A)\right)\partial_{ije}w_{1}, \quad |x| > 2R_{1},$$
$$|f_{3}(x)| \leq C\left(|x|^{-\zeta-1} + |x|^{-1-2\varepsilon_{1}}\right) \leq C|x|^{-1-2\varepsilon_{1}}, |x| > 2R_{1},$$

and

$$\frac{\left|f_{3}\left(x_{1}\right)-f_{3}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq C\left|x_{1}\right|^{-1-2\varepsilon_{1}-\alpha}, \left|x_{1}\right| > 2R_{1}, x_{2} \in B_{\left|x_{1}\right|/2}\left(x_{1}\right)$$

Let  $h_4$  solve  $D_{M_{ij}}F_{\tau}(A)h_4 = f_3$  and the construction of  $h_4$  is similar to that of  $h_2$  in Lemma 6.2. Then we have

$$|D^{j}h_{4}(x)| \leq C|x|^{1-2\varepsilon_{1}-j}, |x| > 2R_{1}, j = 0, 1, 2,$$

and

$$\frac{\left|D^{2}h_{4}\left(x_{1}\right)-D^{2}h_{4}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq C\left|x_{1}\right|^{-1-2\varepsilon_{1}-\alpha}, \left|x_{1}\right| > 2R_{1}, x_{2} \in B_{\left|x_{1}\right|/2}\left(x_{1}\right).$$

Since  $\partial_e w_1 - h_4 \rightarrow 0$  at infinity, we have

$$|\partial_e w_1(x) - h_4(x)| \le C|x|^{2-n}, \quad |x| > R_1.$$

Therefore we have obtained  $|Dw_1(x)| \le C|x|^{1-2\varepsilon_1}$  on  $|x| > R_1$ . Using fundamental theorem of calculus it tells us

$$|w_1(x)| \le C|x|^{2-2\varepsilon_1}, \quad j = 0, 1, \quad |x| > R_1.$$

Lemma 6.1 applied to  $w_1$  gives

$$|D^{j}w_{1}(x)| \leq C|x|^{2-j-2\varepsilon_{1}}, \quad j = 0.., m+1.$$

This provides us a finer estimate on  $f_3$  i.e.

$$|f_3(x)| \le C|x|^{-\zeta-1} + C|x|^{-1-4\varepsilon_1}, |x| > 2R_1,$$

and

$$\frac{|f_3(x_1) - f_3(x_2)|}{|x_1 - x_2|^{\alpha}} \le C\left(|x_1|^{-\zeta - 1 - \alpha} + |x_1|^{-1 - 4\varepsilon_1 - \alpha}\right), |x_1| > 2R_1, x_2 \in B_{|x_1|/2}(x_1).$$

As a consequence, the new estimate of  $h_4$  is

$$|h_4(x)| \le C\left(|x|^{1-\zeta} + |x|^{1-4\varepsilon_1}\right), \quad |x| > 2R_1,$$

and as always, triangle inequality tells us

$$|Dw_1(x)| \le C(|x|^{2-n} + |x|^{1-4\varepsilon_1}) \le C|x|^{-1}, \quad |x| > 2R_1.$$

By Theorem 4 of [17] again, there exists some constant  $\gamma$  such that  $w_1 \rightarrow \gamma$  at infinity. Let

$$w_2(x) = w_1(x) - \gamma = w(x) - \beta \cdot x - \gamma$$

Then we have  $|w_2(x)| \leq C$  for  $|x| > 2R_1$ . Lemma 6.1 applied to  $w_2$  gives

$$|D^k w_2(x)| \le C|x|^{-k}, \quad k = 0, \dots, m+1, \quad |x| > 2R_1$$

The equation satisfied by  $w_2$  can be written as

$$F_{\tau}(A+D^2w_2(x)) = f_{\tau}$$

Taking the difference between this equation and  $F_{\tau}(A) = f(\infty)$  we have

$$\widetilde{a}_{ij}\partial_{ij}w_2 = f - f(\infty), \quad |x| > 2R_1,$$

where

$$\widetilde{a}_{ij}(x):=\int_0^1 D_{M_{ij}}F_\tau(A+tD^2w_2(x))\mathrm{d}t.$$

Note that the convergence of Hessian (Theorem 3.5) tells us  $\widetilde{a_{ij}}$  satisfy

$$\left| D^k \left( \tilde{a}_{ij}(x) - \tilde{a}_{ij}(\infty) \right) \right| \le C |x|^{-2-k}, \quad |x| > 2R_1, \quad k = 0, 1.$$

Write it into the perturbed situation of constant coefficients operator again

$$\widetilde{a}_{ij}(\infty)D_{ij}w_2 = f_4 := f_v - 1 - \left(\widetilde{a}_{ij} - \widetilde{a}_{ij}(\infty)\right)\partial_{ij}w_2, \quad |x| > 2R_1,$$

and we have the following estimates

$$|f_4(x)| \le C\left(|x|^{-\zeta} + |x|^{-4}\right), \quad |x| > 2R_1,$$

and

$$\frac{\left|f_{4}\left(x_{1}\right)-f_{4}\left(x_{2}\right)\right|}{\left|x_{1}-x_{2}\right|^{\alpha}} \leq C\left(\left|x_{1}\right|^{-\zeta-\alpha}+\left|x_{1}\right|^{-4-\alpha}\right), \quad \left|x_{1}\right|>2R_{1}, x_{2}\in B_{\left|x_{1}\right|/2}\left(x_{1}\right).$$

Let  $h_5$  be defined similar to  $h_2$ , which makes  $h_5$  solves  $\tilde{a}_{ij}(\infty)h_5 = f_4$  in  $\mathbb{R}^n \setminus B_{2R_1}$  and satisfies

$$|h_5(x)| \le C\left(|x|^{2-\zeta} + |x|^{-2}\right)$$

As before we have

$$|w_2(x) - h_5(x)| \le C|x|^{2-n}, \quad |x| > 2R_1,$$

which gives

$$|w_2(x)| \le C\left(|x|^{2-n} + |x|^{2-\zeta} + |x|^{-2}\right), \quad |x| > 2R_1.$$
(6.6)

If  $|x|^{-2} > |x|^{2-n} + |x|^{2-\zeta}$  we can apply the same argument as above finite times to remove the  $|x|^{-2}$  from (6.6). Eventually by Lemma 6.1 we have this result.

## 7 Interior Estimates

From the proof of Theorem 3.5, we see that as long as (3.14) holds, then all the asymptotic behavior also holds. In this section, we will reduce the assumption on Hessian (1.10) into assumption on gradient (1.14) or assumption on solution itself (1.16).

This section is organized in the following order. First, we prove that condition (1.14) provides us the desired equivalence (3.14) immediately. Second, based on an important gradient estimate theorem by Y.Y.Li [28], we can furthermore reduce the condition (1.14) by (1.16). These two parts provide us Theorems 1.8 and 1.9 already. Eventually, by the compactness method developed by McGonagle-Song-Yuan [29], we see that the boundeness of Hessian holds true under a weaker assumption of right hand side function than in Theorems 1.8 and 1.9.

**Lemma 7.1** (Linear Growth). Let  $(\tilde{x}, \tilde{u})$  be the function defined as in (2.1), and u satisfy condition (1.14) for some constant  $C_0$ . Then there exist  $\delta = \delta(n, C_0, \tau)$ ,  $M = M(\tau, \max_{x \in \overline{B_1}} |Du(x)|) > 0$  such that

$$|(\widetilde{x} - D\widetilde{u}(\widetilde{x}))| \ge \delta |\widetilde{x}|, \quad \forall |\widetilde{x}| \ge M$$

*Proof.* Firstly, we prove that

$$|2bx| \ge \delta |Du(x) + (a+b)x|, \quad \forall |x| \ge 1.$$

In fact, by triangle inequality

$$|Du(x) + (a+b)x| \le |Du(x)| + (a+b)|x| \le (C_0 + a+b)(|x|+1), \quad \forall x \in \mathbb{R}^n.$$

Thus by taking a sufficiently small  $\delta > 0$  such that

$$(C_0 + a + b)\delta < \frac{2b}{100}$$
, hence  $2b|x| \ge 100\delta(C_0 + a + b)|x|$ .

Thus as long as  $|x| > \frac{1}{99}$ , we have

$$100\delta(C_0 + a + b)|x| \ge \delta(C_0 + a + b)(|x| + 1).$$

Hence there exists a  $\delta > 0$  such that

$$2bx| \ge \delta |Du(x) + (a+b)x|, \quad \forall |x| \ge 1.$$

Secondly, we prove the following result, then the result follows immediately,

$$\{x : |\tilde{x}| = |Du(x) + (a+b)x| \ge M\} \subset \{x : |x| \ge 1\}.$$
(7.1)

In fact, for any given  $u \in C^4(\mathbb{R}^n)$ , |Du(x)| is bounded in  $B_1$ . Let's denote

$$\widetilde{M}:=\max_{x\in\overline{B_1}}|Du(x)|<\infty,\quad\text{depending on }u.$$

Then for any  $|x| \leq 1$ ,

$$|Du(x) + (a+b)x| \le |Du(x)| + (a+b)|x| \le \widetilde{M} + a + b =: M.$$

This tells us (7.1) through an argue by contradiction.

Now we consider solution of (1.9) with  $D^2 u > (-a + b)I$  and we assume that f satisfies conditions (3.2) and (3.3) for some  $f(\infty) < 0$ ,  $\alpha \in (0, 1)$ ,  $\beta > 1$ ,  $\gamma > 0$ . The following gradient estimate by Y.Y.Li [28] plays an important role in our proof.

Theorem 7.2 (Interior Gradient Estimate). Consider the following equation

$$f(\lambda_1,\ldots,\lambda_n) = \psi(x,u(x),Du(x)),$$

where  $\lambda_i$  is the eigenvalue of  $D^2u(x)$ . Suppose f

- (A) is smooth and defined in an open convex cone  $\Gamma \subset \mathbb{R}^n$ , which is different from  $\mathbb{R}^n$ , with vertex at the origin, containing the positive cone  $\{\lambda \in \mathbb{R}^n : each component \lambda_i > 0\}$
- (B) satisfies the following in  $\Gamma$

$$rac{\partial f}{\partial \lambda_i} > 0 \quad orall i, \quad and \ f \ is \ a \ concave \ function.$$

- (C) is invariant under interchange of any two  $\lambda_i s$ .
- (D) there exist  $c_0, c_1 > 0$ , which depend on  $\psi_0$ , such that, for any  $\lambda \in \Gamma$  with  $f(\lambda) \ge \psi_0, \lambda_i \le 0$

$$f_{\lambda_i} \ge c_0 \sum_{j \ne i} \frac{\partial f}{\partial \lambda_j} + c_1, \qquad i = 1, \dots, n.$$

(E) satisfies

$$\lim_{\substack{\lambda \to 0\\ \lambda \in \Gamma}} f(\lambda) > -\infty,$$

and we assume RHS term satisfies  $\psi(x, u, v)$  is assumed to be  $C^1$ , satisfying

$$\psi_u \ge 0, \quad |\psi| \le \psi_1, \quad |\nabla \psi| \le M, \quad \psi \ge \psi_0$$

for some positive constant  $M, \psi_1, \psi_0$  and

$$\lim_{\substack{\lambda \to \lambda_0 \\ \lambda \in \Gamma}} f(\lambda) < \psi_0, \quad \forall \lambda_0 \in \partial \Gamma.$$

If  $u \in C^3(\overline{B_1})$  is a solution of

$$f(\lambda_1,\ldots,\lambda_n) = \psi(x,u(x),Du(x)),$$
 in  $B_1$ ,

with  $\lambda = \lambda (S_u, P) \in \Gamma, u < 0$  on  $\overline{B_1}$  and  $u(0) = -u_0$ . Then there exists  $C = C (n, f, \psi_1, M, \psi_0, u_0)$  such that  $|\nabla u(0)| \leq C$ .

In order to write our equation into a suitable form that satisfies condition (E) of Theorem 7.2, we see from proof of Theorem 3.5 that there exists  $\delta > 0$  such that

$$\lambda_i \left( D^2 u \right) > -a + b + \delta. \tag{7.2}$$

Thus we translate by setting

$$v := u + \frac{a-b-\delta}{2}|x|^2,$$

such that  $\lambda_i(D^2v) = \lambda_i(D^2u) + (a-b) - \delta > 0$ . Then v satisfies

$$G_{\tau}(\lambda(D^2 v)) := \prod_{i=1}^{n} \frac{\lambda_i + \delta}{\lambda_i + \delta + 2b} = \exp\{\frac{2b}{\sqrt{a^2 + 1}}f(x)\} =: g(x), \quad \forall x \in \mathbb{R}^n.$$
(7.3)

**Theorem 7.3** (Global Gradient Estimate). Let v be a smooth and convex solution of (7.3), where g satisfies conditions (3.2) and (3.3) for some  $g(\infty) = \exp\{\frac{2b}{\sqrt{a^2+1}}f(\infty)\} > 0$ ,  $f(\infty) < 0$ ,  $\alpha \in (0,1)$ ,  $\beta > 1$ ,  $\gamma > 0$ . Suppose in addition that there exists some constant  $C_0$  such that (1.15) holds. Then there for some C > 0 such that (1.14) holds.

*Proof.* For any sufficiently large  $|x| \gg 1$ , we consider the equation in  $B_{\frac{|x|}{2}}(x)$  and let

$$\widetilde{v}(y) := \frac{1}{(\frac{|x|}{2})^2} v(x + \frac{|x|}{2}y), \quad y \in B_1(0).$$

Then we have

$$D^{2}\widetilde{v}(y) = D^{2}v(x + \frac{|x|}{2}y), \quad \forall y \in B_{1}(0),$$

hence it satisfies

$$G_{\tau}(D^2 \widetilde{v}(y)) = g(x + \frac{|x|}{2}y) =: \psi(y), \text{ in } B_1(0).$$

We see that the right hand side term satisfies

$$|\psi| \le \psi_1, \quad \psi \ge \psi_0, \quad \text{and} \quad |\nabla \psi(y)| = \frac{|x|}{2} |\nabla g(x + \frac{|x|}{2}y)| \le M,$$

for some constants  $\psi_1, \psi_0, M$  uniform to all  $|x| \gg 1$  from the asymptotic behavior of  $|\nabla g|$ .

Also, we have

$$||\widetilde{v}||_{L^{\infty}(B_1)} \leq \frac{1}{(\frac{|x|}{2})^2} ||v||_{L^{\infty}(B_{\frac{|x|}{2}}(x))} \leq C_n C_0,$$

which is also uniform with respect to  $x \in \mathbb{R}^n$ .

Now we apply the interior gradient estimate theorem i.e. Theorem 7.2 to see that there exists a uniform (to x) constant C such that

$$|D\widetilde{v}(0)| \le C,$$

where C only relies on  $n, f, \psi_1, M, \psi_0, C_0$ . Hence the constant C is uniform to our choice of  $|x| \gg 1$ .

This is exactly telling us the gradient of u has at most 1-order growth. To be more precise, we have

$$|D\widetilde{v}(0)| = \left|\frac{2}{|x|}Dv(x)\right| \le C$$
, uniform to  $|x| \gg 1$ .

Hence there exists some constant C such that condition (1.14) holds true.

Theorems 1.8 and 1.9 follow directly from Lemma 7.1 and Theorem 7.3.

The scaling strategy used here is exactly the same as reducing "bounded Hessian" to "linear growth gradient" condition. Now we finish this section with proving that under the condition (1.14),  $|D^2u(x)|$  is bounded on entire  $x \in \mathbb{R}^n$  using compactness method as in [29]. To be more precise, we have the following result.

**Theorem 7.4** (Global Hessian Estimate). Let u be a smooth solution of (1.9) with

$$D^2 u > (-a+b)I,$$

where f satisfies conditions (3.2) and (3.3) with 1 replaced by some  $f(\infty) < 0$ ,  $\alpha \in (0,1)$ ,  $\beta > 1$ ,  $\gamma > 0$ . Suppose that there exists some constant  $C_0 > 0$  such that (1.14) holds, then there exists  $C = C(n, \alpha, C_0, f(\infty), \tau)$  such that (1.10) holds.

*Proof.* Based on Lemma 7.1, we learn that under the condition that gradient |Du(x)| has at most linear growth, there exist  $\delta = \delta(n, C_0, \tau)$ ,  $M = M(n, C_0, \tau, u) > 0$  such that

$$|(\widetilde{x} - D\widetilde{u}(\widetilde{x}))| \ge \delta |\widetilde{x}|, \quad \forall |\widetilde{x}| \ge M.$$

Now for sufficiently large  $|x| \gg 1$ , we consider in the ball  $B_{\frac{|x|}{2}}(x)$ , by setting

$$v(y) := \frac{4}{|x|^2}u(x + \frac{|x|}{2}y), \quad \forall y \in B_1(0),$$

then v satisfies

$$||Dv||_{C^0(\overline{B_1})} \le C_0 + 1,$$

and

$$F_{\tau}(\lambda(D^2 v)) = \frac{2b}{\sqrt{a^2 + 1}} f(x + \frac{|x|}{2}y) =: g(y), \quad \forall y \in B_1(0).$$

From our decay condition, we can easily prove that

$$||g||_{C^{\alpha}(B_1)} \le M,$$

uniformly.

Now we only need to prove that

$$|D^2v(0)| \le C(M+1),$$

for some constant M uniform to x, v, g. This is proved through argue by contradiction.

Step 1: Argue by Contradiction,  $L^1$ -convergence result

Suppose the result doesn't hold, then there exist sequences of smooth functions  $\{u_k\}_{k=1}^{\infty}, \{g_k\}_{k=1}^{\infty}$  with

$$D^2 u_k > (-a+b)I, \quad \forall k=1,\cdots,$$

satisfying

$$F_{\tau}(\lambda(D^2 u_k)) = g_k(y), \text{ in } B_1,$$
$$||g_k||_{C^{\alpha}(B_1)} \le M,$$

$$|D^2 u_k(0)| > k \left( ||D u_k||_{L^{\infty}(B_1)} + 1 \right),$$

and having uniform (to k) bounded

$$||Du_k||_{L^{\infty}(B_1)} \le ||Du||_{L^{\infty}(B_1)} \le C_0 + 1.$$

Then using integral by parts, we obtain the following  $W^{2,1}$  estimate. Let  $B_1^m$  denote the m dimensional ball  $B_1^m(0) \subset B_1 = B_1^n(0) \subset \mathbb{R}^n$  for all  $m = 1, \dots, n$  and  $B_1^m(x) := B_1^m + x$  is the ball centered at x. Note that we only need to verify that every component of  $D^2u_k$  belongs to  $L^1$ . Hence we use Fubini theorem and for any positive  $n \times n$  matrix we have

$$||A|| := \sup_{|x|=1} |Ax| \le \sum_{i=1}^{n} \lambda_i(A),$$

it follows that

$$\begin{split} \int_{B_1} ||D^2 u_k + (a-b)I|| \mathrm{d}y &= \int_{B_{\sqrt{1-|x|^2}}^{n-m}(x)} \int_{B_r^m} ||D^2 u_k + (a-b)I|| \mathrm{d}x \mathrm{d}r \\ &\leq C_n \int_{B_{\sqrt{1-|x|^2}}^{n-m}(x)} \int_{B_r^m} (\Delta u_k + C_n(a-b)) \mathrm{d}x \mathrm{d}r \\ &\leq C_n(a-b) + \int_{B_1} \Delta u_k \mathrm{d}x. \end{split}$$

Integral by parts to the formula above and since the solutions  $\{u_k\}$  are smooth up to boundary, hence

$$\begin{aligned} |u_k||_{W^{2,1}(B_1)} &\leq C_n(a-b) + \int_{\partial B_1} Du_k \cdot \vec{n} dx \\ &\leq C_n((a-b) + ||Du_k||_{L^{\infty}(B_1)}) \leq C \end{aligned}$$

Thus by the compact Sobolev Embedding  $u_k \in W^{2,1}(B_1^m) \hookrightarrow W^{1,1}(B_1^m)$ , for almost all  $H^m$ -section of  $B_1$ , meaning that for  $H^{n-m}$ -almost all  $(x_{m+1}, \dots, x_n)$  in  $B_1 \cap \mathbb{R}^{n-m}$ , there exists a function  $u_{\infty} \in W^{1,1}(B_1^m)$  such that

$$Du_k \to Du_\infty$$
 in  $L^1(B_1^m)$ , as  $k \to \infty$ .

#### Step 2: Legendre-Lewy Transform

Using Legendre transform as in (2.1), the equation  $F_{\tau}(\lambda(D^2u_k)) = g_k(y)$  becomes a uniformly elliptic Monge-Ampère type equation (3.10). We assume without loss of generality that  $D\overline{u}(0) = 0$  by modifying a suitable linear function from u, which doesn't affect the equation at all.

As in Section 2,  $\widetilde{x} \in Du(B_1) + (a+b)B_1 \supset B_b(0), \widetilde{u_k}$  satisfies

$$\sum_{i=1}^{n} \ln \widetilde{\lambda}_{i}(D^{2}\widetilde{u_{k}}) = g_{k}\left(\frac{1}{2b}(\widetilde{x} - D\widetilde{u_{k}}(\widetilde{x}))\right), \quad \widetilde{x} \in B_{b}(0),$$
(7.4)

and

$$\widetilde{\lambda}_i(D^2 \widetilde{u_k}) = 1 - \frac{2b}{\lambda_i(D^2 u_k) + a + b} \in (0, 1).$$

From (7.2), there exists  $\delta > 0$  such that

$$0 < \delta < \lambda_i < 1.$$

Hence we can easily see that the equation after Legendre transform (7.4) is uniformly elliptic and concave.

#### *Step 3: Uniform* $C^{2,\alpha}$ *Estimate*

By Lemma 7.1, we see that under the condition (1.14), we still have the important equivalence result (3.14). Due to  $||g_k||_{C^{\alpha}(B_1)} \leq M$ , we obtain that

$$\left\|g_k\left(\frac{1}{2b}(\widetilde{x}-D\widetilde{u}_k(\widetilde{x}))\right)\right\|_{C^{\alpha}(B_b)} \le CM,$$

where  $C = C(n, a, b, C_0)$ . By Schauder estimates of fully nonlinear elliptic equations, Theorem 8.1 and the Remark 3 after it in [9] tells us

$$||\widetilde{u_k}||_{C^{2,\alpha}(B_{\frac{b}{2}}(0))} \le C_{\frac{b}{2}}(0)|$$

for some uniformly to k constant C.

Thus by compact embedding for any  $0 < \varepsilon < \alpha$ ,  $C^{2,\alpha} \hookrightarrow C^{2,\alpha-\varepsilon}$ , there exist a subsequence of  $\widetilde{u_k}$ , still denoted by  $\widetilde{u_k}$ , and  $g_{\infty} \in C^{2,\alpha-\varepsilon}(B_{\frac{b}{4}}(0))$ ,  $\widetilde{u_{\infty}} \in C^{2,\alpha-\varepsilon}(B_{\frac{b}{4}}(0))$  such that

$$\sum_{i=1}^{n} \ln \widetilde{\lambda}_{i}(D^{2}\widetilde{u_{\infty}}(x)) = g_{\infty}\left(\frac{1}{2b}(\widetilde{x} - D\widetilde{u_{\infty}}(\widetilde{x}))\right), \quad \widetilde{x} \in B_{\frac{b}{4}}(0).$$

Due to

$$|D^2 u_k(0)| \to \infty$$
, as  $k \to \infty$ ,

hence we learned from Legendre transform that for some direction  $\gamma \in \partial B_1$ ,  $D_{\gamma\gamma} \widetilde{u_{\infty}} = 0$ , we may name this direction as " $x_1$ " direction for simplicity.

Step 4: Contradicts Constant Rank Theorem

The following constant rank theorem by Caffarelli-Guan-Ma [7] plays an important role in proving the result.

**Theorem 7.5.** Let  $\Psi \subset \mathbb{R}^n$  be an open symmetric domain, assume the operator  $F(D^2u) = f(\lambda(D^2u))$  satisfies  $f \in C^2(\Psi)$  symmetric and

$$f_{\lambda_i}(\lambda) = \frac{\partial f}{\partial \lambda_i}(\lambda) > 0, \quad \forall i = 1, \dots, n, \quad \forall \lambda \in \Psi.$$

Define  $\widetilde{F}(A) = F(A^{-1})$  whenever  $A^{-1} \in \widetilde{\Psi}$ , and we assume  $\widetilde{F}$  is locally convex. Assume u is a  $C^3$  convex solution of the following equation in a domain  $\Omega$  in  $\mathbb{R}^n$ 

$$F\left(D^{2}u(x)\right) = \varphi(x, u(x), Du(x)), \quad \forall x \in \Omega$$

for some  $\varphi \in C^{1,1}(\Omega \times \mathbb{R} \times \mathbb{R}^n)$ . If  $\varphi(x, u, p)$  is concave in  $\Omega \times \mathbb{R}$  for any fixed  $p \in \mathbb{R}^n$ , then the Hessian  $D^2u$  has constant rank in  $\Omega$ .

Classical Hessian equations satisfies this condition, for example, all functions of  $f(\lambda) = \sigma_k^{\frac{1}{k}}$ holds. (Remark 1.7 of [7]). Due to the rank of Hessian matrix is a constant inside, hence  $D_{11}\widetilde{u_{\infty}}(y) \equiv 0$  in a neighborhood of 0. Hence on the hypersurface of  $(x_1, y_1)$ , we have

$$(D_1 \widetilde{w_{\infty}}(y_1, y'), y_1) = (c, y_1) = (x_1, D_1 u_{\infty}(x_1, x') + (a - b)x_1)$$
 near  $(0, 0)$ .

This is impossible as Step 1 tells us  $(x_1, D_1 u_{\infty} (x_1, x') + (a - b)x_1)$  is an  $L^1$  graph (for almost all  $x' \in \mathbb{R}^{n-1}$ ).

# 8 Perturbed Results for $\tau = \frac{\pi}{4}$

In this section, we prove the corresponding results for  $\tau = \frac{\pi}{4}$  by similar strategy as in previous sections. The proof is separated into the following two parts. First, as in Section 3, we prove the asymptotic behavior of  $D^2u$  under additional assumption that Hessian  $D^2u$  is bounded. Then the same argument as in Section 5 and Section 6 shows us the desired result. Second, we reduce the assumption from bounded Hessian to linear growth of gradient and quadratic growth of u as in Section 7.

#### 8.1 Limit of Hessian

Let  $u \in C^4(\mathbb{R}^n)$  be a solution of (1.20) with  $\tau = \frac{\pi}{4}$  and satisfy condition (1.21). We apply the same Legendre transform (2.8) as in Section 2, then we see that  $V(\tilde{x})$  satisfies

$$\Delta V = -\frac{\sqrt{2}}{2}f(DV(\widetilde{x})) =: g(\widetilde{x}), \quad \text{in} \quad \mathbb{R}^n.$$

First, we prove the limit of Hessian at infinity result that corresponds to Theorem 3.5.

**Theorem 8.1.** Assume that f satisfies

$$\limsup_{|x|\to\infty} |x|^{\varepsilon_0+k} |D^k(f(x) - f(\infty))| < \infty, \ \forall k = 0, 1,$$
(8.1)

for some  $\varepsilon_0 > 0$ . Then there exist  $A \in \mathcal{A}(f(\infty), -1), \alpha > 0$ ,  $C(n, f, \varepsilon_0)$  and  $R_2(n, f, \varepsilon_0)$  such that

$$|D^2u(x) - A| \le \frac{C}{|x|^{\alpha}}, \quad \forall |x| \ge R_2.$$

*Proof.* By condition (1.21), we see that

$$\widetilde{\lambda}_i(D^2\widetilde{u}) = \frac{1}{\lambda_i(D^2u) + 1} \ge \frac{1}{M+1} > 0, \quad \forall i = 1, 2, \cdots, n.$$

Also note that  $f \in C^0(\mathbb{R}^n)$  has a finite lower bound. Due to  $\lambda_i > 0$  holds for all  $i = 1, 2, \dots, n$ , we also have the following

$$\widetilde{\lambda}_i < \sum_{j=1}^n \widetilde{\lambda}_j = -\frac{\sqrt{2}}{2} f(DV(\widetilde{x})) \le -\frac{\sqrt{2}}{2} \inf_{\mathbb{R}^n} f < \infty.$$

Combine these two results, we see that

$$0 < \frac{1}{M+1} \le \widetilde{\lambda}_i \le -\frac{\sqrt{2}}{2} \inf_{\mathbb{R}^n} f < \infty.$$
(8.2)

As a consequence, there exists a constant  $\delta = \delta(\tau, \inf_{\mathbb{R}^n} f) > 0$  such that  $D^2 u > -I + \delta I$ . More explicitly, we can pick any  $\delta < -\frac{\sqrt{2}}{\inf_{\mathbb{R}^n} f}$ , the result holds. Now we prove a similar equivalence result as formula (3.14). By the definition of Legendre

transform (2.8), we see that for any  $x^1, x^2 \in \mathbb{R}^n$ ,

$$|\widetilde{x^{1}} - \widetilde{x^{2}}| = |Du(x^{1}) - Du(x^{2}) + (x^{1} - x^{2})| \ge \delta |x^{1} - x^{2}|,$$

and similarly we have

$$|\widetilde{x^{1}} - \widetilde{x^{2}}| = |Du(x^{1}) - Du(x^{2}) + (x^{1} - x^{2})| \le (M+1)|x^{1} - x^{2}|.$$

Combine these two formulas above and we see that there exists some constant  $C_0 = C_0(\inf_{\mathbb{R}^n} f, \tau, M)$ such that (3.14) holds true again.

Based on the equivalence (3.14), we see that

$$\limsup_{|\widetilde{x}| \to \infty} g(\widetilde{x}) = -\frac{\sqrt{2}}{2} f(\infty),$$

and

$$\limsup_{|\widetilde{x}| \to \infty} |\widetilde{x}|^{\varepsilon_0 + 1} \cdot |Dg(\widetilde{x})| < \infty.$$
(8.3)

Let  $w(\tilde{x}) := -\frac{\sqrt{2}f(\infty)}{4n}|x|^2$ , which satisfies  $\Delta w = -\frac{\sqrt{2}}{2}f(\infty)$  in  $\mathbb{R}^n$ . Then we consider the difference  $\overline{v}(\tilde{x}) := V - w$ , satisfying

$$\Delta \overline{v} = -\frac{\sqrt{2}}{2}(f(DV(\widetilde{x})) - f(\infty)) =: \overline{g}(\widetilde{x}), \quad \text{in} \quad \mathbb{R}^n.$$

Let  $K(\tilde{x}) := G_{\Delta}$  be the fundamental solution (or Green's function) of Laplacian, then as in Lemma 4.1, we consider the auxiliary function  $v \in C^2(\mathbb{R}^n)$  (by Schauder estimates and the regularity of  $\overline{q}(\widetilde{x})$ )

$$v(\widetilde{x}) := K * \overline{g}(\widetilde{x}) = \int_{\mathbb{R}^n} K(\widetilde{x} - \widetilde{y}) g(\widetilde{y}) \mathrm{d}\widetilde{y},$$

which satisfies  $\Delta v = \overline{g}$  in  $\mathbb{R}^n$  and  $v \to 0$  as  $|\tilde{x}| \to \infty$ .

As in the proof of Lemma 4.1, we separate the domain into  $E_1, E_2, E_3$  and see that  $D^2v \rightarrow 0$ as  $|\widetilde{x}| \to \infty$ . More explicitly, there exists some constant C such that

$$|D^2 v| \le C|x|^{1-(1+\varepsilon_0)} = C|x|^{-\varepsilon_0}, \quad \forall x \in \mathbb{R}^n.$$
(8.4)

Also note that  $v \in C^2(\mathbb{R}^n)$ , hence  $D^2v$  is bounded on entire  $\mathbb{R}^n$ Hence we see that

$$\Delta(V - w - v) = 0, \quad \text{in} \quad \mathbb{R}^n,$$

and due to  $D^2V$ ,  $D^2w$ ,  $D^2v$  are all bounded on  $\mathbb{R}^n$ , hence so is  $D^2(V-w-v)$ . Applying Theorem 2.1, we see that the limit of  $D^2(V-w-v)$  at infinity exists. (Or we can take twice derivatives to both sides and apply Liouville theorem). More explicitly, picking k = 2 in formula (1.6), there exists

$$\widetilde{A} \in \{\widetilde{A} \in \operatorname{Sym}(n) : \sum_{i=1}^n \lambda_i(\widetilde{A}) = 0, \frac{1}{M+1}I + \frac{\sqrt{2}f(\infty)}{2n}I \le \widetilde{A} \le \frac{\sqrt{2}}{2}\inf_{\mathbb{R}^n} fI + \frac{\sqrt{2}f(\infty)}{2n}I\},$$

such that

$$\limsup_{|\widetilde{x}| \to \infty} |\widetilde{x}|^n \left| D^2 (V - w - v) - \widetilde{A} \right| < \infty.$$
(8.5)

Combine formula (8.4), (8.5) and the fact that  $D^2w \equiv -\frac{\sqrt{2}f(\infty)}{2n}I$ , it follows immediately that  $\widetilde{B} := \widetilde{A} - \frac{\sqrt{2}f(\infty)}{2n}I$  belongs to the set of

$$\{\widetilde{B}: \sum_{i=1}^n \lambda_i(\widetilde{B}) = -\frac{\sqrt{2}}{2}f(\infty), \frac{1}{M+1}I \le \widetilde{B} \le \frac{\sqrt{2}}{2}\inf_{\mathbb{R}^n} fI\},\$$

such that

$$\limsup_{|\widetilde{x}|\to\infty} |\widetilde{x}|^{\min\{\varepsilon_0,n\}} |D^2 V - \widetilde{B}| < \infty.$$

By the definition of Legendre transform (2.8) and the equivalence (3.14), we pick  $\alpha := \min\{n, \varepsilon_0\}$ , the desired result follows immediately.

*Proof of Theorems 1.14 and 1.15.* As in the calculus of Theorem 8.1, formula (8.2) and the Legendre transform (2.8),

$$\frac{\sqrt{2}}{\inf_{\mathbb{R}^n} f} \le \lambda_i(D^2 u) \le M.$$

By direct calculus, we have

$$\frac{\partial}{\partial\lambda_i}\left(\sum_{i=1}^n \frac{1}{\lambda_i+1}\right) = -\frac{1}{(1+\lambda_i)^2} \in \left(-\frac{1}{(\frac{\sqrt{2}}{\inf_{\mathbb{R}^n} f}+1)^2}, -\frac{1}{(1+M)^2}\right),$$

meaning that the equation (1.20) is uniformly elliptic. Also, we have

$$\frac{\partial^2}{\partial \lambda_i^2} \left( \sum_{i=1}^n \frac{1}{\lambda_i + 1} \right) = \frac{2}{(1 + \lambda_i)^3} > 0,$$

hence the equation is also convex. Consider

$$-F_{\tau}(D^2u) = -f(x), \quad \text{in} \quad \mathbb{R}^n,$$

then apply the same argument as in the proof of Theorems 1.5 and 1.6, similar results also hold true.  $\hfill\square$ 

#### 8.2 Interior Estimates

In this part, we reduce the strong assumption (1.21) into weaker ones. Similar to the proof of Lemma 7.1, it holds true under this situation as well.

Now we consider solution of (1.20) with  $\tau = \frac{\pi}{4}$ ,  $D^2 u > -I$  and we assume that f satisfies condition (8.1) for some  $\varepsilon_0 > 0$ ,  $f(\infty) < 0$ . The proof is an explicit copy, we only used the ellipticity and concavity of equation. Hence Theorems 1.16 and 1.17 follow immediately.

As in the proof of Theorem 8.1 or Lemma 7.2, we see that there exists a  $\delta = \delta(\tau, \inf_{\mathbb{R}^n} f) > 0$  such that

$$\lambda_i(D^2 u) > -1 + \delta.$$

Then we set

$$v:=u+\frac{1-\delta}{2}|x|^2,\quad \text{then}\quad \lambda_i(D^2v)=\lambda_i(D^2u)+1-\delta>0.$$

And we see that v satisfies

$$\sum_{i=1}^n \frac{1}{\lambda_i(D^2v)+\delta} = -\frac{\sqrt{2}}{2}f(x), \quad \text{in} \quad \mathbb{R}^n.$$

By verifying the conditions in Theorem 7.2, we do scaling as in the proof of Theorem 7.3, Theorem 1.16 follows from similar argument as in Theorem 1.8.

Note that the equation after Legendre-Lewy transform becomes Laplacian operator, standard Schauder estimates still hold true. The estimates to prove Theorem 1.17 follows by similar compactness argument as in Theorem 1.9.

#### References

- [1] Robert A. Adams and John J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [2] Jiguang Bao, Jingyi Chen, Bo Guan, and Min Ji. Liouville property and regularity of a Hessian quotient equation. Amer. J. Math., 125(2):301–316, 2003.
- [3] Jiguang Bao, Haigang Li, and Lei Zhang. Monge-Ampère equation on exterior domains. *Calc. Var. Partial Differential Equations*, 52(1-2):39–63, 2015.
- [4] L. Caffarelli and Yan Yan Li. A Liouville theorem for solutions of the Monge-Ampère equation with periodic data. Ann. Inst. H. Poincaré Anal. Non Linéaire, 21(1):97–120, 2004.
- [5] L. Caffarelli and Yanyan Li. An extension to a theorem of Jörgens, Calabi, and Pogorelov. Comm. Pure Appl. Math., 56(5):549–583, 2003.
- [6] L. Caffarelli, L. Nirenberg, and J. Spruck. The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian. *Acta Math.*, 155(3-4):261–301, 1985.
- [7] Luis Caffarelli, Pengfei Guan, and Xi-Nan Ma. A constant rank theorem for solutions of fully nonlinear elliptic equations. *Comm. Pure Appl. Math.*, 60(12):1769–1791, 2007.
- [8] Luis A. Caffarelli. Interior W<sup>2,p</sup> estimates for solutions of the Monge-Ampère equation. Ann. of Math. (2), 131(1):135–150, 1990.
- [9] Luis A. Caffarelli and Xavier Cabré. Fully nonlinear elliptic equations, volume 43 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1995.
- [10] Eugenio Calabi. Improper affine hyperspheres of convex type and a generalization of a theorem by K. Jörgens. *Michigan Math. J.*, 5:105–126, 1958.

- [11] Li Chen and Ni Xiang. Rigidity theorems for the entire solutions of 2-Hessian equation. J. Differential Equations, 267(9):5202–5219, 2019.
- [12] Shiu Yuen Cheng and Shing-Tung Yau. Complete affine hypersurfaces. I. The completeness of affine metrics. *Comm. Pure Appl. Math.*, 39(6):839–866, 1986.
- [13] Miguel de Guzmán. Differentiation of integrals in R<sup>n</sup>. Lecture Notes in Mathematics, Vol. 481. Springer-Verlag, Berlin-New York, 1975. With appendices by Antonio Córdoba, and Robert Fefferman, and two by Roberto Moriyón.
- [14] Sł awomir Dinew and Sł awomir Koł odziej. Liouville and Calabi-Yau type theorems for complex Hessian equations. Amer. J. Math., 139(2):403–415, 2017.
- [15] L. Ferrer, A. Martínez, and F. Milán. An extension of a theorem by K. Jörgens and a maximum principle at infinity for parabolic affine spheres. *Math. Z.*, 230(3):471–486, 1999.
- [16] Harley Flanders. On certain functions with positive definite Hessian. Ann. of Math. (2), 71:153–156, 1960.
- [17] D. Gilbarg and James Serrin. On isolated singularities of solutions of second order elliptic differential equations. J. Analyse Math., 4:309–340, 1955/56.
- [18] David Gilbarg and Neil S. Trudinger. Elliptic partial differential equations of second order, volume 224 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, second edition, 1983.
- [19] Cristian E. Gutiérrez and Qingbo Huang. A generalization of a theorem by Calabi to the parabolic Monge-Ampère equation. *Indiana Univ. Math. J.*, 47(4):1459–1480, 1998.
- [20] Rongli Huang, Qianzhong Ou, and Wenlong Wang. On the entire self-shrinking solutions to lagrangian mean curvature flow ii, 2019.
- [21] H. Hueber and M. Sieveking. Uniform bounds for quotients of Green functions on C<sup>1,1</sup>-domains. *Ann. Inst. Fourier (Grenoble)*, 32(1):vi, 105–117, 1982.
- [22] Huai-Yu Jian and Xu-Jia Wang. Continuity estimates for the Monge-Ampère equation. *SIAM J. Math. Anal.*, 39(2):608–626, 2007.
- [23] Konrad Jörgens. Über die Lösungen der Differentialgleichung  $rt s^2 = 1$ . Math. Ann., 127:130–134, 1954.
- [24] J. Jost and Y. L. Xin. Some aspects of the global geometry of entire space-like submanifolds. volume 40, pages 233–245. 2001. Dedicated to Shiing-Shen Chern on his 90th birthday.
- [25] An-Min Li and Li Sheng. A Liouville Theorem on the PDE  $det(f_{i\bar{j}}) = 1$ . arXiv e-prints, page arXiv:1809.00824, Sep 2018.
- [26] Dongsheng Li, Zhisu Li, and Yu Yuan. A Bernstein problem for special Lagrangian equations in exterior domains. arXiv e-prints, page arXiv:1709.04727, Sep 2017.
- [27] Ming Li, Changyu Ren, and Zhizhang Wang. An interior estimate for convex solutions and a rigidity theorem. J. Funct. Anal., 270(7):2691–2714, 2016.
- [28] Yan Yan Li. Interior gradient estimates for solutions of certain fully nonlinear elliptic equations. J. Differential Equations, 90(1):172–185, 1991.
- [29] Matt McGonagle, Chong Song, and Yu Yuan. Hessian estimates for convex solutions to quadratic Hessian equation. Ann. Inst. H. Poincaré Anal. Non Linéaire, 36(2):451–454, 2019.
- [30] Saori Nakamori and Kazuhiro Takimoto. A Bernstein type theorem for parabolic k-Hessian equations. Nonlinear Anal., 117:211–220, 2015.
- [31] Tu A. Nguyen and Yu Yuan. A priori estimates for Lagrangian mean curvature flows. *Int. Math. Res. Not. IMRN*, (19):4376–4383, 2011.
- [32] Yehuda Pinchover. On the equivalence of Green functions of second order elliptic equations in  $\mathbb{R}^n$ . *Differential Integral Equations*, 5(3):481–493, 1992.
- [33] A. V. Pogorelov. On the improper convex affine hyperspheres. *Geometriae Dedicata*, 1(1):33–46, 1972.
- [34] Eduardo V. Teixeira and Lei Zhang. Global Monge-Ampére equation with asymptotically periodic data. *Indiana Univ. Math. J.*, 65(2):399–422, 2016.

- [35] Mao-Pei Tsui and Mu-Tao Wang. A Bernstein type result for special Lagrangian submanifolds. *Math. Res. Lett.*, 9(4):529–535, 2002.
- [36] Bo Wang and Jiguang Bao. Asymptotic behavior on a kind of parabolic Monge-Ampère equation. J. Differential Equations, 259(1):344–370, 2015.
- [37] Chong Wang, Rongli Huang, and Jiguang Bao. On the second boundary value problem for lagrangian mean curvature equation, 2018.
- [38] Micah Warren. Calibrations associated to Monge-Ampère equations. *Trans. Amer. Math. Soc.*, 362(8):3947–3962, 2010.
- [39] Micah Warren. Nonpolynomial entire solutions to  $\sigma_k$  equations. Comm. Partial Differential Equations, 41(5):848–853, 2016.
- [40] Micah Warren and Yu Yuan. A Liouville type theorem for special Lagrangian equations with constraints. Comm. Partial Differential Equations, 33(4-6):922–932, 2008.
- [41] Jingang Xiong and Jiguang Bao. On Jörgens, Calabi, and Pogorelov type theorem and isolated singularities of parabolic Monge-Ampère equations. J. Differential Equations, 250(1):367–385, 2011.
- [42] Yu Yuan. A Bernstein problem for special Lagrangian equations. *Invent. Math.*, 150(1):117–125, 2002.
- [43] Yu Yuan. Global solutions to special Lagrangian equations. Proc. Amer. Math. Soc., 134(5):1355– 1358, 2006.
- [44] Wei Zhang and Jiguang Bao. A Calabi theorem for solutions to the parabolic Monge-Ampère equation with periodic data. Ann. Inst. H. Poincaré Anal. Non Linéaire, 35(5):1143–1173, 2018.
- [45] Wei Zhang, Jiguang Bao, and Bo Wang. An extension of Jörgens-Calabi-Pogorelov theorem to parabolic Monge-Ampère equation. *Calc. Var. Partial Differential Equations*, 57(3):Art. 90, 36, 2018.
- [46] William P. Ziemer. *Weakly differentiable functions*, volume 120 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1989. Sobolev spaces and functions of bounded variation.