

SCALED RELATIVE GRAPH OF NORMAL MATRICES

XINMENG HUANG*, ERNEST K. RYU†, AND WOTAO YIN‡

Abstract. The Scaled Relative Graph (SRG) by Ryu, Hannah, and Yin (arXiv:1902.09788, 2019) is a geometric tool that maps the action of a multi-valued nonlinear operator onto the 2D plane, used to analyze the convergence of a wide range of iterative methods. As the SRG includes the spectrum for linear operators, we can view the SRG as a generalization of the spectrum to multi-valued nonlinear operators. In this work, we further study the SRG of linear operators and characterize the SRG of block-diagonal and normal matrices.

Key words. scaled relative graph, non-Euclidean geometry, hyperbolic geometry, normal matrix

AMS subject classifications. 47H05, 47H09, 51M04, 52A55, 90C25

1. Introduction. The Scaled Relative Graph (SRG), recently proposed by Ryu, Hannah, and Yin [12], is a geometric tool that maps the action of a multi-valued nonlinear operator onto the extended complex plane, analogous to how the spectrum maps the action of a linear operator to the complex plane. The SRG can be used to analyze convergence of a wide range of iterative methods expressed as fixed-point iterations.

Scaled relative graph. For a matrix $A \in \mathbb{R}^{n \times n}$, define $z_A: \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{C}$ with

$$z_A(x) = \frac{\|Ax\|}{\|x\|} \exp[i\angle(Ax, x)],$$

where

$$\angle(a, b) = \begin{cases} \arccos\left(\frac{a^T b}{\|a\|\|b\|}\right) & \text{if } a \neq 0, b \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

denotes the angle in $[0, \pi]$ between a and b . The SRG of a matrix $A \in \mathbb{R}^{n \times n}$ is

$$\mathcal{G}(A) = \{z_A(x), \overline{z_A(x)} : x \in \mathbb{R}^n, x \neq 0\}.$$

This definition of the SRG, specific to (single-valued) linear operators, coincides with the more general definition for nonlinear multi-valued operators provided in [12]. Ryu, Hannah, and Yin showed the SRG generalizes spectrum in the following sense.

FACT 1 (Theorem 3.1 of [12]). *If $A \in \mathbb{R}^{n \times n}$ and $n = 1$ or $n \geq 3$, then $\Lambda(A) \subseteq \mathcal{G}(A)$.*

2D geometric illustrations have been used by Eckstein and Bertsekas [4, 5], Giselsson [7, 6], Banjac and Goulart [1], and Giselsson and Moursi [8] to qualitatively understand convergence of optimization algorithms. Ryu, Hannah, and Yin presented the SRG as a rigorous formulation of such illustrations [12].

Contributions. Prior work [12, 9] focused on the SRG of *nonlinear* multi-valued operators. For linear operators, Ryu, Hannah, and Yin [12] established $\mathcal{G}(A)$ includes $\Lambda(A)$, as stated in Fact 1, but did not characterize when and how $\mathcal{G}(A)$ enlarges $\Lambda(A)$. In this work, we further study the SRG of linear operators. In particular, we fully characterize the SRG of block-diagonal and normal matrices as a certain polygon in hyperbolic (non-Euclidean) geometry, under the Poincaré half-plane model.

*School of Mathematical Sciences, University of Science and Technology of China (hxm8888@mail.ustc.edu.cn)

†Department of Mathematics, University of California, Los Angeles (eryu@math.ucla.edu)

‡Department of Mathematics, University of California, Los Angeles (wotaoyin@math.ucla.edu)

Preliminaries. Let $A \in \mathbb{R}^{n \times n}$. Write $\Lambda(A)$ for the spectrum, the set of eigenvalues, of A . A is *normal* if $A^T A = A A^T$. Given matrices A_1, \dots, A_m , write $\text{Diag}(A_1, \dots, A_m)$ for the block-diagonal matrix with m blocks. For $z \in \mathbb{C}$, write \bar{z} for its complex conjugate. For a set $S \subseteq \mathbb{C}$, write $S^+ = \{z \in S \mid \text{Im } z \geq 0\}$. In particular, write $\mathbb{C}^+ = \{z \in \mathbb{C} \mid \text{Im } z \geq 0\}$ and $\mathcal{G}^+(A) = \{z_A(x) : x \in \mathbb{R}^n, x \neq 0\}$. Note $z_A(x) \in \mathbb{C}^+$ for all nonzero $x \in \mathbb{R}^n$. For $z_1, z_2 \in \mathbb{C}$, define

$$[z_1, z_2] = \{\theta z_1 + (1 - \theta)z_2 : \theta \in [0, 1]\},$$

i.e., $[z_1, z_2]$ is the line segment connecting z_1 and z_2 .

2. Arc-edge polygon and arc-convexity. Consider points $z_1, z_2 \in \mathbb{C}^+$. If $\text{Re } z_1 \neq \text{Re } z_2$, let $\text{Circ}(z_1, z_2)$ be the circle in \mathbb{C} through z_1 and z_2 with the center on the real axis. We can construct $\text{Circ}(z_1, z_2)$ by finding the center as the intersection of the perpendicular bisector of $[z_1, z_2]$ and the real axis. If $\text{Re } z_1 = \text{Re } z_2$ but $z_1 \neq z_2$, let $\text{Circ}(z_1, z_2)$ be the line extending $[z_1, z_2]$. If $z_1 = z_2$, then $\text{Circ}(z_1, z_2)$ is undefined. If $\text{Re } z_1 \neq \text{Re } z_2$, let $\text{Arc}_{\min}(z_1, z_2) \subseteq \mathbb{C}^+$ be the arc of $\text{Circ}(z_1, z_2)$ between z_1 and z_2 in the upper-half plane. (If $\text{Im } z_1 > 0$ or $\text{Im } z_2 > 0$, then $\text{Arc}_{\min}(z_1, z_2) \subseteq \mathbb{C}^+$ is the minor arc of $\text{Circ}(z_1, z_2)$ between z_1 and z_2 . If $\text{Im } z_1 = \text{Im } z_2 = 0$, then $\text{Arc}_{\min}(z_1, z_2)$ is a semicircle in \mathbb{C}^+ .) If $\text{Re } z_1 = \text{Re } z_2$ but $z_1 \neq z_2$, let $\text{Arc}_{\min}(z_1, z_2) = [z_1, z_2]$. If $z_1 = z_2$, then $\text{Arc}_{\min}(z_1, z_2) = \{z_1\}$. For $z_1, z_2 \in \mathbb{C}^+$ such that $\text{Re } z_1 \neq \text{Re } z_2$, let $\text{Disk}(z_1, z_2)$ and $\text{Disk}^\circ(z_1, z_2)$ respectively be the closed and open disks enclosed by $\text{Circ}(z_1, z_2)$. Figure 1 illustrates these definitions.




Fig. 1: Illustration of $\text{Circ}(z_1, z_2)$ and $\text{Arc}_{\min}(z_1, z_2)$.

For $m \geq 1$ and $z_1, \dots, z_m \in \mathbb{C}^+$, we call $\text{Poly}(z_1, z_2, \dots, z_m)$ an *arc-edge polygon* and define it as follows. For $m = 1$, let $\text{Poly}(z_1) = \{z_1\}$. For $m \geq 2$, let

$$S = \bigcup_{1 \leq i, j \leq m} \text{Arc}_{\min}(z_i, z_j)$$

and

$$\text{Poly}(z_1, \dots, z_m) = S \cup \{\text{region enclosed by } S\}.$$

Figure 2 illustrates this definition. Note $\text{Poly}(z_1, z_2) = \text{Arc}_{\min}(z_1, z_2)$. The “region enclosed by S ” is the union of all regions enclosed by non-self-intersecting continuous loops (Jordan curves) within S . Since S is a connected set, we can alternatively define $\text{Poly}(z_1, \dots, z_m)$ as the smallest simply connected set containing S .

This construction of Arc_{\min} gives rise to the classical *Poincaré half-plane model* of hyperbolic (non-Euclidean) geometry, where a $\text{Arc}_{\min}(z_1, z_2)$ and $\text{Circ}(z_1, z_2) \cap \mathbb{C}^+$ are, respectively, the “line segment” between z_1 and z_2 and the “line” through z_1 and z_2 in the




Fig. 2: The shaded region illustrates the arc-edge polygon $\text{Poly}(z_1, z_2, z_3, z_4)$ for $z_1 = 1 + i$, $z_2 = 2 + 3i$, $z_3 = 4 + 2i$, and $z_4 = 4 + i$. The solid arcs illustrate $\text{Arc}_{\min}(z_i, z_j)$ and the dashed circles illustrate $\text{Circ}(z_i, z_j)$ for $i, j = 1, \dots, m$.




Fig. 3: Illustration of $f \circ g$ and Lemma 2.1. The one-to-one map $f \circ g$ of (2.1) maps $\text{Poly}(z_1, \dots, z_7)$ (a hyperbolic polygon) into a Euclidean polygon. We denote the mapped points as $w_i = f(g(z_i))$ for $i = 1, \dots, 7$. The equivalent Euclidean geometry tells us that $\text{Poly}(z_1, \dots, z_7)$ is “convex” and can be enclosed by the curve through $z_1 \rightarrow z_2 \rightarrow z_3 \rightarrow z_4 \rightarrow z_6 \rightarrow z_1$. Note that z_5 and z_7 are not necessary in the description of the boundary.

hyperbolic space [3, 11]. The *Beltrami–Klein model* maps the Poincaré half-plane model onto the unit disk and Arc_{\min} to straight line segments [10, 2]. Specifically, the one-to-one map

$$(2.1) \quad f \circ g: \mathbb{C}^+ \rightarrow \{z \in \mathbb{C} : |z| \leq 1, z \neq 1\}$$

defined by

$$f(z) = \frac{2z}{1 + |z|^2}, \quad g(z) = \frac{z - i}{z + i}$$

maps the Poincaré half-plane model to the Beltrami–Klein model while mapping hyperbolic line segments Arc_{\min} into Euclidean straight line segments. The Beltrami–Klein model demonstrates that any qualitative statement about convexity in the Euclidean plane is equivalent to an analogous statement in the Poincaré half-plane model. See Figure 3.

LEMMA 2.1. *Let $z_1, \dots, z_m \in \mathbb{C}^+$ and $m \geq 1$. Then $\text{Poly}(z_1, \dots, z_m)$ is “convex” in the following non-Euclidean sense:*

$$w_1, w_2 \in \text{Poly}(z_1, \dots, z_m) \quad \Rightarrow \quad \text{Arc}_{\min}(w_1, w_2) \subseteq \text{Poly}(z_1, \dots, z_m).$$

If $\text{Poly}(z_1, \dots, z_m)$ has an interior, then there is $\{\zeta_1, \dots, \zeta_q\} \subseteq \{z_1, \dots, z_m\}$ such that

$$\text{Arc}_{\min}(\zeta_1, \zeta_2) \cup \text{Arc}_{\min}(\zeta_2, \zeta_3) \cup \dots \cup \text{Arc}_{\min}(\zeta_{q-1}, \zeta_q) \cup \text{Arc}_{\min}(\zeta_q, \zeta_1)$$

is a Jordan curve, and the region the curve encloses is $\text{Poly}(z_1, \dots, z_m)$.

Proof. Let w_1, \dots, w_m be in the unit complex disk. Consider the construction

$$\tilde{S} = \bigcup_{1 \leq i, j \leq m} [w_i, w_j]$$

and

$$\widetilde{\text{Poly}}(w_1, \dots, w_m) = \tilde{S} \cup \{\text{region enclosed by } \tilde{S}\}.$$

This is the (Euclidean) 2D polyhedron given as the convex hull of w_1, \dots, w_m . The Euclidean convex hull has the properties analogous to those in the Lemma statement, and we use the map $(f \circ g)^{-1}$, where $f \circ g$ is as given by (2.1) to map the properties to our setup. \square

3. SRGs of block-diagonal matrices. We characterize the SRG of block-diagonal matrices as follows.

THEOREM 3.1. *Let A_1, \dots, A_m be square matrices, where $m \geq 1$. Then*

$$\mathcal{G}^+(\text{Diag}(A_1, \dots, A_m)) = \bigcup_{\substack{z_i \in \mathcal{G}^+(A_i) \\ i=1, \dots, m}} \text{Poly}(z_1, \dots, z_m).$$

Proof. When $m = 1$, there is nothing to show. Assume $m \geq 2$.

Step 1. Let $A_i \in \mathbb{R}^{n_i \times n_i}$ and $u_i \in \mathbb{R}^{n_i}$ for $i = 1, \dots, m$. We use the notation $n = n_1 + \dots + n_m$,

$$\mathbf{u} = \begin{bmatrix} u_1 \\ \vdots \\ u_m \end{bmatrix} \in \mathbb{R}^n, \quad \mathbf{u}_i = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ u_i \\ 0 \\ \vdots \\ 0 \end{bmatrix} \in \mathbb{R}^n \quad \text{for } i = 1, \dots, m,$$

and $\mathbf{A} = \text{Diag}(A_1, \dots, A_m) \in \mathbb{R}^{n \times n}$. Then we have

$$\begin{aligned}
 \mathcal{G}^+(\text{Diag}(A_1, \dots, A_m)) &= \bigcup_{\mathbf{u} \in \mathbb{R}^n \setminus \{0\}} z_{\mathbf{A}}(\mathbf{u}) \\
 (3.1) \qquad \qquad \qquad &= \bigcup_{\substack{u_i \in \mathbb{R}^{n_i}, u_i \neq 0 \\ i=1, \dots, m}} z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\})
 \end{aligned}$$

and

$$\begin{aligned}
 \bigcup_{\substack{u_i \in \mathbb{R}^{n_i}, u_i \neq 0 \\ i=1, \dots, m}} \text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)) &= \bigcup_{\substack{u_i \in \mathbb{R}^{n_i}, u_i \neq 0 \\ i=1, \dots, m}} \text{Poly}(z_{A_1}(u_1), \dots, z_{A_m}(u_m)) \\
 (3.2) \qquad \qquad \qquad &= \bigcup_{\substack{z_i \in \mathcal{G}^+(A_i) \\ i=1, \dots, m}} \text{Poly}(z_1, \dots, z_m).
 \end{aligned}$$

To clarify, \mathbf{u}_i depends on u_i for $i = 1, \dots, m$. In the following, we show

$$(3.3) \qquad z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\}) = \text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m))$$

for all \mathbf{u}_i given by $u_i \neq 0$ for $i = 1, \dots, m$. Once (3.3) is proved, (3.1) and (3.2) are equivalent and the proof is complete.

Step 2. We show the following intermediate result: for all nonzero $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ such that

$$(3.4) \qquad \langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{A}\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{A}\mathbf{v} \rangle = \langle \mathbf{A}\mathbf{u}, \mathbf{A}\mathbf{v} \rangle = 0,$$

we have

$$(3.5) \qquad z_{\mathbf{A}}(\text{span}(\mathbf{u}, \mathbf{v}) \setminus \{0\}) = \text{Arc}_{\min}(z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v})).$$

First, consider the case $\text{Re } z_{\mathbf{A}}(\mathbf{u}) \neq \text{Re } z_{\mathbf{A}}(\mathbf{v})$. Let the circle $\text{Circ}(z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v}))$ be centered at $(t, 0)$ with $t \in \mathbb{R}$ and radius $r \geq 0$. Then $z_{\mathbf{A}}(\mathbf{u})$ and $z_{\mathbf{A}}(\mathbf{v})$ satisfy

$$\begin{aligned}
 (\text{Re } z_{\mathbf{A}}(\mathbf{u}) - t)^2 + (\text{Im } z_{\mathbf{A}}(\mathbf{u}))^2 &= r^2 \\
 (\text{Re } z_{\mathbf{A}}(\mathbf{v}) - t)^2 + (\text{Im } z_{\mathbf{A}}(\mathbf{v}))^2 &= r^2.
 \end{aligned}$$

This is equivalent to

$$\begin{aligned}
 \langle \mathbf{A}\mathbf{u}, \mathbf{A}\mathbf{u} \rangle - 2t\langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle + (t^2 - r^2)\langle \mathbf{u}, \mathbf{u} \rangle &= 0 \\
 \langle \mathbf{A}\mathbf{v}, \mathbf{A}\mathbf{v} \rangle - 2t\langle \mathbf{A}\mathbf{v}, \mathbf{v} \rangle + (t^2 - r^2)\langle \mathbf{v}, \mathbf{v} \rangle &= 0.
 \end{aligned}$$

Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $\mathbf{w} = \alpha_1\mathbf{u} + \alpha_2\mathbf{v}$. Assume $\mathbf{w} \neq 0$. Using (3.4) and basic calculations, we have

$$\langle \mathbf{A}\mathbf{w}, \mathbf{A}\mathbf{w} \rangle - 2t\langle \mathbf{A}\mathbf{w}, \mathbf{w} \rangle + (t^2 - r^2)\langle \mathbf{w}, \mathbf{w} \rangle = 0,$$

and this is equivalent to

$$(\text{Re } z_{\mathbf{A}}(\mathbf{w}) - t)^2 + (\text{Im } z_{\mathbf{A}}(\mathbf{w}))^2 = r^2.$$

Therefore

$$z_{\mathbf{A}}(\mathbf{w}) = z_{\mathbf{A}}(\alpha_1\mathbf{u} + \alpha_2\mathbf{v}) \in \text{Circ}(z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v})).$$

Notice that

$$\operatorname{Re} z_{\mathbf{A}}(\mathbf{w}) = \frac{\langle \mathbf{A}\mathbf{w}, \mathbf{w} \rangle}{\langle \mathbf{w}, \mathbf{w} \rangle} = \frac{\alpha_1^2 \langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle + \alpha_2^2 \langle \mathbf{A}\mathbf{v}, \mathbf{v} \rangle}{\alpha_1^2 \langle \mathbf{u}, \mathbf{u} \rangle + \alpha_2^2 \langle \mathbf{v}, \mathbf{v} \rangle}$$

fills the interval $[\operatorname{Re} z_{\mathbf{A}}(\mathbf{u}), \operatorname{Re} z_{\mathbf{A}}(\mathbf{v})]$ as α_1 and α_2 varies. So we have

$$\bigcup_{\substack{\alpha_1, \alpha_2 \in \mathbb{R} \\ \alpha_1 \mathbf{u} + \alpha_2 \mathbf{v} \neq \mathbf{0}}} z_{\mathbf{A}}(\alpha_1 \mathbf{u} + \alpha_2 \mathbf{v}) = \operatorname{Arc}_{\min}(z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v}))$$

and we conclude (3.5).

Next, consider the case $\operatorname{Re} z_{\mathbf{A}}(\mathbf{u}) = \operatorname{Re} z_{\mathbf{A}}(\mathbf{v})$. Note that

$$\operatorname{Re} z_{\mathbf{A}}(\mathbf{u}) = \frac{\langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle}{\langle \mathbf{u}, \mathbf{u} \rangle}, \quad \operatorname{Re} z_{\mathbf{A}}(\mathbf{v}) = \frac{\langle \mathbf{A}\mathbf{v}, \mathbf{v} \rangle}{\langle \mathbf{v}, \mathbf{v} \rangle}.$$

Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $\mathbf{w} = \alpha_1 \mathbf{u} + \alpha_2 \mathbf{v}$. Assume $\mathbf{w} \neq \mathbf{0}$. Using (3.4) and basic calculations, we have

$$\operatorname{Re} z_{\mathbf{A}}(\mathbf{w}) = \frac{\langle \mathbf{A}\mathbf{w}, \mathbf{w} \rangle}{\langle \mathbf{w}, \mathbf{w} \rangle} = \operatorname{Re} z_{\mathbf{A}}(\mathbf{u}) = \operatorname{Re} z_{\mathbf{A}}(\mathbf{v}).$$

Notice that

$$|z_{\mathbf{A}}(\mathbf{w})|^2 = \frac{\langle \mathbf{A}\mathbf{w}, \mathbf{A}\mathbf{w} \rangle}{\langle \mathbf{w}, \mathbf{w} \rangle} = \frac{\alpha_1^2 \langle \mathbf{A}\mathbf{u}, \mathbf{A}\mathbf{u} \rangle + \alpha_2^2 \langle \mathbf{A}\mathbf{v}, \mathbf{A}\mathbf{v} \rangle}{\alpha_1^2 \langle \mathbf{u}, \mathbf{u} \rangle + \alpha_2^2 \langle \mathbf{v}, \mathbf{v} \rangle}$$

fills the interval $[|z_{\mathbf{A}}(\mathbf{u})|^2, |z_{\mathbf{A}}(\mathbf{v})|^2]$ as α_1 and α_2 varies. So $\operatorname{Im} z_{\mathbf{A}}(\mathbf{w})$ fills the interval $[\operatorname{Im} z_{\mathbf{A}}(\mathbf{u}), \operatorname{Im} z_{\mathbf{A}}(\mathbf{v})]$ as α_1 and α_2 varies, and we conclude

$$z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}, \mathbf{v}) \setminus \{0\}) = [z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v})] = \operatorname{Arc}_{\min}(z_{\mathbf{A}}(\mathbf{u}), z_{\mathbf{A}}(\mathbf{v})).$$

Step 3. We show

$$(3.6) \quad z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\}) \subseteq \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m))$$

by induction. Clearly

$$z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1) \setminus \{0\}) = \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1)).$$

Now assume (3.6) holds for $m - 1$. By (3.5), we have

$$z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\}) = \bigcup_{\zeta \in z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_{m-1}) \setminus \{0\})} \operatorname{Arc}_{\min}(\zeta, z_{\mathbf{A}}(\mathbf{u}_m)).$$

By the induction hypothesis, $\zeta \in z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_{m-1}) \setminus \{0\})$, implies

$$\zeta \in \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_{m-1})) \subseteq \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)).$$

By construction,

$$z_{\mathbf{A}}(\mathbf{u}_m) \in \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)).$$

“Convexity” of Lemma 2.1 implies

$$\bigcup_{\zeta \in z_{\mathbf{A}}(\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_{m-1}) \setminus \{0\})} \operatorname{Arc}_{\min}(\zeta, z_{\mathbf{A}}(\mathbf{u}_m)) \subseteq \operatorname{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)),$$

and we conclude (3.6).

Step 4. We show

$$(3.7) \quad z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\}) \supseteq \text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)).$$

First, consider the case where $\text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m))$ has no interior. In 2D Euclidean geometry, a polygon has no interior when it is a single line segment or a point. The Beltrami–Klein model provides us with an equivalent statement in hyperbolic geometry: the “polygon” can be expressed as $\text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)) = \text{Arc}_{\min}(z_{\mathbf{A}}(\boldsymbol{\mu}_1), z_{\mathbf{A}}(\boldsymbol{\mu}_2))$ where $\boldsymbol{\mu}_1, \boldsymbol{\mu}_2 \in \{\mathbf{u}_1, \dots, \mathbf{u}_m\}$. By the reasoning of Step 2, we conclude

$$\begin{aligned} z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\}) &\supseteq z_{\mathbf{A}}(\text{span}(\boldsymbol{\mu}_1, \boldsymbol{\mu}_2) \setminus \{0\}) \\ &= \text{Arc}_{\min}(z_{\mathbf{A}}(\boldsymbol{\mu}_1), z_{\mathbf{A}}(\boldsymbol{\mu}_2)) = \text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m)). \end{aligned}$$

Next, consider the case where $\text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m))$ has an interior. In this case, $\dim \text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \geq 3$ by the arguments of Step 2. Assume for contradiction that there is a $z \in \text{Poly}(z_{\mathbf{A}}(\mathbf{u}_1), \dots, z_{\mathbf{A}}(\mathbf{u}_m))$ but $z \notin z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\})$. Let ζ_1, \dots, ζ_q be vertices provided by Lemma 2.1. There exists corresponding $\{\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_q\} \subseteq \{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ such that $\zeta_i = z_{\mathbf{A}}(\boldsymbol{\mu}_i)$ for $i = 1, \dots, q$. Define the curve

$$\boldsymbol{\eta}(t) : [1, q+1] \rightarrow \text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \cap S^{m-1},$$

where $S^{m-1} \subset \mathbb{R}^m$ is the unit sphere, as

$$\boldsymbol{\eta}(t) = \frac{\cos((t-p)\frac{\pi}{2})}{\|\boldsymbol{\mu}_p\|} \boldsymbol{\mu}_p + \frac{\sin((t-p)\frac{\pi}{2})}{\|\boldsymbol{\mu}_{p+1}\|} \boldsymbol{\mu}_{p+1}, \quad \text{for } p \leq t \leq p+1.$$

where we regard $\boldsymbol{\mu}_{q+1}$ as $\boldsymbol{\mu}_1$. Then $\{\boldsymbol{\gamma}(t)\}_{t \in [1, q+1]} = \{z_{\mathbf{A}}(\boldsymbol{\eta}(t))\}_{t \in [1, q+1]}$ encloses z .

Since $\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \cap S^{m-1}$ is simply connected, we can continuously contract $\{\boldsymbol{\eta}(t)\}_{t \in [1, q+1]}$ to a point in $\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \cap S^{m-1}$ and the curve under the map $z_{\mathbf{A}}$ continuously contracts to a point in $z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\})$. However, this is not possible as $z \notin z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\})$ and $\{\boldsymbol{\gamma}(t)\}_{t \in [1, q+1]}$ has a nonzero winding number around z . We have a contradiction and we conclude $z \in z_{\mathbf{A}}(\text{span}(\mathbf{u}_1, \dots, \mathbf{u}_m) \setminus \{0\})$. \square

4. SRGs of normal matrices. We now use Theorem 3.1 to fully characterize the SRG of normal matrices.





Fig. 4: Illustration of Proposition 4.1



(a) SRG of an $n \times n$ normal matrix with one distinct real eigenvalue and three distinct complex conjugate eigenvalue pairs.

(b) SRG of an $n \times n$ symmetric matrix with distinct eigenvalues $\lambda_1 < \lambda_2 < \dots < \lambda_6$.

Fig. 5: Illustration of Theorem 4.3. For normal matrices, multiplicity of eigenvalues do not affect the SRG.

PROPOSITION 4.1. Let $A = \begin{bmatrix} a_1 & b_1 \\ b_2 & a_2 \end{bmatrix} \in \mathbb{R}^{2 \times 2}$. Then $\mathcal{G}(A)$ consists of two circles centered at $(\frac{a_1+a_2}{2}, \pm \frac{b_1-b_2}{2})$ with radius $\sqrt{(\frac{a_1-a_2}{2})^2 + (\frac{b_1+b_2}{2})^2}$.

Proof. Let

$$x_\theta = \begin{bmatrix} \cos(\theta) \\ \sin(\theta) \end{bmatrix} \in \mathbb{R}^2, \quad T \left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) = \begin{bmatrix} x_1 \\ |x_2| \end{bmatrix}.$$

The stated result follows from

$$\mathcal{G}^+(A) = \{z_A(x_\theta) : \theta \in [0, 2\pi)\}$$

and the calculations

$$\begin{aligned} z_A(x_\theta) &= \left[\frac{1}{2} (a_1 + a_2 + (a_1 - a_2) \cos(2\theta) + (b_1 + b_2) \sin(2\theta)) \right. \\ &\quad \left. \frac{1}{2} |-b_1 + b_2 + (b_1 + b_2) \cos(2\theta) - (a_1 - a_2) \sin(2\theta)| \right] \\ &= T \left(\begin{bmatrix} \frac{a_1+a_2}{2} \\ -\frac{b_1-b_2}{2} \end{bmatrix} + \underbrace{\begin{bmatrix} \cos(-2\theta) & -\sin(-2\theta) \\ \sin(-2\theta) & \cos(-2\theta) \end{bmatrix}}_{\text{rotation by } -2\theta} \begin{bmatrix} \frac{a_1-a_2}{2} \\ \frac{b_1+b_2}{2} \end{bmatrix} \right). \quad \square \end{aligned}$$

PROPOSITION 4.2. A matrix's SRG is invariant under orthogonal similarity transforms.

Proof. Let $A \in \mathbb{R}^{n \times n}$. Let $Q \in \mathbb{R}^{n \times n}$ be orthogonal. For any nonzero $x \in \mathbb{R}^n$, we have

$$\begin{aligned} z_{QAQ^T}(x) &= \frac{\|QAQ^T x\|}{\|x\|} \exp[i\angle(QAQ^T x, x)] \\ &= \frac{\|AQ^T x\|}{\|x\|} \exp[i\angle(AQ^T x, Q^T x)] = z_A(Q^T x). \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{G}(QAQ^T) &= \{z_{QAQ^T}(x), \overline{z_{QAQ^T}}(x) : x \in \mathbb{R}^n, x \neq 0\} \\ &= \{z_A(Qx), \overline{z_A}(Qx) : x \in \mathbb{R}^n, x \neq 0\} \\ &= \{z_A(x), \overline{z_A}(x) : x \in \mathbb{R}^n, x \neq 0\} = \mathcal{G}(A). \end{aligned} \quad \square$$

THEOREM 4.3. *If A is normal, then $\mathcal{G}^+(A) = \text{Poly}(\Lambda(A) \cap \mathbb{C}^+)$.*

Proof. A normal matrix is orthogonally similar to the real block-diagonal matrix

$$\begin{bmatrix} a_1 & b_1 & & & & & & & \\ -b_1 & a_1 & & & & & & & \\ & & \ddots & & & & & & \\ & & & a_k & b_k & & & & \\ & & & -b_k & a_k & & & & \\ & & & & & \lambda_{k+1} & & & \\ & & & & & & \ddots & & \\ & & & & & & & & \lambda_m \end{bmatrix}.$$

Propositions 4.1 tells us

$$\mathcal{G}^+\left(\begin{bmatrix} a_j & b_j \\ -b_j & a_j \end{bmatrix}\right) = \{a_j + |b_j|i\} = \Lambda\left(\begin{bmatrix} a_j & b_j \\ -b_j & a_j \end{bmatrix}\right) \cap \mathbb{C}^+$$

for $j = 1, \dots, k$. We conclude the stated result with Theorem 4.3 and Proposition 4.2. \square

COROLLARY 4.4. *Let $A \in \mathbb{R}^{n \times n}$ be symmetric, and let $\lambda_1 < \lambda_2 < \dots < \lambda_m$ be the distinct (real) eigenvalues of A . If $m = 1$, then $\mathcal{G}^+(A) = \{\lambda_1\}$. If $m \geq 2$, then*

$$\mathcal{G}(A) = \text{Disk}(\lambda_1, \lambda_m) \setminus \bigcup_{i=1}^{m-1} \text{Disk}^\circ(\lambda_i, \lambda_{i+1}).$$

REFERENCES

- [1] G. BANJAC AND P. J. GOULART, *Tight global linear convergence rate bounds for operator splitting methods*, IEEE Transactions on Automatic Control, 63 (2018), pp. 4126–4139.
- [2] E. BELTRAMI, *Saggio di interpretazione della geometria Non-Euclidea*, 1868.
- [3] E. BELTRAMI, *Teoria fondamentale degli spazii di curvatura costante*, Annali di Matematica Pura ed Applicata. Series II., 2 (1868), pp. 232–255.
- [4] J. ECKSTEIN, *Splitting methods for monotone operators with applications to parallel optimization*, PhD thesis, MIT, 1989.
- [5] J. ECKSTEIN AND D. P. BERTSEKAS, *On the Douglas–Rachford splitting method and the proximal point algorithm for maximal monotone operators*, Mathematical Programming, 55 (1992).
- [6] P. GISELSSON, *Lunds universitet, lecture notes: Large-scale convex optimization*, 2015. URL: <http://www.control.lth.se/education/doctorate-program/large-scale-convex-optimization/>. Last visited on 2018/12/01.
- [7] P. GISELSSON AND S. BOYD, *Linear convergence and metric selection for Douglas–Rachford splitting and ADMM*, IEEE Transactions on Automatic Control, 62 (2017), pp. 532–544.
- [8] P. GISELSSON AND W. M. MOURSI, *On compositions of special cases of Lipschitz continuous operators*, arXiv preprint arXiv:1912.13165, (2019).
- [9] X. HUANG, E. K. RYU, AND W. YIN, *Tight coefficients of averaged operators via scaled relative graph*, arXiv preprint arXiv:1912.01593, (2019).
- [10] F. KLEIN, *Ueber die sogenannte nicht-euklidische geometrie*, Mathematische Annalen, 6 (1873), pp. 112–145.
- [11] H. POINCARÉ, *Théorie des groupes fuchsians*, Acta Mathematica, 1 (1882), pp. 1–62.
- [12] E. K. RYU, R. HANNAH, AND W. YIN, *Scaled relative graph: Nonexpansive operators via 2D Euclidean geometry*, arXiv preprint arXiv:1902.09788, (2019).