

ON A NEW TYPE BERTRAND CURVE

ÇETİN CAMCI

ABSTRACT. In this paper, we define a new type Bertrand curve and this curves are said V -Bertrand curve, f -Bertrand curve and a -Bertrand curve. In addition, we give charectarization of the V -Bertrand curve and we define a Bertrand surface.

1. Introduction

Let $\gamma : I \longrightarrow \mathbb{R}^3$ ($s \rightarrow \gamma(s)$) be a unit-speed curve. Formula of the Serret-Frenet equations are given by

$$\begin{pmatrix} T' \\ N' \\ B' \end{pmatrix} = \begin{pmatrix} 0 & \kappa & 0 \\ -\kappa & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix} \begin{pmatrix} T \\ N \\ B \end{pmatrix}$$

where $\{T, N, B, \kappa, \tau\}$ is the Serret-Frenet apparatus of the curves([3], [4])

In 3-Euclidean spaces, Saint-Venant proposed that the ruled surface whose based curve is the principal normals of a curve and another curve which have same principal normals can exist ([9]). Solution of the problem was given by Bertrand([10]). Let M be a unit-speed curve with coordinate neighborhood (I, γ) of the curve. So we can define a curve by

$$\beta(s) = \gamma(s) + \lambda(s)N(s)$$

where $\lambda : I \longrightarrow \mathbb{R}$ ($s \rightarrow \lambda(s)$) is differentiable function. Let $\{\overline{T}, \overline{N}, \overline{B}\}$ be Frenet frame of the curve and $\overline{\kappa}$ and $\overline{\tau}$ be curvatures of the curves K . If $\{N, \overline{N}\}$ is linear dependent ($\overline{N} = \epsilon N, \epsilon = \pm 1$), then it is said that M is Bertrand mate of K ([13],[14]). If there is Bertrand mate of the curve

2000 *Mathematics Subject Classification.* Primary 53C15; Secondary 53C25.

Key words and phrases. Sasakian Space, curve .

M , then it is said that M is Bertrand curve ([13],[14]). For example, any planar curve and circular helix is Bertrand curve. It is well known theorem that M is Bertrand curve if and only if there exist $\lambda, \mu \in \mathbb{R}$ such that $\lambda\kappa + \mu\tau = 1$ where $\lambda \neq 0$, $\mu = \lambda \cot \theta$ and θ is angle between T and \bar{T} ([13],[14]). In fact, M is Bertrand or B -Bertrand curve if and only if there are $\lambda, \mu \in \mathbb{R}$ such that $\lambda\kappa + \mu\tau = 1$.

If the curve is a unit-speed spherical curve, for all $s \in I$, $\gamma(s)$ is perpendicular to $T = \gamma'(s)$. So we have Sabban frame as

$$\{\gamma(s), T(s) = \gamma'(s), Y(s) = \gamma(s) \times T(s)\}$$

([5]). Serret-Frenet formula of the this curve is given by

$$\begin{pmatrix} \gamma' \\ T' \\ Y' \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & \kappa_g \\ 0 & -\kappa_g & 0 \end{pmatrix} \begin{pmatrix} \gamma \\ T \\ Y \end{pmatrix}$$

where $\kappa_g(s) = \det(\gamma, T, T')$ ([5]). Under the above notation, the space curve is defined by

$$(1.1) \quad \tilde{\gamma}(s) = a \int \gamma(s) ds + a \cot \theta \int Y(s) ds + c$$

where $a, \theta \in \mathbb{R}$ and c is constant vector ([5]). Izumiya and Tkeuchi prove that all Bertrand curves can be constructed from equation (1.1) and they gave some Bertrand curve examples by using equation (1.1) ([5]). In this paper, we give two method different from above method. Furthermore, we define a V -Bertrand curve, f -Bertrand curve and a -Bertrand curve where Bertrand curve is T -Bertrand curve and $a \in \mathbb{R}$. Furthermore, we modify charectarization of the Bertrand curve. In addition, we define a Bertrand surface and we give an example Bertrand surface. Moreover, we define a equivalence relation on Bertrand surface set.

2. Preliminaries

In 3-Euclidean spaces, let M be a unit-speed curve and $\{T, N, B, \kappa, \tau\}$ is Serret-Frenet apparatus of the curves. The curve lies on 2-sphere if and

only if below equation is hold

$$(2.1) \quad \left(\left(\frac{1}{\kappa} \right)' \frac{1}{\tau} \right)' + \frac{\tau}{\kappa} = 0$$

([6],[8]). From the solution of the equation (2.1), we have

$$(2.2) \quad \frac{1}{\kappa} = R \cos \left(\int_0^s \tau(u) du + \theta_0 \right).$$

where radius of the sphere is equal to R ([6],[7], [11]). It is well known that if M is spherical curve, then M and osculating circle lies on this sphere. From equation (2.2), we have

$$\frac{1}{\kappa_0} = \cos \theta_0 = \frac{R_0}{R}.$$

If R is equal to 1, then we have $\cos \theta_0 = R_0 = \frac{1}{\kappa_0}$ ([12]).

Choi and Kim defined a unit vector field V which is given by

$$V(s) = u(s)T(s) + v(s)N(s) + w(s)B(s)$$

and integral curve of V is define by

$$\gamma_V(s) = \int V(s) ds$$

where u, v, w are functions from I to \mathbb{R} and $u^2 + v^2 + w^2 = 1$ ([1]). Without loss of generality, we suppose that arc-length parameter of M and integral curve of V are same ([1]). If $u(s) = 1, v(s) = w(s) = 0$, for all $s \in I$, then we have

$$\gamma_T(s) = \int T(s) ds = \gamma(s).$$

If $u(s) = w(s) = 0, v(s) = 1$, for all $s \in I$, then we have

$$\gamma_N(s) = \int N(s) ds$$

and if $u(s) = v(s) = 0, w(s) = 1$, for all $s \in I$, then we have

$$\gamma_B(s) = \int B(s) ds$$

([1]). (γ_N) (resp. (γ_B)) is called principal-direction curve (resp. binormal-direction curve) of γ ([1]). Let

$$\kappa_1 = \sqrt{\kappa^2 + \tau^2}, \tau_1 = \frac{\kappa^2}{\kappa^2 + \tau^2} \left(\frac{\tau}{\kappa} \right)'$$

and $T_1 = N$, $N_1 = \frac{N'}{\|N'\|}$, $B_1 = T_1 \times N_1$ (resp. $\kappa_\beta = |\tau|$, $\tau_\beta = \kappa$ and $T_\beta = B$, $N_\beta = N$, $B_\beta = -T$) be curvatures and Frenet vectors of principal-direction curve (resp. binormal-direction curve) ([1], [2]). If

$$u(s) = 0, v(s) = -\cos \left(\int \tau(s) ds \right) \neq 0, w(s) = \sin \left(\int \tau(s) ds \right)$$

then $\gamma_{pdo}(s) = \int V(s) ds$ is called principal-donor curve of γ , for all $s \in I$ ([1]). So we have

$$\kappa_{pdo} = \kappa \cos \left(\int \tau(s) ds \right), \tau_{pdo} = \kappa \sin \left(\int \tau(s) ds \right)$$

and $T_{pdo} = V$, $N_{pdo} = T$, $B_{pdo} = T_{pdo} \times N_{pdo}$.

3. V-BERTRAND CURVE

Let $\gamma : I \longrightarrow \mathbb{R}^3$ ($s \rightarrow \gamma(s)$) be a unit-speed curve and $\{T, N, B, \kappa, \tau\}$ be Serret-Frenet apparatus of the curve. We can define a curve K by

$$\beta(s) = \int V(s) ds + \lambda(s)N(s)$$

where $\lambda : I \longrightarrow \mathbb{R}$ ($s \rightarrow \lambda(s)$) is differentiable function and $V = uT + vN + wB$ is unit vector field. Let $\{\overline{T}, \overline{N}, \overline{B}\}$ be orthonormal frame of the curve and $\overline{\kappa}$ and $\overline{\tau}$ be curvatures of the curve K . So we can give following definition.

Definition 3.1. If $\{N, \overline{N}\}$ is linear dependent ($\overline{N} = \epsilon N, \epsilon = \pm 1$), then it is said that (M, K) is V-Bertrand mate. If $V = T$, then (M, K) is Bertrand mate.

Theorem 3.1. Let M be a unit-speed curve and $\{T, N, B, \kappa, \tau\}$ be Serret-Frenet apparatus of the curves. M is V-Bertrand curve if and only if it is satisfied that

$$(3.1) \quad \lambda(\kappa \tan \theta + \tau) = u \tan \theta - w$$

and

$$(3.2) \quad \lambda(s) = - \int v(s) ds$$

where θ is constant angle between T and \overline{T} .

Proof. If M is V -Bertrand curve, then V -Bertrand mate of M is equal to

$$(3.3) \quad \beta(s) = \int V(s) ds + \lambda(s) N(s)$$

and $\{N, \overline{N}\}$ is linear dependent. If we derivative equation (3.3), we have

$$\frac{d\overline{s}}{ds} \overline{T} = (u - \lambda\kappa) T + (\lambda' + v) N + (w + \lambda\tau) B.$$

Since $\{N, \overline{N}\}$ is linear dependent, we have

$$\lambda(s) = - \int v(s) ds$$

and we get

$$(3.4) \quad \overline{T} = \frac{ds}{d\overline{s}} (u - \lambda\kappa) T + \frac{ds}{d\overline{s}} (w + \lambda\tau) B = \cos \theta(s) T + \sin \theta(s) B$$

where $\cos \theta(s) = \frac{ds}{d\overline{s}} (u - \lambda\kappa)$ and $\sin \theta(s) = \frac{ds}{d\overline{s}} (w + \lambda\tau)$. So we have

$$\tan \theta = \frac{w + \lambda\tau}{u - \lambda\kappa}$$

or

$$\lambda (\kappa \tan \theta + \tau) = u \tan \theta - w.$$

If we take the derivative of a equation (3.4), we have

$$\frac{d\overline{s}}{ds} \overline{\kappa} \overline{N} = -\theta' \sin \theta T + (\kappa \cos \theta - \tau \sin \theta) N + \theta' \cos \theta B$$

So that $\{N, \overline{N}\}$ is linear dependent, we have $\theta' = 0$ or $\theta = \text{const.}$ Conversely, we define a curve by

$$(3.5) \quad \beta(s) = \int V(s) ds + \lambda(s) N(s)$$

where $\lambda : I \rightarrow \mathbb{R}$ ($s \rightarrow \lambda(s)$) is differentiable function such that $\lambda(s) = - \int v(s) ds$. If we derivative equation (3.5), we have

$$\frac{d\overline{s}}{ds} \overline{T} = (u - \lambda\kappa) T + (\lambda' + v) N + (w + \lambda\tau) B.$$

From equation (3.2), we get

$$(3.6) \quad \overline{T} = \frac{ds}{d\overline{s}}(u - \lambda\kappa)T + \frac{ds}{d\overline{s}}(w + \lambda\tau)B = \cos\phi(s)T + \sin\phi(s)B$$

where $\cos\phi(s) = \frac{ds}{d\overline{s}}(u - \lambda\kappa)$ and $\sin\phi(s) = \frac{ds}{d\overline{s}}(w + \lambda\tau)$. From equation (3.1) and (3.6) we have

$$\tan\phi(s) = \frac{v + \lambda\tau}{u - \lambda\kappa} = \tan\theta.$$

So $\phi = \theta + n\pi$ is angle between T and \overline{T} where $n \in \mathbb{Z}$. If we take the derivative of a equation (3.6), we see that $\{N, \overline{N}\}$ is linear dependent. \square

Definition 3.2. Let M be a unit-speed non-planar curve and $\{T, N, B, \kappa, \tau\}$ is Serret-Frenet apparatus of the curves. If there is $\lambda \neq 0$, $\theta \in \mathbb{R}$ such that

$$\lambda\kappa + \lambda \cot\theta\tau = 1,$$

then we said that M is Bertrand curve (or T -Bertrand curve). If there is $\lambda \neq 0$, $\theta \in \mathbb{R}$ such that

$$\lambda \tan\theta\kappa + \lambda\tau = -1,$$

then we said that M is B -Bertrand curve.

Corollary 3.1. If $u(s) = 1, v(s) = w(s) = 0$, we have Bertrand curve. From equation (3.2), λ is constant. From equation (3.1), we have $\lambda\kappa + \mu\tau = 1$ where $\lambda \neq 0$, $\mu = \lambda \cot\theta$ and θ is constant angle between T and \overline{T} . If $v(s) = 1, u(s) = w(s) = 0$, we have B -Bertrand curve. From equation (3.1), we have $\lambda\kappa + \mu\tau = -1$ where $\mu \neq 0$ is constant, $\lambda = \mu \tan\theta$ and θ is constant angle between T and \overline{T} .

Corollary 3.2. If the curve is a Salkowki curve, then curvature of the curve is equal to 1(or constant). If $\lambda = 1$, $\mu = \lambda \cot\theta = 0$, then we have $\lambda\kappa + \mu\tau = 1$. if $\mu = 0$, $\lambda = \mu \cot\theta = -1$, then we have $\lambda\kappa + \mu\tau = -1$. But this is contradiction. So a Salkowski curve is a Bertrand curve, but it is not B -Bertrand curve. If the curve is anti-Salkowki curve, then torsion of the curve is equal to 1(or constant). If $\mu = -1$, $\lambda = \mu \tan\theta = 0$, then we have $\lambda\kappa + \mu\tau = -1$. if $\lambda = 0$, $\mu = \lambda \cot\theta = -1$, then we have

$\lambda\kappa + \mu\tau = -1$. But this is a contradiction. So a anti-Salkowski curve is a B-Bertrand curve, but it is not a Bertrand curve.

Corollary 3.3. *If the curve is planar curve ($\tau = 0$), then the curve is not only Bertrand but also B-Bertrand curve.*

Corollary 3.4. *If $v(s) = 1, u(s) = w(s) = 0$, we have N-Bertrand curve. From equation (3.2), we have $\lambda = c - s$. From equation (3.1), we have $\frac{\tau}{\kappa} = -\tan \theta$ where θ is constant angle between T and \bar{T} . As a result, M is N-Bertrand curve if and only if M is general helix.*

Corollary 3.5. *If the curve M is Bertrand curve, there is $\lambda \neq 0, \mu \neq 0 \in \mathbb{R}$ such that $\lambda\kappa + \mu\tau = 1$. If we define $\lambda_1 = -\lambda$ and $\mu_1 = -\mu$, then we have $\lambda_1\kappa + \mu_1\tau = -1$. As a result, the curve M is B-Bertrand curve. Conversely, similarly, If the curve M is B-Bertrand curve where there exist $\lambda \neq 0, \mu \neq 0 \in \mathbb{R}$ such that $\lambda\kappa + \mu\tau = -1$, then the curve M is Bertrand curve. In this case, it is said that M is proper Bertrand curve*

Corollary 3.6. *Let (α) be a unit-speed curve and $\{T, N, B, \kappa, \tau\}$ be Serret-Frenet apparatus of the curves. We suppose that $v = 0$, then we can see that λ is non zero constant. So we get $u^2 + w^2 = 1$ and $w = \varepsilon\sqrt{1 - u^2}$ where $\varepsilon = \pm 1$. From equation (3.1), we have*

$$(3.7) \quad u \tan \theta - \varepsilon \sqrt{1 - u^2} = f$$

So, from (3.7), we have

$$(3.8) \quad u^\pm = \frac{f \tan \theta \pm \sqrt{1 + (\tan \theta)^2 - f^2}}{1 + (\tan \theta)^2}.$$

If $u^+ = \frac{f \tan \theta + \sqrt{1 + (\tan \theta)^2 - f^2}}{1 + (\tan \theta)^2}$, then we have

$$w_1^\pm = \pm \sqrt{1 - (u^+)^2}$$

If $u^- = \frac{f \tan \theta - \sqrt{1 + (\tan \theta)^2 - f^2}}{1 + (\tan \theta)^2}$, then we have

$$w_2^\pm = \pm \sqrt{1 - (u^-)^2}$$

So, we have

$$\begin{aligned} V_1^+ &= u^+T + w_1^+B, V_1^- = u^+T + w_1^-B, \\ V_2^+ &= u^-T + w_2^+B, V_2^- = u^-T + w_2^-B \end{aligned}$$

and

$$\begin{aligned} \beta_1^+ &= \int V_1^+ ds + \lambda N, \beta_1^- = \int V_1^- ds + \lambda N \\ \beta_2^+ &= \int V_2^+ ds + \lambda N, \beta_2^- = \int V_2^- ds + \lambda N \end{aligned}$$

In this case, (α) is V_1^+, V_1^-, V_2^+ and V_2^- -Bertrand curve. So we can give following definition.

Definition 3.3. The curves $\beta_1^+, \beta_1^-, \beta_2^+, \beta_2^-$ are called f -Bertrand curve mate of (α) and the curve (α) is called f -Bertrand curve. If $f = r(\text{const.})$, then (α) is called r -Bertrand curve and the curves $\beta_1^+, \beta_1^-, \beta_2^+, \beta_2^-$ are called r -Bertrand curve mate of (α)

Example 3.1. If $\alpha(s) = (a \cos \frac{s}{c}, a \sin \frac{s}{c}, \frac{bs}{c})$, then we have

$$\begin{aligned} T &= \left(-\frac{a}{c} \sin \frac{s}{c}, \frac{a}{c} \cos \frac{s}{c}, \frac{b}{c}\right) \\ N &= \left(-\cos \frac{s}{c}, -\sin \frac{s}{c}, 0\right) \\ B &= \left(\frac{b}{c} \sin \frac{s}{c}, -\frac{b}{c} \cos \frac{s}{c}, \frac{a}{c}\right) \end{aligned}$$

and $\kappa = \frac{a}{c^2}$ and $\tau = \frac{b}{c^2}$ where $c^2 = a^2 + b^2$. If $\bar{\lambda} = \lambda \tan \theta$ and $\bar{\mu} = \lambda$, then we have $\bar{\lambda}\kappa + \bar{\mu}\tau = 1 = f$ where $\tan \theta = \frac{c^2 - \lambda b}{\lambda a}$. From (3.8), we have

$$u^\pm = \frac{\tan \theta \pm \tan \theta}{1 + (\tan \theta)^2}$$

So we have $u^- = 0$ and $u^+ = \frac{2\lambda a(c^2 - \lambda b)}{c^2(a^2 + (\lambda - b)^2)}$. If $u^- = 0$, then we have

$w_2^+ = 1$ and $w_2^- = -1$. If $u^+ = \frac{2\lambda a(c^2 - \lambda b)}{c^2(a^2 + (\lambda - b)^2)}$, then we have

$$w_1^+ = \frac{\lambda^2 a^2 + c^2(2\lambda b - c^2)}{c^2(a^2 + (\lambda - b)^2)}$$

and

$$w_1^- = -\frac{\lambda^2 a^2 + c^2 (2\lambda b - c^2)}{c^2 (a^2 + (\lambda - b)^2)}$$

So we have

$$V_2^+ = B, V_2^- = -B, V_1^+ = u^+ T + w_2^+ B, V_1^- = u^- T + w_2^- B$$

and

$$\begin{aligned}\beta_2^\pm &= \left((\lambda \mp b) \cos \frac{s}{c}, (\lambda \mp b) \sin \frac{s}{c}, \pm \frac{as}{c} \right) \\ \beta_1^\pm &= \left((w_2^\pm b - u_2 a - \lambda) \cos \frac{s}{c}, (w_2^\pm b - u_2 a - \lambda) \sin \frac{s}{c}, \frac{w_2^\pm a + u_2 b}{c} s \right).\end{aligned}$$

So, (α) is 1-Bertrand curve and $\beta_1^+, \beta_1^-, \beta_2^+, \beta_2^-$ curves are all 1-Bertrand curve mate of (α) .

Example 3.2. From (3.8), if $\bar{\lambda}\kappa + \bar{\mu}\tau = \tan \theta = f$, then we obtain $u^+ = 1$ and $u^- = -\cos 2\theta$ where $\tan \theta = \frac{c^2 - \lambda b}{\lambda a}$. If $u^+ = 1$ (resp. $u^- = -\cos 2\theta$), then we have $w_2^\pm = 0$ (resp. $w_1^\pm = \pm \sin 2\theta$). So we have

$$V_1^+ = u^+ T + w_1^+ B, V_1^- = u^- T + w_1^- B, V_2^\pm = T$$

and

$$\begin{aligned}\beta_2^\pm &= \left((\lambda - a) \cos \frac{s}{c}, (\lambda - a) \sin \frac{s}{c}, \frac{bs}{c} \right) \\ \beta_1^\pm &= \left((w_2^\pm b - u_2 a - \lambda) \cos \frac{s}{c}, (w_2^\pm b - u_2 a - \lambda) \sin \frac{s}{c}, \frac{w_2^\pm a + u_2 b}{c} s \right).\end{aligned}$$

(α) is $\tan \theta$ -Bertrand curve and the curves $\beta_1^+, \beta_1^-, \beta_2^+, \beta_2^-$ are all $\tan \theta$ -Bertrand curve mate of (α) .

Let (γ) be Bertrand curve and $\{T, N, B, \kappa, \tau\}$ be Serret-Frenet apparatus of the curves. We suppose that $v = 0$, then we can see that λ is non-zero constant. We can obtain

$$V(s) = uT(s) + wB(s)$$

where $u^2 + w^2 = 1$ and $u, w \in \mathbb{R}$. Integral curve of V is defined as

$$(3.9) \quad \gamma_V(s) = \int V(s) ds$$

Let $\{T_V, N_V, B_V, \kappa_V, \tau_V\}$ be Serret-Frenet apparatus of the curves (γ_V) . From equation (3.9), we have

$$(3.10) \quad T_V = V(s) = uT(s) + wB(s)$$

From equation (3.10), we can see that the curves (γ) and (γ_V) have a same arc-length parameter. From equation (3.10), we have

$$\kappa_V N_V = (u\kappa - w\tau) N$$

So we obtain

$$(3.11) \quad N_V = N$$

and

$$(3.12) \quad \kappa_V = u\kappa - w\tau$$

From equation (3.10) and (3.11), we get

$$(3.13) \quad B_V = T_V \times N_V = -wT(s) + uB(s)$$

From equation (3.13), we have

$$(3.14) \quad -\tau_V N_V = -(w\kappa + u\tau) N$$

So we obtain

$$(3.15) \quad \tau_V = w\kappa + u\tau$$

From equation (3.12) and (3.15), we get

$$(3.16) \quad \kappa = u\kappa_V + w\tau_V$$

and

$$(3.17) \quad \tau = -w\kappa_V + u\tau_V$$

In this case, we can obtain following theorem.

Theorem 3.2. *(γ) is the Bertrand (resp. B-Bertrand) curve if and only if (γ_V) is the Bertrand (resp. B-Bertrand) curve.*

Proof. We suppose that (γ) is the Bertrand curve. There is $\lambda, \mu \in \mathbb{R}$ such that

$$\lambda\kappa + \mu\tau = u - w \cot \theta = 1$$

where $\lambda \neq 0$, $\mu = \lambda \cot \theta$ and θ is constant angle between T and \bar{T} . From equation (3.12) and (3.15), we get

$$(3.18) \quad \lambda\kappa + \mu\tau = \bar{\lambda}\kappa_V + \bar{\mu}\tau_V = 1$$

where

$$(3.19) \quad \bar{\lambda} = \lambda u - \mu w = \lambda \neq 0$$

and

$$(3.20) \quad \bar{\mu} = \lambda w + \mu u = \lambda \cot(-\theta)$$

So, (γ_V) is the Bertrand curve. Similarly, other cases can be proved by the same method. \square

Let (α) be Bertrand curve and $\{T, N, B, \kappa, \tau\}$ is Serret-Frenet apparatus of the curves. There is $\lambda, \mu \in \mathbb{R}$ such that $\lambda\kappa + \mu\tau = 1$ where $\mu = \lambda \tan \theta$. If $\bar{\lambda} = \lambda a$ and $\bar{\mu} = \mu a$, then $\bar{\lambda}\kappa + \bar{\mu}\tau = a$. So we have

$$(3.21) \quad u^\pm(a) = \frac{a \tan \theta \pm \sqrt{1 + (\tan \theta)^2 - a^2}}{1 + (\tan \theta)^2}$$

and $w_1^\pm(a) = \pm \sqrt{1 - (u^+(a))^2}$, $w_2^\pm(a) = \pm \sqrt{1 - (u^-(a))^2}$. In this case, we can give following definition.

Definition 3.4. Let (α) be Bertrand curve. So, we can define a surface as

$$\begin{aligned} \varphi_1^+(t, s) &= \int V_1^+ ds + \lambda N, \varphi_1^-(t, s) = \int V_1^- ds + \lambda N \\ \varphi_2^+(t, s) &= \int V_2^+ ds + \lambda N, \varphi_2^-(t, s) = \int V_2^- ds + \lambda N \end{aligned}$$

where

$$\begin{aligned} V_1^+(t, s) &= u^+(t)T(s) + w_1^+(t)B(s), V_1^-(t, s) = u^+(t)T(s) + w_1^-(t)B(s), \\ V_2^+(t, s) &= u^-(t)T(s) + w_2^+(t)B(s), V_2^-(t, s) = u^-(t)T(s) + w_2^-(t)B(s), \end{aligned}$$

$u^\pm(t) = \frac{t \tan \theta \pm \sqrt{1 + (\tan \theta)^2 - t^2}}{1 + (\tan \theta)^2}$, $w_1^\pm(t) = \pm \sqrt{1 - (u^+(t))^2}$ and $w_2^\pm(t) = \pm \sqrt{1 - (u^-(t))^2}$. These surfaces are said Bertrand surface taken by (α) .

Example 3.3. If $\alpha(s) = (\cos \frac{s}{\sqrt{2}}, \sin \frac{s}{\sqrt{2}}, \frac{s}{\sqrt{2}})$, then we have

$$T = \left(-\frac{1}{\sqrt{2}} \sin \frac{s}{\sqrt{2}}, \frac{1}{\sqrt{2}} \cos \frac{s}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$$

$$N = \left(-\cos \frac{s}{\sqrt{2}}, -\sin \frac{s}{\sqrt{2}}, 0\right)$$

$$B = \left(\frac{1}{\sqrt{2}} \sin \frac{s}{\sqrt{2}}, -\frac{1}{\sqrt{2}} \cos \frac{s}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$$

and $\kappa = \frac{1}{2}$ and $\tau = \frac{1}{2}$. If $\bar{\lambda} = \lambda \tan \theta = 1$ and $\bar{\mu} = \lambda = 1$, then we have $\bar{\lambda}\kappa + \bar{\mu}\tau = 1 = f$ and $\tan \theta = 1$. From (3.21), we have

$$u(t) = \frac{t + \sqrt{2 - t^2}}{2}$$

and

$$w(t) = \frac{1}{\sqrt{2}} \sqrt{1 - t \sqrt{2 - t^2}}$$

So, we have a surface $K(\alpha)$ as

$$\begin{aligned} \varphi(t, s) &= u(t) \int T ds + w(t) \int B ds + \lambda N \\ &= \left(a(t) \cos \frac{s}{\sqrt{2}}, a(t) \sin \frac{s}{\sqrt{2}}, b(t) \frac{s}{\sqrt{2}}\right) \end{aligned}$$

where $a(t) = u(t) - w(t) - 1$ and $b(t) = u(t) + w(t)$. The surface $K(\alpha)$ is Bertrand surface taken by (α) .

Definition 3.5. Let $((\alpha), (\beta))$ be Bertrand curve mate and $K(\alpha)$ and $K(\beta)$ be Bertrand surface. $(K(\alpha), K(\beta))$ is said Bertrand surface mate taken by $((\alpha), (\beta))$ Bertrand mate.

We can define a set as

$$K = \{K(\alpha) / (\alpha) \text{ is a Bertrand curve}\}.$$

So we can define a equivalence relation on K such that $K(\alpha) \sim K(\beta)$ if and only if $((\alpha), (\beta))$ is a Bertrand mate.

In Example 3.3, if (β) is Bertrand mate of (α) , then we have

$$\begin{aligned}\beta(s) &= \alpha(s) + \lambda N \\ &= \left(\cos \frac{s}{\sqrt{2}}, \sin \frac{s}{\sqrt{2}}, \frac{s}{\sqrt{2}}\right) + \left(-\cos \frac{s}{\sqrt{2}}, -\sin \frac{s}{\sqrt{2}}, 0\right) \\ &= \left(0, 0, \frac{s}{\sqrt{2}}\right)\end{aligned}$$

Example 3.4. If $\beta(s) = (0, 0, \frac{s}{\sqrt{2}})$, then we have

$$\begin{aligned}T &= (0, 0, 1) \\ N &= \left(-\cos \frac{s}{r}, -\sin \frac{s}{r}, 0\right) \\ B &= \left(\sin \frac{s}{r}, -\cos \frac{s}{r}, 0\right)\end{aligned}$$

and $\kappa = 0$ and $\tau = r$. If $\bar{\lambda} = \lambda \tan \theta = 0$ and $\bar{\mu} = \lambda = r$, then we have $\bar{\lambda}\kappa + \bar{\mu}\tau = 1 = f$ and $\tan \theta = 0$. From (3.21), we have $u(t) = \sqrt{1-t^2}$ and $w(t) = t$. So, we have a surface $K(\alpha)$ as

$$\begin{aligned}\varphi(t, s) &= u(t) \int T ds + w(t) \int B ds + \lambda N \\ &= \left(-\left(r + \frac{t}{r}\right) \cos \frac{s}{\sqrt{2}}, -\left(r + \frac{t}{r}\right) \sin \frac{s}{\sqrt{2}}, \sqrt{1-t^2}s\right)\end{aligned}$$

The surface $K(\beta)$ is Bertrand surface taken by (β) . Furthermore, $(K(\alpha), K(\beta))$ is Bertrand surface mate taken by $((\alpha), (\beta))$.

4. CONSTRUCTION OF THE BERTRAND AND THE PRINCIPAL-DONOR

CURVES

In 3-Euclidean spaces, let M be a regular spherical curve with coordinate neighborhood (I, γ) . So we can define a curve $K(\alpha : I \rightarrow E^3)$ such that

$$(4.1) \quad \alpha(s) = \int S_M(s) \gamma(s) ds$$

where $S_M : I \rightarrow \mathbb{R}$ ($s \rightarrow S_M(s)$) is differentiable function.

Lemma 4.1. ([12]) *The curve K is spherical curve if and only if*

$$S_M(s) = \|\gamma'(s)\| \cos \left(\int_0^s \frac{\det(\gamma(s), \gamma'(s), \gamma''(s))}{\|\gamma'(s)\|^2} ds + \theta_0 \right).$$

Let M be a unit-speed curve with coordinate neighborhood (I, γ) and $\{T, N, B, \kappa, \tau\}$ be Serret-Frenet apparatus of the curves. Because of unit-speed curve, we have $\|\gamma'(s)\| = 1$. From lemma 4.1, there is $S_T : I \rightarrow \mathbb{R}$ differentiable function such that

$$\left\| \int S_T(s) \gamma'(s) ds \right\| = 1$$

where

$$(4.2) \quad S_T(s) = \kappa(s) \cos \left(\int_0^s \tau(u) du + \theta_0 \right)$$

If we define a curve K with coordinate neighborhood (I, β) such that

$$\beta'(s) = \int S_T(s) \gamma'(s) ds$$

then we have

$$(4.3) \quad \beta''(s) = S_T(s) \gamma'(s)$$

Arc-length parameter of M and K are same. Let $\{\overline{T}, \overline{N}, \overline{B}, \overline{\kappa}, \overline{\tau}\}$ be Serret-Frenet apparatus of the curves where

$$(4.4) \quad S_T(s) = \overline{\kappa}(s) = \kappa(s) \cos \left(\int_0^s \tau(u) du + \theta_0 \right)$$

and

$$(4.5) \quad \overline{\tau}(s) = \kappa(s) \sin \left(\int_0^s \tau(u) du + \theta_0 \right).$$

From (4.3), we have

$$(4.6) \quad \overline{N}(s) = \varepsilon T(s)$$

where $\varepsilon = \pm 1$. From equation (4.3) and (4.6), we can see that principal normal of K and tangent of M is colinear.

Theorem 4.1. *K is principal-donor curve of M if and only if M is principal direction curve of K .*

Proof. From equation of $\beta''(s) = S_T(s)\gamma'(s)$, we have $\overline{N} = T$. So we can write

$$(4.7) \quad \overline{T} = vN + wB$$

where $v^2 + w^2 = 1$. If we take the derivative of equation (4.7), we have

$$(4.8) \quad \overline{\kappa}\overline{N} = -\kappa vT + (v' - w\tau)N + (w' + \tau v)B$$

So we have

$$(4.9) \quad \overline{\kappa} = -\kappa v, \quad v' - w\tau = 0, \quad w' + \tau v = 0$$

From equation (4.9), we have

$$v = -\cos\left(\int_0^s \tau(u)du + \theta_0\right)$$

and

$$w = \sin\left(\int_0^s \tau(u)du + \theta_0\right)$$

Thus, we have

$$(4.10) \quad \overline{T} = -\cos\left(\int_0^s \tau(u)du + \theta_0\right)N + \sin\left(\int_0^s \tau(u)du + \theta_0\right)B$$

From equation (4.10), K is principal-donor curve of M . Conversely, from equation $\beta''(s) = \overline{\kappa}\gamma'(s)$, we have $\overline{N} = T$. So we have $\gamma(s) = \int_0^s \overline{N}(u)du$. \square

Theorem 4.2. *In 3-Euclidean spaces, let M, K be a unit-speed curves with unit coordinate neighborhoods (I, γ) and (I, β) , respectively, such that*

$$\beta''(s) = S_T(s)\gamma'(s)$$

where

$$(4.11) \quad S_T(s) = \kappa(s) \cos \left(\int_0^s \tau(u) du + \theta_0 \right).$$

Following equations are equivalent,

- 1) M is spherical curve
- 2) K is Bertrand curve or B -Bertrand curve.

Proof. If M is spherical curve, from (2.2) we have

$$\frac{1}{\kappa(s)} = A \cos \left(\int_0^s \tau(u) du + \theta_0 \right)$$

where A and θ_0 are constant. If $\theta_0 = 0$, then we have $\overline{\kappa}(s) = A(const.)$. So K is Salkowski curve. If $\theta_0 = -\frac{\pi}{2}$, then we have $\overline{\tau}(s) = A(const.)$. So, K is anti-Salkowski curve. If $\theta_0 \neq 0$ and $\theta_0 \neq -\frac{\pi}{2}$, then we have

$$1 = A \cos(\theta_0) \epsilon \kappa(s) \cos \theta(s) - A \sin(\theta_0 - c) \epsilon \kappa(s) \sin \theta(s)$$

where $\theta(s) = \int_0^s \tau(u) du$. So we have $\lambda \overline{\kappa}(s) + \mu \overline{\tau}(s) = 1$ where

$$\lambda = A \cos(\theta_0) = const. \text{ and } \mu = -A \sin(\theta_0) = const.$$

Furthermore, K is proper Bertrand curve. Conversely, if K is Bertrand curve or B -Bertrand curve, there are $\lambda, \mu \in \mathbb{R}$ such that $\lambda \overline{\kappa}(s) + \mu \overline{\tau}(s) = 1$. So we have

$$1 = \lambda \kappa(s) \cos \left(\int_0^s \tau(u) du + \theta_0 \right) + \mu \kappa(s) \sin \left(\int_0^s \tau(u) du + \theta_0 \right)$$

and

$$\frac{1}{\kappa(s)} = A \left(\cos(\theta_0) \cos \left(\int_0^s \tau(u) du \right) + \sin(\theta_0) \sin \left(\int_0^s \tau(u) du \right) \right)$$

where $\theta_0 \in [0, 2\pi]$, $\cos\theta_0 = \frac{\lambda}{\sqrt{\lambda^2 + \mu^2}}$, $\sin\theta_0 = -\frac{\mu}{\sqrt{\lambda^2 + \mu^2}}$ and $A = \sqrt{\lambda^2 + \mu^2}$.

So we have $\frac{1}{\kappa(s)} = A \cos \left(\int_0^s \tau(u) du + \theta_0 \right)$. \square

Example 4.1. In 3-Euclidean spaces, let M be a circle where $\gamma(s) = (r \cos ws, -r \sin ws, 0)$ and $w = \frac{1}{r}$. So we have $\kappa = w$ and $\tau = 0$. From (4.4) and (4.5), we obtain $\overline{\kappa}(s) = \epsilon w \cos c_0 = A(\text{const.})$ and $\overline{\tau}(s) = \overline{\epsilon} w \sin c_0 = B(\text{const.})$. So, from (4.3), we get

$$\beta''(s) = A(-\sin ws, -\cos ws, 0)$$

If we integrate of the above equation, we have a curve K as

$$\beta(s) = (Ar^2 \cos(\epsilon ws), Ar^2 \sin(\epsilon ws), bws + c_1)$$

Because K is unit speed curve, we have $b = r \sin c_0$. If $a = Ar^2 = \epsilon r \cos c_0$ and $c_1 = 0$, we have

$$\beta(s) = (a \cos(ws), a \sin(ws), bws)$$

where $r = \sqrt{a^2 + b^2}$.

REFERENCES

- [1] Choi, J.H. and Kim Y.H.: Associated curves of Frenet curves and their applications; Applied Mathematics and Computation 218,9116-9124(2012)
- [2] Monterde, J.: Salkowski curves revisited: A family of curves with constant curvature and non-constant torsion, Comput.Aided Geomet. Design 26(2009), 271-278
- [3] Frenet, F.: Sur les courbe à double courbure, Jour. de Math. 17, 437-447 (1852) Extrait d'une These (Toulouse, 1847)
- [4] Serret, J.A.: Sur quelques formules relatives à la théorie de courbes à double courbure, J. De Math., 16(1851)
- [5] Izumiya, S and Tkeuchi N.: Generic properties of helices and Bertrand curves, J. geom., 74, 97-109, (2002).
- [6] Kreyszig, E.: Differential Geometry, Mathematical Exposition, 11.Uni.of Toronto Press (1959).
- [7] Breuer, E. and Gottlieb, D.: Explicit characterization of spherical curves, Proceedings of the American Mathematical Society (1971)27, No.1.
- [8] Struik, D.J.: Lectures on Classical Differential Geometry, Dover, New-York, (1988).

- [9] Saint-Venant J.C.: Memoire sur les lignes courbas non planes, journal d'Ecole polytechnique (1845) 1-76.
- [10] Bertrand J.: Memoire sur la theorie des courbes a double courbure, comptes Rendus 36 (1850); journal de mathematiques press et Appliquees 15(1850) 332-350
- [11] Wong Y. C.: On an explicit characterization of spherical curves. Proc. Amer. Math. Soc., 34(1) 239-242, 1972.
- [12] Camcı, Ç.: How can we construct a k -slant curve from a given spherical curve?, arXiv:1912.13392[math.DG] 23 December 2019.
- [13] do Carmo, M.P.: Diferential Geometry of Curves and Surfaces, Prentice-Hall, Englewood Cliffs. NJ.(1976).
- [14] Hsiung, C.C.: A first course in Diferential Geometry, International Press, Cambridge.MA(1997).

DEPARTMENT OF MATHEMATICS, ONSEKİZ MART UNIVERSITY, 17020 ÇANAKKALE, TURKEY

E-mail address: ccamci@comu.edu.tr