Modern Physics Letters A © World Scientific Publishing Company

Quadratic integrals of a multi-scalar cosmological model

SAMEERAH JAMAL

School of Mathematics, University of the Witwatersrand, Johannesburg, South Africa. Sameerah.Jamal@wits.ac.za

> Received (Day Month Year) Revised (Day Month Year)

In the context of FRW spacetime with zero spatial curvature, we consider a multi-scalar tensor cosmology model under the pretext of obtaining quadratic conservation laws. We propose two new interaction potentials of the scalar field. Integral to this task, is the existence of dynamical Noether symmetries which are Lie-Bäcklund transformations of the physical system. Finally, analytical solutions of the field are found corresponding to each new model. In one of the models, we find that the scale factor mimics A-cosmology in a special case.

Keywords: Dark energy; scalar field; dynamical symmetries; exact solutions.

PACS Nos.: 98.80.-k, 95.35.+d, 95.36.+x.

1. Introduction

Observational cosmological data for the universe suggests an accelerating phase of expansion 1,2 . This phenomenon has posed many new problems. The source of latetime cosmic acceleration 3,4,5 has been attributed to an unidentified matter with a negative equation of state parameter, viz. dark energy. The cosmological constant, leading to Λ -cosmology, is the simplest candidate for the dark energy. In the existing literature, other cosmological models have been proposed to investigate the acceleration phase of the universe. In most approaches, a cosmic fluid is introduced into Einsteins General Relativity 6,7,8,9 , or other studies modify the Einstein-Hilbert Action 10,11,12 . In this work we are interested in two interacting scalar fields in a spatially flat FRW spacetime.

One important spatially homogeneous scalar field model is that containing exponential potential energy ^{13,14}. A number of inflationary models that include scalar fields have been proposed ¹⁵. Such models have an important role in higher dimensional gravity theories and in studying the dynamics of the early universe. In particular, power-law inflationary models (important in solving the horizon and flatness problems) arise in models with a scalar field having an exponential potential ¹⁶. For single scalar fields, a very steep potential will not cause late-time acceleration ¹⁷, while for multi-scalar field models, the universe can evolve to accelerate

era, i.e. assisted inflation ^{18,19} and assisted quintessence ²⁰. We note that a large body of literature exists for the analytic solutions of single scalar fields, for example, 21,22,23,24 . In relation to our study, we mention the Chiral cosmological model ^{25,26} that have important applications in inflation ²⁷ and descriptions for α -attractors 28,29 .

Currently, there are two primary ways to identify the unknown potential function: a) selection by reconstruction of special cosmological behavior according to some well-known model, or b) phenomenological observations from high energy physics. Nevertheless, the functional form of a potential cannot be posed only at a phenomenological level but should also be justified rigorously at a mathematical level. At this juncture, symmetry based methods are one such option. Indeed, the main motivation to apply Noether symmetry methods for cosmological models is that when a potential function is not unique and is an unknown function, one is able to find the explicit forms of the potential function which possesses a symmetry. Additionally, group invariant transformations are vital tools in physical problems as they feature in multiple applications, namely, they may be used to reduce the order of a given dynamical system, construct conservation laws (if they satisfy Noether's theorem), assess the integrability of the physical system and find exact solutions. Practically, there are two classes of Noether symmetries that are of importance. The first is a Noether point symmetry, where the vector field itself contains no derivative terms. The second type of symmetry is called a Noether contact or dynamical symmetry. The latter are the generators of infinitesimal Lie-Bäcklund transformations - linear in first derivatives. Noether point symmetries give rise to linear conserved charges ³⁰ while in contrast, Noether contact symmetries admit quadratic conserved charges. Now, our interest lies in the contact symmetries and hence the quadratic conservation laws. As mentioned above, a significant by-product of this study will be the determination of the interaction potential function.

When it comes to point symmetries, one finds their presence in cosmology in a number of different contexts. In particular, we refer the reader to 31,32,33,34,35,36,37,38,39,40,41,42 for examples where such symmetries have played a role in model determination. On the other hand, applications of contact symmetries in cosmology can be found in, inter alia 43,44 . In particular, Killing tensors have been applied to find quadratic conservation laws in f(R) cosmology with an ideal gas 45 , Brans-Dicke cosmology 46 , f(R) gravity 47 , a discussion in 48 , a generic model 49 and more recently, Brans-Dicke cosmology with a scalar field 50 .

Contact transformations have considerable advantages over point symmetries in the realm of Newtonian physics and General Relativity, i.e they are closely related to physically meaningful conservation laws such as the Runge-Lenz Vector ⁵¹, the Lewis Invariant ⁵² and conserved quantities in the MICZ Problem ^{53,54}. Moreover, the Carter constant in Kerr spacetime ⁵⁵ stems from dynamical symmetries. Yet another property of the integrals of dynamical symmetries is that they possess more degrees of freedom that allows one to consider numerous scenarios in a given

dynamical problem.

In lieu of this discussion, we emphasize the fact that the determination of Noether contact symmetries is simply a geometric problem. The solution of this problem is provided through differential geometry, which relates the Killing tensors of the metric of the space generated by the dynamic fields to the dynamical symmetries 56 .

In gravitational field theories, the key to this idea is that for some minisuperspace defined by a cosmological model, the Killing tensors are constructed from the model itself. Hence, the existence of Noether contact transformations from a cosmological Lagrangian facilitates a theoretical basis to select functional forms of the model - a selection that conforms to the geometric character of gravity. Another way to determine potential functions is to include some ad hoc assumption, see ^{57,58,59}. In most cases however, the complexity of the field equations is considerable and substantial mechanisms for simplifying differential equations are required in the search for exact solutions of the cosmological models.

Back to our study, for the underlying problem, we define the scalar field interaction in the kinetic and potential part, and require that the field equations are invariant under contact transformations. To begin, we shall explore the application of the Killing tensors of the minisuperspace which generate dynamical symmetries (quadratic conservation laws) for the field equations. The requirement that the field equations admit dynamical symmetries linked to quadratic conservation laws results in the discovery of two potential functions. By implication, these fields are integrable. We obtain the corresponding analytical solution of the field equations by first transforming the fields using normal coordinates. Thereafter, for the second potential, we show that in a special case the scale factor resembles Λ -cosmology and follows an exponential law at late-time.

The structure of the paper follows. In section 2 we consider scalar fields interacting both kinematically and potentially in a spatially flat FRW spacetime. We present some of the elemental properties of the dynamical system such as the Action, the Lagrangian and field equations. In Section 3, we very briefly list all pertinent theory regarding Noether contact symmetries. Section 4 contains the Killing vectors and tensors of the metric space governing the field under study. In Section 4, we determine the potentials of the fields that admit quadratic conservation laws. Section 5 discusses and solves the resulting field equations, and Section 6 concludes.

2. The Cosmological Model

We consider a two-scalar field cosmological model in General Relativity with Action Integral 60,61,62

$$S = \int dx^4 \sqrt{-g} \left[R - \frac{1}{2} g_{ij} H_{AB} \left(\Phi^C \right) \Phi^{A,i} \Phi^{B,i} + V \left(\Phi^C \right) \right], \tag{1}$$

where $A, B, C = 1, 2, H_{AB}$ describes the coupling between the two scalar fields and $\Phi^{C} = (w, z)$.

Notably, the action (1) imbibes a multitude of alternative gravitational theories under a conformal transformation ⁶³. We adopt a spatially flat FRW spacetime $diag(-1, a^2(t)\Sigma^3)$ for the interacting fields, where a(t) is the scale factor. Here Σ^3 is the metric of three dimensional Euclidean space. Hence, the point-like Lagrangian becomes

$$L\left(a,\dot{a},\Phi^{C},\dot{\Phi}^{C}\right) = -3a\dot{a}^{2} + \frac{1}{2}H_{AB}\dot{\Phi}^{A}\dot{\Phi}^{B} - a^{3}V\left(\Phi^{C}\right).$$
(2)

The Lagrangian is holonomic and it admits time translation as a trivial symmetry. There are two considerations for the 2D symmetric metric H_{AB} : one may assume that H_{AB} is flat ⁶⁴ and hence admits three Killing vectors which span the E(2) group, or alternatively, that H_{AB} is a space of constant non-vanishing curvature ⁶⁵, once again admitting three Killing vectors, but span instead the SO(3) group. In the work that follows, we opt to consider the latter case.

Suppose that $h_{AB} = \frac{3}{8}H_{AB}$, where $h_{AB} = diag(1, e^{2z})$, and moreover let $a = (\frac{3}{2})^{\frac{1}{3}}u^{\frac{2}{3}}$.

Therefore the Lagrangian (2) may be written as 65

$$L(u, \dot{u}, w, \dot{w}, z, \dot{z}) = -\frac{1}{2} \dot{u}^2 + \frac{1}{2} u^2 \left(\dot{z}^2 + \dot{w}^2 e^{2z} \right) - u^2 V(w, z).$$
(3)

The field equations are the Hamiltonian⁶⁵

$$E = -\frac{1}{2}\dot{u}^{2} + \frac{1}{2}u^{2}\left(\dot{z}^{2} + \dot{w}^{2}e^{2z}\right) + u^{2}V(w, z), \qquad (4)$$

and the Euler-Lagrange equations ⁶⁵

$$\begin{aligned} \ddot{u} &= -e^{2z}u\dot{w}^{2} - u\dot{z}^{2} + 2uV(z,w), \\ \ddot{w} &= -\frac{2e^{2z}u\dot{w}\dot{z} + 2e^{2z}\dot{u}\dot{w} + V_{,w}u}{e^{2z}u}, \\ \ddot{z} &= -\frac{-e^{2z}u\dot{w}^{2} + V_{,z}u + 2\dot{u}\dot{z}}{u}. \end{aligned}$$
(5)

Eventually, we will show that adhering to the aforementioned discussion results in two cases of the interaction potential. Given the importance of analytical solutions to understand the evolution of the universe, we will use the Noether vectors to lead us to the solution of the field equations.

3. Dynamical Noether Symmetries

There is a fundamental relationship between Killing tensors and conservation laws. To view this relation, consider a particle moving in n dimensional Riemannian space with metric $g_{ij}(x^k)$ under the action of the potential $V(x^k)$. Let $\mathcal{L}(x^k, \dot{x}^k) = \frac{1}{2}g_{ij}\dot{x}^i\dot{x}^j - V(x^k)$, where $\dot{x}^i = \frac{dx^i}{dt}$, be the Lagrangian function that defines a given dynamical system. We define the quadratic function

$$I = X^{i}\left(t, x^{k}, \dot{x}^{k}\right) \frac{\partial \mathcal{L}}{\partial \dot{x}^{i}} - B\left(t, x^{i}\right), \qquad (6)$$

as the conserved quantity for the dynamical system, that is $\frac{d}{dt}I = 0$. I is quadratic and clearly time-independent. The vector field $X^i(t, x^k, \dot{x}^k) \partial_i$ that provides the quadratic conservation law (6) is called a dynamical or contact Noether symmetry for the Lagrangian $\mathcal{L}(x^k, \dot{x}^k)$. If the dependence of the vector field X excludes derivative terms such as \dot{x}^k , then X is called a Noether point symmetry and in this case, the integral I will be linear instead. The functional $B(t, x^i)$ is a boundary term introduced to allow for the infinitesimal changes in the value of the Action Integral.

The dynamical symmetry is given by the expression

$$X^{[1]}\mathcal{L} = \dot{B},\tag{7}$$

where $X^{[1]} = X + \dot{X}^i \partial_i$. Alternatively, if we suppose that

$$X = K^{i}_{\ i}\left(t, x^{k}\right) \dot{x}^{i} \partial_{i},\tag{8}$$

then condition (7) is equivalent to the following system ^{66,56}

$$K_{(ij;k)} = 0, (9)$$

$$K_{ij,t} = 0, \quad B_{,t} = 0,$$
 (10)

$$K^{ij}V_{,i} + B_{,i} = 0, (11)$$

where ';' denotes the covariant derivative with respect to the connection coefficients of the metric g_{ij} . From the conditions (10), we have that $K^i_{j}(x^k)$ and $B(x^k)$, condition (9) implies that $K^i_{j}(x^k)$ is a second-rank Killing tensor of the metric g_{ij} , and lastly, condition (11) can be seen as a constraint condition for the potential $V(x^k)$ in relation to the Killing tensor K^i_{j} and boundary term B.

The use of dynamical Noether symmetries provides first integrals which can be used to reduce the order of the dynamical system and possibly lead to analytic solutions of the field equations. In the present paper, we are interested in contact transformations for which the generator X comprises of a second-rank tensor that is symmetric on the indices, that is, $K_{[ij]} = 0$.

4. Killing Tensors and Killing Vectors

The metric

$$ds^{2} = -du^{2} + u^{2} \left(dz^{2} + e^{2z} dw^{2} \right), \qquad (12)$$

is obtained from the Lagrangian (3), and it is easy to show that the metric is flat. Furthermore, (12) admits a seven dimensional homothetic Lie algebra consisting of the following vector fields:

- Three gradient Killing vectors (translations)

$$T^{1} = \left(\mathrm{e}^{z}w^{2} + \mathrm{e}^{-z}\right)\partial_{u} - 2\frac{w\mathrm{e}^{-z}}{u}\partial_{w} + \frac{-\mathrm{e}^{z}w^{2} + \mathrm{e}^{-z}}{u}\partial_{z},$$

$$T^{2} = -e^{z}w\partial_{u} + \frac{e^{-z}}{u}\partial_{w} + \frac{e^{z}w}{u}\partial_{z},$$
$$T^{3} = -e^{z}\partial_{u} + \frac{e^{z}}{u}\partial_{z}.$$

-Three non-gradient Killing vectors (the rotations) which span the SO(3) algebra

$$R^1 = \partial_w, \ R^2 = \partial_z + w \partial_w, \ R^3 = w \partial_z - \frac{1}{2} \left(w^2 - e^{-2z} \right) \partial_w,$$

and the homothetic vector $u\partial_u$.

To solve the determining conditions (9)-(11), it is necessary to obtain the secondrank symmetric Killing tensors of the metric (12). Once this is established, we will define the potential functions V(w, z) that admit quadratic conservation laws.

For the space of dependent variables $\{u, w, z\}$, in order to obtain the dynamical symmetry (8), we find that the general form of the symmetric second-rank Killing tensor is:

$$K_{ij} = \begin{pmatrix} K_{u}^{u}(u, w, z) \ K_{w}^{u}(u, w, z) \ K_{z}^{u}(u, w, z) \\ K_{w}^{u}(u, w, z) \ K_{w}^{w}(u, w, z) \ K_{z}^{w}(u, w, z) \\ K_{z}^{u}(u, w, z) \ K_{z}^{w}(u, w, z) \ K_{z}^{z}(u, w, z) \end{pmatrix},$$

where (all c_i 's are constants) we define

$$K_{u}^{u}(u, w, z) = e^{2z}c_{9}w^{4} - e^{2z}c_{10}w^{3} + \frac{1}{2}e^{2z}c_{12}w^{2} + e^{2z}c_{13}w + 2c_{9}w^{2} + c_{9}e^{-2z} + e^{2z}c_{14} - c_{10}w - c_{11}, \quad (13)$$

$$K_{w}^{u}(u,w,z) = -\frac{1}{8} u \Big(-2 u e^{3z} w^{4} c_{4} - 2 u e^{3z} w^{3} c_{5} + u e^{3z} w^{3} c_{6} - 16 e^{2z} c_{9} w^{3} + 4 u e^{3z} c_{17} w^{2} - 6 u e^{z} w c_{5} - u e^{z} w c_{6} + 12 e^{2z} c_{10} w^{2} - 4 u e^{3z} c_{18} w + 2 u c_{4} e^{-z} + 4 u e^{z} c_{17} - 4 e^{z} c_{7} u - 4 e^{2z} c_{12} w - 4 u e^{3z} c_{19} - 4 e^{2z} c_{13} - 16 c_{9} w + 4 c_{10} \Big),$$

$$(14)$$

$$K_{z}^{u}(u,w,z) = -\frac{1}{4} \left(2 u e^{z} c_{4} w^{3} - 4 e^{2z} c_{9} w^{4} - u e^{z} w^{2} c_{6} + 4 e^{2z} c_{10} w^{3} + 2 e^{-z} c_{4} u w - 2 u e^{z} c_{7} w - 2 e^{2z} c_{12} w^{2} + 2 e^{-z} c_{5} u - 2 u e^{z} c_{8} - 4 e^{2z} c_{13} w + 4 c_{9} e^{-2z} - 4 e^{2z} c_{14} \right) u, \quad (15)$$

$$K_{w}^{w}(u,w,z) = \frac{1}{8} u^{2} \Big(e^{4z} u^{2}c_{1} w^{4} + 4 e^{4z}c_{2} u^{2} w^{3} - 4 e^{4z}c_{15} u^{2} w^{2} - 8 e^{4z}c_{16} u^{2} w + 8 u e^{3z}c_{4} w^{3} + 8 e^{4z}c_{20} u^{2} + 12 u e^{3z} w^{2}c_{5} - 2 u e^{3z} w^{2}c_{6} - 2 e^{2z} u^{2}c_{1} w^{2} - 16 u e^{3z}c_{17} w + 8 e^{3z}c_{7} u w - 4 e^{2z} u^{2}c_{2} w + 8 u e^{3z}c_{18} + 8 u e^{3z}c_{8} + 4 e^{2z} u^{2}c_{15} + 8 e^{2z}c_{3} u^{2} + 32 e^{2z}c_{9} w^{2} - 16 e^{2z}c_{10} w - 8 u e^{z}c_{4} w + 8 e^{2z}c_{11} + 4 e^{2z}c_{12} + 4 u e^{z}c_{5} + 2 u e^{z}c_{6} + c_{1} u^{2} \Big),$$
(16)

$$K_{z}^{w}(u, w, z) = \frac{1}{8}u^{2} \Big(-2u^{2}e^{2z}c_{1}w^{3} + 2ue^{3z}w^{4}c_{4} - 6u^{2}e^{2z}w^{2}c_{2} + 2ue^{3z}w^{3}c_{5} \\ -ue^{3z}w^{3}c_{6} - 12ue^{z}c_{4}w^{2} + 4u^{2}e^{2z}c_{15}w + 16e^{2z}c_{9}w^{3} - 4ue^{3z}c_{17}w^{2} - 6ue^{z}wc_{5} \\ + 3ue^{z}wc_{6} - 12e^{2z}c_{10}w^{2} + 4u^{2}e^{2z}c_{16} \\ + 4ue^{3z}c_{18}w + 2c_{1}u^{2}w + 2uc_{4}e^{-z} + 4ue^{z}c_{17} + 4e^{2z}c_{12}w + 4ue^{3z}c_{19} \\ + 2c_{2}u^{2} + 4e^{2z}c_{13} - 16c_{9}w + 4c_{10}\Big),$$
(17)

$$K_{z}^{z}(u, w, z) = \frac{1}{2} \left(2 e^{2z} c_{9} w^{4} - 2 e^{2z} c_{10} w^{3} - 2 u e^{z} c_{4} w^{3} + e^{2z} c_{12} w^{2} + u e^{z} w^{2} c_{6} \right. \\ \left. + u^{2} c_{1} w^{2} + 2 e^{-z} c_{4} uw + 2 e^{2z} c_{13} w + 2 u e^{z} c_{7} w + 2 u^{2} c_{2} w + 2 c_{9} e^{-2z} + 2 e^{-z} c_{5} u \right. \\ \left. + 2 e^{2z} c_{14} + 2 u e^{z} c_{8} + 2 u^{2} c_{3} - 4 c_{9} w^{2} + 2 c_{10} w + 2 c_{11} \right) u^{2}.$$

$$(18)$$

To perform this analysis, the field equations must admit contact transformations which are one-parameter transformations in the tangent bundle of the dynamical system. In the next section, we give the form of the potential V(w, z), the admitted contact Noether symmetry and the quadratic conservation laws which follow from the symmetry conditions (9)-(11).

5. Quadratic Conservation Laws

We search for those interactions that admit Noether contact symmetries. These symmetries ensure the existence of conservation laws that are linearly independent and in involution with the Hamiltonian. Stated differently, we require that $\{I, H\} = 0$ where $\{,\}$ is the Poisson bracket, and this mandates that the field equations form an integrable dynamical system.

Furthermore, we emphasize that in 65 , linear conservation laws were found and also some cases where systems of linear equations were obtained. In theory, the

latter will admit quadratic conservation laws. However, in that case the various potentials and solutions have already been reported using point symmetries. It is unnecessary to use contact symmetries to re-derive those potentials.

With the aid of (13)-(18), we show that for specific potential functions V(w, z), where the effective potential is defined as $V_{eff} = u^2 V(w, z)$, the Lagrangian (3) is invariant under contact symmetries. For an arbitrary potential function the field equations do not admit dynamical symmetries hence the field equations are not Noether integrable. In particular, we find that for the gravitational field equations (4)-(5), there are two specific choices of the potential function, for which quadratic in momentum conservation laws exist.

5.0.1. Potential I

In the case where $V(w, z) = V_0 w^2 e^{2z}$, the Lagrangian (3) possesses the dynamical symmetry (8) which is explicitly of the form

$$X^{1} = \left(e^{2z}u^{2}w^{2}\dot{w} - 2\frac{e^{2z}wu^{2}}{e^{2z} + 1}\dot{z}\right)\partial_{w} + \left(-2\frac{e^{2z}wz^{2}}{e^{2z} + 1}\dot{w} + u^{2}\dot{z}\right)\partial_{z},\qquad(19)$$

and admits the quadratic conservation law

$$I^{1} = \left(w^{2} \dot{w}^{2} e^{4z} - 2 w \dot{w} \dot{z} e^{2z} + \dot{z}^{2}\right) u^{4}.$$
(20)

A second contact symmetry is

$$X^{2} = \left(\left(\frac{1}{2} w^{4} e^{2z} + w^{2} + \frac{1}{2} e^{-2z} \right) \dot{u} - 2 \frac{ww \left(w^{2} e^{2z} + 1 \right)}{e^{2z} u^{2} - 1} \dot{w} - \frac{e^{-z} \left(e^{5z} u^{2} w^{4} - e^{3z} w^{4} - e^{z} u^{2} + e^{-z} \right) u}{(e^{2z} u^{2} - 1) \left(u^{2} - 1 \right)} \dot{z} \right) \partial_{u} - \left(2 \frac{w \left(w^{2} e^{2z} + 1 \right)}{e^{2z} u^{2} - 1} \dot{u} + 2w^{2} \dot{w} + 2 \frac{w \left(w^{2} e^{2z} - 1 \right)}{e^{2z} + 1} \dot{z} \right) \partial_{w} - \left(\frac{e^{-z} \left(e^{5z} u^{2} w^{4} - e^{3z} w^{4} - e^{z} u^{2} + e^{-z} \right) u}{(e^{2z} u^{2} - 1) \left(u^{2} - 1 \right)} \dot{u} + 2 \frac{w \left(w^{2} e^{2z} - 1 \right)}{e^{2z} + 1} \dot{w} - \left(\frac{1}{2} \left(-w^{4} e^{4z} + 2 w^{2} e^{2z} - 1 \right) e^{-2z} \right) \dot{z} \right) \partial_{z},$$
(21)

and contact Noether integral

$$I^{2} = \frac{1}{2} \frac{1}{e^{2z}u^{2} - 1} \Big(4 e^{2z}u^{2}w^{2}\dot{w}^{2} - 4 e^{4z}u^{4}w^{2}\dot{w}^{2} + 2 e^{2z}u^{4}w^{2}\dot{z}^{2} - e^{4z}u^{4}w^{4}\dot{z}^{2} \\ + e^{2z}u^{2}w^{4}\dot{z}^{2} + 4 u\dot{u}w\dot{w} - 4 u^{2}w\dot{w}\dot{z} - e^{4z}u^{2}\dot{u}^{2}w^{4} - 2 e^{2z}u^{2}\dot{u}^{2}w^{2} - 2 e^{-2z}u\dot{u}\dot{z} \\ - 4 e^{2z}u^{3}\dot{u}w\dot{w} + 4 e^{2z}u\dot{u}w^{3}\dot{w} + 2 e^{2z}u\dot{u}w^{4}\dot{z} - 2 e^{4z}u^{3}\dot{u}w^{4}\dot{z} - 4 e^{4z}u^{4}w^{3}\dot{w}\dot{z} \\ - 4 e^{4z}u^{3}\dot{u}w^{3}\dot{w} + 4 e^{2z}u^{4}w\dot{w}\dot{z} + 4 e^{2z}u^{2}w^{3}\dot{w}\dot{z} + e^{2z}\dot{u}^{2}w^{4} + 2 u^{3}\dot{u}\dot{z} \\ - 2 u^{2}w^{2}\dot{z}^{2} + e^{-2z}u^{2}\dot{z}^{2} - u^{4}\dot{z}^{2} - u^{2}\dot{u}^{2} + 2 \dot{u}^{2}w^{2} + e^{-2z}\dot{u}^{2} \Big).$$

$$(22)$$

When the contact symmetry (8) is

$$X^{3} = \left(\frac{1}{2}w^{2}e^{2z}\dot{u} - \frac{uwe^{2z}}{e^{2z}u^{2} - 1}\dot{w} - \frac{e^{2z}uw^{2}}{u^{2} - 1}\dot{z}\right)\partial_{u} + \left(-\frac{uwe^{2z}}{e^{2z}u^{2} - 1}\dot{u} - \frac{1}{2}\dot{w} - \frac{we^{2z}}{e^{2z} + 1}\dot{z}\right)\partial_{w} + \left(-\frac{e^{2z}uw^{2}}{u^{2} - 1}\dot{u} - \frac{we^{2z}}{e^{2z} + 1}\dot{w} - \frac{1}{2}w^{2}e^{2z}\dot{z}\right)\partial_{z},$$
(23)

we obtain the quadratic integral

$$I^{3} = \frac{1}{2} \left(-u^{2}w^{2}\dot{z}^{2} + 2V_{0}u^{2}w^{2} - 2u^{2}w\dot{w}\dot{z} - 2uw^{2}\dot{u}\dot{z} - u^{2}\dot{w}^{2} - 2uw\dot{u}\dot{w} - w^{2}\dot{u}^{2} \right) e^{2z},$$
(24)

The Lagrangian admits the symmetry

$$X^{4} = \left(e^{2z}\dot{u} - 2\frac{e^{2z}u}{u^{2} - 1}\dot{z}\right)\partial_{u} + \left(-2\frac{e^{2z}u}{u^{2} - 1}\dot{u} - e^{2z}\dot{z}\right)\partial_{z},$$
(25)

with conservation law

$$I^{4} = \left(-u^{2}\dot{z}^{2} - 2\,u\dot{u}\dot{z} - \dot{u}^{2}\right)e^{2\,z}.$$
(26)

Lastly, the following two dynamical symmetries

$$X^{5} = \left(\frac{u^{2}we^{z} \left(w^{2}e^{2z}+1\right)}{e^{2z}u^{2}-1}\dot{w} - \frac{e^{-z} \left(w^{2}e^{2z}+1\right)u^{2}}{u^{2}-1}\dot{z}\right)\partial_{u} + \left(\frac{u^{2}we^{z} \left(w^{2}e^{2z}+1\right)}{e^{2z}u^{2}-1}\dot{u} + 2e^{z}uw^{2}\dot{w} + \frac{w \left(w^{2}e^{2z}-3\right)e^{z}u}{e^{2z}+1}\dot{z}\right)\partial_{w} + \left(-\frac{e^{-z} \left(w^{2}e^{2z}+1\right)u^{2}}{u^{2}-1}\dot{u} + \frac{w \left(w^{2}e^{2z}-3\right)e^{z}u}{e^{2z}+1}\dot{w} - e^{-z}u \left(w^{2}e^{2z}-1\right)\dot{z}\right)\partial_{z},$$

$$(27)$$

$$X^{6} = \left(\frac{u^{2}we^{3z}}{u^{2}e^{2z}-1}\dot{w} - \frac{e^{z}u^{2}}{u^{2}-1}\dot{z}\right)\partial_{u} + \left(\frac{u^{2}we^{3z}}{u^{2}e^{2z}-1}\dot{u} + \frac{wue^{3z}}{e^{2z}+1}\dot{z}\right)\partial_{w} + \left(-\frac{e^{z}u^{2}}{u^{2}-1}\dot{u} + \frac{wue^{3z}}{e^{2z}+1}\dot{w} - e^{z}u\dot{z}\right)\partial_{z}, \quad (28)$$

admit the next two quadratic conservation laws of the field equations,

$$I^{5} = 2 e^{3z} u^{3} w^{2} \dot{w}^{2} - e^{z} u^{3} w^{2} \dot{z}^{2} + e^{3z} u^{3} w^{3} \dot{w} \dot{z} - e^{z} u^{2} \dot{u} w^{2} \dot{z} + e^{3z} u^{2} \dot{u} w^{3} \dot{w} + e^{-z} u^{3} \dot{z}^{2} - 3 e^{z} u^{3} w \dot{w} \dot{z} - e^{-z} u^{2} \dot{u} \dot{z} + e^{z} u^{2} \dot{u} w \dot{w},$$
(29)

$$I^{6} = \left(e^{2z}uw\dot{w}\dot{z} + e^{2z}\dot{u}w\dot{w} - u\dot{z}^{2} - \dot{u}\dot{z}\right)u^{2}e^{z}.$$
(30)

respectively. As we have seen, this case produces six quadratic conservation laws.

5.0.2. Potential II

,

In the second case, let $V(w, z) = V_0$. This case admits X^1 and X^3 and hence I^1 and I^3 . It also admits X^4 , but a difference in the boundary term provides $\bar{I}^4 = I^4 + 2 V_0 u^2 e^{2z}$. We list the remaining contact symmetries and integrals below.

For the dynamical symmetry

$$X^{7} = \left(\left(w^{4} e^{2z} + 2w^{2} + e^{-2z} \right) \dot{u} - 4 \frac{w \left(w^{2} e^{2z} + 1 \right) u}{e^{2z} u^{2} - 1} \dot{w} - 2 \frac{u \left(w^{4} e^{2z} - e^{-2z} \right)}{u^{2} - 1} \dot{z} \right) \partial_{u} + \left(-4 \frac{w \left(w^{2} e^{2z} + 1 \right) u}{e^{2z} u^{2} - 1} \dot{u} - 4w^{2} \dot{w} - 4 \frac{w \left(w^{2} e^{2z} - 1 \right)}{e^{2z} + 1} \dot{z} \right) \partial_{w} + \left(-2 \frac{u \left(w^{4} e^{2z} - e^{-2z} \right)}{u^{2} - 1} \dot{u} - 4 \frac{w \left(w^{2} e^{2z} - 1 \right)}{e^{2z} + 1} \dot{w} + \left(-w^{4} e^{2z} - e^{-2z} + 2w^{2} \right) \dot{z} \right) \partial_{z},$$

$$(31)$$

we determine that the quadratic integral (6) is of the form

$$I^{7} = -e^{2z}u^{2}w^{4}\dot{z}^{2} + 2w^{4}V_{0}u^{2}e^{2z} - 4e^{2z}u^{2}w^{3}\dot{w}\dot{z} - 2e^{2z}u\dot{u}w^{4}\dot{z}$$

$$-4e^{2z}u^{2}w^{2}\dot{w}^{2} - 4e^{2z}u\dot{u}w^{3}\dot{w} - e^{2z}\dot{u}^{2}w^{4} + 2u^{2}w^{2}\dot{z}^{2} - e^{-2z}u^{2}\dot{z}^{2}$$

$$+4w^{2}V_{0}u^{2} + 4u^{2}w\dot{w}\dot{z} + 2u^{2}V_{0}e^{-2z} + 2e^{-2z}u\dot{u}\dot{z} - 4u\dot{u}w\dot{w}$$

$$-2\dot{u}^{2}w^{2} - e^{-2z}\dot{u}^{2}.$$
 (32)

If the generator is

$$X^{8} = \left(\left(w^{3}e^{2z} + w \right) \dot{u} + \frac{\left(3w^{2}e^{2z} + 1 \right)u}{-e^{2z}u^{2} + 1} \dot{w} - 2\frac{uw^{3}e^{2z}}{u^{2} - 1} \dot{z} \right) \partial_{u} + \left(\frac{\left(3w^{2}e^{2z} + 1 \right)u}{-e^{2z}u^{2} + 1} \dot{u} - 2w\dot{w} - \frac{3w^{2}e^{2z} - 1}{e^{2z} + 1} \dot{z} \right) \partial_{w} + \left(-2\frac{uw^{3}e^{2z}}{u^{2} - 1} \dot{u} - \frac{3w^{2}e^{2z} - 1}{e^{2z} + 1} \dot{w} + \left(-w^{3}e^{2z} + w \right) \dot{z} \right) \partial_{z},$$
(33)

then the integral is

$$I^{8} = -e^{2z}u^{2}w^{3}\dot{z}^{2} + 2w^{3}V_{0}u^{2}e^{2z} - 3e^{2z}u^{2}w^{2}\dot{w}\dot{z}$$
$$-2e^{2z}u\dot{u}w^{3}\dot{z} - 2e^{2z}u^{2}w\dot{w}^{2} - 3e^{2z}u\dot{u}w^{2}\dot{w} - e^{2z}\dot{u}^{2}w^{3} + u^{2}w\dot{z}^{2}$$
$$+2wV_{0}u^{2} + u^{2}\dot{w}\dot{z} - u\dot{u}\dot{w} - \dot{u}^{2}w.$$
(34)

The Lagrangian (3), also admits the contact symmetry

$$X^{9} = \left(we^{2z}\dot{u} + \frac{e^{2z}u}{-e^{2z}u^{2}+1}\dot{w} - 2\frac{we^{2z}u}{u^{2}-1}\dot{z}\right)\partial_{u} + \left(\frac{e^{2z}u}{-e^{2z}u^{2}+1}\dot{u} - \frac{e^{2z}}{e^{2z}+1}\dot{z}\right)\partial_{w} + \left(-2\frac{we^{2z}u}{u^{2}-1}\dot{u} - \frac{e^{2z}}{e^{2z}+1}\dot{w} - we^{2z}\dot{z}\right)\partial_{z},$$
(35)

and Noether integral

$$I^{9} = \left(-u^{2}w\dot{z}^{2} + 2V_{0}u^{2}w - u^{2}\dot{w}\dot{z} - 2uw\dot{u}\dot{z} - u\dot{u}\dot{w} - w\dot{u}^{2}\right)e^{2z}.$$
 (36)

Continuing with our list, we have four more dynamical symmetries

$$X^{10} = \left(\left(\frac{1}{8} u^2 \left(w^4 e^{2z} - 2w^2 + e^{-2z} \right) \right) \dot{w} - \frac{1}{2} \frac{u^2 w \left(w^2 e^{2z} - 1 \right)}{e^{2z} + 1} \dot{z} \right) \partial_w + \left(-\frac{1}{2} \frac{u^2 w \left(w^2 e^{2z} - 1 \right)}{e^{2z} + 1} \dot{w} + \frac{1}{2} w^2 u^2 \dot{z} \right) \partial_z, \quad (37)$$

$$X^{11} = \left(\frac{1}{2}u^2w\left(w^2e^{2z} - 1\right)\dot{w} - \frac{1}{2}\frac{u^2\left(3w^2e^{2z} - 1\right)}{e^{2z} + 1}\dot{z}\right)\partial_w + \left(-\frac{1}{2}\frac{u^2\left(3w^2e^{2z} - 1\right)}{e^{2z} + 1}\dot{w} + u^2w\dot{z}\right)\partial_z,$$
(38)

$$X^{12} = \left(-u^2 \left(w^2 e^{2z} - 1\right) \dot{w} + 2 \frac{w e^{2z} u^2}{e^{2z} + 1} \dot{z}\right) \partial_w + \left(2 \frac{w e^{2z} u^2}{e^{2z} + 1} \dot{w}\right) \partial_z, \qquad (39)$$

$$X^{13} = \left(\frac{1}{2}u^2 w e^{2z} \dot{w} - \frac{1}{2}\frac{e^{2z} u^2}{e^{2z} + 1} \dot{z}\right)\partial_w + \left(-\frac{1}{2}\frac{e^{2z} u^2}{e^{2z} + 1} \dot{w}\right)\partial_z.$$
 (40)

Consequently, the associated Noether contact integrals are, respectively,

$$I^{10} = \frac{1}{8} \left(e^{4z} w^4 \dot{w}^2 - 4 e^{2z} w^3 \dot{w} \dot{z} - 2 e^{2z} w^2 \dot{w}^2 + \dot{w}^2 + 4 w^2 \dot{z}^2 + 4 w \dot{w} \dot{z} \right) u^4, \quad (41)$$

$$I^{11} = -\frac{1}{2} \left(-e^{4z} w^3 \dot{w}^2 + 3 e^{2z} w^2 \dot{w} \dot{z} + e^{2z} w \dot{w}^2 - 2 w \dot{z}^2 - \dot{w} \dot{z} \right) u^4, \qquad (42)$$

$$I^{12} = -\left(w^2 \dot{w} e^{2z} - 2w\dot{z} - \dot{w}\right) u^4 e^{2z} \dot{w},\tag{43}$$

and finally,

$$I^{13} = \frac{1}{2} \left(w \dot{w} e^{2z} - \dot{z} \right) u^4 e^{2z} \dot{w}.$$
 (44)

The next natural step is to find analytical solutions for the two different potentials.

6. Analytical Solutions

In order to proceed with the solution of the system of equations corresponding to each of the two potential functions, some manipulation is necessary. We then write the Euler-Lagrange equations for each potential and solve them taking into account the constraint imposed by the associated Hamiltonian. To proceed, we first apply the transformations ⁶⁵, which give the normal coordinates and are found using the invariant functions associated with the Killing vectors:

$$u^{2} = x^{2} - y^{2} - s^{2},$$

$$w = \frac{s}{x + y},$$

$$z = \ln \frac{x + y}{\sqrt{x^{2} - y^{2} - s^{2}}}.$$
(45)

In the new coordinate system $\{x, y, s\}$ the Lagrangian (3) takes on the simpler form

$$L(x, \dot{x}, y, \dot{y}, s, \dot{s}) = -\frac{1}{2}\dot{x}^2 + \frac{1}{2}\dot{y}^2 + \frac{1}{2}\dot{s}^2 - (x^2 - y^2 - s^2)V(x, y, s).$$
(46)

In the case of Potential I, and if we let $\overline{V}_0 = \sqrt{2V_0}$, the analytical solutions of the dynamical system associated with (46) are

$$s(t) = s_0 \sin(V_0 t) + s_1 \cos(V_0 t),$$

$$x(t) = x_0 t + x_1,$$

$$y(t) = y_0 t + y_1,$$
(47)

with Hamiltonian constraint $E = s_1^2 V_0 + V_0 s_0^2 + \frac{1}{2} y_0^2 - \frac{1}{2} x_0^2$. One can easily transform back into $\{u, w, z\}$ if necessary. The scale factor here is

$$a^{3}(t) = \frac{3}{8} \left(\left(x_{0} t + x_{1} \right)^{2} - \left(y_{0} t + y_{1} \right)^{2} - \left(s_{0} \sin \left(\bar{V}_{0} t \right) + s_{1} \cos \left(\bar{V}_{0} t \right) \right)^{2} \right), \quad (48)$$

and using the singularity condition $a(t \to 0) = 0$, we have the constraint for the constants of integration viz. $\frac{3}{8}(x_1^2 - y_1^2 - s_1^2) = 0$. If we set $s_0 = s_1, x_0 = x_1, y_0 = y_1$, the scale factor (48) reduces to

$$a^{3}(t) = \frac{3}{8}s_{0}^{2}\left(t^{2} + 2t - \sin(2\bar{V}_{0}t)\right).$$
(49)

In turn, the Hubble function (defined as $H = \frac{\dot{a}}{a}$) is

$$H(t) = \frac{2\bar{V}_0 \cos\left(2\bar{V}_0 t\right) - 2t - 2}{-3t^2 + 3\sin\left(2\bar{V}_0 t\right) - 6t},\tag{50}$$

and the deceleration parameter is

$$q(t) = \frac{-2 \left(\cos\left(2\bar{V_0}t\right)\right)^2 \bar{V_0}^2 - 8 \bar{V_0} \left(t+1\right) \cos\left(2\bar{V_0}t\right) + 2 \left(-\bar{V_0}\cos\left(2\bar{V_0}t\right) + t+1\right)^2 + \frac{1}{2} \frac{\left(3 + \left(-6t^2 - 12t\right)\bar{V_0}^2\right) \sin\left(2\bar{V_0}t\right) + t^2 + 6\bar{V_0}^2 + 2t + 4}{\left(-\bar{V_0}\cos\left(2\bar{V_0}t\right) + t+1\right)^2}.$$
(51)

In Figure 1, we explore the general dynamical behaviour of this model. It is clear that H(t) has the largest value at early times and then suddenly declines. The deceleration parameter q(t) starts off with a positive value and then begins to oscillate, $-0.249 \leq q \leq 2$. Oscillating cosmological models have appeared in the literature ^{67,68,69} although different from our model, some showcased the cosmic transit phenomenon signified by a so-called flipping behaviour of the deceleration parameter. These models may be important for the coincidence problem due to periods of acceleration ⁷⁰. Here, we observe that q varies periodically between mostly a decelerated phase (q positive) and an accelerating phase (q negative). As for the equation of state parameter w_{eff} , after some initial volatility, we find $w_{eff} \rightarrow -1$, predicting an equation of state behaviour similar to Λ -cosmology (Λ Cold Dark Matter) model.

On the other hand, in terms of Potential II, we have a more interesting solution, viz.

$$s(t) = s_0 \exp(V_0 t) + s_1 \exp(-V_0 t),$$

$$x(t) = x_0 \exp(\bar{V}_0 t) + x_1 \exp(-\bar{V}_0 t),$$

$$y(t) = y_0 \exp(\bar{V}_0 t) + y_1 \exp(-\bar{V}_0 t),$$

(52)

with Hamiltonian constraint $E = -4 (y_0y_1 - x_0x_1 + s_0s_1)V_0$. The scale factor becomes

$$a^{3}(t) = \frac{3}{8} \left(\left(x_{0} e^{\bar{V}_{0}t} + x_{1} e^{-\bar{V}_{0}t} \right)^{2} - \left(y_{0} e^{\bar{V}_{0}t} + y_{1} e^{-\bar{V}_{0}t} \right)^{2} - \left(s_{0} e^{\bar{V}_{0}t} + s_{1} e^{-\bar{V}_{0}t} \right)^{2} \right),$$
(53)

and the singularity condition $a(t \to 0) = 0$, gives the added constraint $-y_0^2 - y_1^2 + x_0^2 + x_1^2 - s_0^2 - s_1^2 + \frac{E}{2V_0} = 0$.

The Hubble parameter in this case is

$$H(t) = \frac{2\left(e^{4\bar{V_0}t}\left(y_0^2 - x_0^2 + s_0^2\right) - y_1^2 + x_1^2 - s_1^2\right)\bar{V_0}}{\left(3\,y_0^2 - 3\,x_0^2 + 3\,s_0^2\right)e^{4\bar{V_0}t} + \left(6\,y_0\,y_1 - 6\,x_0\,x_1 + 6\,s_0\,s_1\right)e^{2\bar{V_0}t} + 3\,y_1^2 - 3\,x_1^2 + 3\,s_1^2}\tag{54}$$



Fig. 1. Evolution of the scale factor (49), the corresponding Hubble function H(t), the deceleration parameter q(t) and the equation of state parameter w_{eff} . Parameter values are : (a) $s_0 = 1$, $\bar{V}_0 = 0.5$; (b) $\bar{V}_0 = 0.470$; (c) $s_0 = 1$, $\bar{V}_0 = 0.5$; and (d) $\bar{V}_0 = 1$, $s_0 = \sqrt{3}$, $x_0 = 2$, $y_0 = 1$.

and the decelation parameter

$$q(t) = \frac{-10 \left(y_1^2 - x_1^2 + s_1^2\right) \left(y_0^2 - x_0^2 + s_0^2\right) e^{4\bar{V}_0 t} + 6 \left(y_1^2 - x_1^2 + s_1^2\right) \frac{E}{4\bar{V}_0} e^{2\bar{V}_0 t}}{\left(\left(y_0^2 - x_0^2 + s_0^2\right) e^{4\bar{V}_0 t} - y_1^2 + x_1^2 - s_1^2\right)^2} + \frac{6 \left(y_0^2 - x_0^2 + s_0^2\right) \frac{E}{4\bar{V}_0} e^{6\bar{V}_0 t} - \left(y_0^2 - x_0^2 + s_0^2\right)^2 e^{8\bar{V}_0 t} - \left(y_1^2 - x_1^2 + s_1^2\right)^2}{\left(\left(y_0^2 - x_0^2 + s_0^2\right) e^{4\bar{V}_0 t} - y_1^2 + x_1^2 - s_1^2\right)^2}.$$
(55)

A standard calculation also reveals that the equation of state parameter is $w_{eff} = -1$. A clear resemblance to Λ -cosmology can be seen in the special case

of $x_0 = -x_1$, $y_0 = -y_1$, $s_0 = -s_1$, where the scale factor becomes

$$a^{3}(t) = \frac{3}{8} \frac{E}{4V_{0}} \sinh^{2}\left(\bar{V}_{0} t\right).$$
(56)

Recall that the exact solution of the scale factor of Λ -cosmology is

$$a_{\Lambda}^{3}(t) = \left(\frac{\Omega_{m0}}{1 - \Omega_{m0}}\right) \sinh^{2}\left(k \ t\right),$$

where $k = \frac{3}{2}H_0\sqrt{1-\Omega_{m0}}$. This special case identifies the scalar fields z and w to be constant, that is $w(t) = w_0$ and $z(t) = z_0$.

7. Conclusion

The extraordinary Noethers Theorem states that the existence of a group invariant transformation of the field equations corresponds to a conservation flow. Therefore such transformations can be used to recognize or select systems in gravity. The main purpose of this work was to search for quadratic forms of the conserved flows in the geometric background of multi-scalar fields interacting in their kinematic and potential parts in a spatially flat FRW spacetime. We have determined the interactions for which the field dynamical system is Liouville integrable, according to the presence of dynamical symmetries. Within the cosmological model, we assumed that the 2D metric H_{AB} that characterizes the interaction in the field space is a space of constant non-vanishing curvature. Specifically, the metric space admits three Killing vectors which span the SO(3) group. We used the the existence of Noether symmetry contact transformations to select the unknown form of the potential functions. A detailed examination of the Killing tensors of the minisuperspace led us to the discovery of two distinct cases of the potential. The first admitted six quadratic conservation laws, and the second case produced a total of ten quadratic conservation laws. These quadratic integrals are in involution with the Hamiltonian function, i.e. by implication this property conveys that the field equations are integrable. Once we established the integrability of the system, the cosmological solutions were found by quadrature for each case. Additionally, we first transformed the dynamical system into simpler but equivalent systems by constructing and applying a set of normal coordinates. In the first case, the scale factor of the universe appeared to be power-law, and for the case of the second potential, exponential-law. Finally, some numerical simulations were performed in order to study the Hubble, deceleration and equation of state parameter.

Acknowledgments: We acknowledge the financial support from the National Research Foundation of South Africa (99279).

References

- 1. A. Riess et al., Astron. J. 116 (1998) 1009.
- 2. S. Perlmutter et al., Astrophys. J. 517 (1999) 565.
- 3. M. Tegmark et al., Astrophys. J. 606 (2004) 702.

- 4. E. Komatsu et al., Astrophys. J. Suppl. Ser. 180 (2009) 330.
- 5. P. A. R. Ade et al., (Planck Collaboration) (2015) arXiv:1502.01589
- 6. J.D. Barrow, P. Saich, Class. Quant. Grav. 10 (1993) 279.
- 7. B. Ratra, P.J.E Peebles, Phys. Rev. D 37 (1988) 3406.
- 8. M.C. Bento, O. Bertolami, A.A. Sen, Phys. Rev. D 66 (2002) 043507.
- S. Pan, S. Bhattacharya, S. Chakraborty, Mon. Not. Roy. Astron. Soc. 452 (2015) 3038.
- 10. R. Ferraro, F. Fiorini, Phys. Rev. D 78 (2008) 124019.
- 11. R. Maartens, Living Rev. Rel. 7 (2004) 7.
- 12. T.P. Sotiriou, V. Faraoni Rev. Mod. Phys. 82 (2010) 451.
- 13. A.B. Burd, J.D. Barrow, Nucl. Phys. B (1988) 929.
- 14. A. A. Coley, J. Ibáñez, R. J. van den Hoogen, J. Math. Phys. 38 (1997) 5256.
- 15. K.A. Olive, Phys. Rep. 190 (1990) 308.
- 16. C. Wetterich, Nucl. Phys. B 302 (1988) 668.
- 17. E. J. Copeland, A. R. Liddle, D. Wands, Phys. Rev. D 57 (1998) 4686.
- 18. Z.-K. Guo, Y.-S. Piao, Y.-Z. Zhang, Phys. Lett. B 568 (2003) 1.
- 19. E. J. Copeland, A. Mazumdar, N. J. Nunes, Phys. Rev. D 60 (1999) 083506.
- 20. S. A. Kim, A. R. Liddle, S. Tsujikawa, Phys. Rev. D 72 (2005) 043506.
- 21. A.G. Muslimov, Class. Quantum Grav. 7 (1990) 231.
- 22. G.F.R. Ellis, M.S. Madsen, Class. Quantum Grav. 8 (1991) 667.
- 23. R. Easther, Class. Quantum Grav. 10 (1993) 2203.
- 24. H.C. Kim, Mod. Phys. Lett. A28 (2013) 1350089.
- 25. S.V. Chervon, R.R. Abbyazov, S.V. Kryukov, Russ. Phys. J. 58 (2015) 597.
- 26. A. Paliathanasis, G. Leon, S. Pan, Gen. Rel. Grav. 51 (2019) 106.
- 27. S.V. Chervon, Quant. Matt. 2(2) (2013) 71.
- 28. P. Christodoulidis, D. Roest, E. I. Sfakianakis, Angular inflation in multi-field α -attractors. Pre-print: arXiv:1803.09841.
- P. Christodoulidis, Probing the inflationary evolution using analytical solutions. Preprint: arXiv:1811.06456.
- H. Stephani, Differential Equations: Their Solutions Using Symmetry, Cambridge University Press, New York, 1989.
- 31. M.C. Kewyama, K.S. Govinder, S.D. Maharaj, J. Math. Phys., 53 (2012) 033707.
- S. Capozziello, E. Piedipalumbo, C. Rubano, P. Scudellaro, Phys. Rev. D. 80 (2009) 104030.
- 33. S. Cotsakis, P.G.L. Leach, H. Pantazi, Grav. Cosm. 4 (1998) 314.
- 34. B. Vakili, Phys. Lett. B 669 (2008) 209.
- 35. S. Basilakos, M. Tsamparlis, A. Paliathanasis, Phys. Rev. D 83 (2011) 103512.
- 36. R. C. de Souza, R. Andre, G. M. Kremer, Phys. Rev. D 87 (2013) 083510.
- A. K. Sanyal, B. Modak, C. Rubano, E. Piedi- palumbo, Gen. Rel. Grav. 37 (2005) 407.
- N. Dimakis, A. Giacomini, S. Jamal, G. Leon, A. Paliathanasis, Phys. Rev. D 95 (2017) 064031.
- 39. S. Jamal, G. Shabbir, Eur. Phys. J. Plus 132 (2017) 70.
- 40. S. Jamal, Gen. Rel. Grav. 49(88) (2017) 1.
- 41. S. Jamal, A. Paliathanasis, J. Geom. Phys. 117 (2017) 50.
- A. Giacomini, S. Jamal, G. Leon, A. Paliathanasis, J. Saavedra, Phys. Rev. D 95 (2017) 124060.
- 43. K. Rosquist, C. Uggla, J. Math. Phys. 32 (1991) 3412.
- 44. A. Paliathanasis, M. Tsamparlis, S. Basilakos, Phys. Rev. D 90 (2014) 103524.
- 45. G. Papagiannopoulos, Spyros Basilakos, J.D. Barrow, A. Paliathanasis, Phys. Rev. D

97 (2018) 024026.

- 46. G. Papagiannopoulos, J.D. Barrow, S. Basilakos, A. Giacomini, A. Paliathanasis, Phys. Rev. D 95 (2017) 024021.
- 47. A. Paliathanasis, Class. Quant. Grav. 33(7) (2016).
- 48. L. Karpathopoulos, M. Tsamparlis, A. Paliathanasis, J. Geom. Phys. 133 (2018) 279.
- 49. M. Tsamparlis, A. Paliathanasis, Arab. J. Math. 8(3) (2019) 201.
- 50. A. Paliathanasis, Gen. Rel. Grav. 51 (2019) 101.
- 51. R. C. OConnell, K. Jagannathan, Am. J. Phys. 71 (2003) 243.
- 52. H. R. Jr. Lewis, Phys. Rev. Lett. 18 (1967) 510.
- 53. D. Zwanziger, Phys. Rev. 176 (1968) 1480.
- 54. H. McIntosh, A. Cisneros, J. Math. Phys. 11 (1970) 896.
- 55. B. Carter, Phys. Rev. 174 (1968) 1559.
- 56. T.M. Kalotas, B.G. Wybourne, J. Phys. A: Math. Gen. 15 (1982) 2077.
- 57. D. Bertacca, S. Matarrese, M. Pietroni, Mod. Phys. Lett. A 22 (2007) 2893.
- 58. J. A. Frieman, C. T. Hill, A Stwebbins, I. Waga, Phys. Rev. Lett. 75 (1995) 2077.
- 59. V. Sahni, L.M. Wang, Phys. Rev. D 62 (2000) 103517.
- 60. N. Tamanini, C.G. Böhmer, Phys. Rev. D 87 (2013) 084031.
- 61. J.G. Bellido, D. Wands, Phys. Rev. D 53 (1996) 5437.
- 62. S. Cremonini, Z. Lalak, K. Turzyński, JCAP 03 (2011) 015.
- 63. A.A. Starobinsky, S. Tsujikawa, J. Yokoyama, Nucl. Phys. B 610 (2001) 383.
- A. Aslam, M. Jamil, D. Momeni, R. Myrzakulov, M. A. Rashid, M. Raza, Astrophys. Space Sci. 348 (2013) 533.
- 65. A. Paliathanasis, M. Tsamparlis, Phys. Rev. D 90 (2014) 043529.
- 66. M. Crampin, Rep. Math. Phys. 20 (1984) 31.
- 67. M. Shen, L. Zhao, Chin. Phys. Lett. 31 (2014) 010401.
- 68. S. Dodelson, M. Kaplinghat, E.Stewart, Phys. Rev. Lett. 85 (2000) 5276.
- 69. S. Nojiri, S. D. Odintsov, Phys. Lett. B 637 (2006) 139.
- 70. P.K. Sahoo, S.K. Tripathy, P. Sahoo, Mod. Phys. Lett. A, 33 (2018) 1850193.