

Is the Hubble tension a hint of AdS phase around recombination?

Gen Ye^{1*} and Yun-Song Piao^{1,2†}

¹ School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China and

² Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100190, China

Anti-de Sitter (AdS) vacua, being theoretically important, might have an unexpected impact on the observable universe. We find that in early dark energy (EDE) scenarios the existence of AdS vacua around recombination can effectively lift the CMB-inferred H_0 value. As an example, we study a phenomenological EDE model with an AdS phase starting at the redshift $z \sim 2000$ and ending shortly after recombination (hereafter the universe will settle down in a $\Lambda > 0$ phase until now), and obtain a best-fit $H_0 = 72.74$ km/s/Mpc without degrading the CMB fit compared with the standard Λ CDM model.

Recently, the tension between the Hubble constant H_0 measured locally and that deduced from the best-fit of Λ CDM to the cosmic microwave background (CMB) observation has acquired extensive attention, e.g. [1, 2] for reviews. Based on the Λ CDM model, the Planck collaboration inferred $H_0 = 67.36 \pm 0.54$ km/s/Mpc [3]. Using Cepheids-calibrated supernovae, Riess et.al (the SH0ES team) reported the Hubble rate $H_0 = 74.03 \pm 1.42$ km/s/Mpc [4], which is at 4.4σ discrepancy compared with that inferred by Planck. The large $H_0 (> 70)$ value is also supported by other local measurements [5–8]. Currently, it is probably not suitable to simply explain this discrepancy by systematic errors in the data [1]. It is thus increasingly likely that new physics beyond the Λ CDM model plays a role in resolution of the Hubble tension.

Theoretically, one possibility is modifying post-recombination physics, such as the dark energy or modified gravity models, e.g. [9–20]. Such solutions are constrained tightly by late-time observations [1]. Another possibility is modifying prerecombination physics (modifying the sound horizon $r_s^* = \int_{z^*}^{\infty} c_s/H(z)dz$ [1, 21, 22]), such as early dark energy [23–29], see also [30], neutrino self-interaction [31, 32], see also [33–35], and decaying dark matter [36–38]. See also attempts concerning CMB non-Gaussianities [39] and fundamental constants [40]. Determination of H_0 requires fitting the integral expression $D_A^* = \int_0^{z^*} \frac{dz}{H(z)} = r_s^*/\theta_s^*$, D_A^* being the angular diameter distance to the last-scattering surface. While $\theta_s^* \equiv r_s^*/D_A^*$ is precisely determined by CMB peak spacing, a smaller r_s^* will eventually lead to a larger H_0 .

It is well known that anti-de Sitter (AdS) vacua are theoretically important. AdS vacua naturally emerge from the string theory. One might uplift AdS to de Sitter (dS) vacua by the KKLT mechanism [41, 42], which inspired the “landscape” idea [43]. The landscape consists of all effective field theories (EFTs) with consistent UV completion (otherwise the EFT is said to be in the swampland), which might be from various compactifications of the string theory. As the swampland criteria for EFTs, the distance conjecture [44] and the dS con-




FIG. 1: A sketch of the potential with an AdS phase. Initially the field is frozen at ϕ_i . It starts rolling down the potential at the redshift $z \sim 3500$ when H drops below its effective mass m_ϕ , and enters an AdS phase at $z \sim 2000$. The field straightly rolls over the AdS region and does not oscillate. It climbs up to the $\Lambda > 0$ region shortly after recombination, hereafter the universe is effectively described by the standard Λ CDM model.

jecture $M_p |\nabla_\phi V|/V > c \sim \mathcal{O}(1)$ [45] (or the refined dS conjecture [46, 47]) have been proposed, which seems to throw dS vacua into the swampland. However, whether metastable dS vacua exists in the landscape or not, AdS vacua should be indispensable. Thus it is significant to ask if AdS vacua have any impact on the observable universe.

We will show this possibility. A novelty of our result is that recombination might happen in the AdS vacuum¹, which may be tested by near-future CMB experiments. In the early dark energy (EDE) scenario [23, 24, 27], modification of r_s is implemented by an EDE scalar field that starts to activate a few decades before recombination. It is the energy injection of this EDE field that results in a reduced r_s^* , so an increased H_0 . We will focus on the EDE scenario with an AdS phase, and find that such an

*yegen14@mailsucas.ac.cn

†yspiao@ucas.ac.cn

¹ The possibility of late-time AdS has also been studied in Ref. [16, 19].

AdS phase will make the EDE injection more efficient while ensuring that it redshifts fast enough around recombination without spoiling the fit to the CMB data.

The scenario we consider is presented in Fig-1. Initially, the scalar field ϕ sits at the hillside of its potential, and its energy density ρ_ϕ is negligible. As the universe expands, radiation and matter are diluted. When $H^2 \simeq \partial_\phi^2 V$, which occurred before recombination, the field starts rolling down the potential, meanwhile ρ_ϕ becomes non-negligible. Then the field will roll over an AdS phase, and during this period ρ_ϕ quickly redshifts away. Hereafter, the field rapidly climbs up to the $\Lambda > 0$ region, so that the universe eventually settles down in the Λ CDM phase until now. See also [48, 49] for the potential with multiple AdS vacua.

One has $\rho_\phi = \dot{\phi}^2/2 + V(\phi)$ and $P_\phi = \dot{\phi}^2/2 - V(\phi)$, respectively. In the AdS phase, $w = P_\phi/\rho_\phi > 1$. When the EDE field rolls down to $V < 0$, we have $w > 1$, so that ρ_ϕ redshifts very rapidly $\rho_\phi \sim a^{-3(1+w)}$ (in Refs.[23, 24, 27] the EDE dissipates less effectively by oscillation with cycle-averaged $w < 1$). This is crucial for getting a larger H_0 , since if ρ_ϕ is non-negligible around recombination, it will interfere with the fit of Λ CDM to the CMB data [2]. Perturbations are also under control since a canonical scalar field always has $c_s^2 = 1$.

As an example, we model Fig-1 as

$$V(\phi) = \begin{cases} V_0 \left(\frac{\phi}{M_p}\right)^4 - V_{ads}, & \frac{\phi}{M_p} < \left(\frac{V_{ads}}{V_0}\right)^{1/4} \\ 0, & \frac{\phi}{M_p} > \left(\frac{V_{ads}}{V_0}\right)^{1/4} \end{cases} \quad (1)$$

where V_{ads} depicts the depth of AdS well. Here, the initial value ϕ_i of ϕ should satisfy $|\phi_i| < M_P$, so that the model is consistent with the swampland conjectures [44, 45]. We also allow for a cosmological constant $\Lambda \simeq (10^{-4} eV)^4 > 0$ (but not included) in (1) to ensure that the universe eventually settles down in the Λ CDM phase. It is possible to replace the positive cosmological constant in Λ CDM with quintessence, or some effective parameterization, e.g.[30]. This may (but does not essentially) change the fit result. When $V_{ads} = 0$, (1) corresponds to a run-away potential, see [26] for the relevant study.

Three new parameters $\{V_0, V_{ads}, \phi_i\}$ are added to the standard six parameters $\{\omega_b, \omega_{cdm}, H_0, \ln(10^{10} A_s), n_s, \tau_{reio}\}$ of Λ CDM, noting initially $\dot{\phi}_i = 0$. Instead of $\{V_0, V_{ads}, \phi_i\}$, we will adopt another set of parameters $\{z_c, \omega_{scf}, \alpha_{ads}\}$ with clearer physical interpretation [23, 24]. z_c is the redshift at which the EDE field starts rolling, which is defined by $\partial_\phi^2 V(\phi_c) = 9H^2(z_c)$, $\phi_c \equiv \phi(z_c)$ [50]. ω_{scf} is the energy fraction of the EDE field at z_c . α_{ads} is related to V_{ads} by $V_{ads} = \alpha_{ads}(\rho_m(z_c) + \rho_r(z_c))$.

In the code, one should search for the $\{V_0, V_{ads}\}$ corresponding to a given set of $\{z_c, \omega_{scf}\}$. A shooting method is used to accomplish this. Exploiting the fact




FIG. 2: Energy fraction f_{EDE} of EDE with respect to redshift z , plotted using the best-fit models. The scalar field energy density quickly redshifts away after the field starts rolling, so the recombination redshift z_{rec} is nearly the same in both EDE models and the standard Λ CDM. Scalar field energy in the $\phi^4 + AdS$ model redshifts much faster due to the AdS phase (shaded region, from $z = 2063$ to $z = 802$ in the best-fit model).

$\dot{\phi}_i = 0$ before z_c , we get the initial guess by solving $\partial_\phi^2 V(\phi_i) = 9H_c^2$ and $V(\phi_i) = 3\omega_{scf}M_p^2H_c^2$, where $H_c^2 \equiv (\rho_m(z_c) + \rho_r(z_c))/3M_p^2$. Then we search for the exact scalar field parameters by iteratively varying $V_{0,MC}$ and $\phi_{i,MC}$ and calculating the corresponding $\omega_{scf,MC}$ with numeric integration.

We modified the MontePython-V3.2 [51, 52] and CLASS codes [53, 54] to perform a Markov chain Monte Carlo (MCMC) analysis on the 6+2 parameter set $\{\omega_b, \omega_{cdm}, H_0, \ln(10^{10} A_s), n_s, \tau_{reio}, z_c, \omega_{scf}\}$. Our datasets include Planck2018 high- l and low- l TT, EE, TE and lensing likelihoods [55]. We follow the convention used by Planck for the three neutrinos species. We use BAO measurement from the CMASS and LOWZ of BOSS DR12 [56] as well as low- z BAO measurements from 6dFGS [57] and MGS of SDSS [58]. The Pantheon [59] dataset with a single nuisance M , which includes luminosity distance of 1048 SN Ia, is also included. We use the latest result $H_0 = 74.03 \pm 1.42$ km/s/Mpc from the SH0ES team [4] for local measurement.

One should also vary α_{ads} in the MCMC analysis. However, variation of α_{ads} will drastically worsen convergence of the chain. This is due to background integration divergence whenever the field fails to climb up the potential. As a consequence, the chain will head to low α_{ads} even though better fit to data favors a higher one. In principle, one should construct a compatible phase space measure to account for this effect and adjust the step length accordingly. Here, we will instead take a shortcut to simply fix α_{ads} to its best-fit value $\alpha_{ads} = 3.79 \times 10^{-4}$, which is enough for our purpose.

The marginalized posterior distributions of $\{H_0, n_s, \omega_{scf}, \ln(1 + z_c)\}$ are shown in Fig-5. As expected, the energy injection ω_{scf} is positively correlated with H_0 . The mean and best-fit values of all model parameters are reported in Table-I. Table-II reports the best-fit χ^2 value of each individual experiment. The




FIG. 3: The 1-sigma contour plot of H_0 versus ω_{scf} and $\ln(1+z_c)$. The models compared are oscillating ϕ^4 (red), $\alpha_{ads} = 0$ (blue) and $\alpha_{ads} = 3.79 \times 10^{-4}$ (green). The colored bands represent the 1-sigma H_0 in Λ CDM (gray) and SH0ES measurement (orange).

AdS model fits the CMB data slightly better than other models. We refer to our model (1) as the ϕ^4 +AdS model for simplicity. We also include a Λ CDM model, an oscillating ϕ^4 model [24] in which the EDE potential is $V(\phi) \sim \phi^4$ and ρ_ϕ redshifts away when ϕ oscillates, and a ϕ^4 +AdS model in the $\alpha_{ads} = 0$ limit (equivalently $V_{ads} = 0$ in (1)) for comparison. In the ϕ^4 +AdS model, the best-fit H_0 has been significantly uplifted (as opposed to other models) to 72.74 km/s/Mpc, in agreement with the local measurements at 1σ level.

In Fig-2, we plot the evolution of $f_{EDE} = \rho_\phi / \rho_{tot}$ with respect to the redshift for the best-fit models. The field thaws at $z = z_c$, quickly reaches the maximum of f_{EDE} , and then ρ_ϕ rapidly redshifts away. Though more energy is injected in the ϕ^4 +AdS model, f_{EDE} at recombination is far smaller, since the existence of an AdS phase makes the dissipation of ρ_ϕ more effective. Our best-fit model suggests that the recombination happened during the AdS phase (or in AdS vacuum). To further illustrate the power of the AdS phase, we compare the oscillating ϕ^4 model, the $\alpha_{ads} = 0$ model and the $\alpha_{ads} = 3.79 \times 10^{-4}$ model in Fig-3. We see that though V_{ads} only takes up a quite small fraction [noting $V_{ads} = \alpha_{ads}(\rho_m + \rho_r)$], its impact on H_0 is quite remarkable.

Here, the spectrum index n_s and the amplitude A_s of primordial perturbations are larger than those in Λ CDM. This is a common phenomenon in EDE scenarios, which has been also observed in Refs.[23, 24, 27]. In particular, n_s seems to be positively correlated with H_0 , see Fig-5. It could be understood, at least partially, by the integrated Sachs-Wolfe (ISW) effect [60]. The gravitational potential Φ can be converted to the density perturbation δ through the Poisson equation $\nabla^2 \Phi = 4\pi G \rho \delta$, thus the ISW contribution to the CMB angular power spectrum is

$$(C_l)_{SW} \propto \int_0^\infty \frac{dk}{k} \mathcal{P}_\Phi(k) j_l^2(kD_A^*), \quad (2)$$

where $\mathcal{P}_\Phi(k)$ is the power spectrum of primordial perturbations in the Newtonian gauge. Considering $\mathcal{P}_\Phi =$




FIG. 4: Difference between various models fitted to full datasets and a reference Λ CDM model obtained using only the Planck2018 data. The upper panel is for the TT spectrum and the lower one for the EE spectrum.

$$A_s \left(\frac{k}{k_{pivot}} \right)^{n_s-1}, \text{ we can explicitly integrate out Eq.(2),} \\ (C_l)_{SW} \propto A_s (D_A^* k_{pivot})^{1-n_s} f(n_s). \quad (3)$$

In particular, $f(n_s)$ is monotonically increasing with respect to n_s . In the EDE models, the reduction in r_s^* causes a reduction in D_A^* , and so the $(D_A^* k_{pivot})^{1-n_s}$ term in Eq.(3) (for $n_s < 1$). Since C_l is fixed by the CMB observation, $A_s f(n_s)$ must be larger accordingly, eventually leading to larger n_s and A_s .

We plot the difference $\Delta C_l / C_{l,ref}$ in Fig-4, where $\Delta C_l = C_{l,model} - C_{l,ref}$ and $C_{l,ref}$ is a reference Λ CDM model obtained using only the Planck2018 dataset. The models compared are Λ CDM (fitted to the full datasets), oscillating ϕ^4 [24] and ϕ^4 +AdS with $\alpha_{ads} = 0$ [equivalently $V_{ads} = 0$ in (1)] and $\alpha_{ads} = 3.79 \times 10^{-4}$. It is observed that as l becomes large, compared with ϕ^4 and Λ CDM models, the TT spectrum in the ϕ^4 +AdS model will go upwards. The near-future ground based CMB experiments [61–64] are expected to probe up to $l = 5000$ with order of magnitude improvement in precision at high l . The improved constraining power from high l will significantly help distinguishing different models (especially those with different n_s and A_s). Another potentially observable signal is the EE spectrum around $l \sim 200$, which shows itself a bump for the ϕ^4 +AdS model. The binned Planck residue at $l = 165$ has 5% relative uncertainty. The $l = 200$ bump in the AdS model shows roughly 5%

| Parameter | Λ CDM | Oscillating ϕ^4 | $\phi^4 + \text{AdS}$ | |
|--------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|
| | | | $\alpha_{ads} = 0$ | $\alpha_{ads} = 3.79 \times 10^{-4}$ |
| 100 ω_b | 2.247(2.224) $^{+0.015}_{-0.014}$ | 2.281(2.29) $^{+0.018}_{-0.02}$ | 2.301(2.289) $^{+0.02}_{-0.022}$ | 2.346(2.354) $^{+0.017}_{-0.016}$ |
| ω_{cdm} | 0.1182(0.1183) $^{+0.0008}_{-0.0013}$ | 0.1267(0.1256) $^{+0.0038}_{-0.0048}$ | 0.1275(0.1262) $^{+0.0031}_{-0.0027}$ | 0.134(0.1322) $^{+0.0019}_{-0.0021}$ |
| H_0 | 68.16(68.23) $^{+0.56}_{-0.4}$ | 70.73(70.78) $^{+0.91}_{-1.3}$ | 70.78(70.27) $^{+0.76}_{-0.71}$ | 72.64(72.74) $^{+0.57}_{-0.64}$ |
| $\ln(10^{10} A_s)$ | 3.049(3.054) $^{+0.013}_{-0.016}$ | 3.064(3.064) $^{+0.017}_{-0.018}$ | 3.066(3.058) $^{+0.014}_{-0.017}$ | 3.077(3.074) $^{+0.015}_{-0.015}$ |
| n_s | 0.9688(0.9696) $^{+0.0039}_{-0.0042}$ | 0.9788(0.9798) $^{+0.006}_{-0.0064}$ | 0.9842(0.9805) $^{+0.005}_{-0.0057}$ | 0.9976(0.9974) $^{+0.0046}_{-0.0045}$ |
| τ_{reio} | 0.0604(0.0636) $^{+0.0066}_{-0.0075}$ | 0.0588(0.0575) $^{+0.007}_{-0.0084}$ | 0.0596(0.0573) $^{+0.0086}_{-0.0084}$ | 0.0574(0.0598) $^{+0.0075}_{-0.0078}$ |
| ω_{scf} | - | 0.0666 $^{+0.029}_{-0.035}$ | 0.067(0.055) $^{+0.018}_{-0.015}$ | 0.113(0.107) $^{+0.005}_{-0.009}$ |
| $\ln(1+z_c)$ | - | 8.347(8.197) $^{+0.11}_{-0.27}$ | 8.28(8.17) $^{+0.12}_{-0.13}$ | 8.22(8.21) $^{+0.072}_{-0.079}$ |
| 100 θ_s | 1.0422(1.0421) $^{+0.0005}_{-0.0004}$ | 1.0415(1.0417) $^{+0.0004}_{-0.0004}$ | 1.0414(1.0415) $^{+0.0006}_{-0.0005}$ | 1.0411(1.0411) $^{+0.0003}_{-0.0003}$ |
| σ_8 | 0.8078(0.81) $^{+0.0054}_{-0.0066}$ | 0.8368(0.8354) $^{+0.011}_{-0.011}$ | 0.835(0.8297) $^{+0.0105}_{-0.0089}$ | 0.8571(0.8514) $^{+0.0079}_{-0.0077}$ |

TABLE I: The mean values and 1σ error of all cosmological and model parameters. Best-fit values are given in the parenthesis. The $\phi^4 + \text{AdS}$ models are labeled by their α_{ads} values. All models are obtained using the same datasets, nuisance priors and precision settings.

| Experiment | Λ CDM | Oscillating ϕ^4 | $\phi^4 + \text{AdS}$ | |
|----------------|---------------|----------------------|-----------------------|----------------------------|
| | | | $\alpha_{ads} = 0$ | $10^4 \alpha_{ads} = 3.79$ |
| χ^2_{CMB} | 2778.7 | 2782.1 | 2777 | 2776.5 |
| BAO low- z | 2.2 | 2.1 | 1.8 | 2.2 |
| BAO high- z | 1.8 | 1.9 | 1.9 | 2.1 |
| Pantheon | 1026.9 | 1027.5 | 1026.9 | 1026.9 |
| SH0ES | 15.4 | 4.9 | 7 | 0.8 |

TABLE II: The best-fit χ^2 per experiment.

difference compared with the blue Λ CDM line thus it is now invisible to the Planck observation. However, the upcoming CMB-S4, for example, is expected to reduce uncertainty by 50% over Planck and hence might be able to probe this signature.

In summary, we showed that the Hubble tension might be telling us the existence of AdS vacua around recombination. Through studying a phenomenological EDE model, we found that such an AdS phase ($|V_{ads}| \sim (0.1eV)^4$) can lift the CMB-inferred H_0 to $H_0 = 72.64^{+0.57}_{-0.64}$ km/s/Mpc, within 1σ range of the local measurement [4], and significantly alleviate the Hubble tension. A novelty of our result is that the recombination happens in the AdS vacuum, which also makes unique predictions accessible to near-future CMB experiments. It should be pointed out that (1) is only applied for illustrating the phenomenology of EDE with an AdS vacuum, and realistic potentials may be more complex, which might fit better to data. Though the model we consider is quite simplified, it highlights an unexpected point that AdS vacua, ubiquitous in consistent UV-complete theories, might also play a crucial role in our observable universe.

In order to have the EDE field start rolling at around

$z \sim 3000$ and be non-negligible, one must fine-tune the potential such that $V_0 \sim (eV)^4$. This fine-tuning guarantees that the field thaws at the right time, otherwise it will not help resolving the Hubble tension. It also should be pointed out that on CMB probes, we only consider the Planck data. The H_0 tension is predominantly only present in Planck data and other CMB observatories do not see it in that magnitude. Our result might be weakened if there really are unknown instrumental effects or systematics present in the Planck experiment. It will be important to test our model against other CMB datasets, e.g. WMAP, SPT and ACT, which will be studied elsewhere. Multiprobe combination constraints on EDE models with constant EDE fraction are also discussed in Ref.[65].

The issues worth studying are as follows. A well-explored conjecture is that AdS vacua is very likely to be accompanied by an infinite tower of ultralight states [44, 66, 67] (this effect also has recently been applied to the Hubble tension [38, 68]). It is quite intriguing to explore whether these additional light states have left any imprints on the last-scattering surface. It has recently been proposed in [49] that a multistage inflation, consisting of multiple inflationary phases separated by AdS vacua, may survive the swampland conjectures. AdS-like potentials also appear in nonsingular cosmological models [69]. Confronting these ideas with the hint from the Hubble tension that the recombination era might happen in an AdS phase, it is quite interesting to wonder if our universe actually has passed through many phases with different AdS vacua.

Acknowledgment We thank Sunny Vagnozzi for valuable comments. G.Y. thanks Yu-Tong Wang for his help on coding. Y.S.P. is supported by NSFC, No. 11575188 and No. 11690021.




FIG. 5: Marginalized posterior distributions of $\{H_0, n_s, \omega_{scf}, \ln(1+z_c)\}$. H_0 is correlated with n_s and ω_{scf} , as explained in the text.

- [2] L. Knox and M. Millea, Phys. Rev. D **101**, no. 4, 043533 (2020) doi:10.1103/PhysRevD.101.043533 [arXiv:1908.03663 [astro-ph.CO]].
- [3] N. Aghanim *et al.* [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO].
- [4] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri and D. Scolnic, Astrophys. J. **876**, no. 1, 85 (2019) [arXiv:1903.07603 [astro-ph.CO]].
- [5] G. C.-F. Chen *et al.*, arXiv:1907.02533 [astro-ph.CO].
- [6] K. C. Wong *et al.*, arXiv:1907.04869 [astro-ph.CO].
- [7] W. L. Freedman *et al.*, arXiv:1907.05922 [astro-ph.CO].
- [8] C. D. Huang *et al.*, arXiv:1908.10883 [astro-ph.CO].
- [9] E. Di Valentino, A. Melchiorri and O. Mena, Phys. Rev. D **96**, no. 4, 043503 (2017) [arXiv:1704.08342 [astro-ph.CO]].
- [10] E. Di Valentino, A. Melchiorri, E. V. Linder and J. Silk, Phys. Rev. D **96**, no. 2, 023523 (2017) [arXiv:1704.00762 [astro-ph.CO]].
- [11] E. O. Colgain, M. H. P. M. van Putten and H. Yavartanoo, Phys. Lett. B **793**, 126 (2019) [arXiv:1807.07451 [hep-th]].
- [12] M. Raveri, arXiv:1902.01366 [astro-ph.CO].

- [13] E. O. Colgain and H. Yavartanoo, *Phys. Lett. B* **797**, 134907 (2019) [arXiv:1905.02555 [astro-ph.CO]].
- [14] X. Li and A. Shafieloo, *Astrophys. J.* **883**, no. 1, L3 (2019) [*Astrophys. J. Lett.* **883**, L3 (2019)] doi:10.3847/2041-8213/ab3e09 [arXiv:1906.08275 [astro-ph.CO]].
- [15] H. Desmond, B. Jain and J. Sakstein, *Phys. Rev. D* **100**, no. 4, 043537 (2019) [arXiv:1907.03778 [astro-ph.CO]].
- [16] L. Visinelli, S. Vagnozzi and U. Danielsson, *Symmetry* **11**, no. 8, 1035 (2019) doi:10.3390/sym11081035 [arXiv:1907.07953 [astro-ph.CO]].
- [17] E. Di Valentino, A. Melchiorri, O. Mena and S. Vagnozzi, arXiv:1908.04281 [astro-ph.CO].
- [18] S. F. Yan, P. Zhang, J. W. Chen, X. Z. Zhang, Y. F. Cai and E. N. Saridakis, arXiv:1909.06388 [astro-ph.CO].
- [19] O. Akarsu, J. D. Barrow, L. A. Escamilla and J. A. Vazquez, arXiv:1912.08751 [astro-ph.CO].
- [20] Q. Ding, T. Nakama and Y. Wang, arXiv:1912.12600 [astro-ph.CO].
- [21] K. Aylor, M. Joy, L. Knox, M. Millea, S. Raghunathan and W. L. K. Wu, *Astrophys. J.* **874**, no. 1, 4 (2019) [arXiv:1811.00537 [astro-ph.CO]].
- [22] J. Evslin, A. A. Sen and Ruchika, *Phys. Rev. D* **97**, no. 10, 103511 (2018) [arXiv:1711.01051 [astro-ph.CO]].
- [23] V. Poulin, T. L. Smith, T. Karwal and M. Kamionkowski, *Phys. Rev. Lett.* **122**, no. 22, 221301 (2019) [arXiv:1811.04083 [astro-ph.CO]].
- [24] P. Agrawal, F. Y. Cyr-Racine, D. Pinner and L. Randall, arXiv:1904.01016 [astro-ph.CO].
- [25] S. Alexander and E. McDonough, *Phys. Lett. B* **797**, 134830 (2019) [arXiv:1904.08912 [astro-ph.CO]].
- [26] M. X. Lin, G. Benevento, W. Hu and M. Raveri, *Phys. Rev. D* **100**, no. 6, 063542 (2019) [arXiv:1905.12618 [astro-ph.CO]].
- [27] T. L. Smith, V. Poulin and M. A. Amin, arXiv:1908.06995 [astro-ph.CO].
- [28] F. Niedermann and M. S. Sloth, arXiv:1910.10739 [astro-ph.CO].
- [29] J. Sakstein and M. Trodden, arXiv:1911.11760 [astro-ph.CO].
- [30] M. Doran and G. Robbers, *JCAP* **0606**, 026 (2006) [astro-ph/0601544].
- [31] C. D. Kreisch, F. Y. Cyr-Racine and O. Dore, arXiv:1902.00534 [astro-ph.CO].
- [32] M. Park, C. D. Kreisch, J. Dunkley, B. Hadzhiyska and F. Y. Cyr-Racine, *Phys. Rev. D* **100**, no. 6, 063524 (2019) [arXiv:1904.02625 [astro-ph.CO]].
- [33] N. Blinov, K. J. Kelly, G. Z. Krnjaic and S. D. McDermott, *Phys. Rev. Lett.* **123**, no. 19, 191102 (2019) [arXiv:1905.02727 [astro-ph.CO]].
- [34] S. Vagnozzi, arXiv:1907.07569 [astro-ph.CO].
- [35] M. Escudero and S. J. Witte, arXiv:1909.04044 [astro-ph.CO].
- [36] K. L. Pandey, T. Karwal and S. Das, arXiv:1902.10636 [astro-ph.CO].
- [37] K. Vattis, S. M. Koushiappas and A. Loeb, *Phys. Rev. D* **99**, no. 12, 121302 (2019) [arXiv:1903.06220 [astro-ph.CO]].
- [38] P. Agrawal, G. Obied and C. Vafa, arXiv:1906.08261 [astro-ph.CO].
- [39] S. Adhikari and D. Huterer, arXiv:1905.02278 [astro-ph.CO].
- [40] L. Hart and J. Chluba, arXiv:1912.03986 [astro-ph.CO].
- [41] S. Kachru, R. Kallosh, A. D. Linde and S. P. Trivedi, *Phys. Rev. D* **68**, 046005 (2003) [hep-th/0301240].
- [42] R. Kallosh, A. Linde, E. McDonough and M. Scalisi, *Phys. Rev. D* **99**, no. 4, 046006 (2019) [arXiv:1809.09018 [hep-th]].
- [43] L. Susskind, In *Carr, Bernard (ed.): Universe or multiverse?* 247-266 [hep-th/0302219].
- [44] H. Ooguri and C. Vafa, *Nucl. Phys. B* **766**, 21 (2007) [hep-th/0605264].
- [45] G. Obied, H. Ooguri, L. Spodyneiko and C. Vafa, arXiv:1806.08362 [hep-th].
- [46] S. K. Garg and C. Krishnan, *JHEP* **1911**, 075 (2019) doi:10.1007/JHEP11(2019)075 [arXiv:1807.05193 [hep-th]].
- [47] H. Ooguri, E. Palti, G. Shiu and C. Vafa, *Phys. Lett. B* **788**, 180 (2019) [arXiv:1810.05506 [hep-th]].
- [48] Y. S. Piao, *Phys. Rev. D* **70**, 101302 (2004) [hep-th/0407258].
- [49] H. H. Li, G. Ye, Y. Cai and Y. S. Piao, arXiv:1911.06148 [gr-qc].
- [50] D. J. E. Marsh and P. G. Ferreira, *Phys. Rev. D* **82**, 103528 (2010) [arXiv:1009.3501 [hep-ph]].
- [51] B. Audren, J. Lesgourgues, K. Benabed and S. Prunet, *JCAP* **1302**, 001 (2013) [arXiv:1210.7183 [astro-ph.CO]].
- [52] T. Brinckmann and J. Lesgourgues, arXiv:1804.07261 [astro-ph.CO].
- [53] J. Lesgourgues, arXiv:1104.2932 [astro-ph.IM].
- [54] D. Blas, J. Lesgourgues and T. Tram, *JCAP* **1107**, 034 (2011) [arXiv:1104.2933 [astro-ph.CO]].
- [55] N. Aghanim *et al.* [Planck Collaboration], arXiv:1907.12875 [astro-ph.CO].
- [56] S. Alam *et al.* [BOSS Collaboration], *Mon. Not. Roy. Astron. Soc.* **470**, no. 3, 2617 (2017) [arXiv:1607.03155 [astro-ph.CO]].
- [57] F. Beutler *et al.*, *Mon. Not. Roy. Astron. Soc.* **416**, 3017 (2011) [arXiv:1106.3366 [astro-ph.CO]].
- [58] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden and M. Manera, *Mon. Not. Roy. Astron. Soc.* **449**, no. 1, 835 (2015) [arXiv:1409.3242 [astro-ph.CO]].
- [59] D. M. Scolnic *et al.*, *Astrophys. J.* **859**, no. 2, 101 (2018) [arXiv:1710.00845 [astro-ph.CO]].
- [60] Sachs, R.K. and Wolfe, A.M., *Astrophysical Journal*, 1967, 147, 73.
- [61] P. Ade *et al.* [Simons Observatory Collaboration], *JCAP* **1902**, 056 (2019) [arXiv:1808.07445 [astro-ph.CO]].
- [62] K. N. Abazajian *et al.* [CMB-S4 Collaboration], arXiv:1610.02743 [astro-ph.CO].
- [63] E. Di Valentino *et al.* [CORE Collaboration], *JCAP* **1804**, 017 (2018) [arXiv:1612.00021 [astro-ph.CO]].
- [64] A. Suzuki *et al.*, *J. Low. Temp. Phys.* **193**, no. 5-6, 1048 (2018) [arXiv:1801.06987 [astro-ph.IM]].
- [65] L. Hollenstein, D. Sapone, R. Crittenden and B. M. Schaefer, *JCAP* **0904**, 012 (2009) [arXiv:0902.1494 [astro-ph.CO]].
- [66] D. Lst, E. Palti and C. Vafa, *Phys. Lett. B* **797**, 134867 (2019) doi:10.1016/j.physletb.2019.134867 [arXiv:1906.05225 [hep-th]].
- [67] D. Klaeuer and E. Palti, *JHEP* **1701**, 088 (2017) [arXiv:1610.00010 [hep-th]].
- [68] L. A. Anchordoqui, I. Antoniadis, D. Lust, J. F. Soriano and T. R. Taylor, arXiv:1912.00242 [hep-th].
- [69] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, *Phys. Rev. D* **64**, 123522 (2001) [hep-th/0103239]. M. C. Johnson and J. L. Lehnert, *Phys. Rev. D* **85**, 103509 (2012) [arXiv:1112.3360 [hep-th]]. Y. Cai,

Y. Wan, H. G. Li, T. Qiu and Y. S. Piao, *JHEP* **1701**, 090 (2017) [arXiv:1610.03400 [gr-qc]]. P. Creminelli, D. Pirtskhalava, L. Santoni and E. Trincherini, *JCAP* **1611**, 047 (2016) [arXiv:1610.04207 [hep-th]]. R. Kolevatov, S. Mironov, N. Sukhov and V. Volkova, *JCAP* **1708**, 038 (2017) [arXiv:1705.06626 [hep-th]]. Y. Cai and Y. S. Piao,

JHEP **1709**, 027 (2017) [arXiv:1705.03401 [gr-qc]]. G. Ye and Y. S. Piao, *Phys. Rev. D* **99**, no. 8, 084019 (2019) [arXiv:1901.08283 [gr-qc]]. Y. Cai and Y. S. Piao, arXiv:1909.12719 [gr-qc].