M.E. De Rossi<sup>1,2</sup> & V. Bromm<sup>3</sup>

- <sup>1</sup> Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales y Ciclo Básico Común. Buenos Aires, Argentina
- <sup>2</sup> CONICET-Universidad de Buenos Aires, Instituto de Astronomía y Física del Espacio (IAFE). Buenos Aires, Argentina
- <sup>3</sup> Department of Astronomy, University of Texas at Austin, 2511 Speedway, Austin, TX 78712, USA

Contact / mariaemilia.dr@gmail.com

**Resumen** / Exploramos la posibilidad de detectar las primeras galaxias con un telescopio genérico en el infrarrojo lejano y bandas submilimétricas aplicando un modelo analítico para la emisión de polvo primordial. Como se muestra en trabajos previos, las galaxias a corrimientos al rojo z > 7 experimentan una corrección K negativa fuerte de forma tal que sistemas de masas similares son más brillantes a mayores z. Más aún, a una dada masa y z, nuestro modelo predice que los flujos luminosos crecen proporcionalmente a los cocientes polvo-metal (D/M) de las fuentes primigenias. Evaluamos la observabilidad de las fuentes del modelo a diferentes z > 7 como función del área de sondeo observada  $(\Delta \Omega)$  y la sensibilidad (S) de un instrumento genérico. Asumiendo  $\Delta \Omega \sim 10 \text{ deg}^2$  y una plausible  $S \sim 1 \mu$ Jy para un relevamiento en el futuro cercano, podríamos asegurar la detección de al menos una fuente típica con  $D/M \sim 5 \times 10^{-3}$  a z > 7. Para  $S \gtrsim 1 \mu$ Jy y  $\Delta \Omega \lesssim 10 \text{ deg}^2$ , valores de D/M mayores que los típicos son requeridos para detectar al menos una fuente individual a z > 7. La observabilidad de las galaxias del modelo es también afectada por la distribución de tamaño de los granos de polvo, especialmente a altos z.

**Abstract** / We explore the possibility of detecting first galaxies with a generic far-infrared/sub-millimeter telescope by applying an analytical model of primordial dust emission. As shown in previous works, galaxies at redshifts z > 7 experience a strong negative K-correction in such a way that systems of similar masses are brighter at higher z. In addition, at a given mass and z, our model predicts that luminosity fluxes increase proportionally to the dust-to-metal ratios (D/M) of primeval sources. We evaluate the observability of model sources at different z > 7 as a function of the observed survey area  $(\Delta \Omega)$  and sensitivity (S) of a generic instrument. Assuming  $\Delta \Omega \sim 10 \text{ deg}^2$  and a plausible  $S \sim 1 \ \mu$ Jy for a near future survey, we could assure the detection of at least one typical source with  $D/M \sim 5 \times 10^{-3}$  at z > 7. For  $S \gtrsim 1 \ \mu$ Jy and  $\Delta \Omega \lesssim 10 \ \text{deg}^2$ , higher than typical D/M are required to detect at least one individual source at z > 7. The observability of model galaxies is also affected by the size distribution of dust grains, specially towards higher z.

Keywords / galaxies: high-redshift — galaxies: evolution — galaxies: formation — galaxies: star formation — cosmology: theory

# 1. Introduction

With the advent of next-generation observational facilities operating at different wavelengths (e.g, the James Webb Space Telescope (JWST), the Square Kilometre Array (SKA) or the planned Origins Space Telescope, among others), the exploration of the early Universe is entering a golden age. In this context, first stars and galaxies constitute unique tools to test galaxy formation models (e.g. Bromm & Larson, 2004; Bromm et al., 2009; Bromm & Yoshida, 2011; Dayal & Ferrara, 2018). Thus, it is crucial to develop theoretical models that can make predictions regarding the nature and properties of primeval luminous sources in order to guide future surveys.

Given that *typical* dwarf galaxies at z > 7 are expected to have very low dust densities and, thus, low FIR fluxes, De Rossi & Bromm (2017) showed that their detection will be very challenging with current observatories. However, primeval galaxies at  $z \gtrsim 7$  are promising targets for upcoming observational facilities. As discussed in De Rossi & Bromm (2019), future space-borne

FIR telescopes could play a crucial role on the exploration of the nature and properties of the first dustemitting galaxies at the very dawn of star formation. In this manuscript, we extend the work by De Rossi & Bromm (2019) by analysing in more detail the dependence of FIR fluxes on the dust-to-metal ratios (D/M)of first galaxies. We also explore in more detail the observability of model sources at z > 7 as a function of the observed sky area  $(\Delta \Omega)$  and sensitivity (S) of a generic FIR telescope.

### 2. Dust model

Dust emission was estimated using the methodology described in De Rossi & Bromm (2017) and De Rossi & Bromm (2019), which has proven to be useful for studying high-z galaxies (De Rossi et al., 2018). We referred the reader to those papers for a detail description of the dust model; here, we only present a brief summary of it.

A model galaxy consists of a dark matter halo hosting a central cluster of Pop II stars, surrounded by a



Figure 1: Impact of variations of the dust-to-metal ratio (D/M) on the  $F_{\rm FIR} - M_{\rm vir}$  relation for the standard (left panel) and shock (right panel) grain size distributions at z = 7 (thin lines) and z = 20 (thick lines). For comparison, top axes show stellar mass. As a reference, horizontal lines indicate some characteristic sensitivity values discussed in De Rossi & Bromm (2019).

mixed phase of gas and dust. Our standard model assumes a dust-to-metal mass ratio  $D/M = 5 \times 10^{-3}$ , a gas metallicity of  $Z_{\rm g} = 5 \times 10^{-3} {\rm ~Z}_{\odot}$  and a star formation efficiency of  $\eta = 0.01$ , which are typical values expected for first galaxies (Greif & Bromm 2006; Mitchell-Wynne et al. 2015; Schneider et al. 2016; see De Rossi & Bromm 2017, 2019 for more details regarding model parameters and the effects of their variations). The spectral energy distribution associated to stars was obtained from YG-GDRASIL model grids (Zackrisson et al., 2011). We considered different silicon-based dust models given in (Cherchneff & Dwek, 2010). However, for the sake of clarity, we only present results corresponding to the socalled UM-ND-20 model (see De Rossi & Bromm 2017, for details). We have checked that other chemical compositions of dust lead to similar general trends. For the grain-size distribution, we adopted the 'standard' and 'shock' prescriptions used in Ji et al. (2014). Dust temperature  $(T_d)$  was determined assuming thermal equilibrium and dust emissivity was estimated by applying the Kirchhoff's law for the estimated  $T_{\rm d}$  profile.

By comparing plausible sensitivities of a generic instrument with the average observed fluxes ( $F_{\rm FIR}$ ) of model galaxies at a reference FIR wavelength band ( $\Delta \lambda = 250 - 750 \ \mu m$ ), we determined the lowest virial mass ( $M_{\rm vir}$ ) that a galaxy should have to be detected at a given z. By combining our dust model for individual sources with the Sheth-Tormen mass function (Sheth et al., 2001), we estimated the projected number of detected sources that are located at redshift higher than z, within a given solid angle  $\Delta \Omega$  (for a detail description of this methodology, see De Rossi & Bromm 2019).

### 3. Results

Fig. 1 shows the  $F_{\rm FIR} - M_{\rm vir}$  relation as a function of D/M at z = 7 (thin lines) and 20 (thick lines). Results for the standard (left panel) and shock (right panel)

grain size distributions are shown. We note that, for a given dust model, higher FIR fluxes are obtained for galaxies located at higher z. This is a consequence of the strong negative K-correction affecting primeval dust-emitting source at  $z \gtrsim 7$  (see De Rossi & Bromm 2019, for a discussion). In addition, at a given mass, FIR fluxes increase almost proportionally to the increase of D/M, with higher fluxes obtained for the shock size distribution. Considering that next-generation FIR telescopes could reach a lowest sensitivity of  $\sim 1 \,\mu$ Jy, we see that a minimum mass of  $M_{\rm vir} \sim 10^{12} {\rm M}_{\odot}$  would be required to detect systems with typical  $D/M = 5 \times 10^{-3}$ (reference case). For a higher D/M = 0.4 (0.06), the minimum mass limit would be  $M_{\rm vir} \sim 10^{10-11'} \,{\rm M_{\odot}}$  $(\sim 10^{10-11} M_{\odot})$ , with the exact value depending on z and the grain size distribution. Such values of  $M_{\rm vir}$  correspond to very rare massive galaxies at  $z \gtrsim 7$ , which might be difficult to be find during blind surveys (De Rossi & Bromm, 2019).

In Fig. 2, we analyse the observability of primeval galaxies in the  $\Delta\Omega$ -S plane. In the left panel, we show the curves above (below) which the probability for detecting one individual source at z > 7 is P(z > 7) = 1 (< 1), considering different D/M and grain size distributions. We can see that, at a given  $\Delta\Omega$ , the minimum sensitivity required to assure at least one individual detection increases almost proportionally to D/M. For  $\Delta\Omega = 0.1 - 10 \text{ deg}^2$  and assuming  $D/M = 5 \times 10^{-3}$ , a sensitivity  $S \lesssim 0.1 - 1 \ \mu$ Jy would be required to assure one detection at z > 7. For  $\Delta\Omega = 0.1 - 10 \text{ deg}^2$  and assuming a higher  $D/M = 0.06 \ (0.4)$ , the required sensitivity would be  $S \lesssim 1 - 10 \ \mu$ Jy ( $S \lesssim 10 - 100 \ \mu$ Jy). In the case of a sensitivity  $S \approx 1 \ \mu$ Jy (which could be a plausible value for a next-generation FIR telescope) and  $D/M = 5 \times 10^{-3}, 0.06 \ \text{and } 0.4$ , surveys areas  $\Delta\Omega \gtrsim 1 - 10 \ \text{deg}^2$ ,  $\gtrsim 10^{-3} - 10^{-2} \ \text{deg}^2$  and  $\gtrsim 10^{-4} - 10^{-3} \ \text{deg}^2$  would be required to assure the



Figure 2: Constraints on the observability of FIR sources in the  $\Delta\Omega$ -sensitivity plane. Left panel:  $\Delta\Omega$ -sensitivity curves above (below) which the probability of detecting one individual source at z > 7 is P(z > 7) = 1 (< 1). Results are shown for different dust-to-metal ratios (D/M) considering the standard (thin lines) and shock (thick lines) size distributions. Middle and right panels: analysis of observability of sources at redshifts higher than z for  $D/M = 5 \times 10^{-3}$  (reference model), considering the standard and shock size distributions, respectively. Results are shown for z = 7 - 19, with successive curves corresponding to variations of  $\Delta z=1$  (i.e. z = 7, 8, 9..., etc).

detection of at least one individual source at z > 7.

In the middle and right panels of Fig. 2, we analyse the observability of sources located at redshifts higher than a given z, for the standard and shock size distributions, respectively. A reference value  $D/M = 5 \times 10^{-3}$ has been adopted for these panels. For the standard size distribution, lower sensitivities and larger surveys areas are required for detecting sources at all analysed z, with such constraints being stronger towards higher z. Considering a  $\Delta \Omega \leq 10 \text{ deg}^2$  and a standard (shock) size distribution,  $S \leq 1$ , 0.1 and 0.01  $\mu$ Jy are required to detect at least one source at z > 7, 10 and 14 (z > 7, 13 and 18).

## 4. Conclusions

We explored the prospects of detecting the dust continuum emission from first galaxy populations with a generic FIR telescope. We analyse the FIR observed fluxes of these systems as a function of mass and redshift, considering also different dust-to-metal ratios and grain size distributions. According to our model, FIR radiation from first galaxies significantly depends on their dust content and dust properties. Assuming a sensitivity of  $S \approx 1 \ \mu$ Jy for a next-generation telescope, we obtained that only rare massive systems could be detected at z > 7.

Combining our dust model for FIR emission of individual galaxies with the Sheth-Tormen mass function, we evaluated the observability of our galaxy population in the  $\Delta\Omega - S$  plane. Assuming a sensitivity of  $S \approx 1 \ \mu$ Jy and a typical  $D/M = 5 \times 10^{-3}$ , a  $\Delta\Omega \gtrsim 10 \ \text{deg}^2$  would be required to detect at least one individual source at  $z \gtrsim 7$ . However, in the case of higher D/M, smaller survey areas are required.

Our results suggest that future FIR surveys could play a fundamental role on constraining the amount of dust in primeval galaxies at z > 7.

More information and results about this work are available in De Rossi & Bromm (2019).

Acknowledgements: MEDR thanks the Asociación Argentina de Astronomía for providing with partial financial support for attending its 61st annual meeting. MEDR is grateful to PICT-2015-3125 of ANPCyT (Argentina) and also to María Sanz and Guadalupe Lucia for their help and support. VB acknowledges support from NSF grant AST-1413501. We thank Alexander Ji for providing tabulated dust opacities for the different dust models used here. This work makes use of the Yggdrasil code (Zackrisson et al., 2011), which adopts Starburst99 SSP models, based on Padova-AGB tracks (Leitherer et al., 1999; Vázquez & Leitherer, 2005) for Population II stars.

#### References

- Bromm V., Larson R.B., 2004, ARA&A, 42, 79
- Bromm V., Yoshida N., 2011, ARA&A, 49, 373
- Bromm V., et al., 2009, Nature, 459, 49
- Cherchneff I., Dwek E., 2010, ApJ, 713, 1
- Dayal P., Ferrara A., 2018, PhR, 780, 1
- De Rossi M.E., Bromm V., 2017, MNRAS, 465, 3668
- De Rossi M.E., Bromm V., 2019, ApJ, 883, 113
- De Rossi M.E., et al., 2018, ApJ, 869, 4
- Greif T.H., Bromm V., 2006, MNRAS, 373, 128
- Ji A.P., Frebel A., Bromm V., 2014, ApJ, 782, 95

Leitherer C., et al., 1999, ApJS, 123, 3

- Mitchell-Wynne K., et al., 2015, Nature Communications, 6, 7945
- Schneider R., Hunt L., Valiante R., 2016, MNRAS, 457, 1842
- Sheth R.K., Mo H.J., Tormen G., 2001, MNRAS, 323, 1
- Vázquez G.A., Leitherer C., 2005, ApJ, 621, 695
- Zackrisson E., et al., 2011, ApJ, 740, 13