

Superconvergent mixed finite element methods for the pseudostress-velocity formulation of the Oseen equation

Xi Chen · Yuwen Li

Received: November 28, 2020 / Accepted: date

Abstract We presents a priori and superconvergence error estimates of mixed finite element methods for the pseudostress-velocity formulation of the Oseen equation. In particular, we derive supercloseness estimates on the velocity and a priori error estimates under unstructured grids. We also obtain supercloseness estimates on the pseudostress under certain structured grids. With the help of supercloseness and postprocessing procedures, postprocessed numerical velocity and pseudostress are shown to be superconvergent. We finally present numerical experiments to verify the theoretical results and illustrate the effectiveness of the superconvergent recovery-based a posteriori error estimator.

Keywords pseudostress, superconvergence, supercloseness, Oseen equation, Stokes equation, postprocessing

Mathematics Subject Classification (2000) 65N30, 65N15

1 Introduction

In fluid mechanics, the Oseen equations describe the flow of a viscous and incompressible fluid at small Reynolds numbers. Instead of dropping the advective term completely, the Oseen approximation linearizes the advective acceleration term $\mathbf{u} \cdot \nabla \mathbf{u}$ to $\mathbf{b} \cdot \nabla \mathbf{u}$, where \mathbf{b} represents the velocity at large distance. As a result it provides a lowest-order solution that is uniformly valid everywhere in the flow field. Since the Oseen equations partly account for the inertia terms (at large distance), they have better approximation in the far field while keeping the same order of accuracy as Stokes approximation near the body, see, e.g., [41]. Mathematically speaking, the Oseen equation can be viewed as

X. Chen: Department of Mechanical Engineering, Pennsylvania State University, University Park, PA 16802. Email: xbc5027@psu.edu

✉ Y. Li: Department of Mathematics, Pennsylvania State University, University Park, PA 16802. Email: yuwenli925@gmail.com

a linearized Navier–Stokes equation arising from fixed point iteration. It is the purpose of this paper to shed light on a priori and superconvergence analysis of pseudostress-velocity mixed methods for the Navier–Stokes equation.

Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain. Let $\mathbf{b}, \mathbf{f} : \Omega \rightarrow \mathbb{R}^d$, $c : \Omega \rightarrow \mathbb{R}$ and $\mathbf{g} : \partial\Omega \rightarrow \mathbb{R}^d$ be given data. Let $\mathbf{u} : \Omega \rightarrow \mathbb{R}^d$ be the velocity field and $p : \Omega \rightarrow \mathbb{R}$ denote the pressure subject to the constraint

$$\int_{\Omega} p dx = 0. \quad (1.1)$$

The Oseen equation under the Dirichlet boundary condition is

$$-\Delta \mathbf{u} + \mathbf{b} \cdot \nabla \mathbf{u} + c\mathbf{u} + \nabla p = \mathbf{f} \text{ in } \Omega, \quad (1.2a)$$

$$\operatorname{div} \mathbf{u} = 0 \text{ in } \Omega, \quad (1.2b)$$

$$\mathbf{u} = \mathbf{g} \text{ on } \partial\Omega. \quad (1.2c)$$

As a default, vectors such as $\mathbf{b}, \mathbf{f}, \mathbf{u}, \mathbf{g}$ are always arranged as columns unless confusion arises. In (1.2a), $\nabla \mathbf{u}$ is the Jacobian matrix of \mathbf{u} , and we adopt the convention $\mathbf{b} \cdot \nabla \mathbf{u} := (\nabla \mathbf{u})\mathbf{b}$. When $\mathbf{b} = \mathbf{0}$ (resp. $\mathbf{b} = \mathbf{0}, c = 0$), (1.2) reduces to the Brinkman equation (resp. Stokes equation). Numerical analysis of Oseen, Brinkman, or Stokes equations based on the velocity-pressure formulation is extensive, see, e.g., [37, 9, 38, 34] for conforming mixed methods, [21, 24, 26, 35] for hybridized discontinuous Galerkin methods, [48, 55, 45] for weak Galerkin methods, and [1] for virtual element methods.

Let $\mathbf{I} \in \mathbb{R}^{d \times d}$ denote the identity matrix. Define the pseudostress

$$\boldsymbol{\sigma} := \nabla \mathbf{u} - p\mathbf{I}$$

which is not symmetric; compared with the symmetric stress

$$\boldsymbol{\sigma}^S := \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T) - p\mathbf{I}.$$

In a series of papers [20, 19, 18], Cai et al. proposed the pseudostress-velocity formulation of the Stokes and Navier–Stokes equations, which allows classical Poisson-type mixed discretization by using Raviart–Thomas or Brezzi–Douglas–Marini elements. Let $\operatorname{tr} \boldsymbol{\sigma}$ denote the trace of the matrix $\boldsymbol{\sigma}$ and

$$\mathcal{A}\boldsymbol{\sigma} := \boldsymbol{\sigma} - \frac{1}{d}(\operatorname{tr} \boldsymbol{\sigma})\mathbf{I}$$

denote the deviatoric part of $\boldsymbol{\sigma}$. Direct calculation shows that the Oseen equation (1.2) is equivalent to the following pseudostress-velocity formulation

$$\begin{aligned} \mathcal{A}\boldsymbol{\sigma} &= \nabla \mathbf{u}, \\ -\operatorname{div} \boldsymbol{\sigma} + \mathbf{b} \cdot \mathcal{A}\boldsymbol{\sigma} + c\mathbf{u} &= \mathbf{f} \end{aligned} \quad (1.3)$$

subject to the constraints

$$\mathbf{u}|_{\partial\Omega} = \mathbf{g}, \quad \int_{\Omega} \operatorname{tr} \boldsymbol{\sigma} dx = 0.$$

Here $\operatorname{div} \boldsymbol{\sigma}$ is the row-wise divergence of the matrix $\boldsymbol{\sigma}$. Comparing to extensive numerical results on velocity-pressure formulation of Stokes-related equations, the numerical analysis of pseudostress-based methods is restricted to classical mixed methods [19, 36], virtual element methods [16, 17], discontinuous Galerkin methods [51] and adaptive mixed methods [23, 22, 40, 42] for the Stokes or Brinkman equation. Very few work is devoted to the mixed formulation of the Oseen equation (1.3), see, e.g., [50] for a lowest order upwinded mixed method on rectangular meshes.

Given a vector space X , let $[X]^n$ denote the Cartesian product of n copies of X and $[X]^{n \times n}$ the space of $n \times n$ matrices whose components are contained in X . Let $H(\operatorname{div}, \Omega) = \{\boldsymbol{v} \in [L^2(\Omega)]^d : \operatorname{div} \boldsymbol{v} \in L^2(\Omega)\}$ be the usual $H(\operatorname{div})$ space. Let

$$\begin{aligned} \mathbf{V} &= [L^2(\Omega)]^d, \\ \boldsymbol{\Sigma} &= \{\boldsymbol{\tau} \in [L^2(\Omega)]^{d \times d} : \boldsymbol{\tau}_i \in H(\operatorname{div}, \Omega), 1 \leq i \leq d, \int_{\Omega} \operatorname{tr} \boldsymbol{\tau} dx = 0\}, \end{aligned}$$

where $\boldsymbol{\tau}_i$ is the i -th row of $\boldsymbol{\tau}$. For scalar-, vector-, or matrix-valued functions, we use (\cdot, \cdot) and $\langle \cdot, \cdot \rangle$ to denote the $L^2(\Omega)$ - and $L^2(\partial\Omega)$ -inner product, respectively. The variational formulation of (1.3) seeks $\boldsymbol{\sigma} \in \boldsymbol{\Sigma}$ and $\boldsymbol{u} \in \mathbf{V}$ such that

$$\begin{aligned} (\mathcal{A}\boldsymbol{\sigma}, \boldsymbol{\tau}) + (\operatorname{div} \boldsymbol{\tau}, \boldsymbol{u}) &= \langle \boldsymbol{g}, \boldsymbol{\tau} \boldsymbol{n} \rangle, \quad \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \\ -(\operatorname{div} \boldsymbol{\sigma}, \boldsymbol{v}) + (\boldsymbol{b} \cdot \mathcal{A}\boldsymbol{\sigma}, \boldsymbol{v}) + (\boldsymbol{c}\boldsymbol{u}, \boldsymbol{v}) &= (\boldsymbol{f}, \boldsymbol{v}), \quad \boldsymbol{v} \in \mathbf{V}, \end{aligned} \quad (1.4)$$

where \boldsymbol{n} is the outward normal to $\partial\Omega$.

Let \mathcal{T}_h be a conforming simplicial or cubical partition of Ω that is shape-regular in the sense that

$$\max_{K \in \mathcal{T}_h} \frac{r_K}{\rho_K} := C_{\text{ratio}} < \infty$$

for some absolute constant C_{ratio} , where r_K, ρ_K are radii of circumscribed and inscribed spheres of K . For each $K \in \mathcal{T}_h$, let h_K denote the diameter of K and $h := \max_{K \in \mathcal{T}_h} h_K < 1$ the mesh size of \mathcal{T}_h . Given an integer $r \geq 0$, let $\tilde{\boldsymbol{\Sigma}}_r(K), V_r(K)$ be suitable finite-dimensional vector spaces defined on K and

$$\begin{aligned} \tilde{\boldsymbol{\Sigma}}_h &:= \{\boldsymbol{\tau} \in H(\operatorname{div}, \Omega) : \boldsymbol{\tau}|_K \in \tilde{\boldsymbol{\Sigma}}_r(K) \text{ for } K \in \mathcal{T}_h\}, \\ V_h &:= \{v \in L^2(\Omega) : v|_K \in V_r(K) \text{ for } K \in \mathcal{T}_h\}. \end{aligned}$$

For instance, when \mathcal{T}_h is a simplicial partition, set

$$\tilde{\boldsymbol{\Sigma}}_r(K) := [\mathcal{P}_r(K)]^d + \mathcal{P}_r(K)\boldsymbol{x}, \quad V_r(K) := \mathcal{P}_r(K), \quad (1.5)$$

where $\mathcal{P}_r(K)$ is the space of polynomials of degree at most r on K , and $\boldsymbol{x} = (x_1, \dots, x_d)^T$ denotes the position vector. In this case, the corresponding $\tilde{\boldsymbol{\Sigma}}_h \times V_h$ is the classical Raviart–Thomas (RT) [52] space. Another possible choice is

$$\tilde{\boldsymbol{\Sigma}}_r(K) := [\mathcal{P}_{r+1}(K)]^d, \quad V_r(K) := \mathcal{P}_r(K). \quad (1.6)$$

In this case, the corresponding $\tilde{\Sigma}_h \times V_h$ becomes the Brezzi–Douglas–Marini (BDM) [15] space.

Given a hypercube $K \subset \mathbb{R}^d$, let $\mathcal{Q}_r(K)$ denote the space of polynomials on K of degree at most r in x_i with $1 \leq i \leq d$. For a cubical mesh \mathcal{T}_h , the pair $\tilde{\Sigma}_h \times V_h$ can be the cubical RT element using the shape functions

$$\tilde{\Sigma}_r(K) := [\mathcal{Q}_r(K)]^d + \mathbf{x}\mathcal{Q}_r(K), \quad V_r(K) := \mathcal{Q}_r(K). \quad (1.7)$$

Let $\tilde{\Pi}_h : [H^1(\Omega)]^d \rightarrow \tilde{\Sigma}_h$ be the canonical interpolation onto $\tilde{\Sigma}_h$ determined by the degrees of freedom of $\tilde{\Sigma}_h$. Let $P_h : L^2(\Omega) \rightarrow V_h$ be the L^2 -projection onto V_h , i.e.,

$$(P_h v, w) = (v, w), \quad v \in L^2(\Omega), \quad w \in V_h. \quad (1.8)$$

Throughout the rest, we assume the commutativity property

$$\operatorname{div} \tilde{\Pi}_h = P_h \operatorname{div}, \quad (1.9)$$

which is satisfied by at least the aforementioned RT, BDM, and cubical RT elements. Readers are referred to e.g., [15, 13, 14] for other cubical or rectangular elements that satisfy (1.9). Let Σ_h be the matrix version of $\tilde{\Sigma}_h$, i.e.,

$$\Sigma_h := \{\boldsymbol{\tau} \in \Sigma : \tau_i \in \tilde{\Sigma}_h \text{ for } 1 \leq i \leq d\}.$$

Let \mathbf{V}_h be the vector-valued broken piecewise polynomial space

$$\mathbf{V}_h := \{\mathbf{v} = (v_1, v_2, \dots, v_d)^T \in \mathbf{V} : v_i \in V_h \text{ for } 1 \leq i \leq d\}.$$

The mixed method for (1.3) seeks $(\boldsymbol{\sigma}_h, \mathbf{u}_h) \in \Sigma_h \times \mathbf{V}_h$ satisfying

$$\begin{aligned} (\mathcal{A}\boldsymbol{\sigma}_h, \boldsymbol{\tau}) + (\operatorname{div} \boldsymbol{\tau}, \mathbf{u}_h) &= \langle \mathbf{g}, \boldsymbol{\tau} \mathbf{n} \rangle, \quad \boldsymbol{\tau} \in \Sigma_h, \\ -(\operatorname{div} \boldsymbol{\sigma}_h, \mathbf{v}) + (\mathbf{b} \cdot \mathcal{A}\boldsymbol{\sigma}_h, \mathbf{v}) + (c\mathbf{u}_h, \mathbf{v}) &= (\mathbf{f}, \mathbf{v}), \quad \mathbf{v} \in \mathbf{V}_h. \end{aligned} \quad (1.10)$$

In this paper, we shall develop a priori and superconvergence error estimates for (1.10). To the best of our knowledge, even a priori error estimates of (1.10) is not available in literature. We emphasize that the presence of lower order terms is a major challenge in error analysis of mixed methods, see, e.g., [30, 28] for elliptic equations with lower order terms and [5] for the perturbed Hodge–Laplace equation. For instance, due to the convection term $\mathbf{b} \cdot \mathcal{A}\boldsymbol{\sigma}$ and possibly negative coefficient c , the construction of the discrete inf-sup condition is far from obvious. Hence we shall use the duality argument to derive a priori error estimates, which also yields improved decoupled error estimates. The main contribution of this paper is summarized as follows.

- We develop a supercloseness estimate on $\|P_h \mathbf{u} - \mathbf{u}_h\|$, which reduces to a result in [23] for the Stokes equation ($\mathbf{b} = \mathbf{0}$, $c = 0$).

- For the mixed method (1.10), we establish optimal a priori error estimates in the energy-norm and L^2 -norm using duality argument. It is noted that mixed methods for the scalar elliptic equation

$$-\operatorname{div}(\mathbf{A}\nabla u + \mathbf{b}u) + cu = f \quad (1.11)$$

have been analyzed in [30, 28], where \mathbf{A} is a uniformly elliptic matrix-valued function. In contrast, the deviatoric operator \mathcal{A} in (1.2) is singular, which is a major difficulty in error analysis. We overcome this difficulty by deriving sharp estimates on $\|\operatorname{div}(\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h)\|_{-1}$ and $\|P_h \mathbf{u} - \mathbf{u}_h\|$ for closing a loop in a priori error estimation, see Section 2 for details.

- We establish supercloseness estimates on $\|\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|$ for the lowest order triangular and general rectangular RT elements. Even for the Stokes equation, such estimates are not known in literature. In [43], Li obtained similar estimates of mixed methods for the elliptic equation (1.11). The novelty of supercloseness analysis here is two-fold. First, (1.4) with a uniformly elliptic \mathcal{A} is still different from the mixed formulation of (1.11). Second, the singular deviatoric operator \mathcal{A} is handled in the framework of finite element exterior calculus, see Section 3 for details.

Built upon the aforementioned supercloseness estimates and available post-processing operators in literature (see, e.g., [59, 23, 8, 57, 7]), we obtain the recovery superconvergence in Theorems 4.1 and 4.2. Regarding superconvergence results of velocity-pressure methods for the Stokes-type equation, readers are referred to e.g., [54, 56, 27] for superconvergence by two-grid L^2 -projections, [49, 32, 44] for superconvergent recovery of lowest-order methods, and [26, 35] for superconvergent hybridized discontinuous Galerkin methods.

The rest of this paper is organized as follows. In section 2, we develop a priori error estimates of the mixed method (1.10) and supercloseness estimate on $\|P_h \mathbf{u} - \mathbf{u}_h\|$. In Section 3, we develop supercloseness estimate on $\|\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|$. Section 4 is devoted to superconvergent postprocessing procedures. Numerical results for smooth, singular, and convection-dominated problems are presented in Section 5.

2 A priori error estimates

Let $\|\cdot\|_r$ denote the $\|\cdot\|_{H^r(\Omega)}$ -norm, $|\cdot|_r$ the $|\cdot|_{H^r(\Omega)}$ -semi norm, and $\|\cdot\| := \|\cdot\|_0$ the $\|\cdot\|_{L^2(\Omega)}$ -norm. Let $\|\|\cdot\|\|$ denote the $H(\operatorname{div})$ -norm,

$$\|\|\boldsymbol{\tau}\|\|^2 := \|\boldsymbol{\tau}\|^2 + \|\operatorname{div} \boldsymbol{\tau}\|^2.$$

In this section, we shall derive a priori error estimates in Theorem 2.1. The same analysis implies that the mixed method (1.10) admits a unique solution provided h is sufficiently small. Throughout the rest, $A \lesssim B$ means that $A \leq CB$, where $C > 0$ is a generic constant dependent solely on \mathbf{b} , c , Ω , C_{ratio} .

The operator \mathcal{A} is singular and satisfies

$$(\mathcal{A}\boldsymbol{\tau}_1, \boldsymbol{\tau}_2) = (\boldsymbol{\tau}_1, \mathcal{A}\boldsymbol{\tau}_2) = (\mathcal{A}\boldsymbol{\tau}_1, \mathcal{A}\boldsymbol{\tau}_2), \quad \boldsymbol{\tau}_1, \boldsymbol{\tau}_2 \in [L^2(\Omega)]^{d \times d}. \quad (2.1)$$

In addition, it holds that (see [20,2])

$$\|\boldsymbol{\tau}\| \lesssim \|\boldsymbol{\tau}\|_{\mathcal{A}} + \|\operatorname{div} \boldsymbol{\tau}\|_{-1}, \quad \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \quad (2.2)$$

where $\|\boldsymbol{\tau}\|_{\mathcal{A}} := (\mathcal{A}\boldsymbol{\tau}, \boldsymbol{\tau})^{\frac{1}{2}}$. (2.2) is a key ingredient in the analysis of pseudostress-based methods. Define $\Pi_h : [H^1(\Omega)]^{d \times d} \rightarrow \boldsymbol{\Sigma}_h$ by

$$\Pi_h \boldsymbol{\tau} := \tilde{\Pi}_h \boldsymbol{\tau} - \frac{1}{d|\Omega|} \left(\int_{\Omega} \operatorname{tr} \tilde{\Pi}_h \boldsymbol{\tau} dx \right) \mathbf{I}, \quad \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \quad (2.3)$$

where $\tilde{\Pi}_h$ is applied to each row of $\boldsymbol{\tau}$. By abuse of notation, we also use $P_h : [L^2(\Omega)]^d \rightarrow \mathbf{V}_h$ to denote the component-wise L^2 -projection. It follows from (1.9) that

$$\operatorname{div} \Pi_h \boldsymbol{\tau} = P_h \operatorname{div} \boldsymbol{\tau}, \quad \boldsymbol{\tau} \in [H^1(\Omega)]^{d \times d}. \quad (2.4)$$

For convenience, we introduce the interpolation errors

$$\boldsymbol{\rho}_h := \mathbf{u} - P_h \mathbf{u}, \quad \boldsymbol{\zeta}_h := \boldsymbol{\sigma} - \Pi_h \boldsymbol{\sigma},$$

which can be easily estimated by (see [18])

$$\|\mathbf{v} - P_h \mathbf{v}\| \lesssim h^s |\mathbf{v}|_s, \quad (2.5a)$$

$$\|\boldsymbol{\tau} - \Pi_h \boldsymbol{\tau}\| \lesssim h^s |\boldsymbol{\tau}|_s, \quad (2.5b)$$

$$\|\operatorname{div}(\boldsymbol{\tau} - \Pi_h \boldsymbol{\tau})\| \lesssim h^s |\operatorname{div} \boldsymbol{\tau}|_s, \quad (2.5c)$$

where $0 \leq s \leq r + 1$, $\int_{\Omega} \operatorname{tr} \boldsymbol{\tau} dx = 0$, and $\mathbf{v}, \boldsymbol{\tau}$ satisfy the regularity indicated by the right hand sides. For the BDM element (1.6), it holds that

$$\|\boldsymbol{\tau} - \Pi_h \boldsymbol{\tau}\| \lesssim h^s |\boldsymbol{\tau}|_s, \quad 0 \leq s \leq r + 2. \quad (2.6)$$

The essential errors to be estimated are

$$\mathbf{e}_h := P_h \mathbf{u} - \mathbf{u}_h, \quad \boldsymbol{\xi}_h := \Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h.$$

Subtracting (1.10) from (1.4), we obtain the error equation

$$(\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \boldsymbol{\tau}_h) + (\operatorname{div} \boldsymbol{\tau}_h, \mathbf{u} - \mathbf{u}_h) = 0, \quad (2.7a)$$

$$-(\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}_h) + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}_h) + (c(\mathbf{u} - \mathbf{u}_h), \mathbf{v}_h) = 0, \quad (2.7b)$$

for all $(\boldsymbol{\tau}_h, \mathbf{v}_h) \in \boldsymbol{\Sigma}_h \times \mathbf{V}_h$.

To estimate $\|\mathbf{e}_h\|$, we consider the dual problem of (1.4): Find $\tilde{\boldsymbol{\sigma}} \in \boldsymbol{\Sigma}$ and $\tilde{\mathbf{u}} \in \mathbf{V}$ such that

$$(\mathcal{A}\tilde{\boldsymbol{\sigma}}, \boldsymbol{\tau}) + (\operatorname{div} \boldsymbol{\tau}, \tilde{\mathbf{u}}) - (\mathbf{b} \cdot \mathcal{A}\boldsymbol{\tau}, \tilde{\mathbf{u}}) = 0, \quad \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \quad (2.8a)$$

$$-(\operatorname{div} \tilde{\boldsymbol{\sigma}}, \mathbf{v}) + (c\tilde{\mathbf{u}}, \mathbf{v}) = (\mathbf{e}_h, \mathbf{v}) \quad \mathbf{v} \in \mathbf{V}. \quad (2.8b)$$

Throughout the rest, it is assumed that (2.8) admits the elliptic regularity

$$\|\tilde{\boldsymbol{\sigma}}\|_1 + \|\tilde{\mathbf{u}}\|_1 \lesssim \|\mathbf{e}_h\|. \quad (2.9)$$

The next lemma is a supercloseness estimate which is crucial for both a priori and superconvergence error analysis.

Lemma 2.1 *It holds that*

$$\|e_h\| \lesssim h(\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + \|\mathbf{u} - \mathbf{u}_h\| + \|\boldsymbol{\rho}_h\|).$$

Proof Taking $\mathbf{v} = e_h$ in (2.8b), we obtain

$$\|e_h\|^2 = -(\operatorname{div} \tilde{\boldsymbol{\sigma}}, e_h) + (c\tilde{\mathbf{u}}, e_h). \quad (2.10)$$

Using the definition (1.8) and the property (2.4), we have

$$\begin{aligned} (\operatorname{div} \tilde{\boldsymbol{\sigma}}, e_h) &= (P_h \operatorname{div} \tilde{\boldsymbol{\sigma}}, P_h \mathbf{u} - \mathbf{u} + \mathbf{u} - \mathbf{u}_h) \\ &= (P_h \operatorname{div} \tilde{\boldsymbol{\sigma}}, \mathbf{u} - \mathbf{u}_h) = (\operatorname{div} \Pi_h \tilde{\boldsymbol{\sigma}}, \mathbf{u} - \mathbf{u}_h). \end{aligned} \quad (2.11)$$

(2.7a) with $\boldsymbol{\tau}_h = \Pi_h \tilde{\boldsymbol{\sigma}}$ implies that

$$\begin{aligned} (\operatorname{div} \Pi_h \tilde{\boldsymbol{\sigma}}, \mathbf{u} - \mathbf{u}_h) &= -(\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \Pi_h \tilde{\boldsymbol{\sigma}}) \\ &= (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\boldsymbol{\sigma}} - \Pi_h \tilde{\boldsymbol{\sigma}}) - (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\boldsymbol{\sigma}}). \end{aligned} \quad (2.12)$$

Using (2.1) and setting $\tilde{\boldsymbol{\tau}} = \boldsymbol{\sigma} - \boldsymbol{\sigma}_h$ in (2.8a), we have

$$\begin{aligned} (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\boldsymbol{\sigma}}) &= (\mathcal{A}\tilde{\boldsymbol{\sigma}}, \boldsymbol{\sigma} - \boldsymbol{\sigma}_h) \\ &= -(\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}}) + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}}) \\ &= -(\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) - (\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), P_h \tilde{\mathbf{u}}) + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}}). \end{aligned}$$

It then follows from the previous equation and (2.7b) with $\mathbf{v}_h = P_h \tilde{\mathbf{u}}$ that

$$\begin{aligned} (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\boldsymbol{\sigma}}) &= -(\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}}) \\ &\quad - (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), P_h \tilde{\mathbf{u}}) - (c(\mathbf{u} - \mathbf{u}_h), P_h \tilde{\mathbf{u}}). \end{aligned} \quad (2.13)$$

As a result of (2.11), (2.12) and (2.13), we obtain

$$\begin{aligned} (\operatorname{div} \tilde{\boldsymbol{\sigma}}, e_h) &= (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\boldsymbol{\sigma}} - \Pi_h \tilde{\boldsymbol{\sigma}}) + (\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) \\ &\quad - (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) + (c(\mathbf{u} - \mathbf{u}_h), P_h \tilde{\mathbf{u}}). \end{aligned} \quad (2.14)$$

On the other hand, the second term on the right hand side of (2.10) is

$$\begin{aligned} (c\tilde{\mathbf{u}}, e_h) &= (c\tilde{\mathbf{u}}, P_h \mathbf{u} - \mathbf{u} + \mathbf{u} - \mathbf{u}_h) \\ &= (c\tilde{\mathbf{u}} - P_h(c\tilde{\mathbf{u}}), P_h \mathbf{u} - \mathbf{u}) + (c(\mathbf{u} - \mathbf{u}_h), \tilde{\mathbf{u}}). \end{aligned} \quad (2.15)$$

Finally with (2.14) and (2.15), the error in (2.10) can be written as

$$\begin{aligned} \|e_h\|^2 &= (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \Pi_h \tilde{\boldsymbol{\sigma}} - \tilde{\boldsymbol{\sigma}}) - (\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) \\ &\quad + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) + (c(\mathbf{u} - \mathbf{u}_h), \tilde{\mathbf{u}} - P_h \tilde{\mathbf{u}}) \\ &\quad + (c\tilde{\mathbf{u}} - P_h(c\tilde{\mathbf{u}}), P_h \mathbf{u} - \mathbf{u}). \end{aligned} \quad (2.16)$$

Using (2.16), (2.5), and the Cauchy–Schwarz inequality, we obtain

$$\|e_h\|^2 \lesssim h\|\tilde{\boldsymbol{\sigma}}\|_1\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + h\|\tilde{\mathbf{u}}\|_1(\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + \|\mathbf{u} - \mathbf{u}_h\| + \|\boldsymbol{\rho}_h\|).$$

Combining the above estimate with (2.9) completes the proof. \square

Lemma 2.1 is a supercloseness estimate, i.e., $P_h \mathbf{u}$ and \mathbf{u}_h are much closer than the distance predicted by standard a priori error estimates. However, a priori error estimates on $\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|$ and $\|\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h)\|$ are not known at the moment. When deriving a priori error estimates, we need the L^2 and negative norm estimates of $\operatorname{div} \boldsymbol{\xi}_h$.

Lemma 2.2 *It holds that*

$$\|\operatorname{div} \boldsymbol{\xi}_h\| \lesssim \|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\xi}_h\| + \|\mathbf{e}_h\| + h\|\boldsymbol{\rho}_h\|, \quad (2.17a)$$

$$\|\operatorname{div} \boldsymbol{\xi}_h\|_{-1} \lesssim h(\|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\xi}_h\| + \|\boldsymbol{\rho}_h\|) + \|\mathbf{e}_h\|. \quad (2.17b)$$

Lemma 2.2 is also essential for proving the supercloseness estimate in Theorem 3.1 and the proof is postponed in the end of this section. Throughout the rest, we say h is sufficiently small provided $h \leq h_0$, where h_0 is a constant dependent solely on Ω , \mathbf{b} , c , C_{ratio} . Now we are in a position to present the main result in this paper.

Theorem 2.1 *For sufficiently small h , it holds that*

$$\|\mathbf{u} - \mathbf{u}_h\| \lesssim \|\boldsymbol{\rho}_h\| + h\|\boldsymbol{\zeta}_h\|, \quad (2.18a)$$

$$\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| \lesssim \|\boldsymbol{\zeta}_h\| + h\|\boldsymbol{\rho}_h\|, \quad (2.18b)$$

$$\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| \lesssim \|\boldsymbol{\zeta}_h\| + h\|\operatorname{div} \boldsymbol{\zeta}_h\| + h\|\boldsymbol{\rho}_h\|. \quad (2.18c)$$

Proof Using Lemma 2.1 and the triangle inequality

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\| &\leq \|\boldsymbol{\rho}_h\| + \|\mathbf{e}_h\|, \\ \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| &\leq \|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\xi}_h\|, \end{aligned}$$

we obtain

$$\|\mathbf{e}_h\| \lesssim h(\|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\xi}_h\| + \|\boldsymbol{\rho}_h\|) + h\|\mathbf{e}_h\|. \quad (2.19)$$

Plugging (2.17a) into (2.19) and kicking $h\|\mathbf{e}_h\|$ with sufficiently small h into the left hand side yields

$$\|\mathbf{e}_h\| \lesssim h(\|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\xi}_h\| + \|\boldsymbol{\rho}_h\|). \quad (2.20)$$

On the other hand, with help of (2.7a) with $\boldsymbol{\tau}_h = \boldsymbol{\xi}_h$, we obtain

$$\begin{aligned} (\mathcal{A}\boldsymbol{\xi}_h, \boldsymbol{\xi}_h) &= (\mathcal{A}(\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}), \boldsymbol{\xi}_h) - (\operatorname{div} \boldsymbol{\xi}_h, \mathbf{u} - \mathbf{u}_h) \\ &= (\mathcal{A}(\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}), \boldsymbol{\xi}_h) - (\operatorname{div} \boldsymbol{\xi}_h, P_h \mathbf{u} - \mathbf{u}_h). \end{aligned} \quad (2.21)$$

A combination of (2.2), (2.21), with a Young's inequality then shows that

$$\begin{aligned} \|\boldsymbol{\xi}_h\|^2 &\lesssim (\mathcal{A}\boldsymbol{\xi}_h, \boldsymbol{\xi}_h) + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1}^2 \\ &\lesssim \|\boldsymbol{\zeta}_h\| \|\boldsymbol{\xi}_h\| + \|\operatorname{div} \boldsymbol{\xi}_h\| \|\mathbf{e}_h\| + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1}^2 \\ &\leq \frac{\varepsilon^2}{2} \|\boldsymbol{\xi}_h\|^2 + \frac{\varepsilon^{-2}}{2} \|\boldsymbol{\zeta}_h\|^2 + \frac{\varepsilon^2}{2} \|\operatorname{div} \boldsymbol{\xi}_h\|^2 + \frac{\varepsilon^{-2}}{2} \|\mathbf{e}_h\|^2 + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1}^2, \end{aligned}$$

or equivalently,

$$\|\boldsymbol{\xi}_h\| \leq C^* (\varepsilon \|\boldsymbol{\xi}_h\| + \varepsilon^{-1} \|\boldsymbol{\zeta}_h\| + \varepsilon \|\operatorname{div} \boldsymbol{\xi}_h\| + \varepsilon^{-1} \|\mathbf{e}_h\| + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1}), \quad (2.22)$$

where $\varepsilon > 0$, C^* is independent of ε . Collecting (2.22), Lemma 2.2 and using the triangle inequality, we deduce that

$$\begin{aligned} \|\boldsymbol{\xi}_h\| &\leq C^* \{ \varepsilon (\|\boldsymbol{\xi}_h\| + \|\boldsymbol{\zeta}_h\| + \|\mathbf{e}_h\| + h\|\boldsymbol{\rho}_h\|) \\ &\quad + \varepsilon^{-1} (\|\boldsymbol{\zeta}_h\| + \|\mathbf{e}_h\|) + h\|\boldsymbol{\zeta}_h\| + h\|\boldsymbol{\xi}_h\| + h\|\boldsymbol{\rho}_h\| + \|\mathbf{e}_h\| \}. \end{aligned}$$

Then we could choose ε and h to be small enough to obtain

$$\|\boldsymbol{\xi}_h\| \leq C (\|\boldsymbol{\zeta}_h\| + h\|\operatorname{div} \boldsymbol{\zeta}_h\| + \|\mathbf{e}_h\| + h\|\boldsymbol{\rho}_h\|). \quad (2.23)$$

Plugging (2.23) into (2.20) with sufficiently small h leads to

$$\|\mathbf{e}_h\| \leq Ch (\|\boldsymbol{\zeta}_h\| + \|\boldsymbol{\rho}_h\|). \quad (2.24)$$

Therefore we close the loop. As a result of (2.23), (2.24), (2.17a), we obtain

$$\begin{aligned} \|\boldsymbol{\xi}_h\| &\leq C (\|\boldsymbol{\zeta}_h\| + h\|\operatorname{div} \boldsymbol{\zeta}_h\| + h\|\boldsymbol{\rho}_h\|), \\ \|\operatorname{div} \boldsymbol{\xi}_h\| &\leq C (\|\boldsymbol{\zeta}_h\| + h\|\operatorname{div} \boldsymbol{\zeta}_h\| + h\|\boldsymbol{\rho}_h\|). \end{aligned} \quad (2.25)$$

Theorem 2.1 then follows from (2.24), (2.25), and the triangle inequality. \square

For sufficiently smooth $(\boldsymbol{\sigma}, \mathbf{u})$, Theorem 2.1 with (2.5) yields

$$\|\mathbf{u} - \mathbf{u}_h\| \lesssim h^{r+1} (|\mathbf{u}|_{r+1} + h|\boldsymbol{\sigma}|_{r+1} + h|\operatorname{div} \boldsymbol{\sigma}|_{r+1}), \quad (2.26a)$$

$$\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| \lesssim h^{r+1} (|\boldsymbol{\sigma}|_{r+1} + |\operatorname{div} \boldsymbol{\sigma}|_{r+1} + h|\mathbf{u}|_{r+1}). \quad (2.26b)$$

When $\boldsymbol{\Sigma}_h$ is based on the BDM element (1.6), the following improved error estimate follows from (2.18c) and (2.6).

$$\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| \lesssim h^{r+2} (|\boldsymbol{\sigma}|_{r+2} + |\mathbf{u}|_{r+1}). \quad (2.27)$$

The well-posedness (1.10) follows from the same analysis in the proof of Theorem 2.1.

Corollary 2.1 *For sufficiently small h , the mixed method (1.10) has a unique solution.*

Proof To establish the existence and uniqueness of the solution to (1.10), it suffices to show the uniqueness because of linearity. Suppose $(\boldsymbol{\sigma}_h, \mathbf{u}_h)$ and $(\hat{\boldsymbol{\sigma}}_h, \hat{\mathbf{u}}_h)$ are both solutions to (1.10), then we have the error equation

$$\begin{aligned} (\mathcal{A}e_\sigma, \boldsymbol{\tau}_h) + (\operatorname{div} \boldsymbol{\tau}_h, \mathbf{e}_u) &= 0, \quad \boldsymbol{\tau}_h \in \boldsymbol{\Sigma}_h, \\ -(\operatorname{div} \mathbf{e}_\sigma, \mathbf{v}_h) + (\mathbf{b} \cdot \mathcal{A}e_\sigma, \mathbf{v}_h) + (ce_u, \mathbf{v}_h) &= 0, \quad \mathbf{v}_h \in \mathbf{V}_h, \end{aligned} \quad (2.28)$$

where

$$\mathbf{e}_\sigma := \boldsymbol{\sigma}_h - \hat{\boldsymbol{\sigma}}_h, \quad \mathbf{e}_u := \mathbf{u}_h - \hat{\mathbf{u}}_h.$$

Consider the dual problem

$$\begin{aligned} (\mathcal{A}\tilde{\boldsymbol{\sigma}}, \boldsymbol{\tau}) + (\operatorname{div} \boldsymbol{\tau}, \tilde{\mathbf{u}}) - (\mathbf{b} \cdot \mathcal{A}\boldsymbol{\tau}, \tilde{\mathbf{u}}) &= 0, \quad \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \\ -(\operatorname{div} \tilde{\boldsymbol{\sigma}}, \mathbf{v}) + (c\tilde{\mathbf{u}}, \mathbf{v}) &= (\mathbf{e}_u, \mathbf{v}), \quad \mathbf{v} \in \mathbf{V}. \end{aligned} \quad (2.29)$$

It then follows from (2.28), (2.29) and the same analysis in Lemma 2.1 that

$$\|\mathbf{e}_u\| \lesssim h\|\mathbf{e}_\sigma\|, \quad (2.30)$$

provided h is sufficiently small. The same argument for proving Lemma 2.2 yields

$$\begin{aligned} \|\operatorname{div} \mathbf{e}_\sigma\| &\lesssim \|\mathbf{e}_\sigma\| + \|\mathbf{e}_u\|, \\ \|\operatorname{div} \mathbf{e}_\sigma\|_{-1} &\lesssim h\|\mathbf{e}_\sigma\| + \|\mathbf{e}_u\|. \end{aligned} \quad (2.31)$$

Following the proof of (2.23) and using (2.30), (2.31), we obtain

$$\|\mathbf{e}_\sigma\| \lesssim \|\mathbf{e}_u\|, \quad (2.32)$$

when h is small enough. Finally combining (2.30), (2.32), (2.31) yields

$$\|\mathbf{e}_u\| \lesssim h\|\mathbf{e}_u\|.$$

Hence $\mathbf{e}_u = \mathbf{0}$ and $\mathbf{e}_\sigma = \mathbf{0}$ provided h is sufficiently small. \square

Proof (Proof of Lemma 2.2) Let $\mathbf{v}_h = \frac{\operatorname{div} \boldsymbol{\xi}_h}{\|\operatorname{div} \boldsymbol{\xi}_h\|}$. Using (2.4) and (2.7b), we arrive at

$$\begin{aligned} \|\operatorname{div} \boldsymbol{\xi}_h\| &= (\operatorname{div} \boldsymbol{\xi}_h, \mathbf{v}_h) = (P_h \operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}_h, \mathbf{v}_h) = (\operatorname{div}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}_h) \\ &= (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}_h) + (c(P_h \mathbf{u} - \mathbf{u}_h), \mathbf{v}_h) + ((c - c_h)(\mathbf{u} - P_h \mathbf{u}), \mathbf{v}_h). \end{aligned} \quad (2.33)$$

where c_h is the piecewise average of c w.r.t. \mathcal{T}_h . (2.17a) then follows from (2.33), the Cauchy–Schwarz and triangle inequalities.

To prove (2.17b), let $\mathbf{v} \in [H_0^1(\Omega)]^d$ with $\|\mathbf{v}\|_1 = 1$. It follows from (2.4), (1.8), and (2.7b) that

$$\begin{aligned} (\operatorname{div} \boldsymbol{\xi}_h, \mathbf{v}) &= (P_h \operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}, \mathbf{v}) + (\operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}_h, \mathbf{v}) \\ &= (P_h \operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}, \mathbf{v} - P_h \mathbf{v}) + (\operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}_h, \mathbf{v} - P_h \mathbf{v}) \\ &\quad + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}) + (\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), P_h \mathbf{v} - \mathbf{v}) \\ &\quad + (c(\mathbf{u} - P_h \mathbf{u}), P_h \mathbf{v}) + (c(P_h \mathbf{u} - \mathbf{u}_h), P_h \mathbf{v}). \end{aligned} \quad (2.34)$$

Using (2.1) and (2.7a), we can rewrite $(\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v})$ as

$$\begin{aligned} &(\mathbf{b} \cdot \mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v}) \\ &= (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathbf{v} \mathbf{b}^T) = (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \mathcal{A}(\mathbf{v} \mathbf{b}^T)) \\ &= (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \Pi_h \mathcal{A}(\mathbf{v} \mathbf{b}^T)) + (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), (\mathbf{I} - \Pi_h) \mathcal{A}(\mathbf{v} \mathbf{b}^T)) \\ &= -(\operatorname{div} \Pi_h \mathcal{A}(\mathbf{v} \mathbf{b}^T), \mathbf{u} - \mathbf{u}_h) + (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), (\mathbf{I} - \Pi_h) \mathcal{A}(\mathbf{v} \mathbf{b}^T)) \\ &= -(\operatorname{div} \Pi_h \mathcal{A}(\mathbf{v} \mathbf{b}^T), P_h \mathbf{u} - \mathbf{u}_h) + (\mathcal{A}(\boldsymbol{\sigma} - \boldsymbol{\sigma}_h), (\mathbf{I} - \Pi_h) \mathcal{A}(\mathbf{v} \mathbf{b}^T)). \end{aligned} \quad (2.35)$$

On the other hand,

$$(c(\mathbf{u} - P_h \mathbf{u}), P_h \mathbf{v}) = ((c - c_h)(\mathbf{u} - P_h \mathbf{u}), P_h \mathbf{v}). \quad (2.36)$$

Combining (2.34) with (2.35) and (2.36) and using (2.5), we obtain

$$\begin{aligned} \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1} &= \sup_{\|\mathbf{v}\|_1=1} (\operatorname{div} \boldsymbol{\xi}_h, \mathbf{v}) \\ &\lesssim h(\|P_h \operatorname{div} \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}\| + \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + \|\mathbf{u} - P_h \mathbf{u}\|) \\ &\quad + (1 + \|\operatorname{div} \Pi_h \mathcal{A}(\mathbf{v}\mathbf{b}^T)\|)\|\mathbf{e}_h\|. \end{aligned} \quad (2.37)$$

It follows from (2.4) and elementary calculation that

$$\|\operatorname{div} \Pi_h \mathcal{A}(\mathbf{v}\mathbf{b}^T)\| = \|P_h \operatorname{div} \mathcal{A}(\mathbf{v}\mathbf{b}^T)\| \leq \|\operatorname{div} \mathcal{A}(\mathbf{v}\mathbf{b}^T)\| \leq C\|\mathbf{v}\|_1 \leq C.$$

Therefore using the previous estimate and (2.37), we obtain

$$\|\operatorname{div} \boldsymbol{\xi}\|_{-1} \lesssim h\|\operatorname{div} \Pi_h \boldsymbol{\sigma} - \operatorname{div} \boldsymbol{\sigma}\| + h\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + h\|\boldsymbol{\rho}_h\| + \|\mathbf{e}_h\|.$$

The proof is complete. \square

3 Superconvergence on pseudostress

Using Lemma 2.1 and (2.26), we obtain the supercloseness estimate on $\mathbf{e}_h = P_h \mathbf{u} - \mathbf{u}_h$.

$$\|\mathbf{e}_h\| \lesssim h^{r+2}(|\boldsymbol{\sigma}|_{r+1} + |\operatorname{div} \boldsymbol{\sigma}|_{r+1} + |\mathbf{u}|_{r+1}). \quad (3.1)$$

In this section, we shall prove a supercloseness estimate on $\boldsymbol{\xi}_h = \Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h$. To that end, $\boldsymbol{\Sigma}_h$ is endowed with the Λ -inner product

$$\Lambda(\boldsymbol{\tau}_1, \boldsymbol{\tau}_2) := (\mathcal{A}\boldsymbol{\tau}_1, \boldsymbol{\tau}_2) + (\operatorname{div} \boldsymbol{\tau}_1, \operatorname{div} \boldsymbol{\tau}_2), \quad \boldsymbol{\tau}_1, \boldsymbol{\tau}_2 \in \boldsymbol{\Sigma}_h.$$

Let $\mathbf{Z}_h := \ker(\operatorname{div}|_{\boldsymbol{\Sigma}_h})$ and \mathbf{Z}_h^\perp be the orthogonal complement of \mathbf{Z}_h w.r.t. $\Lambda(\cdot, \cdot)$ in $\boldsymbol{\Sigma}_h$. We shall decompose $\boldsymbol{\xi}_h \in \boldsymbol{\Sigma}_h$ by the discrete Helmholtz decomposition

$$\boldsymbol{\Sigma}_h = \mathbf{Z}_h \oplus_{\Lambda} \mathbf{Z}_h^\perp, \quad (3.2)$$

which can be analyzed using tools in finite element exterior calculus (FEEC), see, e.g., [3, 4]. In the theory of FEEC, an essential ingredient is the bounded projections that commute with exterior differentiation. In particular, there exist projections $\tilde{\pi}_h : [L^2(\Omega)]^d \rightarrow \tilde{\boldsymbol{\Sigma}}_h$ and $Q_h : L^2(\Omega) \rightarrow V_h$ (see, e.g., [3, 25]) such that

$$\begin{aligned} \tilde{\pi}_h|_{\tilde{\boldsymbol{\Sigma}}_h} &= \operatorname{id}, \quad Q_h|_{V_h} = \operatorname{id}, \\ \operatorname{div} \tilde{\pi}_h \boldsymbol{\tau} &= Q_h \operatorname{div} \boldsymbol{\tau} \text{ for } \boldsymbol{\tau} \in H(\operatorname{div}, \Omega), \\ \|\tilde{\pi}_h \boldsymbol{\tau}\| &\lesssim \|\boldsymbol{\tau}\| \text{ for } \boldsymbol{\tau} \in [L^2(\Omega)]^d, \end{aligned} \quad (3.3)$$

where id is the identity operator. Starting from $\tilde{\pi}_h, Q_h$, one can easily obtain commuting bounded projections onto $\boldsymbol{\Sigma}_h$ and \mathbf{V}_h . Let

$$\pi_h \boldsymbol{\tau} := \tilde{\pi}_h \boldsymbol{\tau} - \frac{1}{d|\Omega|} \left(\int_{\Omega} \operatorname{tr} \tilde{\pi}_h \boldsymbol{\tau} dx \right) \mathbf{I},$$

where $\tilde{\pi}_h$ is applied to each row of $\boldsymbol{\tau}$. Similarly, Q_h can be applied to each component of a vector-valued function in $[L^2(\Omega)]^d$. Using the property of $\tilde{\pi}_h$ in (3.3), we have

$$\pi_h|_{\boldsymbol{\Sigma}_h} = \text{id}, \quad Q_h|_{\mathbf{V}_h} = \text{id}, \quad (3.4a)$$

$$\text{div } \pi_h \boldsymbol{\tau} = Q_h \text{div } \boldsymbol{\tau} \text{ for } \boldsymbol{\tau} \in \boldsymbol{\Sigma}, \quad (3.4b)$$

$$\|\pi_h \boldsymbol{\tau}\| \lesssim \|\boldsymbol{\tau}\| \text{ for } \boldsymbol{\tau} \in [L^2(\Omega)]^d. \quad (3.4c)$$

Then with the help of π_h , Q_h , we obtain the following lemma for estimating the decomposition component living in \mathbf{Z}_h^\perp .

Lemma 3.1 *It holds that*

$$\|\boldsymbol{\tau}_h\|_{\mathcal{A}} \lesssim \|\text{div } \boldsymbol{\tau}_h\|_{-1}, \quad \boldsymbol{\tau}_h \in \mathbf{Z}_h^\perp.$$

Proof Let $\phi \in [H_0^1(\Omega)]^d$ be the weak solution to

$$\Delta \phi = \text{div } \boldsymbol{\tau}_h.$$

Then $\boldsymbol{\tau} := \nabla \phi$ satisfies $\text{div } \boldsymbol{\tau} = \text{div } \boldsymbol{\tau}_h$. Using it and (3.4b), (3.4a), we obtain

$$\begin{aligned} \text{div}(\boldsymbol{\tau}_h - \pi_h \boldsymbol{\tau}) &= \text{div } \boldsymbol{\tau}_h - Q_h \text{div } \boldsymbol{\tau} \\ &= \text{div } \boldsymbol{\tau}_h - Q_h \text{div } \boldsymbol{\tau}_h = \mathbf{0}, \end{aligned}$$

i.e., $\boldsymbol{\tau}_h - \pi_h \boldsymbol{\tau} \in \mathbf{Z}_h$. Recall that $\|\cdot\|_{-1}$ denotes the $H^{-1}(\Omega)$ -norm. The natural H^1 -elliptic regularity of $\Delta \phi = \text{div } \boldsymbol{\tau}_h$ yields

$$\|\boldsymbol{\tau}\| \leq \|\phi\|_1 \lesssim \|\text{div } \boldsymbol{\tau}_h\|_{-1}. \quad (3.5)$$

Hence using $\boldsymbol{\tau}_h - \pi_h \boldsymbol{\tau} \in \mathbf{Z}_h$, $\boldsymbol{\tau}_h \in \mathbf{Z}_h^\perp$, (3.5), and (3.4c), we have

$$\begin{aligned} \|\boldsymbol{\tau}_h\|_{\mathcal{A}}^2 &= A(\boldsymbol{\tau}_h - \pi_h \boldsymbol{\tau}, \boldsymbol{\tau}_h) + (A\pi_h \boldsymbol{\tau}, \boldsymbol{\tau}_h) \\ &= (A\pi_h \boldsymbol{\tau}, \boldsymbol{\tau}_h) \leq \|\pi_h \boldsymbol{\tau}\|_{\mathcal{A}} \|\boldsymbol{\tau}_h\|_{\mathcal{A}} \\ &\lesssim \|\boldsymbol{\tau}\| \|\boldsymbol{\tau}_h\|_{\mathcal{A}} \lesssim \|\text{div } \boldsymbol{\tau}_h\|_{-1} \|\boldsymbol{\tau}_h\|_{\mathcal{A}}. \end{aligned}$$

The proof is complete. \square

Remark 1 It follows from the discrete abstract Poincaré inequality (see [4], Theorem 3.6) that

$$\|\boldsymbol{\tau}_h\|_{\mathcal{A}} \lesssim \|\text{div } \boldsymbol{\tau}_h\| \text{ for } \boldsymbol{\tau}_h \in \mathbf{Z}_h^\perp.$$

Hence Lemma 3.1 is an improved discrete Poincaré inequality. The improvement is achieved by utilizing the special property of the divergence operator.

We emphasize that the improved estimate of ξ_h is dependent on mesh structure, type of finite elements, and quite technical. For simplicity of presentation, first let $\Sigma_h \times \mathbf{V}_h$ be based on the lowest order RT element (1.5) with $r = 0$ in \mathbb{R}^2 although the estimates can be generalized in several ways, see the end of this section. For a scalar-valued function v and a vector-valued function $\mathbf{v} = (v_1, v_2)^T$, let

$$\operatorname{curl} v = (-\partial_{x_2} v, \partial_{x_1} v)^T, \quad \operatorname{rot} \mathbf{v} = \partial_{x_1} v_2 - \partial_{x_2} v_1.$$

The row-wise curl is defined as

$$\operatorname{curl} \mathbf{v} = \begin{bmatrix} (\operatorname{curl} v_1)^T \\ (\operatorname{curl} v_2)^T \end{bmatrix}.$$

It is noted that $\operatorname{rot} \mathbf{v} = \operatorname{tr} \operatorname{curl} \mathbf{v}$. Let

$$W_h = \{w \in H^1(\Omega) : w|_K \in \mathcal{P}_1(K) \text{ for } K \in \mathcal{T}_h\},$$

$$\mathbf{W}_h = \{\mathbf{w} \in [W_h]^2 : \int_{\Omega} \operatorname{rot} \mathbf{w} dx = 0\}.$$

Then we obtain the following discrete sequence in \mathbb{R}^2 .

$$\mathbf{W}_h \xrightarrow{\operatorname{curl}} \Sigma_h \xrightarrow{\operatorname{div}} \mathbf{V}_h \rightarrow 0. \quad (3.6)$$

Since $\int \operatorname{rot} \mathbf{w} dx = \int_{\Omega} \operatorname{tr} \operatorname{curl} \mathbf{w}_h dx = 0$ for all $\mathbf{w}_h \in \mathbf{W}_h$, we have $\operatorname{curl} \mathbf{W}_h \subset \Sigma_h$.

The supercloseness estimate of ξ_h does not hold on general unstructured grids. In [43, 7], Li et al. derived several supercloseness estimates on the vector variable in mixed methods for elliptic equations using RT elements under certain mildly structured grids. Readers are referred to [31, 33, 11, 12] for similar results on Poisson's equation under rectangular, h^2 -uniform quadrilateral, and uniform triangular grids. To avoid lengthy descriptions of different mesh structures, we focus on the following piecewise uniform grids.




Fig. 1 Piecewise uniform grid on a square

Definition 3.1 We say \mathcal{T}_h is a uniform grid provided each pair of adjacent triangles (two triangles sharing a common edge) form an exact parallelogram. Let $\{\Omega_i\}_{i=1}^N$ be a fixed polygonal partition of Ω . We say \mathcal{T}_h is a piecewise uniform grid provided \mathcal{T}_h is aligned with $\{\Omega_i\}_{i=1}^N$ and $\mathcal{T}_h|_{\Omega_i}$ is a uniform grid for each $1 \leq i \leq N$.

For instance, let $\Omega = [0, 1]^2$ be the unit square that is split into the 19 triangles $\{\Omega_i\}_{i=1}^{19}$ given in Figure 1 (left). After 3 consecutive uniform quad-refinement, we obtain the triangulation in Figure 1 (right), which is a piecewise uniform grid (w.r.t. $\{\Omega_i\}_{i=1}^{19}$). The piecewise uniform mesh structure is only used in the next technical lemma.

Lemma 3.2 Let $\mathbf{D} \in \mathbb{R}^{2 \times 2}$ be a constant matrix. Let $\tilde{\Pi}_h$ be the canonical interpolation onto $\tilde{\Sigma}_h$, the lowest order RT element space. For a piecewise uniform \mathcal{T}_h and $\mathbf{v} \in W_\infty^2(\Omega)$, $w_h \in W_h$, it holds that

$$(\mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v}), \operatorname{curl} w_h) \lesssim h^2 |\log h|^{\frac{1}{2}} \|\mathbf{v}\|_{W_\infty^2(\Omega)} \|\operatorname{curl} w_h\|.$$

In [43], Li proved Lemma 3.2 on mildly structured grids when $\mathbf{D} = \mathbf{I}$. For the general anisotropic \mathbf{D} , we could not find a complete proof in literature. In the appendix, we give a detailed proof of Lemma 3.2 using the technique in Theorem 3.2 of [11] and Lemma 3.4 of [39]. Now we present a supercloseness estimate for $\boldsymbol{\xi}_h$, the second main result, in the next theorem.

Theorem 3.1 Let Ω be simply-connected and \mathcal{T}_h be a piecewise uniform grid. For (1.10) using the lowest order RT element and sufficiently small h , it holds that

$$\|\boldsymbol{\xi}_h\| \lesssim h^2 |\log h|^{\frac{1}{2}} (\|\boldsymbol{\sigma}\|_{W_\infty^2(\Omega)} + \|\mathbf{u}\|_2).$$

Proof When Ω is simply connected, (3.6) is an exact sequence, i.e., $\operatorname{curl} \mathbf{W}_h = \mathbf{Z}_h$. It then follows from the exactness and the discrete Helmholtz decomposition (3.2) that

$$\boldsymbol{\xi}_h = \operatorname{curl} \boldsymbol{\phi}_h \oplus_A \boldsymbol{\eta}_h, \quad (3.7)$$

where $\boldsymbol{\phi}_h \in \mathbf{W}_h$, $\boldsymbol{\eta}_h \in \mathbf{Z}_h^\perp$. Using Lemma 3.1 and $\operatorname{div} \operatorname{curl} \boldsymbol{\phi}_h = \mathbf{0}$, we obtain

$$\|\boldsymbol{\eta}_h\|_{\mathcal{A}} \lesssim \|\operatorname{div} \boldsymbol{\eta}_h\|_{-1} = \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1}. \quad (3.8)$$

It remains to estimate $\|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}}$. Note that

$$(\mathcal{A} \operatorname{curl} \boldsymbol{\phi}_h, \boldsymbol{\eta}_h) = \Lambda(\operatorname{curl} \boldsymbol{\phi}_h, \boldsymbol{\eta}_h) = 0.$$

Then using the orthogonality, (2.7a), and (2.1), we have

$$\begin{aligned} \|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}}^2 &= (\mathcal{A}(\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h), \operatorname{curl} \boldsymbol{\phi}_h) \\ &= (\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}, \mathcal{A} \operatorname{curl} \boldsymbol{\phi}_h) = (\tilde{\Pi}_h \boldsymbol{\sigma} - \boldsymbol{\sigma}, \mathcal{A} \operatorname{curl} \boldsymbol{\phi}_h) \\ &= -(\boldsymbol{\sigma} - \tilde{\Pi}_h \boldsymbol{\sigma}, \operatorname{curl} \boldsymbol{\phi}_h) + \frac{1}{2}(\boldsymbol{\sigma} - \tilde{\Pi}_h \boldsymbol{\sigma}, (\operatorname{rot} \boldsymbol{\phi}_h) \mathbf{I}). \end{aligned}$$

Recall that $\boldsymbol{\sigma}_i$ denotes the i -th row of $\boldsymbol{\sigma}$. Let $\mathbf{e}_1 = (1, 0)$, $\mathbf{e}_2 = (0, 1)$, and $\boldsymbol{\phi}_h = (\phi_{h,1}, \phi_{h,2})^T$. It follows from the previous equation and direct calculation that

$$\begin{aligned} \|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}}^2 &= \sum_{i=1}^2 -(\boldsymbol{\sigma}_i - \tilde{\Pi}_h \boldsymbol{\sigma}_i, \operatorname{curl} \phi_{h,i}) \\ &+ \frac{1}{2}(\boldsymbol{\sigma}_i - \tilde{\Pi}_h \boldsymbol{\sigma}_i, (\partial_{x_1} \phi_{h,2}) \mathbf{e}_i) - \frac{1}{2}(\boldsymbol{\sigma}_i - \tilde{\Pi}_h \boldsymbol{\sigma}_i, (\partial_{x_2} \phi_{h,1}) \mathbf{e}_i) \\ &= \sum_{i,j=1}^2 (\mathbf{D}_{ij} (\boldsymbol{\sigma}_i - \tilde{\Pi}_h \boldsymbol{\sigma}_i)^T, \operatorname{curl} \phi_{h,j}), \end{aligned} \quad (3.9)$$

where

$$\mathbf{D}_{11} = \begin{pmatrix} -\frac{1}{2} & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{D}_{12} = \begin{pmatrix} 0 & 0 \\ \frac{1}{2} & 0 \end{pmatrix}, \quad \mathbf{D}_{21} = \begin{pmatrix} 0 & \frac{1}{2} \\ 0 & 0 \end{pmatrix}, \quad \mathbf{D}_{22} = \begin{pmatrix} -1 & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}.$$

Then using (3.9), Lemma 3.2 and (2.2) with $\boldsymbol{\tau} = \operatorname{curl} \boldsymbol{\phi}_h$, we have

$$\begin{aligned} \|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}}^2 &\lesssim h^2 |\log h|^{\frac{1}{2}} \|\boldsymbol{\sigma}\|_{W_\infty^2(\Omega)} \|\operatorname{curl} \boldsymbol{\phi}_h\| \\ &\lesssim h^2 |\log h|^{\frac{1}{2}} \|\boldsymbol{\sigma}\|_{W_\infty^2(\Omega)} \|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}}. \end{aligned} \quad (3.10)$$

Combining (2.2) with (3.8), (3.10), (2.17b), (3.1), we obtain

$$\begin{aligned} \|\boldsymbol{\xi}_h\| &\lesssim \|\boldsymbol{\xi}_h\|_{\mathcal{A}} + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1} \\ &\lesssim \|\operatorname{curl} \boldsymbol{\phi}_h\|_{\mathcal{A}} + \|\boldsymbol{\eta}_h\|_{\mathcal{A}} + \|\operatorname{div} \boldsymbol{\xi}_h\|_{-1} \\ &\lesssim h^2 |\log h|^{\frac{1}{2}} (\|\boldsymbol{\sigma}\|_{W_\infty^2(\Omega)} + \|\mathbf{u}\|_2). \end{aligned}$$

The proof is complete. \square

The supercloseness estimate on $\|\boldsymbol{\xi}_h\|$ can be generalized on rectangular meshes. Throughout the rest of this section, let \mathcal{T}_h be a rectangular mesh,

$$\begin{aligned} W_h &= \{\mathbf{w} \in H^1(\Omega) : w|_K \in \mathcal{Q}_{r+1}(K) \text{ for } K \in \mathcal{T}_h\}, \\ \mathbf{W}_h &= \{\mathbf{w} \in [W_h]^2 : \int_{\Omega} \operatorname{rot} \mathbf{w} dx = 0\}, \end{aligned}$$

and $\boldsymbol{\Sigma}_h \times \mathbf{V}_h$ be based on the rectangular RT element (1.7) with $d = 2$. When Ω is simply-connected, we still have the discrete exact sequence (3.6). Similarly to the case of triangular grids, we need the uniform mesh structure.

Definition 3.2 We say \mathcal{T}_h is a uniformly rectangular mesh if all the rectangles of \mathcal{T}_h are of the same shape and size.

For $w_h \in W_h$ with $(\operatorname{curl} w_h) \cdot \mathbf{n}|_{\partial\Omega} = 0$, Theorem 5.1 of [33] implies

$$(\mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v}), \operatorname{curl} w_h) \lesssim h^{r+2} \|\mathbf{v}\|_{r+2} \|\operatorname{curl} w_h\|,$$

where $\mathbf{v} \in [H^{r+2}(\Omega)]^2$. The previous estimate is not true when $(\operatorname{curl} w_h) \cdot \mathbf{n} \neq 0$ on $\partial\Omega$. As claimed in the remark in Example 6.2 of [33], it holds that

$$(\mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v}), \operatorname{curl} w_h) \leq Ch^{r+1.5} \|\operatorname{curl} w_h\|, \quad w_h \in W_h \quad (3.11)$$

for some absolute constant C independent of h . Following the same argument in the proof of Theorem 3.1 and replacing Lemma 3.2 by (3.11), we actually obtain the supercloseness estimate

$$\|\Pi_h \boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| = \mathcal{O}(h^{r+1.5}).$$

Similar estimates may hold on h^2 -uniform quadrilateral grids described in [33], which is beyond the scope of this paper.

4 Postprocessing

We have derived supercloseness estimates in Lemma 2.1 and Theorem 3.1. However, those results can not be directly used because $P_h \mathbf{u}$ and $\Pi_h \boldsymbol{\sigma}$ are not known in practice. To extract superconvergence information from the smallness of \mathbf{e}_h and $\boldsymbol{\xi}_h$, one may design easy-to-compute postprocessed solutions \mathbf{u}_h^* and $\boldsymbol{\sigma}_h^*$. For instance, following the idea of [53], a element-by-element postprocessing procedure for \mathbf{u}_h in the pseudostress-velocity formulation of the Stokes equation is proposed in [23]. In particular, let

$$\mathcal{P}_{r+1}(K)/\mathcal{P}_r(K) := \{\mathbf{v}_h \in \mathcal{P}_{r+1}(K) : (\mathbf{v}_h, \mathbf{w}_h)_K = 0 \text{ for all } \mathbf{w}_h \in \mathcal{P}_r(K)\}.$$

where $(\cdot, \cdot)_K$ is the $L^2(K)$ -inner product. For each $K \in \mathcal{T}_h$, the postprocessed solution $\mathbf{u}_h^*|_K \in \mathcal{P}_{r+1}(K)$ is the unique solution to

$$\begin{aligned} (\mathbf{u}_h^*, \mathbf{v}_h)_K &= (\mathbf{u}_h, \mathbf{v}_h)_K, \\ (\nabla \mathbf{u}_h^*, \nabla \mathbf{v}_h)_K &= (\boldsymbol{\sigma}_h, \nabla \mathbf{v}_h)_K + (p_h, \operatorname{div} \mathbf{v}_h)_K, \quad \mathbf{v}_h \in \mathcal{P}_{r+1}(K)/\mathcal{P}_r(K), \end{aligned} \quad (4.1)$$

where $p_h := -\frac{1}{d} \operatorname{tr} \boldsymbol{\sigma}_h \approx p$. Since the analysis of the mapping $\mathbf{u}_h \mapsto \mathbf{u}_h^*$ is independent of equations, we can combine the analysis in Theorem 4.1 of [23] for the Stokes equation with the supercloseness on $\|P_h \mathbf{u} - \mathbf{u}_h\|$ in Theorem 2.1 to obtain the following postprocessing superconvergence estimate.

Theorem 4.1 *For sufficiently small h , it holds that*

$$\|\mathbf{u} - \mathbf{u}_h^*\| \lesssim h^{r+2} (\|\mathbf{u}\|_{r+2} + |\boldsymbol{\sigma}|_{r+1} + |\operatorname{div} \boldsymbol{\sigma}|_{r+1}).$$

Proof Let $\hat{\mathbf{u}}$ be the L^2 -projection of \mathbf{u} onto

$$\{\mathbf{v}_h : \mathbf{v}_h|_K \in [\mathcal{P}_{r+1}(K)]^d \text{ for all } K \in \mathcal{T}_h\}.$$

It has been shown in the proof of Theorem 4.1 in [23] that

$$\begin{aligned} h^{-1} \|(\mathbf{I} - P_h)(\hat{\mathbf{u}} - \mathbf{u}_h^*)\| &\lesssim h^{-1} \|P_h(\hat{\mathbf{u}} - \mathbf{u}_h^*)\| \\ &+ \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\| + \left(\sum_{K \in \mathcal{T}_h} \|\nabla(\hat{\mathbf{u}} - \mathbf{u})\|_{L^2(K)}^2 \right)^{\frac{1}{2}}. \end{aligned} \quad (4.2)$$

Using the triangle inequality, $P_h \hat{\mathbf{u}} = P_h \mathbf{u}$, $P_h \mathbf{u}_h^* = \mathbf{u}_h$, and (4.2), we obtain

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h^*\| &\leq \|\mathbf{u} - \hat{\mathbf{u}}\| + \|(\mathbf{I} - P_h)(\hat{\mathbf{u}} - \mathbf{u}_h^*)\| + \|P_h(\hat{\mathbf{u}} - \mathbf{u}_h^*)\| \\ &\lesssim h^{r+2} |\mathbf{u}|_{r+2} + \|P_h \mathbf{u} - \mathbf{u}_h\| + h \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|. \end{aligned}$$

We finally conclude the proof from the previous estimate with (3.1) and (2.26b). \square

Postprocessing technique on the scalar variable in mixed methods for Poisson's equation can be found in e.g., [15, 10, 53, 46].

The postprocessing procedure $\boldsymbol{\sigma}_h \mapsto \boldsymbol{\sigma}_h^*$ can be derived from existing post-processing operator \tilde{R}_h . In particular, $\tilde{R}_h : \tilde{\boldsymbol{\Sigma}}_h \rightarrow Y \subset [L^2(\Omega)]^2$ is a linear mapping onto the space Y of suitable piecewise polynomials and satisfies

$$\tilde{R}_h \mathbf{c} = \mathbf{c}, \quad \mathbf{c} \in \mathbb{R}^2, \quad (4.3a)$$

$$\|\tilde{R}_h \mathbf{v}\| \lesssim \|\mathbf{v}\|, \quad \mathbf{v} \in \tilde{\boldsymbol{\Sigma}}_h, \quad (4.3b)$$

$$\|\boldsymbol{\tau} - \tilde{R}_h \tilde{\Pi}_h \boldsymbol{\tau}\| \lesssim h^{1+\alpha} \|\boldsymbol{\tau}\|_{W_\infty^2(\Omega)}, \quad (4.3c)$$

for some positive constant α and sufficiently smooth $\boldsymbol{\tau}$. When $\tilde{\boldsymbol{\Sigma}}_h$ is the lowest order RT element space, \tilde{R}_h that satisfies (4.3) is given in e.g., [11, 7]. The simple nodal averaging [58] and superconvergent patch recovery [59] are also possible choices.

For $\boldsymbol{\tau} \in \boldsymbol{\Sigma}_h$, let

$$R_h \boldsymbol{\tau} := \tilde{R}_h \boldsymbol{\tau} - \frac{1}{2|\Omega|} \left(\int_\Omega \text{tr} \tilde{R}_h \boldsymbol{\tau} \right) \mathbf{I},$$

where \tilde{R}_h is applied to each row of $\boldsymbol{\tau}$. We have the following super-approximation result of R_h .

Lemma 4.1 *Assume (4.3) holds. For $\boldsymbol{\tau} \in \boldsymbol{\Sigma} \cap [W_\infty^2(\Omega)]^{2 \times 2}$, we have*

$$\|\boldsymbol{\tau} - R_h \Pi_h \boldsymbol{\tau}\| \lesssim h^{1+\alpha} \|\boldsymbol{\tau}\|_{W_\infty^2(\Omega)}.$$

Proof (4.3a) implies $R_h \mathbf{I} = \mathbf{O}$ and thus $R_h \Pi_h \boldsymbol{\tau} = R_h \tilde{\Pi}_h \boldsymbol{\tau}$. Then

$$\begin{aligned} \boldsymbol{\tau} - R_h \Pi_h \boldsymbol{\tau} &= \boldsymbol{\tau} - R_h \tilde{\Pi}_h \boldsymbol{\tau} \\ &= \boldsymbol{\tau} - \tilde{R}_h \tilde{\Pi}_h \boldsymbol{\tau} - \frac{1}{2|\Omega|} \int_\Omega \text{tr}(\boldsymbol{\tau} - \tilde{R}_h \tilde{\Pi}_h \boldsymbol{\tau}) dx, \end{aligned} \quad (4.4)$$

where we use $\int_\Omega \text{tr} \boldsymbol{\tau} dx = 0$. Lemma 4.1 then follows from (4.4), the Cauchy-Schwarz inequality, and (4.3c). \square

Combining Lemma 4.1 and Theorem 3.1, we obtain the following super-convergence estimate by postprocessing.

Theorem 4.2 Let σ_h be the solution to (1.10) based on the lowest order RT element and $\sigma_h^* := R_h \sigma_h$. Assume the assumptions in Theorem 3.1 and Lemma 4.1 hold. We have

$$\|\sigma - \sigma_h^*\| \lesssim \min(h^2 |\log h|^{\frac{1}{2}}, h^{1+\alpha}) (\|\sigma\|_{W_\infty^2(\Omega)} + \|\mathbf{u}\|_2).$$

Proof Using the triangle inequality, Theorem 3.1, and Lemma 4.1, we have

$$\begin{aligned} \|\sigma - \sigma_h^*\| &\leq \|\sigma - R_h \Pi_h \sigma\| + \|R_h(\Pi_h \sigma - \sigma_h)\| \\ &\lesssim \min(h^2 |\log h|^{\frac{1}{2}}, h^{1+\alpha}) (\|\sigma\|_{W_\infty^2(\Omega)} + \|\mathbf{u}\|_2), \end{aligned}$$

which completes the proof. \square

Remark 2 Built upon Theorem 4.2, we can recover very accurate numerical symmetric stress σ_h^{S*} and pressure p_h^* . In fact, let

$$\sigma_h^{S*} := \frac{1}{2}(\sigma_h^* + \sigma_h^{*T}), \quad p_h^* := -\frac{1}{d} \operatorname{tr} \sigma_h^*.$$

Recall σ^S is the symmetric stress. We conclude that σ_h^{S*} superconverges to σ^S and p_h^* superconverges to p in the L^2 -norm from Theorem 4.2 and

$$\begin{aligned} \|\sigma^S - \sigma_h^{S*}\| &\leq \frac{1}{2} \|\sigma - \sigma_h^*\| + \frac{1}{2} \|\sigma^T - \sigma_h^{*T}\| = \|\sigma - \sigma_h^*\|, \\ \|p - p_h^*\| &\leq \frac{1}{d} \|\operatorname{tr}(\sigma - \sigma_h^*)\| \leq \frac{1}{\sqrt{d}} \|\sigma - \sigma_h^*\|. \end{aligned}$$

5 Experiments

In this section, the postprocessing procedure $\sigma_h \mapsto \sigma_h^*$ in Theorem 4.2 is based on the polynomial preserving recovery \tilde{R}_h for the lowest order RT element described in [7]. In the first experiment, we verify our a priori and superconvergence error estimates. We also test the adaptivity performance of the recovery a posteriori error estimator in the other two numerical experiments.

Table 1 Convergence of the lowest order RT element

nt	$\ \mathbf{u} - \mathbf{u}_h\ $	$\ e_h\ $	$\ \mathbf{u} - \mathbf{u}_h^*\ $	$\ \sigma - \sigma_h\ $	$\ \xi_h\ $	$\ \sigma - \sigma_h^*\ $
19	3.210e-1	3.851e-2	9.334e-2	1.372	2.505e-1	2.113
76	1.620e-1	1.027e-2	2.438e-2	6.849e-1	7.172e-2	6.108e-1
304	8.121e-2	2.621e-3	6.176e-3	3.418e-1	1.957e-2	1.724e-1
1216	4.063e-2	6.598e-4	1.550e-3	1.707e-1	5.242e-3	4.353e-2
4864	2.032e-2	1.653e-4	3.880e-4	8.530e-2	1.390e-3	1.088e-2
19456	1.016e-2	4.136e-5	9.703e-5	4.270e-2	3.659e-4	2.719e-3
order	9.990e-1	1.990	1.994	1.001	1.904	1.961

Table 2 Convergence of the lowest order BDM element

nt	$\ \mathbf{u} - \mathbf{u}_h\ $	$\ e_h\ $	$\ \mathbf{u} - \mathbf{u}_h^*\ $	$\ \boldsymbol{\sigma} - \boldsymbol{\sigma}_h\ $	$\ \boldsymbol{\xi}_h\ $
19	3.192e-1	1.779e-2	6.316e-02	4.480e-1	4.755e-1
76	1.618e-1	6.026e-3	1.645e-02	1.249e-1	1.180e-1
304	8.118e-2	1.636e-3	4.155e-03	3.216e-2	2.928e-2
1216	4.062e-2	4.181e-4	1.042e-03	8.104e-3	7.288e-3
4864	2.032e-2	1.051e-4	2.606e-04	2.031e-3	1.818e-3
19456	1.016e-2	2.631e-5	6.515e-05	5.081e-4	4.541e-4
order	9.986e-1	1.964	1.996	1.987	2.005

Problem 1: Consider the Oseen equation (1.2) on the unit square $\Omega = [0, 1]^2$ with the smooth solutions

$$\mathbf{u} = \begin{pmatrix} \sin(\pi(x_1 + x_2)) \\ -\sin(\pi(x_1 + x_2)) \end{pmatrix},$$

$$p = x_1 + x_2 - 1.$$

Set $c = 0$, $\mathbf{b} = (\cos(x_2), \sin(x_1))^T$, $\mathbf{g} = \mathbf{u}|_{\partial\Omega} = \mathbf{0}$. \mathbf{f} is computed from \mathbf{u} and \mathbf{b} . We start with the initial partition in Figure 1 (left). A sequence of piecewise uniform meshes is obtained by uniform quad-refinement, i.e., dividing each triangle into four similar subtriangles by connecting the midpoints of each edge. Numerical results are presented in Tables 1 and 2, where nt is short for “number of triangles”. The order of convergence is computed from the error quantities in those tables by least squares without using the data in the first rows.

The numerical rates of convergence coincide with a priori error estimates (2.26), (2.27) and the supercloseness estimates in (3.1) and Theorem 3.1. It is noted that for the lowest order BDM element, $\|\boldsymbol{\xi}_h\| \approx \mathcal{O}(h^2)$, which is predicted by a priori error estimates and thus not supersmall. Since the recovery procedure in [7] provides the super-approximation rate $\alpha = 1$ in (4.3c) and Lemma 4.1, the recovery superconvergence estimate for the lowest order RT element predicted by Theorem 4.2 is $\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h^*\| = \mathcal{O}(h^2 |\log h|^{\frac{1}{2}})$, numerically confirmed by the last column in Table 1.

Problem 2: Consider (1.2) on the L-shaped domain $\Omega = [-1, 1]^2 \setminus ([0, 1] \times [-1, 0])$ with the smooth pressure $p = x_1 + x_2$ and singular velocity

$$\mathbf{u} = \begin{pmatrix} r^\alpha \sin(\alpha\theta) \\ r^\alpha \cos(\alpha\theta) \end{pmatrix},$$

where $\alpha = \frac{2}{3}$, (r, θ) is the polar coordinate w.r.t. $(0, 0)$. Set $c = 0$, $\mathbf{b} = (1, 2)^T$ and $\mathbf{g} = \mathbf{u}|_{\partial\Omega}$. Direct calculation shows that $\operatorname{div} \mathbf{u} = 0$ and $\mathbf{f} = (1, 1)^T + \mathbf{b} \cdot \nabla \mathbf{u}$. In this experiment, we use the classical adaptive feedback loop (cf. [6, 29, 47])

SOLVE \rightarrow ESTIMATE \rightarrow MARK \rightarrow REFINE

to obtain a sequence of adaptively refined grids $\{\mathcal{T}_{h_\ell}\}_{\ell \geq 0}$ and numerical solutions $(\{\boldsymbol{\sigma}_{h_\ell}, \mathbf{u}_{h_\ell}\})_{\ell \geq 0}$. In particular, the algorithm starts from the initial grid




Fig. 2 Adaptive mesh for the singular problem




Fig. 3 Convergence history of the adaptive method

\mathcal{T}_{h_0} presented in Figure 2(left). The module ESTIMATE computes the superconvergent recovery-based error indicator

$$\mathcal{E}_\ell(K) := \left(\|\sigma_{h_\ell}^* - \sigma_{h_\ell}\|_{L^2(K)}^2 + \|\mathbf{u}_{h_\ell}^* - \mathbf{u}_{h_\ell}\|_{L^2(K)}^2 \right)^{\frac{1}{2}}$$

for each triangle $K \in \mathcal{T}_{h_\ell}$. The module MARK then selects a collection of triangles

$$\mathcal{M}_\ell := \{K \in \mathcal{T}_\ell : \mathcal{E}_\ell(K) \geq 0.7 \max_{K' \in \mathcal{T}_\ell} \mathcal{E}_\ell(K')\} \quad (5.1)$$

to be refined by local quad-refinement. To remove the newly created hanging nodes, minimal number of neighboring elements of \mathcal{M}_ℓ are bisected and the next level triangulation $\mathcal{T}_{h_{\ell+1}}$ is generated. See Figure 2(right) for an adaptively refined triangulation.

It can be observed from Figure 3 that $(\sigma_{h_\ell}, \mathbf{u}_{h_\ell})$ optimally converges to (σ, \mathbf{u}) . In addition, the errors $\|\sigma - \sigma_{h_\ell}^*\|$ and $\|\mathbf{u} - \mathbf{u}_{h_\ell}^*\|$ are apparently super-

convergent to 0. Let

$$\mathcal{E}_\ell := \left(\sum_{K \in \mathcal{T}_{h_\ell}} \mathcal{E}_\ell(K)^2 \right)^{\frac{1}{2}} = \left(\|\boldsymbol{\sigma}_{h_\ell}^* - \boldsymbol{\sigma}_{h_\ell}\|^2 + \|\mathbf{u}_{h_\ell}^* - \mathbf{u}_{h_\ell}\|^2 \right)^{\frac{1}{2}},$$

$$E_\ell := \left(\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_{h_\ell}\|^2 + \|\mathbf{u} - \mathbf{u}_{h_\ell}\|^2 \right)^{\frac{1}{2}}.$$

Due to the observed superconvergence phenomena and the triangle inequality

$$|\mathcal{E}_\ell - E_\ell| \leq \left(\|\boldsymbol{\sigma} - \boldsymbol{\sigma}_{h_\ell}^*\|^2 + \|\mathbf{u} - \mathbf{u}_{h_\ell}^*\|^2 \right)^{\frac{1}{2}},$$

the recovery-based error estimator \mathcal{E}_ℓ is asymptotically exact, i.e.,

$$\lim_{\ell \rightarrow \infty} \frac{\mathcal{E}_\ell}{E_\ell} = 1.$$




Fig. 4 (left) Initial grid. (right) Adaptive grid, 9683 triangles.




Fig. 5 Velocity field of the convection-dominated problem

Problem 3: Consider the Oseen equation (1.2) on the unit square $\Omega = [0, 1]^2$ with $c = 0$, $\mathbf{b} = (500, 1)^T$, $\mathbf{g} = \mathbf{0}$, $\mathbf{f} = 5000(x_2, -x_1)^T$. In the end, we use the same adaptive algorithm in Problem 2 to solve this convection-dominated example. The initial grid is given in Figure 4 (left). The marked set in (5.1) is replaced by

$$\mathcal{M}_\ell = \{K \in \mathcal{T}_\ell : \mathcal{E}_\ell(K) \geq 0.3 \max_{K' \in \mathcal{T}_\ell} \mathcal{E}_\ell(K')\}.$$

Due to the convection coefficient $\mathbf{b} = (500, 1)^T$, the exact solution \mathbf{u} near the axis $x_1 = 1$ is rapidly changing to preserve the homogeneous Dirichlet boundary condition. It can be observed from Figures 4 and 5 that the adaptive mixed method is able to capture the boundary layer without using a uniformly fine grid.

Appendix




Fig. 6 A uniform grid of Ω_i .

Proof (Proof of Lemma 3.2) Let $\{\Omega_i\}_{i=1}^N$ be a polygonal partition of Ω and $\mathcal{T}_h|_{\Omega_i}$ be a uniform grid for each i . Let $\mathbf{f}_1, \mathbf{f}_2$ be fixed unit normals to two edges of an arbitrary but fixed triangle in $\mathcal{T}_h|_{\Omega_i}$. Let $\mathbf{S} = \begin{pmatrix} \mathbf{f}_1^T \\ \mathbf{f}_2^T \end{pmatrix}$ and recall $\mathbf{e}_1 = (1, 0)$, $\mathbf{e}_2 = (0, 1)$. We have

$$\begin{aligned} \int_{\Omega_i} [\mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v})] \cdot \text{curl } w_h dx &= \int_{\Omega_i} \mathbf{S}^{-T} [\mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v})] \cdot \mathbf{S} \text{curl } w_h dx \\ &= \sum_{j=1}^2 \int_{\Omega_i} [\mathbf{e}_j \mathbf{S}^{-T} \mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v})] (\mathbf{f}_j^T \text{curl } w_h) dx := \sum_{j=1}^2 I_j. \end{aligned} \quad (5.2)$$

For $j = 1, 2$, let \mathcal{E}_j° and \mathcal{E}_j^∂ be the set of interior (inside Ω_i) and boundary (on $\partial\Omega_i$) edges orthogonal to \mathbf{f}_j , respectively. Let N_E denote the region which

is the union of triangles sharing E as an edge. To estimate I_1 , let Ω_i be partitioned into parallelograms N_E with $E \in \mathcal{E}_1^o$ and boundary triangles N_E with $E \in \mathcal{E}_1^\partial$, see Figure 6, where $E_1, E_2 \in \mathcal{E}_1^\partial$, $E_3, E_4 \in \mathcal{E}_1^o$. For any $E \in \mathcal{E}_1^o$, note that $\mathbf{f}_1^T \operatorname{curl} w_h$ is single-valued across E and thus constant on N_E . For $E \in \mathcal{E}_1^\partial$, let $P_{E,1}$ and $P_{E,2}$ be the two endpoints of E and h_E denote the length of E . Then

$$\begin{aligned} I_1 &= \sum_{E \in \mathcal{E}_1^o \cup \mathcal{E}_1^\partial} \int_{N_E} [\mathbf{e}_1 \mathbf{S}^{-T} \mathbf{D}(\mathbf{v} - \tilde{\Pi}_h \mathbf{v})] (\mathbf{f}_1^T \operatorname{curl} w_h) dx \\ &= \sum_{E \in \mathcal{E}_1^o} (\mathbf{f}_1^T \operatorname{curl} w_h) \mathbf{e}_1 \mathbf{S}^{-T} \mathbf{D} \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \\ &\quad + \sum_{E \in \mathcal{E}_1^\partial} h_E^{-1} \mathbf{e}_1 \mathbf{S}^{-T} \mathbf{D} \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx (w_h(P_{E,2}) - w_h(P_{E,1})) \\ &:= I_{11} + I_{12}. \end{aligned} \tag{5.3}$$

When N_E is an exact parallelogram, it is shown in Equation (3.15) of [11] that

$$\left| \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \right| \lesssim h^3 |\mathbf{v}|_{H^2(N_E)}. \tag{5.4}$$

Using (5.4) and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} I_{11} &\lesssim \sum_{E \in \mathcal{E}_1^o} h^{-1} \|\operatorname{curl} w_h\|_{L^2(N_E)} \left| \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \right| \\ &\lesssim \sum_{E \in \mathcal{E}_1^o} h^2 \|\operatorname{curl} w_h\|_{L^2(N_E)} |\mathbf{v}|_{H^2(N_E)} \lesssim h^2 \|\operatorname{curl} w_h\| \|\mathbf{v}\|_2. \end{aligned} \tag{5.5}$$

Let \mathcal{N} denote the collection of endpoints of edges in \mathcal{E}_1^∂ ,

$$\mathcal{N}_1 := \{P \in \mathcal{N} : P \text{ is not a corner of } \partial\Omega_i\},$$

and $\mathcal{N}_2 := \mathcal{N} \setminus \mathcal{N}_1$. For instance, $\mathcal{N}_2 = \{P_1, P_2\}$ in Figure 6. For $P \in \mathcal{N}_1$, let E_1, E_2 be the two boundary edges sharing P and ω_P be the union of three triangles having P as a vertex. For $P \in \mathcal{N}_2$, let E denote the unique edge in \mathcal{E}_1^∂ having P as a vertex. Rearranging the summation in I_{12} , we have

$$\begin{aligned} I_{12} &= \sum_{P \in \mathcal{N}_1} h_{E_1}^{-1} \mathbf{e}_1 \mathbf{S}^{-T} \mathbf{D} \left(\int_{N_{E_1}} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx - \int_{N_{E_2}} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \right) w_h(P) \\ &\quad + \sum_{P \in \mathcal{N}_2} h_E^{-1} \mathbf{e}_1 \mathbf{S}^{-T} \mathbf{D} \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx w_h(P). \end{aligned} \tag{5.6}$$

For $P \in \mathcal{N}_1$, Equation (3.15) in [39] shows that

$$\left| \int_{N_{E_1}} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx - \int_{N_{E_2}} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \right| \lesssim h^3 |\mathbf{v}|_{H^2(\omega_P)}. \tag{5.7}$$

Without loss of generality, let $\int_{\Omega} w_h dx = 0$. Then the discrete Sobolev and Poincaré inequalities yield

$$\|w_h\|_{L^\infty(\Omega)} \lesssim |\log h|^{\frac{1}{2}} \|w_h\|_1 \lesssim |\log h|^{\frac{1}{2}} \|\operatorname{curl} w_h\|. \quad (5.8)$$

Standard interpolation error estimate yields

$$\left| \int_{N_E} (\mathbf{v} - \tilde{\Pi}_h \mathbf{v}) dx \right| \lesssim h^2 |\mathbf{v}|_{H^1(N_E)}. \quad (5.9)$$

It then follows from (5.6)–(5.9), and the Cauchy–Schwarz inequality that

$$\begin{aligned} I_{12} &\lesssim \left(\sum_{P \in \mathcal{N}_1} h^2 |\mathbf{v}|_{H^2(\omega_P)} + \sum_{P \in \mathcal{N}_2} h |\mathbf{v}|_{H^1(N_E)} \right) \|w_h\|_{L^\infty(\Omega)} \\ &\lesssim (h^2 |\mathbf{v}|_{W_\infty^2(\Omega)} \sum_{P \in \mathcal{N}_1} |\omega_P|^{\frac{1}{2}} + h |\mathbf{v}|_{W_\infty^1(\Omega)} \sum_{P \in \mathcal{N}_2} |N_E|^{\frac{1}{2}}) |\log h|^{\frac{1}{2}} \|\operatorname{curl} w_h\| \\ &\lesssim h^2 \|\mathbf{v}\|_{W_\infty^2(\Omega)} |\log h|^{\frac{1}{2}} \|\operatorname{curl} w_h\|. \end{aligned} \quad (5.10)$$

We finally conclude the proof from (5.2), (5.5), (5.10), the same analysis for I_2 and all pieces in the partition $\{\Omega_i\}_{i=1}^N$. \square

References

1. Antonietti, P.F., Beirão da Veiga, L., Mora, D., Verani, M.: A stream virtual element formulation of the Stokes problem on polygonal meshes. *SIAM J. Numer. Anal.* **52**(1), 386–404 (2014). DOI 10.1137/13091141X. URL <https://doi.org/10.1137/13091141X>
2. Arnold, D.N., Douglas Jr., J., Gupta, C.P.: A family of higher order mixed finite element methods for plane elasticity. *Numer. Math.* **45**(1), 1–22 (1984). DOI 10.1007/BF01379659. URL <https://doi.org/10.1007/BF01379659>
3. Arnold, D.N., Falk, R.S., Winther, R.: Finite element exterior calculus, homological techniques, and applications. *Acta Numer.* **15**, 1–155 (2006). DOI 10.1017/S0962492906210018. URL <https://doi.org/10.1017/S0962492906210018>
4. Arnold, D.N., Falk, R.S., Winther, R.: Finite element exterior calculus: from Hodge theory to numerical stability. *Bull. Amer. Math. Soc. (N.S.)* **47**(2), 281–354 (2010). DOI 10.1090/S0273-0979-10-01278-4. URL <https://doi.org/10.1090/S0273-0979-10-01278-4>
5. Arnold, D.N., Li, L.: Finite element exterior calculus with lower-order terms. *Math. Comp.* **86**(307), 2193–2212 (2017). DOI 10.1090/mcom/3158. URL <https://doi.org/10.1090/mcom/3158>
6. Babuška, I., Rheinboldt, W.C.: Error estimates for adaptive finite element computations. *SIAM J. Numer. Anal.* **15**(4), 736–754 (1978). DOI 10.1137/0715049. URL <https://doi.org/10.1137/0715049>
7. Bank, R.E., Li, Y.: Superconvergent recovery of Raviart–Thomas mixed finite elements on triangular grids. *J. Sci. Comput.* **81**, 1882–1905 (2019)
8. Bank, R.E., Xu, J.: Asymptotically exact a posteriori error estimators. I. Grids with superconvergence. *SIAM J. Numer. Anal.* **41**(6), 2294–2312 (2003). DOI 10.1137/S003614290139874X. URL <https://doi.org/10.1137/S003614290139874X>
9. Boffi, D., Brezzi, F., Fortin, M.: Mixed finite element methods and applications, *Springer Series in Computational Mathematics*, vol. 44. Springer, Heidelberg (2013). DOI 10.1007/978-3-642-36519-5
10. Bramble, J.H., Xu, J.: A local post-processing technique for improving the accuracy in mixed finite-element approximations. *SIAM J. Numer. Anal.* **26**(6), 1267–1275 (1989). DOI 10.1137/0726073

11. Brandts, J.H.: Superconvergence and a posteriori error estimation for triangular mixed finite elements. *Numer. Math.* **68**(3), 311–324 (1994). DOI 10.1007/s002110050064. URL <https://doi.org/10.1007/s002110050064>
12. Brandts, J.H.: Superconvergence for triangular order $k = 1$ Raviart-Thomas mixed finite elements and for triangular standard quadratic finite element methods. *Appl. Numer. Math.* **34**(1), 39–58 (2000). DOI 10.1016/S0168-9274(99)00034-3. URL [https://doi.org/10.1016/S0168-9274\(99\)00034-3](https://doi.org/10.1016/S0168-9274(99)00034-3)
13. Brezzi, F., Douglas Jr., J., Durán, R., Fortin, M.: Mixed finite elements for second order elliptic problems in three variables. *Numer. Math.* **51**(2), 237–250 (1987). DOI 10.1007/BF01396752. URL <https://doi.org/10.1007/BF01396752>
14. Brezzi, F., Douglas Jr., J., Fortin, M., Marini, L.D.: Efficient rectangular mixed finite elements in two and three space variables. *RAIRO Modél. Math. Anal. Numér.* **21**(4), 581–604 (1987). DOI 10.1051/m2an/1987210405811. URL <https://doi.org/10.1051/m2an/1987210405811>
15. Brezzi, F., Douglas Jr., J., Marini, L.D.: Two families of mixed finite elements for second order elliptic problems. *Numer. Math.* **2**(47), 217–235 (1985)
16. Cáceres, E., Gatica, G.N.: A mixed virtual element method for the pseudostress-velocity formulation of the Stokes problem. *IMA J. Numer. Anal.* **37**(1), 296–331 (2017). DOI 10.1093/imanum/drw002. URL <https://doi.org/10.1093/imanum/drw002>
17. Cáceres, E., Gatica, G.N., Sequeira, F.A.: A mixed virtual element method for the Brinkman problem. *Math. Models Methods Appl. Sci.* **27**(4), 707–743 (2017). DOI 10.1142/S0218202517500142. URL <https://doi.org/10.1142/S0218202517500142>
18. Cai, Z., Tong, C., Vassilevski, P.S., Wang, C.: Mixed finite element methods for incompressible flow: stationary Stokes equations. *Numer. Methods Partial Differential Equations* **26**(4), 957–978 (2010). DOI 10.1002/num.20467. URL <https://doi.org/10.1002/num.20467>
19. Cai, Z., Wang, C., Zhang, S.: Mixed finite element methods for incompressible flow: stationary Navier-Stokes equations. *SIAM J. Numer. Anal.* **48**(1), 79–94 (2010). DOI 10.1137/080718413. URL <https://doi.org/10.1137/080718413>
20. Cai, Z., Wang, Y.: A multigrid method for the pseudostress formulation of Stokes problems. *SIAM J. Sci. Comput.* **29**(5), 2078–2095 (2007). DOI 10.1137/060661429. URL <https://doi.org/10.1137/060661429>
21. Carrero, J., Cockburn, B., Schötzau, D.: Hybridized globally divergence-free LDG methods. I. The Stokes problem. *Math. Comp.* **75**(254), 533–563 (2006). DOI 10.1090/S0025-5718-05-01804-1. URL <https://doi.org/10.1090/S0025-5718-05-01804-1>
22. Carstensen, C., Gallistl, D., Schedensack, M.: Quasi-optimal adaptive pseudostress approximation of the Stokes equations. *SIAM J. Numer. Anal.* **51**(3), 1715–1734 (2013)
23. Carstensen, C., Kim, D., Park, E.J.: A priori and a posteriori pseudostress-velocity mixed finite element error analysis for the Stokes problem. *SIAM J. Numer. Anal.* **49**(6), 2501–2523 (2011)
24. Cesmelioglu, A., Cockburn, B., Nguyen, N.C., Peraire, J.: Analysis of HDG methods for Oseen equations. *J. Sci. Comput.* **55**(2), 392–431 (2013). DOI 10.1007/s10915-012-9639-y. URL <https://doi.org/10.1007/s10915-012-9639-y>
25. Christiansen, S.H., Winther, R.: Smoothed projections in finite element exterior calculus. *Math. Comp.* **77**(262), 813–829 (2008)
26. Cockburn, B., Sayas, F.J.: Divergence-conforming HDG methods for Stokes flows. *Math. Comp.* **83**(288), 1571–1598 (2014). DOI 10.1090/S0025-5718-2014-02802-0. URL <https://doi.org/10.1090/S0025-5718-2014-02802-0>
27. Cui, M., Ye, X.: Superconvergence of finite volume methods for the Stokes equations. *Numer. Methods Partial Differential Equations* **25**(5), 1212–1230 (2009). DOI 10.1002/num.20399. URL <https://doi.org/10.1002/num.20399>
28. Demlow, A.: Suboptimal and optimal convergence in mixed finite element methods. *SIAM J. Numer. Anal.* **39**(6), 1938–1953 (2002). DOI 10.1137/S0036142900376900
29. Dörfler, W.: A convergent adaptive algorithm for Poisson’s equation. *SIAM J. Numer. Anal.* **33**(3), 1106–1124 (1996). DOI 10.1137/0733054. URL <https://doi.org/10.1137/0733054>
30. Douglas Jr., J., Roberts, J.E.: Global estimates for mixed methods for second order elliptic equations. *Math. Comp.* **44**(169), 39–52 (1985). DOI 10.2307/2007791. URL <https://doi.org/10.2307/2007791>

31. Durán, R.: Superconvergence for rectangular mixed finite elements. *Numer. Math.* **58**(3), 287–298 (1990). DOI 10.1007/BF01385626. URL <https://doi.org/10.1007/BF01385626>
32. Eichel, H., Tobiska, L., Xie, H.: Supercloseness and superconvergence of stabilized low-order finite element discretizations of the Stokes problem. *Math. Comp.* **80**(274), 697–722 (2011). DOI 10.1090/S0025-5718-2010-02404-4. URL <https://doi.org/10.1090/S0025-5718-2010-02404-4>
33. Ewing, R.E., Liu, M.M., Wang, J.: Superconvergence of mixed finite element approximations over quadrilaterals. *SIAM J. Numer. Anal.* **36**(3), 772–787 (1999)
34. Falk, R.S., Neilan, M.: Stokes complexes and the construction of stable finite elements with pointwise mass conservation. *SIAM J. Numer. Anal.* **51**(2), 1308–1326 (2013). DOI 10.1137/120888132. URL <https://doi.org/10.1137/120888132>
35. Fu, G., Jin, Y., Qiu, W.: Parameter-free superconvergent $H(\text{div})$ -conforming HDG methods for the Brinkman equations. *IMA J. Numer. Anal.* **39**(2), 957–982 (2019). DOI 10.1093/imanum/dry001. URL <https://doi.org/10.1093/imanum/dry001>
36. Gatica, G.N., Gatica, L.F., Márquez, A.: Analysis of a pseudostress-based mixed finite element method for the Brinkman model of porous media flow. *Numer. Math.* **126**(4), 635–677 (2014). DOI 10.1007/s00211-013-0577-x. URL <https://doi.org/10.1007/s00211-013-0577-x>
37. Girault, V., Raviart, P.A.: Finite element methods for Navier-Stokes equations, *Springer Series in Computational Mathematics*, vol. 5. Springer-Verlag, Berlin (1986). DOI 10.1007/978-3-642-61623-5. Theory and algorithms
38. Guzmán, J., Neilan, M.: Conforming and divergence-free Stokes elements on general triangular meshes. *Math. Comp.* **83**(285), 15–36 (2014). DOI 10.1090/S0025-5718-2013-02753-6. URL <https://doi.org/10.1090/S0025-5718-2013-02753-6>
39. Hu, J., Ma, L., Ma, R.: Optimal Superconvergence Analysis for the Crouzeix-Raviart and the Morley elements. arXiv e-prints arXiv:1808.09810 (2018)
40. Hu, J., Yu, G.: A unified analysis of quasi-optimal convergence for adaptive mixed finite element methods. *SIAM J. Numer. Anal.* **56**(1), 296–316 (2018)
41. Kundu, P.K., Dowling, D.R., Tryggvason, G., Cohen, I.M.: Fluid mechanics, 6th edn. Academic Press (2015)
42. Li, Y.: Quasi-optimal adaptive mixed finite element methods for controlling natural norm errors. arXiv e-prints arXiv:1907.03852 (2019)
43. Li, Y.W.: Global superconvergence of the lowest-order mixed finite element on mildly structured meshes. *SIAM J. Numer. Anal.* **56**(2), 792–815 (2018). DOI 10.1137/17M112587X. URL <https://doi.org/10.1137/17M112587X>
44. Liu, H., Yan, N.: Superconvergence analysis of the nonconforming quadrilateral linear-constant scheme for Stokes equations. *Adv. Comput. Math.* **29**(4), 375–392 (2008). DOI 10.1007/s10444-007-9054-3. URL <https://doi.org/10.1007/s10444-007-9054-3>
45. Liu, X., Li, J., Chen, Z.: A weak Galerkin finite element method for the Oseen equations. *Adv. Comput. Math.* **42**(6), 1473–1490 (2016). DOI 10.1007/s10444-016-9471-2. URL <https://doi.org/10.1007/s10444-016-9471-2>
46. Lovadina, C., Stenberg, R.: Energy norm a posteriori error estimates for mixed finite element methods. *Math. Comp.* **75**(256), 1659–1674 (2006). DOI 10.1090/S0025-5718-06-01872-2
47. Morin, P., Nochetto, R.H., Siebert, K.G.: Data oscillation and convergence of adaptive FEM. *SIAM J. Numer. Anal.* **38**(2), 466–488 (2000)
48. Mu, L., Wang, J., Ye, X.: A stable numerical algorithm for the Brinkman equations by weak Galerkin finite element methods. *J. Comput. Phys.* **273**, 327–342 (2014). DOI 10.1016/j.jcp.2014.04.017. URL <https://doi.org/10.1016/j.jcp.2014.04.017>
49. Pan, J.: Global superconvergence for the bilinear-constant scheme for the Stokes problem. *SIAM J. Numer. Anal.* **34**(6), 2424–2430 (1997). DOI 10.1137/S0036142995286167. URL <https://doi.org/10.1137/S0036142995286167>
50. Park, E.J., Seo, B.: An upstream pseudostress-velocity mixed formulation for the Oseen equations. *Bull. Korean Math. Soc.* **51**(1), 267–285 (2014). DOI 10.4134/BKMS.2014.51.1.267
51. Qian, Y., Wu, S., Wang, F.: A mixed discontinuous Galerkin method with symmetric stress for Brinkman problem based on the velocity-pseudostress formulation. arXiv e-prints arXiv:1907.01246 (2019)

52. Raviart, P.A., Thomas, J.M.: A mixed finite element method for 2nd order elliptic problems. In: *Mathematical aspects of finite element methods* (Proc. Conf., Consiglio Naz. delle Ricerche (C.N.R.), Rome, 1975), pp. 292–315. *Lecture Notes in Math.*, Vol. 606 (1977)
53. Stenberg, R.: Postprocessing schemes for some mixed finite elements. *RAIRO Modél. Math. Anal. Numér.* **25**(1), 151–167 (1991). DOI 10.1051/m2an/1991250101511. URL <https://doi.org/10.1051/m2an/1991250101511>
54. Wang, J., Ye, X.: Superconvergence of finite element approximations for the Stokes problem by projection methods. *SIAM J. Numer. Anal.* **39**(3), 1001–1013 (2001). DOI 10.1137/S003614290037589X. URL <https://doi.org/10.1137/S003614290037589X>
55. Wang, J., Ye, X.: A weak Galerkin finite element method for the stokes equations. *Adv. Comput. Math.* **42**(1), 155–174 (2016). DOI 10.1007/s10444-015-9415-2. URL <https://doi.org/10.1007/s10444-015-9415-2>
56. Ye, X.: Superconvergence of nonconforming finite element method for the Stokes equations. *Numer. Methods Partial Differential Equations* **18**(2), 143–154 (2002). DOI 10.1002/num.1036.abs. URL <https://doi.org/10.1002/num.1036.abs>
57. Zhang, Z., Naga, A.: A new finite element gradient recovery method: superconvergence property. *SIAM J. Sci. Comput.* **26**(4), 1192–1213 (2005). DOI 10.1137/S1064827503402837. URL <https://doi.org/10.1137/S1064827503402837>
58. Zienkiewicz, O.C., Zhu, J.Z.: A simple error estimator and adaptive procedure for practical engineering analysis. *Internat. J. Numer. Methods Engrg.* **24**(2), 337–357 (1987). DOI 10.1002/nme.1620240206. URL <https://doi.org/10.1002/nme.1620240206>
59. Zienkiewicz, O.C., Zhu, J.Z.: The superconvergent patch recovery and a posteriori error estimates. I. The recovery technique. *Internat. J. Numer. Methods Engrg.* **33**(7), 1331–1364 (1992). DOI 10.1002/nme.1620330702. URL <https://doi.org/10.1002/nme.1620330702>