

# Fastest Frozen Temperature for a Thermodynamic System

X. Y. Zhou, Z. Q. Yang, X. R. Tang, X. Wang<sup>1</sup> and Q. H. Liu<sup>1,2,\*</sup>

<sup>1</sup>*School for Theoretical Physics, School of Physics and Electronics, Hunan University, Changsha, 410082, China*

<sup>2</sup>*Synergetic Innovation Center for Quantum Effects and Applications (SICQEA),  
Hunan Normal University, Changsha 410081, China*

For a thermodynamic system obeying both the equipartition theorem in high temperature and the third law in low temperature, the curve showing relationship between the specific heat and the temperature has two common behaviors: it terminates at zero when the temperature is zero Kelvin and converges to a constant value of specific heat as temperature is higher and higher. Since it is always possible to find the characteristic temperature  $T_C$  to mark the excited temperature as the specific heat almost reaches the equipartition value, it is also reasonable to find a temperature in low temperature interval, complementary to  $T_C$ . The present study reports a possibly universal existence of the such a temperature  $\vartheta$ , defined by that at which the specific heat falls *fastest* along with decrease of the temperature. For the Debye model of solids, below the temperature  $\vartheta$  the Debye's law manifest itself.

PACS numbers:

## I. INTRODUCTION

A thermodynamic system usually obeys two laws in opposite limits of temperature; in high temperature the equipartition theorem works and in low temperature the third law of thermodynamics holds. From the shape of a curve showing relationship between the specific heat and the temperature, the behaviors in these two limits are *qualitatively* clear: in high temperature it approaches to a constant whereas it goes over to zero during the temperature is lowering to the zero Kelvin. In statistical mechanics we have usually one characteristic temperature  $T_C$ , at which the specific heat approximately takes the value given by the equipartition theorem, implying that degrees of freedom are freely excited. However, there is no definition of temperature complementary to  $T_C$  to indicate a fact only a few of degrees of freedom are thermally excited so that the third law applies.

Take the specific heat in Debye model of solids for instance, there is the Debye frequency to denote the largest possible frequency  $\omega_D$ . The characteristic temperature  $T_C$  is defined by [1–6]

$$k_B T_C \equiv \hbar \omega_D \quad (1)$$

where  $k_B$  is the Boltzmann's constant and  $\hbar$  is the Planck's constant. When temperature  $T$  reaches  $T_C$ , nearly all vibrational modes are excited so the specific heat  $c(T)$  at the  $T = T_C$  is approximately  $1k_B$  (c.f. Fig. 1)

$$\frac{c(T_C)}{c(T \rightarrow \infty)} \approx 0.952. \quad (2)$$

However, once the temperature is so low that  $T \approx T_C/6$ , the Debye's  $T^3$  law becomes apparent (c.f. Fig. 1)

$$c(T) \sim T^3, T \lesssim T_C/6. \quad (3)$$

In other words, when  $T \lesssim T_C/6$ , the equipartition theorem breaks down and the third law exhibits. It is quite obvious for the Debye model of solids, at  $T_C/6$  where the specific heat falls or rises most rapidly, c.f. Fig. 1.

The key findings of present study are twofold: First, for a thermodynamic system where the third law of thermodynamics takes effect, there is always a temperature at which the specific heat falls or rises most rapidly. Since it near zero Kelvin, we call it fastest frozen temperature (FFT) for the thermodynamic system. Secondly, the FFT can be taken as a mark of temperature at which the third law manifests itself. It must be mentioned that as a thermodynamic system is cooling down from the high temperature to zero Kelvin, there is a common phenomenon known as freeze-out which has been studied in, for instance, condensed matter physics, [1] astroparticle physics, [7–9] heavy ion collisions

---

\*Electronic address: [qianhuiliu@gmail.com](mailto:qianhuiliu@gmail.com)

[10] and statistical mechanics. [2–6] The FFT under study is a manifestation of the freeze-out from the point of view of the specific heat.

To show the possibly universal existence of the FFT, we consider three different kinds of systems. In section II, we study FFT in detail for the Debye model of solids. In order to explore the FFT in a broad class of system, we examine a model system of many independent particles of the same type in section III. The FFT is also present in a mixture of two kinds of particles, which is given in section IV. The final section V concludes this study.

In whole paper, we will use specific heat defined by

$$c = \frac{du}{dT}, \quad (4)$$

where  $u$  is mean energy over a degree of freedom, which will be precisely defined in sections II–IV for different systems, and in our approach, all other parameters remain unchanged except the temperature  $T$ . The FFT  $\vartheta$  is defined by the maximum of the  $dc/dt$  as  $T \rightarrow 0$ ,

$$\frac{dc}{dT} \preceq \frac{dc}{dT} \Big|_{T=\vartheta}. \quad (5)$$

## II. FASTEST FROZEN TEMPERATURE FOR DEBYE MODEL OF SOLIDS

Debye model of solids is an elementary model to investigate the specific heat of solids. The total internal energy  $U$  is [1–6]

$$U = U_0 + 3Nk_B T D(T_C/T) \quad (6)$$

where  $N$  is the number of the atoms in the solid, and  $U_0$  is a constant, and  $D(x)$  denotes the Debye integral,

$$D(x) \equiv \frac{3}{x^3} \int_0^x \frac{y^3}{e^y - 1} dy. \quad (7)$$

The mean energy  $u$  in a vibrational mode is given by

$$u \equiv \frac{U}{3N}. \quad (8)$$

In two opposite extremes the specific heats (4) are, respectively,

$$\lim_{T \rightarrow \infty} c = k_B, \text{ and } \lim_{T \rightarrow 0} c = 0. \quad (9)$$

The numerical calculations give both the FFT  $\vartheta = 0.164T_C \approx T_C/6$  at which  $c(\vartheta) \approx 0.257k_B \approx T_0/4$  and  $c(T_C) \approx 0.952k_B \approx k_B = \lim_{T \rightarrow \infty} c(T)$ . From these results, we see that  $\vartheta$  is really in contrast to  $T_C$  to mark the frozen temperature below which the Debye's  $T^3$  law holds. The specific heat  $c(T)$  and  $dc/dt$  against temperature are plotted in Fig. 1. To note that as the temperature rises above the FFT  $\vartheta$ , Debye's  $T^3$  law fails more and more seriously.

Remember that the shape of  $c(T)$  curve for the Debye model shares two common features of all  $c(T)$  curves for all thermodynamic systems that satisfy both the equipartition theorem at high temperature and the third law at low temperature. The FFT could then be always definable as that complementary to  $T_C$ .

## III. FASTEST FROZEN TEMPERATURE: A THEOREM AND ITS ILLUSTRATIONS

In this section, we first prove a theorem: For a system of many independent particles of the same type, in which the energy levels of a single particle are discrete, it has a FFT during the system is freezing and the FFT can be taken as the frozen temperature. Secondly, we gives two illustrations of the theorem.

### A. A theorem

Construct a model system of many independent particles of the same species, in which the energy levels of a single particle are discrete, i.e.,

$$\epsilon_n = \epsilon_0 < \epsilon_1 < \epsilon_2 < \epsilon_3 < \dots \quad (10)$$

in which the spacing between  $\epsilon_1, \epsilon_2, \epsilon_3, \dots$  is negligible but  $\epsilon_1$  is appreciably different from ground state one  $\epsilon_0$  which can be conveniently chosen to be zero,  $\epsilon_0 = 0$ , and the density of states can be  $A\epsilon^{(D-2)/2}$ , [2–6] in which  $D = 1, 2, 3, 4, 5, 6$  is the number of the spatial dimension, where coefficient  $A$  can be set to be unity. Utilize the Boltzmann statistics, the partition function  $Z_1$  is with  $\beta = (k_B T)^{-1}$  and  $w_n$  standing for the degeneracy in the  $n$ -th level,

$$\begin{aligned} Z_1 &= \sum_{n=0}^{\infty} w_n \exp(-\beta \epsilon_n) \\ &= 1 + \sum_{n=1}^{\infty} w_n \exp(-\beta \epsilon_n) \\ &\approx 1 + A \int_{\epsilon_1}^{\infty} e^{-\beta \epsilon} \epsilon^{(D-2)/2} d\epsilon \\ &= 1 + A \beta^{-\frac{D}{2}} \Gamma\left(\frac{D}{2}, \epsilon_1 \beta\right) \end{aligned} \quad (11)$$

where  $\Gamma(s, x) = \int_x^{\infty} t^{s-1} e^{-t} dt$  is the incomplete gamma function and  $\Gamma(s) = \Gamma(s, 0)$  is the ordinary gamma function. The summation in the partition function can be approximately replaced by an integral, but to get a better result, the first term or the first few terms in the summation are carried out separately and the rest is replaced by an integral and we convert  $\sum_{n=1}^{\infty} w_n \exp(-\beta \epsilon_n)$  into an integral. This problem is analytically tractable, but the relevant expressions are lengthy. The energy per particle and the specific heat  $c$  are, respectively, determined by

$$u = -\frac{\partial \ln Z_1}{\partial \beta}, c = \frac{du}{dT}. \quad (12)$$

We do not explicitly show their expressions. The computations of  $c(T)$  and  $dc(T)/dT$  for  $D = 1, 2, 3, 4, 5$ , and 6 can be easily numerically carried out, and all  $c(T)$  are similar to that given by Fig. 1. Thus, we plot  $c(T)$  and  $dc(T)/dT$  for only  $D = 3$  in Fig. 2. It is clearly that the  $c(T_C) = 1.555k_B \approx 1.5k_B = \lim_{T \rightarrow \infty} c(T)$  where the characteristic temperature

$$T_C \equiv \frac{\epsilon_1 - \epsilon_0}{k_B} = \frac{\epsilon_1}{k_B}. \quad (13)$$

The values of the FFT  $\vartheta$  and ratios of two specific heats  $c(\vartheta)/c(T_C)$  are listed in Table I.

TABLE I:  $\vartheta$  and  $c(\vartheta)/c(T_C)$  for  $D = 1, 2, 3, 4, 5, 6$

	$D = 1$	$D = 2$	$D = 3$	$D = 4$	$D = 5$	$D = 6$
$\vartheta/T_C$	0.332	0.363	0.404	0.448	0.474	0.477
$c(\vartheta)/c(T_C)$	0.312	0.318	0.339	0.380	0.421	0.464

To note that all ratios  $c(\vartheta)/c(T_C)$  are smaller than 1, therefore it is reasonable to take  $\vartheta$  for the frozen temperature, as we have done for Debye model of solids.

Here we give four comments on the theorem above.

1. Replacement of the summation (11) by an integral is an approximate treatment which is a common technique used in calculations of the partition function in statistical mechanics. If the summation can be explicitly carried out, the result can never be qualitatively changed. Two illustrations will be given in next subsection.

2. Once the energy spectrum of a system is totally continuously without bounded from upper,  $Z_1 = A\Gamma(s)\beta^{-\frac{D}{2}}$  and  $c = \text{const.}$  in whole range of temperature, which is not a physical system at low temperature for it violates the third law of thermodynamics, beyond the scope of present study.

3. Once the energy spectrum of a system is continuously in a limited range,  $\epsilon \in [0, \epsilon_{\max}]$  where the  $\epsilon_{\max}$  is the largest one, the system is similar to the Debye model of solids, and the mathematical treatment is almost the same.

4. Once the energy spectrum of a system is partially discrete in the inner-shells and continuously in the rest, the mathematical treatment of the partition function is similar to (11) and the results are similar.

## B. Two illustrations of the theorem

The theorem above applies for the vibrational degrees of freedom and rotational ones of the diatomic gas which is dilute and interaction-free. In this subsection, we deal with these two degrees of freedom, respectively.

Illustration 1: FFT of vibrational degrees of freedom for the diatomic gas.

The energy level is  $\epsilon_n = (n + 1/2)\hbar\omega$ , ( $n = 0, 1, 2, \dots$ ), where  $\omega$  is the vibrational frequency. The partition function is, [3]

$$Z_1 = \sum_{n=0}^{\infty} e^{-\beta(n+1/2)\hbar\omega} = \frac{e^{-\beta\hbar\omega/2}}{1 - e^{-\beta\hbar\omega}} \quad (14)$$

With  $T_C = \hbar\omega/k_B$ , we have the mean vibrational energy  $u$  and specific heat  $c$ , respectively,

$$u = \frac{\hbar\omega}{2} + \frac{\hbar\omega}{e^{T_C/T} - 1}, c = k_B \left( \frac{T_C}{T} \right)^2 \frac{e^{T_C/T}}{(e^{T_C/T} - 1)^2}. \quad (15)$$

The specific heat at  $T_C$  is  $c(T_C) = 0.921k_B \simeq \lim_{T \rightarrow \infty} c(T) = k_B$ . The FFT  $\vartheta = 0.223T_C$  at which  $c(\vartheta) = 0.231k_B$  from which we see that vibrational degrees of freedom are thermally depressed. The value  $T_C$  of different diatomic gases is of order  $10^3$  K, [3] and we have then  $\vartheta \sim (200 - 300)$  K so the vibrational degrees of freedom at room temperature are almost frozen. We plot the specific heat  $c(T)$  and  $dc/dt$  against temperature in Fig. 3.

Illustration 2: FFT of rotational degrees of freedom for a heteronuclear diatomic gas.

The energy level is  $\epsilon_l = l(l+1)\hbar^2/2I$ , ( $l = 0, 1, 2, \dots$ ), where  $I$  is the moment of inertia. The partition function is,

$$Z_1 = \sum_{l=0}^{\infty} (2l+1) e^{-\frac{\beta l(l+1)\hbar^2}{2I}} \quad (16)$$

Note that  $T_C = \hbar^2/(2Ik_B)$ , [2-6] which would be  $\hbar^2/(Ik_B)$  with use of (13). The mean vibrational energy  $u$  and specific heat  $c$  are, respectively, determined by

$$u = \sum_{l=0}^{\infty} \epsilon_l \rho_l, c = \sum_{l=0}^{\infty} \epsilon_l \frac{d\rho_l}{dT}, \quad (17)$$

where

$$\rho_l = \frac{(2l+1)e^{-l(l+1)T_C/T}}{\sum_{l=0}^{\infty} (2l+1)e^{-l(l+1)T_C/T}}. \quad (18)$$

The numerical calculations give both  $c(T_C) = 1.07k_B \approx \lim_{T \rightarrow \infty} c(T) = k_B$  and the FFT  $\vartheta = 0.390T_C$  at which  $c(\vartheta) = 0.452k_B$ . Once  $T$  decreases from  $\vartheta$  the rotational degrees of freedom is rapidly frozen out. The value of  $T_C$ , for example, for HCl is about 15 K, [3] we have  $\vartheta = 12$  K so the rotational degrees of freedom at room temperature are freely excited. We plot the specific heat  $c(T)$  and  $dc/dt$  against temperature in Fig. 4.

In this section, we deal with a system of many independent particles of the same type, and show existence of the FFT and it can be taken as the mark of the frozen temperature.

## IV. FASTEST FROZEN TEMPERATURE FOR A MIXTURE OF TWO KINDS OF ATOM

The theorem in section III is unable to directly apply to a mixture of different kinds of atoms, but in fact the FFT also exists in similar manner. To see it, we study a 1 : 3 mixture of para-hydrogen and ortho-hydrogen gas with consideration of vibrational degrees of freedom only.

The partition functions for ortho- and para-hydrogen are  $Z_1^{ortho}$  and  $Z_1^{para}$ , respectively [2-6]

$$Z_1^{ortho} = \sum_{l=1,3,5,\dots} (2l+1) e^{-\frac{\beta l(l+1)\hbar^2}{2I}}, Z_1^{para} = \sum_{l=0,2,4,\dots} (2l+1) e^{-\frac{\beta l(l+1)\hbar^2}{2I}}. \quad (19)$$

The mean vibrational energy  $u$  and specific heat  $c$  are, respectively, determined by

$$u = -\frac{\partial}{\partial\beta} \left( \frac{3}{4} \ln Z_1^{ortho} + \frac{1}{4} \ln Z_1^{para} \right), c = \frac{du}{dT}.$$

We still follow the convention of taking the rotational characteristic temperature  $T_C = \hbar^2 / (2Ik_B)$ . [2–6] The numerical calculations give both the FFT  $\vartheta = 1.33T_C$  at which  $c(\vartheta) = 0.305k_B$  and  $c(T_C) = 0.117k_B \ll k_B = \lim_{T \rightarrow \infty} c(T)$ . From these results, we see that  $T_C$  is not excited temperature at all. According to the theorem in section III, the characteristic temperature  $T_C$  can only be defined for a given type of particle rather than the a mixture. It is reasonable to define the characteristic temperature (13)  $3T_C = \epsilon_2 - \epsilon_0$  and  $5T_C = \epsilon_3 - \epsilon_1$  for ortho- and para-hydrogen, respectively. One can only take largest one  $5T_C$  to specify the temperature above which the equipartition theorem can apply. However, in our approach the FFT is independent of the choice of  $T_C$ . The specific heat  $c(T)$  and  $dc/dt$  against temperature are plotted in Fig. 5.

Note that at  $T = 0$  K, hydrogen contains mainly para-hydrogen which is more stable, and in general the concentration ratio of the ortho- to para-hydrogen in thermal equilibrium is given by [3]

$$r = 3 \frac{Z_1^{ortho}}{Z_1^{para}}, \quad (20)$$

from which we obtain

$$r(T_C) = 7.44, r(\vartheta) = 4.74, r(300) = 3.01. \quad (21)$$

Since at the room temperature  $T \approx 300$  K  $= 3.53T_C$  the ratio  $r \approx 3$ , and thus "the name ortho- is given to that component which carries the larger statistical weight". [3]

## V. CONCLUSIONS AND DISCUSSIONS

Temperature is a fundamental quantity in physics. It has two unattainable ends; one is the absolute zero and another is the infinitely large. If taking  $\beta = (k_B T)^{-1}$  as an alternative definition of the temperature  $T$  instead, the new temperature  $\beta$  has also two unattainable ends. In physics, a thermodynamic system usually obeys two laws in opposite limits of temperature, however, we have only one characteristic temperature, at which the value of specific heat is approximately that given by the equipartition theorem. We have no such a temperature to mark where the third law comes to play. A well-defined temperature at which the specific heat falls *fastest* along with decrease of the temperature is identified, the so-called FFT, which can be taken as the frozen temperature itself. For Debye model of solids, FFT marks the temperature below which the the Debye's  $T^3$  law comes into play. For the dilute gas composed of noninteracting atoms of the same type or not, FFT marks the temperature the system almost gets stuck in the ground state eigenstate for the rational or vibrational degrees of freedom.

We like to mention the following facts which can be easily demonstrated. There are thermodynamic systems which violate the equipartition theorem in high temperature, photon gas for instance, or violate the third law in low temperature, classical ideal gas for instance, FFT should be used with care. For the former, there is no  $T_C$  and thus no FFT as well; and for the latter, neither  $T_C$  nor FFT is applicable. What is more, for noninteracting Bosons, both  $T_C$  and FFT exist but they are identical.

Therefore the present study shows not only existence of the FFT but also its complementarity to  $T_C$ .

## Acknowledgments

This work is financially supported by National Natural Science Foundation of China under Grant No. 11675051.

- 
- [1] S. H. Simon, The Oxford Solid State Basics, (Oxford: Oxford University Press, 2013)
  - [2] M. W. Zemansky, R. H. Dittman, Heat and Thermodynamics: An Intermediate Textbook, 7th ed., (New York: McGraw-Hill, 1997)
  - [3] R. K. Pathria, P. D. Beale, Statistical Mechanics, 3rd ed., (Oxford: Butterworth-Heinemann, 1996)
  - [4] R. Kubo, Statistical Mechanics, (New York: Wiley, 1965)
  - [5] K. Huang, Statistical Mechanics, 2nd ed., (New York: Wiley, 1986)
  - [6] L. D. Landau and E. M. Lifshitz, Statistical Physics (I), 3rd ed., (Oxford: Pergamon, 1980)
  - [7] Ya. B. Zeldovich, L. B. Okun, S. B. Pikelner, Sov. Phys. Usp., **8** (1966)702
  - [8] B. W. Lee, S. Weinberg, Phys. Rev. Lett., **39** (1977)165
  - [9] M. Cannoni, Eur. Phys. J. C., **75** (2015)106
  - [10] S. V. Akkelin, Y. Hama, Iu. A. Karpenko, and Yu. M. Sinyukov, Phys. Rev. C., **78** (2008)034906




FIG. 1: Curves for specific heat (solid) and its derivative with respect to temperature (dashed) for the Debye model of solid, and the auxiliary dotted lines are guide for the eye. The equipartition value of the specific heat is  $1k_B$ , and  $c(T_C) = 0.952k_B \approx 1k_B$ . The  $dc/dT$  has a global maximum at  $\vartheta = 0.164T_C$  at which the specific heat  $c(\vartheta) = 0.257k_B \approx 1/4k_B$ . The Debye's  $T^3$  law is plotted in circles, showing that it holds true in temperature interval  $T_C \in (0, \vartheta)$ .




FIG. 2: Curves for specific heat (solid) and its derivative with respect to temperature (dashed) for our model system, and the auxiliary dotted lines are guide for the eye. Note that the equipartition value of the specific heat is  $3k_B/2$ , compatible with the number of the dimension  $D = 3$ . The  $dc/dT$  has a global maximum at  $\vartheta = 0.404T_C$  at which the specific heat  $c(\vartheta) = 0.527k_B = 0.339c(T_C)$ .  $c(T_C) = 1.56k_B \approx 1.5k_B$ , the equipartition value.




FIG. 3: Curves for specific heat (solid) and its derivative with respect to temperature (dashed) for the vibrational degrees of freedom of a diatomic gas, and the auxiliary dotted lines are guide for the eye. Note that the equipartition value of the specific heat is  $1k_B$ , and  $c(T_C) = 0.921k_B$  is slightly smaller than  $1k_B$ . The  $dc/dT$  has a global maximum at  $\vartheta = 0.223T_C$  at which the specific heat  $c(\vartheta) = 0.231k_B \approx 1/4k_B$ .




FIG. 4: Curves for specific heat (solid) and its derivative with respect to temperature (dashed) for the rotational degrees of freedom of a heteronuclear diatomic gas, and the auxiliary dotted lines are guide for the eye. Note that the equipartition value of the specific heat is  $1k_B$ , and  $c(T_C) = 1.07k_B \approx 1k_B$ . The  $dc/dT$  has a global maximum at  $\vartheta = 0.390T_C$  at which the specific heat  $c(\vartheta) = 0.452k_B = 0.421c(T_C)$ .




FIG. 5: Curves for specific heat (solid) and its derivative with respect to temperature (dashed) for the rotational degrees of freedom of 1 : 3 mixture of para-hydrogen and ortho-hydrogen molecules, and the auxiliary dotted lines and arrows are guide for the eye. The equipartition value of the specific heat is  $1k_B$ , and  $c(T_C) = 0.117k_B$  is much smaller than  $1k_B$ . The  $dc/dT$  has a global maximum at  $\vartheta = 1.33T_C$  at which the specific heat  $c(\vartheta) = 0.305k_B$ . The conventional definition of characteristic temperature  $T_C = \hbar^2/(2Ik_B)$  is nothing but a unit, and once we use our definition (13), we have for ortho- and para-hydrogen, respectively,  $c(\vartheta_{ortho}) = 0.907k_B \approx 1k_B$  and  $c(\vartheta_{para}) = 0.999k_B \approx 1k_B$ .