


## ON AN INFINITE ORDER CORK AUTOMORPHISMS

SELMAN AKBULUT

ABSTRACT. Here we give a concrete description of the cork automorphism  $f : \partial W \rightarrow \partial W$  of the infinite order loose-cork  $(W, f)$ , defined in [A2]. It is obtained by concatenating the defining ribbon disk of  $W$  in  $B^4$  by an infinite order isotopy of the boundary knot.

## 0. INTRODUCTION

Recall the infinite order loose-cork  $(W, f)$  defined in [A2] and [G]. One definition of  $W$  is that, it is the contractible manifold obtained by blowing down  $B^4$  along the ribbon disk  $D \subset B^4$  bounding the knot  $K\# -K$ , where  $K$  is the figure 8 knot as in Figure 1 (cf. 6.2 of [A1])

FIGURE 1.  $W$ 

What follows from the discussion of [A3] is that, the infinite order cork twisting map  $f_n : \partial W \rightarrow \partial W$  of this cork is induced from the diffeomorphism  $\partial W \rightarrow \partial W_n$ , where  $W_n \approx W$  is the contractible manifold obtained by blowing down  $B^4$  along the canonical disk  $D_n \subset B^4$ , which is obtained by concatenating  $D$  with the concordance of  $K\# -K$  along the collar  $S^3 \times [0, 1]$  of the boundary, induced by order  $n$  “swallow follow” isotopy as shown in Figure 3. This isotopy extends to an ambient isotopy  $F_t : B^3 \times [0, 1] \rightarrow B^3 \times [0, 1]$  which is fixed on  $B^3 \times 0 \cup S^2 \times [0, 1]$ .



**Remark 1.** *If the isotopy  $F_t$  was fixed  $B^3 \times 1$ , then we would get a nontrivial element of  $\pi_0 \text{Diff}(B^4, S^3) = \pi_0 \text{Diff}(S^4)$ , otherwise  $(W, f_n)$  would not be loose-cork. We would like to thank Rob Kirby noticing that this condition was overlooked in the first posting of this paper. Recall that in [W] it was shown  $\pi_k \text{Diff}(S^4) \neq \pi_k SO(5)$  for some  $k > 1$ .*

---

*Date:* December 9, 2021.

1991 *Mathematics Subject Classification.* 58D27, 58A05, 57R65.

Partially supported by NSF grants DMS 0905917.

FIGURE 2.  $f : W \rightarrow W_n$ FIGURE 3.  $W_n$ , for  $n = 2$ 

## REFERENCES

- [A1] S. Akbulut, *4-Manifolds*, Oxford Univ Press. ISBN-13: 978-0198784869
- [A2] S. Akbulut, *On infinite order corks*, PGGT, IP Paperback (2017) 151-157, ISBN: 9781571463401, <https://arxiv.org/pdf/1605.09348.pdf>
- [A3] S. Akbulut, *Homotopy 4-spheres associated to an infinite order loose cork*, <https://arxiv.org/pdf/1901.08299.pdf>
- [G] R. E. Gompf, *Infinite order corks*, G&T, vol.21, no.4 (2017) 2475-2484.
- [W] T. Watanabe, *Some exotic nontrivial elements of the rational homotopy groups of  $\text{Diff}(S^4)$* , <https://arxiv.org/pdf/1812.02448.pdf>

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, MI, 48824  
*E-mail address:* akbulut@msu.edu