

Multi-marginal optimal transportation problem for cyclic costs*

BRENDAN PASS [†] ADOLFO VARGAS-JIMÉNEZ[‡]

November 22, 2021

Abstract

We study a multi-marginal optimal transportation problem with a cost function of the form $c(x_1, \dots, x_m) = \sum_{k=1}^{m-1} |x_k - x_{k+1}|^2 + |x_m - F(x_1)|^2$, where $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$. When $m = 4$, F is a positive multiple of the identity mapping, and the first and last marginals are absolutely continuous with respect to Lebesgue measure, we establish that any solution of the Kantorovich problem is induced by a map; the solution is therefore unique. We go on to show that this result is sharp in a certain sense. Precisely, we exhibit examples showing that Kantorovich solutions may concentrate on a higher dimensional sets if any of the following hold: 1) F is any linear mapping other than a positive scalar multiple of the identity, 2) the last marginal is not absolutely continuous with respect to Lebesgue measure, or 3) the number of marginals $m \geq 5$, even when F is the identity mapping.

1 INTRODUCTION

Let $\mu_k \in P(X_k)$ be Borel probability measures, where $X_k \subseteq \mathbb{R}^n$ are bounded domains and $k = 1, \dots, m$. For a real-valued cost function c on the product

*BP is pleased to acknowledge support from Natural Sciences and Engineering Research Council of Canada Grant 04658-2018. The work of AVJ was completed in partial fulfillment of the requirements for a doctoral degree in mathematics at the University of Alberta.

[†]Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, Alberta, Canada pass@ualberta.ca.

[‡]Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, Alberta, Canada vargasji@ualberta.ca.

space $X := X_1 \times X_2 \times \dots \times X_m$, the multi-marginal version of Kantorovich's optimal transportation problem is to solve:

$$\inf \left\{ \mathcal{F}_c[\mu] : \mu \in \Pi(\mu_1, \dots, \mu_m) \right\} \quad (\text{KP})$$

where

$$\Pi(\mu_1, \dots, \mu_m) = \left\{ \mu \in P(X) : \mu(X_1 \times \dots \times A_k \times \dots \times X_m) = \mu_k(A_k) \right. \\ \left. \text{for all measurable sets } A_k \subseteq X_k, \quad 1 \leq k \leq m \right\}$$

$$\text{and } \mathcal{F}_c[\mu] = \int_X c(x_1, \dots, x_m) d\mu.$$

The Monge formulation of the same problem is to solve:

$$\inf \left\{ \mathcal{G}[(T_1, T_2, \dots, T_m)] : \{T_k\}_{k=1}^m \in \Gamma(\mu_1, \dots, \mu_m) \right\} \quad (\text{MP})$$

where

$$\Gamma(\mu_1, \dots, \mu_m) = \left\{ \{T_k\}_{k=1}^m : T_k : X_1 \mapsto X_k \text{ is measurable, } (T_k)_\# \mu_1 = \mu_k \text{ and } T_1 = Id \right\}$$

$$\text{and } \mathcal{G}[(T_1, T_2, \dots, T_m)] = \int_{X_1} c(x_1, T_2 x_1, \dots, T_m x_1) d\mu_1.$$

As usual, Id denotes the identity map and $(T_k)_\# \mu_1$ the image measure of μ_1 through T_k given by $(T_k)_\# \mu_1(A_k) = \mu_1(T_k^{-1}(A_k))$, for every measurable subset A_k of X_k . For any $\{T_k\}_{k=1}^m \in \Gamma(\mu_1, \dots, \mu_m)$, we can define the measure $\mu = (T_1, T_2, \dots, T_m)_\# \mu_1 \in \Pi(\mu_1, \dots, \mu_m)$. Then $\mathcal{F}_c[\mu] = \mathcal{G}[(T_1, T_2, \dots, T_m)]$; thus, (KP) can be described as a relaxed version of (MP). When $m = 2$, these reduce respectively to the Kantorovich and Monge formulations of the classical optimal transport problem, which has profound connections with many different areas of mathematics, and an extremely wide range of applications in other fields, surveyed in, for example, [27][28] and [29]. The multi-marginal generalization (ie, the case $m \geq 3$) has attracted a great deal of attention recently, largely due to its own substantial collection of applications, in, for instance, physics, economics, finance and image processing (see [25] for an overview).

Under very general conditions (for instance, compactness of the spaces and continuity of the cost is more than enough) there exists a solution for

(KP). In the two marginal case, a simple condition on c , known as the *twist* condition, as well as absolute continuity of the first marginal μ_1 , is well known to ensure existence of Monge solutions, and, consequently, uniqueness in both problems [16][17].

The situation is much more subtle for $m \geq 3$. Beginning with a seminal work by Gangbo and Świąch on the cost $c(x_1, \dots, x_m) = \sum_{k \neq i}^m |x_k - x_i|^2$ [15], many authors have proven that several particular cost functions yield Monge solutions [18][8][26], while a variety of others do not [10][23][24][12]. An important challenge in current research is to fully identify and classify the costs which ensure unique Monge solutions.

A sufficient condition for Monge solutions (known as the *twist on splitting sets*), which encompasses all known examples of costs leading to Monge solutions, is provided in [20]. However, this condition is much more complicated and difficult to check for a given cost than its two marginal counterpart. Second order differential conditions on c which guarantee the twist on splitting sets are known, but they are very strong; of particular relevance to us, they require invertability of certain matrices of second derivatives which does not hold in some situations of interest [22][26].

In this paper, we study the cost

$$c(x_1, \dots, x_m) = \sum_{i=1}^{m-1} |x_i - x_{i+1}|^2 + |x_m - F(x_1)|^2, \quad (1)$$

for a fixed map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$. As highlighted in Section 1.7.4 of [27], this cost measures the discrete time kinetic energy of a cloud of particles whose density at timestep k is μ_k , such that the final position of the particle initially at x_1 is fixed to be $F(x_1)$. In particular, when each $\mu_k = \mathcal{L}^n|_D$ is (normalized) Lebesgue measure on a common bounded domain $X_k = D \subset \mathbb{R}^n$ and $F : D \rightarrow D$ is measure preserving, $F_{\#}\mu_k = \mu_k$, the Monge problem with this cost corresponds to the time discretization of Arnold's variational interpretation of the incompressible Euler equation [1]; the Kantorovich formulation corresponds to a discretization of Brenier's generalization [7]. If $m = 2$ and $I + DF(x)$ is invertible (alternatively it corresponds to the quadratic cost up to a change of variables) where I denotes the identity matrix, the cost is twisted; while for $m = 3$, it is twisted on splitting sets as long as $DF(x) + DF(x)^T > 0$. To the best of our knowledge, little is known analytically about the structure of solutions for $m \geq 4$, though the problem has received a fair bit of attention from a numerical perspective [14][21][4][6].

As we show, cost function (1) is not twisted on splitting sets for $m \geq 4$. Nevertheless, when $m = 4$ and F is a positive scalar multiple of the identity mapping, we are able to prove that all solutions are of Monge type, and therefore unique, under an additional regularity condition on the marginals (in addition to μ_1 , either μ_2 or μ_4 must be absolutely continuous). This result is very special; indeed, as we show later on, it is in some sense impossible to go further. A simple example shows that the extra regularity condition on μ_4 or μ_2 is required. When $m = 4$, and F is any linear mapping other than a positive scalar multiple of the identity, we demonstrate that solutions may not be of Monge type, even for diffuse marginals. Similarly, when $m \geq 5$, we prove that solutions may not be of Monge type, even for $F(x) = x$.

To offer some perspective on these results, we note that generalized incompressible flows (ie, solutions to the infinite marginal version of the Kantorovich problem, when each marginal is uniform and F measure preserving) are not generally unique in dimension $n \geq 2$ [5]; however, unique Monge-type solutions exist when F is close to the identity mapping [13]. It seems reasonable to expect the same to hold for the time discretized problem. Our counterexamples essentially show that this is not the case for $m \geq 4$, at least when the marginals are allowed to differ.

In the next section, we recall some preliminary results and definitions which will be needed in the paper. In Section 3, we prove that solutions are of Monge type when the number of marginals $m = 4$, $F(x) = x$ and the first and last marginal are absolutely continuous. The fourth section is reserved for examples of non-Monge solutions when these conditions are violated.

2 DEFINITIONS AND PRELIMINARIES

For the cost function (1) on $X = X_1 \times X_2 \times \dots \times X_m$, we will approach the problem of minimizing $\mathcal{F}_c[\mu] = \int_X c(x_1, \dots, x_m) d\mu$, by the equivalent problem of maximizing:

$$\mathcal{F}_b[\mu] = \int_X b(x_1, \dots, x_m) d\mu \tag{KPb}$$

where $b(x_1, \dots, x_m) = \sum_{k=1}^{m-1} x_k \cdot x_{k+1} + x_m \cdot F(x_1)$, over $\mu \in \Pi(\mu_1, \dots, \mu_m)$.

The dual of (KPb) is to minimize:

$$\sum_{k=1}^m \int_{X_k} u_k(x_k) d\mu_k \quad (\text{DP})$$

among all m -tuples (u_1, u_2, \dots, u_m) where each $u_k \in L^1(\mu_k)$ and $\sum_{k=1}^m u_k \geq b(x_1, \dots, x_m)$, for all $(x_1, \dots, x_m) \in X$.

We now introduce an important class of functions satisfying the constraint in (DP).

Definition 2.1. *An m -tuple of functions (u_1, u_2, \dots, u_m) is b -conjugate if for all k*

$$u_k(x_k) = \sup_{x_i \in X_i, i \neq k} \left(b(x_1, \dots, x_m) - \sum_{i \neq k} u_i(x_i) \right)$$

It is well known that if (u_1, u_2, \dots, u_m) is b -conjugate, then each u_k in inherits local Lipschitz and semi-convexity properties from b [11].

The following well known duality result captures the connection between (DP) and (KPb). Most of the assertions can be traced back to Kellerer [19]; a proof of the b -conjugacy of the solutions can be found in [15] or [26].

Theorem 2.1. *Assume X_k is compact for every k . Then, there exists a solution μ to the Kantorovich problem and a b -conjugate solution (u_1, u_2, \dots, u_m) to its dual. The maximum and minimum values in (DP) and (KPb) respectively are the same and $\sum_{k=1}^m u_k(x_k) = b(x_1, \dots, x_m)$ for all $(x_1, \dots, x_m) \in \text{spt}(\mu)$, where $\text{spt}(\mu)$ denotes the support of μ .*

We will use the next theorem, established in [2][3], in the construction of counterexamples. For this purpose, we denote by \mathbb{R}^d the set of all 2×2 real matrices that can be expressed as the product of d positive definite real matrices.

Theorem 2.2. *Assume that $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is a 2×2 matrix and $|M| > 0$, where $|M|$ denotes the determinant of M , then:*

1. $M \in \mathbb{R}^2$ iff M is diagonalizable and its eigenvalues are both positive.
2. $M \in \mathbb{R}^3$ iff $\text{tr}(M) > 0$ or $(c - b)^2 > 4|M|$.

We close this section by recalling a couple of well known formulas which will be useful in the construction of counterexamples in Section 4.

For any 2×2 matrices A and B we have:

$$|A + B| = |A| + |B| + \text{trace}(Adj(A)B) \quad (2)$$

where $Adj(A)$ denote the adjugate of A .

Given a convex function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$, its Legendre-transform will be denoted by f^* ; that is, $f^*(y) = \sup_x (x \cdot y - f(x))$. We have special interest in the Legendre-transform of $f(x) = \frac{1}{2}x^T Ax + b \cdot x$ for a given positive definite $n \times n$ matrix A and $b \in \mathbb{R}^n$. For this function, we have:

$$f^*(y) = \frac{1}{2}(y - b)^T A^{-1}(y - b). \quad (3)$$

3 MONGE SOLUTIONS

We now show that under regularity conditions on the first and fourth marginal, we obtain a unique Monge solution for the case $m = 4$ and $F(x) = x$.

In what follows \mathcal{L}^n denotes the Lebesgue measure on \mathbb{R}^n .

Theorem 3.1. *Let μ_k be probability measures over open bounded sets $X_k \subseteq \mathbb{R}^n$, $k = 1, 2, 3, 4$. Take $b(x, y, z, w) = x \cdot y + y \cdot z + z \cdot w + w \cdot x$ and assume μ_1, μ_4 are absolutely continuous with respect to \mathcal{L}^n . Then any solution of the Kantorovich problem (KPb) is induced by a map.*

Proof. Let μ be a solution to (KPb) and (u_1, u_2, u_3, u_4) a b -conjugate solution to its dual. Consider the set

$$S = \left\{ (x, y, z, w) : Du_1(x) \text{ and } Du_4(w) \text{ exist and } b(x, y, z, w) = u_1(x) + u_2(y) + u_3(z) + u_4(w) \right\}.$$

Since the functions $u_1(x)$ and $u_4(w)$ are Lipschitz, they are differentiable \mathcal{L}^n -a.e., and therefore μ_1 and μ_4 a.e. by hypothesis. Hence, $\mu(S) = 1$. By setting $f(x, y) = x \cdot y + \sup_z [y \cdot z - u_3(z) + h(x + z)]$ where $h(x + z) = \sup_w [(x + z) \cdot w - u_4(w)]$, we obtain:

$$b(x, y, z, w) - u_3(z) - u_4(w) \leq f(x, y) \leq u_1(x) + u_2(y)$$

for all x, y, z, w and in particular equality holds on S .

Now, for any fixed x_0 , we will show that there is only one y_0, z_0, w_0 such that in $(x_0, y_0, z_0, w_0) \in S$. Since the function $x \mapsto f(x, y_0)$ is convex and $f(x, y_0) \leq u_1(x) + u_2(y_0)$ for every x , it is subdifferentiable everywhere. For

$(x_0, y_0, z_0, w_0) \in S$ the equality $f(x_0, y_0) = u_1(x_0) + u_2(y_0)$ implies that the subdifferential of $f(x, y_0)$ at x_0 is contained in the subdifferential of $u_1(x)$ at x_0 , which is $\{Du_1(x_0)\}$; that is, $D_x f(x_0, y_0)$ exists and equals $Du_1(x_0)$. By a similar argument $Dh(x_0 + z_0)$ exists, $Dh(x_0 + z_0) = D_x f(x_0, y_0) - y_0 = w_0$, and clearly, $z_0 \in \operatorname{argmax}[y_0 \cdot z - u_3(z) + h(x_0 + z)]$. We claim that the map $(y, z, w) \mapsto D_x f(x_0, y)$ with domain $R := \{(y, z, w) : (x_0, y, z, w) \in S\}$ is injective; this will imply the desired result.

Assume $D_x f(x_0, y_1) = D_x f(x_0, y_2)$ for some $(y_1, z_1, w_1), (y_2, z_2, w_2) \in R$. Note that

$$y_1 + w_1 = y_1 + Dh(x_0 + z_1) = D_x f(x_0, y_1) = D_x f(x_0, y_2) = y_2 + Dh(x_0 + z_2) = y_2 + w_2 \quad (4)$$

and $z_k \in \operatorname{argmax}[y_k \cdot z - u_3(z) + h(x_0 + z)]$, $k = 1, 2$. Therefore

$$y_1 \cdot z_2 - u_3(z_2) + h(x_0 + z_2) \leq y_1 \cdot z_1 - u_3(z_1) + h(x_0 + z_1) \quad (5)$$

$$y_2 \cdot z_1 - u_3(z_1) + h(x_0 + z_1) \leq y_2 \cdot z_2 - u_3(z_2) + h(x_0 + z_2); \quad (6)$$

adding these inequalities gives $(y_1 - y_2) \cdot (z_2 - z_1) \leq 0$, then by (4),

$$(w_2 - w_1) \cdot (z_2 - z_1) \leq 0. \quad (7)$$

Furthermore, since $w_k \in \operatorname{argmax}[(x_0 + z_k) \cdot w - u_4(w)]$,

$$(x_0 + z_1) \cdot w_2 - u_4(w_2) \leq h(x_0 + z_1) = (x_0 + z_1) \cdot w_1 - u_4(w_1) \quad (8)$$

$$(x_0 + z_2) \cdot w_1 - u_4(w_1) \leq h(x_0 + z_2) = (x_0 + z_2) \cdot w_2 - u_4(w_2); \quad (9)$$

after adding and canceling similar terms we obtain

$$(w_2 - w_1) \cdot (z_1 - z_2) \leq 0. \quad (10)$$

Therefore, by (7) and (10), $(w_2 - w_1) \cdot (z_1 - z_2) = 0$ and we must have equality in (5), (6), (8) and (9). This implies that $w_2 \in \operatorname{argmax}[(x_0 + z_2) \cdot w - u_4(w)] \cap \operatorname{argmax}[(x_0 + z_1) \cdot w - u_4(w)]$; additionally, $(y_2, z_2, w_2) \in R$ implies $u_4(w)$ is differentiable at w_2 , and so

$$x_0 + z_1 = Du_4(w_2) = x_0 + z_2; \quad (11)$$

that is, $z_1 = z_2$. The equality $w_k = Dh(x_0 + z_k)$ for $k = 1, 2$ then implies that $w_1 = w_2$, and so $y_1 = y_2$ by (4).

In summary, the equation $D_x f(x_0, y_0) = Du_1(x_0)$, which holds on S and therefore μ almost everywhere, implies that (y_0, z_0, w_0) is uniquely defined from x_0 ; therefore, the 3-tuple (T_2, T_3, T_4) where T_k is the map associating each x_0 to y_0 , z_0 and w_0 respectively, induces μ . \square

Remark 3.1. *In the proof above, T_2 solves a sort of effective, two marginal optimal transportation problem with surplus f and marginals μ_1 and μ_2 . Our argument essentially verifies that f is twisted; that is, $y \mapsto D_x f(x_0, y)$ is injective (this condition is well known to ensure Monge solution for two marginal problems [27]).*

Remark 3.2. *In a similar way, we can prove Theorem 3.1 if we replace $F = I$ by $F = \lambda I$, where $\lambda > 0$ is a scalar.*

A standard argument now implies uniqueness of solutions to (KPb).

Corollary 3.1. *Assume the same conditions as Theorem 3.1. Then the solution to the Kantorovich problem (KPb) is unique.*

Proof. Let μ^1 and μ^2 be distinct solutions of (KPb). By Theorem 3.1, $\mu^1 = (Id, T_2^1, T_3^1, T_4^1)$ and $\mu^2 = (Id, T_2^2, T_3^2, T_4^2)$ for some 3-tuples of measurable maps $(T_2^1, T_3^1, T_4^1) \neq (T_2^2, T_3^2, T_4^2)$. Since the set of solutions of (KPb) is convex, $\mu = \frac{1}{2}\mu^1 + \frac{1}{2}\mu^2$ is also a solution. Hence, applying one more time Theorem 3.1, we conclude that μ is concentrated on a graph. This is clearly not possible, completing the proof. \square

4 NON-MONGE SOLUTIONS.

We now illustrate why the conditions on the marginal μ_4 , the number of variables m and the map F in the definition of b and c of Theorem 3.1 are necessary.

4.1 The regularity condition on μ_4 .

Assuming m and F as in Theorem 3.1 the next example will show that if μ_4 is not absolutely continuous, we can find a solution for (KP) not induced by a map. Furthermore, the uniqueness result of Corollary 3.1 fails.

Example 4.1.1. *Let $X_k = B(0, r) \subseteq \mathbb{R}^n$ be an open ball, $r > 0$. Consider $c(x, y, z, w) = \frac{1}{2}(|x-y|^2 + |y-z|^2 + |z-w|^2 + |w-x|^2)$ and the following measures on X_k : The Dirac measure at the origin $\mu_2 = \mu_4 = \delta_0$ and the renormalized n -dimensional Lebesgue measure $\mu_1 = \mu_3 = \frac{\mathcal{L}^n}{k_n r^n}$, where k_n is the*

volume of the n -dimensional ball of radius 1. Take any μ in $\Pi(\mu_1, \mu_2, \mu_3, \mu_4)$. Since $(x, y, z, w) \in \text{spt}(\mu)$ implies $y = w = 0$, we obtain:

$$\begin{aligned} \int_{X_1 \times X_2 \times X_3 \times X_4} c(x, y, z, w) d\mu &= \int_{X_1 \times X_2 \times X_3 \times X_4} (|x|^2 + |z|^2) d\mu \\ &= \int_{B(0,r)} |x|^2 d\mu_1(x) + \int_{B(0,r)} |z|^2 d\mu_3(z); \end{aligned}$$

that is, $\mathcal{F}[\mu]$ is independent of μ , hence any element in $\Pi(\mu_1, \mu_2, \mu_3, \mu_4)$ is a minimizer. Therefore we can find optimal measures μ for (KP) not concentrated on a graph of a measurable map; for instance, the product measure $\mu = \mu_1 \otimes \mu_2 \otimes \mu_3 \otimes \mu_4$. On the other hand, if we set $\mu = (Id, F, T, F)_\# \mu_1$ where $T_\# \mu_1 = \mu_3$ and $F = 0$, we get solutions for the Monge problem.

4.2 The condition $F = I$.

In this subsection, by assuming F is not a positive multiple of the identity mapping, $m = 4$ and $n = 2$, we will find absolutely continuous marginals in \mathbb{R}^2 such that a solution of (KPb) is concentrated in a 3-dimensional set. Therefore, this solution will not be induced by a map. For this purpose, the following lemma will be needed:

Lemma 4.2.1. *For each 2×2 real matrix F such that $F \neq \lambda I$ for some $\lambda > 0$, there exists $M \in \mathfrak{R}^2$ such that $F + M$ is singular.*

Proof. Let $F = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be 2×2 real matrix such that $F \neq \lambda I$ for any $\lambda > 0$.

We want to show that $|F + M| = 0$ for some $M = \begin{pmatrix} e & f \\ g & h \end{pmatrix} \in \mathfrak{R}^2$. First, note that by (2)

$$|F + M| = |F| + |M| + (de - gb) + (ah - fc).$$

We divide the proof into 3 cases:

1. If $c \neq 0$, take any $e, h \in \mathbb{R}$ with $e \neq h$ and $e, h > 0$. By setting $f = \frac{|F| + eh + de + ah}{c}$ and $g = 0$ we obtain $|M| > 0$ and $|F + M| = 0$. Furthermore, M is triangular with distinct eigenvalues e and h , hence M is diagonalizable. Since $e, h > 0$, we get $M \in \mathfrak{R}^2$ by Theorem 2.2.

2. If $c = 0$ and $b \neq 0$, using a similar argument as in 1. we obtain the same result by taking $f = 0$, $g = \frac{|F|+eh+de+ah}{b}$ and any $e, h > 0$, $e \neq h$.
3. If $b = c = 0$, note that by hypothesis $a \neq d$. Also, we have $a, d \geq 0$, $a, d < 0$ or without loss of generality $a < 0$ and $d \geq 0$. For the second and third case, we can make M diagonalizable with positive determinant and satisfying $|F + M| = 0$, by taking $e = h = -a$ and $f = g = 0$. Hence, $M \in \mathfrak{R}^2$ by Theorem 2.2. For the first case assume without loss of generality $a > d \geq 0$ and consider the matrix:

$$M = \begin{pmatrix} e & f \\ g & h \end{pmatrix} = \begin{pmatrix} \frac{ad}{a-d} + \lambda & \frac{d^2}{a-d} + \frac{a+d}{2a}\lambda \\ -\frac{a^2}{a-d} - \lambda & -\frac{ad}{a-d} - \frac{a+d}{2a}\lambda \end{pmatrix}$$

with $\lambda > 0$. Clearly,

$$|F + M| = \begin{vmatrix} \frac{a^2}{a-d} + \lambda & \frac{d^2}{a-d} + \frac{a+d}{2a}\lambda \\ -\frac{a^2}{a-d} - \lambda & -\frac{ad}{a-d} - \frac{a+d}{2a}\lambda \end{vmatrix} = 0.$$

Since $\frac{d^2}{a-d} + \frac{a+d}{2a}\lambda = -d + \frac{ad}{a-d} + \frac{a+d}{2a}\lambda$ and $\frac{ad}{a-d} + \lambda = -a + \frac{a^2}{a-d} + \lambda$ we have:

$$\begin{aligned} |M| &= -\left(-a + \frac{a^2}{a-d} + \lambda\right)\left(\frac{ad}{a-d} + \frac{a+d}{2a}\lambda\right) + \left(\frac{a^2}{a-d} + \lambda\right)\left(-d + \frac{ad}{a-d} + \frac{a+d}{2a}\lambda\right) \\ &= a\left(\frac{ad}{a-d} + \frac{a+d}{2a}\lambda\right) - d\left(\frac{a^2}{a-d} + \lambda\right) \\ &= \left(\frac{a+d}{2} - d\right)\lambda \\ &= \left(\frac{a-d}{2}\right)\lambda > 0. \end{aligned}$$

Furthermore, $tr(M) = e + h = \frac{a-d}{2a}\lambda > 0$. Hence, $tr(M)^2 - 4|M| > 0$ for big enough λ ; that is, the eigenvalues of M given by

$$\frac{tr(M) \pm \sqrt{tr(M)^2 - 4|M|}}{2}$$

are both positive and different. Then M is diagonalizable and belongs to \mathfrak{R}^2 , by Theorem 2.2.

□

Proposition 4.2.1. *For $b(x, y, z, w) = x \cdot y + y \cdot z + z \cdot w + w \cdot F(x)$, with $(x, y, z, w) \in (\mathbb{R}^2)^4$ and F a linear map such that $F \neq \lambda I$ for any $\lambda > 0$, there are absolutely continuous marginals $\mu_1, \mu_2, \mu_3, \mu_4$ with respect to \mathcal{L}^2 , such that a solution of (KPb) is not concentrated on a graph of a measurable map.*

Proof. Let $F = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be the matrix representation of $F(x)$. By the previous lemma we can choose $M = \begin{pmatrix} e & f \\ g & h \end{pmatrix} \in \mathfrak{R}^2$ such that $F + M$ is singular. Note that $A := M^{-1}F + I$ is also singular and $M^{-1} \in \mathfrak{R}^2$. Let $M_1, M_2 > 0$ be such that $M^{-1} = M_1 M_2$ and $v \in \mathbb{R}^2$ a nonzero vector satisfying $Av = 0$. Decompose each vector $x \in \mathbb{R}^2$ into orthogonal components $x = x_\perp + x_\parallel$ with $x_\perp \perp v$ and $x_\parallel \parallel v$. For all x, y, z, w define:

- i. $u_1(x) = \frac{|x_\perp|^2}{2} + g_1(x) + g_2(x), \quad u_2(y) = \frac{|A^T y|^2}{2} + g(y), \quad \text{where}$
 $g_1(x) = \frac{1}{2}(M_2 F x)^T M_1 (M_2 F x), \quad g_2(x) = \frac{1}{2}(F x)^T M_2 (F x) \quad \text{and} \quad g(y) = \frac{1}{2}y^T M_1 y.$
- ii. $u_3(z) = \frac{1}{2}z^T (M_1^{-1} + M_2)z, \quad u_4(w) = \frac{1}{2}w^T M_2^{-1}w.$
- iii. $\rho(x, y) = \sup_{z, w} [b(x, y, z, w) - u_3(z) - u_4(w)].$

Consider the set:

$$W = \left\{ (x, y, z, w) : x_\perp = A^T y, z = M_1(y + M_2 F x) \quad \text{and} \quad w = M_2(z + F x) \right\}$$

We claim

$$b(x, y, z, w) - u_1(x) - u_2(y) - u_3(z) - u_4(w) \leq 0 \quad (12)$$

for all $(x, y, z, w) \in (\mathbb{R}^2)^4$ and equality holds on W . For the inequality, it suffices to prove $\rho(x, y) \leq u_1(x) + u_2(y)$.

$$\begin{aligned} \rho(x, y) &= x \cdot y + \sup_{z, w} [y \cdot z + z \cdot w + w \cdot (F x) - u_3(z) - u_4(w)] \\ &= x \cdot y + \sup_z [y \cdot z - u_3(z) + \sup_w [(z + F x) \cdot w - u_4(w)]] \\ &= x \cdot y + \sup_z [y \cdot z - u_3(z) + u_4^*(z + F x)] \end{aligned}$$

$$\begin{aligned}
&= x \cdot y + \sup_z \left[y \cdot z - u_3(z) + \frac{1}{2}(z + Fx)^T M_2(z + Fx) \right] && \text{by (3)} \\
&= x \cdot y + \sup_z \left[y \cdot z - \frac{1}{2}z^T M_1^{-1}z + z^T M_2 Fx \right] + g_2(x) \\
&= x \cdot y + \frac{1}{2}(y + M_2 Fx)^T M_1(y + M_2 Fx) + g_2(x) && \text{by (3)} \\
&= x \cdot y + y^T M_1 M_2 Fx + g_1(x) + g_2(x) + g(y) \\
&= y^T \cdot Ax + g_1(x) + g_2(x) + g(y) \\
&= y^T \cdot Ax_\perp + g_1(x) + g_2(x) + g(y) \\
&= A^T y \cdot x_\perp + g_1(x) + g_2(x) + g(y) \\
&\leq \frac{|A^T y|^2}{2} + \frac{|x_\perp|^2}{2} + g_1(x) + g_2(x) + g(y) && \text{by the Cauchy-Schwarz Inequality} \\
&= u_1(x) + u_2(y),
\end{aligned}$$

with equality when $x_\perp = A^T y$. Hence, for any element (x_0, y_0, z_0, w_0) in W , $\rho(x_0, y_0) = u_1(x_0) + u_2(y_0)$. Furthermore, by tracing the cases of equality in the preceding string of inequalities, it is not hard to show that (z_0, w_0) maximizes the map $(z, w) \mapsto y_0 \cdot z + z \cdot w + w \cdot (Fx_0) - u_3(z) - u_4(w)$. Then $b(x_0, y_0, z_0, w_0) - u_3(z_0) - u_4(w_0) = \rho(x_0, y_0)$; that is $b(x_0, y_0, z_0, w_0) - u_3(z_0) - u_4(w_0) = u_1(x_0) + u_2(y_0)$ on W , proving the claim.

Since x_\parallel and y can be chosen freely, $\dim(W) = 3$. Then, if we take any probability measure μ , concentrated on W and absolutely continuous with respect to 3-dimensional Hausdorff measure, $\text{spt}(\mu)$ will not be concentrated on the graph of a measurable map. Now, take the projections of μ as marginals; that is, set $\mu_1 = (\pi_x)_\# \mu$, $\mu_2 = (\pi_y)_\# \mu$, $\mu_3 = (\pi_z)_\# \mu$ and $\mu_4 = (\pi_w)_\# \mu$. It is straightforward to show that M_1 and M_2 can be chosen so that the canonical projections $\pi_x, \pi_y, \pi_z, \pi_w$ from W are surjective, and so the marginals μ_k are absolutely continuous with respect to Lebesgue measure. Hence, inequality (12) together with the fact that equality holds μ almost everywhere implies that μ is a solution to (KPb) which is not induced by a map. \square

4.3 The condition $m = 4$.

In this subsection we show that the hypothesis on the numbers of variables in Theorem 3.1 is necessary. We will follow the ideas behind the proof of Proposition 4.2.1.

In what follows, the presented variables are in \mathbb{R}^2 . For a given $x \in \mathbb{R}^2$ its coordinates will be denoted by x^1, x^2 .

Proposition 4.3.1. *For $b(x_1, \dots, x_m) = \sum_{k=1}^{m-1} x_k \cdot x_{k+1} + x_m \cdot x_1$, $m \geq 5$, there are absolutely continuous marginals μ_k with respect to the Lebesgue measure \mathcal{L}^2 , such that a solution of (KPb) is not concentrated on a graph.*

Proof. By part 2 of Theorem 2.2, $M = \begin{bmatrix} -1 & 3 \\ 0 & -1 \end{bmatrix} \in \mathfrak{R}^3$. Hence, we can choose positive definite matrices $M_1, M_2, M_3 > 0$ such that $M = M_1 M_2 M_3$. For all x_1, \dots, x_m define:

- i. $u_1^m(x_1) = \frac{3(x_1^2)^2}{2} + g_1(x_1) + g_2^m(x_1)$, where $g_1(x_1) = \frac{1}{2}(M_3 x_1)^T M_2 (M_3 x_1) + \frac{1}{2}(M_2 M_3 x_1)^T M_1 (M_2 M_3 x_1)$ and $g_2^m(x_1) = \frac{1}{2}x_1^T M_3 x_1 + \frac{m-5}{2}|x_1|^2$ for all $m \geq 5$.
- ii. $u_2(x_2) = \frac{3(x_2^1)^2}{2} + g(x_2)$ with $g(x_2) = \frac{1}{2}x_2^T M_1 x_2$, $u_3(x_3) = \frac{1}{2}x_3^T (M_1^{-1} + M_2)x_3$ and $u_4(x_4) = \frac{1}{2}x_4^T (M_2^{-1} + M_3)x_4$ for all $m \geq 5$.
- iii. $u_5(x_5) = \frac{1}{2}x_5^T M_3^{-1} x_5$ for $m = 5$.
- iv. For $m > 5$, $u_k(x_k) = \begin{cases} \frac{1}{2}x_5^T (M_3^{-1} + I)x_5 & \text{if } k = 5 \\ |x_k|^2 & \text{if } 5 < k < m \\ \frac{1}{2}|x_m|^2 & \text{if } k = m \end{cases}$
- v. $\rho^m(x_1, x_2) = \sup_{x_3, \dots, x_m} [b(x_1, \dots, x_m) - \sum_{k=3}^m u_k(x_k)]$ for all $m \geq 5$.

Consider the set:

$$W = \left\{ (x_1, \dots, x_m) : x_1^2 = x_2^1, \quad x_3 = M_1(x_2 + M_2 M_3 x_1), \quad x_4 = M_2(x_3 + M_3 x_1), \right. \\ \left. x_5 = M_3(x_4 + x_1) \quad \text{and} \quad x_k = x_1 + x_{k-1}, \quad \text{for } k \geq 6 \right\}$$

We claim $b(x_1, \dots, x_m) - \sum_{k=3}^m u_k(x_k) \leq u_1^m(x_1) + u_2(x_2)$ for all $(x_1, \dots, x_m) \in (\mathbb{R}^2)^m$ and equality holds on W . For the inequality, it suffices to prove $\rho^m(x_1, x_2) \leq u_1^m(x_1) + u_2(x_2)$ for all $m \geq 5$. We divide the proof of the claim into two cases:

1. For m=5

$$\begin{aligned}
\rho^5(x_1, x_2) &= x_1 \cdot x_2 + \sup_{x_3, x_4, x_5} [x_2 \cdot x_3 + x_3 \cdot x_4 + x_4 \cdot x_5 + x_5 \cdot x_1 - u_3(x_3) - u_4(x_4) - u_5(x_5)] \\
&= x_1 \cdot x_2 + \sup_{x_3} \left[x_2 \cdot x_3 - u_3(x_3) + \sup_{x_4} [x_3 \cdot x_4 - u_4(x_4) \right. \\
&\quad \left. + \sup_{x_5} [(x_4 + x_1)x_5 - u_5(x_5)]] \right] \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - u_3(x_3) + \sup_{x_4} [x_3 \cdot x_4 - u_4(x_4) + u_5^*(x_4 + x_1)]] \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - u_3(x_3) + \sup_{x_4} [x_3 \cdot x_4 - u_4(x_4) \\
&\quad + \frac{1}{2}(x_4 + x_1)^T M_3(x_4 + x_1)]] \quad \text{by (3)} \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - u_3(x_3) + \sup_{x_4} [x_3 \cdot x_4 - \frac{1}{2}x_4^T M_2^{-1}x_4 \\
&\quad + x_4^T M_3x_1 + g_2^5(x_1)]] \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - u_3(x_3) + \frac{1}{2}(x_3 + M_3x_1)^T M_2(x_3 + M_3x_1)] \\
&\quad + g_2^5(x_1) \quad \text{by (3)} \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - \frac{1}{2}x_3^T M_1^{-1}x_3 + (M_3x_1)^T M_2x_3] + g_2^5(x_1) \\
&\quad + \frac{1}{2}(M_3x_1)^T M_2(M_3x_1) \\
&= x_1 \cdot x_2 + \sup_{x_3} [x_2 \cdot x_3 - u_3(x_3) + \frac{1}{2}(x_3 + M_3x_1)^T M_2(x_3 + M_3x_1)] + g_2^5(x_1) \\
&= x_1 \cdot x_2 + \frac{1}{2}(x_2 + M_2M_3x_1)^T M_1(x_2 + M_2M_3x_1) + g_2^5(x_1) \\
&\quad + \frac{1}{2}(M_3x_1)^T M_2(M_3x_1) \quad \text{by (3)} \\
&= x_1 \cdot x_2 + x_2^T Mx_1 + g_1(x_1) + g_2^5(x_1) + g(x_2) \\
&= 3x_1^2x_2^1 + g_1(x_1) + g_2^5(x_1) + g(x_2) \\
&\leq \frac{3(x_1^2)^2}{2} + \frac{3(x_2^1)^2}{2} + g_1(x_1) + g_2^5(x_1) + g(x_2) \quad \text{by the Cauchy-Schwarz Ineq.} \\
&= u_1^5(x_1) + u_2(x_2)
\end{aligned}$$

2. The case $m \geq 6$ will be proved using induction. For $m = 6$, note that:

$$x_4 \cdot x_5 - u_5(x_5) + \sup_{x_6} [(x_5 + x_1)x_6 - u_6(x_6)] = (x_1 + x_4)x_5 - \frac{1}{2}x_5^T M_3^{-1}x_5 + \frac{1}{2}|x_1|^2$$

and

$$\sup_{x_5} [(x_1 + x_4)x_5 - \frac{1}{2}x_5^T M_3^{-1}x_5 + \frac{1}{2}|x_1|^2] = \frac{1}{2}(x_4 + x_1)^T M_3(x_4 + x_1) + \frac{1}{2}|x_1|^2.$$

Then

$$\begin{aligned} \rho^6(x_1, x_2) &= \rho^5(x_1, x_2) + \frac{1}{2}|x_1|^2 \\ &\leq u_1^5(x_1) + u_2(x_2) + \frac{1}{2}|x_1|^2 \\ &= u_1^6(x_1) + u_2(x_2) \end{aligned} \tag{13}$$

Assume the statement is true for $m - 1$. Then

$$\begin{aligned} \rho^m(x_1, x_2) &= \sup_{x_3, \dots, x_m} \left[\sum_{k=1}^{m-1} x_k \cdot x_{k+1} + x_m \cdot x_1 - \sum_{k=3}^m u_k(x_k) \right] \\ &= \sup_{x_3, \dots, x_{m-1}} \left[\sum_{k=1}^{m-2} x_k \cdot x_{k+1} - \sum_{k=3}^{m-1} u_k(x_k) + \sup_{x_m} [(x_1 + x_{m-1})x_m - u_m(x_m)] \right] \\ &= \sup_{x_3, \dots, x_{m-1}} \left[\sum_{k=1}^{m-2} x_k \cdot x_{k+1} + \frac{|x_1 + x_{m-1}|^2}{2} - \sum_{k=3}^{m-1} u_k(x_k) \right] \quad \text{by (3)} \\ &= \sup_{x_3, \dots, x_{m-1}} \left[b(x_1, \dots, x_{m-1}) - \sum_{k=3}^{m-2} u_k(x_k) - \frac{|x_{m-1}|^2}{2} + \frac{|x_1|^2}{2} \right] \\ &= \rho^{m-1}(x_1, x_2) + \frac{|x_1|^2}{2} \tag{14} \\ &\leq u_1^{m-1}(x_1) + u_2(x_2) + \frac{|x_1|^2}{2} \quad \text{by induction hypothesis} \\ &= u_1^m(x_1) + u_2(x_2). \end{aligned}$$

If $x_1^2 = x_2^1$, we obtain $\rho^5(x_1, x_2) = u_1^5(x_1) + u_2(x_2)$ and by (13), $\rho^6(x_1, x_2) = u_1^6(x_1) + u_2(x_2)$. Furthermore, by (14) $\rho^m(x_1, x_2) = \rho^{m-1}(x_1, x_2) + \frac{|x_1|^2}{2}$.

Hence, using induction we can easily prove that $\rho^m(x_1, x_2) = u_1^m(x_1) + u_2(x_2)$ for all $m \geq 5$, when $x_1^2 = x_2^1$.
On the other hand, for any element (x_1^0, \dots, x_m^0) in W , (x_3^0, \dots, x_m^0) maximizes the map:

$$(x_3, \dots, x_m) \mapsto x_2^0 \cdot x_3 + \sum_{k=3}^{m-1} x_k \cdot x_{k+1} + x_m \cdot x_1^0 - \sum_{k=3}^m u_k(x_k).$$

Hence $b(x_1^0, \dots, x_m^0) - \sum_{k=3}^m u_k(x_k^0) = \rho^m(x_1^0, x_2^0) = u_1^m(x_1^0) + u_2(x_2^0)$. This proves the claim.

The rest of the proof is identical to the proof of Proposition 4.2.1; since x_1 and x_2^2 can be chosen freely, W is three dimensional, and the claim implies that any probability measure μ supported on W is optimal for its marginals in (KPb).

□

References

- [1] V. Arnold, V, *Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits*. Ann. Inst. Fourier (Grenoble), Volume 16: 319–361, 1966.
- [2] C. S. Ballantine, *Products of positive definite matrices*. I, Pacific J. Math. Volume 23, No 3: 427-433, 1967.
- [3] C. S. Ballantine, *Products of positive definite matrices*. II, Pacific J. Math. Volume 24, No 1: 7-17, 1968.
- [4] J-D. Benamou, G. Carlier, L. Nenna. *Generalized incompressible flows, multi-marginal transport and Sinkhorn algorithm*. Numerische Mathematik. Volume 142, no.1 : pp 33-54, 2019.
- [5] M. Bernot, A. Figalli, F. Santambrogio, *Generalized solutions for the Euler equations in one and two dimensions*. J. Math. Pures Appl. (9) Volume 91: 137–155, 2009.
- [6] Y. Brenier. *Generalized solutions and hydrostatic approximation of the Euler equations* Physica D Volume 237, 14-17: pp 1982-1988, 2008.

- [7] Y. Brenier. *The least action principle and the related concept of generalized flows for incompressible perfect fluids.* J. Am. Math. Soc. 2 225-255 (1989).
- [8] G. Carlier, *On a class of multidimensional optimal transportation problems,* J. Convex Anal. 10, 2010, no.2, pp.517–529.
- [9] Carlier, G., and Ekeland, I. *Matching for teams.* Econom. Theory. 42, 2010, no. 2, pp. 397-418.
- [10] Carlier, G., and Nazaret, B., *Optimal transportation for the determinant* ESAIM Control Optim. Calc. Var. 14, 2008, no.4, pp. 678-698.
- [11] McCann, R. *Polar factorization of maps on Riemannian manifolds.* Geom. Funct. Anal. 11, 2001, pp. 589-608.
- [12] M. Colombo, L. De Pascale and S. Di Marino, *Multimarginal optimal transport maps for one-dimensional repulsive costs,* Canad. J. Math., 67 (2015), no.2, pp. 350–368.
- [13] D.G. Ebin and J. Marsden. *Groups of diffeomorphisms and the motion of an ideal incompressible fluid.* Ann. of Math. (2), 92 (1970), 102-163.
- [14] Gallouët T. and Mérigot Q. *A lagrangian scheme for the incompressible euler equation using optimal transport.* arXiv preprint arXiv:1605.00568, 2016
- [15] Gangbo, W., and Świąch, A., *Optimal maps for the multidimensional Monge-Kantorovich problem,* Comm. Pure Appl. Math. 51, 1998, no.1, pp. 23-45.
- [16] Gangbo, W. *Habilitation Thesis,* Universite de Metz, (1995)
- [17] Gangbo, W. and McCann, R.J. *The geometry of optimal transportation.* Acta Math., 177 (1996), 113-161.
- [18] Heinich, H., *Probleme de Monge pour n probabilités.* C.R. Math. Acad. Sci. Paris 334, 2002, no. 9, pp. 793-795.
- [19] Kellerer, H.G., *Duality theorems for marginal problems.* Z. Wahrsch. Verw. Gebiete 67, 1984, pp. 399-432.

- [20] Y.-H. Kim and B. Pass., *A general condition for Monge solutions in the multi-marginal optimal transport problem.* SIAM J. Math. Anal, 46:15381550, 2014.
- [21] J-M. Mirebeau, *Numerical resolution of Euler equations, through semi-discrete optimal transport*, hal01237356, 2015.
- [22] B. Pass, PhD Thesis. *University of Toronto*, 2011.
- [23] B. Pass. *On the local structure of optimal measures in the multi-marginal optimal transportation problem* Calc. Var. and PDE (2012)43: pp 529-536.
- [24] B.Pass, *Remarks on the semi-classical Hohenberg-Kohn functional* Non-linearity, 26 (2013), no. 9, pp 2731–2744.
- [25] B. Pass. *Multi-marginal optimal transport: theory and applications.* ESAIM: Math. Model. Numer. Anal., 49:1771-1790, 2015.
- [26] B. Pass. *Uniqueness and Monge solutions in the Multi-marginal optimal transportation problem.* SIAM J. Math. Anal. 43:2758-2775, 2015.
- [27] F. Santambrogio, *Optimal transport for Applied Mathematicians: Calculus of Variations, PDEs and Modeling* volume 87 of *Progress in Nonlinear Differential Equations and Their Applications* 87, Birkhauser, Heidelberg, 2015.
- [28] Villani, C., *Topics in optimal transportation*, volume 58 of Graduate Studies in Mathematics. American Mathematical Society, Providence, 2003.
- [29] Villani, C., *Optimal Transport: old and new*, volume 338 of Grundlehren der mathematischen Wissenschaften. Springer, New York, 2009.