AVERAGE–CASE COMPLEXITY OF THE EUCLIDEAN ALGORITHM WITH A FIXED POLYNOMIAL OVER A FINITE FIELD

NARDO GIMÉNEZ¹, GUILLERMO MATERA^{1,2,3}, MARIANA PÉREZ^{3,4}, AND MELINA PRIVITELLI^{3,5}

ABSTRACT. We analyze the behavior of the Euclidean algorithm applied to pairs (g, f) of univariate nonconstant polynomials over a finite field \mathbb{F}_q of q elements when the highestdegree polynomial g is fixed. Considering all the elements f of fixed degree, we establish asymptotically optimal bounds in terms of q for the number of elements f which are relatively prime with g and for the average degree of gcd(g, f). The accuracy of our estimates is confirmed by practical experiments. We also exhibit asymptotically optimal bounds for the average-case complexity of the Euclidean algorithm applied to pairs (g, f)as above.

1. INTRODUCTION

Let \mathbb{F}_q be the finite field of q elements, where q is a prime power, let T be an indeterminate over \mathbb{F}_q and $\mathbb{F}_q[T]$ the ring of univariate polynomials in T with coefficients in \mathbb{F}_q . In this paper we are concerned with the polynomial gcd problem for elements of $\mathbb{F}_q[T]$, namely the problem of computing the greatest common divisor of two nonzero polynomials in $\mathbb{F}_q[T]$.

The fundamental computational tool for this problem is the Euclidean algorithm, and many variants of it are known in the literature (see, e.g., [vzGG99]). It is well-known that the Euclidean algorithm in $\mathbb{F}_q[T]$ requires a number of polynomial divisions which is linear in the degree of the input polynomials. In particular, we are interested in its average-case complexity, which has been the subject of several papers. The paper [MvzG90] establishes the average-case complexity of the Euclidean algorithm and some variants of it based on explicit counting. In [Nor89], the average-case complexity of variants of the Euclidean algorithm is considered using generating functions. Finally, [LV08] and [BNNV14] analyze the average-case complexity and related costs of the Euclidean algorithm and variants using tools of analytic combinatorics such as bivariate generating functions.

All these results consider the average, for fixed degrees e > d > 0, over the set of pairs $(g, f) \in \mathbb{F}_q[T] \times \mathbb{F}_q[T]$ with g monic of degree e and f either of degree at most d, or of degree less than e, assuming the uniform distribution of pairs. Nevertheless, there are important tasks which rely heavily on the computation of gcd's and lie outside the scope of these analyses. For example, a critical step in the standard algorithm for finding the roots in \mathbb{F}_q of a polynomial $f \in \mathbb{F}_q[T]$ with deg f < q consists of computing $gcd(T^q - T, f)$ (see, e.g., [vzGG99]). As the first element in the pair $(T^q - T, f)$ is a fixed polynomial, average-case analyses as before do not contribute to the analysis of the complexity of this problem.

In this paper we consider, for fixed degrees e > d > 0 and a fixed (arbitrary) $g \in \mathbb{F}_q[T]$ monic of degree e, the average-case complexity of the Euclidean algorithm over the set of pairs (g, f) with $f \in \mathbb{F}_q[T]$ monic of degree d, endowed with the uniform probability. We shall be interested in the case $q \gg e$; in this sense, all our results may be regarded as asymptotic in q.

Date: December 26, 2021.

Key words and phrases. Finite fields, rational points, Euclidean algorithm, Schur functions, resultant, average–case complexity.

The authors were partially supported by the grants PIP CONICET 11220130100598 and PIO CONICET-UNGS 14420140100027.

We discuss a number of issues concerning this case of the Euclidean algorithm. Our first result shows that the average degree $E[\mathcal{X}_g]$ of gcd(g, f) for a random element f of $\mathbb{F}_q[T]$, monic of degree d, decreases fast as q tends to infinity. Further, we prove that the decrease rate depends on the factorization pattern of g (see Theorem 4.5 for a precise statement):

(1.1)
$$\left| E[\mathcal{X}_g] - \frac{k\lambda_k^*}{q^k} \right| = \mathcal{O}\left(\frac{1}{q^{k+1}}\right),$$

where λ_k^* denotes the number of distinct monic irreducible factors of g of degree k. The average degree of the gcd of a random pair of elements in $\mathbb{F}_q[T]$ of degrees e and d as above is $(1 - q^{-d})/(q - 1)$ (see [MvzG90, Corollary 2.6]). Our result, although not as precise as the latter, confirms that in our case the average degree of the gcd is $\mathcal{O}(q^{-1})$ (for fixed d, e).

We also show that, with high probability, g and a random monic polynomial f of $\mathbb{F}_q[T]$ of degree d are relatively prime. In fact, we have the following estimate for the probability \mathcal{P}_0 that gcd(g, f) = 1 (see Theorem 4.2):

(1.2)
$$\left| \mathcal{P}_0 - \left(1 - \frac{\lambda_k^*}{q^k} \right) \right| = \mathcal{O}\left(\frac{1}{q^{k+1}} \right)$$

This may be compared with the probability 1-1/q that a random pair of elements of $\mathbb{F}_q[T]$ of degrees e and d are relatively prime (see, e.g., [MvzG90, Proposition 2.4]).

Finally, we analyze the average number $E[t_g^{\text{div}}]$, $E[t_g^{\pm}]$ and $E[t_g^{-,\times}]$ of polynomial divisions, divisions in \mathbb{F}_q , and additions/multiplications in \mathbb{F}_q , performed by the Euclidean algorithm. We have the following bounds (see Theorem 5.4):

$$\left|\frac{E[t_g^{\mathsf{div}}]}{d+1} - 1\right| \le \frac{de}{q}, \qquad \left|\frac{E[t_g^{\div}]}{e+d+1} - 1\right| \le \frac{de}{q}, \qquad \left|\frac{E[t_g^{-,\times}]}{de} - 1\right| \le \frac{de}{q}.$$

The main terms in these bounds agree with those in the corresponding ones for random pairs of polynomials of degree e and d with e > d, according to [MvzG90, Theorem 2.1].

Our approach relies on estimating the number of polynomials f for which gcd(g, f) has a given degree. For this purpose we use classical tools of elimination theory, which are combined with bounds on the number of common zeros with coordinates in \mathbb{F}_q of multivariate polynomials defined over the algebraic closure $\overline{\mathbb{F}}_q$ of \mathbb{F}_q . Another critical point is a lower bound on the number of polynomials f for which the Euclidean algorithm performs the highest possible number of steps. Such a lower bound relies on a description of certain coefficients of the sequence of quotients and remainders determined by the Euclidean algorithm in terms symmetric functions, following [Las03]. Combining such a description with upper bounds on the number of zeros with coordinates in \mathbb{F}_q of multivariate polynomials with coefficients in \mathbb{F}_q , we are able to control the number of polynomials f for which gcd(g, f) has a given degree. Our results are then expressed in terms of these quantities.

The paper is organized as follows. In Section 2 we recall the description of remainders and quotients arising in the Euclidean algorithm applied to a "generic" pair of polynomials of given degrees in terms of symmetric functions. In Section 3 we use this machinery to estimate the degrees of the leading coefficients of the remainders in the generic case and we consider the behavior of the Euclidean algorithm under specializations. In Section 4 we estimate on the number of polynomials f for which gcd(g, f) has a given degree, which are used to prove (1.1) and (1.2). In Section 5 we use the results of Sections 3 and 4 to establish the results on the average-case complexity. Finally, in Section 6 we report on some simulations we perform which show the behavior predicted by the theoretical estimates (1.1) and (1.2).

2. Basic notions and notations

Let \mathbb{F}_q be the finite field of q elements and $\overline{\mathbb{F}}_q$ its algebraic closure. Let X_1, \ldots, X_n be indeterminates over $\overline{\mathbb{F}}_q$. For $\mathbb{K} = \mathbb{F}_q$ or $\mathbb{K} = \overline{\mathbb{F}}_q$, we denote by $\mathbb{K}[X_1, \ldots, X_n]$ the ring of multivariate polynomials in X_1, \ldots, X_n and coefficients in \mathbb{K} . By \mathbb{A}^n we denote the affine *n*-dimensional space $\mathbb{A}^n := \overline{\mathbb{F}}_q^n$, endowed with its Zariski topology over \mathbb{K} , for which a closed set is the zero locus of a set of polynomials of $\mathbb{K}[X_1, \ldots, X_n]$. A subset $V \subset \mathbb{A}^r$ is an affine variety of defined over \mathbb{K} (or an affine \mathbb{K} -variety) if it is the set of common zeros in \mathbb{A}^n of polynomials $F_1, \ldots, F_m \in \mathbb{K}[X_1, \ldots, X_n]$. We shall denote by $\{F_1 = 0, \ldots, F_m = 0\}$ or $V(F_1, \ldots, F_m)$ the affine \mathbb{K} -variety consisting of the common zeros of F_1, \ldots, F_m .

A \mathbb{K} -variety V is *irreducible* if it cannot be expressed as a finite union of proper \mathbb{K} subvarieties of V. Any \mathbb{K} -variety V can be expressed as an irredundant union $V = C_1 \cup \cdots \cup C_s$ of irreducible \mathbb{K} -varieties, unique up to reordering, called the *irreducible* \mathbb{K} -components of V. We say that V has *pure dimension* r if every irreducible \mathbb{K} -component of V has dimension r. A \mathbb{K} -variety of \mathbb{A}^n of pure dimension n-1 is called a \mathbb{K} -hypersurface. A \mathbb{K} hypersurface of \mathbb{A}^n can also be described as the set of zeros of a single nonzero polynomial of $\mathbb{K}[X_1, \ldots, X_n]$.

The degree deg V of an irreducible \mathbb{K} -variety V is the maximum of the cardinality $|V \cap L|$ of $V \cap L$, considering all the linear spaces L of codimension dim V such that $|V \cap L| < \infty$. More generally, following [Hei83] (see also [Ful84]), if $V = C_1 \cup \cdots \cup C_s$ is the decomposition of V into irreducible \mathbb{K} -components, we define the degree of V as

$$\deg V := \sum_{i=1}^{s} \deg \mathcal{C}_i.$$

The degree of a \mathbb{K} -hypersurface V is the degree of a polynomial of minimal degree defining V. In particular, the degree of a linear variety is equal to 1.

Let $\mathbb{A}^n(\mathbb{F}_q)$ be the *n*-dimensional \mathbb{F}_q -vector space \mathbb{F}_q^n . For an affine variety $V \subset \mathbb{A}^n$, the set of \mathbb{F}_q -rational points $V(\mathbb{F}_q)$ of V is defined as $V(\mathbb{F}_q) := V \cap \mathbb{A}^n(\mathbb{F}_q)$. For an affine variety $V \subset \mathbb{A}^n$ of dimension r and degree $d \ge 0$, we have the following bound (see, e.g., [CM06, Lemma 2.1]):

$$(2.1) |V(\mathbb{F}_q)| \le d q^r.$$

2.1. Symmetric functions and the Euclidean algorithm. Next we gather the terminology and results we will use concerning the description of the Euclidean algorithm in terms of symmetric functions, following [Las03].

We call a finite set of indeterminates \mathbb{A} over \mathbb{F}_q an *alphabet* and denote its cardinality by $|\mathbb{A}|$. The *elementary symmetric functions* $\Lambda^i(\mathbb{A})$ and the *complete functions* $S^i(\mathbb{A})$ $(i \ge 0)$ are defined by means of the following identities of formal power series in the variable z:

$$\prod_{a \in \mathbb{A}} (1+za) = \sum_{i \ge 0} \Lambda^i(\mathbb{A}) z^i, \quad \prod_{a \in \mathbb{A}} \frac{1}{1-za} = \sum_{i \ge 0} S^i(\mathbb{A}) z^i.$$

We further define $\Lambda^i(\mathbb{A}) := 0$ and $S^i(\mathbb{A}) := 0$ for i < 0. Observe that $\Lambda^i(\mathbb{A}) = 0$ if $i > |\mathbb{A}|$. Writing $\mathbb{A} + \mathbb{B}$ for the disjoint union of two alphabets \mathbb{A} and \mathbb{B} , we have the following *Cauchy formulas*:

(2.2)
$$\Lambda^{i}(\mathbb{A} + \mathbb{B}) = \sum_{j+k=i} \Lambda^{j}(\mathbb{A})\Lambda^{k}(\mathbb{B}), \quad S^{i}(\mathbb{A} + \mathbb{B}) = \sum_{j+k=i} S^{j}(\mathbb{A})S^{k}(\mathbb{B}),$$

Define $S^i(\mathbb{A} - \mathbb{B})$ $(i \ge 0)$ by means of the identity

$$\frac{\prod_{b\in\mathbb{B}}(1-zb)}{\prod_{a\in\mathbb{A}}(1-za)} = \sum_{i\geq 0} S^i(\mathbb{A}-\mathbb{B})z^i,$$

and set $S^i(\mathbb{A} - \mathbb{B}) := 0$ for i < 0. Define $S^i(-\mathbb{A}) := (-1)^i \Lambda^i(\mathbb{A})$ for any integer *i*. Thus, besides (2.2) we have

(2.3)
$$S^{i}(\mathbb{A} - \mathbb{B}) = \sum_{j+k=i} S^{j}(\mathbb{A})S^{k}(-\mathbb{B}).$$

We shall express polynomials using this terminology. Indeed, let $n := |\mathbb{A}|$ and identify a single indeterminate T with the alphabet $\{T\}$. Since $S^i(T) = T^i$ for $i \ge 0$, according to (2.3) we have that

$$S^{n}(T - \mathbb{A}) = \sum_{i=0}^{n} S^{n-i}(-\mathbb{A})T^{i}$$

is the polynomial in T having \mathbb{A} as its set of roots.

2.2. Schur functions. Let \mathbb{A} be an alphabet of cardinality n. Given $J := (j_1, \ldots, j_n) \in \mathbb{Z}_{>0}^n$, the Schur function $S_J(\mathbb{A})$ is defined as the determinant

$$S_J(\mathbb{A}) := \det \left(S^{j_k + k - h}(\mathbb{A}) \right)_{1 \le h, k \le n}$$

In other words,

$$S_{J}(\mathbb{A}) = \det \begin{pmatrix} S^{j_{1}}(\mathbb{A}) & S^{j_{2}+1}(\mathbb{A}) & \cdots & S^{j_{n}+n-1}(\mathbb{A}) \\ S^{j_{1}-1}(\mathbb{A}) & S^{j_{2}}(\mathbb{A}) & \ddots & \vdots \\ \vdots & \vdots & \cdots & S^{j_{n}+1}(\mathbb{A}) \\ S^{j_{1}-n+1}(\mathbb{A}) & S^{j_{2}-n+2}(\mathbb{A}) & \cdots & S^{j_{n}}(\mathbb{A}) \end{pmatrix}$$

Given n, two sets of alphabets $\{\mathbb{A}_1, \ldots, \mathbb{A}_n\}$, $\{\mathbb{B}_1, \ldots, \mathbb{B}_n\}$ and $J \in \mathbb{Z}_{\geq 0}^n$, we have the *multi-Schur function*

$$S_J(\mathbb{A}_1 - \mathbb{B}_1, \dots, \mathbb{A}_n - \mathbb{B}_n) := \det \left(S^{j_k + k - h} (\mathbb{A}_k - \mathbb{B}_k) \right)_{1 \le h, k \le n},$$

namely

$$S_{J}(\mathbb{A}_{1} - \mathbb{B}_{1}, \dots, \mathbb{A}_{n} - \mathbb{B}_{n}) =$$

$$\det \begin{pmatrix} S^{j_{1}}(\mathbb{A}_{1} - \mathbb{B}_{1}) & S^{j_{2}+1}(\mathbb{A}_{2} - \mathbb{B}_{2}) & \cdots & S^{j_{n}+n-1}(\mathbb{A}_{n} - \mathbb{B}_{n}) \\ S^{j_{1}-1}(\mathbb{A}_{1} - \mathbb{B}_{1}) & S^{j_{2}}(\mathbb{A}_{2} - \mathbb{B}_{2}) & \ddots & \vdots \\ \vdots & \vdots & \cdots & S^{j_{n}+1}(\mathbb{A}_{n} - \mathbb{B}_{n}) \\ S^{j_{1}-n+1}(\mathbb{A}_{1} - \mathbb{B}_{1}) & S^{j_{2}-n+2}(\mathbb{A}_{2} - \mathbb{B}_{2}) & \cdots & S^{j_{n}}(\mathbb{A}_{n} - \mathbb{B}_{n}) \end{pmatrix}$$

Finally, given $H \in \mathbb{Z}_{\geq 0}^{p}$ and $K \in \mathbb{Z}_{\geq 0}^{q}$, and alphabets \mathbb{A} , \mathbb{B} , \mathbb{C} , \mathbb{D} , we shall consider the multi-Schur function $S_{H;K}(\mathbb{A} - \mathbb{B}; \mathbb{C} - \mathbb{D})$ with index (H; K), the concatenation of H and K, and alphabets $\mathbb{A}_{1} = \mathbb{A}, \ldots, \mathbb{A}_{p} = \mathbb{A}$, $\mathbb{B}_{1} = \mathbb{B}, \ldots, \mathbb{B}_{p} = \mathbb{B}$, $\mathbb{A}_{p+1} = \mathbb{C}, \ldots, \mathbb{A}_{p+q} = \mathbb{C}$, $\mathbb{B}_{p+1} = \mathbb{D}, \ldots, \mathbb{B}_{p+q} = \mathbb{D}$. When a tuple $J = (j_{1}, \ldots, j_{n}) \in \mathbb{Z}_{\geq 0}^{n}$ with $j_{i} = m$ for $1 \leq i \leq n$ appears as an index in a Schur function, we denote it as m^{n} . For example, for $\ell \in \mathbb{Z}_{\geq 0}$, we have

$$S_{m^{n};\ell}(\mathbb{A} - \mathbb{B}; T) = \det \begin{pmatrix} S^{m}(\mathbb{A} - \mathbb{B}) & \cdots & S^{m+n-1}(\mathbb{A} - \mathbb{B}) & T^{\ell+n} \\ S^{m-1}(\mathbb{A} - \mathbb{B}) & \ddots & \vdots & \vdots \\ \vdots & \ddots & S^{m}(\mathbb{A} - \mathbb{B}) & T^{\ell+1} \\ S^{m-n}(\mathbb{A} - \mathbb{B}) & \cdots & S^{m-1}(\mathbb{A} - \mathbb{B}) & T^{\ell} \end{pmatrix}$$

We have the following result (see [Las03, equation (1.4.8)]).

Lemma 2.1. Let $J \in \mathbb{Z}_{\geq 0}^n$, $k \in \mathbb{Z}_{\geq 0}$, \mathbb{A} , \mathbb{B} alphabets and T an indeterminate. Then

$$S_J(\mathbb{A} - \mathbb{B} - T)T^k = S_{J;k}(\mathbb{A} - \mathbb{B}; T).$$

2.3. Remainders as Schur functions. Let \mathbb{A} and \mathbb{B} be two alphabets of cardinalities e and d respectively, with e > d. When the Euclidean algorithm is applied to two generic polynomials $S^e(T - \mathbb{A})$ and $S^d(T - \mathbb{B})$ we obtain d remainders $\mathcal{R}_1, \ldots, \mathcal{R}_d$ and quotients q_1, \ldots, q_d satisfying the following identities:

(2.4)

$$S^{e}(T - \mathbb{A}) = q_{1}S^{d}(T - \mathbb{B}) + \mathcal{R}_{1},$$

$$S^{d}(T - \mathbb{B}) = q_{2}\mathcal{R}_{1} + \mathcal{R}_{2},$$

$$\mathcal{R}_{1} = q_{3}\mathcal{R}_{2} + \mathcal{R}_{3},$$

$$\vdots$$

$$\mathcal{R}_{d-2} = q_{d}\mathcal{R}_{d-1} + \mathcal{R}_{d}.$$

Here $\deg_T q_1 = e - d$, $\deg_T q_i = 1$ for $2 \le i \le d$, and $\deg_T \mathcal{R}_i = d - i$ for $1 \le i \le d$. It turns out that all the remainders \mathcal{R}_i are elements of the ring $\mathbb{F}_q[\mathbb{A}, \mathbb{B}][T]$. Further, we may express these remainders in terms of Schur functions [Las03, equation (3.1.5)]:

(2.5)
$$\mathcal{R}_k = (-1)^{d-k+1} S_{(e-d+k)^{k-1}} (\mathbb{B} - \mathbb{A} - T) S^e (T - \mathbb{A}) + S_{k^{e-d+k-1}} (\mathbb{A} - \mathbb{B} - T) S^d (T - \mathbb{B}).$$

3. Degree bounds for the remainders in the generic case

In the sequel, for $1 \leq k \leq d$ we denote by $F_k \in \mathbb{F}_q[\mathbb{A}, \mathbb{B}]$ the leading coefficient of \mathcal{R}_k , considered as an element of $\mathbb{F}_q[\mathbb{A}, \mathbb{B}][T]$. Let $\mathbf{S} := (S^i(-\mathbb{B}) : 1 \leq i \leq d)$ and $\mathbf{T} := (S^i(\mathbb{A}) : 1 \leq i \leq e)$. Observe that both \mathbf{S} and \mathbf{T} are algebraically independent sets over \mathbb{F}_q .

Proposition 3.1. F_k is a nonzero element of $\mathbb{F}_q[\mathbf{T}][\mathbf{S}]$ of degree deg_S $F_k = e - d + k$. Further, it is monic of degree e - d + k in $S^k(-\mathbb{B})$.

Proof. Since $S^d(T - \mathbb{B}) = \sum_{j=0}^d S^{d-j}(-\mathbb{B})T^j$, we can write

$$S_{k^{e-d+k-1}}(\mathbb{A}-\mathbb{B}-T)S^d(T-\mathbb{B}) = \sum_{j=0}^d S^{d-j}(-\mathbb{B})S_{k^{e-d+k-1}}(\mathbb{A}-\mathbb{B}-T)T^j.$$

By Lemma 2.1 we have

$$S_{k^{e-d+k-1}}(\mathbb{A}-\mathbb{B}-T)T^{j}=S_{k^{e-d+k-1};j}(\mathbb{A}-\mathbb{B};T),$$

where (3.1)

$$S_{k^{e-d+k-1};j}(\mathbb{A}-\mathbb{B};T) = \det \begin{pmatrix} S^{k}(\mathbb{A}-\mathbb{B}) & \cdots & S^{e-d+2k-2}(\mathbb{A}-\mathbb{B}) & T^{j+e-d+k-1} \\ S^{k-1}(\mathbb{A}-\mathbb{B}) & \ddots & \vdots & \vdots \\ \vdots & \ddots & S^{k}(\mathbb{A}-\mathbb{B}) & T^{j+1} \\ S^{-(e-d-1)}(\mathbb{A}-\mathbb{B}) & \cdots & S^{k-1}(\mathbb{A}-\mathbb{B}) & T^{j} \end{pmatrix}.$$

Similarly, taking into account that $S^e(T - \mathbb{A}) = \sum_{h=0}^{e} S^{e-h}(-\mathbb{A})T^h$, we see that

$$S_{(e-d+k)^{k-1}}(\mathbb{B} - \mathbb{A} - T)S^{e}(T - \mathbb{A}) = \sum_{h=0}^{e} S^{e-h}(-\mathbb{A})S_{(e-d+k)^{k-1}}(\mathbb{B} - \mathbb{A} - T)T^{h}.$$

Again by Lemma 2.1, we have

$$S_{(e-d+k)^{k-1}}(\mathbb{B}-\mathbb{A}-T)T^h = S_{(e-d+k)^{k-1};h}(\mathbb{B}-\mathbb{A};T),$$

where (3.2)

$$S_{(e-d+k)^{k-1};h}(\mathbb{B}-\mathbb{A};T) = \det \begin{pmatrix} S^{e-d+k}(\mathbb{A}-\mathbb{B}) & \cdots & S^{e-d+2k-2}(\mathbb{A}-\mathbb{B}) & T^{h+k-1} \\ S^{e-d+k-1}(\mathbb{A}-\mathbb{B}) & \ddots & \vdots & \vdots \\ \vdots & \ddots & S^{e-d+k}(\mathbb{A}-\mathbb{B}) & T^{h+1} \\ S^{e-d+1}(\mathbb{A}-\mathbb{B}) & \cdots & S^{e-d+k-1}(\mathbb{A}-\mathbb{B}) & T^{h} \end{pmatrix}.$$

Denote by $A_{j,d-k}$ the coefficient of the monomial T^{d-k} in $S_{k^{e-d+k-1};j}(\mathbb{A} - \mathbb{B};T)$ for $0 \leq j \leq d$, considering $S_{k^{e-d+k-1};j}(\mathbb{A} - \mathbb{B};T)$ as an element of $\mathbb{F}_{q}[\mathbb{A},\mathbb{B}][T]$. Further, denote by $B_{h,d-k}$ the coefficient of the monomial T^{d-k} in $S_{(e-d+k)^{k-1};h}(\mathbb{B} - \mathbb{A};T)$ for $0 \leq h \leq e$. By (2.5) we have

$$F_k = (-1)^{d-k+1} \sum_{h=0}^{e} S^{e-h}(-\mathbb{A}) B_{h,d-k} + \sum_{j=0}^{d} S^{d-j}(-\mathbb{B}) A_{j,d-k}.$$

By (3.1) and (3.2) it is clear that $A_{j,d-k} = 0$ for j > d-k and j + e - d + k - 1 < d - kand $B_{h,d-k} = 0$ for h > d - k and h + k - 1 < d - k. Thus, (3.3)

$$F_{k} = (-1)^{d-k+1} \sum_{h=\max\{d-2k+1,0\}}^{d-k} S^{e-h}(-\mathbb{A})B_{h,d-k} + \sum_{j=\max\{2d-e-2k+1,0\}}^{d-k} S^{d-j}(-\mathbb{B})A_{j,d-k}.$$

Now, $A_{j,d-k}$ and $B_{h,d-k}$ are the determinants of the submatrices obtained by removing the last column and the row corresponding to T^{d-k} in the matrices of (3.1) and (3.2) respectively. More precisely,

$$A_{j,d-k} = \det \begin{pmatrix} S^k(\mathbb{A} - \mathbb{B}) & \cdots & & \\ \vdots & \ddots & & \\ & & S^k(\mathbb{A} - \mathbb{B}) & \cdots & \\ & & & \vdots & S^{k-1}(\mathbb{A} - \mathbb{B}) & \cdots \\ & & & & \vdots & \ddots \\ & & & & & S^{k-1}(\mathbb{A} - \mathbb{B}) \end{pmatrix},$$

with the first j + e - 2d + 2k - 1 columns having $S^k(\mathbb{A} - \mathbb{B})$ in the diagonal and the last d - k - j columns having $S^{k-1}(\mathbb{A} - \mathbb{B})$ in the diagonal, and

$$B_{h,d-k} = \det \begin{pmatrix} S^{e-d+k}(\mathbb{A} - \mathbb{B}) \cdots & & \\ \vdots & \ddots & \\ & S^{e-d+k}(\mathbb{A} - \mathbb{B}) & \cdots \\ & \vdots & S^{e-d+k-1}(\mathbb{A} - \mathbb{B}) & \cdots \\ & & \vdots & \ddots \\ & & & S^{e-d+k-1}(\mathbb{A} - \mathbb{B}) \end{pmatrix}$$

with the first h - d + 2k - 1 columns having $S^{e-d+k}(\mathbb{A} - \mathbb{B})$ in the diagonal and the last d - k - h columns having $S^{e-d+k-1}(\mathbb{A} - \mathbb{B})$ in the diagonal.

Considering $A_{j,d-k}$ and $B_{h,d-k}$ as polynomials in the variables $\mathbf{S} := (S^i(-\mathbb{B}): 1 \leq i \leq d)$ and coefficients in $\mathbb{F}_q[\mathbf{T}]$, by their determinantal expressions we conclude that

(3.4)
$$\deg_{\mathbf{S}} A_{j,d-k} \le e - d + k - 1, \quad \deg_{\mathbf{S}} B_{h,d-k} \le k - 1.$$

Combining these upper bounds and (3.3) we readily see that

$$(3.5) deg_{\mathbf{S}} F_k \le e - d + k.$$

Claim. $A_{d-k,d-k}$ is a nonzero element of $\mathbb{F}_q[\mathbf{T}][\mathbf{S}]$ with $\deg_{\mathbf{S}} A_{d-k,d-k} = e - d + k - 1$. Further, $A_{d-k,d-k}$ is monic of degree e - d + k - 1 in $S^k(-\mathbb{B})$.

Proof of Claim. Let N := e - d + k - 1. Observe that the determinantal expression of $A_{d-k,d-k}$ consists of e - d + k - 1 columns having $S^k(\mathbb{A} - \mathbb{B})$ in the diagonal. More precisely, $A_{d-k,d-k} = \det(a_{ij})_{1 \le i,j \le N}$, where $a_{ij} := S^{k+j-i}(\mathbb{A} - \mathbb{B})$. We remark that

$$S^{h}(\mathbb{A} - \mathbb{B}) = S^{h}(-\mathbb{B}) + S^{1}(\mathbb{A})S^{h-1}(-\mathbb{B}) + \dots + S^{h}(\mathbb{A})$$

is a polynomial of degree one in the variables **S** for $1 \leq h \leq N$. Further, $S^k(\mathbb{A} - \mathbb{B})$ is monic of degree one in $S^k(-\mathbb{B})$ and $S^h(\mathbb{A} - \mathbb{B})$ is of degree zero in $S^k(-\mathbb{B})$ for h < k. Write

$$A_{d-k,\,d-k} = \sum_{\sigma} \pm a_{1\sigma_1} a_{2\sigma_2} \cdots a_{N\sigma_N},$$

where σ runs over all permutations of $(1, 2, \dots, N)$. By the previous remarks we see that

$$\deg_{\mathbf{S}} A_{d-k,\,d-k} \le N.$$

To prove the equality, consider a permutation $(\sigma_1, \ldots, \sigma_N) \neq (1, 2, \ldots, N)$. Then there exists an index *i* with $\sigma_i < i$. For such an index, since $k + \sigma_i - i < k$, the entry $a_{i\sigma_i} = S^{k+\sigma_i-i}(\mathbb{A}-\mathbb{B})$ has degree zero in $S^k(-\mathbb{B})$. Thus $a_{1\sigma_1}a_{2\sigma_2}\cdots a_{N\sigma_N}$ has degree at most N-1 in $S^k(-\mathbb{B})$. On the other hand, the term $a_{11}\cdots a_{NN} = S^k(\mathbb{A}-\mathbb{B})^N$ is monic of degree N in $S^k(-\mathbb{B})$. This implies the claim. \Box

Write the second sum in (3.3) as

$$\sum_{j=\max\{2d-e-2k+1,0\}}^{d-k} S^{d-j}(-\mathbb{B})A_{j,d-k} = S^k(-\mathbb{B})A_{d-k,d-k} + \sum_{j=\max\{2d-e-2k+1,0\}}^{d-k-1} S^{d-j}(-\mathbb{B})A_{j,d-k}.$$

According to the claim, the polynomial $S^k(-\mathbb{B})A_{d-k,d-k}$ is monic of degree e - d + k in $S^k(-\mathbb{B})$. Further, by (3.4), taking into account that $S^{d-j}(-\mathbb{B}) \neq S^k(-\mathbb{B})$ for $0 \leq j \leq d-k-1$ it follows that

$$\deg_{S^{k}(-\mathbb{B})}\left(\sum_{j=\max\{2d-e-2k+1,0\}}^{d-k-1} S^{d-j}(-\mathbb{B})A_{j,d-k}\right) \le e-d+k-1$$

Therefore, $\sum_{j=\max\{2d-e-2k+1,0\}}^{d-k} S^{d-j}(-\mathbb{B})A_{j,d-k}$ is monic of degree e-d+k in $S^k(-\mathbb{B})$. On the other hand, according to (3.4), the first sum in the right-hand side of (3.3) has degree at most k-1, and then less than e-d+k-1, in $S^k(-\mathbb{B})$. We conclude that F_k is monic of degree e-d+k in $S^k(-\mathbb{B})$. This together with (3.5) implies that $\deg_{\mathbf{S}} F_k = e-d+k$, which finishes the proof of the proposition. \Box

3.1. Specialization of the generic case. As expressed in Section 2.3, for generic input elements $S^e(T - \mathbb{A}) \in \mathbb{F}_q[\mathbb{A}][T]$ and $S^d(T - \mathbb{B}) \in \mathbb{F}_q[\mathbb{B}][T]$, the Euclidean algorithm performs d steps. Further, if the Euclidean algorithm is applied to polynomials $g, f \in \mathbb{F}_q[T]$ with deg g = e and deg f = d, and performs d steps, then the degrees of the successive remainders decrease by 1 each step, and the sequences of quotients and remainders associated to g and f coincide with the specialization of the sequence associated to $S^e(T - \mathbb{A})$ and $S^d(T - \mathbb{B})$. The next result shows that, if the sequence of remainders associated to polynomials $g, f \in \mathbb{F}_q[T]$ with deg g = e and deg f = d fails to have the degree pattern of the generic case, the first remainder where such a failure occurs is still a specialization of the corresponding one of the generic case.

Lemma 3.2. For a specialization $\mathbb{A} \mapsto \overline{a}$ and $\mathbb{B} \mapsto \overline{b}$ in \mathbb{F}_q , denote by r_1, \ldots, r_k the first k remainders of the application of the Euclidean algorithm to $S^e(T - \overline{a})$ and $S^d(T - \overline{b})$. If deg $r_i = d - i$ for $1 \leq i \leq k - 1$, then $r_i = \mathcal{R}_i(\overline{a}, \overline{b})$ for $1 \leq i \leq k$. Further, if deg $\mathcal{R}_i(\overline{a}, \overline{b}) = d - i$ for $1 \leq i \leq k - 1$, then $r_i = \mathcal{R}_i(\overline{a}, \overline{b})$ for $1 \leq i \leq k$.

Proof. Substituting \overline{a} for \mathbb{A} and \overline{b} for \mathbb{B} in the first identity of (2.4) we easily see that $\mathcal{R}_1(\overline{a},\overline{b})$ is the remainder in the division of $S^e(T-\overline{a})$ by $S^d(T-\overline{b})$, which proves that $r_1 = \mathcal{R}_1(\overline{a},\overline{b})$. Let j > 1 and assume inductively that $r_i = \mathcal{R}_i(\overline{a},\overline{b})$ for $1 \le i \le j < k$. Thus deg $\mathcal{R}_i(\overline{a},\overline{b}) = d-i$ for $1 \le i \le j$. Taking into account that F_i is the leading coefficient of \mathcal{R}_i for $1 \le i \le k$, we deduce that $F_i(\overline{a},\overline{b}) \ne 0$ for $1 \le i \le j$. Since $q_{j+1} \in \mathbb{F}_q[\mathbb{A},\mathbb{B}]_{F_i}[T]$,

where $\mathbb{F}_{q}[\mathbb{A}, \mathbb{B}]_{F_{j}}$ is the localization of $\mathbb{F}_{q}[\mathbb{A}, \mathbb{B}]$ at F_{j} , we can substitute \overline{a} for \mathbb{A} and \overline{b} for \mathbb{B} in the (j + 1)th equation of (2.4) to obtain

$$r_{j-1} = q_{j+1}(\overline{a}, b)r_j + \mathcal{R}_{j+1}(\overline{a}, b).$$

Since

$$\deg_T \mathcal{R}_{j+1}(\overline{a}, \overline{b}) \le \deg_T \mathcal{R}_{j+1} = d - j - 1 < \deg_T \mathcal{R}_j(\overline{a}, \overline{b}),$$

we conclude that $\mathcal{R}_{j+1}(\overline{a}, \overline{b})$ is the remainder in the division of r_{j-1} by r_j . In other words, $r_{j+1} = \mathcal{R}_{j+1}(\overline{a}, \overline{b})$, which completes the proof of the first assertion of the lemma. The second assertion is proved with a similar argument.

Let $\overline{a} \in \overline{\mathbb{F}}_q^e$ be the tuple of roots of g in any order, so that $g = S^e(T - \overline{a})$. Let $G_k := F_k(\mathbf{S}(\overline{a}), \mathbf{S}(-\mathbb{B}))$ denote the polynomial obtained by substituting \overline{a} for \mathbb{A} in F_k . Since the set $\mathbf{T} := (S^i(\mathbb{A}) : 1 \le i \le e)$ consists of the first e complete symmetric functions in \mathbb{A} , it follows that $\mathbf{T}(\overline{a})$ belongs to \mathbb{F}_q^e , and thus G_k belongs to $\mathbb{F}_q[\mathbf{S}]$. Further, Proposition 3.1 shows that G_k is a nonzero polynomial with $\deg_{\mathbf{S}} G_k = e - d + k$, which is monic in $S^k(-\mathbb{B})$ with $\deg_{S^k(-\mathbb{B})} G_k = e - d + k$.

We end this section with a result which will be crucial to establish lower bounds for the average-case complexity of the Euclidean algorithm. As we shall see in the next section, for a fixed $g \in \mathbb{F}_q[T]$ with deg g = e, a random element $f \in \mathbb{F}_q[T]$ with deg f = d and g are relatively prime with high probability. In this sense, we call a polynomial $f \in \mathbb{F}_q[T]$ with deg f = d generic (with respect to g) if the remainder sequence in the Euclidean algorithm applied to the pair (g, f) has length d. In particular, in such a remainder sequence (r_1, \ldots, r_d) we have deg $(r_k) = d - k$ for $1 \le k \le d$. The next result establishes a lower bound on the number generic monic elements in $\mathbb{F}_q[T]$ of degree d.

Proposition 3.3. Let $\mathcal{G} \subset \mathbb{F}_q[T]$ be the set of monic elements of degree d which are generic in the sense above. Then

$$|\mathcal{G}| \ge q^d \left(1 - \frac{d(2e - d + 1)}{2q}\right).$$

In particular, for q > d(2e - d + 1)/2 the set \mathcal{G} is nonempty.

Proof. Let $f := T^d + s_1 T^{d-1} + \dots + s_d \in \mathcal{G}$ and let (r_1, \dots, r_d) be the sequence of remainders in the Euclidean algorithm applied to the pair (g, f). By hypothesis $\deg(r_j) = d - j$ for $1 \leq j \leq d$, which by Lemma 3.2 is equivalent to the condition $G_j(s_1, \dots, s_d) \neq 0$ for $1 \leq j \leq d$. It follows that

$$\mathcal{G} = \bigcap_{j=1}^{d} \left(\mathbb{F}_{q}^{d} \setminus \mathcal{V}(G_{j})(\mathbb{F}_{q}) \right) = \mathbb{F}_{q}^{d} \setminus \bigcup_{j=1}^{d} \mathcal{V}(G_{j})(\mathbb{F}_{q})$$

As a consequence,

$$|\mathcal{G}| = q^d - \left| \bigcup_{j=1}^d \mathcal{V}(G_j)(\mathbb{F}_q) \right| \ge q^d - \sum_{j=1}^d |\mathcal{V}(G_j)(\mathbb{F}_q)|.$$

According to (2.1), we have

$$\sum_{k=1}^{d} |\mathcal{V}(G_k)(\mathbb{F}_q)| \le q^{d-1} \sum_{k=1}^{d} (e-d+k) = q^{d-1} \frac{d(2e-d+1)}{2}$$

which readily implies the proposition.

Let e, d be positive integers with e > d. For any $m \ge 0$, we denote by $\mathbb{F}_q[T]_m$ the set of monic polynomials of degree m with coefficients in \mathbb{F}_q . For a fixed $g \in \mathbb{F}_q[T]_e$, let $\mathcal{X}_g : \mathbb{F}_q[T]_d \to \{0, \ldots, d\}, \ \mathcal{X}_g(f) = \deg(\gcd(g, f))$ be the random variable defined by the degree of the greatest common divisor $\gcd(g, f)$, where $\mathbb{F}_q[T]_d$ is endowed with the uniform probability. Applying the Euclidean algorithm to a pair (g, f) with $f \in \mathbb{F}_q[T]_d$ we obtain a positive integer k with $1 \le k \le d$, a unique polynomial quotient sequence (q_1, \ldots, q_{k+1}) and a unique polynomial remainder sequence (r_1, \ldots, r_k) , satisfying the following conditions:

$$g = f \cdot q_1 + r_1, \qquad \deg(r_1) < \deg(f), \\ f = r_1 \cdot q_2 + r_2, \qquad \deg(r_2) < \deg(r_1), \\ \vdots \qquad \vdots \\ r_{k-2} = r_{k-1} \cdot q_k + r_k, \qquad \deg(r_k) < \deg(r_{k-1}), \\ r_{k-1} = r_k \cdot q_{k+1}.$$

First we study the average degree of the gcd, namely the expected value of \mathcal{X}_q :

(4.1)
$$E[\mathcal{X}_g] = \sum_{i=0}^d i \, \frac{|B_i|}{q^d} = \sum_{i=1}^d i \, \frac{|B_i|}{q^d} = \sum_{i=1}^d \sum_{j=i}^d \frac{|B_j|}{q^d},$$

where $B_i := \{ f \in \mathbb{F}_q[T]_d : \mathcal{X}_g(f) = i \}$ for $0 \le i \le d$.

r

We start with an estimate on $\sum_{j=1}^{d} |B_j| = |\bigcup_{j=1}^{d} B_j|$. For this purpose, observe that

$$\bigcup_{j=1}^{d} B_j = \{ f \in \mathbb{F}_q[T]_d : \operatorname{res}(g, f) = 0 \},\$$

where $\operatorname{res}(\cdot, \cdot)$ denotes resultant. We recall that g has factorization pattern $(\lambda_1, \ldots, \lambda_e) \in \mathbb{Z}_{\geq 0}^e$, with $\lambda_1 + 2\lambda_2 + \cdots + e\lambda_e = e$, if g has λ_i irreducible factors in $\mathbb{F}_q[T]$ of degree i (counting multiplicities) for $1 \leq i \leq e$. We shall also consider the *reduced* factorization pattern $(\lambda_1^*, \ldots, \lambda_e^*) \in \mathbb{Z}_{\geq 0}^e$ of g, where λ_i^* is the number of distinct irreducible factors of g in $\mathbb{F}_q[T]_i$ for $1 \leq i \leq e$, and denote by g^* the square-free part of g, namely the product of all distinct irreducible factors of g (without multiplicities). In particular, we have that $(\lambda_1^*, \ldots, \lambda_e^*)$ is the factorization pattern of g^* . We have the following result.

Proposition 4.1. Let e, d be integers with e > d > 0. Let g be an element of $\mathbb{F}_q[T]_e, g^*$ its square-free part and $(\lambda_1^*, \ldots, \lambda_e^*)$ the factorization pattern of g^* . Let k be the least integer with $\lambda_k^* > 0$. If $k \leq d$, then

$$\lambda_k^* q^{d-k} - \binom{\lambda_k^*}{2} q^{\max\{d-2k,0\}} \le \left| \bigcup_{j=1}^d B_j \right| \le \lambda_k^* q^{d-k} + \sum_{i=k+1}^d \lambda_i^* q^{d-i}.$$

Proof. For $f \in \mathbb{F}_q[T]_d$ we have $\operatorname{res}(g, f) = 0$ if and only if $\operatorname{res}(g^*, f) = 0$. As a consequence, we shall consider the resultant $\operatorname{res}(g^*, f)$. Denote by $g_i := \prod_{j=1}^{\lambda_i^*} g_{i,j}$ the product of all irreducible factors of g^* of degree i for $1 \leq i \leq e$. Let $\mathbf{S} := (S_{d-1}, \ldots, S_0)$ be a vector of indeterminates and

$$F(\mathbf{S},T) := T^d + S_{d-1}T^{d-1} + \dots + S_0.$$

The product formula for the resultant (see, e.g., [BPR06, Theorem 4.16]) implies

$$\operatorname{res}(g^*, F(\boldsymbol{S}, T)) = \prod_{i=k}^e \mathsf{R}_i := \prod_{i=k}^e \operatorname{res}(g_i, F(\boldsymbol{S}, T)) = \prod_{i=k}^e \prod_{j=1}^{\lambda_i^*} \operatorname{res}(g_{i,j}, F(\boldsymbol{S}, T)).$$

Now, for any i with $k \leq i \leq d$ and $\lambda_i^* > 0$, we have

$$\mathsf{R}_i := \prod_{j=1}^{\lambda_i^*} \mathsf{R}_{i,j}, \quad \mathsf{R}_{i,j} := \operatorname{res}(g_{i,j}, F(\boldsymbol{S}, T)).$$

Since $g_{i,j}$ is an irreducible element of $\mathbb{F}_q[T]$, for $\boldsymbol{s} \in \mathbb{F}_q^d$ we have $\mathsf{R}_{i,j}(\boldsymbol{s}) = 0$ if and only if $g_{i,j}$ divides $F(\boldsymbol{s},T)$. Further, as $\{F(\boldsymbol{s},T): \boldsymbol{s} \in \mathbb{F}_q^d\} \subset \mathbb{F}_q[T]_d$, we conclude that there is a bijection between the set of \mathbb{F}_q -rational zeros of $\mathsf{R}_{i,j}$ and the set of multiples in $\mathbb{F}_q[T]_d$ of $g_{i,j}$. As the latter has cardinality q^{d-i} , we conclude that $|\mathcal{V}(\mathsf{R}_{i,j})(\mathbb{F}_q)| = q^{d-i}$. Therefore,

$$|\mathcal{V}(\mathsf{R}_i)(\mathbb{F}_q)| = \left| \bigcup_{j=1}^{\lambda_i^*} \mathcal{V}(\mathsf{R}_{i,j})(\mathbb{F}_q) \right| \le \lambda_i^* q^{d-i}.$$

On the other hand, for i > d with $\lambda_i^* > 0$, there is no element of $\mathbb{F}_q[T]_d$ having a nontrivial common factor with g_i defined over \mathbb{F}_q . This implies that the set $\mathcal{V}(\mathsf{R}_i)(\mathbb{F}_q)$ is empty, namely

$$|\mathcal{V}(\mathsf{R}_i)(\mathbb{F}_q)| = 0.$$

Now we focuss on the case i = k. If $\mathsf{R}_k := \prod_{j=1}^{\lambda_k^*} \mathsf{R}_{k,j}$, we have

$$|\mathcal{V}(\mathsf{R}_k)(\mathbb{F}_q)| \le |\mathcal{V}(\operatorname{res}(g, F(\boldsymbol{S}, T)))(\mathbb{F}_q)| \le |\mathcal{V}(\mathsf{R}_k)(\mathbb{F}_q)| + \sum_{i=k+1}^d \lambda_i^* q^{d-i}$$

Our previous argument shows that $\mathcal{V}(\mathsf{R}_k)(\mathbb{F}_q)$ is a union of λ_k^* sets $\mathcal{V}(\mathsf{R}_{k,j})(\mathbb{F}_q)$ of cardinality q^{d-k} , which are pairwise distinct. Further, $\mathbf{s} \in \mathcal{V}(\mathsf{R}_{k,j_1})(\mathbb{F}_q) \cap \mathcal{V}(\mathsf{R}_{k,j_2})(\mathbb{F}_q)$ for $j_1 \neq j_2$ if and only if both g_{k,j_1} and g_{k,j_2} divide $F(\mathbf{s},T)$. As g_{k,j_1} and g_{k,j_2} are two distinct irreducible elements of $\mathbb{F}_q[T]$, this holds if and only if $g_{k,j_1} \cdot g_{k,j_2}$ divides $F(\mathbf{s},T)$. It follows that

$$\mathcal{V}(\mathsf{R}_{k,j_1})(\mathbb{F}_q) \cap \mathcal{V}(\mathsf{R}_{k,j_2})(\mathbb{F}_q)| = \begin{cases} q^{d-2k} & \text{for } d \ge 2k, \\ 0 & \text{for } d < 2k. \end{cases}$$

In particular, the Bonferroni inequalities imply

$$\lambda_k^* q^{d-k} - \binom{\lambda_k^*}{2} q^{\max\{d-2k,0\}} \le |\mathcal{V}(\mathsf{R}_k)(\mathbb{F}_q)| = \left| \bigcup_{j=1}^{\lambda_k^*} \mathcal{V}(\mathsf{R}_{k,j})(\mathbb{F}_q) \right| \le \lambda_k^* q^{d-k}.$$

From this the statement of the proposition readily follows.

As an immediate consequence of Proposition 4.1 we obtain an estimate on the probability that a random element of $\mathbb{F}_q[T]_d$ is relatively prime with g.

Theorem 4.2. Let e, d be integers with e > d > 0. Let g be an element of $\mathbb{F}_q[T]_e$, g^* its square-free part and $(\lambda_1^*, \ldots, \lambda_e^*)$ the factorization pattern of g^* . Let k be the least integer with $\lambda_k^* > 0$. If $k \leq d$, then the probability $\mathcal{P}_0 := |B_0|/q^d$ that a random element $f \in \mathbb{F}_q[T]_d$ and g are relatively prime is bounded in the following way:

$$1 - \frac{\lambda_k^*}{q^k} - \sum_{i=k+1}^d \frac{\lambda_i^*}{q^i} \le \mathcal{P}_0 \le 1 - \frac{\lambda_k^*}{q^k} + \binom{\lambda_k^*}{2} \frac{1}{q^{\min\{2k,d\}}}.$$

In particular, for q > 2e we have $\mathcal{P}_0 > \frac{1}{2}$.

Proof. Observe that

$$|B_0| = \left| \{f \in \mathbb{F}_q[T]_d : \gcd(g, f) = 1\} \right| = \left| \mathbb{F}_q^d \setminus \bigcup_{j=1}^d B_j \right| = q^d - \left| \bigcup_{j=1}^d B_j \right|.$$

Then the statement readily follows from Proposition 4.1.

10

If the square-free part g^* of $g \in \mathbb{F}_q[T]_e$ has a factorization pattern $(\lambda_1^*, \ldots, \lambda_e^*)$ as in Theorem 4.2, then all its irreducible factors have degree at least k. It follows that $B_1 \cup \cdots \cup B_{k-1}$ is the empty set, which implies the following corollary.

Corollary 4.3. With hypotheses as in Theorem 4.2, for $1 \le i \le k$, we have

$$\lambda_k^* q^{d-k} - \binom{\lambda_k^*}{2} q^{\max\{d-2k,0\}} \le \left| \bigcup_{j=i}^d B_j \right| \le \lambda_k^* q^{d-k} + \sum_{j=k+1}^d \lambda_j^* q^{d-j}$$

Next we bound the sum of the cardinalities of $\bigcup_{j=i}^{d} B_j$ for $i \ge k+1$.

Proposition 4.4. Let $g \in \mathbb{F}_q[T]_e$ have a factorization pattern $(\lambda_1, \ldots, \lambda_e)$ and let k be the least index with $\lambda_k > 0$. We have

$$\sum_{i=k+1}^{d} \left| \bigcup_{j=i}^{d} B_j \right| \leq \sum_{i=k+1}^{d} (i-k) q^{d-i} \sum_{\substack{h_k \leq \lambda_k, \dots, h_i \leq \lambda_i \\ k h_k + \dots + i h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i}.$$

Proof. Observe that

$$B_i \cup \cdots \cup B_d = \{ f \in \mathbb{F}_q[T]_d : \deg \gcd(g, f) \ge i \}.$$

Fix a factor $m \in \mathbb{F}_q[T]_j$ of degree $j \geq i$ of g. Then the set L_m of multiples $f \in \mathbb{F}_q[T]_d$ of m has cardinality $|L_m| = q^{d-j}$. As a consequence, letting m vary over the set of factors in $\mathbb{F}_q[T]_j$ of g we conclude that

$$\left| \bigcup_{j=i}^{d} B_j \right| \le \sum_{j=i}^{d} \eta_j \, q^{d-j},$$

where η_j is the number of distinct factors of g in $\mathbb{F}_q[T]_j$ for $i \leq j \leq d$. It follows that

$$\sum_{i=k+1}^{d} \left| \bigcup_{j=i}^{d} B_j \right| \le \sum_{i=k+1}^{d} (i-k) \eta_i q^{d-i}.$$

It remains to express the η_i in terms of $\lambda_1, \ldots, \lambda_d$. For this purpose, we observe that

$$\eta_i \leq [X^i] \left(\prod_{j=k}^i (1+X^j)^{\lambda_j} \right) = \sum_{\substack{h_k \leq \lambda_k, \dots, h_i \leq \lambda_i \\ k \, h_k + \dots + i \, h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i},$$

where $[X^i]f$ denotes the coefficient of X^i in the monomial expansion of $f \in \mathbb{K}[X]$. This proves the proposition.

Now we obtain an estimate for the average degree of gcd(g, f) for random $f \in \mathbb{F}_q[T]_d$.

Theorem 4.5. Let e, d be integers with e > d > 0, g an element of $\mathbb{F}_q[T]_e$ with factorization pattern $(\lambda_1, \ldots, \lambda_e)$ and k the least index with $\lambda_k > 0$. Denote by λ_k^* the number of distinct irreducible factors of g in $\mathbb{F}_q[T]_k$. If $k \leq d$, then the average degree $E[\mathcal{X}_g]$ of the greatest common divisor of g and a random element f of $\mathbb{F}_q[T]_d$ is bounded in the following way:

$$\frac{k\,\lambda_k^*}{q^k} - \binom{\lambda_k^*}{2} \frac{k}{q^{\min\{2k,d\}}} \le E[\mathcal{X}_g] \le \frac{k\,\lambda_k^*}{q^k} + \sum_{i=k+1}^d \frac{i}{q^i} \sum_{\substack{h_k \le \lambda_k, \dots, h_i \le \lambda_i \\ k\,h_k + \dots + i\,h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i}.$$

Proof. According to (4.1),

$$E[\mathcal{X}_g] = \sum_{i=1}^d \sum_{j=i}^d \frac{|B_j|}{q^d}.$$

By Proposition 4.1 and Corollary 4.3, for $1 \le i \le k$,

$$\lambda_k^* q^{d-k} - \binom{\lambda_k^*}{2} q^{\max\{d-2k,0\}} \le \left| \bigcup_{j=i}^d B_j \right| \le \lambda_k^* q^{d-k} + \sum_{i=k+1}^d \lambda_i q^{d-i}.$$

By Proposition 4.4, we have

$$\sum_{i=k+1}^{d} \left| \bigcup_{j=i}^{d} B_j \right| \leq \sum_{i=k+1}^{d} (i-k) q^{d-i} \sum_{\substack{h_k \leq \lambda_k, \dots, h_i \leq \lambda_i \\ k h_k + \dots + i h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i}.$$

We conclude that

$$\frac{k\,\lambda_k^*}{q^k} - \binom{\lambda_k^*}{2} \frac{k}{q^{\min\{2k,d\}}} \le E[\mathcal{X}_g] \le \frac{k\,\lambda_k^*}{q^k} + \sum_{i=k+1}^d \frac{k\,\lambda_i}{q^i} + \sum_{\substack{i=k+1\\i=k+1}}^d \frac{i-k}{q^i} \sum_{\substack{h_k \le \lambda_k, \dots, h_i \le \lambda_i\\k\,h_k + \dots + i\,h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i} \\ \le \frac{k\,\lambda_k^*}{q^k} + \sum_{i=k+1}^d \frac{i}{q^i} \sum_{\substack{h_k \le \lambda_k, \dots, h_i \le \lambda_i\\k\,h_k + \dots + i\,h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i},$$

which proves the theorem.

To simplify the upper bound of Theorem 4.5 we recall that the inner sum in such an upper bound is actually an upper bound for the number η_i of distinct factors of g in $\mathbb{F}_q[T]_i$, namely

$$\eta_i \leq [X^i] \left(\prod_{j=k}^i (1+X^j)^{\lambda_j} \right) = \sum_{\substack{h_k \leq \lambda_k, \dots, h_i \leq \lambda_i \\ k \, h_k + \dots + i \, h_i = i}} \binom{\lambda_k}{h_k} \cdots \binom{\lambda_i}{h_i},$$

with equality when g is square-free. Using the generalized Vandermonde identity (see, e.g., [GKP94, page 248]), we have

(4.2)
$$\eta_i \leq \sum_{\substack{h_k \leq \lambda_k, \dots, h_i \leq \lambda_i \\ k h_k + \dots + i h_i = i}} \binom{k \lambda_k}{k h_k} \cdots \binom{i \lambda_i}{i h_i} \leq \binom{k \lambda_k + \dots + i \lambda_i}{i}.$$

On the other hand, taking into account that the expansion of the analytic function $h : \mathbb{C} \to \mathbb{C}$, $h(z) := \prod_{j=k}^{i} (1+z^j)^{\lambda_j}$ has non-negative coefficients at 0, from, e.g., [FS09, Proposition IV.1] we conclude that

(4.3)
$$\eta_i \le h(1) = 2^{\lambda_k + \dots + \lambda_i}.$$

The accuracy of (4.2) and (4.3) depends on the actual factorization pattern $(\lambda_1, \ldots, \lambda_e)$. For example, if $g \in \mathbb{F}_q[T]_e$ is a polynomial with an "equal-degree factorization" (that is, $k\lambda_k = e$), then for large k the bound (4.3) is preferable, while for large λ_k the bound (4.2) is more accurate.

Finally, for the results on the average-case complexity of the Euclidean algorithm we shall use a further upper bound on $E[\mathcal{X}_g]$. This bound, although not as precise as the one of Theorem 4.5, has a simple expression which suffices for the purposes of the next section.

Lemma 4.6. Let e, d be integers with e > d > 0 and let $g \in \mathbb{F}_q[T]_e$. Then

$$E[\mathcal{X}_g] \le \frac{de}{q^k}.$$

Proof. Let $(\lambda_1, \ldots, \lambda_e)$ be the factorization pattern of g and let k be the least index with $\lambda_k > 0$. By Proposition 4.1,

$$E[\mathcal{X}_g] = \sum_{k=1}^d \sum_{j=k}^d \frac{|B_j|}{q^d} \le \sum_{k=1}^d \sum_{j=1}^d \frac{|B_j|}{q^d} \le d\left(\frac{\lambda_k}{q^k} + \sum_{i=k+1}^d \frac{\lambda_i}{q^i}\right) \le \frac{de}{q^k}.$$

5. Average-case analysis of the Euclidean Algorithm

Let e, d be positive integers with q > d(2e - d + 1)/2 and e > d and let $g \in \mathbb{F}_q[T]_e$ be fixed. In this section we analyze the average-case complexity of the Euclidean algorithm applied to pairs (g, f) with $f \in \mathbb{F}_q[T]_d$.

Given positive integers m, n with m > n and $(f_1, f_2) \in \mathbb{F}_q[T]_m \times (\mathbb{F}_q[T]_n \setminus \{0\})$, and an arithmetic operation $\mathsf{w} \in \{\div, -, \times\}$, by $d^{\mathsf{w}}(f_1, f_2)$ we denote the number of operations w used in the "synthetic" polynomial division algorithm applied to (f_1, f_2) (see, e.g., [Knu81]). It turns out that

(5.1)
$$d^{\div}(f_1, f_2) = m - n + 1, \quad d^{-, \times}(f_1, f_2) = n(m - n + 1).$$

Endowing $\mathbb{F}_q[T]_d$ with the uniform probability, for any $\mathsf{w} \in \{\div, -, \times\}$ we consider the random variable $t_g^{\mathsf{w}} : \mathbb{F}_q[T]_d \to \mathbb{N}$ which counts the number of operations w that the Euclidean Algorithm performs on input (g, f) for each $f \in \mathbb{F}_q[T]_d$. Furthermore, $t_g^{\mathsf{div}}(f)$ denotes the number of polynomial divisions involved. Our aim is to study the expected value $E[t_g^{\mathsf{w}}]$ of t_g^{w} for $\mathsf{w} \in \{\div, \mathsf{div}, -, \times\}$, namely

$$E[t_g^{\sf w}] = \frac{1}{q^d} \sum_{f \in \mathbb{F}_q[T]_d} t_g^{\sf w}(f) = \frac{1}{q^d} \sum_{k=0}^d \sum_{f \in B_{d-k}} t_g^{\sf w}(f).$$

As explained before, applying the Euclidean algorithm to an input (g, f) with $f \in \mathbb{F}_q[T]_d$ we obtain a unique polynomial quotient sequence (q_1, \ldots, q_{h+1}) and a unique polynomial remainder sequence (r_1, \ldots, r_h) satisfying the following conditions:

(5.2)

$$\begin{array}{rcl}
g &=& f \cdot q_1 + r_1, & \deg(r_1) &< \deg(f), \\
f &=& r_1 \cdot q_2 + r_2, & \deg(r_2) &< \deg(r_1), \\
\vdots & & \vdots \\
r_{h-2} &=& r_{h-1} \cdot q_h + r_h, & \deg(r_h) &< \deg(r_{h-1}), \\
r_{h-1} &=& r_h \cdot q_{h+1}.
\end{array}$$

We first consider w = div.

Lemma 5.1. The average number $E[t_g^{\text{div}}]$ of polynomial divisions performed by the Euclidean algorithm applied to pairs (g, f) with $f \in \mathbb{F}_q[T]_d$ is bounded as follows:

$$(d+1)\left(1-\frac{d(2e-d+1)}{2q}\right) \le E[t_g^{\mathsf{div}}] \le (d+1)\left(1+\frac{de}{q}\right).$$

Proof. For $f \in B_{d-k}$ with $0 \le k \le d$, we claim that $t_g^{\text{div}}(f) \le k+1$. Indeed, the maximum number of polynomial divisions in (5.2) is achieved from a sequence of remainders of maximum length. Since $f \in B_{d-k}$, in such a sequence the degree of each successive remainder decreases by 1, that is, the sequence has length h = k. Taking into account that there is a further division to perform, to check that r_h divides r_{h-1} , we deduce our claim. As $k \mapsto \frac{k+1}{d-k}$ is an increasing function for $k \in [0, d-1]$, we obtain

$$E[t_g^{\mathsf{div}}] \le \frac{1}{q^d} \sum_{k=0}^d \sum_{f \in B_{d-k}} (k+1) = \frac{1}{q^d} \left(\sum_{k=0}^{d-1} \frac{k+1}{d-k} (d-k) |B_{d-k}| + (d+1) |B_0| \right)$$
$$\le \frac{d}{q^d} \sum_{k=0}^{d-1} (d-k) |B_{d-k}| + (d+1) \frac{|B_0|}{q^d} \le d E[\mathcal{X}_g] + d + 1.$$

Using the bound $E[\mathcal{X}_g] \leq de/q$ of Lemma 4.6, we deduce the upper bound in the statement of the lemma.

Next we show the lower bound. Recall that $f \in \mathbb{F}_q[T]_d$ is generic (with respect to g) if the corresponding remainder sequence is of the form (r_1, \ldots, r_d) , where $\deg(r_j) = d - j$ for $1 \leq j \leq d$. For such an f, the number of polynomial divisions is precisely d + 1. By Proposition 3.3, it follows that

$$E[t_g^{\mathsf{div}}] \ge \frac{1}{q^d}(d+1)|\mathcal{G}| \ge (d+1)\left(1 - \frac{d(2e-d+1)}{2q}\right).$$

This finishes the proof of the lemma.

Next we analyze the case $w = \div$.

Lemma 5.2. Denote by $E[t_{\hat{g}}^{\pm}]$ the average number of divisions performed by the Euclidean algorithm applied to pairs (g, f) with $f \in \mathbb{F}_q[T]_d$. Then

$$(e+d+1)\left(1 - \frac{d(2e-d+1)}{2q}\right) \le E[t_g^{\div}] \le (e+d+1)\left(1 + \frac{de}{q}\right).$$

Proof. Let $f \in B_{d-k}$ with $0 \le k \le d$. According to (5.1), the number of operations \div in each step of (5.2) is

$$d^{\div}(g, f) = \deg(g) - \deg(f) + 1,$$

$$d^{\div}(f, r_1) = \deg(f) - \deg(r_1) + 1,$$

$$\vdots$$

$$d^{\div}(r_{h-1}, r_h), = \deg(r_{h-1}) - \deg(r_h) + 1.$$

Therefore,

(5.3)
$$t_g^{\div}(f) = \deg(g) - \deg(r_h) + h + 1 = e - (d - k) + h + 1 \le e - d + 2k + 1.$$

As $k \mapsto \frac{e - d + 2k + 1}{d - k}$ is increasing for $k \in [0, d - 1]$, from (5.3) we deduce that

$$\begin{split} E[t_g^{\div}] &= \frac{1}{q^d} \sum_{k=0}^d \sum_{f \in B_{d-k}} t_g^{\div}(f) \le \frac{1}{q^d} \sum_{k=0}^{d-1} \frac{e-d+2k+1}{d-k} (d-k) |B_{d-k}| + (e+d+1) \frac{|B_0|}{q^d} \\ &\le (e+d-1) E[\mathcal{X}_g] + e+d+1. \end{split}$$

Combining this with Lemma 4.6 readily implies the upper bound.

To prove the lower bound, we argue as in the proof of Lemma 5.1. For a generic $f \in \mathcal{G}$, the remainder sequence is of length d, and therefore $t_q^{\div}(f) = e + d + 1$. It follows that

$$E[t_g^{\div}] \ge \frac{1}{q^d}(e+d+1)|\mathcal{G}| \ge (e+d+1)\left(1 - \frac{d(2e-d+1)}{2q}\right)$$

This proves the lemma.

Finally, we consider the remaining case $w \in \{-, \times\}$. We have the following result.

Lemma 5.3. Let $E[t_g^{-,\times}]$ be the average number of operations $w \in \{-,\times\}$ performed by the Euclidean algorithm applied to pairs (g, f) with $f \in \mathbb{F}_q[T]_d$. Then

$$de\left(1 - \frac{d(2e - d + 1)}{2q}\right) \le E[t_g^{-,\times}] \le de\left(1 + \frac{de}{q}\right).$$

Proof. For $f \in B_{d-k}$ with $0 \le k \le d$, by (5.1) the number of operations d^{w} with $\mathsf{w} \in \{-, \times\}$ in each step of (5.2) is

$$d^{\mathsf{w}}(g, f) = \deg(f)(\deg(g) - \deg(f) + 1),$$

$$d^{\mathsf{w}}(f, r_1) = \deg(r_1)(\deg(f) - \deg(r_1) + 1),$$

$$\vdots$$

$$d^{\mathsf{w}}(r_{h-1}, r_h) = \deg(r_h)(\deg(r_{h-1}) - \deg(r_h) + 1),$$

Denote $r_0 := f$. We claim that the maximum number of operations w performed in the whole Euclidean algorithm is achieved with a sequence of remainders (r_0, \ldots, r_k) with $\deg(r_{j-1}) - \deg(r_j) = 1$ for $1 \le j \le k$. Indeed, let (r_0, \ldots, r_h) be a remainder sequence such that $\deg(r_{j-1}) - \deg(r_j) > 1$ for a given j. Denote by $(\alpha_0, \ldots, \alpha_h)$ the corresponding sequence of degrees. We compare the number of operations w performed by the Euclidean algorithm to obtain this sequence with that of a remainder sequence with degree pattern $(\alpha_0, \ldots, \alpha_{j-1}, \alpha_j^*, \alpha_j, \ldots, \alpha_h)$, where $\alpha_{j-1} - \alpha_j^* = 1$. Since the number of w operations is determined by the degree pattern of the remainder sequence under consideration, it suffices to compare the cost of the jth step of the first sequence with the sum of those of the jth and (j + 1)th steps of the second sequence. In particular, we see that our claim for this case holds provided that

$$\alpha_j ((\alpha_{j-1} - \alpha_j) + 1) \le \alpha_j^* (\alpha_{j-1} - \alpha_j^* + 1) + \alpha_j (\alpha_j^* - \alpha_j + 1).$$

This can be checked by an easy calculation. Arguing successively in this way, the claim follows.

As a consequence, the maximum number of operations w performed is achieved in a sequence of k remainders (r_1, \ldots, r_k) with $\deg(r_{j-1}) - \deg(r_j) = 1$ for $1 \le j \le k$, namely with $\deg(r_j) = d - j$ for $1 \le j \le k$. It follows that

(5.4)
$$t_g^{\mathsf{w}}(f) \le \deg(f)(\deg(g) - \deg(f) + 1) + \sum_{j=1}^k \deg(r_j)(\deg(r_{j-1}) - \deg(r_j) + 1) = d(e - d + 1) + 2\sum_{j=1}^k (d - j) = d(e - d + 1) + k(2d - k - 1).$$

Since $k \mapsto \frac{d(e-d+1)+k(2d-k-1)}{d-k}$ is increasing for $k \in [0, d-1]$, by (5.4) we obtain

$$E[t_g^{\mathbf{w}}] = \frac{1}{q^d} \sum_{k=0}^d \sum_{f \in B_{d-k}} t_g^{\mathbf{w}}(f)$$

$$\leq \frac{1}{q^d} \sum_{k=0}^{d-1} \frac{d(e-d+1) + k(2d-k-1)}{d-k} (d-k) |B_{d-k}| + de \frac{|B_0|}{q^d}$$

$$\leq de E[\mathcal{X}_g] + de.$$

The upper bound follows easily by Theorem 4.5.

On the other hand, for $f \in \mathcal{G}$, by (5.4) we conclude that $t_q^{\mathsf{w}}(f) = de$. Then we have

$$E[t_g^{\div}] \ge \frac{1}{q^d} de|\mathcal{G}| \ge de\left(1 - \frac{d(2e - d + 1)}{2q}\right),$$

which finishes the proof of the lemma.

Summarizing Lemmas 5.1, 5.2 and 5.3, we have the following result.

Theorem 5.4. Let e, d be positive integers such that q > d(2e - d + 1)/2 and e > d. Let $g \in \mathbb{F}_q[T]_e$ and $w \in \{\div, \operatorname{div}, -, \times\}$. The average cost $E[t_g^w]$ of operations w performed on (uniform distributed) inputs from $\mathbb{F}_q[T]_d$ is bounded in the following way:

$$\left|\frac{E[t_g^{\mathsf{div}}]}{d+1} - 1\right| \le \frac{de}{q}, \qquad \left|\frac{E[t_g^{\div}]}{e+d+1} - 1\right| \le \frac{de}{q}, \qquad \left|\frac{E[t_g^{-,\times}]}{de} - 1\right| \le \frac{de}{q}.$$

6. Simulations on test examples

In this section we report on the simulations we made with the software package Maple. More precisely, for given values of q, e and d with e > d, we executed the Euclidean algorithm on pairs (g, f), where $g \in \mathbb{F}_q[T]_e$ was a fixed polynomial with factorization pattern $(\lambda_1, \ldots, \lambda_e) \in \mathbb{Z}_{\geq 0}^e$ and f ran through all the elements of a random sample $S \subset$ $\mathbb{F}_q[T]_d$. The aim was to analyze to what extent the results of our simulations behaved as predicted by the theoretical results on the average degree of gcd(g, f) (Theorem 4.5), the probability that gcd(g, f) = 1 (Theorem 4.2) and the probability that a random $f \in \mathbb{F}_q[T]_d$ is "generic" with respect to g (Proposition 3.3).

Recall that, given $g \in \mathbb{F}_q[T]_e$, we denote by $E[\mathcal{X}_g]$ the average degree of gcd(g, f) for f running on all the elements of $\mathbb{F}_q[T]_d$. Further, the probability that gcd(g, f) = 1 is denoted by \mathcal{P}_0 . Finally, we denote by $\mathcal{P}_{\mathcal{G}}$ the probability that $f \in \mathbb{F}_q[T]_d$ is generic with respect to g. According to Theorems 4.5 and 4.2 and Proposition 3.3, if the square-free part g^* of g has factorization pattern $(\lambda_1^*, \ldots, \lambda_e^*)$ and $k \leq d$ is the least index with $\lambda_k^* > 0$, then

$$E[\mathcal{X}_g] \approx \mathbf{E}_g := \frac{k\lambda_k^*}{q^k}, \quad \mathcal{P}_0 \approx \mathbf{P}_0 := 1 - \frac{\lambda_k^*}{q^k}, \quad \mathcal{P}_{\mathcal{G}} \geq \mathbf{P}_{\mathcal{G}} := 1 - \frac{d(2e-d+1)}{2q}$$

The simulations we exhibit were aimed to test whether the right-hand side in the previous expressions approximates the left-hand side on the random samples under consideration. For this purpose, given a random sample $\mathcal{S} \subset \mathbb{F}_q[T]_d$, we computed the sample means

$$\mu := \frac{1}{|\mathcal{S}|} \sum_{f \in \mathcal{S}} \deg \gcd(g, f), \quad \beta := \frac{|B_{0,s}|}{|\mathcal{S}|}, \quad \gamma := \frac{|\mathcal{G}_s|}{|\mathcal{S}|}$$

where $B_{0,S} := \{f \in S : \gcd(g, f) = 1\}$ and $\mathcal{G}_S := \{f \in S : f \text{ is generic}\}$. Furthermore, we considered the corresponding relative errors

$$\varepsilon_1 := \frac{|\mu - \mathbf{E}_g|}{\mathbf{E}_q}, \quad \varepsilon_2 = \frac{|\beta - \mathbf{P}_0|}{\mathbf{P}_0}$$

6.1. Examples for q = 67, e = 7, d = 3 with $\lambda_1^* > 0$. Our first simulations concerned random samples S of 300000 polynomials $f \in \mathbb{F}_{67}[T]$ of degree at most d := 3 and polynomials $g \in \mathbb{F}_{67}[T]$ of degree e := 7 with distinct values of $\lambda_1^* > 0$, listed in the first column of Table 1.

λ_1^*	μ	${\tt E}_g$	β	P_0	γ	$P_{\mathcal{G}}$	ε_1	ε_2
1	0.015273	0.014925	0.984963	0.985075	0.941560	0.731343	0.023312	0.000011
2	0.030590	0.029851	0.969647	0.970149	0.927223	0.731343	0.024756	0.000049
3	0.045027	0.044776	0.956067	0.955224	0.914517	0.731343	0.000561	0.000083
4	0.059633	0.059701	0.942077	0.940299	0.900570	0.731343	0.000114	0.000189
5	0.074163	0.074627	0.928270	0.925373	0.887490	0.731343	0.000622	0.000313
6	0.089130	0.089552	0.914273	0.910448	0.873877	0.731343	0.000471	0.000420
7	0.103867	0.104478	0.900650	0.895522	0.860893	0.731343	0.000585	0.000573

TABLE 1. Examples with q = 67, e = 7 and d = 3.

6.2. Examples for q = 127, e = 9, d = 4 with $\lambda_1^* > 0$. Next we considered random samples S of 10000000 polynomials $f \in \mathbb{F}_{127}[T]$ of degree at most d := 4. We considered polynomials $g \in \mathbb{F}_{127}[T]$ of degree e := 9 with $\lambda_1^* > 0$ distinct roots. The corresponding results are listed in Table 2.

 λ_1^* E_q β ${\tt P}_0$ $P_{\mathcal{G}}$ ε_1 μ γ ε_2 $0.\overline{007874}$ 0.008006 0.9920610.992126 0.9610860.7637790.0243840.000006 1 20.015706 0.0157480.9843570.9842520.9536890.763779 0.000267 0.000011 3 0.0236530.023622 0.946121 0.763779 0.0000160.9765390.9763780.0235910.0314740.031496 0.9688910.9685040.938736 0.7637790.000069 0.0000394 0.039370 0.763779 0.000003 50.0393710.961239 0.960629 0.931346 0.000064 0.047185 0.047244 0.9537430.9527560.924084 0.763779 0.000125 0.0001046 0.055216 0.055118 0.9461350.9448820.916689 0.763779 0.000178 0.00013370.062906 0.062992 0.9370080.909605 0.7637790.000137 0.0001990.9388738 9 0.070742 0.070866 0.931443 0.9291330.902504 0.7637790.000175 0.010884

TABLE 2. Examples with q = 127, e = 9 and d = 4.

6.3. Examples for q = 409, e = 9, d = 4 with $\lambda_1^* > 0$. For our third family of examples we considered random samples S of 10000000 polynomials $f \in \mathbb{F}_{409}[T]$ of degree at most d := 4 and polynomials $g \in \mathbb{F}_{409}[T]$ of degree e := 9 with $\lambda_1^* > 0$. The corresponding results are listed in Table 3.

λ_1^*	μ	E_g	β	P_0	γ	$P_{\mathcal{G}}$	ε_1	ε_2
1	0.002457	0.002445	0.997549	0.997555	0.987825	0.926650	0.000491	0.000001
1	0.002474	0.002445	0.997526	0.997555	0.987809	0.926650	0.011861	0.000003
2	0.004941	0.004889	0.995065	0.995110	0.985414	0.926650	0.010636	0.044678
2	0.004902	0.004889	0.995109	0.995110	0.985340	0.926650	0.000268	$1 \cdot 10^{-6}$
3	0.007329	0.007335	0.992688	0.992665	0.983014	0.926650	0.000082	0.000002
3	0.007309	0.007335	0.992710	0.992665	0.983031	0.926650	0.00035	0.000005
4	0.009772	0.009779	0.99026	0.990220	0.980597	0.926650	0.000072	0.00004
4	0.009761	0.009779	0.990277	0.990220	0.980693	0.926650	0.000184	0.000006
5	0.012245	0.012225	0.987821	0.987776	0.978268	0.926650	0.000164	0.002785
5	0.012241	0.012225	0.987836	0.987776	0.978206	0.926650	0.000131	0.000006
6	0.014649	0.014669	0.985448	0.985331	0.975835	0.926650	0.000136	0.000012
7	0.017035	0.017115	0.983098	0.982885	0.973513	0.926650	0.000467	0.000022
8	0.019524	0.019552	0.980654	0.980440	0.971072	$0.92\overline{6650}$	0.000143	0.000022
9	0.021948	0.022005	0.978267	0.977995	0.968726	0.926650	0.000278	0.000028

TABLE 3. Examples with q = 409, e = 9 and d = 4.

6.4. Examples for q = 67, e = 7, d = 3 with $\lambda_k^* > 0$. Now we report on random samples S of 300000 polynomials $f \in \mathbb{F}_{67}[T]$ of degree at most d := 3, and polynomials $g \in \mathbb{F}_{67}[T]$ of degree e := 7 having different values for the least index k with $\lambda_k^* > 0$. In the first column of Table 4 we show the different values of k considered, while the second column exhibits the corresponding values of λ_k^* . As the sample mean μ and the asymptotic estimates \mathbb{E}_g were close to zero, instead of the relative error we considered the absolute error

$$\varepsilon_1 := |\mu - \mathbf{E}_q|,$$

listed in the ninth column of Table 4.

TABLE 4.	Examples	s with	q = 67	, e = 7	and d	= 3.
----------	----------	--------	--------	---------	---------	------

k	λ_k^*	μ	E_g	β	P ₀	γ	$\mathbb{P}_{\mathcal{G}}$	ε_1	ε_2
2	1	0.000453	0.000445	0.999773	0.999777	0.955343	0.731343	0.000008	0.000004
2	2	0.000836	0.000891	0.999586	0.999555	0.0955466	0.731343	0.000055	0.000003
3	1	0.000030	0.000001	0.999990	0.999997	0.955866	0.731343	0.000029	0.000007
3	1	0.000030	0.000001	0.999990	0.999997	0.956393	0.731343	0.000029	0.000009

6.5. Examples for q = 127, e = 9, d = 4 with $\lambda_k^* > 0$. The next family of examples concerned random samples S of 10000000 polynomials $f \in \mathbb{F}_{127}[T]$ of degree at most d := 3. We considered polynomials $g \in \mathbb{F}_{127}[T]$ of degree e := 9 with different values for the least index k with $\lambda_k^* > 0$. The corresponding results are summarized in Table 5. As the sample mean μ and the asymptotic estimates \mathbb{E}_g were close to zero, we considered the absolute error

$$\varepsilon_1 := |\mu - \mathbf{E}_g|$$

k	λ_k^*	μ	\mathtt{E}_g	β	P_0	γ	$P_{\mathcal{G}}$	ε_1	ε_2
2	1	0.000135	0.000124	0.999932	0.999938	0.968791	0.763779	0.000011	0.000006
2	2	0.000262	0.000248	0.999869	0.999876	0.968705	0.763779	0.000014	0.000007
3	1	0.000006	0.000001	0.999999	0.9999999	0.968915	0.763779	0.000005	0
4	1	0	$2 \cdot 10^{-9}$	1	0.999999	0.968926	0.763779	$2 \cdot 10^{-9}$	0.000001

TABLE 5. Examples with q = 127, e = 9 and d = 4.

6.6. Examples for q = 211, e = 17, d = 7 with $\lambda_k^* > 0$. Now we report on simulations with random samples S of 10000000 polynomials $f \in \mathbb{F}_{211}[T]$ of degree at most d := 3. We considered polynomials $g \in \mathbb{F}_{211}[T]$ of degree e := 17 having different values for the least index k with $\lambda_k^* > 0$. As the sample mean μ and the asymptotic estimates \mathbb{E}_g were close to zero, we considered the absolute error

$$\varepsilon_1 := |\mu - \mathbf{E}_g|$$

The results are listed in Table 6.

TABLE 6. Examples with q = 211, e = 17 and d = 7.

k	λ_k^*	μ	E_g	β	P ₀	γ	$P_{\mathcal{G}}$	ε_1	ε_2
2	2	0.000087	0.000089	0.999956	0.999955	0.967239	0.535545	0.000002	0.000001
2	1	0.000042	0.000045	0.999979	0.999978	0.967284	0.535545	0.000003	0.000001
3	1	0	$3 \cdot 10^{-7}$	1	0.999999	0.967244	0.535545	$3 \cdot 10^{-7}$	0.000001
3	2	0	$6 \cdot 10^{-7}$	1	0.9999999	0.967236	0.535545	$6 \cdot 10^{-7}$	0.000001

6.7. Examples for q = 409, e = 9, d = 4 with $\lambda_k^* > 0$. The last family of examples involves random samples S of 10000000 polynomials $f \in \mathbb{F}_{409}[T]$ of degree at most d := 4. We considered polynomials $g \in \mathbb{F}_{409}[T]$ of degree e := 9 having different values for the least index k with $\lambda_k^* > 0$. As the sample mean μ and the asymptotic estimates \mathbb{E}_g were close to zero, we considered the absolute error

$$\varepsilon_1 := |\mu - \mathbf{E}_g|.$$

The results are exhibited in Table 7.

TABLE 7. Examples with q = 409, e = 9 and d = 4.

k	λ_k^*	μ	${ t E}_g$	β	P_0	γ	$P_{\mathcal{G}}$	ε_1	ε_2
2	2	0.000026	0.000024	0.999987	0.999988	0.931219	0.926650	0.000002	0.000001
2	1	0.000015	0.000002	0.999993	0.999994	0.990212	0.926650	0.000013	0.000001
3	2	0	$8 \cdot 10^{-8}$	1	0.999999	0.946044	0.926650	$8 \cdot 10^{-8}$	0.000001
3	1	0	$4 \cdot 10^{-8}$	1	0.999999	0.946146	0.926650	$4 \cdot 10^{-8}$	0.000001

6.8. Conclusions. Summarizing, the results of Tables 1–7 show that the numerical experiments we performed behave as predicted by the asymptotic estimates of Theorems 4.5 and 4.2. On the other hand, it seems that the estimate on the number generic polynomials of Proposition 3.3 is somewhat pessimistic. Our numerical experiments suggest that the number of generic polynomials depends on the factorization pattern of g, while the lower bound of Proposition 3.3 depends only on q, e and d.

References

- [BNNV14] V. Berthé, H. Nakada, R. Natsui, and B. Vallée, Fine costs for Euclid's algorithm on polynomials and Farey maps, Adv. Appl. Math. 54 (2014), 27–65.
- [BPR06] S. Basu, R. Pollack, and M.-F. Roy, Algorithms in real algebraic geometry, 2nd ed., Algorithms Comput. Math., vol. 10, Springer, Berlin, 2006.
- [CM06] A. Cafure and G. Matera, Improved explicit estimates on the number of solutions of equations over a finite field, Finite Fields Appl. **12** (2006), no. 2, 155–185.
- [FS09] P. Flajolet and R. Sedgewick, Analytic combinatorics, Cambridge Univ. Press, Cambridge, 2009.
 [Ful84] W. Fulton, Intersection theory, Springer, Berlin Heidelberg New York, 1984.
- [GKP94] R. Graham, D. Knuth, and O. Patashnik, *Concrete mathematics: a foundation for computer science*, 2nd ed., Addison–Wesley, Reading, Massachusetts, 1994.
- [Hei83] J. Heintz, Definability and fast quantifier elimination in algebraically closed fields, Theoret. Comput. Sci. 24 (1983), no. 3, 239–277.
- [Knu81] D.E. Knuth, The art of computer programming II: Semi-numerical algorithms, vol. 2, Addison-Wesley, Reading, Massachusetts, 1981.
- [Las03] A. Lascoux, Symmetric functions and combinatorial operators on polynomials, CBMS Reg. Conf. Ser. Math., no. 99, American Mathematical Society, Providence, RI, 2003.
- [LV08] L. Lhote and B. Vallée, Gaussian laws for the main parameters of the Euclid algorithms, Algorithmica 50 (2008), no. 4, 497–554.

[MvzG90] K. Ma and J. von zur Gathen, Analysis of Euclidean algorithms for polynomials over finite fields, J. Symb. Comput. 9 (1990), no. 4, 429–455.

[Nor89] G. Norton, Precise analyses of the right- and left-shift greatest common divisor algorithms for GF(q)/x, SIAM J. Comput. **18** (1989), no. 3, 608–624.

[vzGG99] J. von zur Gathen and J. Gerhard, Modern computer algebra, Cambridge Univ. Press, Cambridge, 1999.

¹ Universidad Nacional de General Sarmiento, Instituto del Desarrollo Humano, J.M. Gutiérrez 1150 (B1613GSX) Los Polvorines, Buenos Aires, Argentina

E-mail address: agimenez@ungs.edu.ar, gmatera@ungs.edu.ar

 2 Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Matemática, Ciudad Universitaria, Pabellón I (1428) Buenos Aires, Argentina

³ CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS (CONICET), ARGENTINA

⁴ UNIVERSIDAD NACIONAL DE HURLINGHAM, INSTITUTO DE TECNOLOGÍA E INGENIERÍA, AV. GDOR. VERGARA 2222 (B1688GEZ), VILLA TESEI, BUENOS AIRES, ARGENTINA *E-mail address*: mariana.perez@unahur.edu.ar

⁵UNIVERSIDAD NACIONAL DE GENERAL SARMIENTO, INSTITUTO DE CIENCIAS, J.M. GUTIÉRREZ 1150 (B1613GSX) LOS POLVORINES, BUENOS AIRES, ARGENTINA *E-mail address:* mprivite@ungs.edu.ar