

The generation of matter-antimatter asymmetries and hypermagnetic fields by the chiral vortical effect of transient fluctuations

S. Abbaslu^{*1}, S. Rostam Zadeh^{†2}, M. Mehraeen^{‡1,3} and S. S. Gousheh^{§1}

¹*Department of Physics, Shahid Beheshti University, Tehran, Iran*

²*School of Particles and Accelerators, Institute for Research in Fundamental Sciences (IPM),
P.O.Box 19395-5531, Tehran, Iran*

³*Department of Physics, Case Western Reserve University, 10900 Euclid Avenue, Cleveland,
Ohio 44106, USA*

February 18, 2022

Abstract: We study the contribution of the temperature-dependent chiral vortical effect to the generation and evolution of hypermagnetic fields and matter-antimatter asymmetries, in the symmetric phase of the early Universe, in the temperature range $100 \text{ GeV} \leq T \leq 10 \text{ TeV}$. Our most important result is that, due to the chiral vortical effect, small overlapping transient fluctuations in the vorticity field in the plasma and temperature of matter degrees of freedom can lead to the generation of strong hypermagnetic fields and matter-antimatter asymmetries, all starting from zero initial values. We show that, either an increase in the amplitudes of the fluctuations of vorticity or temperature, or a decrease in their widths, leads to the production of stronger hypermagnetic fields, and therefore, larger matter-antimatter asymmetries. We have the interesting result that fluctuating vorticity fields are more productive, by many orders of magnitude, as compared to vorticities that are constant in time.

1 Introduction

Anomalous transport effects play important roles in particle physics and cosmology, particularly in the early Universe [1]. One important effect of this kind is the so-called Chiral Vortical Effect (CVE), which refers to the generation of an electric current parallel to the vorticity field in the chiral plasma [2]. This effect was discovered by Vilenkin who showed that a neutrino current density can result from a rotating black hole [2]. He obtained the neutrino current density in the direction of the rotation axis as

$$J(0) = -\frac{\Omega}{12}T^2 - \frac{\Omega}{4\pi^2}\mu^2 - \frac{\Omega^3}{48\pi^2}, \quad (1.1)$$

where Ω is the angular velocity, μ is the chiral chemical potential of the neutrino, and T is its temperature. Thirty years after its discovery, the CVE appeared in the relativistic hydrodynamic equations as an interesting manifestation of anomalies in quantum field theory [3]. This effect

^{*}s_abbasluo@sbu.ac.ir

[†]sh_rostamzadeh@ipm.ir

[‡]mxm1289@case.edu

[§]ss-gousheh@sbu.ac.ir

has attracted much attention and has been investigated extensively in recent years, leading to a deeper understanding of the subject [4–10]. In particular, it has been established that in single species chiral plasma in the broken phase at high temperatures, the CVE shows up in the vector current as $\vec{J}_{\text{cv}} = \frac{1}{4\pi^2}(\mu_R^2 - \mu_L^2)\vec{\Omega}$, and in the axial current as $\vec{J}_{\text{cv}}^5 = \left[\frac{T^2}{6} + \frac{1}{4\pi^2}(\mu_R^2 + \mu_L^2)\right]\vec{\Omega}$, where μ_R and μ_L are the right-handed and the left-handed chemical potentials of the species, respectively [2–10]¹. Interestingly, the term proportional to T^2 indicates that there can be an axial current, even if $\mu_R = \mu_L = 0$.

In this study, we present the correct form of the chiral vortical current in the symmetric phase. Then, we show the prominent effects of the temperature-dependent part of this current in the symmetric phase of the early Universe close to the electroweak phase transition (EWPT). In particular, we show that even very small, but overlapping, transient fluctuations in the vorticity field and temperature of matter degrees of freedom can have important consequences, including the generation of hypermagnetic field in the absence of initial matter asymmetries. The vorticity fluctuations that we consider are about the zero background value, while the temperature fluctuations are about the finite equilibrium temperature of the plasma. The most important role of the CVE in this context is to produce the magnetic fields, either through the chiralities, or through the temperature fluctuation, the latter of which is the main focus of this work. Henceforth, we shall refer to transient fluctuations, which we take to be in the form of short pulses, simply as fluctuations.

Another anomalous transport effect is the chiral magnetic effect (CME), which refers to the generation of an electric current parallel to the magnetic field in the imbalanced chiral plasma [11–14]. It is known that, in single species chiral plasma in the broken phase at high temperatures, the CME appears in the vector current as $\vec{J}_{\text{cm}} = \frac{Q}{4\pi^2}(\mu_R - \mu_L)\vec{B}$, and in the axial current as $\vec{J}_{\text{cm}}^5 = \frac{Q}{4\pi^2}(\mu_R + \mu_L)\vec{B}$, where Q is the electric charge [10, 11, 15, 16] of the species.¹ Later, we will present the correct form of the chiral magnetic current, in the symmetric phase. The chiral magnetic current originating from the electroweak Abelian anomaly, and the chiral vortical current are both non-dissipative currents which can strongly affect the generation and the evolution of the magnetic fields and the matter-antimatter asymmetries in the early Universe [17–21].

Observations clearly show that our Universe is magnetized on all scales [22, 23]. Various models have been proposed to explain the origin of these magnetic fields [20, 24–30], among which the one relying on the electroweak Abelian anomaly has attracted much attention and has been considerably investigated [17–19, 31]. There exists a relationship between the generation and the evolution of the hypermagnetic fields and the fermion number densities in this model, which is due to the chiral coupling of the hypercharge gauge fields to the fermions before the EWPT [27]. In case there is a preexisting asymmetry of the right-handed electrons, their number density is almost conserved far from the EWPT, i.e. $T > 10\text{TeV}$, due to their tiny Yukawa coupling. For lower temperatures, this asymmetry can be converted to the hypermagnetic helicity according to the Abelian anomaly equation, $\partial_\mu j_{e_R}^\mu \sim \vec{E}_Y \cdot \vec{B}_Y$ [17–19, 27, 31, 32]. The anomaly equation shows that, in a reverse process, a strong helical hypermagnetic field can generate the matter-antimatter asymmetries in the Universe, as well [33–35].

Another challenge in particle physics and cosmology is the excess of matter over antimatter, with the measured baryon asymmetry of the Universe being of the order of $\eta_B \sim 10^{-10}$ [36–38]. The three Sakharov conditions² should be satisfied in any CPT invariant model used to explain

¹There are additional contributions of $O(\mu/T)$, where μ is the chemical potential, which are significant at lower temperatures. See Appendix A for details.

²i- baryon number violation, ii- C and CP violation, iii- a departure from thermal equilibrium.

this asymmetry from an initially symmetric Universe [39].

In previous studies based on the electroweak Abelian anomalous model, it has been assumed that there is either a significant amount of matter-antimatter asymmetries to produce the hypermagnetic field, or a strong hypermagnetic field to produce the matter-antimatter asymmetries. The most important result of this study is that the matter-antimatter asymmetries and the hypermagnetic field can all be generated simultaneously from zero initial values, by considering the temperature-dependent CVE before the EWPT. To obtain this interesting result, we take into account the effects of the temperature-dependent term of the chiral vortical current on the evolution of the hypermagnetic fields and the matter-antimatter asymmetries, by considering simultaneous small fluctuations, about the background values, in temperature of the right-handed electrons and the vorticity field, close to the EWPT. We also show that fluctuations in the vorticity field are much more productive than vorticity fields that are constant in time. To be more precise, sharp fluctuations yield results comparable to constant vorticities whose amplitudes are many orders of magnitude larger.

As mentioned above, an underlying assumption of this work is the presence of fluctuations in the plasma of the early Universe, containing all elementary particles and gauge fields. A general description of this plasma in the context of hydrodynamics, as an effective field theory, has been presented [40, 41]. In such a physical system, hydrodynamic variables naturally fluctuate around their statistical averages. Such stochastic and frequent fluctuations are a commonplace in any plasma, including that of the early Universe. These fluctuations, which may stem from a sum of weakly-correlated random local events, can occur for physical observables such as the number density, velocity, and temperature [40, 42–44]. Even though the gauge interactions are strong and their rates are fast in this epoch, the extreme temperatures of the primordial plasma imply that fluctuations could still occur for all species. Moreover, owing to the different Yukawa and gauge couplings, and in particular the chiral nature of the electroweak sector, different chiral fermions could, in principle, experience different fluctuations. All that is required here is that the fluctuations for at least one of the species be non-identical to the rest, at least once.

Another justification for our hypothesis is the following. Realistic fluctuations of density, temperature and vorticity are usually local in space and time. Meanwhile, in a multi-component plasma local and independent density fluctuations can occur for different matter components. The concurrent occurrence of density, temperature and vorticity fluctuations leads to the possibility of occurrence of non-identical fluctuations for different species of particles. To be concrete, consider a local density fluctuation leading to a local excess of a particle species, e.g. e_R , in the central region and rarefaction in the surrounding region, followed immediately by a temperature fluctuation which increases the temperature of the central region. The concurrent occurrence of these two fluctuations leads to a temporary local, and to a much lesser degree global average, increase of temperature of that particle species relative to quasi-equilibrium temperature of the plasma. All that is required now is an overlapping vorticity fluctuation to have the necessary conditions for our model. In this work, we make the simplifying assumption, as is usually done, of considering these fluctuations to be global rather than local. It should be noted that the amplitude and duration of these fluctuations are usually small enough such that the equilibrium energy density, pressure and entropy of the system can be defined. Nevertheless, as we shall show, even very small and brief fluctuations can trigger the mechanism that we propose here and affect significantly the dynamical evolution of the system, as is formulated within the equations of anomalous magnetohydrodynamics (AMHD).

This paper is organized as follows: In Sec. 2, the anomalous magnetohydrodynamics equations and the evolution equations for the matter-antimatter asymmetries are derived in the ex-

panding Universe. In Sec. 3, the set of coupled differential equations are solved numerically. In Sec. 4, the results are summarized and the conclusion is presented.

2 Evolution Equations

2.1 Anomalous Magnetohydrodynamics Equations

In this section, the anomalous magnetohydrodynamics (AMHD) equations are obtained in the Landau-Lifshitz frame in the symmetric phase of the expanding Universe. Taking the CVE and the CME into account, the Maxwell's equations for the hypercharge-neutral plasma in the expanding Universe are given as [3, 20, 45–52]

$$\frac{1}{R}\vec{\nabla}\cdot\vec{E}_Y = 0, \quad \frac{1}{R}\vec{\nabla}\cdot\vec{B}_Y = 0, \quad (2.1)$$

$$\frac{1}{R}\vec{\nabla}\times\vec{E}_Y + \left(\frac{\partial\vec{B}_Y}{\partial t} + 2H\vec{B}_Y\right) = 0, \quad (2.2)$$

$$\begin{aligned} \frac{1}{R}\vec{\nabla}\times\vec{B}_Y - \left(\frac{\partial\vec{E}_Y}{\partial t} + 2H\vec{E}_Y\right) &= \vec{J} \\ &= \vec{J}_{\text{Ohm}} + \vec{J}_{\text{cv}} + \vec{J}_{\text{cm}}, \end{aligned} \quad (2.3)$$

$$\vec{J}_{\text{Ohm}} = \sigma \left(\vec{E}_Y + \vec{v} \times \vec{B}_Y \right), \quad (2.4)$$

$$\vec{J}_{\text{cv}} = c_v \vec{\omega}, \quad (2.5)$$

$$\vec{J}_{\text{cm}} = c_B \vec{B}_Y, \quad (2.6)$$

where R is the scale factor, $H = \dot{R}/R$ is the Hubble parameter, σ is the electrical hyperconductivity of the plasma, and \vec{v} and $\vec{\omega} = \frac{1}{R}\vec{\nabla}\times\vec{v}$ are the bulk velocity and vorticity of the plasma, respectively. Furthermore, the chiral vorticity and helicity coefficients c_v and c_B are as follows (see Appendix A)³,

$$\begin{aligned} c_v(t) &= \sum_{i=1}^{n_G} \left[\frac{g'}{48} \left(-Y_R T_{R_i}^2 + Y_L T_{L_i}^2 N_w - Y_{d_R} T_{d_{R_i}}^2 N_c - Y_{u_R} T_{u_{R_i}}^2 N_c + Y_Q T_{Q_i}^2 N_c N_w \right) \right. \\ &\quad \left. + \frac{g'}{16\pi^2} \left(-Y_R \mu_{R_i}^2 + Y_L \mu_{L_i}^2 N_w - Y_{d_R} \mu_{d_{R_i}}^2 N_c - Y_{u_R} \mu_{u_{R_i}}^2 N_c + Y_Q \mu_{Q_i}^2 N_c N_w \right) \right], \end{aligned} \quad (2.7)$$

$$\begin{aligned} c_B(t) &= -\frac{g'^2}{8\pi^2} \sum_{i=1}^{n_G} \left[-\left(\frac{1}{2}\right) Y_R^2 \mu_{R_i} - \left(\frac{-1}{2}\right) Y_L^2 \mu_{L_i} N_w - \left(\frac{1}{2}\right) Y_{d_R}^2 \mu_{d_{R_i}} N_c - \left(\frac{1}{2}\right) Y_{u_R}^2 \mu_{u_{R_i}} N_c \right. \\ &\quad \left. - \left(\frac{-1}{2}\right) Y_Q^2 \mu_{Q_i} N_c N_w \right], \end{aligned} \quad (2.8)$$

³The temperature-independent parts of these coefficients were presented in [20]. There are additional terms of $O(\mu/T)$ which, as we shall show explicitly in Sec. 3, are negligible within the initial conditions and results of our model (see Appendix A).

where n_G is the number of generations, and $N_c = 3$ and $N_w = 2$ are the ranks of the non-Abelian $SU(3)$ and $SU(2)$ gauge groups, respectively. Moreover, μ_{L_i} (μ_{R_i}), μ_{Q_i} , and $\mu_{u_{R_i}}$ ($\mu_{d_{R_i}}$) are the common chemical potentials of left-handed (right-handed) leptons, left-handed quarks with different colors, and up (down) right-handed quarks with different colors, respectively. Furthermore, ‘ i ’ is the generation index, and the relevant hypercharges are

$$\begin{aligned} Y_L &= -1, & Y_R &= -2, \\ Y_Q &= \frac{1}{3}, & Y_{u_R} &= \frac{4}{3}, & Y_{d_R} &= -\frac{2}{3}. \end{aligned} \quad (2.9)$$

After substituting the hypercharges in Eqs. (2.7) and (2.8), we obtain

$$c_v(t) = \sum_{i=1}^{n_G} \left[\frac{g'}{24} \left(T_{R_i}^2 - T_{L_i}^2 + T_{d_{R_i}}^2 - 2T_{u_{R_i}}^2 + T_{Q_i}^2 \right) + \frac{g'}{8\pi^2} \left(\mu_{R_i}^2 - \mu_{L_i}^2 + \mu_{d_{R_i}}^2 - 2\mu_{u_{R_i}}^2 + \mu_{Q_i}^2 \right) \right], \quad (2.10)$$

$$c_B(t) = \frac{-g'^2}{8\pi^2} \sum_{i=1}^{n_G} \left[-2\mu_{R_i} + \mu_{L_i} - \frac{2}{3}\mu_{d_{R_i}} - \frac{8}{3}\mu_{u_{R_i}} + \frac{1}{3}\mu_{Q_i} \right]. \quad (2.11)$$

Let us make the same assumptions as in our previous studies, and simplify c_v and c_B accordingly [18, 20]. We assume that all quark Yukawa processes are in equilibrium and, because of the flavor mixing in the quark sector, all up or down quarks belonging to different generations with distinct handedness have the same chemical potential [18, 53]. For simplicity, we also assume that the Higgs asymmetry is zero and obtain [18, 54]

$$\mu_{u_R} = \mu_{d_R} = \mu_Q. \quad (2.12)$$

Furthermore, we assume that only the contributions of the baryonic and the first-generation leptonic chemical potentials to c_v and c_B are significant. As for the temperature fluctuations, it suffices to consider fluctuations in only one of the matter components, which we take to be e_R . Using Eq. (2.12) and the aforementioned assumptions, we simplify Eqs. (2.10) and (2.11) to obtain

$$c_v(t) = \frac{g'}{24} (\Delta T^2) + \frac{g'}{8\pi^2} (\mu_{e_R}^2 - \mu_{e_L}^2), \quad (2.13)$$

$$c_B(t) = -\frac{g'^2}{8\pi^2} \left(-2\mu_{e_R} + \mu_{e_L} - \frac{3}{4}\mu_B \right), \quad (2.14)$$

where $\mu_B = 12\mu_Q$, and $\Delta T^2 = T_{e_R}^2 - T^2$ is the temperature fluctuation, and T is the equilibrium temperature of the thermal bath, which includes all other components of the plasma. We set $\Delta T^2 = T^2 \beta[x(T)]$, where $\beta[x(T)]$ is an arbitrary profile function to be specified later, and $x(T) = t(T)/t_{EW} = (T_{EW}/T)^2$ is given by the Friedmann law.

In the Landau-Lifshitz frame, the continuity and Navier-Stokes equations are given as follows [20, 47–49]

$$\frac{\partial \rho}{\partial t} + \frac{1}{R} \vec{\nabla} \cdot [(\rho + p) \vec{v}] + 3H(\rho + p) = 0, \quad (2.15)$$

$$\left[\frac{\partial}{\partial t} + \frac{1}{R} (\vec{v} \cdot \vec{\nabla}) + H \right] \vec{v} + \frac{\vec{v}}{\rho + p} \frac{\partial p}{\partial t} = -\frac{1}{R} \frac{\vec{\nabla} p}{\rho + p} + \frac{\vec{J} \times \vec{B}_Y}{\rho + p} + \frac{\nu}{R^2} \left[\nabla^2 \vec{v} + \frac{1}{3} \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right], \quad (2.16)$$

where ν is the kinematic viscosity, and ρ and p are the energy density and the pressure of the plasma, respectively. Combining the fluid incompressibility condition in the lab frame, $\partial_t \rho + 3H(\rho + p) = 0$ or equivalently $H\vec{v} + \vec{v}\partial_t p/(\rho + p) = 0$, with Eq. (2.15) leads to the condition $\vec{\nabla} \cdot \vec{v} = 0$ [20, 49, 55].

In the following, we choose a simple monochromatic Chern-Simons configuration for the hypermagnetic field $\vec{B}_Y = (1/R)\vec{\nabla} \times \vec{A}_Y$, and the velocity field $\vec{v} = (1/R)\vec{\nabla} \times \vec{S}$ [18, 20]. To do this, we choose $\vec{A}_Y = \gamma(t)(\cos kz, \sin kz, 0)$, and $\vec{S} = r(t)(\cos kz, \sin kz, 0)$, for their corresponding vector potentials [56–58]. Note that we have chosen a fully helical form with the negative helicity for both, the reason for which will be stated later. Let us now obtain the evolution equation for the velocity field. Neglecting the displacement current in the lab frame in Eq. (2.3), the total current becomes $\vec{J} = (1/R)\vec{\nabla} \times \vec{B}_Y$, and as a result, $\vec{J} \times \vec{B}_Y$ vanishes in Eq. (2.16). Then, using $\vec{\nabla} \cdot \vec{v} = 0$ and $H\vec{v} + \vec{v}\partial_t p/(\rho + p) = 0$, as stated earlier, and neglecting the gradient terms in Eq. (2.16), the evolution equation for the velocity field becomes ⁴

$$\frac{\partial \vec{v}}{\partial t} = -\nu k'^2 \vec{v}, \quad (2.17)$$

where the kinematic viscosity $\nu \simeq 1/(5\alpha_Y^2 T)$ [60, 61]. Neglecting the displacement current in the lab frame and using the aforementioned configurations, the hyperelectric field and the evolution equation for the hypermagnetic field are obtained, as follows:

$$\vec{E}_Y = -\frac{k'}{\sigma} \vec{B}_Y + \frac{c_v}{\sigma} k' \vec{v} - \frac{c_B}{\sigma} \vec{B}_Y, \quad (2.18)$$

$$\frac{dB_Y(t)}{dt} = \left[-\frac{1}{t} - \frac{k'^2}{\sigma} - \frac{c_B k'}{\sigma} \right] B_Y(t) + \frac{c_v}{\sigma} k'^2 \langle \vec{v}(t) \cdot \hat{B}_Y(t) \rangle, \quad (2.19)$$

where $\vec{\omega} = -k'\vec{v}$, $\sigma = 100T$, angle brackets denote spatial average, and $k' = k/R = kT$. The latter shows the increase of the hypermagnetic length scale, due to the expansion of the Universe. Note that with the choice of vector potentials for $\vec{v}(t)$ and $\vec{B}_Y(t)$, the advection term $\vec{v} \times \vec{B}_Y$ has been set to zero in the above, and we have the following simplification: $\langle \vec{v}(t) \cdot \hat{B}_Y(t) \rangle \rightarrow v(t)$. In the next subsection we obtain the evolution equations for the matter-antimatter asymmetries.

2.2 Evolution equations for the matter-antimatter asymmetries

Before the EWPT, the gauge fields of $U_Y(1)$ couple to the fermions chirally, in contrast to those of $U_{em}(1)$ in the broken phase, leading to the non-conservation of the matter currents. This shows up in the Abelian anomaly equations [62], which, for the first-generation leptons, are

$$\begin{aligned} \nabla_\mu j_{e_R}^\mu &= -\frac{1}{4}(Y_R^2) \frac{g'^2}{16\pi^2} Y_{\mu\nu} \tilde{Y}^{\mu\nu} = \frac{g'^2}{4\pi^2} \vec{E}_Y \cdot \vec{B}_Y, \\ \nabla_\mu j_{e_L}^\mu &= \frac{1}{4}(Y_L^2) \frac{g'^2}{16\pi^2} Y_{\mu\nu} \tilde{Y}^{\mu\nu} = -\frac{g'^2}{16\pi^2} \vec{E}_Y \cdot \vec{B}_Y, \end{aligned} \quad (2.20)$$

⁴The term $(\vec{v} \cdot \vec{\nabla})\vec{v}$ is neglected because of being next to leading order. Furthermore, the magnetic pressure $B^2/8\pi$ is negligible compared to the fluid (radiation) pressure p which is homogeneous and isotropic. Indeed, the maximum value of their ratio at the onset of the EWPT is $B^2/8\pi p \approx 10^{-7} \ll 1$. Therefore, to a good approximation, the homogeneity and isotropy conditions remain valid and the pressure variations in the fluid $\vec{\nabla} p$ can be neglected [59].

where ∇_μ is the covariant derivative with respect to Friedmann-Robertson-Walker (FRW) metric $ds^2 = dt^2 - R^2(t)\delta_{ij}dx^i dx^j$, t is the physical time, and x^i s are the comoving coordinates. Integrating the above equations over all space and considering the perturbative chirality flip reactions for the leptons, we obtain (see Refs. [17, 18, 20, 63] and Appendix B for details),

$$\begin{aligned}\frac{d\eta_{e_R}}{dt} &= \frac{g'^2}{4\pi^2 s} \langle \vec{E}_Y \cdot \vec{B}_Y \rangle + \left(\frac{\Gamma_0}{t_{EW}} \right) \left(\frac{1-x}{\sqrt{x}} \right) (\eta_{e_L} - \eta_{e_R}) \\ &\quad - \frac{d}{dt} \left[\frac{g' \mu_{e_R}}{4\pi^2 s} \langle \vec{v} \cdot \vec{B}_Y \rangle \right] - \frac{d}{dt} \left[\left(\frac{\mu_{e_R}^2}{8\pi^2 s} + \frac{T^2}{24s} \right) \langle \vec{v} \cdot \vec{\omega} \rangle \right] + \frac{d}{dt} \left[\frac{2\sigma_{e_R}}{g' Y_{RS}} \langle \vec{v} \cdot \vec{E}_Y \rangle \right] \\ \frac{d\eta_{\nu_e^L}}{dt} &= \frac{d\eta_{e_L}}{dt} = -\frac{g'^2}{16\pi^2 s} \langle \vec{E}_Y \cdot \vec{B}_Y \rangle + \left(\frac{\Gamma_0}{2t_{EW}} \right) \left(\frac{1-x}{\sqrt{x}} \right) (\eta_{e_R} - \eta_{e_L}) \\ &\quad + \frac{d}{dt} \left[\frac{g' \mu_{e_L}}{8\pi^2 s} \langle \vec{v} \cdot \vec{B}_Y \rangle \right] + \frac{d}{dt} \left[\left(\frac{\mu_{e_L}^2}{8\pi^2 s} + \frac{T^2}{24s} \right) \langle \vec{v} \cdot \vec{\omega} \rangle \right] + \frac{d}{dt} \left[\frac{2\sigma_{e_L}}{g' Y_{LS}} \langle \vec{v} \cdot \vec{E}_Y \rangle \right].\end{aligned}\quad (2.21)$$

In the equations above, $\eta_f = (n_f/s)$ with $f = e_R, e_L, \nu_e^L$ is the fermion asymmetry and n_f is the charge density of the f th species of fermion, $s = 2\pi^2 g^* T^3/45$ is the entropy density and $g^* = 106.75$ is the effective number of relativistic degrees of freedom, $x = (t/t_{EW}) = (T_{EW}/T)^2$ is given by the Friedmann law, $\Gamma_0 = 121$, $t_{EW} = (M_0/2T_{EW}^2)$, and $M_0 = (M_{Pl}/1.66\sqrt{g^*})$, where M_{Pl} is the Plank mass. Furthermore, the term $\frac{\Gamma_0}{t_{EW}} \left(\frac{1-x}{\sqrt{x}} \right)$ appearing in the equations is the chirality flip rate of the right-handed electrons. In a similar manner, the evolution equation for the baryon asymmetry can also be obtained as (see Appendix B)

$$\frac{d\eta_B}{dt} = \frac{3g'^2}{8\pi^2 s} \langle \vec{E}_Y \cdot \vec{B}_Y \rangle + \frac{d}{dt} \left[\frac{2}{sg'} \left(\frac{\sigma_{d_R}}{Y_{d_R}} + \frac{\sigma_{u_R}}{Y_{u_R}} + 2\frac{\sigma_Q}{Y_Q} \right) \langle \vec{v} \cdot \vec{E}_Y \rangle \right]. \quad (2.22)$$

Using $\mu_f = (6s/T^2)\eta_f$ with Eq. (2.18) we obtain

$$\begin{aligned}\langle \vec{E}_Y \cdot \vec{B}_Y \rangle &= \frac{B_Y^2(t)}{100} \left[-\frac{k'}{T} - \frac{6sg'^2}{4\pi^2 T^3} \left(\eta_{e_R} - \frac{\eta_{e_L}}{2} + \frac{3}{8}\eta_B \right) \right] \\ &\quad + \left[\frac{g'}{24}\beta[x(T)] + \frac{36s^2 g'}{8\pi^2 T^6} (\eta_{e_R}^2 - \eta_{e_L}^2) \right] \frac{k'T}{100} \langle \vec{v}(t) \cdot \vec{B}_Y(t) \rangle.\end{aligned}\quad (2.23)$$

With the helical configurations chosen, $\langle \vec{v}(t) \cdot \vec{B}_Y(t) \rangle \rightarrow v(t)B_Y(t)$. Using $1\text{Gauss} \simeq 2 \times 10^{-20}\text{GeV}^2$, and setting the kinematic viscosity ν to zero for simplicity, we obtain the complete set of evolution equations for the matter-antimatter asymmetries and the amplitudes of the hypermagnetic and velocity fields as

$$\begin{aligned}\frac{d\eta_{e_R}}{dx} &= \frac{1}{\lambda_R(x)} \left[\Lambda_R(x) + [-C_1 - C_2\eta_T(x)] \left(\frac{B_Y(x)}{10^{20}G} \right)^2 x^{3/2} \right. \\ &\quad \left. + [C_3\beta(x) + C_4\Delta\eta^2(x)] v(x) \left(\frac{B_Y(x)}{10^{20}G} \right) \sqrt{x} \right] - \Gamma_0 \frac{1-x}{\sqrt{x}} [\eta_{e_R}(x) - \eta_{e_L}(x)],\end{aligned}\quad (2.24)$$

$$\begin{aligned}\frac{d\eta_{e_L}}{dx} &= \frac{1}{\lambda_L(x)} \left[\Lambda_L(x) - \frac{1}{4} [-C_1 - C_2\eta_T(x)] \left(\frac{B_Y(x)}{10^{20}G} \right)^2 x^{3/2} \right. \\ &\quad \left. - \frac{1}{4} [C_3\beta(x) + C_4\Delta\eta^2(x)] v(x) \left(\frac{B_Y(x)}{10^{20}G} \right) \sqrt{x} \right] + \Gamma_0 \frac{1-x}{2\sqrt{x}} [\eta_{e_R}(x) - \eta_{e_L}(x)],\end{aligned}\quad (2.25)$$

$$\frac{dB_Y}{dx} = \frac{1}{\sqrt{x}} [-C_5 - C_6\eta_T(x)] B_Y(x) - \frac{1}{x} B_Y(x) + [C_7\beta(x) + C_8\Delta\eta^2(x)] \frac{v(x)}{x^{3/2}}, \quad (2.26)$$

$$\begin{aligned} \frac{d\eta_B}{dx} = & \frac{3}{2} \left[-C_1 - C_2\eta_T(x) \left(\frac{B_Y(x)}{10^{20}G} \right)^2 x^{3/2} + [C_3\beta(x) + C_4\Delta\eta^2(x)] v(x) \left(\frac{B_Y(x)}{10^{20}G} \right) \sqrt{x} \right] \\ & + \frac{81x}{100Mg'10^{20}G} \langle \vec{E}_Y(x) \cdot \partial_x \vec{v}(x) \rangle \end{aligned} \quad (2.27)$$

$$\begin{aligned} \Delta\eta^2(x) &= \eta_{e_R}^2(x) - \eta_{e_L}^2(x), \\ \eta_T(x) &= \eta_{e_R}(x) - \frac{\eta_{e_L}(x)}{2} + \frac{3}{8}\eta_B(x), \\ \lambda_R(x) &= 1 - \frac{6g'Y_R}{8\pi^2} \frac{\langle \vec{v}(x) \cdot \vec{B}_Y(x) \rangle}{10^{20}G} \frac{x}{5000} - \frac{36M}{4\pi^2} \eta_{e_R}(x) k v^2(x), \\ \lambda_L(x) &= 1 + \frac{6g'Y_L}{8\pi^2} \frac{\langle \vec{v}(x) \cdot \vec{B}_Y(x) \rangle}{10^{20}G} \frac{x}{5000} + \frac{36M}{4\pi^2} \eta_{e_L}(x) k v^2(x), \\ \Lambda_R(x) &= \frac{6g'Y_R}{8\pi^2} \frac{x}{5000} \frac{\eta_{e_R}(x)}{10^{20}G} \left[\langle \vec{v}(x) \cdot \partial_x \vec{B}_Y(x) \rangle + \left\langle \frac{\vec{v}(x) \cdot \vec{B}_Y(x)}{x} \right\rangle + \langle \vec{B}_Y(x) \cdot \partial_x \vec{v}(x) \rangle \right] \\ &+ \left[\frac{36M}{4\pi^2} \eta_{e_R}^2(x) + \frac{1}{12M} \right] k \vec{v}(x) \cdot \partial_x \vec{v}(x) + \frac{x}{25Mg'Y_R 10^{20}G} \langle \vec{E}_Y(x) \cdot \partial_x \vec{v}(x) \rangle, \\ \Lambda_L(x) &= -\frac{6g'Y_L}{8\pi^2} \frac{x}{5000} \frac{\eta_{e_L}(x)}{10^{20}G} \left[\langle \vec{v}(x) \cdot \partial_x \vec{B}_Y(x) \rangle + \left\langle \frac{\vec{v}(x) \cdot \vec{B}_Y(x)}{x} \right\rangle + \langle \vec{B}_Y(x) \cdot \partial_x \vec{v}(x) \rangle \right] \\ &- \left[\frac{36M}{4\pi^2} \eta_{e_L}^2(x) + \frac{1}{12M} \right] k \vec{v}(x) \cdot \partial_x \vec{v}(x) + \frac{x}{25Mg'Y_L 10^{20}G} \langle \vec{E}_Y(x) \cdot \partial_x \vec{v}(x) \rangle, \end{aligned} \quad (2.28)$$

where $M = 2\pi^2 g^*/45$, and the coefficients $C_i, i = 1, \dots, 8$, are

$$\begin{aligned} C_1 &= 0.00096 \left(\frac{k}{10^{-7}} \right) \alpha_Y, \\ C_2 &= 865688 \alpha_Y^2, \\ C_3 &= 0.71488 \left(\frac{k}{10^{-7}} \right) \alpha_Y^{3/2}, \\ C_4 &= 17152.7 \left(\frac{k}{10^{-7}} \right) \alpha_Y^{3/2}, \\ C_5 &= 0.356 \left(\frac{k}{10^{-7}} \right)^2, \\ C_6 &= 3.18373 \times 10^8 \alpha_Y \left(\frac{k}{10^{-7}} \right), \\ C_7 &= 262.9 \times 10^{20} \sqrt{\alpha_Y} \left(\frac{k}{10^{-7}} \right)^2, \\ C_8 &= 63 \times 10^{25} \sqrt{\alpha_Y} \left(\frac{k}{10^{-7}} \right)^2, \end{aligned} \quad (2.29)$$

and $\alpha_Y = g'^2/4\pi \simeq 0.01$ is the fine-structure constant for the $U_Y(1)$. We now choose the profile of temperature fluctuation $\beta[x(T)] = \Delta T^2/T^2$, as defined in Eq. (2.13) and the paragraph below it, to be a Gaussian function of x :

$$\beta(x) = \frac{\beta_0}{b\sqrt{2\pi}} \exp \left[-\frac{(x - x_0)^2}{2b^2} \right], \quad (2.30)$$

where β_0 is the amplitude multiplying the normalized Gaussian distribution, and $x = (t/t_{EW}) = (T_{EW}/T)^2$, as defined before. The profile of the vorticity fluctuation must have an overlap with that of temperature fluctuation, in order to produce any effect. For simplicity, we choose the two profiles to be identical. That is,

$$\omega(x) = k'v(x) = \frac{k'v_0}{b\sqrt{2\pi}} \exp \left[-\frac{(x - x_0)^2}{2b^2} \right], \quad (2.31)$$

where v_0 is the amplitude of the velocity fluctuation. We should mention that the occurrence of any fluctuation in a plasma in a quasi-equilibrium state would normally trigger a restoring response originating from dissipative effects, such as viscous effects. Here, for simplicity we assume that the combined results of the original fluctuations and the ensuing dissipative effects have the profiles given by Eqs. (2.30,2.31).

The majority of analysis presented in the following section is for a single pulse for temperature and vorticity, as stated above, which, as we shall show, produce matter-antimatter asymmetries and helical hypermagnetic field. However, at the end of the next section we present the results for two sets of successive pulses, the latter one having the same temperature profile but with negative amplitude, showing that the first pulse is the main determinant of the outcome.

3 Numerical Solution

In this section, we obtain the numerical solutions of the evolution equations. As mentioned earlier, we investigate the effects of the temperature fluctuations of right-handed electrons, in the presence of vorticity, on the generation and evolution of the hypermagnetic field and the matter-antimatter asymmetries, in the temperature range $100 \text{ GeV} \leq T \leq 10 \text{ TeV}$. We consider the temperature fluctuations as small Gaussian distributions in x , as shown in Eq. (2.30), that occur close to the EWPT. As for the vorticity field, we consider small fluctuations, whose profiles, as shown in Eq. (2.31), coincide with those of the temperature fluctuations. We also investigate the cases with constant vorticity fields for comparison. As we shall show, the former is much more interesting and will be the focus of our work, since it is not only physically more realistic, but also could yield orders of magnitude larger results for the asymmetries and the hypermagnetic field. In the following, we solve the evolution equations by considering the comoving wave number as $k = 10^{-7}$, and setting the initial values of the hypermagnetic field amplitude and the matter-antimatter asymmetries to zero, i.e. $B_Y^{(0)} = 0$, and $\eta_{eR}^{(0)} = \eta_{eL}^{(0)} = \eta_B^{(0)} = 0$.

For our first case, we solve the coupled differential equations with the initial conditions, $v_0 = 10^{-5}$, $b = 2 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$, for various values of the amplitude of temperature fluctuations β_0 , and present the results in Fig. 1. As can be seen in the figure, the simultaneous occurrence of small vorticity fluctuation and temperature fluctuation for the right handed electrons leads to the generation of strong hypermagnetic fields which then produce the matter-antimatter asymmetries, all starting from zero initial values. It can be seen that by increasing the amplitude of the temperature fluctuation, the maximum and the final values of

the hypermagnetic field amplitude, as well as the matter-antimatter asymmetries, increase. We have found that in our model signs of the matter-antimatter asymmetries produced and the helicity of hypermagnetic and vorticity fields, are always opposite. This is a manifestation of the generalized charge conservation as stated in Eq. (6.9) of Appendix B.⁵ The Chern-Simons configuration that we have chosen for the hypermagnetic and vorticity fields has negative helicity. Figure 1 also shows that the hypermagnetic field amplitude grows to its maximum value of about 10^{21} G, then decreases due to the expansion of the Universe. We have also investigated the effects of changing the amplitude of the vorticity fluctuation, and have found similar results.




Figure 1: Time plots of: (a) the right-handed electron asymmetry η_{e_R} , (b) the left-handed electron asymmetry η_{e_L} , (c) the baryon asymmetry η_B , and (d) the hypermagnetic field amplitude B_Y , for various values of the amplitude of temperature fluctuation of e_R . The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{e_R}^{(0)} = \eta_{e_L}^{(0)} = \eta_B^{(0)} = 0$, $v_0 = 10^{-5}$, $b = 2 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$. The dashed line is for $\beta_0 = 3 \times 10^{-4}$, the solid line is for $\beta_0 = 5 \times 10^{-4}$, and the dotted line is for $\beta_0 = 7 \times 10^{-4}$.

For our second case we solve the set of evolution equations with the initial conditions, $v_0 = 10^{-5}$, $\beta_0 = 5 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$, for various values of the width or duration of both fluctuations b , and show the results in Fig. 2. As can be seen, by decreasing the width of the Gaussian function, the maximum and the final values of the hypermagnetic field amplitude, and the baryon asymmetry increase.⁶

For our third case, we solve the coupled equations with the initial conditions, $v_0 = 10^{-5}$, $b = 2 \times 10^{-4}$, and $\beta_0 = 5 \times 10^{-4}$, for various values of center time of the fluctuations x_0 , and present the results in Fig. 3. As can be seen, when the fluctuations occur at an earlier time or

⁵In fact, the second and fourth terms in Eq. (6.9) are negligible in this study.

⁶We have also used a few other profiles, the results of which we can summarize as follows. For smooth profiles with the same normalization, the results are mainly dependent on the widths and not on the precise functional form of the profiles. However, the results usually increase by an order of magnitude for profiles functions which have discontinuities.




Figure 2: Time plots of: (a) the baryon asymmetry η_B , and (b) the hypermagnetic field amplitude B_Y , for various values of the width of fluctuations. The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{e_R}^{(0)} = \eta_{e_L}^{(0)} = \eta_B^{(0)} = 0$, $v_0 = 10^{-5}$, $\beta_0 = 5 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$. The dotted line is obtained for $b = 3 \times 10^{-4}$, the solid line for $b = 2 \times 10^{-4}$, the dashed line for $b = 10^{-4}$.

higher temperature, the maxima and the final amplitudes of the hypermagnetic fields increase, and as a result, the matter-antimatter asymmetries increase as well.




Figure 3: Time plots of: (a) the baryon asymmetry η_B , and (b) the hypermagnetic field amplitude B_Y , for various values of the time of fluctuations. The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{e_R}^{(0)} = \eta_{e_L}^{(0)} = \eta_B^{(0)} = 0$, $v_0 = 10^{-5}$, $\beta_0 = 5 \times 10^{-4}$, $b = 2 \times 10^{-4}$. The dotted line is obtained for $x_0 = 55 \times 10^{-5}$, the solid line for $x_0 = 45 \times 10^{-5}$, and the dashed line for $x_0 = 35 \times 10^{-5}$.

For our fourth and final case, we solve the set of evolution equations, with the initial conditions $\beta_0 = 5 \times 10^{-4}$, $b = 2 \times 10^{-4}$ and $x_0 = 45 \times 10^{-5}$, for two different vorticity configurations. First configuration is a vorticity fluctuation with amplitude $v_0 = 10^{-5}$ and $x_0 = 45 \times 10^{-5}$. Second configuration is a constant vorticity with amplitude $v_0 = 10^{-2}$. The results are presented in Fig. 4. As can be seen from the figure, the general trends of the evolution curves are similar. The prominent feature of this comparison is the surprising result that a fluctuation with amplitude smaller by three orders of magnitude produces results comparable with the constant vorticity configuration.

Now we can address our assertion that $\mu/T \ll 1$ within our model, i.e., our initial conditions and results. First, note that upon using the relations $\mu_f = (6s/T^2)\eta_f$ and $s = (2\pi^2 g^*/45)T^3$, we obtain $(\mu_f/T) = (12\pi^2 g^*/45)\eta_f = 280.95\eta_f$. The largest asymmetry in our results is obtained for η_B and is shown in the Fig. 2(a). This figure shows that $(\eta_B)_{\max} \sim 5 \times 10^{-9}$, leading to $(\mu/T)_{\max} \sim 10^{-6}$. The condition $\mu/T \ll 1$ has been used, for example, to justify neglect-




Figure 4: Time plots of: (a) the baryon asymmetry η_B , and (b) the hypermagnetic field amplitude B_Y , for two different vorticity configurations. The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{e_R}^{(0)} = \eta_{e_L}^{(0)} = \eta_B^{(0)} = 0$, $\beta_0 = 5 \times 10^{-4}$, $b = 2 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$. The solid line is for vorticity fluctuation with $v_0 = 10^{-5}$, and the dashed line is for constant vorticity with $v_0 = 10^{-2}$.

ing terms of $O(\mu/T)$ in Eqs. (5.5-5.8) in Appendix A, leading to the expressions for c_v and c_B shown in Eqs. (2.7,2.8), respectively. Moreover, as shown in Appendix B, this condition, along with the assumption of non-relativistic velocity of the plasma, imply that the CME and CVE contributions to the temporal components of the four-currents in the AMHD equations are negligible within our model.

We finally address the issue of two successive pulses with opposite temperature profiles to see by how much can the second pulse negate the results of the first. For this purpose it suffices to assume that the profile of the vorticities are unchanged. To be specific, we assume

$$\beta(x) = \beta_+(x) + \beta_-(x), \quad (3.1)$$

$$v(x) = v_+(x) + v_-(x), \quad (3.2)$$

where

$$\beta_{\pm}(x) = \frac{\pm\beta_0}{b\sqrt{2\pi}} \exp\left[-\frac{(x-x_{0,\pm})^2}{2b^2}\right], \quad (3.3)$$

and

$$v_{\pm}(x) = \frac{v_0}{b\sqrt{2\pi}} \exp\left[-\frac{(x-x_{0,\pm})^2}{2b^2}\right]. \quad (3.4)$$

We consider three cases in which the time separation of the pulses $\Delta x_0 = x_{0,+} - x_{0,-}$ are $5b$, b and $0.1b$, where b denotes the width of the pulses. The results are shown in Fig. 5, where we also show our results for a single pulse for comparison. As can be seen, the final values of the asymmetries generated are reduced, as compared to the single pulse case, by a factor of about 5, 50 and 1000, respectively. The final values of the hypermagnetic field generated are reduced by square root of values stated above. It is interesting to note that even in the case $\Delta x_0 = 0.1b$, the model produced $\eta_B \simeq 10^{-13}$, a value which can be increased easily by increasing β_0 , v_0 , and/or decreasing b .




Figure 5: Time plots of: (a) the baryon asymmetry η_B , (b) the hypermagnetic field amplitude B_Y for two sets of successive and opposing fluctuations. The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{e_R}^{(0)} = \eta_{e_L}^{(0)} = \eta_B^{(0)} = 0$, $v_{0,+} = v_{0,-} = 10^{-5}$, $b = 2 \times 10^{-4}$, $\beta_{0,+} = -\beta_{0,-} = 5 \times 10^{-4}$, and $x_{0,+} = 4.5 \times 10^{-4}$. The large dashed line is for $x_{0,-} = 1.45 \times 10^{-3} = 5b + x_{0,+}$, the medium dashed line is for $x_{0,-} = 6.5 \times 10^{-4} = b + x_{0,+}$, the dotted line is for $x_{0,-} = 4.7 \times 10^{-4} = 0.1b + x_{0,+}$, and the solid line is obtained in the absence of the second set of fluctuations.

4 Conclusion

In this study, we have investigated the contribution of the temperature-dependent CVE to the generation and evolution of the hypermagnetic fields and the matter-antimatter asymmetries, in the symmetric phase of the early Universe and in the temperature range $100 \text{ GeV} \leq T \leq 10 \text{ TeV}$. The CVE has two possible sources in a vortical plasma, one from the chiralities and the other from the temperature of the particles. The former has been investigated in the literature much more than the latter. Here, we have focused on the latter in the form of transient temperature fluctuations, and have shown its important role in the production and evolution of the hypermagnetic fields and the matter-antimatter asymmetries. The transient fluctuations that we have considered are in the form of sharp Gaussian shaped pulses. In particular, we have shown that small simultaneous and transient fluctuations of vorticity about zero background value and temperature of some matter degrees of freedom about the equilibrium temperature of the plasma, close to the EWPT, can generate strong hypermagnetic fields and large matter-antimatter asymmetries, even in the absence of any initial seed for the hypermagnetic field or any initial matter-antimatter asymmetries. Furthermore, we have shown that, an increase in the amplitude of temperature or vorticity fluctuations leads to the production of stronger hypermagnetic fields, and therefore, larger matter-antimatter asymmetries. This outcome has not been observed in any of the previous studies. In some studies which only take the CME into account, either an initially strong hypermagnetic field produces matter-antimatter asymmetries, or initial large matter-antimatter asymmetries strengthen a preexisting seed of hypermagnetic field [17–19]. In some other studies which also include the CVE and assume large initial chiralities in the vortical plasma, a seed of the hypermagnetic field is produced which then grows due to the CME [20].

In this work, we have considered a simple monochromatic helical configuration for the vorticity and hypermagnetic fields with a negative helicity, which ensures the production of the desired positive matter-antimatter asymmetries. Furthermore, we have shown that, either an increase in the amplitude of the temperature or vorticity fluctuations, or a decrease in their widths leads to the production of stronger hypermagnetic fields, and therefore, larger matter-antimatter asymmetries. We have also shown that fluctuations in vorticity are several orders

of magnitude more productive than constant vorticity. Within our model, the temperature-dependent CVE is the dominant effect as compared to the CME. We have also shown that when there are two sets of successive pulses with opposite temperature profiles, the first pulse is the main determinant of the outcome. A generalization of this work would be a stochastic analysis of fluctuations of both temperature and vorticity.

5 APPENDIX A

In this appendix we present the expressions for the chiral vorticity and helicity coefficients, c_v and c_B given by Eqs. (2.7), and (2.8). First, we start with the relevant and well known expressions in the broken phase. In relativistic hydrodynamics, the energy and number currents can flow separately in the presence of dissipative processes, therefore the definition of the flow will not be trivial [64]. Some common hydrodynamic frames in which the equations of AMHD may be formulated include the Landau-Lifshitz (or energy) frame [65], the Eckart (or conserved charge/particle) frame [66], and the more recently introduced anomalous “no-drag” frame [67]. In the latter, as its name suggests, a stationary obstacle experiences no drag, even when the energy and charge currents are present. In the Landau-Lifshitz frame, the energy-momentum tensor $T^{\mu\nu}$ and the total electric current J^μ for a plasma consisting of a single species of massless fermions of both chiralities are given by

$$T^{\mu\nu} = (\rho + p)u^\mu u^\nu - pg^{\mu\nu} + \frac{1}{4}g^{\mu\nu}F^{\alpha\beta}F_{\alpha\beta} - F^{\nu\sigma}F^\mu{}_\sigma + \tau^{\mu\nu}, \quad (5.1)$$

$$J^\mu = \rho_{\text{el}}u^\mu + J_{\text{cm}}^\mu + J_{\text{cv}}^\mu + \nu^\mu, \quad (5.2)$$

$$J_{\text{cm}}^\mu = (Q_R\xi_{\text{B,R}} + Q_L\xi_{\text{B,L}})B^\mu = c_B B^\mu, \quad (5.3)$$

$$J_{\text{cv}}^\mu = (Q_R\xi_{\text{v,R}} + Q_L\xi_{\text{v,L}})\omega^\mu = c_v \omega^\mu, \quad (5.4)$$

where $F_{\alpha\beta} = \nabla_\alpha A_\beta - \nabla_\beta A_\alpha$ is the field strength tensor, p and ρ are the pressure and the energy density of the plasma, ρ_{el} is the electric charge density, $u^\mu = \gamma(1, \vec{v}/R)$ is the four-velocity of the plasma normalized such that $u^\mu u_\mu = 1$, and γ is the Lorentz factor.⁷ In the above equations, ν^μ and $\tau^{\mu\nu}$ denote the electric diffusion current and viscous stress tensor, respectively [3], Q_R (Q_L) denotes the electric charges of the right-handed (left-handed) fermions, $B^\mu = (\epsilon^{\mu\nu\rho\sigma}/2R^3)u_\nu F_{\rho\sigma}$ is the magnetic field four-vector, and $\omega^\mu = (\epsilon^{\mu\nu\rho\sigma}/R^3)u_\nu \nabla_\rho u_\sigma$ is the vorticity four-vector, with the totally anti-symmetric Levi-Civita tensor density specified by $\epsilon^{0123} = -\epsilon_{0123} = 1$.

Furthermore, in the Landau-Lifshitz frame, the CME and CVE coefficients for chiral fermions are given as [3, 45–49]

$$\xi_{\text{B,R}} = \frac{Q_R \mu_R}{4\pi^2} \left[1 - \frac{1}{2} \frac{n_R \mu_R}{\rho + p} \right] - \frac{1}{24} \frac{n_R T^2}{\rho + p}, \quad (5.5)$$

$$\xi_{\text{B,L}} = -\frac{Q_L \mu_L}{4\pi^2} \left[1 - \frac{1}{2} \frac{n_L \mu_L}{\rho + p} \right] + \frac{1}{24} \frac{n_L T^2}{\rho + p}, \quad (5.6)$$

⁷Note that the self-consistency of our calculation in which the diagonal Einstein tensor obtained from the FRW metric is used implies that not only should the electromagnetic field density be small compared to the energy density of the Universe [68], but also the bulk velocity should obey the condition $|\vec{v}| \ll 1$, or equivalently $\gamma \simeq 1$ and $u^\mu \simeq (1, \vec{v}/R)$.

$$\xi_{v,R} = \frac{\mu_R^2}{8\pi^2} \left[1 - \frac{2}{3} \frac{n_R \mu_R}{\rho + p} \right] + \frac{1}{24} T^2 \left[1 - \frac{2n_R \mu_R}{\rho + p} \right], \quad (5.7)$$

$$\xi_{v,L} = -\frac{\mu_L^2}{8\pi^2} \left[1 - \frac{2}{3} \frac{n_L \mu_L}{\rho + p} \right] - \frac{1}{24} T^2 \left[1 - \frac{2n_L \mu_L}{\rho + p} \right]. \quad (5.8)$$

where T is the temperature and n_R (n_L) is the right-handed (left-handed) charge density.⁸ As we shall show explicitly in Sec. 3, $\mu_{R,L}/T \ll 1$ within our model, i.e., our initial conditions and results. Hence, $n_{R,L} \simeq \frac{1}{6} \mu_{R,L} T^2$ and $\rho = 3p \simeq \frac{\pi^2}{30} g^* T^4$, and Eqs. (5.5-5.8) may be simplified as follows

$$\xi_{B,R} \simeq \frac{Q_R \mu_R}{4\pi^2}, \quad (5.9)$$

$$\xi_{B,L} \simeq -\frac{Q_L \mu_L}{4\pi^2}, \quad (5.10)$$

$$\xi_{v,R} \simeq \frac{\mu_R^2}{8\pi^2} + \frac{1}{24} T^2, \quad (5.11)$$

$$\xi_{v,L} \simeq -\frac{\mu_L^2}{8\pi^2} - \frac{1}{24} T^2. \quad (5.12)$$

Moreover, since $\mu/T \ll 1$, we will consider only the Ohmic part of the diffusion current $\nu^\mu = \sigma[E^\mu + T(u^\mu u^\nu - g^{\mu\nu})\nabla_\nu(\frac{\mu}{T})]$ given by the first term, where $\sigma = \sigma_R + \sigma_L$ is the electrical conductivity, and $E^\mu = F^{\mu\nu} u_\nu$ is the electric field four-vector. In order to carry over these results to the symmetric phase of the early Universe, it suffices to replace the electromagnetic field by the hypercharge gauge field and the electric charges of different particle species by their relevant hypercharges. Taking into account all three generations of leptons and quarks, we can easily obtain the chiral vorticity and helicity coefficients c_v and c_B , given by Eqs. (2.7), and (2.8), using Eqs. (5.9-5.12). The four-vectors B^μ , ω^μ , a^μ , and E^μ in the limit $v \ll 1$ are given below,

$$B^\mu = \gamma \left(\vec{v} \cdot \vec{B}, \frac{\vec{B} - \vec{v} \times \vec{E}}{R} \right) \simeq \left(\vec{v} \cdot \vec{B}, \frac{\vec{B}}{R} \right) \quad (5.13)$$

$$\omega^\mu = \gamma \left(\vec{v} \cdot \vec{\omega}, \frac{\vec{\omega} - \vec{v} \times \vec{a}}{R} \right) \simeq \left(\vec{v} \cdot \vec{\omega}, \frac{\vec{\omega}}{R} \right) \quad (5.14)$$

$$a^\mu = \gamma \left(\vec{v} \cdot \vec{a}, \frac{\vec{a} + \vec{v} \times \vec{\omega}}{R} \right) \simeq \left(\vec{v} \cdot \vec{a}, \frac{\vec{a}}{R} \right) \quad (5.15)$$

$$E^\mu = \gamma \left(\vec{v} \cdot \vec{E}, \frac{\vec{E} + \vec{v} \times \vec{B}}{R} \right) \simeq \left(\vec{v} \cdot \vec{E}, \frac{\vec{E} + \vec{v} \times \vec{B}}{R} \right), \quad (5.16)$$

where $a^\mu = \Omega^{\mu\nu} u_\nu$ is acceleration four-vector, $\Omega_{\mu\nu} = \nabla_\mu u_\nu - \nabla_\nu u_\mu$ is vorticity tensor, $a^i = R\Omega^{0i}$ is three vector acceleration. We have also used the assumption $\partial_t \sim \vec{\nabla} \cdot \vec{v}$ in the derivative expansion of the hydrodynamics, so $\vec{v} \times \vec{E} \simeq v^2 \vec{B}$ and we have ignored the terms of $O(v^2)$.

⁸The charge density n is the difference between the particle and anti-particle charge densities.

6 APPENDIX B

In this appendix we start with the basic expression for the anomaly given by Eq. (2.20) and obtain its explicit form which eventually leads to Eqs. (2.21, 2.24, 2.25). The equations of AMHD consist of energy-momentum conservation, Maxwell's equations and the anomaly relations. These equations may be expressed covariantly as

$$\nabla_\mu T^{\mu\nu} = 0, \quad (6.1)$$

$$\begin{aligned} \nabla_\mu F^{\mu\nu} &= J^\nu \\ \nabla_\mu \tilde{F}^{\mu\nu} &= 0 \end{aligned} \quad (6.2)$$

$$\nabla_\mu j_{R,L}^\mu = C_{R,L} E_\mu B^\mu, \quad (6.3)$$

where $\tilde{F}^{\mu\nu} = \frac{1}{2R^3} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$ is the dual field strength tensor, $C_{R,L}$ are the corresponding right- and left- handed anomaly coefficients, and $j_{R,L}^\mu$ are the fermionic currents given as follows

$$\begin{aligned} j_R^\mu &= n_R u^\mu + \xi_{B,R} B^\mu + \xi_{v,R} \omega^\mu + V_R^\mu, \\ j_L^\mu &= n_L u^\mu + \xi_{B,L} B^\mu + \xi_{v,L} \omega^\mu + V_L^\mu, \end{aligned} \quad (6.4)$$

where $n_{R,L}$ and $V_{R,L}^\mu$ are the chiral charge density and chiral particle diffusion current, respectively [3, 47–49]. The anomaly equations given above can be written out as

$$\partial_t j_{(R,L)}^0 + \frac{1}{R} \vec{\nabla} \cdot \vec{j}_{(R,L)} + 3H j_{(R,L)}^0 = C_{(R,L)} E_\mu B^\mu, \quad (6.5)$$

where

$$j_{(R,L)}^0 = n_{(R,L)} + \xi_{B,(R,L)} \vec{v} \cdot \vec{B} + \xi_{v,(R,L)} \vec{v} \cdot \vec{\omega} + V_{R,L}^0. \quad (6.6)$$

Upon taking the spatial average of Eq. (6.5), the boundary term vanishes and we obtain⁹

$$\begin{aligned} \partial_t n_{(R,L)} + 3H n_{(R,L)} &= - \left[\partial_t + 3H \right] \left[\xi_{B,(R,L)} \langle \vec{v} \cdot \vec{B} \rangle + \xi_{v,(R,L)} \langle \vec{v} \cdot \vec{\omega} \rangle + \frac{\sigma_{R,L}}{Q_{R,L}} \langle \vec{v} \cdot \vec{E} \rangle \right] \\ &\quad - C_{R,L} \langle \vec{E} \cdot \vec{B} \rangle. \end{aligned} \quad (6.7)$$

Using the relation $\dot{s}/s = -3H$, the anomaly equation reduces to the simplified form

$$\partial_t \left(\frac{n_{(R,L)}}{s} \right) = - \partial_t \left[\frac{\xi_{B,(R,L)}}{s} \langle \vec{v} \cdot \vec{B} \rangle + \frac{\xi_{v,(R,L)}}{s} \langle \vec{v} \cdot \vec{\omega} \rangle + \frac{\sigma_{R,L}}{Q_{R,L}s} \langle \vec{v} \cdot \vec{E} \rangle \right] - \frac{C_{R,L}}{s} \langle \vec{E} \cdot \vec{B} \rangle. \quad (6.8)$$

It is worth mentioning that since $(\partial_t + 3H) \langle \vec{A} \cdot \vec{B} \rangle = -2 \langle \vec{E} \cdot \vec{B} \rangle$, Eq. (6.8) can be rewritten in the following form

$$\partial_t \left[\eta_{R,L} + \frac{\xi_{B,(R,L)}}{s} \langle \vec{v} \cdot \vec{B} \rangle + \frac{\xi_{v,(R,L)}}{s} \langle \vec{v} \cdot \vec{\omega} \rangle + \frac{\sigma_{(R,L)}}{Q_{(R,L)}s} \langle \vec{v} \cdot \vec{E} \rangle - \frac{C_{R,L}}{2s} \langle \vec{A} \cdot \vec{B} \rangle \right] = 0, \quad (6.9)$$

which clearly reveals the conservation of a generalized charge [7, 47].

⁹Note that $E_\mu B^\mu = (\vec{v} \cdot \vec{E})(\vec{v} \cdot \vec{B}) - (\vec{E} + \vec{v} \times \vec{B}) \cdot (\vec{B} - \vec{v} \times \vec{E}) = -\vec{E} \cdot \vec{B} + O(v^2)$, and we have ignored the terms of $O(v^2)$.

Using Eqs. (6.8, 5.9-5.12) and $x = (T_{\text{EW}}/T)^2 = t/t_{\text{EW}}$, we obtain

$$\begin{aligned} \frac{d\eta_R}{dx} = & \frac{1}{\left[1 + \frac{6Q_R}{4\pi^2} \frac{\langle \vec{v} \cdot \vec{B} \rangle}{10^{20}G} \frac{x}{5000} - \frac{36M}{4\pi^2} \eta_R k v^2\right]} \left[-\frac{6Q_R}{4\pi^2} \frac{x}{5000} \frac{\eta_R}{10^{20}G} \left[\langle \vec{v} \cdot \partial_x \vec{B} \rangle + \left\langle \frac{\vec{v} \cdot \vec{B}}{x} \right\rangle + \langle \vec{B} \cdot \partial_x \vec{v} \rangle \right] \right. \\ & + \left[\frac{36M}{4\pi^2} \eta_R^2 + \frac{1}{12M} \right] k \vec{v} \cdot \partial_x \vec{v} - \frac{x}{50MQ_R 10^{20}G} \left[\langle \vec{v} \cdot \partial_x \vec{E} \rangle + \left\langle \frac{\vec{v} \cdot \vec{E}}{x} \right\rangle + \langle \vec{E} \cdot \partial_x \vec{v} \rangle \right] \\ & \left. - \frac{t_{\text{EW}} C_R}{s} \langle \vec{E} \cdot \vec{B} \rangle \right], \end{aligned} \quad (6.10)$$

$$\begin{aligned} \frac{d\eta_L}{dx} = & \frac{1}{\left[1 - \frac{6Q_L}{4\pi^2} \frac{\langle \vec{v} \cdot \vec{B} \rangle}{10^{20}G} \frac{x}{5000} + \frac{36M}{4\pi^2} \eta_L k v^2\right]} \left[\frac{6Q_L}{4\pi^2} \frac{x}{5000} \frac{\eta_L}{10^{20}G} \left[\langle \vec{v} \cdot \partial_x \vec{B} \rangle + \left\langle \frac{\vec{v} \cdot \vec{B}}{x} \right\rangle + \langle \vec{B} \cdot \partial_x \vec{v} \rangle \right] \right. \\ & - \left[\frac{36M}{4\pi^2} \eta_L^2 + \frac{1}{12M} \right] k \vec{v} \cdot \partial_x \vec{v} - \frac{x}{50MQ_L 10^{20}G} \left[\langle \vec{v} \cdot \partial_x \vec{E} \rangle + \left\langle \frac{\vec{v} \cdot \vec{E}}{x} \right\rangle + \langle \vec{E} \cdot \partial_x \vec{v} \rangle \right] \\ & \left. - \frac{t_{\text{EW}} C_L}{s} \langle \vec{E} \cdot \vec{B} \rangle \right], \end{aligned} \quad (6.11)$$

where $M = 2\pi^2 g^*/45$. Here the electric field \vec{E} is not an independent variable and is determined by Eq. (2.18). From the approximation of a vanishing displacement current, we have $\partial_x \vec{E} = -\vec{E}/x$.

As is well known, in the symmetric phase of the early Universe $T > 100$ GeV, and in Sec. 3 we show that the condition $\mu/T \leq 10^{-6}$ holds within our model. These conditions, in conjunction with the low-velocity limit considered in this work, highly suppress the contributions of the temporal components of the CVE, CME and diffusion currents. To ascertain this claim, we have solved Eqs. (2.24)-(2.26) both with and without the additional temporal components and present the differences in Fig. 6. Comparing the scales of this figure and Fig. 1, we observe the following for the differences: $\Delta\eta/\eta \sim 10^{-4}$ and $\Delta B_Y/B_Y \sim 10^{-9}$.

References

- [1] D. E. Kharzeev, “Topology, magnetic field, and strongly interacting matter,” *Ann. Rev. Nucl. Part. Sci.* **65**, 193-214 (2015) [[arXiv:1501.01336 \[hep-ph\]](#)].
- [2] A. Vilenkin, Macroscopic Parity Violating Effects: Neutrino Fluxes From Rotating Black Holes And In Rotating Thermal Radiation, *Phys. Rev. D* **20**, 1807 (1979), [DOI: 10.1103/PhysRevD.20.1807];
- [3] D.T. Son, P. Surowka, Hydrodynamics with Triangle Anomalies, *Phys.Rev.Lett.* **103** (2009) 191601, [[arXiv:0906.5044 \[hep-th\]](#)].
- [4] A. V. Sadofyev, V. I. Shevchenko, V. I. Zakharov, Notes on chiral hydrodynamics within effective theory approach, *Phys.Rev.D* **83**, 105025, (2011), [[arXiv:1012.1958 \[hep-th\]](#)].
- [5] Shi Pu, Jian-hua Gao and Qun Wang, A consistent description of kinetic equation with triangle anomaly, *Phys.Rev. D* **83**, 094017 (2011), [[arXiv:1008.2418 \[nucl-th\]](#)].




Figure 6: Time plots of the differences between the results with and without the inclusion of the temporal components of the CVE, CME and diffusion currents given by the last three terms of Eq. (6.6): (a) the right-handed electron asymmetry $\Delta\eta_{eR}$, (b) the left-handed electron asymmetry $\Delta\eta_{eL}$, (c) the baryon asymmetry $\Delta\eta_B$, and (d) the hypermagnetic field amplitude ΔB_Y . The initial conditions are: $k = 10^{-7}$, $B_Y^{(0)} = 0$, $\eta_{eR}^{(0)} = \eta_{eL}^{(0)} = \eta_B^{(0)} = 0$, $\beta_0 = 5 \times 10^{-4}$, $v_0 = 10^{-5}$, $b = 2 \times 10^{-4}$, and $x_0 = 45 \times 10^{-5}$.

- [6] V.P. Kirilin, A.V. Sadofyev, V.I. Zakharov. Chiral Vortical Effect in Superfluid, Phys.Rev. **D86**, 025021, (2012), [\[arXiv:1203.6312 \[hep-th\]\]](#).
- [7] A. Avdoshkin, V. P. Kirilin, A. V. Sadofyev and V. I. Zakharov, “On consistency of hydrodynamic approximation for chiral media,” Phys. Lett. B **755**, 1-7 (2016), [\[arXiv:1402.3587 \[hep-th\]\]](#).
- [8] A. Avkhadiev and A. V. Sadofyev, Chiral Vortical Effect for Bosons, Phys. Rev. D **96**, no.4, 045015 (2017), [\[arXiv:1702.07340 \[hep-th\]\]](#).
- [9] V. P. Kirilin, A. V. Sadofyev, Anomalous Transport and Generalized Axial Charge, Phys.Rev. **D96** (2017) no.1, 016019, [\[arXiv:1703.02483 \[hep-th\]\]](#).
- [10] Ö. F. Dayi and E. Kiliçarslan, “Quantum Kinetic Equation in the Rotating Frame and Chiral Kinetic Theory,” Phys. Rev. D **98**, no.8, 081701 (2018), [\[arXiv:1807.05912 \[hep-th\]\]](#).
- [11] A. Vilenkin, Equilibrium Parity Violating Current In A Magnetic Field, Phys. Rev. D **22**, 3080 (1980), [DOI: 10.1103/PhysRevD.22.3080].
- [12] D. Kharzeev, Parity violation in hot QCD: Why it can happen, and how to look for it, Phys. Lett. B **633**, 260 (2006), [\[hep-ph/0406125\]](#).
- [13] D. Kharzeev and A. Zhitnitsky, Charge separation induced by P-odd bubbles in QCD matter, Nucl. Phys. A **797**, 67 (2007), [\[arXiv:0706.1026 \[hep-ph\]\]](#).
- [14] D. E. Kharzeev, L. D. McLerran and H. J. Warringa, The Effects of topological charge change in heavy ion collisions: ‘Event by event P and CP violation’, Nucl. Phys. A **803**, 227 (2008), [\[arXiv:0711.0950 \[hep-ph\]\]](#).

- [15] Jian-Hua Gao, Zuo-Tang Liang, Shi Pu, Qun Wang, Xin-Nian Wang, Chiral Anomaly and Local Polarization Effect from Quantum Kinetic Approach, *Phys.Rev.Lett.* **109**, (2012) ,32301, [[arXiv:1203.0725 \[hep-ph\]](#)].
- [16] D.E. Kharzeev, J. Liao, S.A. Voloshin, G. Wang, Chiral magnetic and vortical effects in high-energy nuclear collisions-A status report, *Prog.Part.Nucl.Phys.* **88**, (2016), 1-28, [[arXiv:1511.04050 \[hep-ph\]](#)].
- [17] S. Rostam Zadeh and S. S. Gousheh, Contributions to the $U_Y(1)$ Chern-Simons term and the evolution of fermionic asymmetries and hypermagnetic fields, *Phys. Rev. D* **94**, 056013 (2016), [[arXiv:1512.01942 \[hep-ph\]](#)].
- [18] S. Rostam Zadeh and S. S. Gousheh, Effects of the $U_Y(1)$ Chern-Simons term and its baryonic contribution on matter asymmetries and hypermagnetic fields, *Phys. Rev. D* **95**, 056001 (2017), [[arXiv:1607.00650 \[hep-ph\]](#)].
- [19] S. Rostam Zadeh and S. S. Gousheh, A Minimal System Including Weak Sphalerons for Investigating the Evolution of Matter Asymmetries and Hypermagnetic Fields, *Phys. Rev. D* **99**, 096009, (2019), [[arXiv:1812.10092 \[hep-ph\]](#)].
- [20] Abbaslu, S. Rostam Zadeh and S. S. Gousheh, “Contribution of the chiral vortical effect to the evolution of the hypermagnetic field and the matter-antimatter asymmetry in the early Universe,” *Phys. Rev. D* **100**, no.11, 116022 (2019) , [[arXiv:1908.10105 \[hep-ph\]](#)].
- [21] F. Elahi and S. R. Zadeh, “Flavon magnetobaryogenesis,” *Phys. Rev. D* **102**, no.9, 096018 (2020), [[arXiv:2008.04434 \[hep-ph\]](#)].
- [22] L. M. Widrow, Origin of galactic and extragalactic magnetic fields, *Rev. Mod. Phys.* **74**, 775 (2002), [[arXiv:astro-ph/0207240](#)].
- [23] P. P. Kronberg, Extragalactic magnetic fields, *Rep. Prog. Phys.***57**, 325 (1994), [DOI: 10.1088/0034-4885/57/4/001] .
- [24] R. M. Kulsrud, R. Cen, J. P. Ostriker, and D. Ryu, The Protogalactic Origin for Cosmic Magnetic Fields, *Astrophys.J.* **480**, 481 (91997), [[arXiv:astro-ph/9607141](#)].
- [25] T. Vachaspati, Magnetic fields from cosmological phase transitions, *Phys.Lett. B* **265**, 258261 (1991), [DOI: 10.1016/0370-2693(91)90051-Q].
- [26] K. Enqvist and P. Olesen, On primordial magnetic fields of electroweak origin, *Phys. Lett. B* **319**, 178 (1993), [[arXiv:hep-ph/9308270](#)].
- [27] M. Joyce and M. Shaposhnikov, Primordial magnetic fields, right-handed electrons, and the Abelian anomaly, *Phys. Rev. Lett.* **79**, 1193 (1997), [[arXiv:astro-ph/9703005](#)].
- [28] J. M. Cornwall, Speculations on primordial magnetic helicity, *Phys.Rev. D* **56**, 6146 (1997), [[hep-th/9704022](#)].
- [29] S. M. Carroll, G. B. Field, and R. Jackiw, Limits on a Lorentz and Parity Violating Modification of Electrodynamics, *Phys. Rev. D* **41**, 1231 (1990), [DOI: 10.1103/PhysRevD.41.1231].
- [30] M. M. Anber, E. Sabancilar, Hypermagnetic Fields and Baryon Asymmetry from Pseudoscalar Inflation, *Phys.Rev. D* **92** (2015) no.10, 101501,[[arXiv:1507.00744](#)].
- [31] V. B. Semikoz, A. Yu. Smirnov, and D. D. Sokoloff, Generation of hypermagnetic helicity and leptogenesis in the early Universe, *Phys. Rev. D* **93**, 103003 (2016), [[arXiv:1604.02273 \[hep-ph\]](#)].

- [32] A. J. Long, E. Sabancilar, and T. Vachaspati, Leptogenesis and primordial magnetic fields, *J. Cosmol. Astropart. Phys.* 02 (2014) 036, [[arXiv:1309.2315 \[astro-ph.CO\]](#)].
- [33] M. Giovannini and M. E. Shaposhnikov, Primordial hypermagnetic fields and triangle anomaly, *Phys. Rev. D* **57**, 2186 (1998), [[arXiv:hep-ph/9710234](#)].
- [34] M. Giovannini, Primordial Hypermagnetic Knots, *Phys. Rev. D* **61**, 063004 (2000), [[arXiv:hep-ph/9905358](#)].
- [35] K. Kamada and A. J. Long, Baryogenesis from decaying magnetic helicity, *Phys. Rev. D* **94**, 123509 (2016), [[arXiv:1606.08891 \[astro-ph.CO\]](#)].
- [36] K. Bamba, C. Q. Geng and S. H. Ho, “Hypermagnetic Baryogenesis,” *Phys. Lett. B* **664**, 154-156 (2008), [[arXiv:0712.1523 \[hep-ph\]](#)].
- [37] B. Fields and S. Sarkar, Big-Bang nucleosynthesis (2006 Particle Data Group mini-review), *J. Phys. G* 33 (2006) 1 [[arXiv:astro-ph/0601514](#)]; G. Steigman, Primordial Nucleosynthesis: The Predicted and Observed Abundances and Their Consequences, *PoS NICXI* (2010) 001, [[arXiv:1008.4765 \[astro-ph.CO\]](#)].
- [38] V. Simha and G. Steigman, Constraining The Early-Universe Baryon Density And Expansion Rate, *JCAP* 0806 (2008) 016, [[arXiv:0803.3465 \[astro-ph\]](#)].
- [39] A. D. Sakharov, Violation of CP Invariance, C Asymmetry, and Baryon Asymmetry of the Universe, *Pisma Zh. Eksp. Teor. Fiz.* 5 (1967) 32 [*JETP Lett.* 5 (1967) 24] [*Sov. Phys. Usp.* 34 (1991) 392] [*Usp. Fiz. Nauk* 161 (1991) 61], [DOI: 10.1070/PU1991v034n05ABEH002497].
- [40] J. I. Kapusta, B. Muller and M. Stephanov, “Relativistic Theory of Hydrodynamic Fluctuations with Applications to Heavy Ion Collisions,” *Phys. Rev. C* **85**, 054906 (2012), [DOI:10.1103/PhysRevC.85.054906], [[arXiv:1112.6405 \[nucl-th\]](#)].
- [41] S. Jeon and U. Heinz, “Introduction to Hydrodynamics,” *Int. J. Mod. Phys. E* **24**, no.10, 1530010 (2015) [DOI:10.1142/S0218301315300106], [[arXiv:1503.03931 \[hep-ph\]](#)].
- [42] J. de Zarate and J. Sengers, *Hydrodynamic Fluctuations in Fluids and Fluid Mixtures* (Elsevier Science, 2006).
- [43] T. Bhattacharyya, P. Garg, R. Sahoo and P. Samantray, “Time Evolution of Temperature Fluctuation in a Non-Equilibrated System,” *Eur. Phys. J. A* **52**, no.9, 283 (2016) doi:10.1140/epja/i2016-16283-x, [[arXiv:1510.03154 \[hep-ph\]](#)].
- [44] M. N. Chernodub, “Chiral Heat Wave and mixing of Magnetic, Vortical and Heat waves in chiral media,” *JHEP* **01**, 100 (2016) [DOI:10.1007/JHEP01(2016)100] [[arXiv:1509.01245 \[hep-th\]](#)].
- [45] Y. Neiman and Y. Oz, “Relativistic Hydrodynamics with General Anomalous Charges,” *JHEP* **03**, 023 (2011), [DOI:10.1007/JHEP03(2011)023], [[arXiv:1011.5107 \[hep-th\]](#)].
- [46] K. Landsteiner, E. Megias and F. Pena-Benitez, ”Anomalous Transport from Kubo Formulae,” *Lect. Notes Phys.* **871**, 433-468 (2013) [DOI:10.1007/978-3-642-37305-3-17], [[arXiv:1207.5808 \[hep-th\]](#)].
- [47] N. Yamamoto, “Chiral transport of neutrinos in supernovae: Neutrino-induced fluid helicity and helical plasma instability,” *Phys. Rev. D* **93**, no.6, 065017 (2016), [DOI:10.1103/PhysRevD.93.065017], [[arXiv:1511.00933 \[astro-ph.HE\]](#)].
- [48] K. Landsteiner, “Notes on Anomaly Induced Transport,” *Acta Phys. Polon. B* **47**, 2617 (2016), [DOI:10.5506/APhysPolB.47.2617], [[arXiv:1610.04413 \[hep-th\]](#)].

- [49] S. Anand, J. R. Bhatt and A. K. Pandey, “Chiral Battery, scaling laws and magnetic fields,” JCAP **07**, 051 (2017) [DOI:10.1088/1475-7516/2017/07/051], [arXiv:1705.03683 \[astro-ph.CO\]](#).
- [50] C. P. Dettmann, N. E. Frankel, and V. Kowalenko, Plasma electrodynamics in the expanding Universe, Phys Rev. D **48**, 12 (1993), [DOI: 10.1103/PhysRevD.48.5655].
- [51] K. Subramanian, J. D. Barrow, Magnetohydrodynamics in the Early Universe and the Damping of Non-linear Alfvén Waves, Phys. Rev. D **58**, 083502 (1998), [\[arXiv:astro-ph/9712083\]](#).
- [52] R. Durrer and A. Neronov, Cosmological magnetic fields: Their generation, evolution and observation, Astron. Astrophys. Rev. **21**, 62 (2013), [\[arXiv:1303.7121 \[astro-ph.CO\]\]](#).
- [53] J. M. Cline, K. Kainulainen, and K. A. Olive, Erasure and Regeneration of the Primordial Baryon Asymmetry by Sphalerons, Phys. Rev. Lett. **71**, 2372 (1993), [\[arXiv:hep-ph/9304321\]](#); J. M. Cline, K. Kainulainen and K. A. Olive, “Protecting the primordial baryon asymmetry from erasure by sphalerons,” Phys. Rev. D **49**, 6394-6409 (1994) [\[arXiv:hep-ph/9401208 \[hep-ph\]\]](#).
- [54] M. Dvornikov and V. B. Semikoz, Leptogenesis via hypermagnetic fields and baryon asymmetry, J. Cosmol. Astropart. Phys. 1202 (2012) 040; Erratum: JCAP **1208** (2012) E01, [\[arXiv:1111.6876 \[hep-ph\]\]](#).
- [55] P. Pavlotic, N. Leite, G. Sigl, Chiral Magnetohydrodynamic Turbulence, Phys Rev. D **96**, 023504 (2017), [\[arXiv:1612.07382 \[astro-ph.CO\]\]](#).
- [56] V. Rubakov and A. Tavkhelidze, Stable anomalous states of superdense matter in gauge theories, Phys. Lett. B **165**, 109 (1985), [DOI: 10.1016/0370-2693(85)90701-4].
- [57] V. Rubakov, On the electroweak theory at high fermion density, Prog. Theor. Phys. **75**, 366 (1986), [DOI: 10.1143/PTP.75.366].
- [58] M. Giovannini, Spectrum of anomalous magnetohydrodynamics, Phys. Rev. D **93**, 103518 (2016), [\[arXiv:1509.02126 \[hep-th\]\]](#).
- [59] P. Pavlovic, N. Leite, G. Sigl, Chiral Magnetohydrodynamic Turbulence, Phys Rev. D **96**, 023504 (2017), [\[arXiv:1612.07382 \[astro-ph.CO\]\]](#).
- [60] S. Weinberg, Gravitation and Cosmology (John Wiley & Sons, New York, 1972).
- [61] R. Banerjee, K. Jedamzik, The Evolution of cosmic magnetic fields: From the very early universe, to recombination, to the present, Phys Rev. D **70**, 123003 (2004), [\[arXiv:astro-ph/0410032\]](#).
- [62] M. E. Shaposhnikov, Structure of the High Temperature Gauge Ground State and Electroweak Production of the Baryon Asymmetry, Nucl. Phys. B **299**, 797 (1988);
- [63] M. Dvornikov and V. B. Semikoz, Lepton asymmetry growth in the symmetric phase of an electroweak plasma with hypermagnetic fields versus its washing out by sphalerons, Phys. Rev. D **87**, 025023 (2013), [\[arXiv:1212.1416 \[astro-ph.CO\]\]](#).
- [64] A. Monnai, “Landau and Eckart frames for relativistic fluids in nuclear collisions,” Phys. Rev. C **100**, no.1, 014901 (2019) [DOI:10.1103/PhysRevC.100.014901] [\[arXiv:1904.11940 \[nucl-th\]\]](#).
- [65] L. D. Landau and E. M. Lifshitz, Fluid Mechanics (Pergamon Press, New York, 1959).
- [66] C. Eckart, “The Thermodynamics of Irreversible Processes. 1. The Simple Fluid,” Phys. Rev. **58**, 267-269 (1940), [DOI:10.1103/PhysRev.58.267].
- [67] M. A. Stephanov and H. U. Yee, “No-Drag Frame for Anomalous Chiral Fluid,” Phys. Rev. Lett. **116**, no.12, 122302 (2016) [DOI:10.1103/PhysRevLett.116.122302] [\[arXiv:1508.02396 \[hep-th\]\]](#).

- [68] Banerjee R (2002) Evolution of primordial magnetic fields in the early Universe. PhD thesis, Ludwig-Maximilians-Universität, München.