

PATH INDEPENDENCE OF THE ADDITIVE FUNCTIONALS FOR STOCHASTIC DIFFERENTIAL EQUATIONS DRIVEN BY G-LÉVY PROCESSES*

HUIJIE QIAO^{1,2} AND JIANG-LUN WU³

1. *School of Mathematics, Southeast University
Nanjing, Jiangsu 211189, China*

2. *Department of Mathematics, University of Illinois at Urbana-Champaign
Urbana, IL 61801, USA
hjiaogean@seu.edu.cn*

3. *Department of Mathematics, Computational Foundry, Swansea University
Bay Campus, Swansea SA1 8EN, UK
j.l.wu@swansea.ac.uk*

ABSTRACT. In the paper, we consider a type of stochastic differential equations driven by G-Lévy processes. We prove that a kind of their additive functionals has path independence and extend some known results.

1. INTRODUCTION

Recently, development of mathematical finance forces the appearance of a type of processes-G-Brownian motions([5]). And then the related theory, such as stochastic calculus and stochastic differential equations (SDEs in short) driven by G-Brownian motions, are widely studied([1, 5, 6, 8]). However, in some financial models, volatility uncertainty makes G-Brownian motions insufficient for simulating these models. One important reason lies in the continuity of their paths with respect to the time variable. So, Hu-Peng [2] solved the problem by introducing G-Lévy processes. And the type of processes has discontinuous (right continuous with left limits) paths. Later, Paczka [3] defined the Itô-Lévy stochastic integrals, deduced the Itô formula, established SDEs driven by G-Lévy processes and stated the existence and uniqueness of solutions for these equations under lipschitz conditions. Most recently, under non-lipschitz conditions, Wang-Gao [14] proved well-definedness of SDEs driven by G-Lévy processes and investigated exponential stability of their solutions. Here, we follow up the line in [14], define the additive functionals of SDEs driven by G-Lévy processes and study their path independence.

Concretely speaking, we consider the following SDEs on \mathbb{R}^d :

$$dY_t = b(t, Y_t)dt + h_{ij}(t, Y_t)d\langle B^i, B^j \rangle_t + \sigma(t, Y_t)dB_t + \int_{\mathbb{R}^d \setminus \{0\}} f(t, Y_t, u)L(dt, du), \quad (1)$$

AMS Subject Classification(2010): 60H10, 60G51.

Keywords: Path independence, additive functionals, G-Lévy processes, stochastic differential equations driven by G-Lévy processes.

*This work was partly supported by NSF of China (No. 11001051, 11371352, 11671083) and China Scholarship Council under Grant No. 201906095034.

where B is a G-Brownian motion, $\langle B^i, B^j \rangle_t$ is the mutual variation process of B^i and B^j for $i, j = 1, 2, \dots, d$ and $L(dt, du)$ is a G-random measure (See Subsection 2.4). The coefficients $b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $h_{ij} = h_{ji} : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times d}$ and $f : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}) \mapsto \mathbb{R}^d$ are Borel measurable. Here and hereafter we use the convention that the repeated indices stand for the summation. Thus, under $(\mathbf{H}_{b,h,\sigma,f}^1)$ -($\mathbf{H}_{b,h,\sigma,f}^2$) in Subsection 2.5, by [14, Theorem 3.1], we know that Eq.(1) has a unique solution Y_t . And then we introduce the additive functionals of Y_t and define path independence of these functionals. Finally, we prove that these functionals have path independence under some assumption.

Next, we say our motivations. First, we mention that Ren-Yang [12] proved path independence of additive functionals for SDEs driven by G-Brownian motions. Since these equations can not satisfy the actual demand very well, to extend them becomes one of our motivations. Second, by analyzing some special cases, we surprisingly find that, we can express explicitly these additive functionals. However, in our known results ([9, 10, 11]), it is difficult to express explicitly additive functionals. Therefore, this is the other of our motivations.

This paper is arranged as follows. In Section 2, we introduce G-Lévy processes, the Itô-Lévy stochastic integrals, SDEs driven by G-Lévy processes, additive functionals, path independence and some related results. The main results and their proofs are placed in Section 3. Moreover, we analysis some special cases and compare our result with some known results ([9, 10, 11, 12]) in Subsection 3.3.

2. PRELIMINARY

In the section, we introduce some concepts and results used in the sequel.

2.1. Notation. In the subsection, we introduce notations used in the sequel.

For convenience, we shall use $|\cdot|$ and $\|\cdot\|$ for norms of vectors and matrices, respectively. Furthermore, let $\langle \cdot, \cdot \rangle$ denote the scalar product in \mathbb{R}^d . Let Q^* denote the transpose of the matrix Q .

Let $lip(\mathbb{R}^n)$ be the set of all Lipschitz continuous functions on \mathbb{R}^n and $C_{b,lip}(\mathbb{R}^d)$ be the collection of all bounded and Lipschitz continuous functions on \mathbb{R}^d . Let $C_b^3(\mathbb{R}^d)$ be the space of bounded and three times continuously differentiable functions with bounded derivatives of all orders less than or equal to 3.

2.2. G-Lévy processes. In the subsection, we introduce G-Lévy processes.(c.f.[2])

Let Ω be a given set and \mathcal{H} be a linear space of real functions defined on Ω such that if $X_1, \dots, X_n \in \mathcal{H}$, then $\phi(X_1, \dots, X_n) \in \mathcal{H}$ for each $\phi \in lip(\mathbb{R}^n)$. If $X \in \mathcal{H}$, we call X as a random variable.

Definition 2.1. If a functional $\bar{\mathbb{E}} : \mathcal{H} \mapsto \mathbb{R}$ satisfies: for $X, Y \in \mathcal{H}$,

- (i) $X \geq Y, \bar{\mathbb{E}}[X] \geq \bar{\mathbb{E}}[Y]$,
- (ii) $\bar{\mathbb{E}}[X + Y] \leq \bar{\mathbb{E}}[X] + \bar{\mathbb{E}}[Y]$,
- (iii) for all $\lambda \geq 0, \bar{\mathbb{E}}[\lambda X] = \lambda \bar{\mathbb{E}}[X]$,
- (iv) for all $c \in \mathbb{R}, \bar{\mathbb{E}}[X + c] = \bar{\mathbb{E}}[X] + c$,

we call $\bar{\mathbb{E}}$ as a sublinear expectation on \mathcal{H} and $(\Omega, \mathcal{H}, \bar{\mathbb{E}})$ as a sublinear expectation space.

Next, we define the distribution of a random vector on $(\Omega, \mathcal{H}, \bar{\mathbb{E}})$. For a n -dimensional random vector $X = (X_1, X_2, \dots, X_n)$ for $X_i \in \mathcal{H}$, $i = 1, 2, \dots, n$, set

$$F_X(\phi) := \bar{\mathbb{E}}(\phi(X)), \quad \phi \in \text{lip}(\mathbb{R}^n),$$

and then we call F_X as the distribution of X .

Definition 2.2. Assume that X_1, X_2 are two n -dimensional random vectors defined on different sublinear expectation spaces. If for all $\phi \in \text{lip}(\mathbb{R}^n)$,

$$F_{X_1}(\phi) = F_{X_2}(\phi),$$

we say that the distributions of X_1, X_2 are the same.

Definition 2.3. For two random vectors $Y = (Y_1, Y_2, \dots, Y_m)$ for $Y_j \in \mathcal{H}$ and $X = (X_1, X_2, \dots, X_n)$ for $X_i \in \mathcal{H}$, if for all $\phi \in \text{lip}(\mathbb{R}^n \times \mathbb{R}^m)$,

$$\bar{\mathbb{E}}[\phi(X, Y)] = \bar{\mathbb{E}}[\bar{\mathbb{E}}[\phi(x, Y)]_{x=X}],$$

we say that Y is independent from X .

Here, we use two above concepts to define Lévy processes on $(\Omega, \mathcal{H}, \bar{\mathbb{E}})$.

Definition 2.4. Let $X = (X_t)_{t \geq 0}$ be a d -dimensional càdlàg process on $(\Omega, \mathcal{H}, \bar{\mathbb{E}})$. If X satisfies

- (i) $X_0 = 0$;
 - (ii) for $t, s \geq 0$, the increment $X_{s+t} - X_t$ is independent from $(X_{t_1}, X_{t_2}, \dots, X_{t_n})$, for any n and $0 \leq t_1 < t_2 < \dots < t_n \leq t$;
 - (iii) the distribution of $X_{s+t} - X_t$ does not depend on t ;
- we call X as a Lévy process.

Definition 2.5. Assume that X is a d -dimensional Lévy process. If there exists a decomposition $X_t = X_t^c + X_t^d$ for $t \geq 0$, where (X_t^c, X_t^d) is a $2d$ -dimensional Lévy process satisfying

$$\lim_{t \downarrow 0} \frac{\bar{\mathbb{E}}|X_t^c|^3}{t} = 0, \quad \bar{\mathbb{E}}|X_t^d| \leq Ct, \quad t \geq 0, \quad C \geq 0,$$

we call X as a G -Lévy process.

In the following, we characterize G -Lévy processes by partial differential equations.

Theorem 2.6. Assume that X is a d -dimensional G -Lévy process. Then for $g \in C_b^3(\mathbb{R}^d)$ with $g(0) = 0$, set

$$G_X[g(\cdot)] := \lim_{t \downarrow 0} \frac{\bar{\mathbb{E}}[g(X_t)]}{t},$$

and then G_X has the following Lévy-Khintchine representation

$$G_X[g(\cdot)] = \sup_{(\nu, \zeta, Q) \in \mathcal{U}} \left\{ \int_{\mathbb{R}^d \setminus \{0\}} g(u) \nu(du) + \langle \partial_x g(0), \zeta \rangle + \frac{1}{2} \text{tr}[\partial_x^2 g(0) Q Q^*] \right\}, \quad (2)$$

where \mathcal{U} is a subset of $\mathcal{M}(\mathbb{R}^d \setminus \{0\}) \times \mathbb{R}^d \times \mathbb{R}^{d \times d}$, $\mathcal{M}(\mathbb{R}^d \setminus \{0\})$ is the collection of all measures on $(\mathbb{R}^d \setminus \{0\}, \mathcal{B}(\mathbb{R}^d \setminus \{0\}))$, $\mathbb{R}^{d \times d}$ is the set of all $d \times d$ matrices and \mathcal{U} satisfies

$$\sup_{(\nu, \zeta, Q) \in \mathcal{U}} \left\{ \int_{\mathbb{R}^d \setminus \{0\}} |u| \nu(du) + |\zeta| + \frac{1}{2} \text{tr}[Q Q^*] \right\} < \infty. \quad (3)$$

Theorem 2.7. *Suppose that X is a d -dimensional G -Lévy process. Then for $\phi \in C_{b, \text{lip}}(\mathbb{R}^d)$, $v(t, x) := \bar{\mathbb{E}}[\phi(x + X_t)]$ is the unique viscosity solution of the following partial integro-differential equation:*

$$\begin{aligned} 0 &= \partial_t v(t, x) - G_X[v(t, x + \cdot) - v(t, x)] \\ &= \partial_t v(t, x) - \sup_{(\nu, \zeta, Q) \in \mathcal{U}} \left\{ \int_{\mathbb{R}^d \setminus \{0\}} [v(t, x + u) - v(t, x)] \nu(du) + \langle \partial_x v(t, x), \zeta \rangle \right. \\ &\quad \left. + \frac{1}{2} \text{tr}[\partial_x^2 v(t, x) Q Q^*] \right\} \end{aligned}$$

with the initial condition $v(0, x) = \phi(x)$.

Conversely, if we have a set \mathcal{U} satisfying (3), is there a d -dimensional G -Lévy process having the Lévy-Khintchine representation (2) with the same set \mathcal{U} ? The answer is affirmed. We take $\Omega := D_0(\mathbb{R}^+, \mathbb{R}^d)$, where $D_0(\mathbb{R}^+, \mathbb{R}^d)$ is the space of all càdlàg functions $\mathbb{R}_+ \ni t \mapsto \omega_t \in \mathbb{R}^d$ with $\omega_0 = 0$, equipped with the Skorokhod topology.

Theorem 2.8. *Suppose that \mathcal{U} satisfies (3). Then there exists a sublinear expectation $\bar{\mathbb{E}}$ on Ω such that the canonical process X is a d -dimensional G -Lévy process having the Lévy-Khintchine representation (2) with the same set \mathcal{U} .*

2.3. A capacity. In the subsection, we introduce a capacity and related definitions.

First of all, fix a set \mathcal{U} satisfying (3) and $T > 0$ and take $\Omega_T := D_0([0, T], \mathbb{R}^d)$ and the sublinear expectation $\bar{\mathbb{E}}$ in Theorem 2.8. Thus, we know that $(\Omega_T, \mathcal{H}, \bar{\mathbb{E}})$ is a sublinear expectation space. Here, we work on the space. Let $L_G^p(\Omega_T)$ be the completion of $\text{lip}(\Omega_T)$ under the norm $\|\cdot\|_p := (\bar{\mathbb{E}}|\cdot|^p)^{1/p}$, $p \geq 1$.

Let

$$\mathcal{V} := \{\nu \in \mathcal{M}(\mathbb{R}^d \setminus \{0\}) : \exists (\zeta, Q) \in \mathbb{R}^d \times \mathbb{R}^{d \times d} \text{ such that } (\nu, \zeta, Q) \in \mathcal{U}\}$$

and let \mathcal{G} be the set of all the Borel measurable functions $g : \mathbb{R}^d \mapsto \mathbb{R}^d$ with $g(0) = 0$.

Assumption:

(i) There exists a measure $\mu \in \mathcal{M}(\mathbb{R}^d)$ such that

$$\int_{\mathbb{R}^d \setminus \{0\}} |z| \mu(dz) < \infty, \quad \mu(\{0\}) = 0$$

and for all $\nu \in \mathcal{V}$ there exists a function $g_\nu \in \mathcal{G}$ satisfying

$$\nu(A) = \mu(g_\nu^{-1}(A)), \quad \forall A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}).$$

(ii) There exists $0 < q < 1$ such that

$$\sup_{\nu \in \mathcal{V}} \int_{0 < |z| < 1} |z|^q \nu(dz) < \infty.$$

(iii)

$$\sup_{\nu \in \mathcal{V}} \nu(\mathbb{R}^d \setminus \{0\}) < \infty.$$

Let $(\tilde{\Omega}, \mathcal{F}, \mathbb{P})$ be a probability space supporting a Brownian motion W and a Poisson random measure $N(dt, dz)$ with the intensity measure $\mu(dz)dt$. Let

$$\mathcal{F}_t := \sigma\{W_s, N((0, s], A) : 0 \leq s \leq t, A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})\} \vee \mathcal{N}, \quad \mathcal{N} := \{U \in \mathcal{F}, \mathbb{P}(U) = 0\}.$$

We introduce the following set.

Definition 2.9. $\mathcal{A}_{0,T}^{\mathcal{U}}$ is a set of all the processes $\theta_t = (\theta_t^d, \theta_t^{1,c}, \theta_t^{2,c})$ for $t \in [0, T]$ satisfying

(i) $(\theta_t^{1,c}, \theta_t^{2,c})$ is a \mathcal{F}_t -adapted process and θ^d is a \mathcal{F}_t -predictable random field on $[0, T] \times \mathbb{R}^d$,

(ii) For \mathbb{P} -a.s. ω and a.e. $t \in [0, T]$,

$$(\theta^d(t, \cdot)(\omega), \theta_t^{1,c}(\omega), \theta_t^{2,c}(\omega)) \in \{(g_\nu, \zeta, Q) \in \mathcal{G} \times \mathbb{R}^d \times \mathbb{R}^{d \times d} : (\nu, \zeta, Q) \in \mathcal{U}\},$$

(iii)

$$\mathbb{E}^{\mathbb{P}} \left[\int_0^T \left(|\theta_t^{1,c}| + \|\theta_t^{2,c}\|^2 + \int_{\mathbb{R}^d \setminus \{0\}} |\theta^d(t, z)| \mu(dz) \right) dt \right] < \infty.$$

For $\theta \in \mathcal{A}_{0,T}^{\mathcal{U}}$, set

$$B_t^{0,\theta} := \int_0^t \theta_s^{1,c} ds + \int_0^t \theta_s^{2,c} dW_s + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \theta^d(s, z) N(ds, dz), \quad t \in [0, T],$$

and by Corollary 14 in [3], it holds that for $\xi \in L_G^1(\Omega_T)$

$$\bar{\mathbb{E}}[\xi] = \sup_{\theta \in \mathcal{A}_{0,T}^{\mathcal{U}}} \mathbb{E}^{\mathbb{P}^\theta}[\xi], \quad \mathbb{P}^\theta = \mathbb{P} \circ (B^{0,\theta})^{-1}.$$

And then define

$$\bar{C}(D) := \sup_{\theta \in \mathcal{A}_{0,T}^{\mathcal{U}}} \mathbb{P}^\theta(D), \quad D \in \mathcal{B}(\Omega_T),$$

and \bar{C} is a capacity. For $D \in \mathcal{B}(\Omega_T)$, if $\bar{C}(D) = 0$, we call D as a polar set. So, if a property holds outside a polar set, we say that the property holds quasi-surely (q.s. in short).

2.4. The Itô integrals with respect to G-Lévy processes. In the subsection, we introduce the Itô integrals with respect to G-Lévy processes under the framework of the above subsection.

Let X denote the canonical process on the space, i.e. $X_t(\omega) = \omega_t, t \in [0, T]$. So, X is a d -dimensional G-Lévy process. Although the Itô integrals with respect to G-Brownian motions have been introduced in [8], we need to introduce two related spaces used in the sequel. Take $0 = t_0 < t_1 < \dots < t_N = T$. Let $p \geq 1$ be fixed. Define

$$\mathcal{M}_G^{p,0}(0, T) := \left\{ \eta_t(\omega) = \sum_{j=1}^N \xi_{j-1}(\omega) 1_{[t_{j-1}, t_j)}(t); \xi_{j-1}(\omega) \in L_G^p(\Omega_{t_{j-1}}) \right\}.$$

Let $\mathcal{M}_G^p(0, T)$ and $\mathcal{H}_G^p(0, T)$ denote the completion of $\mathcal{M}_G^{p,0}(0, T)$ under the norm

$$\|\eta\|_{\mathcal{M}_G^p(0, T)} = \left(\int_0^T \bar{\mathbb{E}}|\eta_t|^p dt \right)^{\frac{1}{p}} \quad \text{and} \quad \|\eta\|_{\mathcal{H}_G^p(0, T)} = \left(\bar{\mathbb{E}} \left(\int_0^T |\eta_t|^2 dt \right)^{\frac{p}{2}} \right)^{\frac{1}{p}},$$

respectively. Let $\mathcal{M}_G^p([0, T], \mathbb{R}^d)$ and $\mathcal{H}_G^p([0, T], \mathbb{R}^d)$ be the collection of all the processes

$$\eta_t = (\eta_t^1, \eta_t^2, \dots, \eta_t^d), \quad t \in [0, T], \quad \eta^i \in \mathcal{M}_G^p(0, T) \text{ and } \mathcal{H}_G^p(0, T),$$

respectively.

Next, we introduce the Itô integrals with respect to random measures. First, define a random measure: for any $0 \leq t \leq T$ and $A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$,

$$\kappa_t := X_t - X_{t-}, \quad L((0, t], A) := \sum_{0 < s \leq t} I_A(\kappa_s), \quad q.s..$$

And then we define the Itô integral with respect to the random measure $L(dt, du)$. Let $\mathcal{H}_G^S([0, T] \times (\mathbb{R}^d \setminus \{0\}))$ be the collection of all the processes defined on $[0, T] \times (\mathbb{R}^d \setminus \{0\}) \times \Omega$ with the form

$$f(s, u)(\omega) = \sum_{k=1}^{n-1} \sum_{l=1}^m \phi_{k,l}(X_{t_1}, X_{t_2} - X_{t_1}, \dots, X_{t_k} - X_{t_{k-1}}) I_{[t_k, t_{k+1})}(s) \psi_l(u), \quad n, m \in \mathbb{N},$$

where $0 \leq t_1 < \dots < t_n \leq T$ is a partition of $[0, T]$, $\phi_{k,l} \in C_{b,lip}(\mathbb{R}^{d \times k})$ and $\{\psi_l\}_{l=1}^m \subset C_{b,lip}(\mathbb{R}^d)$ are functions with disjoint supports and $\psi_l(0) = 0$.

Definition 2.10. For any $f \in \mathcal{H}_G^S([0, T] \times (\mathbb{R}^d \setminus \{0\}))$, set

$$\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, u) L(ds, du) := \sum_{0 < s \leq t} f(s, \kappa_s) I_{\mathbb{R}^d \setminus \{0\}}(\kappa_s), \quad q.s..$$

By Theorem 28 in [3], we have that $\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, u) L(ds, du) \in L_G^2(\Omega_T)$. Let $\mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}))$ be the completion of $\mathcal{H}_G^S([0, T] \times (\mathbb{R}^d \setminus \{0\}))$ with respect to the norm $\|\cdot\|_{\mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}))}$, where

$$\|f\|_{\mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}))} := \mathbb{E} \left[\int_0^T \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} |f(s, u)|^2 \nu(du) ds \right]^{1/2}, \quad f \in \mathcal{H}_G^S([0, T] \times (\mathbb{R}^d \setminus \{0\})).$$

Thus, Corollary 29 in [3] admits us to get that for $f \in \mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}))$,

$$\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, u) L(ds, du) = \sum_{0 < s \leq t} f(s, \kappa_s) I_{\mathbb{R}^d \setminus \{0\}}(\kappa_s), \quad q.s.. \quad (4)$$

Let $\mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}), \mathbb{R}^d)$ be the space of all the processes

$$f(t, u) = (f^1(t, u), f^2(t, u), \dots, f^d(t, u)), \quad f^i \in \mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\})).$$

2.5. Stochastic differential equations driven by G-Lévy processes. In the subsection, we introduce stochastic differential equations driven by G-Lévy processes and related additive functionals.

First, we introduce some notations. Let \mathbb{S}^d be the space of all $d \times d$ symmetric matrices. For $A \in \mathbb{S}^d$, set

$$G(A) := \frac{1}{2} \sup_{Q \in \mathcal{Q}} \text{tr}[QQ^* A],$$

where \mathcal{Q} is a nonempty, bounded, closed and convex subset of $\mathbb{R}^{d \times d}$. And then $G : \mathbb{S}^d \mapsto \mathbb{R}$ is a monotonic, sublinear and positive homogeneous functional([6]). We choose $\mathcal{U} \subset \mathcal{M}(\mathbb{R}^d \setminus \{0\}) \times \{0\} \times \mathcal{Q}$ satisfying (3) and still work under the framework of Subsection 2.3. So, the canonical process X_t can be represented as $X_t = B_t + X_t^d$, where B_t is a G-Brownian motion associated with \mathcal{Q} and X_t^d is a pure jump G-Lévy process associated with $\mathcal{M}(\mathbb{R}^d \setminus \{0\})$.

Next, we consider Eq.(1) and assume:

($\mathbf{H}_{b,h,\sigma,f}^1$) There exists a constant $C_1 > 0$ such that for any $t \in [0, T]$ and $x, y \in \mathbb{R}^d$,

$$|b(t, x) - b(t, y)|^2 + |h_{ij}(t, x) - h_{ij}(t, y)|^2 + \|\sigma(t, x) - \sigma(t, y)\|^2 \\ + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} |f(t, x, u) - f(t, y, u)|^2 \nu(du) \leq C_1 \rho(|x - y|^2),$$

where $\rho : (0, +\infty) \mapsto (0, +\infty)$ is a continuous, increasing and concave function so that

$$\rho(0+) = 0, \quad \int_0^1 \frac{dr}{\rho(r)} = +\infty.$$

($\mathbf{H}_{b,h,\sigma,f}^2$) There exists a constant $C_2 > 0$ such that for any $t \in [0, T]$

$$|b(t, 0)|^2 + |h_{ij}(t, 0)|^2 + \|\sigma(t, 0)\|^2 + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} |f(t, 0, u)|^2 \nu(du) \leq C_2.$$

By [14, Theorem 3.1], we know that under ($\mathbf{H}_{b,h,\sigma,f}^1$)-($\mathbf{H}_{b,h,\sigma,f}^2$), Eq.(1) has a unique solution Y_t with

$$\bar{\mathbb{E}} \left[\sup_{t \in [0, T]} |Y_t|^2 \right] < \infty. \quad (5)$$

And then we introduce the following additive functional

$$F_{s,t} := \alpha \int_s^t G(g_1)(r, Y_r) dr + \beta \int_s^t g_1^{ij}(r, Y_r) d\langle B^i, B^j \rangle_r + \int_s^t \langle g_2(r, Y_r), dB_r \rangle \\ + \int_s^t \int_{\mathbb{R}^d \setminus \{0\}} g_3(r, Y_r, u) L(dr, du) + \int_s^t \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} \gamma g_3(r, Y_r, u) \nu(du) dr, \\ 0 \leq s < t \leq T, \quad (6)$$

where $\alpha, \beta, \gamma \in \mathbb{R}$ are three constants and

$$g_1 : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times d}, \quad g_1^{ij} = g_1^{ji}, \\ g_2 : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d, \\ g_3 : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}) \mapsto \mathbb{R},$$

are Borel measurable so that $F_{s,t}$ is well-defined.

Definition 2.11. *The additive functional $F_{s,t}$ is called path independent, if there exists a function*

$$V : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R},$$

such that for any $s \in [0, T]$ and $Y_s \in L_G^2(\Omega_T)$, the solution $(Y_t)_{t \in [s, T]}$ of Eq.(1) satisfies

$$F_{s,t} = V(t, Y_t) - V(s, Y_s). \quad (7)$$

3. MAIN RESULTS AND THEIR PROOFS

In the section, we state and prove the main results under the framework of Subsection 2.5. And then we analysis some special cases and compare our result with some known results.

3.1. Main results. In the subsection, we state and prove the main results. Let us begin with a key lemma.

Lemma 3.1. *Assume that \mathcal{Q} is bounded away from 0 and Z_t is a 1-dimensional G-Itô-Lévy process, i.e.*

$$Z_t = \int_0^t \Gamma_s ds + \int_0^t \Phi_{ij}(s) d\langle B^i, B^j \rangle_s + \int_0^t \langle \Psi_s, dB_s \rangle + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} K(s, u) L(ds, du), \quad (8)$$

where $\Gamma \in \mathcal{M}_G^1(0, T)$, $\Phi_{ij} \in \mathcal{M}_G^1(0, T)$, $\Phi_{ij} = \Phi_{ji}$, $i, j = 1, 2, \dots, d$, $\Psi \in \mathcal{H}_G^1([0, T], \mathbb{R}^d)$, $K \in \mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\}))$. Then $Z_t = 0$ for all $t \in [0, T]$ q.s. if and only if $\Gamma_t = 0$, $\Phi_{ij}(t) = 0$, $\Psi_t = 0$ a.e. \times q.s. on $[0, T] \times \Omega_T$ and $K(t, u) = 0$ a.e. \times a.e. \times q.s. on $[0, T] \times (\mathbb{R}^d \setminus \{0\}) \times \Omega_T$.

Proof. Sufficiency is direct if one inserts $\Gamma_t = 0$, $\Phi_{ij}(t) = 0$, $\Psi_t = 0$, $K(t, u) = 0$ into (8). Let us prove necessity. If $Z_t = 0$ for any $t \in [0, T]$, we get that

$$0 = \int_0^t \Gamma_s ds + \int_0^t \Phi_{ij}(s) d\langle B^i, B^j \rangle_s + \int_0^t \Psi_s^* dB_s + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} K(s, u) L(ds, du). \quad (9)$$

By taking the quadratic process with $\int_0^t \langle \Psi_s, dB_s \rangle$ on two sides of (9), it holds that

$$\begin{aligned} 0 &= \left\langle \int_0^t \Psi_s^* dB_s, \int_0^t \Psi_s^* dB_s \right\rangle_t = \int_0^t \Psi_s^i \Psi_s^j d\langle B^i, B^j \rangle_s = \int_0^t \text{tr}(\Psi_s \Psi_s^* d\langle B \rangle_s) \\ &= \int_0^t \text{tr}(d\langle B \rangle_s \Psi_s \Psi_s^*) = \int_0^t \langle d\langle B \rangle_s \Psi_s, \Psi_s \rangle, \end{aligned}$$

where

$$\langle B \rangle := \begin{pmatrix} \langle B^1, B^1 \rangle & \langle B^1, B^2 \rangle & \dots & \langle B^1, B^d \rangle \\ \vdots & \vdots & & \vdots \\ \langle B^d, B^1 \rangle & \langle B^d, B^2 \rangle & \dots & \langle B^d, B^d \rangle \end{pmatrix}.$$

Note that \mathcal{Q} is bounded away from 0. Thus, there exists a constant $\iota > 0$ such that $\langle B \rangle_s \geq \iota s I_d$ and then

$$0 = \int_0^t \langle d\langle B \rangle_s \Psi_s, \Psi_s \rangle \geq \iota \int_0^t \langle \Psi_s, \Psi_s \rangle ds.$$

From this, we know that $\Psi_t = 0$ a.e. \times q.s.. So, (9) becomes

$$0 = \int_0^t \Gamma_s ds + \int_0^t \Phi_{ij}(s) d\langle B^i, B^j \rangle_s + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} K(s, u) L(ds, du).$$

Next, set

$$\tau_0 = 0, \quad \tau_n := \inf\{t > \tau_{n-1} : \kappa_t \neq 0\}, \quad n = 1, 2, \dots,$$

and $\{\tau_n\}$ is a stopping time sequence with respect to $(\mathcal{B}_t)_{t \geq 0}$ and $\tau_n \uparrow \infty$ as $n \rightarrow \infty$ q.s.(c.f. [3, Proposition 16]). So, by (4) it holds that for $t \in [0, \tau_1 \wedge T)$,

$$0 = \int_0^{\tau_1 \wedge T} \Gamma_s ds + \int_0^{\tau_1 \wedge T} \Phi_{ij}(s) d\langle B^i, B^j \rangle_s,$$

i.e.

$$-\int_0^{\tau_1 \wedge T} \Gamma_s ds = \int_0^{\tau_1 \wedge T} \Phi_{ij}(s) d\langle B^i, B^j \rangle_s.$$

Thus, by the similar deduction to that in [12, Corollary 1] one can have that

$$\bar{\mathbb{E}} \int_0^{\tau_1 \wedge T} (\text{tr}[\Phi_s \Phi_s])^{1/2} ds = \bar{\mathbb{E}} \int_0^{\tau_1 \wedge T} |\Gamma_s| ds = 0.$$

Based on this, we know that $\Phi_t = 0, \Gamma_t = 0$ for $t \in [0, \tau_1 \wedge T)$. If $\tau_1 \geq T$, the proof is over; if $\tau_1 < T$, we continue. For $t = \tau_1$, (9) goes to

$$\Phi_t = 0, \quad \Gamma_t = 0, \quad K(t, \kappa_t) = 0.$$

For $t \in [\tau_1, \tau_2 \wedge T)$, by the same means to the above for $t \in [0, \tau_1 \wedge T)$, we get that $\Phi_t = 0, \Gamma_t = 0$ for $t \in [\tau_1, \tau_2 \wedge T)$. If $\tau_2 \geq T$, the proof is over; if $\tau_2 < T$, we continue till $T \leq \tau_n$. Thus, we obtain that $\Gamma_t = 0, \Phi_{ij}(t) = 0, \Psi_t = 0$ a.e. \times q.s. on $[0, T] \times \Omega_T$ and $K(t, u) = 0$ a.e. \times a.e. \times q.s. on $[0, T] \times (\mathbb{R}^d \setminus \{0\}) \times \Omega_T$. The proof is complete. \square

The main result in the section is the following theorem.

Theorem 3.2. *Assume that \mathcal{Q} is bounded away from 0 and b, h, σ, f satisfy $(\mathbf{H}_{b,h,\sigma,f}^1)$ - $(\mathbf{H}_{b,h,\sigma,f}^2)$. Then for $V \in C_b^{1,2}([0, T] \times \mathbb{R}^d)$, $F_{s,t}$ is path independent in the sense of (7) if and only if (V, g_1, g_2, g_3) satisfies the partial integral-differential equation*

$$\begin{cases} \partial_t V(t, x) + \langle \partial_x V(t, x), b(t, x) \rangle = \alpha G(g_1)(t, x) + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} \gamma g_3(t, x, u) \nu(du), \\ \langle \partial_x V(t, x), h_{ij}(t, x) \rangle + \frac{1}{2} \langle \partial_x^2 V(t, x) \sigma^i(t, x), \sigma^j(t, x) \rangle = \beta g_1^{ij}(t, x), \\ (\sigma^T \partial_x V)(t, x) = g_2(t, x), \\ V(t, x + f(t, x, u)) - V(t, x) = g_3(t, x, u), \\ t \in [0, T], x \in \mathbb{R}^d, u \in \mathbb{R}^d \setminus \{0\}. \end{cases} \quad (10)$$

Proof. First, we prove necessity. On one hand, since $F_{s,t}$ is path independent in the sense of (7), by Definition 2.11 it holds that

$$\begin{aligned} V(t, Y_t) - V(s, Y_s) &= \alpha \int_s^t G(g_1)(r, Y_r) dr + \beta \int_s^t g_1^{ij}(r, Y_r) d\langle B^i, B^j \rangle_r + \int_s^t \langle g_2(r, Y_r), dB_r \rangle \\ &\quad + \int_s^t \int_{\mathbb{R}^d \setminus \{0\}} g_3(r, Y_r, u) L(dr, du) + \int_s^t \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} \gamma g_3(r, Y_r, u) \nu(du) dr. \end{aligned} \quad (11)$$

On the other hand, applying the Itô formula for G-Itô-Lévy processes ([3, Theorem 32]) to $V(t, Y_t)$, one can obtain that

$$\begin{aligned} V(t, Y_t) - V(s, Y_s) &= \int_s^t \partial_r V(r, Y_r) dr + \int_s^t \partial_k V(r, Y_r) b^k(r, Y_r) dr \\ &\quad + \int_s^t \partial_k V(r, Y_r) h_{ij}^k(r, Y_r) d\langle B^i, B^j \rangle_r + \int_s^t \langle (\sigma^T \partial_x V)(r, Y_r), dB_r \rangle \\ &\quad + \int_s^t \int_{\mathbb{R}^d \setminus \{0\}} (V(r, Y_r + f(r, Y_r, u)) - V(r, Y_r)) L(dr, du) \\ &\quad + \frac{1}{2} \int_s^t \partial_{kl} V(r, Y_r) \sigma^{ki}(r, Y_r) \sigma^{lj}(r, Y_r) d\langle B^i, B^j \rangle_r. \end{aligned} \quad (12)$$

By (5) $(\mathbf{H}_{b,h,\sigma,f}^1)$ - $(\mathbf{H}_{b,h,\sigma,f}^2)$, one can verify that

$$\partial_r V(r, Y_r) + \partial_k V(r, Y_r) b^k(r, Y_r) \in \mathcal{M}_G^1(0, T),$$

$$\begin{aligned}
& \partial_k V(r, Y_r) h_{ij}^k(r, Y_r) + \frac{1}{2} \partial_{kl} V(r, Y_r) \sigma^{ki}(r, Y_r) \sigma^{lj}(r, Y_r) \in \mathcal{M}_G^1(0, T), \\
& (\sigma^T \partial_x V)(r, Y_r) \in \mathcal{H}_G^1([0, T], \mathbb{R}^d), \\
& V(r, Y_r + f(r, Y_r, u)) - V(r, Y_r) \in \mathcal{H}_G^2([0, T] \times (\mathbb{R}^d \setminus \{0\})).
\end{aligned}$$

Thus, by (11) (12) and Lemma 3.1 we know that

$$\begin{cases}
\partial_r V(r, Y_r) + \langle \partial_x V(r, Y_r), b(r, Y_r) \rangle = \alpha G(g_1)(r, Y_r) + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} \gamma g_3(r, Y_r, u) \nu(du), \\
\langle \partial_x V(r, Y_r), h_{ij}(r, Y_r) \rangle + \frac{1}{2} \langle \partial_x^2 V(r, Y_r) \sigma^i(r, Y_r), \sigma^j(r, Y_r) \rangle = \beta g_1^{ij}(r, Y_r), \\
(\sigma^T \partial_x V)(r, Y_r) = g_2(r, Y_r), \\
V(r, Y_r + f(r, Y_r, u)) - V(r, Y_r) = g_3(r, Y_r, u), \quad a.e. \times q.s..
\end{cases}$$

Now, we insert $r = s, Y_s = x \in \mathbb{R}^d$ into the above equalities and get that

$$\begin{cases}
\partial_s V(s, x) + \langle \partial_x V(s, x), b(s, x) \rangle = \alpha G(g_1)(s, x) + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} \gamma g_3(s, x, u) \nu(du), \\
\langle \partial_x V(s, x), h_{ij}(s, x) \rangle + \frac{1}{2} \langle \partial_x^2 V(s, x) \sigma^i(s, x), \sigma^j(s, x) \rangle = \beta g_1^{ij}(s, x), \\
(\sigma^T \partial_x V)(s, x) = g_2(s, x), \\
V(s, x + f(s, x, u)) - V(s, x) = g_3(s, x, u), \\
s \in [0, T], x \in \mathbb{R}^d, u \in \mathbb{R}^d \setminus \{0\}.
\end{cases}$$

Since s, x are arbitrary, we have (10).

Next, we treat sufficiency. By the Itô formula for G-Itô-Lévy processes to $V(t, Y_t)$, we have (12). And then one can apply (10) to (12) to get (11). That is, $F_{s,t}$ is path independent in the sense of (7). The proof is complete. \square

3.2. Some special cases. In the subsection, we analysis some special cases.

If $b(t, x) = 0, h_{ij}(t, x) = 0, \sigma(t, x) = I_d, f(t, x, u) = u$, Eq.(1) becomes

$$dY_t = dB_t + \int_{\mathbb{R}^d \setminus \{0\}} u L(dt, du) = dX_t.$$

We take $\alpha = 1, \beta = \frac{1}{2}, \gamma = 1$. Thus, by Theorem 3.2, it holds that $F_{s,t}$ is path independent in the sense of (7) if and only if (V, g_1, g_2, g_3) satisfies the partial integral-differential equation

$$\begin{cases}
\partial_t V(t, x) = G(\partial_x^2 V)(t, x) + \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} (V(t, x + u) - V(t, x)) \nu(du), \\
\partial_x^2 V(t, x) = g_1(t, x), \\
\partial_x V(t, x) = g_2(t, x), \\
V(t, x + u) - V(t, x) = g_3(t, x, u), \\
t \in [0, T], x \in \mathbb{R}^d, u \in \mathbb{R}^d \setminus \{0\}.
\end{cases}$$

Besides, by Theorem 2.7, it holds that for $\phi \in C_{b, lip}(\mathbb{R}^d)$, $V(t, x) = \bar{\mathbb{E}}[\phi(x + X_t)]$ is the unique viscosity solution of the following partial integro-differential equation:

$$\partial_t V(t, x) - G(\partial_x^2 V)(t, x) - \sup_{\nu \in \mathcal{V}} \int_{\mathbb{R}^d \setminus \{0\}} (V(t, x + u) - V(t, x)) \nu(du) = 0$$

with the initial condition $V(0, x) = \phi(x)$. So,

$$\begin{aligned}
g_1(t, x) &= \partial_x^2 \bar{\mathbb{E}}[\phi(x + X_t)], \\
g_2(t, x) &= \partial_x \bar{\mathbb{E}}[\phi(x + X_t)],
\end{aligned}$$

$$g_3(t, x, u) = \bar{\mathbb{E}}[\phi(x + u + X_t)] - \bar{\mathbb{E}}[\phi(x + X_t)].$$

That is, we can find g_1, g_2, g_3 .

If $d = 1, \alpha = 1, \beta = 0, \gamma = 0$, it follows from Theorem 3.2 that $F_{s,t}$ is path independent in the sense of (7) if and only if (V, g_1, g_2, g_3) satisfies the partial integral-differential equation

$$\begin{cases} \partial_t V(t, x) + \partial_x V(t, x)b(t, x) = G(g_1)(t, x), \\ \partial_x V(t, x)h(t, x) + \frac{1}{2}\partial_x^2 V(t, x)\sigma^2(t, x) = 0, \\ (\sigma\partial_x V)(t, x) = g_2(t, x), \\ V(t, x + f(t, x, u)) - V(t, x) = g_3(t, x, u), \\ t \in [0, T], x \in \mathbb{R}, u \in \mathbb{R} \setminus \{0\}. \end{cases}$$

And then if $\sigma(t, x) \neq 0$, the unique solution of the above second equation is

$$V(t, x) = V(t, 0) + \partial_x V(t, 0) \int_0^x e^{-2 \int_0^z \frac{h(t,v)}{\sigma^2(t,v)} dv} dz.$$

Besides, since \mathcal{Q} is bounded away from 0, G is invertible. Thus,

$$\begin{aligned} g_1(t, x) &= G^{-1} \left(\partial_t V(t, x) + b(t, x) \partial_x V(t, 0) e^{-2 \int_0^x \frac{h(t,v)}{\sigma^2(t,v)} dv} \right), \\ g_2(t, x) &= \sigma(t, x) \partial_x V(t, 0) e^{-2 \int_0^x \frac{h(t,v)}{\sigma^2(t,v)} dv}, \\ g_3(t, x, u) &= \partial_x V(t, 0) \int_x^{x+f(t,x,u)} e^{-2 \int_0^z \frac{h(t,v)}{\sigma^2(t,v)} dv} dz. \end{aligned}$$

That is, we also give out g_1, g_2, g_3 in the case.

Remark 3.3. *In the above special cases, we can describe concretely g_1, g_2, g_3 . This is interesting and also is one of our motivations.*

3.3. Comparison with some known results. In the subsection, we compare our result with some known results.

First, if we take $f(t, x, u) = 0$ in Eq.(1) and $g_3(t, x, u) = 0$ in (6), Theorem 3.2 becomes [12, Theorem 2]. Therefore, our result is more general.

Second, we take $\mathcal{M}(\mathbb{R}^d \setminus \{0\}) = \{\nu\}$, $\mathcal{Q} = \{I_d\}$ in Subsection 2.5. Thus, B is a classical Brownian motion with $\langle B^i, B^j \rangle_t = I_{i=j}t$ and $L(dt, du)$ is a classical Poisson random measure. And then Eq.(1) goes into

$$dY_t = b(t, Y_t)dt + \sum_{i=1}^d h_{ii}(t, Y_t)dt + \sigma(t, Y_t)dB_t + \int_{\mathbb{R}^d \setminus \{0\}} f(t, Y_t, u)L(dt, du). \quad (13)$$

Note that $\nu(\mathbb{R}^d \setminus \{0\}) < \infty$. So, by (5) $(\mathbf{H}_{b,h,\sigma,f}^1)$ -($\mathbf{H}_{b,h,\sigma,f}^2$), [4, Theorem 13] admits us to obtain that

$$\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, Y_s, u) \tilde{L}(ds, du) := \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, Y_s, u) L(ds, du) - \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(s, Y_s, u) \nu(du) ds$$

is a \mathcal{B}_t -martingale, where $\mathcal{B}_t := \sigma\{\omega_s, 0 \leq s \leq t\}$, $0 \leq t \leq T$. Therefore, we can rewrite Eq.(13) to get that

$$dY_t = b(t, Y_t)dt + \sum_{i=1}^d h_{ii}(t, Y_t)dt + \int_{\mathbb{R}^d \setminus \{0\}} f(t, Y_t, u) \nu(du)dt + \sigma(t, Y_t)dB_t$$

$$+ \int_{\mathbb{R}^d \setminus \{0\}} f(t, Y_t, u) \tilde{L}(dt, du).$$

This is a classical stochastic differential equation with jumps. Thus, by [7, Theorem 1.2] it holds that under $(\mathbf{H}_{b,h,\sigma,f}^1)-(\mathbf{H}_{b,h,\sigma,f}^2)$, the above equation has a unique solution.

In the following, note that $G(g_1) = \frac{1}{2}\text{tr}(g_1) = \frac{1}{2} \sum_{i=1}^d g_1^{ii}$. And then $F_{s,t}$ can be represented as

$$\begin{aligned} F_{s,t} = & \int_s^t \left(\frac{\alpha}{2} + \beta \right) \sum_{i=1}^d g_1^{ii}(r, Y_r) dr + \int_s^t \langle g_2(r, Y_r), dB_r \rangle + \int_s^t \int_{\mathbb{R}^d \setminus \{0\}} g_3(r, Y_r, u) \tilde{L}(dr, du) \\ & + \int_s^t \int_{\mathbb{R}^d \setminus \{0\}} (1 + \gamma) g_3(r, Y_r, u) \nu(du) dr. \end{aligned}$$

This is just right [11, (3)] without the distribution of Y_r for $r \in [s, t]$. So, in the case Definition 2.11 and Theorem 3.2 are Definition 2.1 and Theorem 3.2 in [11] without the distribution of Y_r for $r \in [s, t]$, respectively. Therefore, our result overlaps [11, Theorem 3.2] in some sense.

Acknowledgements:

The authors are very grateful to Professor Xicheng Zhang for valuable discussions. The first author also thanks Professor Renming Song for providing her an excellent environment to work in the University of Illinois at Urbana-Champaign.

REFERENCES

- [1] F. Gao: Pathwise properties and hoemomorphic flows for stochastic differential equations driven by G-Brownian motion, *Stochastic Processes and their Applications*, 119(2009)3356-3382.
- [2] M. Hu and S. Peng: G-Lévy processes under sublinear expectations. arXiv:0911.3533v1.
- [3] K. Paczka: It calculus and jump diffusions for G-Lévy processes. arXiv:1211.2973v3.
- [4] K. Paczka: G-martingale representation in the G-Lévy setting. arXiv:1404.2121v1.
- [5] S. Peng: Multi-dimensional G-Brownian motion and related stochastic calculus under G-expectation, *Stochastic Processes and their Applications*, 118(2008)2223-2253.
- [6] S. Peng: *Nonlinear Expectations and Stochastic Calculus under Uncertainty with Robust CLT and G-Brownian Motion*. Probability Theory and Stochastic Modelling 95, Springer, 2019.
- [7] H. Qiao: Euler-Maruyama approximation for SDEs with jumps and non-Lipschitz coefficients, *Osaka Journal of Mathematics*, 51(2014)47-66.
- [8] H. Qiao: The cocycle property of stochastic differential equations driven by G-Brownian motion, *Chinese Annals of Mathematics, Series B*, 36(2015)147-160.
- [9] H. Qiao and J.-L. Wu: Characterising the path-independence of the Girsanov transformation for non-Lipschitz SDEs with jumps, *Statistics and Probability Letters*, 119(2016)326-333.
- [10] H. Qiao and J.-L. Wu: On the path-independence of the Girsanov transformation for stochastic evolution equations with jumps in Hilbert spaces, *Discrete and Continuous Dynamical Systems-B*, 24(2019)1449-1467.
- [11] H. Qiao and J.-L. Wu: Path independence of the additive functionals for McKean-Vlasov stochastic differential equations with jumps, <http://arxiv.org/abs/1911.03830>.
- [12] P. Ren and F. Yang: Path independence of additive functionals for stochastic differential equations under G-framework, *Front. Math. China* 14 (2019), no. 1, 135-148.
- [13] A. Truman, F.-Y. Wang, J.-L. Wu, W. Yang: A link of stochastic differential equations to nonlinear parabolic equations, *SCIENCE CHINA Mathematics*, 55 (2012) 1971-1976.
- [14] B. Wang and H. Gao: Exponential stability of solutions to stochastic differential equations driven by G-Lévy process, appear in *Applied Mathematics & Optimization*.

- [15] J.-L. Wu and W. Yang: On stochastic differential equations and a generalised Burgers equation, pp 425-435 in *Stochastic Analysis and Its Applications to Finance: Essays in Honor of Prof. Jia-An Yan* (eds. T S Zhang, X Y Zhou), Interdisciplinary Mathematical Sciences, Vol. 13, World Scientific, Singapore, 2012.