

The bounds for the number of linear extensions via chain and antichain coverings

I. A. Bochkov*, F. V. Petrov†

April 9, 2024

Abstract

Let (\mathcal{P}, \leq) be a finite poset. Define the numbers a_1, a_2, \dots (respectively, c_1, c_2, \dots) so that $a_1 + \dots + a_k$ (respectively, $c_1 + \dots + c_k$) is the maximal number of elements of \mathcal{P} which may be covered by k antichains (respectively, k chains.) Then the number $e(\mathcal{P})$ of linear extensions of poset \mathcal{P} is not less than $\prod a_i!$ and not more than $n! / \prod c_i!$. A corollary: if \mathcal{P} is partitioned onto disjoint antichains of sizes b_1, b_2, \dots , then $e(\mathcal{P}) \geq \prod b_i!$.

1 Introduction

Let (\mathcal{P}, \leq) be a finite poset.

In a recent paper [6] the following double inequality for the number $e(\mathcal{P})$ of linear extensions of \mathcal{P} is applied. Partition \mathcal{P} onto disjoint antichains $\mathcal{P} = A_1 \sqcup A_2 \sqcup A_3 \dots$, where A_i is the antichain of elements with rank i . Also partition \mathcal{P} in arbitrary way onto disjoint chains $\mathcal{P} = C_1 \sqcup C_2 \sqcup C_3 \dots$. Then

$$\frac{n!}{\prod |C_i|!} \geq e(\mathcal{P}) \geq \prod |A_i|! \quad (1)$$

To quote [6]: “These bounds are probably folklore; for the lower bound see e.g [5].”

We prove that the right inequality in (1) holds for arbitrary antichain partition. We also improve both inequalities in terms of Greene–Kleitman–Fomin parameters, which we define now.

The *antichain Greene–Kleitman–Fomin parameters* $a_1 \geq a_2 \geq \dots$ of a finite poset \mathcal{P} are defined as follows: $a_1 + \dots + a_k$ is the maximal number of elements of \mathcal{P} which may be covered by k antichains ($k = 1, 2, \dots$). The fact that the sequence (a_i) is weakly decreasing is a part of Greene–Kleitman–Fomin theorem [1, 2]. Another claim of this theorem is that for the partition $n = c_1 + c_2 + \dots$ conjugate to the partition $n = a_1 + a_2 + \dots$ the sum $c_1 + \dots + c_k$ is the maximal number of elements of \mathcal{P} which may be covered by k chains. The numbers c_1, c_2, \dots are called *chain Greene–Kleitman–Fomin parameters* of poset \mathcal{P} .

The main result of this paper is

Theorem 1.

$$\frac{n!}{\prod c_i!} \geq e(\mathcal{P}) \geq \prod a_i! \quad (2)$$

*St. Petersburg State University, bycha@yandex.ru

†St. Petersburg State University, St. Petersburg Department of V. A. Steklov Mathematical Institute RAS, f.v.petrov@spbu.ru. The work was supported by the Foundation for the Advancement of Theoretical Physics and Mathematics “BASIS”.

2 Majorization lemmata

Let X be a finite multiset consisting of non-negative numbers. For non-negative integer k define $s_k(X)$ as the sum of $\min(k, |X|)$ maximal elements of multiset X ; also denote $s(X) = s_{|X|}(X)$ the sum of all elements of X . We say that multiset X *majorizes* another multiset Y of non-negative numbers and write $X \succ Y$ if $s(X) = s(Y)$ and $s_k(X) \geq s_k(Y)$ for all k .

We need the following version of Karamata's majorization inequality.

Theorem (Karamata inequality). *Let $f(x) : \{0, 1, \dots\} \rightarrow (0, +\infty)$ be a log-convex function, i.e., $f(0) = 1$ and $f(x+1)/f(x)$ is an increasing function of x . Next, let $X, Y \subset \{0, 1, \dots\}$ be two finite multisets. Then X majorizes Y if and only if $\prod_X f \geq \prod_Y f$ for any log-convex function f .*

This theorem immediately yields

Proposition 2. *For multisets $X, Y, Z \subset \{0, 1, \dots\}$ the conditions $X \succ Y$ and $X \cup Z \succ Y \cup Z$ are equivalent.*

Further we use Karamata inequality for the log-convex function $f(x) = x!$

We also use the following simple fact

Proposition 3. *Let X, Y be two finite multisets consisting of non-negative integers, and the sum of elements in Y is by 1 less than the sum of elements in X : $s(Y) = s(X) - 1$. Denote by x_i and y_i there i -th largest elements, respectively. Also denote $x_i = 0$ for $i > |X|$, $y_i = 0$ for $i > |Y|$. (In these notations we have $s_k(X) = \sum_{i \leq k} x_i$, $s_k(Y) = \sum_{i \leq k} y_i$.)*

Assume that $s_k(X) \geq s_k(Y) \geq s_k(X) - 1$ for all k , and there exists m such that $s_k(X) = s_k(Y)$ for all $k = 1, 2, \dots, m-1$. Then $x_m > 0$ and $Y \succ (X \cup \{x_m - 1\}) \setminus x_m$.

Proof. We get $y_i = x_i$ for $i = 1, 2, \dots, m-1$. Consider the minimal j such that $y_j \neq x_j$. Then $j \geq m$ and denoting $\varepsilon = s_j(X) - s_j(Y) \in \{0, 1\}$ we have

$$y_j = s_j(Y) - s_{j-1}(Y) = s_j(Y) - s_{j-1}(X) = s_j(X) - \varepsilon - s_{j-1}(X) = x_j - \varepsilon,$$

therefore $\varepsilon = 1$ and $y_j = x_j - 1$. So we get $x_m \geq x_j = y_j + 1 > 0$.

Assume that $x_{j+1} = x_j$. Then $y_{j+1} \leq y_j = x_j - 1 = x_{j+1} - 1$ and $s_{j+1}(Y) \leq s_{j+1}(X) - 2$, a contradiction. Therefore $x_{j+1} \leq x_j - 1$, and Y majorizes multiset $X_j := (X \cup \{x_j - 1\}) \setminus x_j$: indeed, $s_k(Y) = s_k(X_j) = s_k(X)$ for $k \leq j-1$, and $s_k(Y) \geq s_k(X) - 1 = s_k(X_j)$ for $k \geq j$.

Since $j \geq m$, we have $\{x_j - 1, x_m\} \succ \{x_m - 1, x_j\}$, and by Proposition 2 we get

$$Y \succ X_j = (X \setminus \{x_m, x_j\}) \cup \{x_j - 1, x_m\} \succ (X \setminus \{x_m, x_j\}) \cup \{x_m - 1, x_j\} = X_m,$$

as needed. □

3 Lower bound

Let (\mathcal{P}, \leq) be a finite poset, $A \subset \mathcal{P}$ be an antichain. The proof of the main result of [3] implies the inequality

$$e(\mathcal{P}) \geq \sum_{x \in A} e(\mathcal{P} - x) \quad (3)$$

for the number of linear extensions $e(\cdot)$. Inequality (3) is also proved differently in [4]. In [3] it was proved that (3) turns into equality if antichain A have non-empty intersection with any maximal chain.

For sake of completeness we prove (3). Define an injection from linear extensions of posets $\mathcal{P} - x$, where $x \in A$, to linear extensions of poset \mathcal{P} . A linear extension of $\mathcal{P} - x$, $x \in \mathcal{P}$, is understood as an order-preserving bijection $f : \mathcal{P} - x \mapsto \{2, 3, \dots, n\}$, where $n = |\mathcal{P}|$. For such f we construct an order-preserving bijection $\phi f : \mathcal{P} \mapsto \{1, 2, \dots, n\}$ as follows. Consider the *greedy falling chain from x* : put $x_0 = x$, define x_{i+1} as an element of \mathcal{P} which is less than x_i for which $f(x_i)$ is maximal possible. We get a chain $x_0 > x_1 > \dots > x_t$.

Let's shift the values along this chain. Namely, put $\phi f(x_i) = f(x_{i+1})$ for $i = 0, 1, \dots, t-1$, $\phi f(x_t) = 1$. For $y \notin \{x_0, \dots, x_t\}$ put $\phi f(y) = f(y)$. Note that ϕf is order-preserving due to the greedy property. Indeed, if $y \{x_0, \dots, x_t\}$ and $x_i > y$, then by greediness we get $f(y) \leq f(x_{i+1}) = \phi f(y)$, other inequalities are obvious.

It is straightforward that for the map ϕf the chain $x_t < x_{t-1} < \dots < x_0$ is the *greedy increasing chain*: $x_t = (\phi f)^{-1}(1)$, and each next element realizes the minimal possible value of ϕf .

Therefore, if the maps $f : \mathcal{P} - x \mapsto \{2, 3, \dots, n\}$ and $g : \mathcal{P} - y \mapsto \{2, 3, \dots, n\}$ satisfy $\phi f = \phi g$, then x, y belong to the greedy increasing chain of ϕf . If x, y also belong to the antichain A , then we get $x = y$ and $f = g$.

Injectivity of φ is proved, it implies inequality (3).

Proof of lower bound in (2). Induction on $n = |\mathcal{P}|$. Base $n = 1$ is obvious.

Step from $n - 1$ to n .

Let A be a maximal antichain in \mathcal{P} , then $|A| = a_1$. Fix $x \in A$. Let $r_1 \geq r_2 \dots$ denote antichain Greene–Kleitman–Fomin parameters for $\mathcal{P} - x$. Due to Proposition 3 the multiset $\{r_1, r_2, \dots\}$ majorizes the multiset $\{a_1 - 1, a_2, a_3, \dots\}$. By Karamata inequality for log-convex function $f(x) = x!$ we get

$$\prod_{i \geq 1} r_i! \geq (a_1 - 1)! \prod_{i \geq 1} a_i! = \frac{1}{a_1} \prod_i a_i!$$

Therefore by induction proposition we have $e(\mathcal{P} - x) \geq \frac{1}{a_1} \prod_i a_i!$ For all $x \in A$. Summing up over all $x \in A$ and using (3) we see that indeed $e(\mathcal{P}) \geq \prod_i a_i!$, as needed. \square

Corollary. *Let \mathcal{P} be partitioned onto antichains of sizes c_1, c_2, \dots . Then $e(\mathcal{P}) \geq \prod_i c_i!$.*

Proof. It suffices to note that multiset $\{c_1, c_2, \dots\}$ is majorized by the multiset $\{a_1, a_2, \dots\}$ and apply Karamata inequality for factorial. Alternatively, we may prove this claim directly by induction using (3). \square

4 Upper bound

For bounding the number of linear extensions from above we need

Lemma 4. *Let \mathcal{P} be a finite poset, $A \subset \mathcal{P}$ be the antichain of maximal elements in \mathcal{P} , $c_1 \geq c_2 \geq \dots$ be the chain Greene–Kleitman–Fomin parameters of \mathcal{P} . Then the elements of A may be enumerated as $x_1, x_2, \dots, x_{|A|}$ so that for all $i = 1, 2, \dots, |A|$ there exist i chains whose maximal elements belong to the set $\{x_1, \dots, x_i\} \cup (\mathcal{P} \setminus A)$ with total size $c_1 + \dots + c_i$.*

Proof. Assume that the elements x_1, \dots, x_i are already chosen and satisfy the conditions of Lemma. Contract the set $A \setminus \{x_1, \dots, x_i\}$ to a new one element t , denote the new poset \mathcal{Q} . The first i chain Greene–Kleitman–Fomin parameters of \mathcal{Q} are the same as for \mathcal{P} . Denote the $(i+1)$ -th parameter by α . We should prove that $\alpha = c_{i+1}$: it allows to choose appropriate x_{i+1} . Assume that on the contrary $\alpha < c_{i+1}$. Then by Greene–Kleitman–Fomin duality we may find i chains in \mathcal{Q} of total size $c_1 + \dots + c_i$ (not containing t) and α antichains so that they cover all elements of \mathcal{Q} and each chain intersects each antichain. Note that the antichain containing t remains an antichain if we replace \mathcal{Q} back to \mathcal{P} (by splitting the element t). But then $i+1$ chains in \mathcal{P} may cover at most $c_1 + \dots + c_i + \alpha$ elements: at most $\alpha(i+1)$ elements may be covered by α antichains, and exactly $c_1 + \dots + c_i - i\alpha$ elements remain. A contradiction. \square

Proof of the upper bound in (2). By induction, we suppose the inequality proved for posets with $n-1$ elements (the base $n=1$ is obvious). Enumerate antichain A of maximal elements as in Lemma 4. Let $r_1 \geq r_2 \geq \dots$ be chain Greene–Kleitman–Fomin parameters of the poset $\mathcal{P} - x_i$. It is clear that for all j we have $c_1 + \dots + c_j \geq r_1 + \dots + r_j \geq c_1 + \dots + c_j - 1$, and for $j \leq i-1$ the equality $r_1 + \dots + r_j = c_1 + \dots + c_j$ holds. Therefore using Proposition 3 we conclude that the multiset $\{r_1, r_2, \dots\}$ majorizes the multiset $\{c_1, c_2, \dots, c_{j-1}, c_j - 1, c_{j+1}, \dots\}$. By Karamata inequality for factorial we have $\prod r_j! \geq \frac{1}{c_j} \prod_j c_j!$, and using induction proposition we get

$$e(\mathcal{P} - x_i) \leq \frac{(n-1)!}{\prod r_j!} \leq \frac{c_j(n-1)!}{\prod c_j!}.$$

Sum up this by all i and apply the obvious equality $e(\mathcal{P}) = \sum_i e(\mathcal{P} - x_i)$ we complete the induction step. \square

5 Accuracy of the bounds

The upper and lower bounds in (2) are close enough: their ratio is always $e^{O(n \log \log n)} = n^{o(1)}$ (but alas worse than exponential in n). This follows from the following general inequality.

Theorem 5. *For any two conjugate partitions $a_1 \geq a_2 \geq \dots$ and $c_1 \geq c_2 \geq \dots$ of the positive integer n onto positive integer parts the inequality*

$$\prod_i a_i! \prod_i c_i! \geq \left(\frac{n}{eH_n} \right)^n = n! e^{-n \log \log n + O(\log n)}$$

holds (here $H_n = 1 + 1/2 + \dots + 1/n = \log n + O(1)$ is a Harmonic sum).

Proof. We have

$$\prod_i c_i! = \prod_k k^{|i: c_i \geq k|} = \prod_k k^{a_k},$$

thus the inequality to prove is

$$\prod_k k^{a_k} a_k! \geq \left(\frac{n}{eH_n} \right)^n, \quad (4)$$

which we now prove for arbitrary non-negative integers a_1, a_2, \dots, a_n which sum up to n . Without loss of generality a_1, a_2, \dots, a_n are chosen so that the left hand side of (4) is

minimal possible. Choose two indices k, ℓ such that $a_\ell > 0$ and try to replace a_k to $a_k + 1$, a_ℓ to $a_\ell - 1$. Left hand side of (4) is multiplied by $k(a_k + 1)/(\ell a_\ell)$, thus $k(a_k + 1) \geq \ell a_\ell$. This last inequality holds for $a_\ell = 0$ too. Sum up the inequalities $k(a_k + 1) \cdot \frac{1}{\ell} \geq a_\ell$ over $\ell = 1, 2, \dots, n$ we get $k(a_k + 1)H_n \geq a_1 + \dots + a_n = n$. Next, using the inequality $a! \geq (\frac{a+1}{e})^a$ which holds for all non-negative integer a (it may be proved by induction, for example) we get

$$\prod_k k^{a_k} a_k! \geq \prod_k \left(\frac{k(a_k + 1)}{e} \right)^{a_k} \geq \prod_k \left(\frac{n}{eH_n} \right)^{a_k} = \left(\frac{n}{eH_n} \right)^n,$$

as needed. □

References

- [1] C. Greene, D. J. Kleitman. The structure of Sperner k-families. J. Combin. Th. A, 20 (1976), pp.41-68.
- [2] S. V. Fomin, Finite partially ordered sets and Young tableaux, Soviet Math. Dokl. 19 (1978), 1510-1514.
- [3] Paul H. Edelman, Takayuki Hibi, Richard Stanley. A recurrence for linear extensions. Order 6(1989), no. 1, pp. 15-18.
- [4] A. Sidorenko. Inequalities for the number of linear extensions. Order 8 (1992), no. 4, pp. 331-340.
- [5] G.R. Brightwell. The number of linear extensions of ranked posets. LSE CDAM Res. Report 18 (2003), p. 6.
- [6] A. H. Morales, I. Pak, G. Panova. Asymptotics of the number of standard Young tableaux of skew shape. European Journal of Combinatorics 70 (2018), pp. 26-49.