

Optical properties of periodically-driven open nonequilibrium quantum systems

Gabriel Cabra,¹ Ignacio Franco,^{2,3} and Michael Galperin^{1,*}

¹*Department of Chemistry & Biochemistry, University of California San Diego, La Jolla, CA 92093, USA*

²*Department of Chemistry, University of Rochester, Rochester, NY 14627, USA*

³*Department of Physics, University of Rochester, Rochester, NY 14627, USA*

Characterization and control of matter by optical means is at the forefront of research both due to fundamental insights and technological promise. Theoretical modeling of periodically driven systems is a prerequisite to understanding and engineering nanoscale quantum devices for quantum technologies. Here, we develop a theory for transport and optical response of molecular junctions, open nonequilibrium quantum systems, under external periodic driving. Periodic driving is described using the Floquet theory combined with nonequilibrium Green's function description of the system. Light-matter interaction is modeled employing the self-consistent Born approximation. Generic three-level model is utilized to illustrate effect of the driving on optical and transport properties of junctions.

I. INTRODUCTION

Recent progress in laser techniques combined with technological advances at the nanoscale resulted in the possibility of optical measurements in current carrying single-molecule junctions. In particular, bias induced fluorescence [1–3] and phosphorescence [4] were reported in the literature. Electroluminescence allowed to visualize intra-molecular interactions [5, 6], explore real space energy transfer [7], and probe charge fluctuations in biased molecular junctions [8]. Raman spectroscopy yields information on vibrational structure and electron flux induced heating of vibrational and electronic degrees of freedom in single-molecule junctions [9–12]. Recently, experimental measurements of ultra-strong light-matter interaction in single molecule cavities (so far without current) were reported in the literature [13, 14]. Combination of molecular electronics and optical spectroscopy evolved into new direction of research coined molecular optoelectronics [15, 16].

Control of response in nanoscale systems by external driving is an active area of research both experimentally and theoretically due to technological promise of engineering and control in quantum devices. For example, periodic radio frequency potential modulations in a suspended carbon nanotube junction facilitated experimental studies of strong coupling between tunneling electron and nanomechanical motion [17]. AC-driven charging and discharging of quantum dot was employed in experimental verification of existence of quantum stochastic resonance, where intrinsic fluctuations lead to amplification and optimization of a weak signal [18]. In single molecule junctions, periodic driving by set of laser pulse pairs was suggested as a tool to study sub-picosecond intra-molecular dynamics [19, 20].

Here we consider a molecular junction driven by a time-periodic external field. The effective electronic properties of this driven system are characterized by de-

veloping a theory of its optical and transport response in this highly nonequilibrium setting. Theoretical studies of periodic driving in junctions usually employ quantum master equation [21–25] and nonequilibrium Green's function [26–28] approaches. Periodicity in external driving allows mapping of the original time-dependent problem into effective time-independent formulation in an extended Floquet space [29]. The Floquet theory was used in numerous theoretical studies of external driving on transport in junctions. Studies employing combination of the theory with Schrödinger equation [30], scattering matrix approach [31], quantum master equation [32–35], and Green's functions [36–40] are available in the literature. For strongly correlated systems combinations of the Floquet with slave boson [41], dynamical mean field theory [42], and functional renormalization group [43] were also formulated.

Note that for molecular junctions characteristic strength of molecular coupling to contacts $\Gamma \sim 0.01 - 0.1$ eV [44] is of the same order of magnitude or stronger than thermal energy $k_B T \sim 0.01$ eV. Thus, Redfield/Lindblad quantum master equation, which relies on assumption $\Gamma \ll k_B T$, is not suitable and utilization of nonequilibrium Green's function (NEGF) technique is preferable. We consider a case where light-matter interaction is smaller than the system-bath coupling, so that diagrammatic perturbation theory can be employed, and treat the interaction within the self-consistent Born approximation (SCBA). Periodicity of driving makes the Floquet theory useful. Here, we formulate a Floquet-NEGF-SCBA theory of transport and optical response of single-molecule junctions under external periodic driving. The formulation complements study of optical properties of current carrying junctions [45] by accounting for periodic external driving and extends recent study on optical absorption properties of driven matter [46] to realm of open nonequilibrium molecular systems.

The structure of the paper is the following. In Section II we introduce junction model and formulate Floquet version of the NEGF-SCBA treatment of electron and photon fluxes in the junction. We use the Floquet-NEGF-SCBA to illustrate effects of quantum coherence

* migalperin@ucsd.edu

on junction responses in Section III. Section IV concludes and outlines directions for future research.

II. THEORY

Here we present a model of a junction and briefly describe the nonequilibrium Green's function (NEGF) approach to simulation of photon and electron fluxes, where we treat light-matter interaction at the level of the self-consistent Born approximation (SCBA). After this we present the Floquet formulation of the NEGF-SCBA problem.

A. Model

We consider a junction consisting of a molecule M coupled to two contacts L and R and to quantum modes $\{\alpha\}$ of radiation field pt . For simplicity, molecule is modeled as a non-interacting (mean-field or DFT) system. Note, this is the usual [47–49] although not always reliable [50] level of modeling in first principles simulations. In addition, the molecule is subjected to external driving by monochromatic classical field $\vec{E}(t)$

$$\vec{E}(t) = \vec{E}_0 \cos(\omega_0 t + \phi_0) \quad (1)$$

Contacts are modeled as reservoirs of free electrons each at its own equilibrium (see Fig. 1). The external driving distorts the electronic structure of the junction leading to effective properties that can in principle be very different from those observed at equilibrium.

Hamiltonian of the model is (here and below $\hbar = e = k_B = 1$)

$$\hat{H}(t) = \hat{H}_M(t) + \sum_{B=L,R,pt} \left(\hat{H}_B + \hat{V}_{MB} \right) \quad (2)$$




FIG. 1. Sketch of a molecular junction subjected to external driving.

where $\hat{H}_M$ and $\hat{H}_B$ ($B = L, R, pt$) represent decoupled molecule and baths (contacts and radiation field), while $\hat{V}_{MB}$ are the couplings. Explicit expressions are

$$\begin{aligned} \hat{H}_M(t) &= \sum_{m_1, m_2 \in M} \left(H_{m_1 m_2}^M - \vec{\mu}_{m_1 m_2} \cdot \vec{E}(t) \right) \hat{d}_{m_1}^\dagger \hat{d}_{m_2} \\ \hat{H}_{L(R)} &= \sum_{k \in L(R)} \varepsilon_k \hat{c}_k^\dagger \hat{c}_k; \quad \hat{H}_{pt} = \sum_{\alpha} \omega_{\alpha} \hat{a}_{\alpha}^\dagger \hat{a}_{\alpha} \\ \hat{V}_{ML(R)} &= \sum_{m \in M} \sum_{k \in L(R)} \left(V_{km} \hat{c}_k^\dagger \hat{d}_m + H.c. \right) \\ \hat{V}_{M,pt} &= \sum_{m_1, m_2 \in M} \sum_{\alpha} \left(V_{\alpha, m_1 m_2}^{pt} \hat{a}_{\alpha}^\dagger \hat{d}_{m_1}^\dagger \hat{d}_{m_2} + H.c. \right) \end{aligned} \quad (3)$$

Here $\hat{d}_m^\dagger$ ($\hat{d}_m$) and $\hat{c}_k^\dagger$ ($\hat{c}_k$) create (annihilate) electron in orbital m of the molecule and state k of the contacts, respectively. $\hat{a}_{\alpha}^\dagger$ ($\hat{a}_{\alpha}$) creates (destroys) excitation quanta in mode α of the radiation field. $\mathbf{H}^M$ is molecular (or Kohn-Sham) orbitals basis representation of the effective single particle Hamiltonian of the molecule. ε_k and ω_{α} are energy of free electron in state k in contacts and frequency of radiation field mode α , respectively. V_{km} is the matrix element for electron transfer from molecular orbital m to state k in contacts. $V_{\alpha, m_1 m_2}^{pt}$ is the matrix element for intra-molecular (optical) electron transfer from orbital m_2 to orbital m_1 while creating excitation quanta (photon) in mode α . We stress that coupling to the radiation field in (3) is completely general, i.e. it is not restricted to the rotating wave approximation.

Below for simplicity we assume

$$V_{m_1 m_2, \alpha}^{pt} = U_{m_1 m_2} [V_{\alpha}^{pt}]^* \quad (4)$$

where $\mathbf{U}$ is the matrix in molecular orbital space with 1 for allowed optical transitions and 0 otherwise.

B. Photon and electron fluxes within NEGF-SCBA

The nonequilibrium Green's function (NEGF) in the subspace of molecular orbitals is defined on the Keldysh contour as

$$G_{m_1 m_2}(\tau_1, \tau_2) = -i \langle T_c \hat{d}_{m_1}(\tau_1) \hat{d}_{m_2}^\dagger(\tau_2) \rangle \quad (5)$$

where T_c is the contour ordering operator and $\tau_{1,2}$ are the contour variables. Retarded (r) and lesser ($<$)/greater ($>$) projections of the Greens' function (5) are usually obtained by solving differential (or Kadanoff-Baym) and integral (or Keldysh) forms of the Dyson equation, re-

spectively [51, 52].

$$\begin{aligned} & \left(i \mathbf{I} \frac{\partial}{\partial t_1} - \mathbf{H}^M - \vec{\mu} \cdot \vec{E}(t_1) \right) \mathbf{G}^r(t_1, t_2) \\ & - \int dt \Sigma^r(t_1, t) \mathbf{G}^r(t, t_2) = \mathbf{I} \delta(t_1 - t_2) \quad (6) \\ \mathbf{G}^{\lessgtr}(t_1, t_2) = & \\ & \int dt_3 \int dt_4 \mathbf{G}^r(t_1, t_3) \Sigma^{\lessgtr}(t_3, t_4) \mathbf{G}^a(t_4, t_2) \end{aligned}$$

Here, Hamiltonian $\mathbf{H}^M$, molecular dipole moment μ , Green's function $\mathbf{G}$ and self-energy Σ projections are written as matrices in the subspace of molecular orbitals. $\mathbf{I}$ is the unity matrix in the subspace and $\mathbf{G}^a(t_4, t_2) \equiv [\mathbf{G}^r(t_2, t_4)]^\dagger$ is the advanced projection of the Green's function (5). The self-energy accounts for contributions of the baths (contacts and radiation field) into molecular dynamics

$$\begin{aligned} \Sigma_{m_1 m_2}(\tau_1, \tau_2) &= \sum_{B=L, R, rad} \Sigma_{m_1 m_2}^B(\tau_1, \tau_2) \\ \Sigma_{m_1 m_2}^{L(R)}(\tau_1, \tau_2) &= \sum_{k \in L(R)} V_{m_1 k} g_k(\tau_1, \tau_2) V_{k m_2} \quad (7) \\ \Sigma_{m_1 m_2}^{pt}(\tau_1, \tau_2) &= \sum_{m_3, m_4 \in M} G_{m_3 m_4}(\tau_1, \tau_2) \\ &\times \left(U_{m_3 m_1} F(\tau_1, \tau_2) U_{m_4 m_2} + U_{m_4 m_2} F(\tau_2, \tau_1) U_{m_1 m_3} \right) \end{aligned}$$

where

$$F(\tau_1, \tau_2) \equiv \sum_{\alpha} |V_{\alpha}^{pt}|^2 f_{\alpha}(\tau_1, \tau_2), \quad (8)$$

$g_k(\tau_1, \tau_2) \equiv -i \langle T_c \hat{c}_k(\tau_1) \hat{c}_k^\dagger(\tau_2) \rangle_0$ is the Green's function of free electron in state k , and $f_{\alpha}(\tau_1, \tau_2) \equiv -i \langle T_c \hat{a}_{\alpha}(\tau_1) \hat{a}_{\alpha}^\dagger(\tau_2) \rangle_0$ is the Green's function of photon in mode α of the radiation field.

Note that while expressions for self-energies due to contacts L and R are exact, because of non-quadratic molecule-radiation field coupling in (3) self-energy due to radiation field can be treated only approximately. We utilize second order diagrammatic expansion (the self-consistent Born approximation) for Σ^{pt} (for more details see, e.g., Ref. 53). Note also that for simplicity we disregard effect of molecule on radiation field modes.

Once Green's function is known, it can be used for evaluation of molecular responses to external perturbations. Below we are interested in simulating photon flux $I_{pt}(t)$ [53] and electron current $I_{L(R)}(t)$ through interface

$L(R)$ [54]

$$\begin{aligned} I_{pt}(t) &= 2 \operatorname{Re} \int_{-\infty}^t dt' \left[F^<(t, t') \Pi^>(t', t) \right. \\ &\quad \left. - F^>(t, t') \Pi^<(t', t) \right] \quad (9) \\ I_{L(R)}(t) &= 2 \operatorname{Re} \int_{-\infty}^t dt' \operatorname{Tr} \left[\Sigma^{L(R)<}(t, t') \mathbf{G}^>(t', t) \right. \\ &\quad \left. - \Sigma^{L(R)>}(t, t') \mathbf{G}^<(t', t) \right] \end{aligned}$$

Here, trace is over molecular subspace and $\Pi^{\lessgtr}$ is the lesser/greater projection of the photon self-energy due to coupling to electrons evaluated within the self-consistent Born approximation

$$\Pi(\tau_1, \tau_2) = -i \operatorname{Tr} \left[\mathbf{U} \mathbf{G}(\tau_1, \tau_2) \mathbf{U}^\dagger \mathbf{G}(\tau_2, \tau_1) \right] \quad (10)$$

Note that in (9) we follow the convention where positive photon flux goes from molecule to radiation field, while positive electron flux goes from contact into the molecule.

C. Floquet formulation for the NEGF-SCBA

While for general time-dependent drivings one has to solve the Dyson equation (6) numerically on a two-dimensional time grid [55], periodic driving allows to formulate an effective time-independent problem in an extended Floquet space [29]. Because of time periodicity of the Hamiltonian, one-sided Fourier transform of the Green's function retarded projection can be expanded in discrete series of Floquet modes

$$\mathbf{G}^r(t_1, t_2) = \sum_{f=-\infty}^{+\infty} \int \frac{dE}{2\pi} \mathbf{G}^r(f; E) e^{-iE(t_1-t_2)+if\omega_0 t_1} \quad (11)$$

Substituting this expansion into the Dyson equations (6) leads to

$$\begin{aligned} & \left[\mathbf{I}(E - f\omega_0) - \mathbf{H}^M \right] \mathbf{G}^r(f; E) \\ & + \frac{1}{2} \vec{\mu} \cdot \vec{E}_0 \left[e^{i\phi_0} \mathbf{G}^r(f-1; E) + e^{-i\phi_0} \mathbf{G}^r(f+1; E) \right] \\ & - \sum_{f_1, f_2=-\infty}^{+\infty} \Sigma^r(f_1, f_2; E - [f + f_1]\omega_0) \\ & \quad \times \mathbf{G}^r(f + f_1 + f_2; E) = \mathbf{I} \delta_{f,0} \quad (12) \\ \mathbf{G}^{\lessgtr}(f_1, f_2; E) &= \sum_{f_3, f_4=-\infty}^{+\infty} \mathbf{G}^r(f_1 - f_3; E - f_3\omega_0) \\ & \quad \times \Sigma^{\lessgtr}(f_3, f_4; E) \mathbf{G}^a(f_2 - f_4; E - f_4\omega_0) \end{aligned}$$

Here $\mathbf{G}^a(f; E) \equiv [\mathbf{G}^r(f; E)]^\dagger$,

$$\mathbf{G}^\lessgtr(t_1, t_2) = \sum_{f_1, f_2=-\infty}^{+\infty} \int \frac{dE}{2\pi} \mathbf{G}^\lessgtr(f_1, f_2; E) \quad (13)$$

$$\times e^{-iE(t_1-t_2)+if_1\omega_0 t_1-if_2\omega_0 t_2},$$

and explicit expressions for the self-energy projections in the Floquet space are given in Appendix A. Equations (12) present effective time-independent formulation of the NEGF-SCBA.

Once Floquet space projections of the Green's function are known, we use them to evaluate dc (period averaged) photon and electron fluxes. Substituting (11) and (13) to (9) yields

$$I_{pt}^{dc} = \sum_{f=-\infty}^{+\infty} \int \frac{d\omega}{2\pi} \left[F^<(\omega) \Pi^>(f, f; \omega + f\omega_0) - F^>(\omega) \Pi^<(f, f; \omega + f\omega_0) \right]$$

$$I_{L(R)}^{dc} = \sum_{f=-\infty}^{+\infty} \int \frac{dE}{2\pi} \quad (14)$$

$$\text{Tr} \left[\Sigma^{L(R)<}(E - f\omega_0) \mathbf{G}^>(f, f; E) - \Sigma^{L(R)>}(E - f\omega_0) \mathbf{G}^<(f, f; E) \right]$$

Here, $\mathbf{G}^<(f, f; E)$ and $\mathbf{G}^>(f, f; E)$ are calculated from (12), expressions for $\Pi^<(f, f; \omega)$ and $\Pi^>(f, f; \omega)$ are given in (A7),

$$F^<(\omega) = -i\gamma(\omega) N_{pt}(\omega)$$

$$F^>(\omega) = -i\gamma(\omega) [N_{pt}(\omega) + 1] \quad (15)$$

$$\Sigma^{L(R)<}(E) = i\Gamma^{L(R)} f_{L(R)}(E)$$

$$\Sigma^{L(R)>}(E) = -i\Gamma^{L(R)} [1 - f_{L(R)}(E)]$$

where

$$\gamma(\omega) \equiv 2\pi \sum_{\alpha} |V_{\alpha}^{pt}|^2 \delta(\omega - \omega_{\alpha})$$

$$\Gamma_{m_1 m_2}^{L(R)}(E) \equiv 2\pi \sum_{k \in L(R)} V_{m_1 k} V_{k m_2} \delta(E - \varepsilon_k) \quad (16)$$

are the radiative energy dissipation rate and electronic decay rate, respectively, $f_{L(R)}(E)$ is the Fermi-Dirac distribution, and $N_{pt}(\omega)$ is population of radiation field modes at frequency ω .

We note that optical absorption theory presented in Ref. 46 is zero bias quasiparticle limit of present formulation with focus on incoming flux and neglect of self-consistency in treatment of light-matter interaction (see Appendix B for details). In the case of quantum treatment of radiation field, the latter assumption is known to violate conservation laws [56, 57] and may lead to qualitative failures [58]. Thus, it is not applicable in optical

studies of open quantum systems [59]. Below we present numerical illustrations of the developed Floquet-NEGF-SCBA formulation.

III. NUMERICAL RESULTS AND DISCUSSION

Simulations start from evaluation of Green's functions employing (12) in the absence of radiation field (i.e. taking self-energy Σ^{pt} to be zero). Resulting expressions are utilized to evaluate self-energy due to coupling to the radiation field Σ^{pt} - last two lines in Eq. (7). Thus obtained self-energy is used in simulation of the updated Green's functions employing (12), etc. This self-consistent procedure continues until convergence. The solution was assumed to reach convergence when difference of level populations for each Floquet mode, $n_m(f) \equiv -i \int dE G_{mm}^<(f, f; E)/2\pi$, at subsequent steps, s and $s+1$, of the iterative procedure is less than 10^{-6} : $|n_m^{(s+1)}(f) - n_m^{(s)}(f)| < 10^{-6}$ for each f .

In our simulations we use a generic three level model with each level independently coupled to the contacts

$$H_{m_1 m_2}^M = \delta_{m_1, m_2} \varepsilon_{m_1} \quad (17)$$

$$\Gamma_{m_1 m_2}^{L(R)} = \delta_{m_1, m_2} \Gamma_{m_1}^{L(R)}$$

For simplicity we also assume the wide band approximation for contacts and radiation bath, i.e. $\Gamma^{L(R)}$ and γ are energy independent and corresponding Lamb shifts, $\Lambda^{L(R)}$ and λ are zero. Parameters and results of the simulations are presented in terms of arbitrary unit of energy U_0 and corresponding unit of flux $I_0 = U_0/\hbar$. Unless stated otherwise, parameters are the following: $k_B T = 0.03$, electron escape rates are $\Gamma_m^L = \Gamma_m^R = 0.01$ and energy dissipation rate is $\gamma = 10^{-4}$. Population of the modes corresponding to the pumping frequency ω_p is $N(\omega_p) = 1$, other modes are assumed to be unpopulated. Phase of the driving field $\phi_0 = 0$. Fermi energy is taken as origin, $E_F = 0$, and bias V_{sd} is applied symmetrically $\mu_L = E_F + |e|V_{sd}/2$ and $\mu_R = E_F - |e|V_{sd}/2$. Floquet space in the simulations was restricted to 21 modes, i.e. $f \in \{-10, \dots, 0, \dots, 10\}$. Simulations are performed on energy grid spanning the range from -2 to 2 with step 10^{-3} .

We start from the model considered in Ref. 46. Here, $\varepsilon_0 = -0.5$, $\varepsilon_1 = 0.5$ and $\varepsilon_2 = 1.1$. Radiation field is assumed to be coupled to optical transition between orbitals 0 and 1, while external driving couples orbitals 1 and 2 and is set to be in resonance with the transition, $\omega_0 = 0.6$ (see Fig. 2a). As discussed in Ref. 46, at zero applied bias voltage ($V_{sd} = 0$) single peak in absorption spectrum splits into two for stronger driving (compare solid and dashed lines in Fig. 2b). The splitting starts at $\vec{\mu} \cdot \vec{E}_0 \sim \sum_{m=1}^2 \Gamma_m$ (here $\Gamma_m \equiv \Gamma_m^L + \Gamma_m^R$) and increases with strength of the driving (see Fig. 2c).

It is easy to understand the effect by going to the rotating frame of the driving field. Because driving is




FIG. 2. Three-level junction responses to external drivings. Shown are (a) sketch of the junction; (b) photon flux vs. pumping frequency ω_p for $\vec{\mu} \cdot \vec{E}_0 = 0$ (solid line, blue) and 0.1 (dashed line, blue) at $V_{sd} = 0$; (c) map of the photon flux vs. ω_p and $\vec{\mu} \cdot \vec{E}_0$ at $V_{sd} = 0$; (d) photon flux for $\vec{\mu} \cdot \vec{E}_0 = 0.1$ at $V_{sd} = 0$ (dashed line, blue) and 1 (dotted line, blue); (e) electron density of states for $\vec{\mu} \cdot \vec{E}_0 = 0$ (solid line, blue) and 0.2 (dashed line, red); (f) electron flux vs. bias V_{sd} for $\vec{\mu} \cdot \vec{E}_0 = 0$ (solid line, blue) and 0.2 (dashed line, red). Red circles in (b) and (d) show results of the Floquet-NEGF-SCBA simulations. Other results are obtained by going to rotational frame of the field. See text for parameters and further details.

at resonance, $\omega_0 = \varepsilon_2 - \varepsilon_1$, utilization of the rotating wave approximation (RWA) allows to map original time-dependent problem $\hat{H}(t)$, Eq. (2), onto an effective Hamiltonian $\hat{\tilde{H}}(t)$

$$\begin{aligned} \hat{\tilde{H}}(t) &= i \left[\frac{d}{dt} e^{\hat{S}(t)} \right] e^{-\hat{S}(t)} + e^{\hat{S}(t)} \hat{H}(t) e^{-\hat{S}(t)} \\ \hat{S}(t) &= -i \frac{\omega_0}{2} t \left(\hat{n}_0 + \hat{n}_1 - \hat{n}_2 \right) \end{aligned} \quad (18)$$

where $\hat{n}_m \equiv \hat{d}_m^\dagger \hat{d}_m$. The transformation leads to

$$\begin{aligned} \hat{\tilde{H}}(t) &= \sum_{m=0}^2 \bar{\varepsilon}_m \hat{d}_m^\dagger \hat{d}_m - \frac{1}{2} \vec{\mu} \cdot \vec{E}_0 \left(\hat{d}_1^\dagger \hat{d}_2 e^{i\phi_0} + \hat{d}_2^\dagger \hat{d}_1 e^{-i\phi_0} \right) \\ &+ \sum_k \varepsilon_k \hat{c}_k^\dagger \hat{c}_k + \sum_\alpha \omega_\alpha \hat{a}_\alpha^\dagger \hat{a}_\alpha \\ &+ \sum_{m=0}^2 \sum_k \left(\bar{V}_{km}(t) \hat{c}_k^\dagger \hat{d}_m + H.c. \right) \\ &+ \sum_\alpha \left([V_\alpha^{pt}]^* \hat{a}_\alpha^\dagger \hat{d}_0^\dagger \hat{d}_1 + H.c. \right) \end{aligned} \quad (19)$$

Here $\bar{\varepsilon}_{0,1} = \varepsilon_{0,1} + \omega_0/2$, $\bar{\varepsilon}_2 = \varepsilon_2 - \omega_0/2$, $\bar{V}_{k0(1)}(t) = V_{k0(1)} e^{-i\omega_0 t/2}$ and $\bar{V}_{k2}(t) = V_{k2} e^{i\omega_0 t/2}$. Note that in (19)

the external driving enters as a time-independent electron hopping between renormalized degenerate levels 1 and 2, $\bar{\varepsilon}_1 = \bar{\varepsilon}_2$. Moreover, because the original levels are well separated, $|\varepsilon_{m_1} - \varepsilon_{m_2}| \gg \Gamma_{m_1}, \Gamma_{m_2}$, bath induced coherences among them can be safely disregarded. In this regime, the transition to the rotating frame of the field allows mapping onto an effective time-independent problem. Indeed, for diagonal electron dissipation matrix Γ time factors in $\bar{V}_{km}(t)$ will result in shift of chemical potentials compensating renormalization of the orbital levels, i.e. $\mu_{L(R)}$ is shifted up by $\omega_0/2$ for levels 0 and 1 and down by $\omega_0/2$ for level 2. This time-independent problem can be treated within the standard NEGF-SCBA [60].

In the rotating frame of the field, the splitting, $\vec{\mu}_0 \cdot \vec{E}_0$, is manifestation of absorption into two eigenlevels resulting from diagonalization of the 1 – 2 block of the molecular Hamiltonian. The two peaks in the spectrum become distinguishable when the splitting is bigger than broadening of the eigenlevels. Note that for the parameters of the simulation, the RWA is quite a reasonable approximation. Effects beyond the RWA, such as, e.g., difference absorption peak heights (dips in Fig. 2b), are small - compare Floquet-NEGF-SCBA treatment of the original problem (circles) with NEGF-SCBA results for effective time independent model (dashed line). At finite




FIG. 3. Photon flux vs. pumping frequency ω_p in a three-level junction model at $V_{sd} = 0$. Shown are (a) sketch of the junction (b) photon flux for several driving strengths $\vec{\mu} \cdot \vec{E}_0$ at $\omega_0 = 0.2$; (c) photon flux for several driving frequencies ω_0 at $\vec{\mu} \cdot \vec{E}_0 = 0.2$. Simulations are performed with the Floquet-NEGF-SCBA approach. See text for parameters and further details.

bias, when the depopulation of level 0 and the population of level 1 by electron transfer between molecule and contacts becomes possible, absorption is substituted by emission, dips become less pronounced and depths differ due to different population of the eigenlevels at $V_{sd} = 1$ (compare dashed and dotted lines in Fig. 2d).

Similarly, splitting can be observed in transport characteristics of the junction. Fig. 2e shows splitting in the density of states A (conductance) induced by $\vec{\mu} \cdot \vec{E}_0 = 0.2$ external driving (compare solid and dashed lines), while Fig. 2f demonstrates consequences of the splitting on current-voltage characteristics of the junction. The difference is due to tunneling via driven levels of the model. The calculations are performed in the rotated frame of the field.

When bath-induced coherences are non-negligible, considerations becomes more involved. In this case one cannot formulate an effective time-independent problem, and Floquet space consideration becomes important. For example, such situation happens when the radiation field is coupled to two optical transitions with external driving in between levels in the transitions. We consider a three-level model with $\epsilon_0 = -0.1$, $\epsilon_1 = 0.1$ and $\epsilon_2 = 1$. Radiation field is assumed to be coupled to optical transitions between orbitals 0 and 2 and between 1 and 2, while external driving couples orbitals 0 and 1 (see Fig. 3a). As previously, driving frequency is taken at resonance with its optical transition, $\omega_0 = \epsilon_1 - \epsilon_0 = 0.2$. Contrary to the model in Fig. 2a, where single peak splits into two on increase of driving strength, absorption in Fig. 3b show transition from split to one peak to split spectrum (dashed to solid to dotted lines in the panel b).

Because radiation field couples orbitals 0 and 1 formulation of effective time-independent problem becomes impossible. Nevertheless, transition to rotating frame similar to the one above yields qualitative understanding of the observed effects of quantum coherence on molecular absorption spectrum. One can show that absorption spectrum of the model in Fig. 3a having single peak (dip in Fig. 3) at $\epsilon_2 - \epsilon_0$ in the absence of driving, when the driving is present can have four peaks at frequencies (see Appendix C for details)

$$\omega = \epsilon_2 - \frac{\epsilon_0 + \epsilon_1 \pm \omega_0}{2} \pm \frac{1}{2} \sqrt{(\epsilon_1 - \epsilon_0 - \omega_0)^2 + (\vec{\mu} \cdot \vec{E}_0)^2} \quad (20)$$

In particular, for the parameters of calculation one expects three peaks for $\vec{\mu} \cdot \vec{E}_0 = \omega_0$, and four peaks otherwise (two additional peaks for each $\vec{\mu} \cdot \vec{E}_0$ are outside of ω_p range in Fig. 3b). Heights of the peaks and their broadenings can only be obtained from the simulations. Another experimentally easily accessible parameter is driving frequency. Also here qualitatively similar behavior is observed (see Fig. 3c).

Finally, we demonstrate coherent control of absorption and transport in a junction beyond RWA considerations. The molecule is presented by its HOMO $\epsilon_0 = -0.5$, LUMO $\epsilon_1 = 0.4$ and LUMO+1 $\epsilon_2 = 0.6$ levels. External




FIG. 4. Coherent control in a three-level junction model at $V_{sd} = 0.5$. Shown are (a) sketch of the junction and (b) photon I_{pt} and (c) electron I_R fluxes, Eq. (14) for $\phi_0 = 0$ (solid line, triangles, blue) and $\phi_0 = \pi/2$ (dashed line, circles, red). Simulations are performed with the Floquet-NEGF-SCBA approach. See text for parameters and further details.

driving couples LUMO and LUMO+1 and is chosen to be in resonance with the optical transition, $\omega_0 = 0.2$. Its coupling strength is taken as $\vec{\mu} \cdot \vec{E}_0 = 0.2$. Radiation field couples ground state (HOMO) with the two excited states (LUMO and LUMO+1). Energy dissipation rate is $\gamma = 0.01$. Coherence between radiation and driving fields results in control of the junction responses. In particular, changing driving field phase ϕ_0 one can affect optical absorption which in turn affects current through the junction (compare solid and dashed lines in Figs. 4b for absorption and in Fig. 4c for current). Note that negative sign of photon flux indicates absorption, which negative sign of electron flux indicates electron transfer from molecule to the contact R . Note also that possibility of coherent control of optical properties was first discussed in Ref. 61.

IV. CONCLUSION

Characterization and control of matter by optical means is at the forefront of research both due to fundamental insights and technological promise. For example, in molecular junctions periodic driving of the system by laser pulse pairs was proposed as a tool to study intramolecular sub-picosecond dynamics in biased junctions. Theoretical modeling of periodically driven systems is a prerequisite to understanding and engineering nanoscale quantum devices for quantum technologies.

We consider open nonequilibrium optoelectronic system under external periodic driving. The system was treated within the nonequilibrium Green's function (NEGF) approach and light-matter interaction was assumed to be small enough to allow treatment at the second order of the diagrammatic expansion - the self-consistent Born approximation (SCBA). Periodic driving is accounted for within the Floquet theory, which maps original time-dependent Hamiltonian onto effective time-independent problem in the extended Floquet space.

Combined Floquet-NEGF-SCBA expressions were formulated and the approach was illustrated with generic three-level junction model simulations demonstrating coherent control of optical response in such devices. The formulation complemented study of optical properties of current carrying junctions proposed in Ref 45 by accounting for periodic external driving and extended recent study on optical absorption properties of matter put forward in Ref. 46 to realm of open nonequilibrium molecular systems. Incorporation of strong intra-system interactions and application of the methodology in ab initio simulations are goals for future research.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation under Grant No. CHE-1565939 (M.G.) and CHE-1553939 (I.F.). G.C. acknowl-

edges the UCSD Physical Sciences Undergraduate Summer Research Award.

Appendix A: Self-energies in the Floquet space

Here we give expressions for the self-energies utilized in (12). The expressions are obtained by substituting (11)

Retarded projection is

$$\Sigma^r(f_1, f_2; E) = \delta_{f_1,0} \delta_{f_2,0} \Sigma^{L(R)r}(E) + \Sigma^{pt r}(f_1, f_2; E) \quad (\text{A1})$$

$$\Sigma^{L(R)r}(E) = \Lambda^{L(R)}(E) - \frac{i}{2} \Gamma^{L(R)}(E) \quad (\text{A2})$$

$$\begin{aligned} \Sigma^{pt r}(f_1, f_2; E) = i \int \frac{d\omega}{2\pi} & \left(\mathbf{U}^\dagger \left[\mathbf{G}^>(f_1, f_2; E - \omega) F^r(\omega) + \delta_{f_2,0} \mathbf{G}^r(f_1; E - \omega) F^<(\omega) \right] \mathbf{U} \right. \\ & \left. + \mathbf{U} \left[\mathbf{G}^<(f_1, f_2; E + \omega) F^a(\omega) + \delta_{f_2,0} \mathbf{G}^r(f_1; E + \omega) F^<(\omega) \right] \mathbf{U}^\dagger \right) \end{aligned} \quad (\text{A3})$$

where $F^<(\omega)$ is defined in (15),

$$F^r(\omega) = \lambda(\omega) - \frac{i}{2} \gamma(\omega), \quad F^a(\omega) = [F^r(\omega)]^*, \quad (\text{A4})$$

$\gamma(\omega)$ and $\Gamma^{L(R)}(E)$ are defined in (16), and $\lambda(\omega)$ and $\Lambda^{L(R)}(E)$ are the real parts of the corresponding retarded projections associated with $\gamma(\omega)$ and $\Gamma^{L(R)}(E)$ via the Kramers-Kronig relations.

Lesser and greater projections are

$$\begin{aligned} \Sigma^{\lessgtr}(f_1, f_2; E) &= \delta_{f_1,0} \delta_{f_2,0} \Sigma^{L(R)\lessgtr}(E) \\ &+ \Sigma^{pt \lessgtr}(f_1, f_2; E) \end{aligned} \quad (\text{A5})$$

where $\Sigma^{L(R)\lessgtr}(E)$ are defined in (15) and

$$\begin{aligned} \Sigma^{pt \lessgtr}(f_1, f_2; E) &= i \int \frac{d\omega}{2\pi} \\ & \left(\mathbf{U}^\dagger \mathbf{G}^{\lessgtr}(f_1, f_2; E - \omega) F^{\lessgtr}(\omega) \mathbf{U} \right. \\ & \left. + \mathbf{U} \mathbf{G}^{\lessgtr}(f_1, f_2; E + \omega) F^{\lessgtr}(\omega) \mathbf{U}^\dagger \right) \end{aligned} \quad (\text{A6})$$

Finally, substituting (13) into lesser and greater projections of (10) leads to

$$\begin{aligned} \Pi^{\lessgtr}(f_1, f_2; \omega) &= -i \sum_{\substack{f_3, f_4 \\ f_5, f_6}}^{+\infty} \delta_{f_1, f_3 - f_5} \delta_{f_2, f_4 - f_6} \\ & \times \int \frac{dE}{2\pi} \text{Tr} \left[\mathbf{U} \mathbf{G}^{\lessgtr}(f_3, f_4; E + \frac{\omega}{2}) \mathbf{U}^\dagger \mathbf{G}^{\lessgtr}(f_6, f_5; E - \frac{\omega}{2}) \right] \end{aligned} \quad (\text{A7})$$

Appendix B: Comparison to the theory of Ref. 46

Here we compare our consideration to the theory of optical absorption presented in Ref. 46.

and (13) into corresponding projections of (7).

We start from the expression for absorption rate given in Eq. (14) of the latter reference. In terms of our definitions the expression is

$$\begin{aligned} I(\omega_p) &= \lim_{t \rightarrow \infty} \frac{|V_p^{pt}|^2}{2\hbar^2} \frac{1}{t - t_0} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 \\ & \text{Re} \left[\left(e^{-i\omega_p(t_1 - t_2)} + e^{-i\omega_p(t_1 + t_2)} \right) \langle \hat{\mu}(t_2) \hat{\mu}(t_1) \rangle \right] \end{aligned} \quad (\text{B1})$$

where

$$\hat{\mu} = \sum_{m_1, m_2 \in M} U_{m_1 m_2} \hat{d}_{m_1}^\dagger \hat{d}_{m_2}, \quad (\text{B2})$$

V_p^{pt} and $U_{m_1 m_2}$ are defined in Eq. (4) and time-dependence in correlation function of (B1) is with respect to Hamiltonian excluding coupling to radiation field (interaction picture). Note that the latter results in non-conserving expression and is applicable only in cases when coupling to radiation field is small (relative to other energy scales of the problem) so that mistake in the treatment is numerically negligible. Note also that factor of 2 in the denominator in (B1) comes from assumed cosine time dependence of the classical consideration of the radiation field in Ref. 46 which is effectively included in V_p^{pt} in our quantum consideration. Thus, it will be dropped below. Similarly, we drop $e^{-i\omega_p(t_1 + t_2)}$ because, as was discussed in Ref. 46, the term does not contribute to the absorption spectrum. Finally, we take $\hbar = 1$ in our consideration.

Implementing the adjustments and using the fact that integrand is symmetric function of t_1 and t_2 , Eq. (B1)

can be rewritten as

$$I(\omega_p) = \lim_{t \rightarrow \infty} |V_p^{pt}|^2 \frac{1}{t - t_0} \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \quad (B3)$$

$$2 \operatorname{Re} \left[e^{-i\omega_p(t_1 - t_2)} \langle \hat{\mu}(t_2) \hat{\mu}(t_1) \rangle \right]$$

Below we show that this expressions yields time-averaging of in-scattering (absorption) photon flux, first term in Eq. (9), i.e. we show that

$$I(\omega_p) = - \lim_{t \rightarrow \infty} \frac{1}{t - t_0} \int_{t_0}^t dt_1 I_{pt}^{abs}(t_1) \quad (B4)$$

where

$$I_{pt}^{abs}(t_1) = 2 \operatorname{Re} \int_{t_0}^{t_1} dt_2 F_p^<(t_1, t_2) \Pi^>(t_2, t_1) \quad (B5)$$

Note that in our consideration t_0 was taken at $-\infty$ and convention of negative sign for in-coming photon flux was followed. Note also that for dc part of the flux (the only part considered here and in Ref. 46) averaging (B4) is redundant.

Taking into account that $F^<$ in (9) is defined as sum over modes of radiation field, see Eq. (8), to get (B5) one has to consider a single mode of the radiation field with frequency ω_p . From definition (15) and assuming singly populated mode $N_{pt}(\omega_p) = 1$, Green's function $F_p^<$ for the mode can be written in time domain as

$$F_p^<(t_1, t_2) = -i |V_p^{pt}|^2 e^{-i\omega_p(t_1 - t_2)} \quad (B6)$$

Thus, comparing (B3) to (B5) and (B6) one concludes that

$$\Pi^>(t_2, t_1) = -i \langle \hat{\mu}(t_2) \hat{\mu}^\dagger(t_1) \rangle \quad (B7)$$

Right-hand side of this expression is definition of the two-particle Green's function. When coupling to radiation field is neglected (as is done in Ref. 46) and taking into account that the rest of the Hamiltonian (3) does not contain many-body interactions, one can employ Wick's theorem expressing two-particle Green's function in terms of

product of single-particle GFs. In particular, employing Wick's theorem in the right side of (B7) results in an expression similar to greater projection of (10). The crucial difference is that the resulting expression contains zero-order single-particle GFs, while Eq. (10) employs full (dressed in coupling to radiation field) GFs. This completes derivation of theory presented in Ref. 46 from the Floquet-NEGF-SCBA consideration.

Appendix C: Derivation of Eq.(20)

Here we present qualitative analysis leading to Eq. (20).

Model shown in Fig. 3a has Hamiltonian (2)-(3) with $H_{m_1 m_2}^M = \delta_{m_1, m_2} \varepsilon_{m_1}$ ($m \in \{0, 1, 2\}$), $\vec{\mu}_{m_1 m_2} = \vec{\mu}(\delta_{m_1, 0} \delta_{m_2, 1} + \delta_{m_1, 1} \delta_{m_2, 0})$, $\Gamma_{m_1 m_2}^{L(R)} = \delta_{m_1, m_2} \Gamma_{m_1}^{L(R)}$ and $U_{m_1 m_2} = (\delta_{m_1, 0} + \delta_{m_1, 1}) \delta_{m_2, 2}$.

Performing transformation to the field rotational frame, Eq. (18), with

$$\hat{S}(t) = -i \frac{\omega_0}{2} t \left(\hat{n}_0 - \hat{n}_1 \right) \quad (C1)$$

leads to Hamiltonian similar to (19) with $\bar{\varepsilon}_0 = \varepsilon_0 + \omega_0/2$, $\bar{\varepsilon}_1 = \varepsilon_1 - \omega_0/2$ and $\bar{\varepsilon}_2 = \varepsilon_2$; with $\bar{V}_{k0}(t) = V_{k0} e^{i\omega_0 t/2}$, $\bar{V}_{k1}(t) = V_{k1} e^{-i\omega_0 t/2}$ and $\bar{V}_{k2}(t) = V_{k2}$; with $\vec{\mu} \cdot \vec{E}_0$ inducing electron transfer between orbitals 0 and 1

$$-\frac{1}{2} \vec{\mu} \cdot \vec{E}_0 \left(\hat{d}_1^\dagger \hat{d}_0 e^{-i\phi_0} + H.c. \right) \quad (C2)$$

and with coupling to radiation field - last line in Eq. (19) - becoming time-dependent

$$\sum_{\alpha} \left(V_{\alpha}^{pt} \hat{d}_2^\dagger \left[\hat{d}_0 e^{i\omega_0 t/2} + \hat{d}_1 e^{-i\omega_0 t/2} \right] \hat{a}_{\alpha} + H.c. \right) \quad (C3)$$

Diagonalizing the 0 - 1 block of the molecular Hamiltonian yields egen-orbitals ε_+ and ε_- and leads to expressions for system-baths (contacts and radiation field) couplings in the form

$$\sum_k \left(\left[i V_{k1} \sin \theta e^{i(\omega_0 t + \phi_0)/2} - i V_{k0} \cos \theta e^{-i(\omega_0 t + \phi_0)/2} \right] \hat{c}_k^\dagger \hat{d}_+ \right. \quad (C4)$$

$$\left. + \left[i V_{k1} \cos \theta e^{i(\omega_0 t + \phi_0)/2} + i V_{k0} \sin \theta e^{-i(\omega_0 t + \phi_0)/2} \right] \hat{c}_k^\dagger \hat{d}_- + V_{k2} \hat{c}_k^\dagger \hat{d}_2 + H.c. \right)$$

$$\sum_{\alpha} \left(i V_{\alpha}^{pt} \hat{d}_2^\dagger \left[(\sin \theta e^{i(\omega_0 t + \phi_0)/2} - \cos \theta e^{-i(\omega_0 t + \phi_0)/2}) \hat{d}_+ + (\cos \theta e^{i(\omega_0 t + \phi_0)/2} + \sin \theta e^{-i(\omega_0 t + \phi_0)/2}) \hat{d}_- \right] + H.c. \right) \quad (C5)$$

Analyzing expression (C4) within RWA-type consid-

eration used in Ref. 62 and neglecting off-diagonal ele-

ments in the electronic dissipation rate matrix Γ leads to diagonal structure of the Green's function $G_{m_1 m_2}^{(0)} = \delta_{m_1 m_2} G_{m_1 m_1}^{(0)}$ at zero order in the molecule-radiation field coupling.

Thus, dc part of the absorption spectrum (i.e., dc part of the incoming photon flux) takes the form

$$2 \operatorname{Re} \int_{-\infty}^t dt' F^{<}(t, t') G_{22}^{(0)>}(t', t) \quad (\text{C6})$$

$$\times \left[G_{++}^{(0)<}(t, t') \left(\sin^2 \theta e^{i\omega_0(t-t')/2} + \cos^2 \theta e^{-i\omega_0(t-t')/2} \right) \right. \\ \left. + G_{--}^{(0)<}(t, t') \left(\cos^2 \theta e^{i\omega_0(t-t')/2} + \sin^2 \theta e^{-i\omega_0(t-t')/2} \right) \right]$$

Taking integral in t' leads to Eq. (20)

-
- [1] S. W. Wu, G. V. Nazin, and W. Ho, *Phys. Rev. B* **77**, 205430 (2008).
 - [2] X. H. Qiu, G. V. Nazin, and W. Ho, *Science* **299**, 542 (2003).
 - [3] Z.-C. Dong, X.-L. Guo, A. S. Trifonov, P. S. Dorozhkin, K. Miki, K. Kimura, S. Yokoyama, and S. Mashiko, *Phys. Rev. Lett.* **92**, 086801 (2004).
 - [4] K. Kimura, K. Miwa, H. Imada, M. Imai-Imada, S. Kawahara, J. Takeya, M. Kawai, M. Galperin, and Y. Kim, *Nature* **570**, 210 (2019).
 - [5] C. Chen, P. Chu, C. A. Bobisch, D. L. Mills, and W. Ho, *Phys. Rev. Lett.* **105**, 217402 (2010).
 - [6] Y. Zhang, Y. Luo, Y. Zhang, Y.-J. Yu, Y.-M. Kuang, L. Zhang, Q.-S. Meng, Y. Luo, J.-L. Yang, Z.-C. Dong, and J. G. Hou, *Nature* **531**, 623 (2016).
 - [7] H. Imada, K. Miwa, M. Imai-Imada, S. Kawahara, K. Kimura, and Y. Kim, *Nature* **538**, 364 (2016).
 - [8] N. L. Schneider, J. T. Lü, M. Brandbyge, and R. Berndt, *Phys. Rev. Lett.* **109**, 186601 (2012).
 - [9] Z. Ioffe, T. Shamai, A. Ophir, G. Noy, I. Yutsis, K. Kfir, O. Cheshnovsky, and Y. Selzer, *Nature Nanotech.* **3**, 727 (2008).
 - [10] D. R. Ward, N. J. Halas, J. W. Ciszek, J. M. Tour, Y. Wu, P. Nordlander, and D. Natelson, *Nano Lett.* **8**, 919 (2008).
 - [11] D. R. Ward, D. A. Corley, J. M. Tour, and D. Natelson, *Nature Nanotech.* **6**, 33 (2011).
 - [12] D. Natelson, Y. Li, and J. B. Herzog, *Phys. Chem. Chem. Phys.* **15**, 5262 (2013).
 - [13] R. Chikkaraddy, B. de Nijs, F. Benz, S. J. Barrow, O. A. Scherman, E. Rosta, A. Demetriadou, P. Fox, O. Hess, and J. J. Baumberg, *Nature* **535**, 127 (2016).
 - [14] N. Kongsuwan, A. Demetriadou, R. Chikkaraddy, F. Benz, V. A. Turek, U. F. Keyser, J. J. Baumberg, and O. Hess, *ACS Photonics* **5**, 186 (2018).
 - [15] M. Galperin and A. Nitzan, *Phys. Chem. Chem. Phys.* **14**, 9421 (2012).
 - [16] M. Galperin, *Chem. Soc. Rev.* **46**, 4000 (2017).
 - [17] G. A. Steele, A. K. Hüttel, B. Witkamp, M. Poot, H. B. Meerwaldt, L. P. Kouwenhoven, and H. S. J. v. d. Zant, *Science* **325**, 1103 (2009).
 - [18] T. Wagner, P. Talkner, J. C. Bayer, E. P. Rugeramigabo, P. Hänggi, and R. J. Haug, *Nature Phys.* **15**, 330 (2019).
 - [19] Y. Selzer and U. Peskin, *J. Phys. Chem. C* **117**, 22369 (2013).
 - [20] M. A. Ochoa, Y. Selzer, U. Peskin, and M. Galperin, *J. Phys. Chem. Lett.* **6**, 470 (2015).
 - [21] F. Zhan, N. Li, S. Kohler, and P. Hnggi, *Phys. Rev. E* **80**, 061115 (2009).
 - [22] Donarini, A., Yar, A., and Grifoni, M., *Eur. Phys. J. B* **85**, 316 (2012).
 - [23] U. Peskin, *Fortschritte der Physik* **65**, 1600048 (2017).
 - [24] H. Al Hussein, S. Abdalah, K. Al Naimee, R. Meucci, and F. Arecchi, *Nanomaterials and Nanotechnology* **8**, 184798041878238 (2018).
 - [25] V. Leyton, S. Weiss, and M. Thorwart, *Phys. Rev. B* **97**, 214303 (2018).
 - [26] T.-H. Park and M. Galperin, *Europhys. Lett.* **95**, 27001 (2011).
 - [27] T.-H. Park and M. Galperin, *Phys. Rev. B* **84**, 075447 (2011).
 - [28] M. I. Sena-Junior, L. R. F. Lima, and C. H. Lewenkopf, *J. Phys. A* **50**, 435202 (2017).
 - [29] S. Kohler, J. Lehmann, and P. Hänggi, *Phys. Rep.* **406**, 379 (2005).
 - [30] D. Thuberg, E. Muñoz, S. Eggert, and S. A. Reyes, *Phys. Rev. Lett.* **119**, 267701 (2017).
 - [31] B. H. Wu and J. C. Cao, *Phys. Rev. B* **73**, 245412 (2006).
 - [32] T.-S. Ho, K. Wang, and S.-I. Chu, *Phys. Rev. A* **33**, 1798 (1986).
 - [33] J. Lehmann, S. Kohler, P. Hnggi, and A. Nitzan, *Phys. Rev. Lett.* **88**, 228305 (2002).
 - [34] J. Lehmann, S. Kohler, P. Hänggi, and A. Nitzan, *J. Chem. Phys.* **118**, 3283 (2003).
 - [35] B. H. Wu and C. Timm, *Phys. Rev. B* **81**, 075309 (2010).
 - [36] S. Kohler, J. o. Lehmann, and P. H. anggi, *Superlattices and Microstructures* **34**, 419 (2003).
 - [37] S. Kohler, S. Camalet, M. Strass, J. Lehmann, G.-L. Ingold, and P. Hänggi, *Chem. Phys.* **296**, 243 (2004).
 - [38] G. Stefanucci, S. Kurth, A. Rubio, and E. K. U. Gross, *Phys. Rev. B* **77**, 075339 (2008).
 - [39] B. H. Wu and J. C. Cao, *J. Phys.: Condens. Matter* **20**, 085224 (2008).
 - [40] D. Rai and M. Galperin, *J. Phys. Chem. C* **117**, 13730 (2013).
 - [41] B. H. Wu and J. C. Cao, *Phys. Rev. B* **81**, 085327 (2010).
 - [42] N. Tsuji, T. Oka, and H. Aoki, *Phys. Rev. B* **78**, 235124 (2008).

- [43] A. K. Eissling, V. Meden, and D. M. Kennes, Phys. Rev. B **94**, 245116 (2016).
- [44] I. Kinoshita, A. Misu, and T. Munakata, J. Chem. Phys. **102**, 2970 (1995).
- [45] M. Galperin and A. Nitzan, J. Chem. Phys. **124**, 234709 (2006).
- [46] B. Gu and I. Franco, Phys. Rev. A **98**, 063412 (2018).
- [47] N. Sergueev, D. Roubtsov, and H. Guo, Phys. Rev. Lett. **95**, 146803 (2005).
- [48] T. Frederiksen, M. Paulsson, M. Brandbyge, and A.-P. Jauho, Phys. Rev. B **75**, 205413 (2007).
- [49] R. Avriller and T. Frederiksen, Phys. Rev. B **86**, 155411 (2012).
- [50] A. Baratz, M. Galperin, and R. Baer, J. Phys. Chem. C **117**, 10257 (2013).
- [51] H. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors*, second, substantially revised edition ed., Vol. 123 (Springer, Berlin Heidelberg, 2008).
- [52] G. Stefanucci and R. van Leeuwen, *Nonequilibrium Many-Body Theory of Quantum Systems. A Modern Introduction*. (Cambridge University Press, 2013).
- [53] Y. Gao and M. Galperin, J. Chem. Phys. **144**, 174113 (2016).
- [54] A.-P. Jauho, N. S. Wingreen, and Y. Meir, Phys. Rev. B **50**, 5528 (1994).
- [55] P. Myöhänen, A. Stan, G. Stefanucci, and R. van Leeuwen, Phys. Rev. B **80**, 115107 (2009).
- [56] G. Baym and L. P. Kadanoff, Phys. Rev. **124**, 287 (1961).
- [57] G. Baym, Phys. Rev. **127**, 1391 (1962).
- [58] A. Nitzan and M. Galperin, J. Phys. Chem. Lett. **9**, 4886 (2018).
- [59] S. Mukamel and M. Galperin, J. Phys. Chem. C **123**, 29015 (2019).
- [60] M. Galperin, M. A. Ratner, and A. Nitzan, J. Chem. Phys. **121**, 11965 (2004).
- [61] M. O. Scully, Phys. Rev. Lett. **104**, 207701 (2010).
- [62] M. O. Scully, S.-Y. Zhu, and A. Gavrielides, Phys. Rev. Lett. **62**, 2813 (1989).