

Fractional matching preclusion of fault Hamiltonian graphs

Huiqing Liu* Shunzhe Zhang[†] Xinyuan Zhang[‡]

Abstract

Matching preclusion is a measure of robustness in the event of edge failure in inter-connection networks. As a generalization of matching preclusion, the fractional matching preclusion number (FMP number for short) of a graph is the minimum number of edges whose deletion results in a graph that has no fractional perfect matchings, and the fractional strong matching preclusion number (FSMP number for short) of a graph is the minimum number of edges and/or vertices whose deletion leaves a resulting graph with no fractional perfect matchings. A graph G is said to be f -fault Hamiltonian if there exists a Hamiltonian cycle in $G - F$ for any set F of vertices and/or edges with $|F| \leq f$. In this paper, we establish the FMP number and FSMP number of $(\delta - 2)$ -fault Hamiltonian graphs with minimum degree $\delta \geq 3$. As applications, the FMP number and FSMP number of some well-known networks are determined.

Keywords: fractional perfect matching; fractional matching preclusion number; fractional strong matching preclusion number; f -fault Hamiltonian graph

1 Introduction

Let $G = (V(G), E(G))$ be a simple, undirected and finite graph. We denote $VE(G) = V(G) \cup E(G)$ and simply write $|V(G)|$ by $|G|$. For $v \in V(G)$, the set of all edges incident with v is denoted by $E_G(v)$ and the *minimum degree* of G , denoted by $\delta(G)$, is the minimum size of $|E_G(v)|$. For $S \subseteq V(G)$, $G[S]$ denotes the subgraph induced by S and $G - S$ denotes the subgraph induced by $V(G) \setminus S$. For $F \subseteq E(G)$, $G - F$ denotes the resulting graph by deleting all edges of F from G . For $f \in VE(G)$, we simplify $G - \{f\}$ to $G - f$. Denote

*Hubei Key Laboratory of Applied Mathematics, Faculty of Mathematics and Statistics, Hubei University, Wuhan 430062, PR China. E-mail: hqliu@hubu.edu.cn. This author's work was partially supported by NNSFC (Nos. 11571096, 61373019).

[†]Hubei Key Laboratory of Applied Mathematics, Faculty of Mathematics and Statistics, Hubei University, Wuhan 430062, PR China. E-mail: shunzhezhang@hubu.edu.cn.

[‡]Artificial intelligence School, Wuchang University of Technology, Wuhan 430223, PR China. E-mail: xinyuan0715@163.com.

$F_V = F \cap V(G)$ and $F_E = F \cap E(G)$ for $F \subseteq VE(G)$. Two graphs are *vertex disjoint* if they have no vertex in common. A k -*cycle* is a cycle with k vertices. A cycle is called a *Hamiltonian cycle* if it contains all vertices of the graph. A graph is said to be *Hamiltonian* if it contains a Hamiltonian cycle. A graph G is said to be an f -*fault Hamiltonian graph* if there exists a Hamiltonian cycle in $G - F$ for any set F of vertices and/or edges with $|F| \leq f$. A graph is said to be *even* if it has even number of vertices, otherwise, it is said to be *odd*.

A *matching* in a graph is a set of edges no two of which are adjacent. With any matching M of a graph G , we may associate a $\{0, 1\}$ -valued function f that assigns to each edge of G a number in $\{0, 1\}$ such that $\sum_{e \in E_G(v)} f(e) \leq 1$ for each vertex $v \in V(G)$. A matching is *perfect* if $\sum_{e \in E_G(v)} f(e) = 1$ for each vertex v . A matching is *almost-perfect* if there exists exactly one vertex u such that $\sum_{e \in E_G(u)} f(e) = 0$ and $\sum_{e \in E_G(v)} f(e) = 1$ for each vertex $v \in V(G - u)$.

A *matching preclusion set* (MP set for short) is an edge subset F of G if $G - F$ has neither perfect matchings nor almost-perfect matchings. The *MP number* of G , denoted by $mp(G)$, is the minimum size of MP sets of G . In 2005, Brigham *et al.* [3] first introduced the matching preclusion problem which offers a way of measuring the robustness of a given graph as a network topology with respect to link failures. That is, in the situation in which each node of a communication network is demanded to have a special partner at any time, one that has a larger matching preclusion number may be considered as more robust in the event of possible link failures. Since then, the matching preclusion problem of various networks was studied, see [5, 17, 21, 22, 34]. For any $v \in V(G)$, the set of edges incident with v forms a MP set of G if $|G|$ is even. Thus $mp(G) \leq \delta(G)$.

Another type of failure in a communication network occurs through nodes, which is in fact more offensive, is through node failures. As a more general matching preclusion problem, the strong matching preclusion deals with the corresponding matching problem that has also been analyzed under vertex deletions, see [1, 13]. Park and Ihm [30] considered the following extended form of matching preclusion. A *strong matching preclusion set* (SMP set for short) is a set F of edges and/or vertices of G if $G - F$ has neither perfect matchings nor almost-perfect matchings. The *SMP number* of G , denoted by $smp(G)$, is the minimum size of SMP sets of G . According to the definition of $mp(G)$ and $smp(G)$, we have $smp(G) \leq mp(G)$.

By utilizing the definition of matching with the continuous unit interval $[0, 1]$ instead of the “discrete unit interval” $\{0, 1\}$, we get the following generalization of matching introduced in [31].

A *fractional matching* is a function f that assigns to each edge a number in $[0, 1]$ such

that $\sum_{e \in E_G(v)} f(e) \leq 1$ for each vertex $v \in V(G)$. A *fractional perfect matching* is a fractional matching f so that $\sum_{e \in E_G(v)} f(e) = 1$ for every $v \in V(G)$. Note that a perfect matching is also a fractional perfect matching.

Let G be a graph and $S \subseteq V(G)$, we use $i(G - S)$ and $c(G - S)$ to denote the number of isolated vertices and the number of components of $G - S$, respectively. It is obvious that $i(G - S) \leq c(G - S)$. The following proposition is a necessary and sufficient condition for a graph to have a fractional perfect matching.

Proposition 1.1 [31] *A graph G has a fractional perfect matching if and only if $i(G - S) \leq |S|$ for every set $S \subseteq V(G)$.*

As a generalization of matching preclusion, the concept of the fractional matching preclusion number was introduced by Liu *et al.* [23]. A *fractional matching preclusion set* (FMP set for short) is an edge subset F of G if $G - F$ has no fractional perfect matchings. The *FMP number* of G , denoted by $fmp(G)$, is the minimum size of FMP sets of G . Obviously, $fmp(G) \leq \delta(G)$. By the definition of $fmp(G)$, $mp(G) \leq fmp(G)$ if $|G|$ is even. So we have the following proposition.

Proposition 1.2 [23] *Let G be a graph of even order. If $mp(G) = \delta(G)$, then $mp(G) = fmp(G) = \delta(G)$.*

A *fractional strong matching preclusion set* (FSMP set for short) is a set F of edges and/or vertices of G if $G - F$ has no fractional perfect matchings. The *FSMP number* of G , denoted by $fsmp(G)$, is the minimum size of FSMP sets of G . By the definition of $fmp(G)$ and $fsmp(G)$, we have the following proposition.

Proposition 1.3 [23] *Let G be a graph. Then $fsmp(G) \leq fmp(G) \leq \delta(G)$.*

In 2017, Liu and Liu [23] considered the FMP number and FSMP number of complete graphs, Petersen graph and twisted cubes. Later, Ma *et al.* [24] obtained the FMP number and FSMP number of (burnt) pancake graphs. Ma *et al.* [25] determined the FMP number and FSMP number of arrangement graphs. Recently, Zhang *et al.* [36] established the FMP number and FSMP number of the n -dimensional restricted HL-graphs, respectively.

In this paper, we establish the FMP number and FSMP number of $(\delta - 2)$ -fault Hamiltonian graphs with minimum degree $\delta \geq 3$. As applications, the FMP number and FSMP number of some well-known networks are determined.

The rest of the paper is organized as follows: Section 2 provides some useful lemmas. In Section 3, we investigate the FMP number and FSMP number of $(\delta - 2)$ -fault Hamiltonian graphs with minimum degree $\delta \geq 3$. In Section 4, we determine the FMP number and FSMP number of some well-known networks. Our conclusions are given in Section 5.

2 Lemmas

Mao *et al.* [26] gave a sufficient condition to determine the MP number and SMP number of fault Hamiltonian graphs.

Lemma 2.1 [26] *Let G be a $(\delta - 2)$ -fault Hamiltonian graph with minimum degree δ . Then $\text{smp}(G) = \text{mp}(G) = \delta$.*

The following lemma shows a necessary condition for the existence of a Hamiltonian cycle.

Lemma 2.2 [2] *Let S be a set of vertices of a Hamiltonian graph G . Then*

$$c(G - S) \leq |S|. \quad (1)$$

Moreover, if equality holds in (1), then each of the $|S|$ components of $G - S$ has a Hamiltonian path.

The following is a necessary and sufficient condition for a graph to have a fractional perfect matching.

Lemma 2.3 [31] *A graph G has a fractional perfect matching if and only if there is a partition $\{V_1, V_2, \dots, V_s\}$ of the vertex set $V(G)$ such that, for each i , the graph $G[V_i]$ is either K_2 or Hamiltonian.*

An *independent set* in a graph is a set of vertices no two of which are adjacent. The cardinality of a maximum independent set in a graph G is called the *independent number* of G and is denoted by $\alpha(G)$. A *covering* of a graph G is a set of vertices which together meet all edges of G . The minimum number of vertices in a covering of a graph G is called the *covering number* of G and is denoted by $\beta(G)$. Gallai [11] showed the relationship between the independent number $\alpha(G)$ and the covering number $\beta(G)$ of a graph G .

Lemma 2.4 [11] $\alpha(G) + \beta(G) = |G|$ for a graph G . Furthermore, $\alpha(G) \geq |G| - |E(G)|$, where the equality holds if and only if $E(G)$ is a matching of G .

Lemma 2.5 *Let G be a graph with $F \subseteq V E(G)$. If $G - F$ is an independent set, then*

$$\alpha(G) \geq |G - F| - |F_E|, \quad (2)$$

where the equality holds if and only if $E(G - F_V) = F_E$ and F_E is a matching of $G - F_V$.

Proof. Note that $G - F$ is an independent set, then $E(G - F_V) \subseteq F_E$. By Lemma 2.4, $\alpha(G) \geq \alpha(G - F_V) \geq |G - F_V| - |E(G - F_V)| \geq |G - F| - |F_E|$, where the equality holds if and only if $E(G - F_V) = F_E$ and F_E is a matching of $G - F_V$. ■

Lemma 2.6 *Let G be a $(\delta - 2)$ -fault Hamiltonian graph with minimum degree δ and F a subset of $VE(G)$ with $|F| = \delta - 1$. If there exists $S \subseteq V(G - F)$ such that $i(G - (F \cup S)) \geq |S| + 1$, then $G - (F \cup S)$ is an independent set with $\frac{|G| + |F_E| - \delta}{2} + 1$ vertices and $\alpha(G) \geq |G - (F \cup S)| - |F_E|$.*

Proof. Let $F' = F - \beta$ for some $\beta \in F$, and thus $|F'| = \delta - 2$. Since G is $(\delta - 2)$ -fault Hamiltonian, $G - F'$ is Hamiltonian. By Lemma 2.2, for any $S' \subseteq V(G - F')$,

$$c(G - (F' \cup S')) \leq |S'|. \quad (3)$$

Note that $S \subseteq V(G - F)$ and $G - (F \cup S) = G - (F' \cup (S \cup \{\beta\}))$, where $\beta \in F$. If $\beta \in F_V$, then $S \cup \{\beta\} \subseteq V(G - F')$. This together with (3), we have

$$c(G - (F \cup S)) = c(G - (F' \cup (S \cup \{\beta\}))) \leq |S \cup \{\beta\}| = |S| + 1.$$

If $\beta \in F_E$, then $S \subseteq V(G - F')$, and thus, by (3),

$$c(G - (F \cup S)) = c(G - (F' \cup (S \cup \{\beta\}))) \leq c(G - (F' \cup S)) + 1 \leq |S| + 1.$$

Then $c(G - (F \cup S)) \leq |S| + 1$. Note that $i(G - (F \cup S)) \geq |S| + 1$. Hence,

$$|S| + 1 \leq i(G - (F \cup S)) \leq c(G - (F \cup S)) \leq |S| + 1,$$

which means $i(G - (F \cup S)) = c(G - (F \cup S)) = |S| + 1$, and thus $G - (F \cup S)$ is an independent set and $|G - (F \cup S)| = |S| + 1$. Note that $|G - (F \cup S)| = |G| - |F_V| - |S|$ and $|F_V| + |F_E| = |F| = \delta - 1$. Then $|G - (F \cup S)| = \frac{|G| + |F_E| - \delta}{2} + 1$. Therefore, by Lemma 2.5, $\alpha(G) \geq |G - (F \cup S)| - |F_E|$. ■

3 The fractional (strong) matching preclusion of G

The following theorem investigates the FSMP number of regular bipartite graphs.

Theorem 3.1 *Let G be a regular bipartite graph. Then $f_{smp}(G) = 1$.*

Proof. Let $G = G[X, Y]$ be a regular bipartite graph. Then $|X| = |Y|$ and G has a perfect matching (see [2]). By Lemma 2.3, G has a fractional perfect matching. Thus $fsm(p)(G) \geq 1$. Let $x \in X$ and $X' = X - x$. Then $i((G - x) - X') = |Y| > |X| - 1 = |X'|$, and thus, by Proposition 1.1, $G - x$ has no fractional perfect matchings. Then $fsm(p)(G) \leq 1$. Hence $fsm(p)(G) = 1$. ■

In the following, we always assume that G is $(\delta - 2)$ -fault Hamiltonian with minimum degree $\delta \geq 3$. By Lemma 2.1 and Proposition 1.2, we have the following result directly.

Lemma 3.2 *If $|G|$ is even, then $fmp(G) = \delta$.*

The following lemma shows the upper and lower bound of the FMP number and FSMP number of G .

Lemma 3.3 $\delta - 1 \leq fsm(p)(G) \leq fmp(G) \leq \delta$.

Proof. By Proposition 1.3, $fsm(p)(G) \leq fmp(G) \leq \delta$. Let $F \subseteq VE(G)$ with $|F| \leq \delta - 2$. Since G is $(\delta - 2)$ -fault Hamiltonian, $G - F$ has a Hamiltonian cycle, and thus $G - F$ has a fractional perfect matching by Lemma 2.3. It follows that $fsm(p)(G) \geq \delta - 1$. ■

Lemma 3.4 *Let F be a subset of $VE(G)$ with $|F| = \delta - 1$. If $G - F$ has no fractional perfect matchings, then $|G| - |F_V|$ is odd. Furthermore, $|F_E| \leq \delta - 2$ when $|G|$ is even.*

Proof. Suppose that $|G| - |F_V|$ is even. Then $G - F$ is even order. By Lemma 2.1, $smp(G) = \delta > |F|$, and hence, $G - F$ has a perfect matching. Then by Lemma 2.3, $G - F$ has a fractional perfect matching, a contradiction. Furthermore, if $|G|$ is even, then $|F_V| \neq 0$. Note that $|F_V| + |F_E| = |F| = \delta - 1$, then $|F_E| \leq \delta - 2$. ■

Next, we first give the definitions of two different graph classes, then show some sufficient conditions to determine the FMP number and FSMP number of fault Hamiltonian graphs.

A graph G is called \mathcal{H} -free if G does not contain H as an induced subgraph for any $H \in \mathcal{H}$, and we call each H a *forbidden subgraph*. Let $\mathcal{G}_1(k) = \{G \mid G \text{ is a } k\text{-regular odd graph or } \{K_4 - e\}\text{-free even graph in which every edge lies in at least a 3-cycle and a 4-cycle}\}$ and $\mathcal{G}_2(4) = \{G \mid G \text{ is a 4-regular } \{K_4, K_4 - e, K_{2,3}\}\text{-free odd graph in which every edge lies in at least a 3-cycle and two 4-cycles}\}$.

Theorem 3.5 *Let G be a $(\delta - 2)$ -fault Hamiltonian graph with minimum degree $\delta \geq 3$.*

- (i) *If $\alpha(G) \leq \lceil \frac{|G|+1}{2} \rceil - \delta$, then $fsm(p)(G) = fmp(G) = \delta$;*
- (ii) *If $G \in \mathcal{G}_1(\delta)$ and $\alpha(G) \leq \lceil \frac{|G|+1}{2} \rceil - \delta + 1$, then $fsm(p)(G) = fmp(G) = \delta$;*
- (iii) *If $G \in \mathcal{G}_2(4)$ and $\alpha(G) \leq \frac{|G|+1}{2} - 2$, then $fsm(p)(G) = fmp(G) = 4$.*

Proof. By Lemma 3.3, $\delta - 1 \leq f_{\text{smp}}(G) \leq f_{\text{mp}}(G) \leq \delta$. In the following, we will show that, for any $F \subseteq VE(G)$ with $|F| = \delta - 1$, $G - F$ has a fractional perfect matching. Suppose, to the contrary, that $G - F$ has no fractional perfect matchings. By Proposition 1.1, there exists $S \subseteq V(G - F)$ such that $i(G - (F \cup S)) \geq |S| + 1$. By Lemma 2.6,

$$|G - (F \cup S)| = \frac{|G| + |F_E| - \delta}{2} + 1. \quad (4)$$

and

$$\alpha(G) \geq |G - (F \cup S)| - |F_E| = \frac{|G| - |F_E| - \delta}{2} + 1. \quad (5)$$

(i) Note that $|F_E| \leq |F| = \delta - 1$, then by the inequality (5), $\alpha(G) \geq \frac{|G| - |F_E| - \delta}{2} + 1 \geq \frac{|G| + 1}{2} - \delta + 1$, a contradiction to the assumption.

(ii) By the assumption $\alpha(G) \leq \lceil \frac{|G| + 1}{2} \rceil - \delta + 1$ and the inequality (5), we have

$$\frac{|G| - |F_E| - \delta}{2} + 1 \leq |G - (F \cup S)| - |F_E| \leq \alpha(G) \leq \lceil \frac{|G| + 1}{2} \rceil - \delta + 1, \quad (6)$$

which implies $|F_E| = \delta - 1$ when $|G|$ is odd and $\delta - 2 \leq |F_E| \leq \delta - 1$ when $|G|$ is even, and thus $|F_E| = \delta - 2$ as Lemma 3.4. Combining this with the inequality (6), we have

$$\alpha(G) = |G - (F \cup S)| - |F_E| = \lceil \frac{|G| + 1}{2} \rceil - \delta + 1. \quad (7)$$

Consider a partition $\{V(G) - (F_V \cup S), F_V \cup S\}$ of $V(G)$. Since G is δ -regular, we can deduce $|E(G[F_V \cup S])| = |E(G)| - (\delta|G - (F \cup S)| - |F_E|)$.

If $|G|$ is odd, then $|F_E| = \delta - 1 \geq 2$ as $\delta \geq 3$. By (4), $|G - (F \cup S)| = \frac{|G| + 1}{2}$, and thus $|F_V \cup S| = \frac{|G| - 1}{2}$. Then $|E(G[F_V \cup S])| = \frac{\delta|G|}{2} - (\frac{\delta(|G| + 1)}{2} - (\delta - 1)) = \frac{\delta}{2} - 1$. This together with (7), we have $|G[F_V \cup S]| - |E(G[F_V \cup S])| = \frac{|G| - 1}{2} - (\frac{\delta}{2} - 1) = \frac{|G| + 1}{2} - \frac{\delta}{2} > \lceil \frac{|G| + 1}{2} \rceil - \delta + 1 = \alpha(G) \geq \alpha(G[F_V \cup S])$, a contradiction to Lemma 2.4.

If $|G|$ is even, then $|F_E| = \delta - 2 \geq 1$ as $\delta \geq 3$. By (4), $|G - (F \cup S)| = \frac{|G|}{2}$, and thus $|F_V \cup S| = \frac{|G|}{2}$. Then $|E(G[F_V \cup S])| = \frac{\delta|G|}{2} - (\frac{\delta|G|}{2} - (\delta - 2)) = \delta - 2$. Note that $G - (F \cup S)$ is an independent set as Lemma 2.6. By the equality (7) and Observation 2.5, $E(G - (F_V \cup S)) = F_E$ and F_E is a matching of $G - (F_V \cup S)$. This implies every edge of $G - (F_V \cup S)$ lies in a 4-cycle which must contain an edge of $G[F_V \cup S]$. Suppose that $uvxy$ is a 4-cycle with $uv \in E(G - (F_V \cup S))$ and $xy \in E(G[F_V \cup S])$. By the equality (7) and Lemma 2.4, $\frac{|G| + 2}{2} - \delta + 1 = \alpha(G) \geq \alpha(G[F_V \cup S]) \geq |G[F_V \cup S]| - |E(G - (F_V \cup S))| = \frac{|G|}{2} - (\delta - 2) = \frac{|G| + 2}{2} - \delta + 1$. Then, by Lemma 2.4, $E(G[F_V \cup S])$ is a matching of $G[F_V \cup S]$. Since G is $\{K_4 - e\}$ -free, then there is no 3-cycles containing vx in G , a contradiction to $G \in \mathcal{G}_1(\delta)$.

(iii) By the assumption $\alpha(G) \leq \frac{|G|+1}{2} - 2$ and the inequality (5), we have

$$\frac{|G| - |F_E|}{2} - 1 \leq \alpha(G) \leq \frac{|G| - 1}{2} - 1, \quad (8)$$

which implies $|F_E| \geq 1$. Note that $|F_E| \leq |F| = 3$ and $|G|$ is odd, then by Lemma 3.4, $|F_V| \neq 1$, and thus $|F_E| \neq 2$. This implies $|F_E| = 1$ or $|F_E| = 3$. Since G is 4-regular, we can deduce $|E(G[F_V \cup S])| = |E(G)| - (4|G - (F \cup S)| - |E(G - (F_V \cup S))|)$.

If $|F_E| = 1$, then $|E(G - (F_V \cup S))| \leq |F_E| = 1$. Assume that $|E(G - (F_V \cup S))| = 0$. Then by (4), $\alpha(G) \geq |G - (F_V \cup S)| = \frac{|G|+|F_E|-\delta}{2} + 1 = \frac{|G|-1}{2}$, a contradiction to (8). Thus $|E(G - (F_V \cup S))| = |F_E| = 1$. By (4), we have $|G - (F \cup S)| = \frac{|G|-1}{2}$. Then $|E(G[F_V \cup S])| = \frac{4|G|}{2} - (\frac{4(|G|-1)}{2} - 1) = 3$. Note that $|E(G - (F_V \cup S))| = 1$, then every edge of $E(G - (F_V \cup S))$ lies in a 4-cycle which must contain an edge of $G[F_V \cup S]$. Let $u_1v_1x_1y_1$ and $u_1v_1w_1z_1$ be two 4-cycles with $u_1v_1 \in E(G - (F_V \cup S))$ and $x_1y_1, w_1z_1 \in E(G[F_V \cup S])$, where x_1, y_1, w_1, z_1 are four distinct vertices as G is $\{K_4, K_4 - e, K_{2,3}\}$ -free. Note that v_1x_1 lies in a 3-cycle. Let v_1x_1v' be a 3-cycle with $x_1v' \in E(G[F_V \cup S])$ and $v' \neq y_1$ as G is $\{K_4 - e\}$ -free. Recall that $|E(G[F_V \cup S])| = 3$, there is no 3-cycles containing u_1y_1 in G , a contradiction to $G \in \mathcal{G}_2(4)$.

If $|F_E| = 3$, then $|E(G - (F_V \cup S))| \leq |F_E| = 3$. By (4), we have $|G - (F \cup S)| = \frac{|G|+1}{2}$, and thus $|F_V \cup S| = |G| - |G - (F \cup S)| = \frac{|G|-1}{2}$. Then $|E(G[F_V \cup S])| \leq \frac{4|G|}{2} - (\frac{4(|G|+1)}{2} - 3) = 1$. Assume that $|E(G[F_V \cup S])| = 0$. Then $\alpha(G) \geq |F_V \cup S| = \frac{|G|-1}{2}$, a contradiction to (8). Thus $|E(G[F_V \cup S])| = 1$. This implies every edge of $G[F_V \cup S]$ lies in a 4-cycle which must contain an edge of $E(G - (F_V \cup S))$. Let $u_2v_2x_2y_2$ and $u_2v_2w_2z_2$ be two 4-cycles with $u_2v_2 \in E(G[F_V \cup S])$ and $x_2y_2, w_2z_2 \in E(G - (F_V \cup S))$, where x_2, y_2, w_2, z_2 are four distinct vertices as G is $\{K_4, K_4 - e, K_{2,3}\}$ -free. Note that v_2x_2 lies in a 3-cycle. Let v_2x_2v'' be a 3-cycle with $x_2v'' \in E(G - (F_V \cup S))$ and $v'' \neq y_2$ as G is $\{K_4 - e\}$ -free. Since $|E(G - (F_V \cup S))| \leq 3$, there is no 3-cycles containing u_2y_2 in G , a contradiction to $G \in \mathcal{G}_2(4)$. ■

4 Applications to some networks

In the following, we will determine the FMP number and FSMP number of some well-known networks by the conclusions in Section 3.

4.1 Restricted HL-graphs

The restricted HL-graph is defined using a special graph construction operator. Given two graphs G_0 and G_1 , consider a set $\Phi(G_0, G_1)$, made of all bijections from $V(G_0)$ to $V(G_1)$.

Then, given a bijection $\phi \in \Phi(G_0, G_1)$, we denote by $G_0 \oplus_\phi G_1$ a graph whose vertex set is $V(G_0) \cup V(G_1)$ and edge set is $E(G_0) \cup E(G_1) \cup \{(v, \phi(v)) : v \in V(G_0)\}$. Based on the graph constructor, Vaidya *et al.* [33] gave a recursive definition of a class of graphs as follows.

Definition 4.1 [33] Let $RHL_0 = \{K_1\}$, $RHL_1 = \{K_2\}$, $RHL_2 = \{C_4\}$, $RHL_3 = \{G(8, 4)\}$ and $RHL_n = \{G_0 \oplus_\phi G_1 : G_0, G_1 \in RHL_{n-1}, \phi \in \Phi(G_0, G_1)\}$ for $n \geq 4$. A graph that belongs to RHL_n , denoted by G^n , is called an n -dimensional restricted HL-graph.




Fig. 1 $G(8, 4)$

The graph $G(8, 4)$ is shown in Fig. 1. G^n is n -regular with 2^n vertices. Many of the non-bipartite hypercube-like interconnection networks such as crossed cube [10], Möbius cube [7], twisted cube [14], multiply twisted cube [9], generalized twisted cube [4], locally twisted cube [35], the twisted hypercubes [37] etc. proposed in the literature are restricted HL-graphs.

Lemma 4.2 $\alpha(G^n) \leq 3 \times 2^{n-3}$ for $n \geq 3$.

Proof. Clearly, $\alpha(G^3) = 3$ (see Fig. 1, the green dots represent the vertices of a maximum independent set of G^3). By induction, suppose that $\alpha(G^{n-1}) \leq 3 \times 2^{n-4}$ for $n \geq 4$. Note that G^n can be decomposed into two vertex disjoint subgraphs each of which is isomorphic to G_{n-1} , and thus every independent set of G^n contains at most $2 \times (3 \times 2^{n-4}) = 3 \times 2^{n-3}$ vertices. Then $\alpha(G^n) \leq 3 \times 2^{n-3}$ for $n \geq 3$. ■

Park *et al.* [29] considered Hamiltonian properties in faulty restricted HL-graphs.

Lemma 4.3 [29] G^n is $(n - 2)$ -fault Hamiltonian for $n \geq 3$.

Now we can determine the FMP number and FSMP number of G^n , which was also obtained in [36].

Theorem 4.4 $fsm p(G^n) = fsm p(G^n) = n$ for $n \geq 5$.

Proof. Note that G^n is n -regular with 2^n vertices, and thus $|G^n|$ is even. By Lemmas 4.2, 4.3 and Theorem 3.5(i), it suffices to show that

$$3 \times 2^{n-3} \leq \frac{2^n + 2}{2} - n,$$

which implies $2^{n-3} - n + 1 \geq 0$. It is obvious that the inequality $2^{n-3} - n + 1 \geq 0$ holds if $n \geq 5$. Therefore, we complete the proof of Theorem 4.4. ■

4.2 Torus networks

Torus networks have been proved to be a viable choice for the interconnection networks, such as ease of implementation, low latency, and high-bandwidth inter-processor communication.

Definition 4.5 [33] Given k_1, \dots, k_n with $k_i \geq 3$, the n -dimensional torus, denoted by T_{k_1, \dots, k_n} , has the set $\{v_{x_1, x_2, \dots, x_n} \mid 0 \leq x_i \leq k_i - 1, 1 \leq i \leq n\}$ as its vertex set. Two vertices x_{a_1, a_2, \dots, a_n} and y_{b_1, b_2, \dots, b_n} are adjacent if there exists an integer i with $1 \leq i \leq n$ such that $a_i - b_i = \pm 1 \pmod{k_i}$, and $a_j = b_j$ for $1 \leq j \neq i \leq n$.




Fig. 2 Two 2-dimensional torus networks

T_{k_1, \dots, k_n} is $(2n)$ -regular with $k_1 \cdots k_n$ vertices and vertex transitive. T_{k_1, \dots, k_n} is bipartite if and only if all k_1, \dots, k_n are even. Fig. 2 shows the 2-dimensional torus $T_{3,4}$ which is non-bipartite, and $T_{4,4}$ which is bipartite. In particular, the n -dimensional torus T_{k_1, \dots, k_n} is said to be a k -ary n -cube Q_n^k if $k_i = k$ for each $1 \leq i \leq n$. Let $T_{k_1, \dots, k_{n-1}}^i$ be the subgraph

of T_{k_1, \dots, k_n} induced by the vertices with i in the n -th position where $0 \leq i \leq k_n - 1$. Then T_{k_1, \dots, k_n} can be decomposed into k_n vertex disjoint such subgraphs, each of which is isomorphic to $T_{k_1, \dots, k_{n-1}}$ (see Fig. 2, each red cycle in $T_{3,4}$ represents T_3^i for $0 \leq i \leq 3$ and each red cycle in $T_{4,4}$ represents T_4^j for $0 \leq j \leq 3$).

Lemma 4.6 $\alpha(T_{k_1, \dots, k_n}) \leq \frac{(k_1-1)k_2 \cdots k_n}{2}$ for $n \geq 1$.

Proof. Clearly, $\alpha(T_{k_1}) = \frac{k_1-1}{2}$. By induction, suppose that $\alpha(T_{k_1, \dots, k_{n-1}}) \leq \frac{(k_1-1)k_2 \cdots k_{n-1}}{2}$ for $n \geq 2$. Note that T_{k_1, \dots, k_n} can be decomposed into k_n vertex disjoint subgraphs, each of which is isomorphic to $T_{k_1, \dots, k_{n-1}}$, and thus every independent set of T_{k_1, \dots, k_n} contains at most $k_n \times \frac{(k_1-1)k_2 \cdots k_{n-1}}{2} = \frac{(k_1-1)k_2 \cdots k_n}{2}$ vertices. Then $\alpha(T_{k_1, \dots, k_n}) \leq \frac{(k_1-1)k_2 \cdots k_n}{2}$ for $n \geq 1$. ■

Kim and Park [20] considered Hamiltonian properties in n -dimensional torus networks with faults.

Lemma 4.7 [20] T_{k_1, \dots, k_n} is $(2n-2)$ -fault Hamiltonian for $n \geq 2$.

By Theorem 3.1, Lemmas 3.2 and 4.7, we can determine the FMP number and FSMP number of bipartite n -dimensional torus networks easily.

Theorem 4.8 Let T_{k_1, \dots, k_n} be bipartite with $n \geq 2$. Then

$$fmp(T_{k_1, \dots, k_n}) = 2n \quad \text{and} \quad fsmf(T_{k_1, \dots, k_n}) = 1.$$

In the following, we always assume T_{k_1, \dots, k_n} is non-bipartite, and thus k_i is odd for some $1 \leq i \leq n$. By symmetry, we can choose k_1, \dots, k_n such that

(T1) k_i is odd for $1 \leq i \leq t$ with t as large as possible.

(T2) $k_1 \leq \dots \leq k_t$, subject to (T1).

Therefore, k_1 is odd. The following lemma gives the FMP number and FSMP number of $T_{k_1, 4}$.

Lemma 4.9 $fsmf(T_{k_1, 4}) = 3$ and $fmp(T_{k_1, 4}) = 4$.

Proof. Let $F = \{v_{0,0}, v_{1,0}, v_{2,0}, v_{1,2}, v_{2,2}\}$ and $S_1 = \{v_{0,2}, v_{1,1}, v_{1,3}, v_{2,1}, v_{2,3}\}$. Then $i(T_{3,4} - F - S_1) = 6 > 5 = |S_1|$ (see Fig. 3(a), the red dots represent the vertices of S_1 , the black dots represent the isolated vertices of $T_{3,4} - F - S_1$), and thus, by Proposition 1.1, $T_{3,4} - F$ has no fractional perfect matchings. It follows that F is an FSMP set of $T_{3,4}$, then $fsmf(T_{3,4}) \leq 3$. Note that $fsmf(T_{3,4}) \geq 3$ by Lemma 3.3. Hence $fsmf(T_{3,4}) = 3$.




Fig. 3 $T_{k_1,4}$ with $k_1 \geq 3$ and k_1 is odd.

Let $S_2 = \{v_{0,2}, v_{1,1}, v_{1,3}, v_{2,1}, v_{2,3}\} \cup \{v_{3,0}, v_{3,2}\} \cup \dots \cup \{v_{k_1-1,1}, v_{k_1-1,3}\}$. Then $i(T_{k_1,4} - F - S_2) = 2k_1 > 2k_1 - 1 = |S_2|$ (see Fig. 3(b), the red dots represent the vertices of S_2 , the black dots represent the isolated vertices of $T_{k_1,4} - F - S_2$), and thus, by Proposition 1.1, $T_{k_1,4} - F$ has no fractional perfect matchings. It follows that F is an $FSMP$ set of $T_{k_1,4}$, then $fsmpt(T_{k_1,4}) \leq 3$. Note that $fsmpt(T_{k_1,4}) \geq 3$ by Lemma 3.3. Hence $fsmpt(T_{k_1,4}) = 3$.

Note that $T_{k_1,4}$ is 4-regular graph with $4k_1$ vertices. By Lemmas 3.2 and 4.7, $fmp(T_{k_1,4}) = 4$. Therefore, we complete the proof of Lemma 4.9. ■

Lemma 4.10 $fsmpt(T_{k_1,k_2}) = fmp(T_{k_1,k_2}) = 4$ for $k_2 \neq 4$.

Proof. If k_2 is even, then $|T_{k_1,k_2}| = k_1k_2$ is even and $k_2 \geq 6$ as $k_2 \neq 4$. By Lemma 4.6, $\alpha(T_{k_1,k_2}) \leq \frac{(k_1-1)k_2}{2} \leq \frac{k_1k_2+2}{2} - 4$. Then $fsmpt(T_{k_1,k_2}) = fmp(T_{k_1,k_2}) = 4$ as Theorem 3.5(i).

If k_2 is odd, then $|T_{k_1,k_2}| = k_1k_2$ is odd. By Lemma 4.6, $\alpha(T_{3,3}) \leq 3 = \frac{9+1}{2} - 4 + 2$. Note that $T_{3,3} \in \mathcal{G}_2(4)$, then $fsmpt(T_{3,3}) = fmp(T_{3,3}) = 4$ as Theorem 3.5(iii). Now, suppose that $T_{k_1,k_2} \neq T_{3,3}$, and thus $k_2 \geq 5$. By Lemma 4.6, $\alpha(T_{k_1,k_2}) \leq \frac{(k_1-1)k_2}{2} \leq \frac{k_1k_2+1}{2} - 4 + 1$. Note that $T_{k_1,k_2} \in \mathcal{G}_1(4)$, then $fsmpt(T_{k_1,k_2}) = fmp(T_{k_1,k_2}) = 4$ as Theorem 3.5(ii). ■

Lemma 4.11 $fsmpt(T_{k_1,\dots,k_n}) = fmp(T_{k_1,\dots,k_n}) = 2n$ for $n \geq 3$.

Proof. Note that T_{k_1,\dots,k_n} is $(2n)$ -regular with $k_1 \cdots k_n$ vertices. By Lemmas 4.6, 4.7 and Theorem 3.5(i), it suffices to show that

$$\frac{(k_1 - 1)k_2 \cdots k_n}{2} \leq \frac{k_1 \cdots k_n + 1}{2} - 2n,$$

which implies $k_2 \cdots k_n - 4n + 1 \geq 0$. It is obvious that the inequality $k_2 \cdots k_n - 4n + 1 \geq 0$ holds if $n \geq 4$. Particularly, if $n = 3$, then the inequality $k_2 k_3 - 4 \times 3 + 1 \geq 0$ holds if $(k_2, k_3) \neq (3, 3)$. By the choice of k_i , we have $k_1 \neq 3$. By Lemma 4.6, $\alpha(T_{3,3,3}) \leq 9 = \frac{27+1}{2} - 6 + 1$. Note that $T_{3,3,3} \in \mathcal{G}_1(6)$, then $fsm p(T_{3,3,3}) = fmp(T_{3,3,3}) = 6$ as Theorem 3.5(ii). Therefore, we complete the proof of Lemma 4.11. ■

Combining with Lemmas 4.9, 4.10 and 4.11, we obtain the FMP number and FSMP number of non-bipartite n -dimensional torus networks.

Theorem 4.12 *Let T_{k_1, \dots, k_n} be non-bipartite with $n \geq 2$. Then $fmp(T_{k_1, \dots, k_n}) = 2n$ and*

$$fsm p(T_{k_1, \dots, k_n}) = \begin{cases} 3, & \text{if } (n, k_2) = (2, 4); \\ 2n, & \text{otherwise.} \end{cases}$$

4.3 Recursive circulant graphs

The recursive circulant graph has many nice properties, such as, vertex transitive, strongly hierarchical, higher connectivity which increases the fault tolerance, smaller diameter which reduces the transmission delay, etc. Park and Chwa [27] first introduced the concept of the recursive circulant graph $G(cd^n, d)$ with $n \geq 1$ and $1 \leq c < d$ as follows.

Definition 4.13 [27] *The recursive circulant graph $G(cd^n, d)$ with $n \geq 1$ and $1 \leq c < d$, has the vertex set $V = \{0, 1, \dots, cd^n - 1\}$, and the edge set $E = \{ (u, v) \mid u - v \equiv cd^i \pmod{cd^n} \text{ and } 0 \leq i \leq \lceil \log_d c \rceil + n - 1 \}$.*




Fig. 4 Two recursive circulant graphs $G(3^2, 3)$

4.3.1 The recursive circulant graph $G(d^n, d)$

By the definition of 4.13, when $c = 1$, $G(d^n, d)$ with $n \geq 1$, $d \geq 3$ is $(2n)$ -regular with d^n vertices and vertex transitive. The recursive circulant graph $G(3^2, 3)$ is shown in Fig. 4(a). Let $V_{d,i}^{n-1} = \{v \in V(G(d^n, d)) \mid v \equiv i \pmod{d}\}$ and $G_{d,i}^{n-1}$ be the subgraph of $G(d^n, d)$ induced by the vertices of $V_{d,i}^{n-1}$. Park and Chwa [28] proved that $G(d^n, d)$ can be decomposed into d vertex disjoint such subgraphs, each of which is isomorphic to $G(d^{n-1}, d)$ (see Fig. 4(a), each red odd cycle in $G(3^2, 3)$ represents $G_{3,i}^1$ for $0 \leq i < 3$).

In order to be convenient to study the fractional (strong) matching preclusion of $G(d^2, d)$, we relabel vertices of $G(d^2, d)$ such that the vertex $i \times d + j$ corresponds to $v_{i,j}$ for $0 \leq i, j < d$. Therefore, the vertex set of $G(d^2, d)$ is represented as

$$V(G(d^2, d)) = \{v_{i,j} : 0 \leq i, j < d\}.$$

The edge set of $G(d^2, d)$ is classified into two sets:

$$\begin{aligned} E_1 &= \{v_{i,k}v_{j,k} : j \equiv i + 1 \pmod{d} \text{ and } 0 \leq i, j, k < d\}; \\ E_2 &= \{v_{i,j}v_{i,k} : k = j + 1 \text{ and } 0 \leq i, j, k < d\} \cup \\ &\quad \{v_{i,0}v_{j,d-1} : j \equiv i - 1 \pmod{d} \text{ and } 0 \leq i, j < d\}. \end{aligned}$$

In Fig. 4(b), The vertex $3i + j$ corresponds to $v_{i,j}$ for $0 \leq i, j < 3$.

Lemma 4.14 $\alpha(G(d^2, d)) = \frac{d^2-d}{2}$.

Proof. See Appendix. ■

Lemma 4.15 $\alpha(G(d^n, d)) \leq \frac{d^n-d^{n-1}}{2}$ for $n \geq 2$.

Proof. The conclusion holds for $n = 2$ as Lemma 4.14. By induction on n , suppose that $\alpha(G(d^{n-1}, d)) \leq \frac{d^{n-1}-d^{n-2}}{2}$ for $n \geq 3$. Note that $G(d^n, d)$ can be decomposed into d vertex disjoint subgraphs, each of which is isomorphic to $G(d^{n-1}, d)$, and thus every independent set of $G(d^n, d)$ contains at most $d \times \frac{d^{n-1}-d^{n-2}}{2} = \frac{d^n-d^{n-1}}{2}$ vertices. Then $\alpha(G(d^n, d)) \leq \frac{d^n-d^{n-1}}{2}$ for $n \geq 2$ and $d \geq 3$. ■

Tsai *et al.* [32] researched Hamiltonian properties of faulty recursive circulant graphs.

Lemma 4.16 [32] $G(d^n, d)$ is $(2n - 2)$ -fault Hamiltonian for $n \geq 2$.

The following two lemmas determine the FMP number and FSMP number of $G(3^2, 3)$ and $G(4^2, 4)$, respectively.




Fig. 5 $G(3^2, 3) - F_i$ with $i = 1, 2$.

Lemma 4.17 $fsm p(G(3^2, 3)) = fmp(G(3^2, 3)) = 3$.

Proof. First we show that $fsm p(G(3^2, 3)) = 3$. Let $F_1 = \{v_{0,0}v_{1,0}, v_{0,0}v_{2,0}, v_{0,1}v_{1,1}, v_{0,1}v_{2,1}, v_{0,2}v_{1,2}, v_{0,2}v_{2,2}\}$ and $S_1 = \{v_{0,1}, v_{1,2}, v_{2,1}\}$. Then $i(G(3^2, 3) - F_1 - S_1) = 4 > 3 = |S_1|$ (see Fig. 5(a), the red dots represent the vertices of S_1 , the black dots represent the isolated vertices of $G(3^2, 3) - F_1 - S_1$), and thus, by Proposition 1.1, $G(3^2, 3) - F_1$ has no fractional perfect matchings. It follows that $fsm p(G(3^2, 3)) \leq 3$. Note that $fsm p(G(3^2, 3)) \geq 3$ by Lemma 3.3. Hence $fsm p(G(3^2, 3)) = 3$.

Let $F_2 = \{v_{0,0}v_{2,0}, v_{0,0}v_{2,2}, v_{0,1}v_{1,0}, v_{0,1}v_{1,2}, v_{0,2}v_{1,1}, v_{0,2}v_{2,1}\}$ and $S_2 = \{v_{0,1}, v_{1,0}, v_{1,2}, v_{2,1}\}$. Then $i(G(3^2, 3) - F_2 - S_2) = 5 > 4 = |S_2|$ (see Fig. 5(b), the red dots represent the vertices of S_2 , the black dots represent the isolated vertices of $G(3^2, 3) - F_2 - S_2$), and thus, by Proposition 1.1, $G(3^2, 3) - F_2$ has no fractional perfect matchings. It follows that $fmp(G(3^2, 3)) \leq 3$. Note that $fmp(G(3^2, 3)) \geq 3$ by Lemma 3.3. Hence $fmp(G(3^2, 3)) = 3$. ■

Lemma 4.18 $fsm p(G(4^2, 4)) = 3$ and $fmp(G(4^2, 4)) = 4$.

Proof. By Lemmas 3.2 and 4.16, $fmp(G(4^2, 4)) = 4$. Now, we show that $fsm p(G(4^2, 4)) = 3$. Let $F_3 = \{v_{0,0}v_{1,0}v_{0,3}, v_{0,0}v_{2,0}v_{2,3}, v_{0,1}v_{1,1}v_{1,3}, v_{0,1}v_{2,1}v_{2,3}, v_{0,2}v_{1,2}v_{1,3}, v_{0,2}v_{2,2}v_{2,3}\}$ and $S_3 = \{v_{0,2}, v_{1,1}, v_{1,3}, v_{2,0}, v_{2,2}, v_{3,1}, v_{3,3}\}$. Then $i(G(4^2, 4) - F_3 - S_3) = 8 > 7 = |S_3|$ (see Fig. 6, the red dots represent the vertices of S_3 , the black dots represent the isolated vertices of $G(4^2, 4) - F_3 - S_3$), and thus, by Proposition 1.1, $G(4^2, 4) - F_3$ has no fractional perfect matchings. It follows that $fsm p(G(4^2, 4)) \leq 3$. Note that $fsm p(G(4^2, 4)) \geq 3$ by Lemma 3.3. Hence $fsm p(G(4^2, 4)) = 3$. ■




Fig. 6 $G(4^2, 4) - \{v_{0,0}, v_{1,0}v_{0,3}, v_{3,0}v_{2,3}\}$

Lemma 4.19 $fsm(p(G(d^2, d)) = fsm(p(G(d^2, d)) = 4$ for $d \geq 5$.

Proof. If d is even, then $|G(d^2, d)| = d^2$ is even and $d \geq 6$ as $d \geq 5$. By Lemma 4.15, $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2} \leq \frac{d^2+2}{2} - 4$. Then $fsm(p(T_{k_1, k_2})) = fsm(p(T_{k_1, k_2})) = 4$ as Theorem 3.5(i). If d is odd, then $|G(d^2, d)| = d^2$ is odd. By Lemma 4.15, $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2} \leq \frac{d^2+1}{2} - 4 + 1$ for $d \geq 5$. Note that $G(d^2, d) \in \mathcal{G}_1(4)$, then $fsm(p(T_{k_1, k_2})) = fsm(p(T_{k_1, k_2})) = 4$ as Theorem 3.5(ii). ■

Lemma 4.20 $fsm(p(G(d^n, d)) = fsm(p(G(d^n, d)) = 2n$ for $n \geq 3$.

Proof. Note that $G(d^n, d)$ is $(2n)$ -regular with d^n vertices. By Lemmas 4.15, 4.16 and Theorem 3.5(i), it suffices to show that

$$\frac{d^n - d^{n-1}}{2} \leq \frac{d^n + 1}{2} - 2n,$$

which implies $d^{n-1} - 4n + 1 \geq 0$. It is obvious that the inequality $d^{n-1} - 4n + 1 \geq 0$ holds if $n \geq 4$. Particularly, if $n = 3$, then the inequality $d^2 - 12 + 1 \geq 0$ holds if $d \geq 4$. By Lemma 4.15, $\alpha(G(3^3, 3)) \leq 9 = \frac{3^3+1}{2} - 6 + 1$. Note that $G(3^3, 3) \in \mathcal{G}_1(6)$, then $fsm(p(G(3^3, 3))) = fsm(p(G(3^3, 3))) = 6$ as Theorem 3.5(ii). Therefore, we complete the proof of Lemma 4.20. ■

Combining with Lemmas 4.17, 4.18, 4.19 and 4.20, we obtain the FMP number and FSMP number of $G(d^n, d)$.

Theorem 4.21 *Let $n \geq 2$ and $d \geq 3$ be integers. Then*

$$fmp(G(d^n, d)) = \begin{cases} 3, & \text{if } (n, d) = (2, 3); \\ 2n, & \text{otherwise,} \end{cases}$$

and

$$fsm(G(d^n, d)) = \begin{cases} 3, & \text{if } (n, d) \in \{(2, 3), (2, 4)\}; \\ 2n, & \text{otherwise.} \end{cases}$$

4.3.2 The recursive circulant graph $G(2^n, 4)$

Park and Chwa [27] introduced an interesting recursive circulant graph $G(2^n, 4)$, which is n -regular with 2^n vertices. By Lemma 3.2, $fmp(G(2^n, 4)) = n$.

In the following, we establish the FSMP number of $G(2^n, 4)$. Park and Ihm [30] showed that $G(2^n, 4)$ with odd n is an n -dimensional restricted HL-graph, whose fractional (strong) matching preclusion properties were analyzed in [36].

Lemma 4.22 [36] *Let $n \geq 3$ be an odd. Then $fsm(G(2^3, 4)) = 2$ and $fsm(G(2^n, 4)) = n$ for $n \geq 5$.*

If n is even, then by Theorem 4.21, $fsm(G(2^4, 4)) = 3$ and $fsm(G(2^n, 4)) = n$ for $n \geq 6$. Combining this with Lemma 4.22, we obtain fractional (strong) matching preclusion of $G(2^n, 4)$ with $n \geq 3$ as follows.

Theorem 4.23 *Let $n \geq 3$ be an integer. Then $fmp(G(2^n, 4)) = n$ and*

$$fsm(G(2^n, 4)) = \begin{cases} n - 1, & \text{if } n \in \{3, 4\}; \\ n, & \text{otherwise.} \end{cases}$$

4.4 (n, k) -arrangement graphs and (n, k) -star graphs

The (n, k) -arrangement graph [8] and (n, k) -star graph [6] are two generalization versions of the star graph S_n . The two parameters n and k can be tuned to make a suitable choice for the number of nodes in the network and for the degree/diameter tradeoff.

4.4.1 (n, k) -arrangement graphs

Definition 4.24 [8] *The (n, k) -arrangement graph, denoted by $A_{n,k}$, is defined for positive integers n and k such that $1 \leq k \leq n - 1$. The vertex set of the graph is all the permutations on k elements of the set $\{1, 2, \dots, n\}$. Two vertices $a_1 a_2 \dots a_k$ and $b_1 b_2 \dots b_k$ are adjacent if there exists an integer s with $1 \leq s \leq k$ such that $a_s \neq b_s$ and for any $i \neq s$, $1 \leq i \leq k$, we have $a_i = b_i$.*




Fig. 7 The $(4, 2)$ -arrangement graph $A_{4,2}$

The $(4, 2)$ -arrangement graph $A_{4,2}$ is shown in Fig. 7. $A_{n,1}$ is isomorphic to the complete graph K_n , $A_{n,n-2}$ are isomorphic to the n -alternating group graph AG_n and $A_{n,n-1}$ is isomorphic to the n -dimensional star graph S_n . $A_{n,k}$ is $k(n-k)$ -regular with $P_n^k = \frac{n!}{(n-k)!}$ vertices. In [8], Day and Tripathi proved that $A_{n,k}$ can be decomposed into n vertex disjoint subgraphs, each of which is isomorphic to $A_{n-1,k-1}$ (see Fig. 7, each red triangle in $A_{4,2}$ is isomorphic to $A_{3,1} = K_3$).

Lemma 4.25 $\alpha(A_{n,k}) \leq P_n^{k-1}$ for $1 \leq k \leq n-2$.

Proof. Clearly, $\alpha(A_{n,1}) = 1$. By induction on k , suppose that $\alpha(A_{n-1,k-1}) \leq P_{n-1}^{k-2}$ for $2 \leq k \leq n-2$. Note that $A_{n,k}$ can be decomposed into n vertex disjoint subgraphs, each of which is isomorphic to $A_{n-1,k-1}$, and thus every independent set of $A_{n,k}$ contains at most $n \times P_{n-1}^{k-2} = P_n^{k-1}$ vertices. Then $\alpha(A_{n,k}) \leq P_n^{k-1}$. ■

Hsu *et al.* [16] investigated fault Hamiltonicity of the arrangement graphs.

Lemma 4.26 [16] $A_{n,k}$ is $(k(n-k)-2)$ -fault Hamiltonian for $2 \leq k \leq n-2$.

Now we can determine the FMP number and FSMP number of $A_{n,k}$, which was also obtained in [25].

Theorem 4.27 $fsm p(A_{n,k}) = fmp(A_{n,k}) = k(n-k)$ for $2 \leq k \leq n-2$.

Proof. Note that $A_{n,k}$ is $k(n-k)$ -regular with P_n^k vertices, and thus $|A_{n,k}|$ is even. By Lemmas 4.25, 4.26 and Theorem 3.5(i), it suffices to show that

$$P_n^{k-1} \leq \frac{P_n^k + 2}{2} - k(n-k),$$

which implies $(n-k-1)P_n^{k-1} - 2k(n-k) + 2 \geq 0$. It is obvious that the inequality $(n-k-1)P_n^{k-1} - 2k(n-k) + 2 \geq 0$ holds if $A_{n,k} \neq A_{4,2}$. By Lemma 4.25, $\alpha(A_{4,2}) \leq 4 = \frac{12+2}{2} - 4 + 1$. Note that $A_{4,2} \in \mathcal{G}_1(4)$, then $fsm p(A_{4,2}) = fmp(A_{4,2}) = 4$ as Theorem 3.5(ii). Therefore, we complete the proof of Theorem 4.27. ■

4.4.2 (n, k) -star graphs

Definition 4.28 [6] The (n, k) -star graph, denoted by $S_{n,k}$, is defined for positive integers n and k such that $1 \leq k \leq n-1$. The vertex set of the graph is all the permutations on k elements of the set $\{1, 2, \dots, n\}$. Two vertices corresponding to the permutations $a_1 a_2 \dots a_k$ and $b_1 b_2 \dots b_k$ are adjacent if and only if either:

- (1) There exists an integer $2 \leq s \leq k$ such that $a_1 = b_s$ and $b_1 = a_s$ and for any $i \neq s$, $2 \leq i \leq k$, we have $a_i = b_i$. That is, $b_1 b_2 \dots b_k$ is obtained from $a_1 a_2 \dots a_k$ by swapping a_1 and a_s .
- (2) For all $2 \leq i \leq k$, we have $a_i = b_i$ and $a_1 \neq b_1$. That is, $b_1 b_2 \dots b_k$ is obtained from $a_1 a_2 \dots a_k$ by replacing a_1 by an element in $\{1, 2, \dots, n\} - \{a_1, a_2, \dots, a_k\}$.




Fig. 8 The $(4, 2)$ -star graph $S_{4,2}$

The $(4, 2)$ -star graph $S_{4,2}$ is depicted in Fig. 8. $S_{n,1}$ is isomorphic to the complete graph K_n and $S_{n,n-1}$ is isomorphic to the n -dimensional star graph S_n . $S_{n,k}$ is $(n-1)$ -regular

with $P_n^k = \frac{n!}{(n-k)!}$ vertices. In [6], Chiang and Chen proved that $S_{n,k}$ can be decomposed into n vertex disjoint subgraphs, each of which is isomorphic to $S_{n-1,k-1}$ (see Fig. 8, each red triangle in $S_{4,2}$ is isomorphic to $S_{3,1} = K_3$).

Lemma 4.29 $\alpha(S_{n,k}) \leq P_n^{k-1}$ for $1 \leq k \leq n-2$.

Proof. Clearly, $\alpha(S_{n,1}) = 1$. By induction on k , suppose that $\alpha(S_{n-1,k-1}) \leq P_{n-1}^{k-2}$ for $2 \leq k \leq n-2$. Note that $S_{n,k}$ can be decomposed into n vertex disjoint subgraphs, each of which is isomorphic to $S_{n-1,k-1}$, and thus every independent set of $S_{n,k}$ contains at most $n \times P_{n-1}^{k-2} = P_n^{k-1}$ vertices. Then $\alpha(S_{n,k}) \leq P_n^{k-1}$. ■

Hsu *et al.* [15] considered Hamiltonian properties of faulty (n, k) -star graphs.

Lemma 4.30 [15] $S_{n,k}$ is $(n-3)$ -fault Hamiltonian for $2 \leq k \leq n-2$.

By Lemmas 4.29, 4.30 and Theorem 3.5(i), we can determine the FMP number and FSMP number of $S_{n,k}$.

Theorem 4.31 $fsm_p(S_{n,k}) = fmp(S_{n,k}) = n-1$ for $2 \leq k \leq n-2$.

4.5 (Burnt) pancake graphs

Pancake graphs and burnt pancake graphs, introduced by Gates and Papadimitriou [12], are two well-studied interconnection networks such as ring embedding, super connectivity, broadcasting, fault-tolerant Hamiltonicity.

4.5.1 Pancake graphs

Definition 4.32 [12] The pancake graph of dimension n , denoted by PG_n , has the set of all $n!$ permutations on $\{1, 2, 3, \dots, n\}$ as its vertex set. Two vertices $a_1a_2 \dots a_n$ and $b_1b_2 \dots b_n$ are adjacent if there exists an integer k with $2 \leq k \leq n$ such that $a_i = b_{k+1-i}$ for every i with $1 \leq i \leq n$, and $a_i = b_i$ for $k+1 \leq i \leq n$.

Note that PG_3 is a 6-cycle and PG_4 is given in Fig. 9. PG_n is $(n-1)$ -regular with $n!$ vertices and vertex transitive. Let PG_{n-1}^i be the subgraph of PG_n induced by the vertices with i in the n -th position where $1 \leq i \leq n$. Then PG_n can be decomposed into n vertex disjoint such subgraphs, each of which is isomorphic to PG_{n-1} (see Fig. 9, each red cycle represents PG_3^i for $1 \leq i \leq 4$).

Lemma 4.33 $\alpha(PG_4) = 10$.




Fig. 9 The pancake graph PG_4

Proof. See Appendix. ■

Lemma 4.34 $\alpha(PG_n) \leq \frac{5}{12}n!$ for $n \geq 4$.

Proof. The conclusion holds for $n = 4$ as Lemma 4.33. By induction, suppose that $\alpha(PG_{n-1}) \leq \frac{5}{12}(n-1)!$ for $n \geq 5$. Note that PG_n can be decomposed into n vertex disjoint subgraphs each of which is isomorphic to PG_{n-1} , and thus every independent set of PG_n contains at most $n \times \frac{5}{12}(n-1)! = \frac{5}{12}n!$ vertices. Then $\alpha(PG_n) \leq \frac{5}{12}n!$ for $n \geq 4$. ■

Hung *et al.* [18] considered Hamiltonian properties in faulty pancake graphs.

Lemma 4.35 [18] PG_n is $(n-3)$ -fault Hamiltonian for $n \geq 4$.

By Lemmas 4.34, 4.35 and Theorem 3.5(i), we can determine the FMP number and FSMP number of PG_n , which was also obtained in [24].

Theorem 4.36 $fsm(p)(PG_n) = fmp(PG_n) = n - 1$ for $n \geq 4$.

4.5.2 Burnt pancake graphs

We say the list $a_1a_2 \dots a_n$ is a signed permutation on $\{1, 2, 3, \dots, n\}$ if $|a_1||a_2| \dots |a_n|$ is a permutation on $\{1, 2, 3, \dots, n\}$. For notational simplicity, we use the notation \bar{a} instead of $-a$ and $[n]$ instead of $\{1, 2, 3, \dots, n\} \cup \{\bar{1}, \bar{2}, \bar{3}, \dots, \bar{n}\}$,

Definition 4.37 [12] The burnt pancake graph of n -dimension, denoted by BP_n , has the set of signed permutations on $\{1, 2, 3, \dots, n\}$ as its vertex set. Two vertices $a_1a_2 \dots a_n$ and

$b_1 b_2 \dots b_n$ are adjacent if there exists an integer k with $1 \leq k \leq n$ such that $a_i = b_{\overline{k+1-i}}$ for every i with $1 \leq i \leq k$, and $a_i = b_i$ for $k+1 \leq i \leq n$.




Fig. 10 the burnt pancake graph BP_3

Note that BP_2 is an 8-cycle and BP_3 is given in Fig. 10. BP_n is n -regular with $n!2^n$ vertices and vertex transitive. Let BP_{n-1}^i be the subgraph of BP_n induced by the vertices with i in the n -th position where $i \in [n]$. Then BP_n can be decomposed into $2n$ vertex disjoint such subgraphs, each of which is isomorphic to BP_{n-1} (see Fig. 10, each red even cycle represents BP_2^i for $i \in [3]$).

Lemma 4.38 $\alpha(BP_3) = 20$.

Proof. See Appendix. ■

Lemma 4.39 $\alpha(BP_n) \leq \frac{5}{12}n!2^n$ for $n \geq 3$.

Proof. The conclusion holds for $n = 3$ as Lemma 4.38. By induction, suppose that $\alpha(BP_{n-1}) \leq \frac{5}{12}(n-1)!2^{n-1}$ for $n \geq 4$. Note that BP_n can be decomposed into $2n$ vertex disjoint subgraphs, each of which is isomorphic to BP_{n-1} , and thus every independent set of BP_n contains at most $2n \times \frac{5}{12}(n-1)!2^{n-1} = \frac{5}{12}n!2^n$ vertices. Then $\alpha(BP_n) \leq \frac{5}{12}n!2^n$ for $n \geq 3$. ■

Kaneko [19] considered Hamiltonian properties in faulty burnt pancake graphs.

Lemma 4.40 [19] BP_n is $(n - 2)$ -fault Hamiltonian for $n \geq 3$.

By Lemmas 4.39, 4.40 and Theorem 3.5(i), we can determine the FMP number and FSMP number of BP_n , which was also obtained in [24].

Theorem 4.41 $fsm p(BP_n) = fmp(BP_n) = n$ for $n \geq 3$.

5 Conclusions

In this paper, we establish the FMP number and FSMP number of fault Hamiltonian graphs. Let G be a $(\delta - 2)$ -fault Hamiltonian graph with minimum degree $\delta \geq 3$. If $\alpha(G) \leq \lceil \frac{|G|+1}{2} \rceil - \delta$, then $fsm p(G) = fmp(G) = \delta$; If $G \in \mathcal{G}_1(\delta)$ and $\alpha(G) \leq \lceil \frac{|G|+1}{2} \rceil - \delta + 1$, then $fsm p(G) = fmp(G) = \delta$; If $G \in \mathcal{G}_2(4)$ and $\alpha(G) \leq \frac{|G|+1}{2} - 2$, then $fsm p(G) = fmp(G) = 4$. As applications, the FMP number and FSMP number of some well-known networks, such as the restricted HL-graph G^n , the n -dimensional torus T_{k_1, \dots, k_n} , the recursive circulant graphs $G(d^n, d)$ and $G(2^n, 4)$, the (n, k) -arrangement graph $A_{n,k}$, the (n, k) -star graph $S_{n,k}$, the pancake graph PG_n and the burnt pancake graph BP_n , are determined (see Table 1).

| Networks | the FMP number | the FSMP number |
|--|--------------------------------|--|
| $G^n (n \geq 5)$ | n | n |
| Bipartite $T_{k_1, \dots, k_n} (n \geq 2, k_i \geq 3)$ | $2n$ | 1 |
| Non-bipartite $T_{k_1, \dots, k_n} (n \geq 2, k_i \geq 3)$ | $2n$ | 3, if $(n, k_2) = (2, 4)$ |
| | | $2n$, if $(n, k_2) \neq (2, 4)$ |
| $G(d^n, d) (n \geq 2, d \geq 3)$ | 3, if $(n, d) = (2, 3)$ | 3, if $(n, d) \in \{(2, 3), (2, 4)\}$ |
| | $2n$, if $(n, d) \neq (2, 3)$ | $2n$, if $(n, d) \notin \{(2, 3), (2, 4)\}$ |
| $G(2^n, 4) (n \geq 3)$ | n | $n - 1$, if $n \in \{3, 4\}$ |
| | | n , if $n \notin \{3, 4\}$ |
| $A_{n,k} (2 \leq k \leq n - 2)$ | $k(n - k)$ | $k(n - k)$ |
| $S_{n,k} (2 \leq k \leq n - 2)$ | $n - 1$ | $n - 1$ |
| $PG_n (n \geq 4)$ | $n - 1$ | $n - 1$ |
| $BP_n (n \geq 3)$ | n | n |

Table 1: The FMP number and FSMP number of some well-known networks

References

- [1] R.E.L. Aldred, R.P. Anstee and S.C. Locke, Perfect matchings after vertex deletions, *Discrete Math.*, 307(2007) 3048-3054.
- [2] J.A. Bondy and U.S.R. Murty, Graph Theory, *Springer*, 2008.
- [3] R.C. Brigham, F. Harary, E.C. Violin and J. Yellen, Perfect-matching preclusion, *Congr Numer.*, 174(2005) 185-192.
- [4] F.B. Chedid, On the generalized twisted cube, *Inform. Proc. Lett.*, 55(1995) 49-52.
- [5] E. Cheng and L. Liptak, Matching preclusion and conditional matching preclusion problems for tori and related cartesian products, *Discrete Appl. Math.*, 160(12)(2012) 1699-1716.
- [6] W.K. Chiang and R.J. Chen, The (n, k) -star graph: a generalized star graph, *Inform. Proc. Lett.*, 56(1995) 259-264.
- [7] P. Cull and S. Larson, The Möbius cubes, in: *Proc. of the 6th IEEE Distributed Memory Computing Conf.*, 1991, pp. 699-702.
- [8] K. Day and A. Tripathi, Arrangement graphs: a class of generalized star graphs, *Inform. Proc. Lett.*, 42(1992) 235-241.
- [9] K. Efe, A variation on the hypercube with lower diameter, *IEEE Trans. Comput.*, 40(1991) 1312-1316.
- [10] K. Efe, The crossed cube architecture for parallel computation, *IEEE Trans. Parallel Distributed Syst.*, 3(1992) 513-524.
- [11] T. Gallai, Über extreme Punkt- und Kantenmengen, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, 2(1959) 133-138.
- [12] W.H. Gates and C.H. Papadimitriou, Bounds for sorting by prefix reversal, *Discrete Math.*, 27(1979) 47-57.
- [13] D.R. Guichard, Perfect matchings in pruned grid graphs, *Discrete Math.*, 308(2008) 6552-6557.
- [14] P.A.J. Hilbers, M.R.J. Koopman and J.L.A. van de Snepscheut, The Twisted cube, *PARLE: Parallel Architectures and Languages Europe Volume 1: Parallel Architectures*, Springer, 1987, pp. 152-159.

- [15] H.C. Hsu, Y.L. Hsieh, J.J.M. Tan and L.H. Hsu, Fault Hamiltonicity and fault Hamiltonian connectivity of the (n, k) -star graphs, *Networks*, 42(2003) 189-201.
- [16] H.C. Hsu, T.K. Li, J. Tan and L.H. Hsu, Fault Hamiltonicity and fault Hamiltonian connectivity of the arrangement graphs, *IEEE Trans. Comput.*, 53(2004) 39-53.
- [17] X.L. Hu and H.Q. Liu, The (conditional) matching preclusion for burnt pancake graphs, *Discrete Appl. Math.*, 161(2013) 1481-1489.
- [18] C.N. Hung, H.C. Hsu, K.Y. Liang and L.H. Hsu, Ring embedding in faulty pancake graphs, *Inform. Proc. Lett.*, 86(2003) 271-275.
- [19] K. Kaneko, Hamiltonian cycles and Hamiltonian paths in faulty burnt pancake graphs, *IEICE-Trans. Inform. Sys.*, 90(2007) 716-721.
- [20] H.C. Kim and J.H. Park, Paths and cycles in d -dimensional tori with faults, in: *Proc. of Workshop on Algorithms and Computation, WAAC01, Pusan, Korea*, June 2001, pp. 67-74.
- [21] Q.L. Li, J.H. He and H.P. Zhang, Matching preclusion for vertex-transitive networks, *Discrete Appl. Math.*, 207(2016) 90-98.
- [22] Q.L. Li, W.C. Shiu and H.Y. Yao, Matching preclusion for cube-connected cycles, *Discrete Appl. Math.*, 190-191(2015) 118-126.
- [23] Y. Liu and W.W. Liu, Fractional matching preclusion of graphs, *J. Comb. Optim.*, 34(2017) 522-533.
- [24] T.L. Ma, Y.P. Mao, E. Cheng and C. Melekian, Fractional matching preclusion for (burnt) pancake graphs. *15th International Symposium on Pervasive Systems, Algorithms and Networks (I-SPAN)*, (2018) 133-141.
- [25] T.L. Ma, Y.P. Mao, E. Cheng and J.L. Wang, Fractional matching preclusion for arrangement graphs, *Discrete Appl. Math.*, <https://doi.org/10.1016/j.dam.2019.07.014>.
- [26] Y.P. Mao, Z. Wang, E. Cheng and C. Melekian, Strong matching preclusion number of graphs, *Theoret. Comput. Sci.*, 713(2018) 11-20.
- [27] J.H. Park and K.Y. Chwa, Recursive circulant: a new topology for multicomputer networks, In *Proc. Internat. Symp. Parallel Architectures, Algorithms and Networks (ISPAN)*, (1994) 73-80.

- [28] J.H. Park and K.Y. Chwa, Recursive circulants and their embeddings among hypercubes, *Theoret. Comput. Sci.*, 244(2000) 35-62.
- [29] J.H. Park, H.C. Kim and H.S. Lim, Fault-Hamiltonicity of hypercube-like interconnection networks, in: *Proc. IEEE International Parallel and Distributed Processing Symposium IPDPS 2005*, Denver, April 2005.
- [30] J.H. Park and I.S. Ihm, Strong matching preclusion, *Theoret. Comput. Sci.*, 412(2011) 6409-6419.
- [31] E.R. Scheinerman and D.H. Ullman, Fractional graph theory: a rational approach to the theory of graphs, *New York: Wiley*, 1997.
- [32] C.H. Tsai, J.J.M. Tan, Y.C. Chuang and L.H. Hsu, Hamiltonian properties of faulty recursive circulant graphs, *J. Interconnection Networks*, 3(2002) 273-289.
- [33] A.S. Vaidya, P.S.N. Rao and S.R. Shankar, A class of hypercube-like networks, in: *Proc. of the 5th IEEE Symposium on Parallel and Distributed Processing SPDP 1993*, December 1993, pp. 800-803.
- [34] S.Y. Wang, R.X. Wang, S.W. Lin and J. Li, Matching preclusion for k -ary n -cubes, *Discrete Appl. Math.*, 158(2010) 2006-2070.
- [35] X. Yang, D.J. Evans and G.M. Megson, The locally twisted cubes, *Int. J. Comput. Math.*, 82(2005) 401-413.
- [36] S.Z. Zhang, H.Q. Liu, D. Li and X.L. Hu, Fractional matching preclusion of the restricted HL-graphs, *J. Comb. Optim.*, <https://doi.org/10.1007/s10878-019-00441-x>.
- [37] X.D. Zhu, A hypercube variant with small diameter, *J. Graph Theory*, 85(2017) 651-660.

Appendix

Proof of Lemma 4.14. First we show that $\alpha(G(d^2, d)) \geq \frac{d^2-d}{2}$. Denote $I = I_1 \cup I_2$ if d is odd, where $I_1 = \{v_{2i,2j} : 0 \leq i \leq \frac{d-3}{2} \text{ and } 0 \leq j \leq \frac{d-1}{2}\}$ and $I_2 = \{v_{2i-1,2j-1} : 1 \leq i, j \leq \frac{d-1}{2}\}$. Otherwise, $I = I'_1 \cup I'_2$ if d is even, where $I'_1 = \{v_{2i,2j} : 0 \leq i, j \leq \frac{d-2}{2}\}$ and $I'_2 = \{v_{2i-1,2j-1} : 1 \leq i \leq \frac{d}{2} \text{ and } 1 \leq j \leq \frac{d-2}{2}\}$. It follows that I is an independent set, and thus $\alpha(G(d^2, d)) \geq |I| = \frac{d^2-d}{2}$.

Next we prove $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2}$. Recall that $G_{d,i}^1$ is isomorphic to $G(d, d)$ which is a cycle with d vertices for $0 \leq i < d$. If d is odd, then $G(d^2, d)$ has a spanning subgraph




Fig. 11 $G(d^2, d)$ can be decomposed into $d - 1$ vertex disjoint $(d + 1)$ -cycles and a vertex $v_{d-1,d-1}$.

consisting of d vertex disjoint odd cycles $G_{d,0}^1, \dots, G_{d,d-1}^1$. Note that every independent set of the odd cycle $G_{d,i}^1$ contains at most $\frac{d-1}{2}$ vertices for $0 \leq i < d$, then every independent set of $G(d^2, d)$ contains at most $d \times \frac{d-1}{2} = \frac{d^2-d}{2}$ vertices. Thus $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2}$.

If d is even, then we denote $C_k = v_{k-1,k-1} \dots v_{k-1,d-1} v_{k,0} \dots v_{k,k-1} v_{k-1,k-1}$ for $1 \leq k < d$, and thus C_k is a $(d + 1)$ -cycle. It follows that $G(d^2, d)$ has a spanning subgraph consisting of $d - 1$ vertex disjoint odd cycles C_1, \dots, C_{d-1} and a vertex $v_{d-1,d-1}$ (see Fig. 11, each red cycle represents odd cycle C_k for $1 \leq k < d$). Since every independent set of the odd cycle C_k for $1 \leq k < d$ contains at most $\frac{d}{2}$ vertices, every independent set of $G(d^2, d)$ contains at most $(d - 1) \times \frac{d}{2} + 1 = \frac{d^2-d}{2} + 1$ vertices. Thus $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2} + 1$. Now, suppose that $\alpha(G(d^2, d)) = \frac{d^2-d}{2} + 1$. Then there exists an independent set I' of $G(d^2, d)$ such that $|I'| = \frac{d^2-d}{2} + 1$. This implies that $v_{d-1,d-1} \in I'$ and $|V(C_k) \cap I'| = \frac{d}{2}$ for $1 \leq k < d$. Denote $P_1 := C_1 - \{v_{0,0}, v_{0,d-1}, v_{1,0}\}$. Since P_1 is an even path with order $d - 2$, we have $|V(P_1) \cap I'| \leq \frac{d}{2} - 1$. This together with $|V(C_1) \cap I'| = \frac{d}{2}$, we can deduce that $|\{v_{0,0}, v_{0,d-1}, v_{1,0}\} \cap I'| \geq 1$. Note that $v_{0,0}v_{d-1,d-1}, v_{0,d-1}v_{d-1,d-1} \in E(G(d^2, d))$ and $v_{d-1,d-1} \in I'$. This implies $v_{1,0} \in I'$. By the similar argument above, we have $v_{l,l-1} \in I'$ for $1 \leq l \leq d - 2$ and $v_{d-2,d-2} \in I'$ (see Fig. 11, the green dots represent the vertices in I'). But $v_{d-2,d-3}v_{d-2,d-2} \in E(G(d^2, d))$, a contradiction. Then $\alpha(G(d^2, d)) \neq \frac{d^2-d}{2} + 1$, and thus $\alpha(G(d^2, d)) \leq \frac{d^2-d}{2}$. ■

Proof of Lemma 4.33. First we show that $\alpha(PG_4) \geq 10$. Denote $I = \{1234, 3124, 3412, 1342, 4132, 2413, 4123, 4231, 2341, 3421\}$ (see Fig. 9, the green dots represent the vertices of

I). It follows that I is an independent set of PG_4 , and thus $\alpha(PG_4) \geq |I| = 10$.




Fig. 12 PG_4 can be decomposed into four vertex disjoint 6-cycles.

Next we prove $\alpha(PG_4) \leq 10$. Recall that PG_4 can be decomposed into four vertex disjoint $PG_3^1, PG_3^2, PG_3^3, PG_3^4$, each of which is isomorphic to a 6-cycle (see Fig. 12, each red cycle represents PG_3^i for $1 \leq i \leq 4$). Thus $\alpha(PG_3^i) = 3$ for $1 \leq i \leq 4$. Let I' be a maximum independent set of PG_4 . Suppose that $\alpha(PG_4) \geq 11$. Then there exists at least three elements of $\{1, 2, 3, 4\}$ such that $|I' \cap V(PG_3^i)| = 3$ for $1 \leq i \leq 4$. By symmetry, assume that $|I' \cap V(PG_3^i)| = 3$ for $1 \leq i \leq 3$. Then $I' \cap V(PG_3^1) = \{4231, 2341, 3421\}$ or $\{3241, 2431, 4321\}$. Without loss of generality, assume that $I' \cap V(PG_3^1) = \{4231, 2341, 3421\}$. Note that 2341 and 3421 are adjacent to 1432 and 1243, respectively. Thus $I' \cap V(PG_3^2) = \{3412, 1342, 4132\}$ and $I' \cap V(PG_3^3) = \{1423, 4213, 2143\}$ (see Fig. 12, the green dots represent the vertices in I'). But 3412 is adjacent to 2143 in PG_4 , a contradiction. Then $\alpha(PG_4) \leq 10$.

Proof of Lemma 4.38. First we show that $\alpha(BP_3) \geq 20$. Denote $I = \{2\bar{1}\bar{3}, \bar{2}1\bar{3}, 3\bar{1}2, \bar{1}32, \bar{1}\bar{3}2, \bar{3}\bar{1}2, \bar{2}\bar{3}\bar{1}, \bar{3}\bar{2}\bar{1}, 23\bar{1}, \bar{3}\bar{2}\bar{1}, 123, 213, \bar{1}\bar{3}\bar{2}, \bar{3}\bar{1}\bar{2}, 13\bar{2}, \bar{3}\bar{1}\bar{2}, 32\bar{1}, \bar{2}\bar{3}\bar{1}, \bar{2}\bar{3}\bar{1}\}$ (see Fig. 10, the green dots represent the vertices of I). It follows that I is an independent set of BP_3 , and thus $\alpha(BP_3) \geq |I| = 20$.

Next we prove $\alpha(BP_3) \leq 20$. Recall that BP_3 can be decomposed into six vertex disjoint BP_2^i with $i \in [3]$, each of which is isomorphic to an 8-cycle (see Fig. 13, each red cycle represents PG_3^i for $i \in [3]$). Thus $\alpha(BP_2^i) = 4$ for $i \in [3]$. Let I' be a maximum independent set of BP_3 . Suppose that $\alpha(BP_3) \geq 21$. Then there exists at least three elements of $[3]$ such that $|I' \cap V(BP_2^i)| = 4$ for $i \in [3]$. Without loss of generality, assume that $|I' \cap V(BP_2^1)| = 4$ and $I' \cap V(BP_2^1) = \{321, \bar{2}\bar{3}\bar{1}, \bar{3}\bar{2}\bar{1}, \bar{2}\bar{3}\bar{1}\}$. Now, we consider the




Fig. 13 BP_3 can be decomposed into six vertex disjoint 8-cycles.

following two cases.

Case 1. $|I' \cap V(BP_2^i)| = |I' \cap V(BP_2^{\bar{i}})| = 4$ for some $i \in \{1, 2, 3\}$.

By symmetry, assume that $|I' \cap V(BP_2^1)| = |I' \cap V(BP_2^{\bar{1}})| = 4$. Recall that there exists at least three elements of $[3]$ such that $|I' \cap V(BP_2^i)| = 4$ for $i \in [3]$. Without loss of generality, assume that $|I' \cap V(BP_2^{\bar{2}})| = 4$. Since $2\bar{3}1$ is adjacent to $\bar{1}3\bar{2}$, $I' \cap V(BP_2^{\bar{2}}) = \{3\bar{1}\bar{2}, 13\bar{2}, 3\bar{1}2, \bar{1}3\bar{2}\}$. Note that $13\bar{2}$ is adjacent to $2\bar{3}1$ and $|I' \cap V(BP_2^{\bar{2}})| = 4$, then $I' \cap V(BP_2^{\bar{1}}) = \{2\bar{3}1, 3\bar{2}1, 23\bar{1}, \bar{3}2\bar{1}\}$. This implies $\bar{1}2\bar{3}, 21\bar{3}, 12\bar{3} \notin I' \cap V(BP_2^{\bar{3}})$, $\bar{1}3\bar{2}, 13\bar{2} \notin I' \cap V(BP_2^2)$ and $\bar{1}2\bar{3}, 2\bar{1}3, \bar{1}23 \notin I' \cap V(BP_2^3)$ (see Fig. 13, the green dots represent the vertices in I' and the black crosses represent the vertices out of I'). Thus $|I' \cap V(BP_2^{\bar{3}})| \leq 3$ and $|I' \cap V(BP_2^3)| \leq 3$. Since $\alpha(BP_3) = |I'| \geq 21$, we have $3 \leq |I' \cap V(BP_2^2)| \leq 4$.

Suppose that $|I' \cap V(BP_2^2)| = 4$. Then $I' \cap V(BP_2^2) = \{312, \bar{1}3\bar{2}, \bar{3}1\bar{2}, 1\bar{3}2\}$. This implies $\bar{2}1\bar{3} \notin I' \cap V(BP_2^{\bar{3}})$ and $123 \notin I' \cap V(BP_2^3)$. Hence, $|I' \cap V(BP_2^{\bar{3}})| \leq 2$ and $|I' \cap V(BP_2^3)| \leq 2$. Then $|I'| \leq 4 \times 4 + 2 \times 2 = 20$, a contradiction. Thus $|I' \cap V(BP_2^2)| = 3$. Recall that $|I' \cap V(BP_2^{\bar{3}})| \leq 3$ and $|I' \cap V(BP_2^3)| \leq 3$. Then $|I' \cap V(BP_2^{\bar{3}})| = |I' \cap V(BP_2^3)| = 3$ as $\alpha(BP_3) \geq 21$. It follows that $\bar{2}1\bar{3} \in I' \cap V(BP_2^{\bar{3}})$ and $\bar{2}13 \in I' \cap V(BP_2^3)$, and thus $312, \bar{3}1\bar{2} \notin I' \cap V(BP_2^2)$. But $|I' \cap V(BP_2^2)| \leq 2$, a contradiction.

Case 2. $|I' \cap V(BP_2^i)| \leq 3$ or $|I' \cap V(BP_2^{\bar{i}})| \leq 3$ for any $i \in \{1, 2, 3\}$.

Recall that there exists at least three elements of $[3]$ such that $|I' \cap V(BP_2^i)| = 4$ for $i \in [3]$. Suppose that $|I' \cap V(BP_2^{\bar{2}})| = 4$ or $|I' \cap V(BP_2^{\bar{3}})| = 4$. Without loss of generality, assume that $|I' \cap V(BP_2^{\bar{2}})| = 4$. Since $2\bar{3}1$ is adjacent to $\bar{1}3\bar{2}$, $I' \cap V(BP_2^{\bar{2}}) = \{\bar{3}1\bar{2}, 13\bar{2}, 3\bar{1}\bar{2}, \bar{1}3\bar{2}\}$. Note that 321 and $3\bar{1}\bar{2}$ are adjacent to $\bar{1}2\bar{3}$ and $21\bar{3}$. Thus $|I' \cap V(BP_2^{\bar{3}})| \leq 3$. This implies $|I' \cap V(BP_2^3)| = 4$. Since $\bar{3}1\bar{2}$ is adjacent to $2\bar{1}3$, $I' \cap V(BP_2^3) = \{\bar{1}2\bar{3}, \bar{2}1\bar{3}, \bar{1}23, 213\}$. But $\bar{3}21$ is adjacent to $\bar{1}23$ in BP_3 , a contradiction. Then $|I' \cap V(BP_2^{\bar{2}})| \leq 3$ and $|I' \cap V(BP_2^{\bar{3}})| \leq 3$. Hence, $|I' \cap V(BP_2^2)| = |I' \cap V(BP_2^3)| = 4$. Since $2\bar{3}1$ is adjacent to $\bar{1}3\bar{2}$, $I' \cap V(BP_2^2) = \{312, \bar{1}32, \bar{3}12, \bar{1}3\bar{2}\}$. Since $\bar{3}21$ is adjacent to $\bar{1}23$, $I' \cap V(BP_2^3) = \{2\bar{1}3, 123, \bar{2}13, \bar{1}2\bar{3}\}$. But $\bar{2}13$ is adjacent to $\bar{3}12$ in BP_3 , a contradiction.