arXiv:2001.03729v1 [math.AG] 11 Jan 2020

EFFECTIVE CONES OF CYCLES ON PRODUCTS OF PROJECTIVE BUNDLES OVER CURVES

RUPAM KARMAKAR

ABSTRACT. Let $X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$ where C is a smooth curve and E_1, E_2 are vector bundles over C.In this paper we compute the pseudo effective cones of higher codimension cycles on X.

1. Introduction

The cones of divisors and curves on projective varieties have been extensively studied over the years and by now are quite well understood. However, more recently the theory of cones of cycles of higher dimension has been the subject of increasing interest(see [F], [DELV], [DJV], [CC] etc). Lately, there has been significant progress in the theoretical understanding of such cycles, due to [FL1], [FL2] and others. But the the number of examples where the cone of effective cycles have been explicitly computed is relatively small till date [F], [CLO], [PP] etc.

Let E_1 and E_2 be two vector bundles over a smooth curve C and consider the fibre product $X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$. Motivated by the results in [F], in this paper, we compute the cones of effective cycles on X in the following cases.

Case I: When both E_1 and E_2 are semistable vector bundles of rank r_1 and r_2 respectively, the cone of effective codimension k-cycles are described in theorem 3.2.

Case II: When Neither E_1 nor E_2 is semistable, the cone of low dimension effective cycles are computed in theorem 3.3 and the remaining cases in therem 3.5.

2. Preliminaries

Let X be a smooth projective varity of dimension n. $N_k(X)$ is the real vector space of k-cycles on X modulo numerical equivalence. For each k, $N_k(X)$ is a real vector space of finite dimension. Since X is smooth, we can identify $N_k(X)$ with the abstract dual $N^{n-k}(X) := N_{n-k}(X)^\vee$ via the intersection pairing $N_k(X) \times N_{n-k}(X) \longrightarrow \mathbb{R}$.

For any k-dimensional subvariety Y of X, let $[Y]$ be its class in $N_k(X)$. A class $\alpha \in N_k(X)$ is said to be effective if there exists subvarities $Y_1, Y_2, ..., Y_m$ and non-negetive real numbers $n_1, n_2, ..., n_m$ such that α can be written as $\alpha = \sum n_i Y_i$. The *pseudo-effective cone* $\overline{\text{Eff}}_k(X) \subset N_k(X)$ is the closure of the cone generated by classes of effective cycles. It is full-dimensional and does not contain any nonzero linear subspaces. The pseudo-effective dual classes form a closed cone in $N^k(X)$ which we denote by $\overline{\text{Eff}}^k(X)$.

2010 *Mathematics Subject Classification.* Primary 14J60; Secondary 14H60, 14J10.

For smooth varities Y and Z, a map $f : N^k(Y) \longrightarrow N^K(Z)$ is called pseudo-effective if $f(\overline{\text{Eff}}^k(Y)) \subset \overline{\text{Eff}}^k(Z)$.

The *nef cone* $\text{Nef}^k(X) \subset N^k(X)$ is the dual of $\overline{\text{Eff}}^k(X) \subset N^k(X)$ via the pairing $N^k(X) \times N_k(X) \longrightarrow \mathbb{R}$, i.e,

$$\text{Nef}^k(X) := \left\{\alpha \in N_k(X) | \alpha \cdot \beta \geq 0 \forall \beta \in \overline{\text{Eff}}_k(X)\right\}$$

3. Cone of effective cycles

Let E_1 and E_2 be two vector bundles over a smooth curve C of rank r_1, r_2 and degrees d_1, d_2 respectively. Let $\mathbb{P}(E_1) = \textbf{Proj}\ (\oplus_{d\geq 0} Sym^d(E_1))$ and $\mathbb{P}(E_2) = \textbf{Proj}\ (\oplus_{d\geq 0} Sym^d(E_2))$ be the associated projective bundle together with the projection morphisms $\pi_1 : \mathbb{P}(E_1) \longrightarrow C$ and $\pi_2 : \mathbb{P}(E_2) \longrightarrow C$ respectively. Let $X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$ be the fibre product over C. Consider the following commutative diagram:

$$\begin{array}{ccc} X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2) & \xrightarrow{p_2} & \mathbb{P}(E_2) \\ \downarrow{\scriptstyle p_1} & & \downarrow{\scriptstyle \pi_2} \\ \mathbb{P}(E_1) & \xrightarrow{\pi_1} & C \end{array} \tag{1}$$

Let f_1, f_2 and F denote the numerical equivalence classes of the fibres of the maps π_1, π_2 and $\pi_1 \circ p_1 = \pi_2 \circ p_2$ respectively. Note that, $X \cong \mathbb{P}(\pi_1^*(E_2)) \cong \mathbb{P}(\pi_2^*(E_1))$. We first fix the following notations for the numerical equivalence classes,

$$\eta_1 = [\mathcal{O}_{\mathbb{P}(E_1)}(1)] \in N^1(\mathbb{P}(E_1)) \quad , \quad \eta_2 = [\mathcal{O}_{\mathbb{P}(E_2)}(1)] \in N^1(\mathbb{P}(E_2)),$$
$$\xi_1 = [\mathcal{O}_{\mathbb{P}(\pi_1^*(E_2))}(1)] \quad , \quad \xi_2 = [\mathcal{O}_{\mathbb{P}(\pi_2^*(E_1))}(1)] \quad , \quad \zeta_1 = p_1^*(\eta_1) \quad , \quad \zeta_2 = p_2^*(\eta_2)$$
$$\zeta_1 = \xi_2,\ \zeta_2 = \xi_1 \quad , \quad F = p_1^*(f_1) = p_2^*(f_2)$$

We here summarise some results that has been discussed in [KMR] (See section 3 in [KMR] for more details) :

$$N^1(X)_{\mathbb{R}} = \mathbb{R}\zeta_1 \oplus \mathbb{R}\zeta_2 \oplus \mathbb{R}F,$$
$$\zeta_1^{r_1} \cdot F = 0\ ,\ \zeta_1^{r_1+1} = 0\ ,\ \zeta_2^{r_2} \cdot F = 0\ ,\ \zeta_2^{r_2+1} = 0\ ,\ F^2 = 0\ ,$$
$$\zeta_1^{r_1} = (\deg(E_1))F \cdot \zeta_1^{r_1-1}\ ,\ \zeta_2^{r_2} = (\deg(E_2))F \cdot \zeta_2^{r_2-1} \quad ,$$
$$\zeta_1^{r_1} \cdot \zeta_2^{r_2-1} = \deg(E_1) \quad , \quad \zeta_2^{r_2} \cdot \zeta_1^{r_1-1} = \deg(E_2).$$

Also, The dual basis of $N_1(X)_{\mathbb{R}}$ is given by $\{\delta_1, \delta_2, \delta_3\}$, where

$$\delta_1 = F \cdot \zeta_1^{r_1-2} \cdot \zeta_2^{r_2-1},$$
$$\delta_2 = F \cdot \zeta_1^{r_1-1} \cdot \zeta_2^{r_2-2},$$
$$\delta_3 = \zeta_1^{r_1-1} \cdot \zeta_2^{r_2-1} - \deg(E_1)F \cdot \zeta_1^{r_1-2} \cdot \zeta_2^{r_2-1} - \deg(E_2)F \cdot \zeta_1^{r_1-1} \cdot \zeta_2^{r_2-2}.$$

Theorem 3.1. *Let $r_1 = \text{rank}(E_1)$ and $r_2 = \text{rank}(E_2)$ and without loss of generality assume that $r_1 \leq r_2$. Then the bases of $N^k(X)$ are given by*

$$N^k(X) = \begin{cases} \left(\{\zeta_1^i \cdot \zeta_2^{k-i}\}_{i=0}^{k}, \{F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}\}_{j=0}^{k-1}\right) & if \quad k < r_1 \\ \left(\{\zeta_1^i \cdot \zeta_2^{k-i}\}_{i=0}^{r_1-1}, \{F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}\}_{j=0}^{r_1-1}\right) & if \quad r_1 \leq k < r_2 \\ \left(\{\zeta_1^i \cdot \zeta_2^{k-i}\}_{i=t+1}^{r_1-1}, \{F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}\}_{j=t}^{r_1-1}\right) & if \quad k = r_2 + t \quad where \quad t \in \{0, 1, 2, ..., r_1 - 2\}. \end{cases}$$

Proof. To begin with consider the case where $k < r_1$. We know that $X \cong \mathbb{P}(\pi_2^* E_1)$ and the natural morphism $\mathbb{P}(\pi_2^* E_1) \longrightarrow \mathbb{P}(E_2)$ can be identified with p_2. With the above identifications in place the chow group of X has the following isomorphism [see theorem 3.3 , page 64 [Ful]]

$$A(X) \cong \bigoplus_{i=0}^{r_1-1} \zeta_1^i A(\mathbb{P}(E_2)) \tag{2}$$

Choose i_1, i_2 suct that $0 \leq i_1 < i_2 \leq k$. Consider the k- cycle $\alpha := F \cdot \zeta_1^{r_1-i_1-1} \cdot \zeta_2^{r_2+i_1-k-1}$.

Then $\zeta_1^{i_1} \cdot \zeta_2^{k-i_1} \cdot \alpha = 1$ but $\zeta_1^{i_2} \cdot \zeta_2^{k-i_2} \cdot \alpha = 0$. So, $\{\zeta_1^{i_1} \cdot \zeta_2^{k-i_1}\}$ and $\{\zeta_1^{i_2} \cdot \zeta_2^{k-i_2}\}$ can not be numerically equivalent.

Similarly, take j_1, j_2 such that $0 \leq j_1 < j_2 \leq k$ and consider the k-cycle $\beta := \zeta_1^{r_1-j_1-1} \cdot \zeta_2^{r_2+j_1-k}$.

Then as before it happens that $F \cdot \zeta_1^{j_1} \cdot \zeta_2^{k-j_1-1} \cdot \beta = 1$ but $F \cdot \zeta_1^{j_2} \cdot \zeta_2^{k-j_2-1} \cdot \beta = 0$. So $\{F \cdot \zeta_1^{j_1} \cdot \zeta_2^{k-j_1-1}\}$ and $\{F \cdot \zeta_1^{j_2} \cdot \zeta_2^{k-j_2-1}\}$ can not be numerically equivalent.

For the remaining case lets assume $0 \leq i \leq j \leq k$ and consider the k-cycle $\gamma := F \cdot \zeta_1^{r_1-i-1} \cdot \zeta_2^{r_2+i-1-k}$.

Then $\{\zeta_1^i \cdot \zeta_2^{k-i}\} \cdot \gamma = 1$ and $\{F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}\} \cdot \gamma = 0$. So, they can not be numerically equivalent. From these observations and (2) we obtain a basis of $N^k(X)$ which is given by

$$N^k(X) = \left(\{\zeta_1^i \cdot \zeta_2^{k-i}\}_{i=0}^{k}, \{F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}\}_{j=0}^{k-1}\right)$$

For the case $r_1 \leq k < r_2$ observe that $\zeta_1^{r_1+1} = 0$, $F \cdot \zeta_1^{r_1} = 0$ and $\zeta_1^{r_2} = deg(E_1) F \cdot \zeta_1^{r_1-1}$.

When $k \geq r_2$ we write as $k = r_2 + t$ where t ranges from 0 to $r_1 - 1$. In that case the observations like $\zeta_2^{r_2+1} = 0$, $F \cdot \zeta_2^{r_2} = 0$ and $\zeta_2^{r_2} = \deg(E_2) F \cdot \zeta_2^{r_2-1}$ proves our case. □

Now we are ready to treat the case where both E_1 and E_2 are semistable vector bundles over C.

Theorem 3.2. *Let E_1 and E_2 be two semistable vector bundles over C of rank r_1 and r_2 respectively with $r_1 \leq r_2$ and $X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$. Then for all $k \in \{1, 2, ..., r_1 + r_2 - 1\}$*

$$\overline{\text{Eff}}^k(X) = \begin{cases} \left\langle \left\{(\zeta_1-\mu_1F)^i(\zeta_2-\mu_2F)^{k-i}\right\}_{i=0}^{k}, \left\{F\cdot\zeta_1^j\cdot\zeta_2^{k-j-1}\right\}_{j=0}^{k-1}\right\rangle & if \quad k<r_1 \\ \left\langle \left\{(\zeta_1-\mu_1F)^i(\zeta_2-\mu_2F)^{k-i}\right\}_{i=0}^{r_1-1}, \left\{F\cdot\zeta_1^j\cdot\zeta_2^{k-j-1}\right\}_{j=0}^{r_1-1}\right\rangle & if \quad r_1\le k<r_2 \\ \left\langle \left\{(\zeta_1-\mu_1F)^i(\zeta_2-\mu_2F)^{k-i}\right\}_{i=t+1}^{r_1-1}, \left\{F\cdot\zeta_1^j\cdot\zeta_2^{k-j-1}\right\}_{j=t}^{r_1-1}\right\rangle & if \quad k=r_2+t, \quad t=0,...,r_1-1. \end{cases}$$

where $\mu_1=\mu(E_1)$ *and* $\mu_2=\mu(E_2)$.

Proof. Firstly, $(\zeta_1-\mu_1F)^i\cdot(\zeta_2-\mu_2F)^{k-i}$ and $F\cdot\zeta_1^i\cdot\zeta_2^{k-j-1}[=F\cdot(\zeta_1-\mu_1F)^i\cdot(\zeta_2-\mu_2F)^{k-j-1}]$ are intersections of nef divisors. So, they are pseudo-effective for all $i\in\{0,1,2,...,k\}$. conversely, when $k<r_1$ notice that we can write any element C of $\overline{\text{Eff}}^k(X)$ as

$$C=\sum_{i=0}^{k}a_i(\zeta_1-\mu_1F)^i\cdot(\zeta_2-\mu_2F)^{k-i}+\sum_{j=0}^{k}b_jF\cdot\zeta_1^j\cdot\zeta_2^{k-j-1}$$

where $a_i,b_i\in\mathbb{R}$.

For a fixed i_1 intersect C with $D_{i_1}:=F\cdot(\zeta_1-\mu_1F)^{r_1-i_1-1}\cdot(\zeta_2-\mu_2F)^{r_2-k+i_1-1}$ and for a fixed j_1 intersect C with $D_{j_1}:=(\zeta_1-\mu_1F)^{r_1-j_1-1}\cdot(\zeta_2-\mu_2F)^{r_2+j_1-k}$. These intersections lead us to

$$C\cdot D_{i_1}=a_{i_1}\quad and\quad C\cdot D_{j_1}=b_{j_1}$$

Since $C\in\overline{\text{Eff}}^k(X)$ and D_{i_1},D_{j_1} are intersection of nef divisors, a_{i_1} and b_{j_1} are non-negetive. Now running i_1 andj_1 through $\{0,1,2,...,k\}$ we get all the a_i's and b_i's are non-negetive and that proves our result for $k<r_1$. The cases where $r_1\le k<r_2$ and $k\ge r_2$ can be proved very similarly after the intersection products involving ζ_1 and ζ_2 in page 2 are taken into count. □

Next we study the more interesting case where E_1 and E_2 are two unstable vector bundles of rank r_1 and r_2 and degree d_1 and d_2 respectively over a smooth curve C.

Let E_1 be the unique Harder-Narasimhan filtration

$$E_1=E_{10}\supset E_{11}\supset...\supset E_{1l_1}=0$$

with $Q_{1i}:=E_{1(i-1)}/E_{1i}$ being semistable for all $i\in[1,l_1-1]$. Denote $n_{1i}=\text{rank}(Q_{1i})$, $d_{1i}=\deg(Q_{1i})$ and $\mu_{1i}=\mu(Q_{1i}):=\frac{d_{1i}}{n_{1i}}$ for all i.

Similarly, E_2 also admits the unique Harder-Narasimhan filtration

$$E_2=E_{20}\supset E_{21}\supset...\supset E_{2l_2}=0$$

with $Q_{2i} := E_{2(i-1)}/E_{2i}$ being semistable for $i \in [1, l_2 - 1]$. Denote $n_{2i} = \text{rank}(Q_{2i})$, $d_{2i} = \deg(Q_{2i})$ and $\mu_{2i} = \mu(Q_{2i}) := \frac{d_{2i}}{n_{2i}}$ for all i.

Consider the natural inclusion $\overline{i} = i_1 \times i_2 : \mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21}) \longrightarrow \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$, which is induced by natural inclusions $i_1 : \mathbb{P}(Q_{11}) \longrightarrow \mathbb{P}(E_1)$ and $i_2 : \mathbb{P}(Q_{21}) \longrightarrow \mathbb{P}(E_2)$. In the next theorem we will see that the cycles of $\mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$ of dimension at most $n_{11} + n_{21} - 1$ can be tied down to cycles of $\mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21})$ via $\overline{i}$.

Theorem 3.3. *Let E_1 and E_2 be two unstable bundle of rank r_1 and r_2 and degree d_1 and d_2 respectively over a smooth curve C and $r_1 \leq r_2$ without loss of generality and $X = \mathbb{P}(E_1) \times_C \mathbb{P}(E_2)$.*

Then for all $k \in \{1, 2, ..., \mathbf{n}\}$ ($\mathbf{n} := n_{11} + n_{21} - 1$)

Case(1): $n_{11} \leq n_{21}$

$$\overline{\text{Eff}}_k(X) = \begin{cases} \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_1 - \mu_{11}F)^i(\zeta_2 - \mu_{21}F)^{\mathbf{n}-k-i} \right\}_{i=t+1}^{n_{11}-1}, \left\{ F \cdot \zeta_1^{r_1-n_{11}+j} \cdot \zeta_2^{r_2+n_{11}-k-j-2} \right\}_{j=t}^{n_{11}-1} \right\rangle \\ \qquad if \quad k < n_{11} \quad and \quad t = 0, 1, 2, ..., n_{11} - 2 \\ \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_1 - \mu_{11}F)^i(\zeta_2 - \mu_{21}F)^{\mathbf{n}-k-i} \right\}_{i=0}^{n_{11}-1}, \left\{ F \cdot \zeta_1^{r_1-n_{11}+j} \cdot \zeta_2^{r_2+n_{11}-k-j-2} \right\}_{j=0}^{n_{11}-1} \right\rangle \\ \qquad if \quad n_{11} \leq k < n_{21}. \\ \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_1 - \mu_{11}F)^i(\zeta_2 - \mu_{21}F)^{\mathbf{n}-k-i} \right\}_{i=0}^{\mathbf{n}-k}, \left\{ F \cdot \zeta_1^{r_1-n_{11}+j} \cdot \zeta_2^{r_2+n_{11}-k-j-2} \right\}_{j=0}^{\mathbf{n}-k} \right\rangle \\ \qquad if \quad k \geq n_{21}. \end{cases}$$

Case(2): $n_{21} \leq n_{11}$

$$\overline{\text{Eff}}_k(X) = \begin{cases} \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_2 - \mu_{21}F)^i(\zeta_1 - \mu_{11}F)^{\mathbf{n}-k-i} \right\}_{i=t+1}^{n_{21}-1}, \left\{ F \cdot \zeta_2^{r_2-n_{21}+j} \cdot \zeta_1^{r_1+n_{21}-k-j-2} \right\}_{j=t}^{n_{21}-1} \right\rangle \\ \qquad if \quad k < n_{21} \quad and \quad t = 0, 1, 2, ..., n_{21} - 2 \\ \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_2 - \mu_{21}F)^i(\zeta_1 - \mu_{11}F)^{\mathbf{n}-k-i} \right\}_{i=0}^{n_{21}-1}, \left\{ F \cdot \zeta_2^{r_2-n_{21}+j} \cdot \zeta_1^{r_1+n_{21}-k-j-2} \right\}_{j=0}^{n_{21}-1} \right\rangle \\ \qquad if \quad n_{21} \leq k < n_{11}. \\ \left\langle \left\{ [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_2 - \mu_{21}F)^i(\zeta_1 - \mu_{11}F)^{\mathbf{n}-k-i} \right\}_{i=0}^{\mathbf{n}-k}, \left\{ F \cdot \zeta_2^{r_2-n_{21}+j} \cdot \zeta_1^{r_1+n_{21}-k-j-2} \right\}_{j=0}^{\mathbf{n}-k} \right\rangle \\ \qquad if \quad k \geq n_{11}. \end{cases}$$

Thus in both cases $\overline{i}_*$ *induces an isomorphism between* $\overline{\text{Eff}}_k([\mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21})])$ *and* $\overline{\text{Eff}}_k(X)$ *for* $k \leq \mathbf{n}$.

Proof. to begin with consider $Case(1)$ and then take $k \geq n_{21}$. Since $(\zeta_1 - \mu_{11}F)$ and $(\zeta_2 - \mu_{21}F)$ are nef

$$\phi_i := [\mathbb{P}(Q_{11}) \times \mathbb{P}(Q_{21})](\zeta_1 - \mu_{11}F)^i(\zeta_2 - \mu_{21}F)^{\mathbf{n}-k-i} \in \overline{\text{Eff}}_k(X).$$

for all $i \in \{0, 1, 2, ..., \mathbf{n} - k\}$.

Now The result in [Example 3.2.17, [Ful]] adjusted to bundles of quotients over curves shows that

$$[\mathbb{P}(Q_{11})] = \eta_1^{r_1-n_{11}} + (d_{11} - d_1)\eta_1^{r_1-n_{11}-1}f_1$$

and

$$[\mathbb{P}(Q_{21})] = \eta_2^{r_2-n_{21}} + (d_{21} - d_2)\eta_2^{r_2-n_{21}-1}f_2$$

Also, $p_1^*[\mathbb{P}(Q_{11})] \cdot p_2^*[\mathbb{P}(Q_{21})] = [\mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21})]$. With little calculations it can be shown that

$\phi_i \cdot (\zeta_1 - \mu_{11}F)^{n_{11}-i} \cdot (\zeta_2 - \mu_{21}F)^{k+i+1-n_{11}}$
$= (\zeta_1^{r_1-n_{11}} + (d_{11} - d_1)F \cdot \zeta_1^{r_1-n_{11}-1})(\zeta_2^{r_2-n_{21}} + (d_{21} - d_2)F \cdot \zeta_2^{r_2-n_{21}-1})(\zeta_1 - \mu_{11}F)^{n_1 1-i} \cdot (\zeta_2 - \mu_{21}F)^{k+i+1-n_{11}}$
$= (\zeta_1^{r_1} - d_1F \cdot \zeta_1^{r_1-1})(\zeta_2^{r_2} - d_2F \cdot \zeta_2^{r_2-1}) = 0.$

So, ϕ_i 's are in the boundary of $\overline{\text{Eff}}_k(X)$ for all $i \in \{0, 1, ..., \mathbf{n}-k\}$. The fact that $F \cdot \zeta_1^{r_1-n_{11}+j} \cdot \zeta_2^{r_2+n_{11}-k-j-2}$'s are in the boundary of $\overline{\text{Eff}}_k(X)$ for all $i \in \{0, 1, ..., \mathbf{n} - k\}$ can be deduced from the proof of Theorem 2.2. The other cases can be proved similarly.

The proof of $Case(2)$ is similar to the proof of $Case(1)$.

Now, to show the isomorphism between pseudo-effective cones induced by $\overline{i}_*$ observe that Q_{11} and Q_{21} are semi-stable bundles over C. So, Theorem 2.2 gives the expressions for $\overline{\text{Eff}}_k([\mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21})])$. Let $\zeta_{11} = \mathcal{O}_{\mathbb{P}(\tilde{\pi}_2^*(Q_{11}))}(1) = \tilde{p}_1^*(\mathcal{O}_{\mathbb{P}(Q_{11})}(1))$ and $\zeta_{21} = \mathcal{O}_{\mathbb{P}(\tilde{\pi}_1^*(Q_{21})}(1) = \tilde{p}_2^*(\mathcal{O}_{\mathbb{P}(Q_{21})}(1))$, where $\tilde{\pi_2} = \pi_2|_{\mathbb{P}(Q_{21})}$, $\tilde{\pi_1} = \pi_1|_{\mathbb{P}(Q_{11})}$ and $\tilde{p}_1 : \mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21}) \longrightarrow \mathbb{P}(Q_{11})$, $\tilde{p}_2 : \mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21}) \longrightarrow \mathbb{P}(Q_{21})$ are the projection maps. Also notice that $\overline{i}^*\zeta_1 = \zeta_{11}$ and $\overline{i}^*\zeta_1 = \zeta_{21}$.

Using the above relations and projection formula the isomorphism between $\overline{\text{Eff}}_k([\mathbb{P}(Q_{11}) \times_C \mathbb{P}(Q_{21})])$ and $\overline{\text{Eff}}_k(X)$ for $k \leq \mathbf{n}$ can be proved easily.

□

Next we want to show that higher dimension pseudo effective cycles on X can be related to the pseudo effective cycles on $\mathbb{P}(E_{11}) \times_C \mathbb{P}(E_{21})$. More precisely there is a isomorphism between $\overline{\text{Eff}}^k(X)$ and $\overline{\text{Eff}}^k([\mathbb{P}(E_{11}) \times_C \mathbb{P}(E_{21})])$ for $k < r_1 + r_2 - 1 - \mathbf{n}$. Useing the coning construction as in [ful] we show this in two steps, first we establish an isomorphism between $\overline{\text{Eff}}^k([\mathbb{P}(E_1) \times_C \mathbb{P}(E_2)])$ and $\overline{\text{Eff}}^k([\mathbb{P}(E_{11}) \times_C \mathbb{P}(E_2)])$ and then an isomorphism between $\overline{\text{Eff}}^k([\mathbb{P}(E_{11}) \times_C \mathbb{P}(E_2)])$ and

$\overline{\text{Eff}}^k([\mathbb{P}(E_{11})\times_C\mathbb{P}(E_{21})])$ in similar fashion. But before proceeding any further we need to explore some more facts.

Let E be an unstable vector bundle over a non-singular projective variety V. There is a unique filtration

$$E = E^0 \supset E^1 \supset E^2 \supset ... \supset E^l = 0$$

which is called the harder-Narasimhan filtration of E with $Q^i := E^{i-1}/E^i$ being semistable for $i \in [1, l-1]$. Now the following short-exact sequence

$$0 \longrightarrow E^1 \longrightarrow E \longrightarrow Q^1 \longrightarrow 0$$

induced by the harder-narasimhan filtration of E gives us the natural inclusion $j : \mathbb{P}(Q^1) \hookrightarrow \mathbb{P}(E)$. Considering $\mathbb{P}(Q^1)$ as a subscheme of $\mathbb{P}(E)$ we obtain the commutative diagram below by blowing up $\mathbb{P}(Q^1)$.

$$\begin{array}{ccc} \tilde{Y} = Bl_{\mathbb{P}(Q^1)}\mathbb{P}(E) & \xrightarrow{\Phi} & \mathbb{P}(E^1) = Z \\ \downarrow{\scriptstyle \Psi} & & \downarrow{\scriptstyle q} \\ Y = \mathbb{P}(E) & \xrightarrow{\quad p \quad} & V \end{array} \tag{3}$$

where Ψ is blow-down map.

Theorem 3.4. *With the above notation,there exists a locally free sheaf G on Z such that $\tilde{Y} \simeq \mathbb{P}_Z(G)$ and $\nu : \mathbb{P}_Z(G) \longrightarrow Z$ it's corresponding bundle map.*

In particular if we place $V = \mathbb{P}(E_2)$, $E = \pi_2^*E_1$, $E^1 = \pi_2^*E_{11}$ and $Q^1 = \pi_2^*Q_{11}$ then the above commutative diagram becomes

$$\begin{array}{ccc} \tilde{Y'} = Bl_{\mathbb{P}(\pi_2^*Q_{11})}\mathbb{P}(\pi_2^*E_1) & \xrightarrow{\Phi'} & \mathbb{P}(\pi_2^*E_{11}) = Z' \\ \downarrow{\scriptstyle \Psi'} & & \downarrow{\scriptstyle \overline{p}_2} \\ Y' = \mathbb{P}(\pi_2^*E_1) & \xrightarrow{\quad p_2 \quad} & \mathbb{P}(E_2) \end{array} \tag{4}$$

where $p_2 : \mathbb{P}(\pi_2^*E_1) \longrightarrow \mathbb{P}(E_2)$ and $\overline{p}_2 : \mathbb{P}(\pi_2^*E_{11}) \longrightarrow \mathbb{P}(E_2)$ are projection maps.

and there exists a locally free sheaf G' on Z' such that $\tilde{Y'} \simeq \mathbb{P}_{Z'}(G')$ and $\nu' : \mathbb{P}_{Z'}(G') \longrightarrow Z'$ it's bundle map.

Now let $\zeta_{Z'} = \mathcal{O}_{Z'}(1)$, $\gamma = \mathcal{O}_{\mathbb{P}_{Z'}(G')}(1)$, F the numerical equivalence class of a fibre of $\pi_2 \circ p_2$, F_1 the numerical equivalence class of a fibre of $\pi_2 \circ \overline{p}_2$, $\tilde{E}$ the class of the exceptional divisor of Ψ' and $\zeta_1 = p_1^*(\eta_1) = \mathcal{O}_{\mathbb{P}(\pi_2^*E_1)}(1)$. Then we have the following relations:

$$\gamma = (\Psi')^*\zeta_1, \quad (\Phi')^*\zeta_{Z'} = (\Psi')^*\zeta_1 - \tilde{E}, \quad (\Phi')^*F_1 = (\Psi')^*F \tag{5}$$

$$\tilde{E} \cdot (\Psi')^*(\zeta_1 - \mu_{11}F)^{n_{11}} = 0 \tag{6}$$


Additionaly, if we also denote the support of the exceptional divisor of $\tilde{Y}'$ by $\tilde{E}$, then $\tilde{E} \cdot N(\tilde{Y}') = (j_{\tilde{E}})_* N(\tilde{E})$, where $j_{\tilde{E}} : \tilde{E} \longrightarrow \tilde{Y}'$ is the canonical inclusion.

Proof. With the above hypothesis the following commutative diagram is formed:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & q^*E^1 & \longrightarrow & q^*E & \longrightarrow & q^*Q^1 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \| & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}(E^1)}(1) & \longrightarrow & G & \longrightarrow & q^*Q^1 & \longrightarrow & 0 \end{array}$$

where G is the push-out of morphisms $q^*E^1 \longrightarrow q^*E$ and $q^*E^1 \longrightarrow \mathcal{O}_{\mathbb{P}(E^1)}(1)$ and the first vertical map is the natural surjection. Now let $W = \mathbb{P}_Z(G)$ and $\nu : W \longrightarrow Z$ be it's bundle map. So there is a cannonical surjection $\nu^*G \longrightarrow \mathcal{O}_{\mathbb{P}_Z(G)}(1)$. Also note that $q^*E \longrightarrow G$ is surjective by snake lemma. Combining these two we obtain a surjective morphism $\nu^*q^*E \longrightarrow \mathcal{O}_{\mathbb{P}_Z(G)}(1)$ which determines $\omega : W \longrightarrow Y$. We claim that we can identify $(\tilde{Y}, \Phi, \Psi)$ and (W, ν, ω). Now Consider the following commutative diagram:

(7)



where $\mathbf{i}$ is induced by the universal property of the fiber product. Since $\mathbf{i}$ can also be obtained from the surjective morphism $q^*E \longrightarrow G$ it is a closed immersion. Let $\mathcal{T}$ be the $\mathcal{O}_Y$ algebra $\mathcal{O}_Y \oplus \mathcal{I} \oplus \mathcal{I}^2 \oplus ...$, where $\mathcal{I}$ is the ideal sheaf of $\mathbb{P}(Q^1)$ in Y. We have an induced map of $\mathcal{O}_Y$- algebras $Sym(p^*E^1) \longrightarrow \mathcal{T} * \mathcal{O}_Y(1)$ which is onto because the image of the composition $p^*E^1 \longrightarrow p^*E \longrightarrow \mathcal{O}_Y(1)$ is $\mathcal{T} \otimes \mathcal{O}_Y(1)$. This induces a closed immersion

$$\mathbf{i}' : \tilde{Y} = Proj(\mathcal{T} * \mathcal{O}_Y(1)) \longrightarrow Proj(Sym(p^*E^1) = Y \times_V Z.$$

$\mathbf{i}'$ fits to a similar commutative diagram as (5) and as a result Φ and Ψ factor through pr_2 and pr_1. Both W and $\tilde{Y}$ lie inside $Y \times_V Z$ and ω and Ψ factor through pr_1 and ν and Φ factor through pr_2. So to prove the identification between $(\tilde{Y}, \Phi, \Psi)$ and (W, ν, ω) , it is enough to show that $\tilde{Y} \cong W$. This can be checked locally. So, after choosing a suitable open cover for V it is enough to prove $\tilde{Y} \cong W$ restricted to each of these open sets. Also we know that

$p^{-1}(U) \cong \mathbb{P}_U^{rk(E)-1}$ when $E_{|U}$ is trivial and $\mathbb{P}_U^n = \mathbb{P}_{\mathbb{C}}^n \times U$. Now the the isomorphism follows from [proposition 9.11, [EH]] after adjusting the the definition of projectivization in terms of [H].

We now turn our attention to the diagram (3). observe that if we fix the notations $W' = \mathbb{P}'_Z(G')$ with $\omega' : W' \longrightarrow Y'$ as discussed above then we have an identification between $(\tilde{Y}', \Phi', \Psi')$ and (W', ν', ω').

$\omega' : W' \longrightarrow Y'$ comes with $(\omega')^*\mathcal{O}_{Y'}(1) = \mathcal{O}_{\mathbb{P}_{Z'}(G')}(1)$. So, $\gamma = (\Psi')^*\,\zeta_1$ is achieved. $(\Phi')^*F_1 = (\Psi')^*F$ follows from the commutativity of the diagram (3).

The closed immersion $\mathbf{i}'$ induces a relation between the $\mathcal{O}(1)$ sheaves of $Y \times_V Z$ and $\tilde{Y}$. For $Y \times_V Z$ the $\mathcal{O}(1)$ sheaf is $pr_2^*\mathcal{O}_Z(1)$ and for $Proj(\mathcal{T} * \mathcal{O}_Y(1)$ the $\mathcal{O}(1)$ sheaf is $\mathcal{O}_{\tilde{Y}}(-\tilde{E}) \otimes (\Psi)^*\mathcal{O}_Y(1)$. Since Φ factors through pr_2, $(\Phi)^*\mathcal{O}_Z(1) = \mathcal{O}_{\tilde{Y}}(-\tilde{E}) \otimes (\Psi)^*\mathcal{O}_Y(1)$. In the particular case (see diagram (3)) $(\Phi')^*\mathcal{O}_{Z'}(1) = \mathcal{O}_{\tilde{Y}'}(-\tilde{E}) \otimes (\Psi')^*\mathcal{O}_{Y'}(1)$ i. e. $(\Phi')^*\,\zeta_{Z'} = (\Psi')^*\,\zeta_1 - \tilde{E}$.

Next consider the short exact sequence:

$$0 \longrightarrow \mathcal{O}_{Z'}(1) \longrightarrow G' \longrightarrow \overline{p}_2^*\pi_2^*Q_{11} \longrightarrow 0$$

We wish to calculate below the total chern class of G' through the chern class relation obtained from the above short exact sequence.

$$c(G') = c(\mathcal{O}_{Z'}(1)) \cdot c(\overline{p}_2^*\pi_2^*Q_{11}) = (1 + \zeta_{Z'}) \cdot \overline{p}_2^*\pi_2^*(1 + d_{11}[pt]) = (1 + \xi_{Z'})(1 + d_{11}F_1)$$

From the grothendieck relation for G' we have

$$\begin{aligned}
&\gamma^{n_{11}+1} - \Phi'^*(\zeta_{Z'} + d_{11}F_1) \cdot \gamma^{n_{11}} + \Phi'^*(d_{11}F_1 \cdot \zeta_{Z'}) \cdot \gamma^{n_{11}-1} = 0 \\
\Rightarrow\ &\gamma^{n_{11}+1} - (\Psi'^*\zeta_1 - \tilde{E}) + d_{11}\Psi'^*F) \cdot \gamma^{n_{11}} + d_{11}(\Psi'^*\zeta_1 - \tilde{E}) \cdot \Psi'^*F) \cdot \gamma^{n_{11}-1} = 0 \\
\Rightarrow\ &\tilde{E} \cdot \gamma^{n_{11}} - d_{11}\tilde{E} \cdot \Psi'^*F \cdot \gamma^{n_{11}-1} = 0 \\
\Rightarrow\ &\tilde{E} \cdot \Psi'^*(\zeta_1 - \mu_{11}F)^{n_{11}} = 0
\end{aligned}$$

For the last part note that $\tilde{E} = \mathbb{P}(\pi_2^*Q_{11}) \times_{\mathbb{P}(E_2)} Z'$. Also $N(\tilde{Y}')$ and $N(\tilde{E})$ are free $N(Z')$-module. Using these informations and projection formula, the identity $\tilde{E} \cdot N(\tilde{Y}') = (j_{\tilde{E}})_*N(\tilde{E})$ is obtained easily.

□

Now we are in a position to prove the next theorem.

Theorem 3.5. *$\overline{\mathrm{Eff}}^k(X) \cong \overline{\mathrm{Eff}}^k(Y') \cong \overline{\mathrm{Eff}}^k(Z')$ and $\overline{\mathrm{Eff}}^k(Z') \cong \overline{\mathrm{Eff}}^k(Z'')$. So, $\overline{\mathrm{Eff}}^k(X) \cong \overline{\mathrm{Eff}}^k(Z'')$ for $k < r_1 + r_2 - 1 - \mathbf{n}$*

where $Z' = \mathbb{P}(E_{11}) \times_C \mathbb{P}(E_2)$ and $Z'' = \mathbb{P}(E_{11}) \times_C \mathbb{P}(E_{21})$

Proof. Since $Y' = \mathbb{P}(\pi_2^*E_1) \cong \mathbb{P}(E_1) \times_C \mathbb{P}(E_2) = X$, $\overline{\mathrm{Eff}}^k(X) \cong \overline{\mathrm{Eff}}^k(Y')$ is followed at once. To prove that $\overline{\mathrm{Eff}}^k(X) \cong \overline{\mathrm{Eff}}^k(Z')$ we first define the the map: $\theta_k : N^k(X) \longrightarrow N^k(Z')$ by

$$\zeta_1^i \cdot \zeta_2^{k-i} \mapsto \bar{\zeta}_1^{\,i} \cdot \bar{\zeta}_2^{\,k-i}, \quad F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1} \mapsto F_1 \cdot \bar{\zeta}_1^j \cdot \bar{\zeta}_2^{k-j-1}$$

where $\bar{\zeta}_1 = \overline{p}_1^*(\mathcal{O}_{\mathbb{P}(E_{11})}(1))$ and $\bar{\zeta}_2 = \overline{p}_2^*(\mathcal{O}_{\mathbb{P}(E_2)}(1))$. $\overline{p}_1 : \mathbb{P}(E_{11}) \times_C \mathbb{P}(E_2) \longrightarrow \mathbb{P}(E_{11})$ and

$\overline{p}_2 : \mathbb{P}(E_{11}) \times_C \mathbb{P}(E_2) \longrightarrow \mathbb{P}(E_2)$ are respective projection maps.

It is evident that the above map is in isomorphism of abstract groups. We claim that this induces an isomorphism between $\overline{\text{Eff}}^k(X)$ and $\overline{\text{Eff}}(Z')$. First we construct an inverse for θ_k. Define $\Omega_k : N^k(Z') \longrightarrow N^k(X)$ by

$$\Omega_k(l) = {\Psi'}_*{\Phi'}^*(l)$$

Ω_k is well defined since Φ' is flat and Ψ' is birational. Ω_k is also pseudo-effective. Now we need to show that Ω_k is the inverse of θ_k.

$$\begin{aligned}
\Omega_k(\bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i}) &= {\Psi'}_*((\Phi'^*\bar{\zeta_1})^i \cdot (\Phi'^*\bar{\zeta_2})^{k-i}) \\
&= {\Psi'}_*((\Phi'^*\zeta_{Z'})^i \cdot (\Phi'^*\bar{\zeta_2})^{k-i}) \\
&= {\Psi'}_*((\Psi'^*\zeta_1 - \tilde{E})^i \cdot (\Psi'^*\zeta_2)^{k-i}) \\
&= {\Psi'}_*((\sum_{0\leq c\leq i} (-1)^i \tilde{E}^c (\Psi'^*\zeta_1)^{i-c}) \cdot (\Psi'^*\zeta_2)^{k-i})
\end{aligned}$$

Similarly,

$$\Omega_k(F_1 \cdot \bar{\zeta_1}^j \cdot \bar{\zeta_2}^{k-j-1}) = {\Psi'}_*((\sum_{0\leq d\leq j} (-1)^j \tilde{E}^d (\Psi'^*\zeta_1)^{j-d}) \cdot (\Psi'^*\zeta_2)^{k-j-1})$$

So,

$$\begin{aligned}
&\Omega_k\Big(\textstyle\sum_i a_i\, \bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i} + \sum_j b_j\, F_1 \cdot \bar{\zeta_1}^j \cdot \bar{\zeta_2}^{k-j-1}\Big) \\
&= \Big(\sum_i a_i\, {\zeta_1}^i \cdot {\zeta_2}^{k-i} + \sum_j b_j\, F \cdot {\zeta_1}^j \cdot {\zeta_2}^{k-j-1}\Big) + {\Psi'}_*\Big(\sum_i \sum_{1\leq c\leq i} \tilde{E}^c \Psi'^*(\alpha_{i,c}) + \sum_j \sum_{1\leq d\leq j} \tilde{E}^d \Psi'^*(\beta_{j,d})\Big)
\end{aligned}$$

for some cycles $\alpha_{i,c}, \beta_{j,d} \in N(X)$. But, $\Psi'^*(\tilde{E}^t) = 0$ for all $1 \leq t \leq i \leq r_1 + r_2 - 1 - \mathbf{n}$ for dimensional reasons. Hence, the second part in the right hand side of the above equation vanishes and we make the conclusion that $\Omega_k = \theta_k^{-1}$.

Next we seek an inverse of Ω_k which is pseudo-effective and meet our demand of being equal to θ_k. Define $\eta_k : N^k(X) \longrightarrow N^k(Z')$ by

$$\eta_k(s) = {\Phi'}_*(\delta \cdot \Psi'^* s)$$

where $\delta = \Psi'^*(\xi_2 - \mu_{11}F)^{n_{11}}$.

By the relations (5) and (6) , $\Psi'^*((\zeta_1^i \cdot \zeta_2^{k-i})$ is $\Phi'^*(\bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i})$ modulo $\tilde{E}$ and $\delta \cdot \tilde{E} = 0$. Also ${\phi'}_*\delta = [Z']$ which is derived from the fact that ${\Phi'}_*\gamma^{n_{11}} = [Z']$ and the same relations (5) and (6). Therefore

$$\eta_k(\zeta_1^i \cdot \zeta_2^{k-i}) = {\Phi'}_*(\delta \cdot \Phi'^*(\bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i})) = (\bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i}) \cdot [Z'] = \bar{\zeta_1}^i \cdot \bar{\zeta_2}^{k-i}$$

In a similar way, $\Psi'^*(F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1})$ is $\Phi'^*(F_1 \cdot \bar{\zeta_1}^j \cdot \bar{\zeta_2}^{k-j-1})$ modulo $\tilde{\mathbf{E}}$ and as a result of this

$$\eta_k(F \cdot \zeta_1^j \cdot \zeta_2^{k-j-1}) = F_1 \cdot \bar{\zeta_1}^j \cdot \bar{\zeta_2}^{k-j-1}$$

So, $\eta_k = \theta_k$.

Next we need to show that η_k is a pseudo- effective map. Notice that $\Psi'^* s = \bar{s} + \mathbf{j}_* s'$ for any effective cycle s on X, where $\bar{s}$ is the strict transform under Ψ' and hence effective. Now δ is intersection of nef classes. So, $\delta \cdot \bar{s}$ is pseudo-effective. Also $\delta \cdot \mathbf{j}_* s' = 0$ from theorem 2.4 and ${\Phi'}_*$ is pseudo-effective. Therefore η_k is pseudo-effective and first part of the theorem is proved. We will sketch the prove for the second part i.e. $\overline{\mathrm{Eff}}^k(Z') \cong \overline{\mathrm{Eff}}^k(Z'')$ which is similar to the proof of the first part. Consider the following diagram:

$$\begin{array}{ccc} Z'' = \mathbb{P}(E_{11}) \times_C \mathbb{P}(E_{21}) & \xrightarrow{\hat{p}_2} & \mathbb{P}(E_{21}) \\ \downarrow{\scriptstyle \hat{p}_1} & & \downarrow{\scriptstyle \hat{\pi}_2} \\ \mathbb{P}(E_{11}) & \xrightarrow{\hat{\pi}_1} & C \end{array} \tag{8}$$

Define $\hat{\theta}_k : N^k(Z') \longrightarrow N^k(Z'')$ by

$$\bar{\zeta_1}^{\,i} \cdot \bar{\zeta_2}^{\,k-i} \mapsto \hat{\zeta_1}^{\,i} \cdot \hat{\zeta_2}^{\,k-i}, \quad F \cdot \bar{\zeta_1}^{\,j} \cdot \bar{\zeta_2}^{\,k-j-1} \mapsto F_2 \cdot \hat{\zeta_1}^{\,j} \cdot \hat{\zeta_2}^{\,k-j-1}$$

where $\hat{\zeta_1} = \hat{p_1}^*(\mathcal{O}_{\mathbb{P}(E_{11})}(1)), \hat{\zeta_2} = \hat{p_2}^*(\mathcal{O}_{\mathbb{P}(E_{21})}(1))$ and F_2 is the class of a fibre of $\hat{\pi_1} \circ \hat{p_1}$.

This is a isomorphism of abstract groups and behaves exactly the same as θ_k. The methods applied to get the result for θ_k can also be applied successfully here.

Acknowledgement. The author would like to thank Prof. D.S. Nagaraj, IISER Tirupati for suggestions and discussions at every stage of this work. This work is supported financially by a fellowship from IMSc,Chennai (HBNI), DAE, Government of India. □

References

[CC] D. Chen, I. Coskun *Extremal higher codimension cycles on modulii spaces of curves*, Proc. Lond. Math. Soc. 111 (2015) 181- 204.

[CLO] I. Coskun, J. Lesieutre, J. C. Ottem *Effective cones of cycles on blow-ups of projective space*, Algebra Number Theory, 10(2016) 1983-2014.

[DELV] O. Debarre, L. Ein, R. Lazarsfeld, C. Voisin *Pseudoeffective and nef classes on abelian varieties*, Compos. Math. 147 (2011) 1793-1818.

[DJV] O. Debarre, Z. Jiang, C. Voisin *Pseudo-effective classes and pushforwards*, Pure Appl. Math. Q. 9 (2013) 643-664.

[EH] D. Eisenbud, J. Harris *3264 and all that- a second course in algebraic geometry*, Cambridge University Press, Cambridge, 2016.

[F] Mihai Fulger *The cones of effective cycles on projective bundles over curves*, Math. Z. 269, (2011) 449-459.

[FL1] M. Fulger, B. Lehmann *Positive cones of dual cycle classes*, Algebr. Geom. 4(2017) 1- 28.

[FL2] M. Fulger, B. Lehmann *Kernels of numerical pushforwards*, Adv. Geom. 17(2017) 373-378.

[Ful] William Fulton *Intersection Theory, 2nd ed.*, Ergebnisse der Math. und ihrer Grenzgebiete(3), vol 2, Springer, Berlin(1998)

[H] Robin Hartshorne *Algebraic Geometry. Graduate Texts in Mathematics*, Springer- Verlag, New York Heidelberg, (1977)

[KMR] R. Karmakar, S. Misra, N. Ray *Nef and Pseudoeffective cones of product of projective bundles over a curve* Bull. Sci. math. (2018), https://doi.org/10.1016/j.bulsci.2018.12.002

[PP] N. Pintye, A. Prendergast-Smith *Effective cycles on some linear blow ups of projective spaces*, 2018, https://arxiv.org/abs/1812.08476.

Institute of Mathematical Sciences, CIT Campus, Taramani, Chennai 600113, India and Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400094, India
E-mail address, Rupam Karmakar: rupamk@imsc.res.in