

Hamiltonian and Diffeomorphism Constraints Generalized for Timelike and Spacelike 3+1 Foliation

Leonid Perlov,
Department of Physics, University of Massachusetts, Boston, USA
leonid.perlov@umb.edu

November 8, 2021

Abstract

The form of Hamiltonian and Diffeomorphism constraints in Sen-Ashtekar-Barbero-Immirzi variables is well known for the spacelike 3+1 ADM foliation. It is also known that Sen-Ashtekar-Barbero-Immirzi connection can be introduced only in 3 dimensional space and does not work for $D > 3$. The reason it works in $D = 3$ is due to existence of isomorphism between $so(3)$ algebra and R^3 space with a vector product. It turns out that similar isomorphism exists between $so(2, 1)$ algebra and $R_{2,1}^3$ space algebra with respect to its vector product. By using this isomorphism we find both analog of Sen-Ashtekar-Barbero-Immirzi connection for timelike 3+1 foliation and the corresponding forms of Gauss, Diffeomorphism and Hamiltonian constraints. We then combine spacelike and timelike foliation constraints into the generalized form of the Hamiltonian and Diffeomorphism constraints using generalized Sen-Ashtekar-Barbero-Immirzi connection variables. We prove that Immirzi parameter is covariant with respect to timelike-spacelike ADM foliation change as in both cases in self-dual Ashtekar case it disappears in Hamiltonian constraint keeping it polynomial.

1 Introduction

It is known that Sen-Ashtekar-Barbero-Immirzi connection [8], [9] and flux variables can be introduced only in 3 dimensional space and do not work for $D > 3$, see [1]. The reason it works in $D = 3$ is due to isomorphism between $so(3)$ algebra and R^3 space with a vector product. Such isomorphism does not exist for $D > 3$, and, therefore it is impossible to introduce Sen-Ashtekar-Barbero-Immirzi connection.

We have noticed that similar isomorphism also exists in $D = 3$ between $so(2, 1)$ algebra and algebra of vectors in $R_{2,1}^3$ space with its vector product [2]. By using this isomorphism we derive Sen-Ashtekar-Barbero-Immirzi formalism in timelike foliation ADM with $SO(2, 1)$ structure group. We obtain a new connection in that case and corresponding rotational, Gauss, Diffeomorphism and Hamiltonian constraints. We then

combine $so(3)$ connection for spacelike foliation ADM with $so(2, 1)$ connection for timelike foliation ADM into one expression, which we call a generalized connection. We continue by combining rotational, Gauss, Diffeomorphism and Hamiltonian set of constraints into one set of generalized constraints for both cases.

Recently 3+1 timelike foliation obtained much attention due to an attempt to make the next step towards covariant theory. A timelike foliation was considered in [3], [4], [5], [7], [6], providing new variables.

Our result is novel as we obtain the timelike case constraints in a much simpler form and directly from the original ADM variables by using isomorphism between $so(2, 1)$ and $R_{2,1}^3$ algebras. Moreover, we were able to combine spacelike and timelike cases together into generalized form by using the relation between $so(2, 1)$ and $so(3)$ algebra structure coefficients.

The paper is organized as follows. In section 2 we discuss $so(2, 1) \rightarrow R_{2,1}^3$ isomorphism. Then in 3 we remind the formalism of the spacetime foliation by the time-like surfaces in original variables P^{ab}, q_{ab} . In the following section 4 we introduce $SO(2, 1)$ rotational constraints and extrinsic curvature variables K_a^i . We then write $SO(3)$ and $SO(2, 1)$ rotational constraints in a combined generalized form. In section 5 we obtain a generalized Sen-Ashtekar-Barbero-Immirzi connection for both time-like and spacelike ADM foliation as well as a generalized covariant derivative and a generalized Gauss constraint. In section 6 we introduce a generalized canonical transformation between variables (q_{ab}, P^{cd}) and Sen-Ashtekar-Barbero-Immirzi variables (A_a^i, E_a^i) . By calculating Poisson brackets we show that the symplectic structure is preserved, i.e the transformation is canonical. In section 7 we derive a diffeomorphism constraint for timelike ADM foliation. We then combine it with the spacelike ADM foliation into one generalized equation. In section 8 we derive the generalized form of the Hamiltonian constraint. A number of appendices below show all calculation details. Everywhere below we use the following index convention for symmetrization and anti-symmetrization: $A_{(a}A_{b)} = \frac{1}{2}(A_a + A_b)$, $A_{[a}A_{b]} = \frac{1}{2}(A_a - A_b)$

2 $so(2, 1) \rightarrow R_{2,1}^3$ Isomorphism and Structure coefficients

Before considering $so(2, 1) \rightarrow R_{2,1}^3$ isomorphism we will first remind how the similar isomorphism is written in spacelike ADM with $SO(3)$ structure group. The dynamics equations can be written in Sen-Ashtekar-Barbero-Immirzi polynomial form in that case only because of isomorphism between $so(3)$ algebra and algebra R^3 with vector product. This isomorphism makes $so(3)$ regular and adjoint representations the same, which is not the case for any dimension higher than 3, as such isomorphism exists only in $D = 3$. The isomorphism can be written in the following form:

$$\Gamma_{ai}^l = \epsilon_{kiso(3)}^l \Gamma_a^k \quad (1)$$

,where $\epsilon_{kiso(3)}^l = \epsilon_{likso(3)} = \epsilon^{lki}_{so(3)}$ - fully antisymmetric $so(3)$ tensor, which is also $so(3)$ algebra structure coefficients with $\epsilon^{012}_{so(3)} = 1$, $\epsilon_{012so(3)} = 1$, while Γ_a^k

are vectors in R^3 . The covariant derivative can be written as:

$$D_a E_i^a = [D_a E^a]_i + \Gamma_{ai}^l E_l^a = \partial_a E_i^a + \epsilon_{kiso(3)}^l \Gamma_a^k E_l^a \quad (2)$$

Similar isomorphism exists between $so(2, 1)$ algebra and algebra of $R_{2,1}^3$ vectors with respect to its vector product:

$$\Gamma_{ai}^l = \epsilon_{kiso(2,1)}^l \Gamma_a^k = \epsilon_{ikso(2,1)}^l \Gamma_a^k \quad (3)$$

,where $\epsilon_{kiso(2,1)}^l$ is $so(2, 1)$ antisymmetric tensor and algebra $so(2, 1)$ structure coefficients with $\epsilon_{so(2,1)}^{012} = 1, \epsilon_{012so(2,1)} = -1$, while Γ_a^k are vectors in $R_{2,1}^3$.

Since $\epsilon_{kiso(2,1)}^l = \eta_{ij} \epsilon_{kso(3)}^{jl}$, where $\eta_{ij} = \text{Diag}(-1, 1, 1)$, we can also rewrite it as:

$$\Gamma_{ai}^l = \eta_{ij} \epsilon_{kso(3)}^{jl} \Gamma_a^k \quad (4)$$

By using this isomorphism the covariant derivative can be written as:

$$D_a E_i^a = [D_a E^a]_i + \Gamma_{ai}^l E_l^a = \partial_a E_i^a + \epsilon_{ikso(2,1)}^l \Gamma_a^k E_l^a = \partial_a E_i^a + \eta_{ij} \epsilon_{kso(3)}^{jl} \Gamma_a^k E_l^a \quad (5)$$

3 Space-time foliation with timelike surfaces

The ADM foliation with spacelike surfaces is well known, see for example [1], [12]. The ADM foliation with timelike surfaces is less known, even though, it exists, see for example [11], [3], [6]. In the last two papers it is obtained from the Holst action [10], while in the first one it's obtained in the original variables even for the more general case of Einstein-Gauss-Bonnet gravity. If one ingores the Gauss-Bonnet additional term, then it becomes the ADM timelike foliation of Einstein gravity.

We will repeate in brief this formalism in the original variables. The projected metric differs in spacelike and timelike foliations cases only by a sign s:

$$q_{\mu\nu} = g_{\mu\nu} - s n_\mu n_\nu \quad (6)$$

,where n_μ is a unit normal vector to a foliation surface Σ with $n_\mu n^\mu = s$. The vector n_μ is timelike and foliation is by spacelike surfaces when $s = -1$, and, correspondingly, n_μ is spacelike and the foliation is by timelike surfaces when $s = 1$. Lagrangian expressed via projected metric is:

$$L = {}^{(4)} R \sqrt{-\det(g)} = |q|^{1/2} N {}^{(4)} R = |q|^{1/2} N (R - s(K_{ab} K^{ab} - K^2)) \quad (7)$$

,where N is the normal projection of the time coordinate in spacelike foliation and a normal projection of the space coordinate in the timelike foliation.

$$K_{ab} = \frac{1}{2N} (\dot{q}_{ab} - (L_{\vec{N}} q)_{ab}) \quad (8)$$

ADM action for both timelike and spacelike case in P^{ab}, q_{ab} variables is

$$S = \int d^4x |q|^{1/2} N (R - s(K_{ab}K^{ab} - K^2)) \quad (9)$$

$$P^{ab} = \frac{\partial L}{\partial \dot{q}_{ab}} = -s|q|^{1/2} (K^{ab} - \text{tr}(K)q^{ab}) \quad (10)$$

The Legendre transform then produces for both cases:

$$S = \int d^4x (\dot{q}_{ab}P^{ab} + N^a H_a + NH) \quad (11)$$

, where

$$H_a = -2q_{ac}D_b P^{bc} \quad (12)$$

$$H = \frac{-s}{\sqrt{|q|}} \left[q_{ac}q_{bd} - \frac{1}{D-1} q_{ab}q_{cd} \right] P^{ab}P^{cd} - \sqrt{|q|} R \quad (13)$$

4 Generalized SO(2,1)-SO(3) Rotational Constraint

In this section we will obtain a new rotational constraint for timelike ADM foliation with $SO(2,1)$ structure group. We begin with (q_{ab}, P^{cd}) and introduce the $SO(2,1)$ Sen-Ashtekar-Barbero-Immirzi variables in a canonical way. On four dimensional Lorentz manifold with 3+1 timelike foliation we introduce a bundle space with triads invariant with respect to $SO(2,1)$ rotation.

$$q_{ab} = e_a^i e_b^j \eta_{ij} \quad (14)$$

, where η_{ij} is Minkowski $R_{1,2}^3$ metric $\eta_{ij} = \text{Diag}(-1, 1, 1)$

We introduce an electric flux variable as a weight one density:

$$E_j^a = |\det(e_a^i)| e_j^a, \quad E_a^j = e_a^j / |\det(e_a^i)| \quad (15)$$

We will use the notation $q = -|\det(e_a^i)|^2$

We then introduce K_a^i one-form in a little different way than in a spacelike case (notice η_{ij}):

$$K_{ab} := K_{(a}^i e_{b)}^j \eta_{ij} \quad (16)$$

satisfying modified rotational constraint, again notice η_{ij} :

$$G_{ab} := K_{[a}^i e_{b]}^j \eta_{ij} = 0 \quad (17)$$

By using (15) we can rewrite it as:

$$G_{ab} := K_{[a}^i E_{b]}^j \eta_{ij} = 0 \quad (18)$$

or by raising indices a and b we obtain the form:

$$G^{ab} := q^{at}q^{be}K_{[t}^iE_{e]}^j\eta_{ij} = 0 \quad (19)$$

Notice that (14), (15), (16), (17), (18) and (19) differ from corresponding $SO(3)$ expressions by Minkowski metric η_{ij} presence.

The rotational constraint (18) can be converted into a different, although equivalent form, by contracting it with $e_k^a e_m^b$:

$$G_{ab}e_k^a e_m^b = K_a^i e_b^j \eta_{ij} e_k^a e_m^b - K_b^i e_a^j \eta_{ij} e_k^a e_m^b = K_{aj} e_k^a \delta_m^j - K_{bj} e_m^b \delta_k^j = 2K_{a[m} e_{k]}^a = 0 \quad (20)$$

,where we used $e_b^j e_m^b = \delta_m^j$ and $K_a^i \eta_{ij} = K_{aj}$. By using (15) and changing index m to j , it can also be written as:

$$G_{jk} = K_{a[j} E_{k]}^a = 0 \quad (21)$$

or by using $SO(3)$ antisymmetric tensor as:

$$G^i = \epsilon^{ijk}{}_{so(3)} K_{aj} E_k^a = 0 \quad (22)$$

Finally we can rewrite these constraints once again by contracting each G^i with η_{ij} .

$$G_i = \eta_{ij} \epsilon^{jkl}{}_{so(3)} K_{ak} E_l^a = \epsilon_i{}^{kl}{}_{so(2,1)} K_{ak} E_l^a = \epsilon_{ik}{}^l{}_{so(2,1)} K_a^k E_l^a = 0 \quad (23)$$

We define now $\bar{\eta}_{ij}$ to be $Diag(-1, 1, 1)$ in $SO(2, 1)$ case, and $Diag(1, 1, 1)$ in $SO(3)$ case, in order to write $SO(3)$ rotational constraint (22) and $SO(2, 1)$ rotational constraint (23) in a generalized form:

$$G_i = \bar{\eta}_{ij} \epsilon^{jkl}{}_{so(3)} K_{ak} E_l^a = \bar{\eta}_{ij} \epsilon_k{}^{jl}{}_{so(3)} K_a^k E_l^a = 0 \quad (24)$$

5 Generalized $SO(2,1)$ - $SO(3)$ Sen-Ashtekar-Barbero-Immirzi Connection and Generalized Gauss Constraint

By using (2) and (5) and a generalized metric $\bar{\eta}_{ij}$ defined at the end of the previous section, we can write the generalized covariant derivative as:

$$D_a E_i^a = [D_a E^a]_i + \Gamma_{ai}^l E_l^a = \partial_a E_i^a + \bar{\eta}_{ij} \epsilon_k{}^{jl}{}_{so(3)} \Gamma_a^k E_l^a \quad (25)$$

,where Γ_a^k are vectors correspondingly in $R_{2,1}^3$ for $SO(2, 1)$ timelike foliation, and in R^3 for $SO(3)$ spacelike foliation.

One can easily see that $R_{2,1}^3$ vectors Γ_a^i are invariant under Weyl canonical transformation in $SO(2, 1)$ timelike case similar to $SO(3)$ spacelike case:

$$(K_a^i, E_i^a) \rightarrow^{(\beta)} K_a^i = \beta K_a^i, \quad {}^{(\beta)} E_i^a = E_i^a / \beta \quad (26)$$

,where $\beta \in C$ is Barbero-Immirzi parameter.

An invariance follows from the explicit formula for Γ_a^i expressed via $\Gamma_{a,j}^l$ in (4), when the latter is expressed via triads. Thus in both $SO(3)$ and $SO(2, 1)$ case we can write it in a generalized form:

$$\begin{aligned}\Gamma_a^i &= \frac{1}{2} \bar{\eta}^{im} \epsilon_{m \text{ } so(3)}^{jk} e_k^b [e_{a,b}^j - e_{b,a}^j + e_j^c e_a^l e_{c,b}^l] \\ &= \frac{1}{2} \bar{\eta}^{im} \epsilon_{m \text{ } so(3)}^{jk} E_k^b [E_{a,b}^j - E_{b,a}^j + E_j^c E_a^l E_{c,b}^l] \\ &= \frac{1}{4} \bar{\eta}^{im} \epsilon_{m \text{ } so(3)}^{jk} E_k^b \left[2 E_a^j \frac{(\det(E))_{,b}}{\det(E)} - E_b^j \frac{(\det(E))_{,a}}{\det(E)} \right] \quad (27)\end{aligned}$$

We can see that Γ_a^i is a homogeneous function of degree zero, therefore $^{(\beta)}\Gamma_a^j = \Gamma_a^j$ and a covariant derivative D_a does not depend on β and $D_a(^{(\beta)}E_j^a) = 0$ in both $SO(3)$ [1] and $SO(2, 1)$ cases. It is easy to see that the generalized rotational constraint (24) also does not depend on β .

Therefore, by using (24) and the generalized covariant derivative (25), we can write:

$$\begin{aligned}G_i &= 0 + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} (^{\beta}K_a^k) (^{\beta}E_l^a) \\ &= D_a(^{\beta}E_i^a) + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} (^{\beta}K_a^k) (^{\beta}E_l^a) \\ &= \partial_a E_i^a + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} \Gamma_a^k (^{\beta}E_l^a) + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} (^{\beta}K_a^k) (^{\beta}E_l^a) \\ &= \partial_a E_i^a + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} \Gamma_a^k (^{\beta}E_l^a) + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} (^{\beta}K_a^k) (^{\beta}E_l^a) \\ &= \partial_a (^{\beta}E_i^a) + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} [\Gamma_a^k + (^{\beta}K_a^k)] (^{\beta}E_l^a) = ^{\beta}\mathcal{D}_a(^{\beta}E_i^a) = 0 \quad (28)\end{aligned}$$

or by introducing notation:

$$^{\beta}A_{ai}^l = \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} (\Gamma_a^k + (^{\beta}K_a^k)) \quad (29)$$

and notations:

$$^{\beta}A_a^k = \Gamma_a^k + (^{\beta}K_a^k), \quad ^{\beta}A_{ai}^l = \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} A_a^k \quad (30)$$

we can rewrite (28) as a generalized Gauss constraint:

$$\begin{aligned}^{\beta}\mathcal{D}_a(^{\beta}E_i^a) &= \partial_a (^{\beta}E_i^a) + (^{\beta}A_{ai}^l) (^{\beta}E_l^a) = \\ &= \partial_a (^{\beta}E_i^a) + \bar{\eta}_{ij} \epsilon_{k \text{ } so(3)}^{jl} [\Gamma_a^k + (^{\beta}K_a^k)] (^{\beta}E_l^a) = 0 \quad (31)\end{aligned}$$

6 Generalized Metric and Momentum Transformation to Sen-Ashtekar-Barbero-Immirzi Variables. Symplectic Structure

Before we go over to generalized $SO(2, 1) - SO(3)$ Diffeomorphism and Hamiltonian constraints we have to define a new transformation between q_{ab}, P^{cd} and Sen-Ashtekar-Barbero-Immirzi variables E_i^a, K_a^j for timelike ADM foliation with $SO(2, 1)$ structure

group, and prove that symplectic structure is preserved, i.e in new variables we have the same dynamics. Such transformation is well known for $SO(3)$ case [1]:

$$q_{ab} = E_a^i E_b^j |\det E_i^c|^{2/D-1}, \quad P^{cd} = |\det(E_e^c)|^{-2/(D-1)} (E_k^c E_k^t K_t^i E_i^d - E_k^c E_k^d K_t^i E_i^t) \quad (32)$$

Preserving symplectic structure

$$\{E_j^a(x), E_k^b(y)\} = \{K_a^j(x), K_b^k(y)\} = 0, \quad \{E_i^a(x), K_b^j(y)\} = \frac{k}{2} \delta_b^a \delta_i^j \delta(x, y) \quad (33)$$

,where $k = 16\pi G/c^3$ - gravitational coupling constant.

In order to define similar transformation for timelike $SO(2, 1)$ foliation we need to modify (32) in the following way:

$$q_{ab} = E_a^i E_b^j \eta_{ij} (-|\det E_i^c|^{2/D-1}), \quad P^{cd} = (-|\det(E_e^c)|^{-2/(D-1)}) (E_k^c E_m^t \eta^{mk} K_t^i E_i^d - E_k^c E_m^d \eta^{mk} K_t^i E_i^t) \quad (34)$$

Notice η_{ij} presence in several places.

We have to prove that this transformation preserves the same symplectic structure (33). Coordinate-coordinate Poisson bracket $\{q_{ab}, q_{cd}\}$ is zero, since q_{ab} contains only electric fluxes E_i^a and $\{E_j^a(x), E_k^b(y)\}$ is zero, as it follows from (33) and (34). We calculate momentum-momentum Poisson bracket by using momentum formula (34):

$$P^{ab}(x) = (-|\det(E_e^c)|^{-2/(D-1)}) (E_k^a E_m^t \eta^{mk} K_t^i E_i^b - E_k^a E_m^b \eta^{mk} K_t^i E_i^t) \quad (35)$$

$$P^{cd}(x) = (-|\det(E_e^c)|^{-2/(D-1)}) (E_k^c E_m^t \eta^{mk} K_t^i E_i^d - E_k^c E_m^d \eta^{mk} K_t^i E_i^t) \quad (36)$$

and notation $q := -|\det(E_e^c)|^{2/(D-1)}$ as follows:

$$\begin{aligned} \{P^{ab}(x), P^{cd}(y)\} = & \left\{ \frac{1}{q} (E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b - E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^t), \right. \\ & \left. \frac{1}{q} (E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d - E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^e) \right\} \quad (37) \end{aligned}$$

After lengthy calculations (see Appendix C) we obtain:

$$\{P^{ab}(x), P^{cd}(y)\} = 2kq^{ac} G^{bd} \quad (38)$$

,where

$$G^{bd} = q^{bt} q^{dp} K_t^j E_p^i \eta_{ij} - q^{dt} q^{bp} K_t^j E_p^i \eta_{ij} = K^{bj} E^{di} \eta_{ij} - K^{dj} E^{bi} \eta_{ij} = 2K^{j[b} E^{d]i} \eta_{ij} = 2e^{bt} e^{de} K_{[t}^j E_{e]}^i \eta_{ij}$$

is $SO(2, 1)$ rotational constraint (19).

When rotational constrain is zero, the Poisson bracket (38) is also zero. Therefore momentum-momentum Poisson bracket remains on shell the same as in original variables (q_{ab}, P^{cd}) . Exactly as in spacelike case.

Finally we consider coordinate-momentum bracket:

$$P^{ab}(x) = (-|\det(E_e^c)|^{-2/(D-1)})(E_k^a E_m^t \eta^{mk} K_t^i E_i^b - E_k^a E_m^b \eta^{mk} K_t^i E_i^t) \quad (39)$$

$$q_{cd}(y) = E_c^j E_d^m \eta_{mj} (-|\det(E_e^c)|)^{2/(D-1)} \quad (40)$$

$$\{P^{ab}(x), q_{cd}(y)\} = |\det(E_e^c)|^{-2/(D-1)} \{(E_k^a E_m^t \eta^{mk} K_t^i E_i^b - E_k^a E_m^b \eta^{mk} K_t^i E_i^t), E_c^j E_d^m \eta_{mj} |\det(E_e^c)|^{2/(D-1)}\} \quad (41)$$

After lengthy calculations (see Appendix C) we obtain:

$$\{P^{ab}(x), q_{cd}(y)\} = k \delta_{(c}^b \delta_{d)}^a \delta(x, y) \quad (42)$$

so, the symplectic structure is preserved, i.e. new variables E_i^a and K_a^i are canonical in $SO(2, 1)$ timelike foliation case as well.

What remains is to write generalized transformations in both $SO(3)$ and $SO(2, 1)$ cases. By looking at (34) we see that it turns into (32), when instead of Minkowski metric $\eta_{ij} = \text{Diag}(-1, 1, 1)$ we use Euclidean $\text{Diag}(1, 1, 1)$. Therefore, by using the generalized metric $\bar{\eta}_{ij}$ we write generalized transformations:

$$q_{ab} = E_a^i E_b^j \bar{\eta}_{ij} (-|\det E_i^c|^{2/D-1}), \quad P^{cd} = (-|\det(E_e^c)|^{-2/(D-1)})(E_k^c E_m^t \bar{\eta}^{mk} K_t^i E_i^d - E_k^c E_m^d \bar{\eta}^{mk} K_t^i E_i^t) \quad (43)$$

The symplectic structure (33) is the same in both cases so it can also be called generalized.

7 Generalized SO(2,1)-SO(3) Diffeomorphism Constraint

The diffeomorphism constraint in the original ADM variables can be written as in [1] (1.2.6):

$$H_a = -2s q_{ac} D_b P^{bc} \quad (44)$$

where $s = -1$ for $SO(3)$ spacelike foliation case, while $s = 1$ in $SO(2, 1)$ timelike case.

By substituting generalized variables (43) into (44) we obtain:

$$H_a = -2s q_{ac} D_b (-|\det(E_e^c)|^{-2/(D-1)} (E_k^b E_m^t \bar{\eta}^{mk} K_t^i E_i^c - E_k^b E_m^c \bar{\eta}^{mk} K_t^i E_i^t)) \quad (45)$$

or rewriting it by using metric expression: $q^{bt} = E_k^b E_m^t \bar{\eta}^{mk}$ as

$$H_a = -(2s/q) D_b (q_{ac} q^{bt} q K_t^i E_i^c - q_{ac} q^{bc} q K_t^i E_i^t) = -2s D_b (K^{bi} E_{ia} - \delta_a^b K_t^i E_i^t) \quad (46)$$

we obtain a diffeomorphism constraint in variables K_t^i and E_i^t :

$$H_a = -2s D_b (K^{bi} E_{ia} - \delta_a^b K_t^i E_i^t) = -2s D_b (K_b^i E_i^a - \delta_a^b K_t^i E_i^t) \quad (47)$$

,where we at the same time lowered b and raised a in the first term.

By using an equality that follows from the rotational constraint (18): we can show that (see Appendix G):

$$K_b^i E_i^a = K_a^i E_i^b \quad (48)$$

by substituting it into the first term of (47) we rewrite it as:

$$H_a = -2sD_b(K_a^i E_i^b - \delta_a^b K_t^i E_i^t) \quad (49)$$

In order to express this constraint via generalized connection A_a^i and electric flux E_i^a we introduce generalized curvatures:

$$R_{ab}^i = 2\partial_{[a}\Gamma_{b]}^i + \bar{\eta}^{ij}\epsilon_{jkl\,so(3)}\Gamma_a^k\Gamma_b^l \quad (50)$$

and

$$^{(\beta)}F_{ab}^i = 2\partial_{[a}^{(\beta)}A_{b]}^i + \bar{\eta}^{ij}\epsilon_{jkl\,so(3)}^{(\beta)}A_a^k A_b^l \quad (51)$$

,where $A_a^k = \Gamma_a^k + \beta K_a^k$ is $so(3)$ connection in spacelike foliation, and $so(2,1)$ connection in timelike foliation. See (50) and (51) derivation in Appendix D.

By expressing $^{(\beta)}F_{ab}^j$ via R_{ab}^j , (see derivation in Appendix E) we obtain:

$$^{(\beta)}F_{ab}^i = R_{ab}^i + 2\beta D_{[a}K_{b]}^i + \beta^2 \bar{\eta}^{ij}\epsilon_{jkl\,so(3)}K_a^k K_b^l \quad (52)$$

By contracting (52) with $^{(\beta)}E_i^b = E_i^b/\beta$ (see all calculations in Appendix F) we obtain:

$$^{(\beta)}F_{ab}^i {}^{(\beta)}E_i^b = \frac{R_{ab}^i E_i^b}{\beta} + 2D_{[a}K_{b]}^i E_i^b + \beta K_a^i G_j \quad (53)$$

As in $so(3)$ case the first term on the right hand side is zero (see Appendix K). The last term is also zero on a shell, where the rotational constraint is zero. As for the second term, we write it as:

$$\begin{aligned} 2D_{[a}K_{b]}^i E_i^b &= D_a(K_b^i)E_i^b - D_b(K_a^i)E_i^b = -(D_b(K_a^i)E_i^b - D_a(K_b^i)E_i^b) = \\ &= -(D_b(K_a^i)E_i^b - D_a(K_b^i)E_i^b) = \\ &= -(D_b(K_a^i)E_i^b - \delta_b^a D_a(K_b^i)E_i^b) = \\ &= -(D_b(K_a^i)E_i^b - \delta_b^a D_a(K_b^i)E_i^b) = \\ &= -(D_b(K_a^i E_i^b) - \delta_b^a D_a(K_b^i E_i^b)) = \\ &= -(D_b(K_a^i E_i^b) - \delta_b^a D_a(K_b^i E_i^b)) = (-s/2)H_a \end{aligned} \quad (54)$$

,where in order to go from the forth to the fifth line we used Lemma1 from Appendix A, while in the last line we used (49): $H_a = -2sD_b(K_a^i E_i^b - \delta_a^b K_t^i E_i^t)$.

If then follows from (53) and (54) that diffeomorphism constraint has the following form:

$$^{(\beta)}F_{ab}^i {}^{(\beta)}E_i^b = -(s/2)H_a + \beta K_a^i G_j \quad (55)$$

or on shell:

$$H_a = -2s {}^{(\beta)}F_{ab}^j {}^{(\beta)}E_j^b \quad (56)$$

8 Generalized SO(2,1)-SO(3) Hamiltonian Constraint

Let us derive a Hamiltonian constraint. We will remind first how it was derived in $SO(3)$ spacelike ADM foliation [1] case, and then we will derive it for $SO(2,1)$ time-like ADM foliation case.

$SO(3)$ Spacelike Foliation Case:

By contracting (52) with $\epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b$ we obtain:

$$\begin{aligned} {}^{(\beta)}F_{ab}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b = & -q \frac{R_{ab}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} e_k^a e_l^b}{\beta^2} + 2\beta D_{[a} K_{b]}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b + \\ & \beta^2 \epsilon_{mnso(3)}^j K_a^m K_b^n \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b \end{aligned} \quad (57)$$

,where minus in the first term on the right hand side is because we have moved index j by one position into the middle.

By using $R_{ab}^{kl} = R_{ab}^j \epsilon_j^{kl}$ (see derivation in Appendix D) we rewrite the first term.

By using (Appendix I) we rewrite the second term, and by using (Appendix J) we rewrite the third term as:

$${}^{(\beta)}F_{ab}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b = -q \frac{R_{ab}^{kl} e_k^a e_l^b}{\beta^2} - 2 \frac{E_j^a D_a G_j}{\beta} + (K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b) \quad (58)$$

or, when contracting Riemann tensor with triads $R_{ab}^{kl} e_k^a e_l^b = R$ we obtain:

$${}^{(\beta)}F_{ab}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b = -q \frac{R}{\beta^2} - 2 {}^{(\beta)}E_j^a D_a G_j + (K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b) \quad (59)$$

or

$$\begin{aligned} {}^{(\beta)}F_{ab}^j \epsilon_j^{kl} \epsilon_{so(3)}^{(\beta)} E_k^{a(\beta)} E_l^b + 2 {}^{(\beta)}E_j^a D_a G_j \\ = \frac{\sqrt{q}}{\beta^2} \left[-\sqrt{q} R - \beta^2 \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{q}} \right] \\ = \frac{\sqrt{q}}{\beta^2} \left[H + (s - \beta^2) \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{q}} \right] \end{aligned} \quad (60)$$

,where in order to go from the second to the third line, we substituted the expression for $-\sqrt{q}R$ from the expression for Hamiltonian in $SO(3)$ case (see [1] 4.2.7):

$$H = -\frac{s}{\sqrt{q}}(K_a^l K_b^j - K_a^j K_b^l) E_j^a E_l^b - \sqrt{q} R \quad (61)$$

It can be easily done when regrouping terms as follows:

$$H = -\frac{s}{\sqrt{q}}(K_a^l E_l^b K_b^j E_j^a - K_a^j E_j^a K_b^l E_l^b) - \sqrt{q} R = -\frac{s}{\sqrt{q}}((K_a^j E_j^b)(K_b^k E_k^a) - (K_a^j E_j^a)^2) - \sqrt{q} R \quad (62)$$

thus we obtain:

$$-\sqrt{q}R = H + -\frac{s}{\sqrt{q}}((K_a^j E_j^b)(K_b^k E_k^a) - (K_a^j E_j^a)^2) \quad (63)$$

We now express H in (60):

$$H = \frac{\beta^2}{\sqrt{q}} \left[{}^{(\beta)}F_{ab}^j \epsilon_j^{kl} {}^{so(3)} E_k^a {}^{(\beta)} E_l^b + 2 {}^{(\beta)} E_j^a D_a G_j \right] + (\beta^2 - s) \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{q}} \quad (64)$$

or

$$H = [\beta^2 {}^{(\beta)}F_{ab}^j - (\beta^2 - s) \epsilon_{jmn} {}^{so(3)} K_a^m K_b^n] \frac{\epsilon_j^{kl} {}^{so(3)} E_k^a E_l^b}{\sqrt{q}} \quad (65)$$

We have repeated [1] $SO(3)$ spacelike ADM foliation case for instructional aim. We will now derive a new Hamiltonian constraint for timelike ADM foliation with $SO(2, 1)$ structure group:

$SO(2, 1)$ Timelike Foliation Case:

By contracting (52) now with $\epsilon_j^{kl} {}^{so(2,1)} E_k^a {}^{(\beta)} E_l^b$, we obtain (see details in Appendices D, I and J):

$${}^{(\beta)}F_{ab}^j \epsilon_j^{kl} {}^{so(2,1)} E_k^a {}^{(\beta)} E_l^b = -q \frac{R}{\beta^2} - 2 \frac{E_j^a D_a G_j}{\beta} - ((K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b)) \quad (66)$$

By comparing (66) with (59), we see only one, however crucial difference: a different sign in the third term.

By using (66)

$$\begin{aligned} & {}^{(\beta)}F_{ab}^j \epsilon_j^{kl} {}^{so(2,1)} E_k^a {}^{(\beta)} E_l^b + 2 {}^{(\beta)} E_j^a D_a G_j \\ &= \frac{\sqrt{|q|}}{\beta^2} \left[-\sqrt{|q|}R + \beta^2 \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{|q|}} \right] \\ &= \frac{\sqrt{|q|}}{\beta^2} \left[H + (s + \beta^2) \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{|q|}} \right] \quad (67) \end{aligned}$$

,where in order to go from the second to the third line, we substituted the expression for $\sqrt{q}R$ from Hamiltonian expression in case of timelike foliation (see Appendix B for details):

$$H = -\frac{s}{\sqrt{|q|}}(K_a^l K_b^j - K_a^j K_b^l) E_j^a E_l^b - \sqrt{|q|}R \quad (68)$$

Notice the only, however crucial change in (67) compared to (60). We have $(s + \beta^2)$ instead of $(s - \beta^2)$. This is due to the sign difference in the third term of (66) and (59),

which is in turn a consequence of the Riemann tensor contraction with $\epsilon_{jkl\,so(2,1)}$ vs $\epsilon^{jkl}_{\,so(3)}$, since:

$$\epsilon_{jkl\,so(3)}\epsilon^{jmn}_{\,so(3)} = 2\delta_k^{[m}\delta_l^{n]} \quad (69)$$

while

$$\epsilon_{jkl\,so(2,1)}\epsilon^{jmn}_{\,so(2,1)} = -2\delta_k^{[m}\delta_l^{n]} \quad (70)$$

By expressing H in (67) we obtain:

$$H = \frac{\beta^2}{\sqrt{|q|}} \left[{}^{(\beta)}F_{ab}^j \epsilon_j{}^{kl}_{\,so(2,1)} {}^{(\beta)}E_k^a E_l^b + 2 {}^{(\beta)}E_j^a D_a G_j \right] - (\beta^2 + s) \frac{(K_b^j E_j^a)(K_a^k E_k^b) - (K_a^j E_j^a)^2}{\sqrt{|q|}} \quad (71)$$

or

$$H = [\beta^{2(\beta)} F_{ab}^j - (\beta^2 + s) \epsilon_{mn\,so(2,1)}^j K_a^m K_b^n] \frac{\epsilon_j{}^{kl}_{\,so(2,1)} E_k^a E_l^b}{\sqrt{|q|}} \quad (72)$$

We can now rewrite Hamiltonian constraints in (65) and (72) for $SO(3)$ and $SO(2,1)$ foliations in a general form:

$$H = [\beta^{2(\beta)} F_{ab}^j - (\beta^2 + 1) \bar{\eta}^{ji} \epsilon_{imn\,so(3)} K_a^m K_b^n] \frac{\bar{\eta}_{ji} \epsilon^{ikl}_{\,so(3)} E_k^a E_l^b}{\sqrt{|q|}} \quad (73)$$

,where, by remembering that $s = 1$ in $SO(2,1)$ and $s = -1$ in $SO(3)$ case, we have combined both $(\beta^2 - s)$, $s = -1$ in (65) and $(\beta^2 + s)$, $s = 1$ in (72) into one expression $(\beta^2 + 1)$. We also used the following identities: $\bar{\eta}^{ji} \epsilon_{imn\,so(3)} = \epsilon_{mn\,so(2,1)}^j$ in $SO(2,1)$ case and $\bar{\eta}_{ji} \epsilon^{ikl}_{\,so(3)} = \epsilon_j{}^{kl}_{\,so(3)}$ in $SO(3)$ case.

We can see in (73) that the self-dual Ashtekar case for Immirzi $\beta = \pm i$ is preserved in both $SO(3)$ and $SO(2,1)$ cases, making general formula very simple:

$$H = [F_{ab}^j \bar{\eta}_{ji} \epsilon^{ikl}_{\,so(3)} E_k^a E_l^b] = 0 \quad (74)$$

This fact is very important as it proves that Immirzi parameter is covariant with respect to spacelike - timelike foliation change.

Finally we need to check that the hamiltonian and diffeomorphism constraints in new variables commute with the smeared rotational constraint. Like in [1] we introduce the smeared constraint by using G_{ik} form (21)

$$G(\Lambda) = \int_{\sigma} d^3 x \Lambda^{jk} K_{aj} E_k^a \quad (75)$$

, where $\Lambda \in so(3)$ for spacelike foliation and $\in so(2, 1)$ for timelike foliation. The constraint satisfy the Poisson algebra:

$$\{G(\Lambda), G(\Lambda')\} = \frac{k}{2} G([\Lambda, \Lambda']) \quad (76)$$

Since coordinate and momentum in (43) are $so(3)$ invariant in $SO(3)$ spacelike foliation case and $so(2, 1)$ invariant in $SO(2, 1)$ timelike foliation case, they will commute with the corresponding smeared rotational constraint. Also both diffeomorphism and hamiltonian constraints commute with the smeared rotational constraint, since they are both functions of q_{ab} and P^{cd} . So the whole system of constraints is still first class.

9 All Generalized Constraints

To summarize we write the system of the generalized $SO(3) - SO(2, 1)$ constraints together:

$${}^\beta \mathcal{D}_a ({}^\beta E_i^a) = \partial_a ({}^\beta E_i^a) + \bar{\eta}_{ij} \epsilon_{so(3)}^{jl} [\Gamma_a^k + ({}^\beta K_a^k)] ({}^\beta E_l^a) = 0$$

$$H_a = -2s^{(\beta)} F_{ab}^{(i)} E_i^b = 0$$

$$H = [\beta^{2(\beta)} F_{ab}^i - (\beta^2 + 1) \bar{\eta}^{ij} \epsilon_{jmn so(3)} K_a^m K_b^n] \frac{\bar{\eta}_{ij} \epsilon_{so(3)}^{jkl} E_k^a E_l^b}{\sqrt{|q|}} = 0 \quad (77)$$

10 Discussion

The existence of isomorphism between $so(2, 1)$ algebra and algebra of vectors in $R_{2,1}^3$ space with vector product as algebra operation was noticed and used to derive Sen-Ashtekar-Barbero-Immirzi formalism for timelike foliation with $SO(2, 1)$ structure group. A new $so(2, 1)$ connection along with the Gauss, Diffeomorphism and Hamiltonian constraints have been obtained in $SO(2, 1)$ case. The constraints in $SO(3)$ spacelike and in $SO(2, 1)$ timelike ADM were combined into one set of generalized constraints using the generalized connection. In addition, it's been proved that Immirzi parameter is covariant with respect to timelike-spacelike ADM foliation change as in both cases in self-dual Ashtekar case it disappears in Hamiltonian constraint keeping it polynomial.

11 Appendix A Lemma 1 for Diffeomorphism Constraints

Lemma 1

$$(D_a(K_{jb}) - D_b(K_{ja}))E_j^b = D_a(K_{jb}E_j^b) - D_b(K_{ja}E_j^b) \quad (78)$$

Proof:

$$\begin{aligned} D_a(K_{jb}E_j^b) - D_b(K_{ja}E_j^b) &= \delta_a^b D_b(K_{jb})E_j^b + \delta_a^b K_{jb} D_b(E_j^b) - D_b(K_{ja})E_j^b - K_{ja} D_b(E_j^b) = \\ &= \delta_a^b D_b(K_{jb})E_j^b - D_b(K_{ja})E_j^b = (D_a(K_{jb}) - D_b(K_{ja}))E_j^b \end{aligned} \quad (79)$$

above in the first identity on the right hand side the second and the forth terms cancel.

12 Appendix B Hamiltonian Constraint in SO(2,1) Case

We would like to express Hamiltonian constraint first in K_a^i and E_i^a variables and then in A_a^i , E_i^a variables. We begin with Hamiltonian constraint ADM expression (13):

$$H = \frac{-s}{\sqrt{|q|}} \left[q_{ac}q_{bd} - \frac{1}{D-1}q_{ab}q_{cd} \right] P^{ab}P^{cd} - \sqrt{|q|}R \quad (80)$$

by substituting into it the metric q_{ab} and momentum P^{ab} expressions from (34) we obtain:

$$\begin{aligned} H &= \frac{-s}{\sqrt{|q|}} \left[q_{ac}q_{bd} - \frac{1}{D-1}q_{ab}q_{cd} \right] \frac{1}{q} (qq^{at_1}K_{t_1}^{i_1}E_{i_1}^b - qq^{ab}K_{t_2}^{i_2}E_{i_2}^{t_2}) \\ &\quad - \frac{1}{q} (qq^{ct_3}K_{t_3}^{i_3}E_{i_3}^d - qq^{cd}K_{t_4}^{i_4}E_{i_4}^{t_4}) - \sqrt{|q|}R = \\ &= \frac{-s}{\sqrt{|q|}} \left[q_{ac}q_{bd} - \frac{1}{D-1}q_{ab}q_{cd} \right] (K^{ai_1}E_{i_1}^b - q^{ab}K_{t_2}^{i_2}E_{i_2}^{t_2}) \\ &\quad (K^{ci_3}E_{i_3}^d - q^{cd}K_{t_4}^{i_4}E_{i_4}^{t_4}) - \sqrt{|q|}R \end{aligned} \quad (81)$$

by opening parentheses:

$$\begin{aligned} H &= \frac{-s}{\sqrt{|q|}} \left[q_{ac}q_{bd} - \frac{1}{D-1}q_{ab}q_{cd} \right] (K^{ai_1}E_{i_1}^b K^{ci_3}E_{i_3}^d - q^{cd}K^{ai_1}E_{i_1}^b K_{t_4}^{i_4}E_{i_4}^{t_4} \\ &\quad - q^{ab}K_{t_2}^{i_2}E_{i_2}^{t_2} K^{ci_3}E_{i_3}^d + q^{ab}q^{cd}K_{t_2}^{i_2}E_{i_2}^{t_2} K_{t_4}^{i_4}E_{i_4}^{t_4}) - \sqrt{|q|}R = \\ &= \frac{-s}{\sqrt{|q|}} ((K_c^{i_1}E_{di_1}K^{ci_3}E_{i_3}^d - q_{ab}K^{ai_1}E_{i_1}^b K_{t_4}^{i_4}E_{i_4}^{t_4} - q_{cd}K_{t_2}^{i_2}E_{i_2}^{t_2} K^{ci_3}E_{i_3}^d + \\ &\quad q_{cd}q^{cd}K_{t_2}^{i_2}E_{i_2}^{t_2} K_{t_4}^{i_4}E_{i_4}^{t_4}) - \frac{1}{D-1}(K_b^{i_1}E_{i_1}^b K_d^{i_3}E_{i_3}^d - DK_b^{i_1}E_{i_1}^b K_{t_4}^{i_4}E_{i_4}^{t_4} - DK_{t_2}^{i_2}E_{i_2}^{t_2} K_d^{i_3}E_{i_3}^d - D^2 K_{t_2}^{i_2}E_{i_2}^{t_2} K_{t_4}^{i_4}E_{i_4}^{t_4})) \\ &\quad - \sqrt{|q|}R = \frac{-s}{\sqrt{|q|}} ((K_c^{i_1}K^{ci_3}E_{di_1}E_{i_3}^d - K_b^{i_1}E_{i_1}^b K_{t_4}^{i_4}E_{i_4}^{t_4} - K_{t_2}^{i_2}E_{i_2}^{t_2} K_d^{i_3}E_{i_3}^d + DK_{t_2}^{i_2}E_{i_2}^{t_2} K_{t_4}^{i_4}E_{i_4}^{t_4}) \\ &\quad - \frac{(1-D)^2}{D-1}(K_b^{i_1}E_{i_1}^b K_d^{i_3}E_{i_3}^d)) - \sqrt{|q|}R = \frac{-s}{\sqrt{|q|}} (K_c^{i_1}K_{i_3}^c E_d^{i_1}E_{i_3}^d + (D-2)K_b^{i_1}E_{i_1}^b K_{t_4}^{i_4}E_{i_4}^{t_4} - (D-1)K_b^{i_1}E_{i_1}^b K_d^{i_3}E_{i_3}^d) \\ &\quad - \sqrt{|q|}R = \frac{-s}{\sqrt{|q|}} (K_c^{i_1}K_{i_3}^c E_d^{i_1}E_{i_3}^d - K_b^{i_1}E_{i_1}^b K_d^{i_3}E_{i_3}^d) - \sqrt{|q|}R \end{aligned} \quad (82)$$

We rewrite the final result by using the same indices as in the book:

$$H = \frac{-s}{\sqrt{|q|}} (K_a^l K_j^a E_l^b E_j^b - K_a^l K_b^j E_l^a E_j^b) - \sqrt{|q|}R \quad (83)$$

13 Appendix C Timelike $SO(2, 1)$ Symplectic Structure Calculations

Momentum-Momentum Poisson Bracket

$$P^{ab}(x) = -|\det(E_e^c)|^{-2/(D-1)}(E_k^a E_m^t \eta^{mk} K_t^i E_i^b - E_k^a E_m^b \eta^{mk} K_t^i E_i^t) = \frac{1}{q}(E_k^a E_m^t \eta^{mk} K_t^i E_i^b - E_k^a E_m^b \eta^{mk} K_t^i E_i^t) \quad (84)$$

$$P^{cd}(x) = -|\det(E_e^c)|^{-2/(D-1)}(E_k^c E_m^t \eta^{mk} K_t^i E_i^d - E_k^c E_m^d \eta^{mk} K_t^i E_i^t) = \frac{1}{q}(E_k^c E_m^t \eta^{mk} K_t^i E_i^d - E_k^c E_m^d \eta^{mk} K_t^i E_i^t) \quad (85)$$

$$\begin{aligned} \{P^{ab}(x), P^{cd}(y)\} &= \left\{ \frac{1}{q}(E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b - E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^t), \right. \\ &\quad \left. \frac{1}{q}(E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d - E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}) \right\} \quad (86) \end{aligned}$$

By introducing the following notations:

$$\begin{aligned} a &= 1/q \\ b &= E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b \\ c &= E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^t \\ d &= 1/q \\ e &= E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d \\ f &= E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} \end{aligned}$$

We can rewrite (86) as:

$$\{P^{ab}(x), P^{cd}(y)\} = \{a(b - c), d(e - f)\} \quad (87)$$

or by using the Leibniz rule for the Poisson brackets:

$$\begin{aligned} \{P^{ab}(x), P^{cd}(y)\} &= (a(\{b, d\} - \{c, d\}))(e - f) + \\ &\quad d(\{a, e\} - \{a, f\})(b - c) + \\ &\quad ad\{b, e\} - ad\{c, e\} - ad\{b, f\} + ad\{c, f\} + \\ &\quad \{a, d\}(e - f)(b - c) \quad (88) \end{aligned}$$

The last term is zero since $\{a, d\} = \{(\det E)^{\frac{-2}{D-1}}, (\det E)^{\frac{-2}{D-1}}\} = 0$, as $\{E_j^a(x), E_k^b(y)\} = 0$

Let's calculate separately $\{b, f\}, \{c, f\}, \{c, e\}, \{b, e\}, \{a, e\}, \{a, f\}, \{b, d\}, \{c, d\}$

$$\begin{aligned} \{b, e\} = \{ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d \} = \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} \{K_{t_1}^{i_1}, E_{m_1}^c\} E_{i_1}^b E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d + \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_1}^c \{K_{t_1}^{i_1}, E_{p_3}^{e_1} \eta^{p_3 m_1}\} E_{i_1}^b K_{e_1}^{j_1} E_{j_1}^d + \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} \{K_{t_1}^{i_1}, E_{j_1}^d\} E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} \{E_{k_1}^a, K_{e_1}^{j_1}\} E_{j_1}^d E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} E_{k_1}^a \{E_{p_1}^{t_1} \eta^{p_1 k_1} K_{e_1}^{j_1}\} E_{j_1}^d K_{t_1}^{i_1} E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} \{E_{i_1}^b, K_{e_1}^{j_1}\} E_{j_1}^d \quad (89) \end{aligned}$$

$$\begin{aligned} \{b, e\} = \{ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d \} = \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} (-\frac{k}{2} \delta_{m_1}^{i_1} \delta_{t_1}^c) E_{i_1}^b E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d + \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_1}^c (-\frac{k}{2} \delta_{m_1}^{i_1} \delta_{p_3}^{e_1} \eta^{p_3 t_1}) E_{i_1}^b K_{e_1}^{j_1} E_{j_1}^d + \\ & E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} (-\frac{k}{2} \delta_{j_1}^{i_1} \delta_{t_1}^d) E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} (\frac{k}{2} \delta_{k_1}^{j_1} \delta_{e_1}^a) E_{j_1}^d E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} E_{k_1}^a (\frac{k}{2} \delta_{e_1}^{t_1} \delta_{p_1}^{j_1}) \eta^{p_1 k_1} E_{j_1}^d K_{t_1}^{i_1} E_{i_1}^b + \\ & E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} (\frac{k}{2} \delta_{i_1}^{j_1} \delta_{e_1}^b) E_{j_1}^d = \\ & q^2 \frac{k}{2} (-q^{ac} q^{be_1} K_{e_1}^{j_1} E_{j_1}^d - q^{ae_1} q^{bc} K_{e_1}^{j_1} E_{j_1}^d - q^{ad} q^{ce_1} K_{e_1}^{j_1} E_{j_1}^b + q^{ca} q^{dt_1} K_{t_1}^{i_1} E_{i_1}^b \\ & + q^{ct_1} q^{da} K_{t_1}^{i_1} E_{i_1}^b + q^{cb} q^{at_1} K_{t_1}^{i_1} E_{i_1}^d) = \\ & = q^2 \frac{k}{2} q (-q^{ac} \hat{G}^{bd} - q^{bc} \hat{G}^{ad} - q^{ad} \hat{G}^{cb} + q^{ac} \hat{G}^{db} + q^{da} \hat{G}^{cb} + q^{cb} \hat{G}^{ad}) = \\ & = \frac{k}{2} q^3 q^{ac} (\hat{G}^{db} - \hat{G}^{bd}) = \frac{k}{2} q^3 q^{ac} G^{db} \quad (90) \end{aligned}$$

,where

$$G^{db} := \hat{G}^{db} - \hat{G}^{bd} \quad (91)$$

and we have introduced the notations for \hat{G} with various indices:

$$\hat{G}^{db} = q^{de_1} K_{e_1}^{j_1} q^{bp} E_p^{j_1} \quad (92)$$

Thus

$$\{b, e\} = \frac{k}{2} q^3 q^{ac} G^{db} \quad (93)$$

The next Poisson bracket is:

$$\begin{aligned}
\{a, e\} &= \{(\det E)^{\frac{-2}{D-1}}, E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d\} = \\
&= E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} \{(\det E)^{\frac{-2}{D-1}}, K_{e_1}^{j_1}\} E_{j_1}^d = \\
&= E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} \frac{-2}{D-1} (\det E)^{\frac{-2}{D-1}} \frac{\{(\det E), K_{e_1}^{j_1}\}}{\det E} E_{j_1}^d = \\
&= E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} \frac{-2}{D-1} \frac{1}{q} E_r^n \{E_n^r, K_{e_1}^{j_1}\} E_{j_1}^d = \\
&= q^{ce_1} q \frac{-2}{D-1} \frac{1}{q} E_r^n \left(\frac{k}{2} \delta_n^{j_1} \delta_{e_1}^r \right) E_{j_1}^d = \\
&= q^{ce_1} E_{j_1}^d \frac{-k}{D-1} E_{e_1}^{j_1} = q^{ce_1} \delta_{e_1}^d \frac{-k}{D-1} = q^{cd} \frac{-k}{D-1} \quad (94)
\end{aligned}$$

We obtain:

$$\{a, e\} = q^{cd} \frac{-k}{D-1} \quad (95)$$

The next bracket can be obtained from (94), by changing the sign and making the following index replacement:

$$c \rightarrow a, d \rightarrow b, a \rightarrow c, b \rightarrow d \quad (96)$$

$$\{b, d\} = \{E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, (\det E)^{\frac{-2}{D-1}}\} \quad (97)$$

$$\{b, d\} = q^{ab} \frac{k}{D-1} \quad (98)$$

The next bracket goes as follows:

$$\begin{aligned}
\{b, f\} &= \{E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = \\
&= E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} \{K_{t_1}^{i_1}, E_{m_2}^c\} E_{i_1}^b E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&= E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_2}^c \{K_{t_1}^{i_1}, E_{p_4}^d\} E_{i_1}^b K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&= E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} \{K_{t_1}^{i_1}, E_{j_2}^{e_2}\} E_{i_1}^b + \\
&= E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \{E_{k_1}^a, K_{e_2}^{j_2}\} E_{j_2}^{e_2} E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b + \\
&= E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_1}^a \{E_{p_1}^{t_1} \eta^{p_1 k_1}, K_{e_2}^{j_2}\} E_{j_2}^{e_2} K_{t_1}^{i_1} E_{i_1}^b + \\
&= E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} \{E_{i_1}^b, K_{e_2}^{j_2}\} E_{j_2}^{e_2} \quad (99)
\end{aligned}$$

or

$$\begin{aligned}
\{b, f\} &= \{E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = \\
&E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} \left(-\frac{k}{2} \delta_{m_2}^{i_1} \delta_{t_1}^c\right) E_{i_1}^b E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_2}^c \left(-\frac{k}{2} \delta_{p_4}^{i_1} \delta_{t_1}^d\right) \eta^{p_4 m_2} E_{i_1}^b K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} \left(-\frac{k}{2} \delta_{j_2}^{i_1} \delta_{t_1}^{e_2}\right) E_{i_1}^b + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \left(\frac{k}{2} \delta_{e_2}^a \delta_{k_1}^{j_2}\right) E_{j_2}^{e_2} E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_1}^a \left(\frac{k}{2} \delta_{e_2}^{t_1} \delta_{p_1}^{j_2}\right) \eta^{p_1 k_1} E_{j_2}^{e_2} K_{t_1}^{i_1} E_{i_1}^b + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} \left(\frac{k}{2} \delta_{e_2}^b \delta_{i_1}^{j_2}\right) E_{j_2}^{e_2} = \\
&\frac{k}{2} (-q^{ac} q q^{bd} q K_{e_2}^{j_2} E_{j_2}^{e_2} - q q^{ad} q q^{bc} K_{e_2}^{j_2} E_{j_2}^{e_2} - q^{ae_2} q q^{cd} q K_{e_2}^{j_2} E_{j_2}^b + \\
&q^{cd} q q^{at_1} q K_{t_1}^{i_1} E_{i_1}^b + q^{cd} q q^{at_1} q K_{t_1}^{i_1} E_{i_1}^b + q^{cd} q q^{at_1} q K_{t_1}^{i_1} E_{i_1}^b) = \\
&\frac{-k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + k q^2 q^{cd} q^{at_1} K_{t_1}^{i_1} E_{i_1}^b \quad (100)
\end{aligned}$$

To summarize

$$\begin{aligned}
\{b, f\} &= \{E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} \\
&= -\frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + k q^2 q^{cd} q^{at_1} K_{t_1}^{i_1} E_{i_1}^b \\
&= -\frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + k q^2 q^{cd} q^{at_1} K_{t_1}^{i_1} q q^{bp} E_p^{i_1} \\
&= -\frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + k q^3 q^{cd} K^{ai_1} E^{bi_1} \quad (101)
\end{aligned}$$

The bracket $\{c, e\}$ is similar to $\{f, b\} = -\{b, f\}$ above in (100) with the following index replacement:

$$a \rightarrow c, b \rightarrow d, c \rightarrow a, d \rightarrow b \quad (102)$$

$$\begin{aligned}
\{c, e\} &= \{E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}, E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d\} = -\{E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d, E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}\} \\
&= \frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ca} q^{db} + q^{da} q^{cb}) - k q^2 q^{ab} q^{ct_1} K_{t_1}^{i_1} E_{i_1}^d \\
&= \frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ca} q^{db} + q^{cb} q^{da}) - k q^3 q^{ab} K^{ci_1} E^{di_1} \quad (103)
\end{aligned}$$

$$\begin{aligned}
\{a, f\} &= \{(\det E)^{\frac{-2}{D-1}}, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \frac{-2}{D-1} \frac{1}{q} \frac{\{(\det E), K_{e_2}^{j_2}\}}{(\det E)} E_{j_2}^{e_2} = \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \frac{-2}{D-1} \frac{1}{q} E_{e_3}^{j_3} \{E_{j_3}^{e_3}, K_{e_2}^{j_2}\} E_{j_2}^{e_2} = E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \frac{-2}{D-1} \frac{1}{q} E_{e_3}^{j_3} \frac{k}{2} \delta_{j_3}^{j_2} \delta_{e_3}^{e_2} E_{j_2}^{e_2} = \\
&\frac{-k}{D-1} q q^{cd} \frac{1}{q} E_{e_2}^{j_2} E_{j_2}^{e_2} = \frac{-k}{D-1} q^{cd} D \quad (104)
\end{aligned}$$

The bracket $\{c, d\}$ can be calculated from (104) by changing the sign and making the following index replacement:

$$c \rightarrow a, d \rightarrow b, a \rightarrow c, b \rightarrow d \quad (105)$$

We obtain:

$$\{c, d\} = \{E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}, (\det E)^{\frac{-2}{D-1}}\} = \frac{k}{D-1} q^{ab} D \quad (106)$$

Finally we need to calculate the last bracket $\{c, f\}$:

$$\begin{aligned}
\{c, f\} &= \{E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = \\
&E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} \{K_{t_2}^{i_2}, E_{m_2}^c\} E_{i_2}^{t_2} E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} E_{m_2}^c \{K_{t_2}^{i_2}, E_{p_4}^d \eta^{p_4 m_2}\} E_{i_2}^{t_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
&E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} \{K_{t_2}^{i_2}, E_{j_2}^{e_2}\} E_{i_2}^{t_2} + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \{E_{k_2}^a, K_{e_2}^{j_2}\} E_{j_2}^{e_2} E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2} + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_2}^a \{E_{p_2}^b \eta^{p_2 k_2}, K_{e_2}^{j_2}\} E_{j_2}^{e_2} K_{t_2}^{i_2} E_{i_2}^{t_2} + \\
&E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} \{E_{i_2}^{t_2}, K_{e_2}^{j_2}\} E_{j_2}^{e_2} \quad (107)
\end{aligned}$$

or

$$\begin{aligned}
\{c, f\} = & \{E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = \\
& E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} E_{i_2}^{t_2} \left(-\frac{k}{2} \delta_{m_2}^{i_2} \delta_{t_2}^c\right) E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
& E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} E_{i_2}^{t_2} E_{m_2}^c \left(-\frac{k}{2} \delta_{p_4}^{i_2} \delta_{t_2}^d\right) \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2} + \\
& E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} E_{i_2}^{t_2} E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} \left(-\frac{k}{2} \delta_{j_2}^{i_2} \delta_{t_2}^{e_2}\right) + \\
& E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} \left(\frac{k}{2} \delta_{k_2}^{j_2} \delta_{e_2}^a\right) E_{j_2}^{e_2} E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2} + \\
& E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_2}^a \left(\frac{k}{2} \delta_{p_2}^{j_2} \delta_{e_2}^b\right) E_{j_2}^{e_2} \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2} + \\
& E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} \left(\frac{k}{2} \delta_{e_2}^{t_2} \delta_{i_2}^{j_2}\right) E_{j_2}^{e_2} = \\
& \frac{k}{2} (-q^{ab} q q^{cd} q K_{e_2}^{j_2} E_{j_2}^{e_2} - q^{ab} q q^{cd} q K_{e_2}^{j_2} E_{j_2}^{e_2} - q^{ab} q q^{cd} q K_{e_2}^{j_2} E_{j_2}^{e_2} \\
& + q^{cd} q q^{ab} q K_{t_2}^{i_2} E_{i_2}^{t_2} + q^{cd} q q^{ab} q K_{t_2}^{i_2} E_{i_2}^{t_2} + q^{cd} q q^{ab} q K_{t_2}^{i_2} E_{i_2}^{t_2}) = 0 \quad (108)
\end{aligned}$$

Thus

$$\{c, f\} = \{E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}, E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}\} = 0 \quad (109)$$

To summarize we have obtained:

$$\begin{aligned}
\{b, d\} &= \frac{k}{D-1} q^{ab} \\
\{c, d\} &= \frac{k}{D-1} q^{ab} D \\
\{a, e\} &= \frac{-k}{D-1} q^{cd} \\
\{a, f\} &= \frac{-k}{D-1} q^{cd} D \\
\{b, e\} &= \frac{k}{2} q q^{ac} G^{db} \\
\{c, e\} &= \frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ca} q^{db} + q^{da} q^{cb}) - k q^3 q^{ab} K^{ci_1} E^{di_1} \\
\{b, f\} &= -\frac{k q^2}{2} K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + k q^3 q^{cd} K^{ai_1} E^{bi_1} \\
\{c, f\} &= 0
\end{aligned}$$

By substituting into (88) :

$$\begin{aligned}
\{P^{ab}(x), P^{cd}(y)\} &= (a(\{b, d\} - \{c, d\}))(e - f) + \\
& d(\{a, e\} - \{a, f\})(b - c) + \\
& ad\{b, e\} - ad\{c, e\} - ad\{b, f\} + ad\{c, f\} \quad (110)
\end{aligned}$$

we obtain:

$$\begin{aligned}
\{P^{ab}(x), P^{cd}(y)\} = & \frac{1}{q} \left(\frac{k}{D-1} q^{ab} - \frac{k}{D-1} q^{ab} D \right) (E_{m_1}^c E_{p_3}^{e_1} \eta^{p_3 m_1} K_{e_1}^{j_1} E_{j_1}^d - E_{m_2}^c E_{p_4}^d \eta^{p_4 m_2} K_{e_2}^{j_2} E_{j_2}^{e_2}) + \\
& \frac{1}{q} \left(\frac{-k}{D-1} q^{cd} - \frac{-k}{D-1} q^{cd} D \right) (E_{k_1}^a E_{p_1}^{t_1} \eta^{p_1 k_1} K_{t_1}^{i_1} E_{i_1}^b - E_{k_2}^a E_{p_2}^b \eta^{p_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}) \\
& + \frac{k}{2} \frac{1}{q^2} q^3 q^{ac} G^{db} \\
& - \frac{k}{2q^2} (q^2 K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ca} q^{db} + q^{da} q^{cb}) - 2q^2 q^{ab} q^{ct_1} K_{t_1}^{i_1} E_{i_1}^d) \\
& - \frac{k}{2q^2} (-q^2 K_{e_2}^{j_2} E_{j_2}^{e_2} (q^{ac} q^{bd} + q^{ad} q^{bc}) + 2q^2 q^{cd} q^{at_1} K_{t_1}^{i_1} E_{i_1}^b) \quad (111)
\end{aligned}$$

$$\begin{aligned}
\{P^{ab}(x), P^{cd}(y)\} = & \frac{1}{q} (qq^{ce_1} K_{e_1}^{j_1} E_{j_1}^d - qq^{cd} K_{e_2}^{j_2} E_{j_2}^{e_2}) (-kq^{ab}) + \frac{1}{q} (qq^{at_1} K_{t_1}^{i_1} E_{i_1}^b - qq^{ab} K_{t_2}^{i_2} E_{i_2}^{t_2}) (kq^{cd}) \\
& + 2kqq^{ac} G^{db} \\
& + \frac{k}{q^2} (q^2 q^{ab} K^{ci_1} E_{i_1}^d - q^2 q^{cd} K^{ai_1} E_{i_1}^b) \quad (112)
\end{aligned}$$

$$\begin{aligned}
\{P^{ab}(x), P^{cd}(y)\} = & -kq^{ab} K^{cj_1} E_{j_1}^d + kq^{cd} K^{ai_1} E_{i_1}^b + 2kqq^{ac} G^{db} \\
& + k(q^{ab} K^{ci_1} E_{i_1}^d - q^{cd} K^{ai_1} E_{i_1}^b) = 2kqq^{ac} G^{bd} \quad (113)
\end{aligned}$$

$$\{P^{ab}(x), P^{cd}(y)\} = 2kqq^{ac} G^{bd} \quad (114)$$

Coordinate-Momentum Poisson Bracket

We mark each line by the label L(line number) and provide the detailed comments underneath the formula on how we move from one line to the next in our calculations.

$$P^{ab}(x) = -|\det(E_e^c)|^{-2/(D-1)} (E_{k_1}^a E_{m_1}^{t_1} \eta^{m_1 k_1} K_{t_1}^{i_1} E_{i_1}^b - E_{k_2}^a E_{m_2}^b \eta^{m_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}) \quad (115)$$

$$q_{cd}(y) = E_c^{i_3} E_d^{j_3} \eta_{i_3 j_3} (-|\det(E_e^a)|^{2/(D-1)}) \quad (116)$$

$$\begin{aligned}
\text{L1: } \{P^{ab}(x), q_{cd}(y)\} &= |\det(E_e^c)|^{-2/(D-1)} \{ (E_{k_1}^a E_{m_1}^{t_1} \eta^{m_1 k_1} K_{t_1}^{i_1} E_{i_1}^b - E_{k_2}^a E_{m_2}^b \eta^{m_2 k_2} K_{t_2}^{i_2} E_{i_2}^{t_2}), \\
&\quad E_c^{i_3} E_d^{j_3} \eta_{i_3 j_3} (\det(E))^{2/(D-1)} \} \\
\text{L2: } &= \frac{1}{q} \{ (E_{k_1}^a E_{m_1}^t \eta^{m_1 k_1} K_{t_1}^{i_1} q q^{be} E_e^{j_1} \eta_{i_1 j_1} - E_{k_2}^a E_{m_2}^b \eta^{m_2 k_2} K_{t_2}^{i_2} q q^{t_2 e} E_e^{j_2} \eta_{i_2 j_2}), E_c^{i_3} E_d^{j_3} \eta_{i_3 j_3} (\det(E))^{2/(D-1)} \} \\
&= \frac{1}{q} q^2 (q^{at} q^{be} - q^{ab} q^{te}) E_e^j \eta_{ij} \{ K_t^i, E_c^{i_3} E_d^{j_3} \eta_{i_3 j_3} (\det(E))^{2/(D-1)} \} \\
\text{L3: } &= q (q^{at} q^{be} - q^{ab} q^{te}) E_e^j \eta_{ij} (q \{ K_t^i E_c^{i_3} \} E_d^{j_3} \eta_{i_3 j_3} + q \{ K_t^i E_d^{j_3} \} E_c^{i_3} \eta_{i_3 j_3} + \frac{2}{D-1} \frac{q_{cd}}{q} q \frac{\{ K_t^i, \det(E) \}}{\det(E)}) = \\
\text{L4: } & q (q^{at} q^{be} - q^{ab} q^{te}) E_e^j \eta_{ij} (q (-E_c^j) \{ K_t^i, E_j^p \} E_p^{i_3} E_d^{j_3} \eta_{i_3 j_3} + q (-E_d^j) \{ K_t^i, E_j^p \} E_p^{i_3} E_c^{i_3} \eta_{i_3 j_3} + \\
&\quad \frac{2}{D-1} q_{cd} \{ K_t^i, E_j^m \} E_m^j) = \\
\text{L5: } & q (q^{at} q^{be} - q^{ab} q^{te}) E_e^j \eta_{ij} (q (-E_c^j) (-\frac{k}{2} \delta_j^i \delta_t^p) E_p^{i_3} E_d^{j_3} \eta_{i_3 j_3} + q (-E_d^j) (-\frac{k}{2} \delta_j^i \delta_t^p) E_p^{i_3} E_c^{i_3} \eta_{i_3 j_3} + \\
&\quad \frac{2}{D-1} q_{cd} (-\frac{k}{2} \delta_j^i \delta_t^m) E_m^j) = \\
\text{L6: } & k q (q^{at} q^{be} - q^{ab} q^{te}) E_e^j \eta_{ij} (q E_c^i E_t^{i_3} E_d^{j_3} \eta_{i_3 j_3} + q E_d^i E_t^{j_3} E_c^{i_3} \eta_{i_3 j_3} - \frac{2}{D-1} q_{cd} E_t^i) = \\
\text{L7: } & k q (q^{at} q^{be} - q^{ab} q^{te}) (q \frac{q_{ec}}{q} \frac{q_{dt}}{q} + q \frac{q_{ed}}{q} \frac{q_{ct}}{q} - \frac{2}{D-1} q_{cd} \frac{q_{et}}{q}) = \\
\text{L8: } & k (q^{at} q^{be} - q^{ab} q^{te}) (q_{ec} q_{dt} + q_{ed} q_{ct} - \frac{2}{D-1} q_{cd} q_{et}) = \\
\text{L9: } & k (\delta_c^b \delta_d^a + \delta_d^b \delta_c^a) - 2 q^{ab} q_{cd} q - \frac{2}{D-1} (q_{cd} q^{ab} - D q_{cd} q^{ab}) = \\
\text{L10: } & k (\delta_c^b \delta_d^a + \delta_d^b \delta_c^a) - 2 q^{ab} q_{cd} q - \frac{2}{D-1} (1-D) q_{cd} q^{ab} = \\
\text{L11: } & k (\delta_c^b \delta_d^a + \delta_d^b \delta_c^a) - 2 q^{ab} q_{cd} q + 2 q^{ab} q_{cd} q = \\
\text{L12: } & k \delta_{(c}^b \delta_{d)}^a \delta(x, y) \quad (117)
\end{aligned}$$

$$\{P^{ab}(x), q_{cd}(y)\} = k \delta_{(c}^b \delta_{d)}^a \delta(x, y) \quad (118)$$

,where

in the line L2: we used $E_i^a E_j^b \eta^{ij} = q q^{ab}$ and $q := (\det(E))^{2/(D-1)}$

in the line L3: we used Leibniz rule and $\{E_j^a(x), E_k^b(y)\} = 0$

in the line L4: we used: $\delta E_a^i = -E_a^i \delta E_i^b E_b^i$ and $[\delta(E)]/\det(E) = E_d^j \delta E_j^a$

in the line L5: we calculated the Poisson brackets: $\{E_i^a(x), K_b^j(y)\} = \frac{k}{2} \delta_b^a \delta_i^j \delta(x, y)$

in the line L7: we used: $E_e^i E_c^j \eta_{ij} = q_{ec}/q$, etc,

in the line L9: we have opened the parentheses and used: $q^{at} q_{td} = \delta_d^a$ and $q^{et} q_{et} = D$

in the line L10: $D-1$ cancels.

in the line L11: the last two terms are the same and mutually cancel.

14 Appendix D Timelike $SO(2, 1)$ Riemann Curvature Expression in $SO(2, 1)$ case

In this Appendix we derive the following two formulas:

$$R_{ab}^i = 2\partial_{[a}\Gamma_{b]}^i + \bar{\eta}^{ij}\epsilon_{jkl\ so(3)}\Gamma_a^k\Gamma_b^l \quad (119)$$

and

$$^{(\beta)}F_{ab}^i = 2\partial_{[a}^{(\beta)}A_{b]}^i + \bar{\eta}^{ij}\epsilon_{jkl\ so(3)}^{(\beta)}A_a^kA_b^l \quad (120)$$

, where $\bar{\eta}_{ij} = \text{Diag}(1, 1, 1)$ in $SO(3)$ case, and $\bar{\eta}_{ij} = \text{Diag}(-1, 1, 1)$ in $SO(2, 1)$ case. We begin with the curvature tensor definition:

$$R_{bcd}^a = \partial_b\Gamma_{cd}^a - \partial_c\Gamma_{bd}^a + \Gamma_{bt}^a\Gamma_{cd}^t - \Gamma_{ct}^a\Gamma_{bd}^t \quad (121)$$

$SO(3)$ case:

In $SO(3)$ case (119) becomes:

$$R_{ab}^i = 2\partial_{[a}\Gamma_{b]}^i + \epsilon_{ikl\ so(3)}^i\Gamma_a^k\Gamma_b^l \quad (122)$$

We then use $so(3) \rightarrow R^3$ isomorphism:

$$\Gamma_{ai}^l = \epsilon_{ki\ so(3)}^l\Gamma_a^k \quad (123)$$

By using the torsion-free condition: $\Gamma_{ai}^l = \Gamma_{ia}^l$, we can write it also as:

$$\Gamma_{ai}^l = \Gamma_{ia}^l = \epsilon_{ki\ so(3)}^l\Gamma_a^k = \epsilon_{ka\ so(3)}^l\Gamma_i^k \quad (124)$$

By substituting (124) into (121), we obtain:

$$R_{bcd}^a = \partial_b(\epsilon_{pd\ so(3)}^a\Gamma_c^p) - \partial_c(\epsilon_{pd\ so(3)}^a\Gamma_b^p) + \epsilon_{pt\ so(3)}^a\Gamma_b^p\epsilon_{sd\ so(3)}^t\Gamma_c^s - \epsilon_{mt\ so(3)}^a\Gamma_c^m\epsilon_{ld\ so(3)}^t\Gamma_b^l \quad (125)$$

We rewrite the third term by using the antisymmetric tensor properties:

$$\epsilon_{pt\ so(3)}^a\epsilon_{sd\ so(3)}^t = \epsilon^{tap\ so(3)}\epsilon_{tsd\ so(3)} = 2\delta_s^{[a}\delta_d^{p]} = \delta_s^a\delta_d^p - \delta_s^p\delta_d^a \quad (126)$$

and the fourth term by:

$$\epsilon_{mt\ so(3)}^a\epsilon_{ld\ so(3)}^t = \epsilon^{tam\ so(3)}\epsilon_{tld\ so(3)} = 2\delta_l^{[a}\delta_d^{m]} = \delta_l^a\delta_d^m - \delta_l^m\delta_d^a \quad (127)$$

we obtain:

$$\begin{aligned} R_{bcd}^a &= \epsilon_{pd\ so(3)}^a(\partial_b\Gamma_c^p - \partial_c\Gamma_b^p) + (\delta_s^a\delta_d^p - \delta_s^p\delta_d^a)\Gamma_b^p\Gamma_c^s - (\delta_l^a\delta_d^m - \delta_l^m\delta_d^a)\Gamma_c^m\Gamma_b^l = \\ &= \epsilon_{pd\ so(3)}^a(\partial_b\Gamma_c^p - \partial_c\Gamma_b^p) + (\Gamma_c^a\Gamma_b^d - \Gamma_b^s\Gamma_c^s) - (\Gamma_c^d\Gamma_b^a - \Gamma_b^l\Gamma_c^l) = \\ &= \epsilon_{pd\ so(3)}^a(\partial_b\Gamma_c^p - \partial_c\Gamma_b^p) + \Gamma_c^a\Gamma_b^d - \Gamma_b^a\Gamma_c^d \quad (128) \end{aligned}$$

By renaming index $p \rightarrow i$:

$$R_{bcd}^a = \epsilon_{ido(3)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) + \Gamma_b^d \Gamma_c^a - \Gamma_b^a \Gamma_c^d \quad (129)$$

In order to prove formula (122) we will show that if we take R_{bc}^i in that form and contract it with $\epsilon_{ido(3)}^a$ we will obtain R_{bcd}^a as in (129):

$$\begin{aligned} \epsilon_{ido(3)}^a R_{bc}^i &= \epsilon_{ido(3)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) + \epsilon_{ido(3)}^a \epsilon_{klso(3)}^i \Gamma_b^k \Gamma_c^l = \\ &= \epsilon_{ido(3)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) - \epsilon_{ido(3)}^a \epsilon_{klso(3)}^i \Gamma_b^k \Gamma_c^l = \\ &= \epsilon_{ido(3)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) - 2\delta_k^{[a} \delta_l^{d]} \Gamma_b^k \Gamma_c^l = \\ &= \epsilon_{ido(3)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) - \Gamma_b^a \Gamma_c^d + \Gamma_b^d \Gamma_c^a \quad (130) \end{aligned}$$

By comparing it to (129) we see that it equals to R_{bcd}^a . So, we have proved that:

$$R_{bcd}^a = \epsilon_{ido(3)}^a R_{bc}^i \quad (131)$$

, where $R_{ab}^i = 2\partial_{[a} \Gamma_{b]}^i + \epsilon_{klso(3)}^i \Gamma_a^k \Gamma_b^l$

Let us now consider $SO(2, 1)$ case:

We use the $so(2, 1) \rightarrow R_{2,1}^3$ isomorphism:

$$\Gamma_{ai}^l = \epsilon_{kiso(2,1)}^l \Gamma_a^k \quad (132)$$

So we are getting $SO(2, 1)$ case from $SO(3)$ case very easily by replacing everywhere $\epsilon_{kiso(3)}^l$ tensors with $\epsilon_{kiso(2,1)}^l$. The only difference that we will encounter is a different sign compared to (126) and (127):

$$\epsilon_{ptso(2,1)}^a \epsilon_{sdso(2,1)}^t = \epsilon^{tap}_{so(2,1)} \epsilon_{tsdso(2,1)} = -2\delta_s^{[a} \delta_d^{p]} = -(\delta_s^a \delta_d^p - \delta_s^p \delta_d^a) \quad (133)$$

and the fourth term by:

$$\epsilon_{mtso(2,1)}^a \epsilon_{ldso(2,1)}^t = \epsilon^{tam}_{so(2,1)} \epsilon_{tldso(2,1)} = -2\delta_l^{[a} \delta_d^{m]} = -(\delta_l^a \delta_d^m - \delta_l^m \delta_d^a) \quad (134)$$

It only changes the sign of the last two terms in the $so(2, 1)$ analog of (129) and of (130)

$$R_{bcd}^a = \epsilon_{ido(2,1)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) + \Gamma_b^a \Gamma_c^d - \Gamma_b^d \Gamma_c^a \quad (135)$$

$$\epsilon_{ido(2,1)}^a R_{bc}^i = \epsilon_{ido(2,1)}^a (\partial_b \Gamma_c^i - \partial_c \Gamma_b^i) + \Gamma_b^a \Gamma_c^d - \Gamma_b^d \Gamma_c^a \quad (136)$$

So again we obtain:

$$R_{bcd}^a = \epsilon_{ido(2,1)}^a R_{bc}^i \quad (137)$$

, where $R_{ab}^i = 2\partial_{[a} \Gamma_{b]}^i + \epsilon_{klso(2,1)}^i \Gamma_a^k \Gamma_b^l$

By combining two cases and remembering that we can express: $\epsilon_{klso(2,1)}^i = \bar{\eta}^{ij} \epsilon_{jklso(3)}$ we can write it in a general case:

$$R_{ab}^i = 2\partial_{[a} \Gamma_{b]}^i + \bar{\eta}^{ij} \epsilon_{jklso(3)} \Gamma_a^k \Gamma_b^l \quad (138)$$

, which finally proves formula (119). In addition the general form of (137) is as follows:

$$R_{bcd}^a = \bar{\eta}^{at} \epsilon_{tid\,so(3)} R_{bc}^i \quad (139)$$

If we use generalized Sen-Ashtekar connection from (30):

$${}^\beta A_a^k = \Gamma_a^k + ({}^\beta K_a^k), \quad {}^\beta A_{ai}^l = \bar{\eta}_{ij} \epsilon_{k\,so(3)}^{jl} A_a^k \quad (140)$$

and repeat all the steps using A_a^k instead of Γ_a^k , we will obtain (120).

15 Appendix E Expressing ${}^{(\beta)}F_{ab}^j$ via R_{ab}^j

$$R_{ab}^i = 2\partial_{[a}\Gamma_{b]}^i + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} \Gamma_a^k \Gamma_b^l \quad (141)$$

We want to prove that:

$${}^{(\beta)}F_{ab}^i = 2\partial_{[a}{}^{(\beta)}A_{b]}^i + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} {}^{(\beta)}A_a^k {}^{(\beta)}A_b^l \quad (142)$$

Proof:

We remind that ${}^{(\beta)}E_i^b = E_i^b/\beta$ and ${}^{(\beta)}K_a^k = \beta K_a^k$

By substituting ${}^{(\beta)}A_a^k = \Gamma_a^k + \beta K_a^k$ into (142) we obtain:

$$\begin{aligned} {}^{(\beta)}F_{ab}^i &= 2\partial_{[a}\Gamma_{b]}^i + 2\partial_{[a}{}^{(\beta)}K_{b]}^i + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} (\Gamma_a^k + \beta K_a^k)(\Gamma_b^l + \beta K_b^l) = \\ &= (2\partial_{[a}\Gamma_{b]}^i + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} \Gamma_a^k \Gamma_b^l) + (2\partial_{[a}{}^{(\beta)}K_{b]}^i + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} \Gamma_a^k \beta K_b^l + \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} \Gamma_b^l \beta K_a^k) \\ &\quad + (\beta^2 \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} K_a^k K_b^l) = R_{ab}^i + 2\beta D_{[a}K_{b]}^i + \beta^2 \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} K_a^k K_b^l \end{aligned} \quad (143)$$

Therefore:

$${}^{(\beta)}F_{ab}^i = R_{ab}^i + 2\beta D_{[a}K_{b]}^i + \beta^2 \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} K_a^k K_b^l \quad (144)$$

16 Appendix F Timelike $SO(2, 1)$ Diffeomorphism Contraction Calculations

We would like to derive the following expression:

$${}^{(\beta)}F_{ab}^i {}^{(\beta)}E_i^b = \frac{R_{ab}^i E_i^b}{\beta} + 2D_{[a}K_{b]}^i E_i^b + \beta K_a^i G_i \quad (145)$$

We begin with (144):

$${}^{(\beta)}F_{ab}^i = R_{ab}^i + 2\beta D_{[a}K_{b]}^i + \beta^2 \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} K_a^k K_b^l \quad (146)$$

contracting it with ${}^{(\beta)}E_j^b$

$${}^{(\beta)}F_{ab}^i {}^{(\beta)}E_i^b = \frac{R_{ab}^i E_i^b}{\beta} + 2\beta D_{[a}K_{b]}^i \frac{E_i^b}{\beta} + \beta^2 \bar{\eta}^{ij} \epsilon_{jkl\,so(3)} K_a^k K_b^l \frac{E_i^b}{\beta} \quad (147)$$

by simplifying it we obtain:

$${}^{(\beta)}F_{ab}{}^{(\beta)}E_i^b = \frac{R_{ab}^i E_i^b}{\beta} + 2D_{[a}K_{b]}^i E_i^b + \beta \bar{\eta}^{ij} \epsilon_{jkl}{}_{so(3)} K_a^k K_b^l E_i^b \quad (148)$$

So, we only need to prove that the last term is a rotational constraint, i.e:

$$\bar{\eta}^{ij} \epsilon_{jkl}{}_{so(3)} K_a^k K_b^l E_i^b = K_a^i G_i \quad (149)$$

We do it in the following steps:

$$\begin{aligned} \bar{\eta}^{ij} \epsilon_{jkl}{}_{so(3)} K_a^k K_b^l E_i^b &= K_a^k (\epsilon_{jkl}{}_{so(3)} K_b^l E^{bj}) = K_a^k (\epsilon_{k}^{lj}{}_{so(3)} K_{lb} E_j^b) \\ &= K_a^k (\bar{\eta}_{km} \epsilon^{mlj}{}_{so(3)} K_{lb} E_j^b) = K_a^k G_k = K_a^i G_i \end{aligned} \quad (150)$$

,where in the first step above we used generalized metric to raise index j , in the next step we raised and lowered at the same time indices lj , and permutated twice $jkl \rightarrow klj$, so the sign stays the same, then we raised index m by using generalized metric, since in $so(3)$ case it is all the same, while in $so(2, 1)$ case even $\epsilon_{jkl}{}_{so(3)}$ indices should be lowered and raised by using Minkowski tensor η_{ij} . Finally we used the following rotational constraint expression obtained in (23):

$$G_i = \bar{\eta}_{ij} \epsilon^{jkl}{}_{so(3)} K_{ak} E_l^a \quad (151)$$

17 Appendix G Identity from Rotational Constraint

$$G_{ab} := K_{[a}^i E_{b]}^j \bar{\eta}_{ij} = 1/2 (K_{ai} E_b^i - K_{bi} E_a^i) = 0 \quad (152)$$

therefore:

$$K_{bi} E_a^i = K_{ai} E_b^i \quad (153)$$

or, by raising index i

$$K_b^i E_a^i = K_a^i E_b^i \quad (154)$$

it follows that

$$K_b^i E_i^a = K_a^i E_i^b \quad (155)$$

since

$$K_a^i E_i^b = K_a^i E_t^i q^{tb} q \quad (156)$$

,while

$$K_b^i E_i^a = K_b^i E_t^i q^{ta} q \quad (157)$$

and the right hand sides of (156) and (157) are equal, as

$$K_a^i E_t^i q^{tb} = K_b^i E_t^i q^{ta} \quad (158)$$

since

$$K_a^i = K_b^i q^{ta} q_{tb} = K_b^i \delta_b^a \quad (159)$$

18 Appendix I Contracting Riemann Curvature with Triads Second Term

We prove the following identity first in $SO(3)$ spacelike foliation ADM case:

$$2\beta D_{[a} K_{b]}^j \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b = -2^{(\beta)} E_k^a D_a G^k \quad (160)$$

it can be rewritten as:

$$\begin{aligned} 2\beta D_{[a} K_{b]}^j \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ \beta (D_a K_b^j - D_b K_a^j) \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ {}^{(\beta)} E_k^a (D_a (K_b^j E_l^b) - {}^{(\beta)} E_l^b D_b (K_a^j E_k^a)) \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ - {}^{(\beta)} E_k^a (D_a (K_b^j E_l^b) \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b - {}^{(\beta)} E_l^b D_b (K_a^j E_k^a) \epsilon_j^{lk} {}^{(\beta)} E_k^{a(\beta)} E_l^b) &= \\ - {}^{(\beta)} E_k^a D_a (G^k) - {}^{(\beta)} E_l^b D_b (G^l) &= \\ - 2^{(\beta)} E_k^a D_a (G^k) \end{aligned} \quad (161)$$

,where in the fourth line for the first and second terms we used rotational constraint definition (24): $G^k = \epsilon_j^{kl} K_b^j E_l^b$ and moved index k by one position for the first term: $\epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b = -\epsilon_j^{lk} {}^{(\beta)} E_k^{a(\beta)} E_l^b$ and index l by two positions for the second term $\epsilon_j^{lk} {}^{(\beta)} E_k^{a(\beta)} E_l^b = \epsilon_j^{kl} {}^{(\beta)} E_k^{a(\beta)} E_l^b$. It is easy to pass to generalized form:

$$2\beta D_{[a} K_{b]}^j \bar{\eta}_{ji} \epsilon^{ikl} {}^{(\beta)} E_k^{a(\beta)} E_l^b = -2^{(\beta)} E_k^a D_a G^k \quad (162)$$

Similar to the above:

$$\begin{aligned} 2\beta D_{[a} K_{b]}^j \bar{\eta}_{ji} \epsilon^{ikl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ \beta (D_a K_b^j - D_b K_a^j) \bar{\eta}_{ji} \epsilon^{ikl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ {}^{(\beta)} E_k^a (D_a (K_b^j E_l^b) - {}^{(\beta)} E_l^b D_b (K_a^j E_k^a)) \bar{\eta}_{ji} \epsilon^{ikl} {}^{(\beta)} E_k^{a(\beta)} E_l^b &= \\ - {}^{(\beta)} E_k^a (D_a (K_b^j E_l^b) \bar{\eta}_{ji} \epsilon^{kil} {}^{(\beta)} E_k^{a(\beta)} E_l^b - {}^{(\beta)} E_l^b D_b (K_a^j E_k^a) \bar{\eta}_{ji} \epsilon^{lik} {}^{(\beta)} E_k^{a(\beta)} E_l^b) &= \\ - {}^{(\beta)} E_k^a D_a (G^k) - {}^{(\beta)} E_l^b D_b (G^l) &= \\ - 2^{(\beta)} E_k^a D_a (G^k) \end{aligned} \quad (163)$$

, where we used generalized rotational constraint (24):

$$G_i = \bar{\eta}_{ij} \epsilon^{jkl} K_{ak} E_l^a = \bar{\eta}_{ij} \epsilon_k^{jl} K_a^k E_l^a = 0 \quad (164)$$

and

$$G_i = \bar{\eta}_{ij} \epsilon^{jkl} K_{ak} E_l^a = \bar{\eta}_{ij} \epsilon_k^{jl} K_a^k E_l^a = 0 \quad (165)$$

since $G_i = 0$, $G^k = \bar{\eta}^{ki} G_i = 0$

$$G^k = \bar{\eta}^{ki} G_i = K_b^j E_l^b \bar{\eta}_{ji} \epsilon^{kil} {}^{(\beta)} E_k^{a(\beta)} E_l^b = 0 \quad (166)$$

19 Appendix J Contracting Riemann Curvature with Triads Third Term

We would like to prove that in $SO(3)$ spacelike ADM case:

$$\beta^2 \epsilon_{mnso(3)}^j K_a^m K_b^n \epsilon_j^{kl} {}^{so(3)} E_k^{a(\beta)} E_l^b = (K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b) \quad (167)$$

The proof is straightforward. β cancels on the left hand side right away and we use $\epsilon_{mnso(3)}^j \epsilon_j^{kl} {}^{so(3)} = 2\delta_m^{[k} \delta_n^{l]}$:

$$\begin{aligned} \beta^2 \epsilon_{mnso(3)}^j K_a^m K_b^n \epsilon_j^{kl} {}^{so(3)} E_k^{a(\beta)} E_l^b &= K_a^m K_b^n 2\delta_m^{[k} \delta_n^{l]} E_k^a E_l^b = \\ &= (\delta_m^k \delta_n^l - \delta_n^k \delta_m^l) K_a^m K_b^n E_k^a E_l^b = (K_a^k K_b^l E_k^a E_l^b - K_a^l K_b^k E_k^a E_l^b) = \\ &= ((K_a^j E_j^a)^2 - (K_b^l E_l^a)(K_a^k E_k^b)) = ((K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b)) \end{aligned} \quad (168)$$

On the other hand, in $SO(2,1)$ case we have an opposite sign in:

$$\epsilon_{mnso(2,1)}^j \epsilon_j^{kl} {}^{so(2,1)} = -2\delta_k^{[m} \delta_l^{n]}:$$

Therefore in $SO(2,1)$ timelike ADM case the sign also becomes opposite:

$$\beta^2 \epsilon_{mnso(2,1)}^j K_a^m K_b^n \epsilon_j^{kl} {}^{so(2,1)} E_k^{a(\beta)} E_l^b = -((K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b)) \quad (169)$$

The generalized formula for both cases would look like this:

$$\beta^2 \bar{\eta}^{ji} \epsilon_{imnso(3)} K_a^m K_b^n \bar{\eta}_{jp} \epsilon^{pkl} {}^{so(2,1)} E_k^{a(\beta)} E_l^b = -s((K_a^j E_j^a)^2 - (K_b^j E_j^a)(K_a^k E_k^b)) \quad (170)$$

,where $s = -1$ in $SO(3)$ spacelike ADM case and $s = 1$ in $SO(2,1)$ timelike foliation ADM case, $\bar{\eta}^{ij}$ - generalized metric : $\text{Diag}(1,1,1)$ in spacelike and $\text{Diag}(-1,1,1)$ in timelike foliation cases.

20 Appendix K Contracting Riemann Curvature with one Triad

We need to prove that:

$$R_{ab}^i E_i^b = 0 \quad (171)$$

The proof for $SO(3)$ spacelike foliation can be found in [1] (4.2.35): The Bianchi identity can be written in the form:

$$\epsilon_{ijkso(3)} \epsilon^{efc} {}^{so(3)} R_{ef}^j e_c^k = 0 \rightarrow \frac{1}{2} \epsilon_{ijkso(3)} \epsilon^{efc} {}^{so(3)} R_{ef}^j e_c^k e_a^i = \frac{1}{2} E_j^b \epsilon_{cabso(3)} \epsilon^{efc} {}^{so(3)} R_{ae}^j = R_{ab}^j E_j^b = 0 \quad (172)$$

It is still true in $SO(2,1)$ timelike ADM foliation case, as what changes are antisymmetric tensors $\epsilon_{ijkso(3)} \rightarrow \epsilon_{ijkso(2,1)}$, and it causes only a sign change in the last term

when contracting $so(2,1)$ antisymmetric tensors: $\epsilon_{cabso(2,1)}\epsilon^{efc}_{so(2,1)} = -2\delta_a^{[e}\delta_b^{f]}$
instead of $so(3)$ version $\epsilon_{cabso(3)}\epsilon^{efc}_{so(3)} = 2\delta_a^{[e}\delta_b^{f]}$

$$\epsilon_{ijkso(2,1)}\epsilon^{efc}_{so(2,1)}R_{ef}^j e_c^k = 0 \rightarrow \frac{1}{2}\epsilon_{ijkso(2,1)}\epsilon^{efc}_{so(2,1)}R_{ef}^j e_c^k e_a^i = \frac{1}{2}E_j^b \epsilon_{cabso(2,1)}\epsilon^{efc}_{so(2,1)}R_{ae}^j = -R_{ab}^j E_j^b = 0 \quad (173)$$

References

- [1] T. Thiemann (2007). "Introduction to Modern Canonical Quantum General Relativity", Cambridge University Press, Dataset <https://doi.org/10.1017/CBO9780511755682>
- [2] Dubrovin, Fomenko, Novikov (1990). "Modern Geometry—Methods and Applications", Springer, Dataset <https://doi.org/10.1007/978-1-4684-9946-9>
- [3] S. Alexandrov, Z. Kadar (2005). "Timelike surfaces in Lorentz covariant loop gravity and spin foam models", Class. Quantum Gravity, vol 22, 17, Dataset <https://doi.org/10.1088/0264-9381/22/17/010>
- [4] S. Alexandrov (2000). "SO(4,C)-covariant Ashtekar–Barbero gravity and the Immirzi parameter," Class. Quantum Grav. 17, 4255. <https://doi.org/10.1088/0264-9381/17/20/307>
- [5] S. Alexandrov (2002). "On choice of connection in loop quantum gravity," Phys. Rev. D 65, 024011. <https://doi.org/10.1103/PhysRevD.65.024011>
- [6] H Liu, K. Noui (2017). "Gravity as an $su(1,1)$ gauge theory in four dimensions", Class.Quant.Grav. 34 no.13, 135008 <https://doi.org/10.1088/1361-6382/aa7348>
- [7] M. Montesinos, J. Romero, R. Escobedo, M. Celada (2018). "SU(1,1) Barbero-like variables derived from Holst action", Phys. Rev. D 98, 124002. <https://doi.org/10.1103/PhysRevD.98.124002>
- [8] J. F. Barbero (1994). "Real-polynomial formulation of general relativity in terms of connections", Phys. Rev. D 49, 6935. <https://doi.org/10.1103/PhysRevD.49.6935>
- [9] G. Immirzi. "Quantum gravity and Regge calculus" (1997) Nucl. Phys. Proc. Suppl.57, 65 (1997). [https://doi.org/10.1016/S0920-5632\(97\)00354-X](https://doi.org/10.1016/S0920-5632(97)00354-X)
- [10] S. Holst. (1996) "Barbero's Hamiltonian derived from a generalized Hilbert-Palatini action", Phys. Rev. D 53, 5966. <https://doi.org/10.1103/PhysRevD.53.5966>
- [11] T. Torii, H. Shinkai (2008). "N+1 formalism in Einstein-Gauss-Bonnet gravity", Phys. Rev. D 78 (08) 4037 <https://doi.org/10.1103/PhysRevD.78.084037>
- [12] C. Rovelli. (2004) "Quantum Gravity", Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9780511755804>