

# Contravariant Pseudo-Hessian manifolds and their associated Poisson structures

Abdelhak Abouqateb<sup>a</sup>, Mohamed Boucetta<sup>b</sup>, Charif Bourzik<sup>c</sup>

<sup>a</sup>Université Cadi-Ayyad  
Faculté des sciences et techniques  
BP 549 Marrakech Maroc  
e-mail: a.abouqateb@uca.ac.ma

<sup>b</sup>Université Cadi-Ayyad  
Faculté des sciences et techniques  
BP 549 Marrakech Maroc  
e-mail: m.boucetta@uca.ac.ma

<sup>c</sup>Université Cadi-Ayyad  
Faculté des sciences et techniques  
BP 549 Marrakech Maroc  
e-mail: bourzikcharif@gmail.com

---

## Abstract

A contravariant pseudo-Hessian manifold is a manifold  $M$  endowed with a pair  $(\nabla, h)$  where  $\nabla$  is a flat connection and  $h$  is a symmetric bivector field satisfying a contravariant Codazzi equation. When  $h$  is invertible we recover the known notion of pseudo-Hessian manifold. Contravariant pseudo-Hessian manifolds have properties similar to Poisson manifolds and, in fact, to any contravariant pseudo-Hessian manifold  $(M, \nabla, h)$  we associate naturally a Poisson tensor on  $TM$ . We investigate these properties and we study in details many classes of such structures in order to highlight the richness of the geometry of these manifolds.

**Keywords:** Affine manifolds, Poisson manifolds, pseudo-Hessian manifolds, Associative commutative algebras

**2000 MSC:** 53A15,

**2000 MSC:** 53D17,

**2000 MSC:** 17D25

---

## 1. Introduction

A contravariant pseudo-Hessian manifold is an affine manifold  $(M, \nabla)$  endowed with a symmetric bivector field  $h$  such that, for any  $\alpha, \beta, \gamma \in \Omega^1(M)$ ,

$$(\nabla_{h_\#(\alpha)}h)(\beta, \gamma) = (\nabla_{h_\#(\beta)}h)(\alpha, \gamma), \quad (1.1)$$

where  $h_\# : T^*M \longrightarrow TM$  is the contraction. We will refer to (1.1) as contravariant Godazzi equation. These manifolds were introduced in [3] as a generalization of pseudo-Hessian manifolds. Recall that a pseudo-Hessian manifold is an affine manifold  $(M, \nabla)$  with a pseudo-Riemannian metric  $g$  satisfying the Godazzi equation

$$\nabla_X g(Y, Z) = \nabla_Y g(X, Z), \quad (1.2)$$

for any  $X, Y, Z \in \Gamma(TM)$ . The book [13] is devoted to the study of Hessian manifolds which are pseudo-Hessian manifolds with a Riemannian metric.

In this paper, we study contravariant pseudo-Hessian manifolds. The passage from pseudo-Hessian manifolds to contravariant pseudo-Hessian manifolds is similar to the passage from symplectic manifolds to Poisson manifolds and this similarity will guide our study. Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold. We will show that  $T^*M$  has a Lie algebroid structure,  $M$  has a singular foliation whose leaves are pseudo-Hessian manifolds and  $TM$  has a Poisson tensor whose symplectic leaves are pseudo-Kählerian manifolds. We investigate an analog of Darboux-Weinstein's theorem and we show that it is not true in general but holds in some cases. We will study in details the correspondence which maps a contravariant pseudo-Hessian bivector field on  $(M, \nabla)$  to a Poisson bivector field on  $TM$ . We study affine, linear and quadratic contravariant pseudo-Hessian structures on vector spaces and we show that an affine contravariant pseudo-Hessian structure on a vector space  $V$  is equivalent to an associative commutative algebra product and a 2-cocycle on  $V^*$ . We study right invariant contravariant pseudo-Hessian structures on a Lie group  $G$  and we show that  $TG$  has a structure of Lie group (different from the one associated to the adjoint action) for which the associated Poisson tensor is right invariant. We show that a right invariant contravariant pseudo-Hessian structure on a Lie group is equivalent to a  $S$ -matrix on the associated left symmetric algebra (see [2, 4]) and we associate to any  $S$ -matrix on a left symmetric algebra  $\mathfrak{g}$  a solution of the classical Yang-Baxter equation on  $\mathfrak{g} \times \mathfrak{g}$ . Finally, we show that an action of a left symmetric algebra  $\mathfrak{g}$  on an affine manifold  $(M, \nabla)$  transforms a  $S$ -matrix on  $\mathfrak{g}$  to a contravariant pseudo-Hessian bivector field on  $(M, \nabla)$ . Since the Lie algebra of affine vector fields of  $(M, \nabla)$  has a natural structure of finite dimensional associative algebra, we have a mean to define contravariant pseudo-Hessian structures on any affine manifold. The paper contains many examples of contravariant pseudo-Hessian structures.

The paper is organized as follows. In Section 2, we give the definition of a contravariant pseudo-Hessian manifold and we investigate its properties. In Section 3, we study in details the Poisson structure of the tangent bundle of a contravariant pseudo-Hessian manifold. Section 4 is devoted to the study of linear and affine contravariant pseudo-Hessian structures. Quadratic contravariant pseudo-Hessian structures will be studied in Section 5. In Section 6, we study right invariant pseudo-Hessian structures on Lie groups.

## 2. Contravariant pseudo-Hessian manifolds: definition and principal properties

### 2.1. Definition of a contravariant pseudo-Hessian manifold

Recall that an affine manifold is a  $n$ -manifold  $M$  endowed with a maximal atlas such that all transition functions are restrictions of elements of the affine group  $\text{Aff}(\mathbb{R}^n)$ . This is equivalent to the existence on  $M$  of a flat connection  $\nabla$ , i.e., torsionless and with vanishing curvature (see [13] for more details). An affine coordinates system on an affine manifold  $(M, \nabla)$  is a coordinates system  $(x_1, \dots, x_n)$  satisfying  $\nabla \partial_{x_i} = 0$  for any  $i = 1, \dots, n$ .

Let  $g$  be a pseudo-Riemannian metric on an affine manifold  $(M, \nabla)$ . The triple  $(M, \nabla, g)$  is called a *pseudo-Hessian manifold* if  $g$  can be locally expressed in any affine coordinates system  $(x_1, \dots, x_n)$  as

$$g_{ij} = \frac{\partial^2 \phi}{\partial x_i \partial x_j}.$$

That is equivalent to  $g$  satisfying the *Codazzi equation* (1.2). When  $g$  is Riemannian, we call  $(M, \nabla, g)$  a Hessian manifold. The geometry of Hessian manifolds was studied intensively in [13].

We consider now a more general situation.

**Definition 2.1** ([3]). *Let  $h$  be a symmetric bivector field on an affine manifold  $(M, \nabla)$  and  $h_{\sharp} : T^*M \rightarrow TM$  the associated contraction given by  $\beta(h_{\sharp}(\alpha)) = h(\alpha, \beta)$ . The triple  $(M, \nabla, h)$  is called a contravariant pseudo-Hessian manifold if  $h$  satisfies the contravariant Codazzi equation*

$$(\nabla_{h_{\sharp}(\alpha)} h)(\beta, \gamma) = (\nabla_{h_{\sharp}(\beta)} h)(\alpha, \gamma), \quad (2.1)$$

for any  $\alpha, \beta, \gamma \in \Omega^1(M)$ . We call such  $h$  a pseudo-Hessian bivector field.

One can see easily that if  $(M, \nabla, g)$  is a pseudo-Hessian manifold then  $(M, \nabla, g^{-1})$  is a contravariant pseudo-Hessian manifold.

The following proposition is obvious and gives the local expression of the equation (2.1) in affine charts.

**Proposition 2.2.** *Let  $(M, \nabla, h)$  be an affine manifold endowed with a symmetric bivector field. Then  $h$  satisfies (2.1) if and only if, for any  $m \in M$ , there exists an affine coordinates system  $(x_1, \dots, x_n)$  around  $m$  such that for any  $1 \leq i < j \leq n$  and any  $k = 1, \dots, n$*

$$\sum_{l=1}^n [h_{il} \partial_{x_l}(h_{jk}) - h_{jl} \partial_{x_l}(h_{ik})] = 0, \quad (2.2)$$

where  $h_{ij} = h(dx_i, dx_j)$ .

**Example 2.3.** 1. Take  $M = \mathbb{R}^n$  endowed with its canonical affine structure and consider

$$h = \sum_{i=1}^n f_i(x_i) \partial_{x_i} \otimes \partial_{x_i},$$

where  $f_i : \mathbb{R} \rightarrow \mathbb{R}$  for  $i = 1, \dots, n$ . Then one can see easily that  $h$  satisfies (2.2) and hence defines a contravariant pseudo-Hessian structure on  $\mathbb{R}^n$ .

2. Take  $M = \mathbb{R}^n$  endowed with its canonical affine structure and consider

$$h = \sum_{i,j=1}^n x_i x_j \partial_{x_i} \otimes \partial_{x_j}.$$

Then one can see easily that  $h$  satisfies (2.2) and hence defines a contravariant pseudo-Hessian structure on  $\mathbb{R}^n$ .

3. Let  $(M, \nabla)$  be an affine manifold,  $(X_1, \dots, X_r)$  a family of parallel vector fields and  $(a_{i,j})_{1 \leq i,j \leq n}$  a symmetric  $n$ -matrix. Then

$$h = \sum_{i,j} a_{i,j} X_i \otimes X_j$$

defines a contravariant pseudo-Hessian structure on  $M$ .

## 2.2. The Lie algebroid of a contravariant pseudo-Hessian manifold

We show that associated to any contravariant pseudo-Hessian manifold there is a Lie algebroid structure on its cotangent bundle and a Lie algebroid flat connection. The reader can consult [10, 12] for more details on Lie algebroids and their connections.

Let  $(M, \nabla, h)$  be an affine manifold endowed with a symmetric bivector field. We associate to this triple a bracket on  $\Omega^1(M)$  by putting

$$[\alpha, \beta]_h := \nabla_{h_\#(\alpha)}\beta - \nabla_{h_\#(\beta)}\alpha, \quad (2.3)$$

and a map  $\mathcal{D} : \Omega^1(M) \times \Omega^1(M) \longrightarrow \Omega^1(M)$  given by

$$\langle \mathcal{D}_\alpha \beta, X \rangle := (\nabla_X h)(\alpha, \beta) + \langle \nabla_{h_\#(\alpha)}\beta, X \rangle, \quad (2.4)$$

for any  $\alpha, \beta \in \Omega^1(M)$  and  $X \in \Gamma(TM)$ . This bracket is skew-symmetric and satisfies obviously

$$[\alpha, \beta]_h = \mathcal{D}_\alpha \beta - \mathcal{D}_\beta \alpha \quad \text{and} \quad [\alpha, f\beta]_h = f[\alpha, \beta]_h + h_\#(\alpha)(f)\beta,$$

where  $f \in C^\infty(M)$ ,  $\alpha, \beta \in \Omega^1(M)$ .

**Theorem 2.4.** *With the hypothesis and notations above, the following assertions are equivalent:*

- (i)  *$h$  is a pseudo-Hessian bivector field.*
- (ii)  *$(T^*M, h_\#, [\ , \ ]_h)$  is a Lie algebroid.*

*In this case,  $\mathcal{D}$  is a connection for the Lie algebroid structure  $(T^*M, h_\#, [\ , \ ]_h)$  satisfying*

$$h_\#(\mathcal{D}_\alpha \beta) = \nabla_{h_\#(\alpha)}h_\#(\beta) \quad \text{and} \quad R_{\mathcal{D}}(\alpha, \beta) := \mathcal{D}_{[\alpha, \beta]_h} - \mathcal{D}_\alpha \mathcal{D}_\beta + \mathcal{D}_\beta \mathcal{D}_\alpha = 0,$$

*for any  $\alpha, \beta \in \Omega^1(M)$ .*

*Proof.* According to [3, Proposition 2.1],  $(T^*M, h_\#, [\ , \ ]_h)$  is a Lie algebroid if and only if, for any affine coordinates system  $(x_1, \dots, x_n)$ ,

$$h_\#([dx_i, dx_j]_h) = [h_\#(dx_i), h_\#(dx_j)] \quad \text{and} \quad \bigoplus_{i,j,k} [dx_i, [dx_j, dx_k]_h]_h = 0,$$

for  $1 \leq i < j < k \leq n$ . Since  $[dx_i, dx_j]_h = 0$  this is equivalent to  $[h_\#(dx_i), h_\#(dx_j)] = 0$  for any  $1 \leq i < j \leq n$  which is equivalent to (2.2).

Suppose now that (i) or (ii) holds. For any,  $\alpha, \beta, \gamma \in \Omega^1(M)$ ,

$$\begin{aligned} \langle \mathcal{D}_\alpha \beta, h_\#(\gamma) \rangle &= \nabla_{h_\#(\gamma)}h(\alpha, \beta) + h(\nabla_{h_\#(\alpha)}^*\beta, \gamma) \\ &= \nabla_{h_\#(\alpha)}h(\gamma, \beta) + h(\nabla_{h_\#(\alpha)}^*\beta, \gamma) \\ &= h_\#(\alpha).h(\beta, \gamma) - h(\nabla_{h_\#(\alpha)}^*\gamma, \beta) \\ &= \langle \gamma, \nabla_{h_\#(\alpha)}h_\#(\beta) \rangle. \end{aligned}$$

This shows that  $h_\#(\mathcal{D}_\alpha \beta) = \nabla_{h_\#(\alpha)}h_\#(\beta)$ .

Let us show now that the curvature of  $\mathcal{D}$  vanishes. Since  $[dx_i, dx_j]_h = 0$ , it suffices to show that, for any  $i, j, k \in \{1, \dots, n\}$  with  $i < j$ ,  $\mathcal{D}_{dx_i}\mathcal{D}_{dx_j}dx_k = \mathcal{D}_{dx_j}\mathcal{D}_{dx_i}dx_k$ . We have

$$\langle \mathcal{D}_{dx_i}dx_k, \frac{\partial}{\partial x_l} \rangle = \frac{\partial h_{ik}}{\partial x_l}$$

and hence

$$\mathcal{D}_{dx_i}dx_k = \sum_{l=1}^n \frac{\partial h_{ik}}{\partial x_l} dx_l$$

and then

$$\begin{aligned}
\mathcal{D}_{dx_j} \mathcal{D}_{dx_i} dx_k &= \sum_{l=1}^n \mathcal{D}_{dx_j} \left( \frac{\partial h_{ik}}{\partial x_l} dx_l \right) \\
&= \sum_{l=1}^n \left( h_{\#}(dx_j) \left( \frac{\partial h_{ik}}{\partial x_l} \right) dx_l + \frac{\partial h_{ik}}{\partial x_l} \left( \sum_{s=1}^n \frac{\partial h_{jl}}{\partial x_s} dx_s \right) \right) \\
&= \sum_{l,r} h_{jr} \left( \frac{\partial^2 h_{ik}}{\partial x_r \partial x_l} \right) dx_l + \sum_{s,l} \frac{\partial h_{ik}}{\partial x_l} \frac{\partial h_{jl}}{\partial x_s} dx_s \\
&= \sum_{l,r} h_{jr} \left( \frac{\partial^2 h_{ik}}{\partial x_r \partial x_l} \right) dx_l + \sum_{l,r} \frac{\partial h_{ik}}{\partial x_r} \frac{\partial h_{jr}}{\partial x_l} dx_l \\
&= \sum_{l,r} \left( h_{jr} \left( \frac{\partial^2 h_{ik}}{\partial x_r \partial x_l} \right) + \frac{\partial h_{ik}}{\partial x_r} \frac{\partial h_{jr}}{\partial x_l} \right) dx_l \\
&= \sum_l \frac{\partial}{\partial x_l} \left( \sum_r h_{jr} \frac{\partial h_{ik}}{\partial x_r} \right) dx_l.
\end{aligned}$$

So

$$\mathcal{D}_{dx_i} \mathcal{D}_{dx_j} dx_k - \mathcal{D}_{dx_j} \mathcal{D}_{dx_i} dx_k = d \left( \sum_r \left( h_{jr} \frac{\partial h_{ik}}{\partial x_r} - h_{ir} \frac{\partial h_{jk}}{\partial x_r} \right) \right) \stackrel{(2.2)}{=} d(0) = 0.$$

□

The following result is an important consequence of Theorem 2.4.

**Proposition 2.5.** ([4, Theorem 6.7]) *Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold. Then:*

1. *The distribution  $\text{Im}h_{\#}$  is integrable and defines a singular foliation  $\mathcal{L}$  on  $M$ .*
2. *For any leaf  $L$  of  $\mathcal{L}$ ,  $(L, \nabla|_L, g_L)$  is a pseudo-Hessian manifold where  $g_L$  is given by  $g_L(h_{\#}(\alpha), h_{\#}(\beta)) = h(\alpha, \beta)$ .*

*We will call the foliation defined by  $\text{Im}h_{\#}$  the affine foliation associated to  $(M, \nabla, h)$ .*

**Remark 2.6.** *This proposition shows that contravariant pseudo-Hessian bivector fields can be used either to build examples of affine foliations on affine manifolds or to build examples of pseudo-Hessian manifolds.*

For the reader familiar with Poisson manifolds what we have established so far shows the similarities between Poisson manifolds and contravariant pseudo-Hessian manifolds. One can consult [8] for more details on Poisson geometry. Poisson manifolds have many relations with Lie algebras and we will see now and in Section 4 that contravariant pseudo-Hessian manifolds are related to commutative associative algebras.

Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold and  $\mathcal{D}$  the connection given in (2.4). Let  $x \in M$  and  $\mathfrak{g}_x = \ker h_{\#}(x)$ . For any  $\alpha, \beta \in \Omega^1(M)$ ,  $h_{\#}(\mathcal{D}_{\alpha}\beta) = \nabla_{h_{\#}(\alpha)}h_{\#}(\beta)$ . This shows that if  $h_{\#}(\alpha)(x) = 0$  then  $h_{\#}(\mathcal{D}_{\alpha}\beta)(x) = 0$ . Moreover,  $\mathcal{D}_{\alpha}\beta - \mathcal{D}_{\beta}\alpha = \nabla_{h_{\#}(\alpha)}\beta - \nabla_{h_{\#}(\beta)}\alpha$ . This implies that if  $h_{\#}(\alpha)(x) = h_{\#}(\beta)(x) = 0$  then  $\mathcal{D}_{\alpha}\beta(x) = \mathcal{D}_{\beta}\alpha(x)$ . For any  $a, b \in \mathfrak{g}_x$  put

$$a \bullet b = (\mathcal{D}_{\alpha}\beta)(x),$$

where  $\alpha, \beta$  are two differential 1-forms satisfying  $\alpha(x) = a$  and  $\beta(x) = b$ . This defines a commutative product on  $\mathfrak{g}_x$  and moreover, by using the vanishing of the curvature of  $\mathcal{D}$ , we get:

**Proposition 2.7.**  $(\mathfrak{g}_x, \bullet)$  is a commutative associative algebra.

Near a point where  $h$  vanishes, the algebra structure of  $\mathfrak{g}_x$  can be made explicit.

**Proposition 2.8.** We consider  $\mathbb{R}^n$  endowed with its canonical affine connection,  $h$  a symmetric bivector field on  $\mathbb{R}^n$  such that  $h(0) = 0$  and  $(\mathbb{R}^n, \nabla, h)$  is a contravariant pseudo-Hessian manifold. Then the product on  $(\mathbb{R}^n)^*$  given by

$$e_i^* \bullet e_j^* = \sum_{k=1}^n \frac{\partial h_{ij}}{\partial x_k}(0) e_k^*$$

is associative and commutative.

*Proof.* It is a consequence of the relation  $\mathcal{D}_{dx_i} dx_j = dh_{ij}$  true by virtue of (2.4).  $\square$

### 2.3. The product of contravariant pseudo-Hessian manifolds and the splitting theorem

As the product of two Poisson manifolds is a Poisson manifold [15], the product of two contravariant pseudo-Hessian manifolds is a contravariant pseudo-Hessian manifold.

Let  $(M_1, \nabla^1, h^1)$  and  $(M_2, \nabla^2, h^2)$  be two contravariant pseudo-Hessian manifolds. We denote by  $p_i : M = M_1 \times M_2 \rightarrow M_i, i = 1, 2$  the canonical projections. For any  $X \in \Gamma(TM_1)$  and  $Y \in \Gamma(TM_2)$ , we denote by  $X+Y$  the vector field on  $M$  given by  $(X+Y)(m_1, m_2) = (X(m_1), Y(m_2))$ . The product of the affine atlases on  $M_1$  and  $M_2$  is an affine atlas on  $M$  and the corresponding affine connection is the unique flat connection  $\nabla$  on  $M$  satisfying  $\nabla_{X_1+Y_1}(X_2+Y_2) = \nabla_{X_1}^1 Y_1 + \nabla_{X_2}^2 Y_2$ , for any  $X_1, X_2 \in \Gamma(TM_1)$  and  $Y_1, Y_2 \in \Gamma(TM_2)$ . Moreover, the product of  $h_1$  and  $h_2$  is the unique symmetric bivector field  $h$  satisfying

$$h(p_1^* \alpha_1, p_1^* \alpha_2) = h^1(\alpha_1, \alpha_2) \circ p_1, \quad h(p_2^* \beta_1, p_2^* \beta_2) = h^2(\beta_1, \beta_2) \circ p_2 \quad \text{and} \quad h(p_1^* \alpha_1, p_2^* \beta_1) = 0,$$

for any  $\alpha_1, \beta_1 \in \Omega^1(M_1), \alpha_2, \beta_2 \in \Omega^1(M_2)$ ,

**Proposition 2.9.**  $(M, \nabla, h)$  is a contravariant pseudo-Hessian manifold.

*Proof.* Let  $(m_1, m_2) \in M$ . Choose an affine coordinates system  $(x_1, \dots, x_{n_1})$  near  $m_1$  and an affine coordinates system  $(y_1, \dots, y_{n_2})$  near  $m_2$ . Then

$$h = \sum_{i,j} h_{ij}^1 \circ p_1 \partial_{x_i} \otimes \partial_{x_j} + \sum_{l,k} h_{lk}^2 \circ p_2 \partial_{y_l} \otimes \partial_{y_k}$$

and one can check easily that  $h$  satisfies (2.2).  $\square$

If we pursue the exploration of the analogies between Poisson manifolds and contravariant pseudo-Hessian manifolds we can ask naturally if there is an analog of the Darboux-Weinstein's theorem (see[15]) in the context of contravariant pseudo-Hessian manifolds. More precisely, let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold and  $m \in M$  where  $\text{rank} h_\#(m) = r$ . One can ask if there exists an affine coordinates system  $(x_1, \dots, x_r, y_1, \dots, y_{n-r})$  such that

$$h = \sum_{i,j=1}^r h_{ij}(x_1, \dots, x_r) \partial_{x_i} \otimes \partial_{x_j} + \sum_{i,j=1}^{n-r} f_{ij}(y_1, \dots, y_{n-r}) \partial_{y_i} \otimes \partial_{y_j},$$

where  $(h_{ij})_{1 \leq i, j \leq r}$  is invertible and its the inverse of  $\left(\frac{\partial^2 \phi}{\partial x_i \partial x_j}\right)_{1 \leq i, j \leq r}$  and  $f_{ij}(m) = 0$  for any  $i, j$ .

Moreover, if the rank of  $h_{\#}$  is constant near  $m$  then the functions  $f_{ij}$  vanish.

The answer is no in general for a geometric reason. Suppose that  $m$  is regular, i.e., the rank of  $h$  is constant near  $m$  and suppose that there exists an affine coordinates system  $(x_1, \dots, x_r, y_1, \dots, y_{n-r})$  such that

$$h = \sum_{i,j=1}^r h_{ij}(x_1, \dots, x_r) \partial_{x_i} \otimes \partial_{x_j}.$$

This will have a strong geometric consequence, namely that  $\text{Im}h_{\#} = \text{span}(\partial_{x_1}, \dots, \partial_{x_r})$  and the associated affine foliation is parallel, i.e., if  $X$  is a local vector field and  $Y$  is tangent to the foliation then  $\nabla_X Y$  is tangent to the foliation. We give now an example of a regular contravariant pseudo-Hessian manifold whose associated affine foliation is not parallel which shows that the analog of Darboux-Weinstein is not true in general.

**Example 2.10.** We consider  $M = \mathbb{R}^4$  endowed with its canonical affine connection  $\nabla$ , denote by  $(x, y, z, t)$  its canonical coordinates and consider

$$X = \cos(t)\partial_x + \sin(t)\partial_y + \partial_z, \quad Y = -\sin(t)\partial_x + \cos(t)\partial_y \quad \text{and} \quad h = X \otimes Y + Y \otimes X.$$

We have  $\nabla_X X = \nabla_Y X = \nabla_X Y = \nabla_Y Y = 0$  and hence  $h$  is a pseudo-Hessian bivector field,  $\text{Im}h_{\#} = \text{span}\{X, Y\}$  and the rank of  $h$  is constant equal to 2. However, the foliation associated to  $\text{Im}h_{\#}$  is not parallel since  $\nabla_{\partial_t} Y = -X + \partial_z \notin \text{Im}h_{\#}$ .

However, when  $h$  has constant rank equal to  $\dim M - 1$ , we have the following result and its important corollary.

**Theorem 2.11.** Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold and  $m \in M$  such that  $m$  is a regular point and the rank of  $h_{\#}(m)$  is equal to  $n - 1$ . Then there exists an affine coordinates system  $(x_1, \dots, x_n)$  around  $m$  and a function  $f(x_1, \dots, x_n)$  such that

$$h = \sum_{i,j=1}^{n-1} h_{ij} \partial_{x_i} \otimes \partial_{x_j},$$

and the matrix  $(h_{ij})_{1 \leq i, j \leq n-1}$  is invertible and its inverse is the matrix  $\left(\frac{\partial^2 f}{\partial x_i \partial x_j}\right)_{1 \leq i, j \leq n-1}$ .

**Corollary 2.12.** Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold with  $h$  of constant rank equal to  $\dim M - 1$ . Then the affine foliation associated to  $\text{Im}h_{\#}$  is  $\nabla$ -parallel.

In order to prove this theorem, we need the following lemma.

**Lemma 2.13.** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a differentiable function such that  $\partial_x(f) + f\partial_y(f) = 0$ . Then  $f$  is a constant.

*Proof.* Let  $f$  be a solution of the equation above. We consider the vector field  $X_f = \partial_x + f\partial_y$ . The integral curve  $(x(t), y(t))$  of  $X_f$  passing through  $(a, b) \in \mathbb{R}^2$  satisfies

$$x'(t) = 1, \quad y'(t) = f(x(t), y(t)) \quad \text{and} \quad (x(0), y(0)) = (a, b).$$

Now

$$y''(t) = \partial_x(f)(x(t), y(t)) + y'(t)\partial_y(f)(x(t), y(t)) = 0$$

and hence, the flow of  $X_f$  is given by  $\phi(t, (x, y)) = (t+x, f(x, y)t+y)$ . The relation  $\phi(t+s, (x, y)) = \phi(t, \phi(s, (x, y)))$  implies that the map  $F(x, y) = (1, f(x, y))$  satisfies

$$F(u + tF(u)) = F(u), \quad u \in \mathbb{R}^2, t \in \mathbb{R}.$$

Let  $u, v \in \mathbb{R}^2$  such that  $F(u)$  and  $F(v)$  are linearly independent. Then there exists  $s, t \in \mathbb{R}$  such that  $u - v = tF(u) + sF(v)$  and hence  $F(u) = F(v)$  which is a contradiction. So  $F(x, y) = \alpha(x, y)(a, b)$ , i.e.,  $(1, f(x, y)) = (\alpha(x, y)a, \alpha(x, y)b)$  and  $\alpha$  must be constant and hence  $f$  is constant.  $\square$

*Proof of Theorem 2.11.*

*Proof.* Let  $(x_1, \dots, x_n)$  be an affine coordinates system near  $m$  such that  $(X_1, \dots, X_{n-1})$  are linearly independent in a neighborhood of  $m$ , where  $X_i = h_{\#}(dx_i)$ ,  $X_n = \sum_{j=1}^{n-1} f_j X_j$  and, by virtue of the proof of Theorem 2.4, for any  $1 \leq i < j \leq n$ ,  $[X_i, X_j] = 0$ . For any  $i = 1, \dots, n-1$ , the relation  $[X_i, X_n] = 0$  is equivalent to

$$X_i(f_j) = h_{in}\partial_{x_n}(f_j) + \sum_{l=1}^{n-1} h_{il}\partial_{x_l}(f_j) = 0, \quad j = 1, \dots, n-1.$$

But  $h_{in} = X_n(x_i) = \sum_{l=1}^{n-1} f_l h_{il}$  and hence, for any  $i, j = 1, \dots, n-1$ ,

$$\sum_{l=1}^{n-1} h_{il}(f_l \partial_{x_n}(f_j) + \partial_{x_l}(f_j)) = 0.$$

Or the matrix  $(h_{ij})_{1 \leq i, j \leq n-1}$  is invertible so we get

$$f_l \partial_{x_n}(f_j) + \partial_{x_l}(f_j) = 0, \quad l, j = 1, \dots, n-1. \quad (2.5)$$

For  $l = j$  we get that  $f_j$  satisfies  $f_j \partial_{x_n}(f_j) + \partial_{x_j}(f_j) = 0$  so, according to Lemma 2.13,  $\partial_{x_n}(f_j) = \partial_{x_j}(f_j) = 0$  and, from (2.5),  $f_j = \text{constant}$ . We consider  $y = f_1 x_1 + \dots + f_{n-1} x_{n-1} - x_n$ , we have  $h_{\#}(dy) = 0$  and  $(x_1, \dots, x_{n-1}, y)$  is an affine coordinates system around  $m$ .

On the other hand, there exists a coordinates system  $(z_1, \dots, z_n)$  such that

$$h_{\#}(dx_i) = \partial_{z_i}, \quad i = 1, \dots, n-1.$$

We deduce that

$$\partial_{x_i} = \sum_{j=1}^{n-1} h^{ij} \partial_{z_j}, \quad i = 1, \dots, n-1,$$

with  $h^{ij} = \frac{\partial z_j}{\partial x_i}$ . We consider  $\sigma = \sum_{j=1}^{n-1} z_j dx_j$ . We have  $d\sigma = 0$  so according to the foliated Poincaré Lemma (see[6, p.56]) there exists a function  $f$  such that  $h^{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$ .  $\square$



#### 2.4. The divergence and the modular class of a contravariant pseudo-Hessian manifold

We define now the divergence of a contravariant pseudo-Hessian structure. We recall first the definition of the divergence of multivector fields associated to a connection on a manifold.

Let  $(M, \nabla)$  be a manifold endowed with a connection. We define  $\text{div}_\nabla : \Gamma(\otimes^p TM) \rightarrow \Gamma(\otimes^{p-1} TM)$  by

$$\text{div}_\nabla(T)(\alpha_1, \dots, \alpha_{p-1}) = \sum_{i=1}^n \nabla_{e_i}(T)(e_i^*, \alpha_1, \dots, \alpha_{p-1}),$$

where  $\alpha_1, \dots, \alpha_{p-1} \in T_x^*M$ ,  $(e_1, \dots, e_n)$  a basis of  $T_x M$  and  $(e_1^*, \dots, e_n^*)$  its dual basis. This operator respects the symmetries of tensor fields.

Suppose now that  $(M, \nabla, h)$  is a contravariant pseudo-Hessian manifold. The divergence of this structure is the vector field  $\text{div}_\nabla(h)$ . This vector field is an invariant of the pseudo-Hessian structure and has an important property. Indeed, let  $\mathbf{d}_h : \Gamma(\wedge^\bullet TM) \rightarrow \Gamma(\wedge^{\bullet+1} TM)$  be the differential associated to the Lie algebroid structure  $(T^*M, h_\#, [\ , \ ]_h)$  and given by

$$\begin{aligned} \mathbf{d}_h Q(\alpha_1, \dots, \alpha_p) &= \sum_{j=1}^p (-1)^{j+1} h_\#(\alpha_j).Q(\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_p) \\ &\quad + \sum_{1 \leq i < j \leq p} (-1)^{i+j} Q([\alpha_i, \alpha_j]_h, \alpha_1, \dots, \hat{\alpha}_i, \dots, \hat{\alpha}_j, \dots, \alpha_p). \end{aligned}$$

**Proposition 2.14.**  $\mathbf{d}_h(\text{div}_\nabla(h)) = 0$ .

*Proof.* Let  $(x_1, \dots, x_n)$  be an affine coordinates system. We have

$$\begin{aligned} \mathbf{d}_h \text{div}_\nabla(h)(\alpha, \beta) &= \sum_{i=1}^n \left( h_\#(\alpha). \nabla_{\partial_{x_i}}(h)(dx_i, \beta) - h_\#(\beta). \nabla_{\partial_{x_i}}(h)(dx_i, \alpha) - \nabla_{\partial_{x_i}}(h)(dx_i, \nabla_{h_\#(\alpha)}\beta) + \nabla_{\partial_{x_i}}(h)(dx_i, \nabla_{h_\#(\beta)}\alpha) \right) \\ &= \sum_{i=1}^n \left( \nabla_{h_\#(\alpha)} \nabla_{\partial_{x_i}}(h)(dx_i, \beta) - \nabla_{h_\#(\beta)} \nabla_{\partial_{x_i}}(h)(dx_i, \alpha) \right) \\ &\stackrel{(2.1)}{=} \sum_{i=1}^n \left( \nabla_{[h_\#(\alpha), \partial_{x_i}]}(h)(dx_i, \beta) - \nabla_{[h_\#(\beta), \partial_{x_i}]}(h)(dx_i, \alpha) \right). \end{aligned}$$

If we take  $\alpha = dx_l$  and  $\beta = dx_k$ , we have

$$[\partial_{x_i}, h_\#(dx_l)] = \sum_{m=1}^n \partial_{x_i}(h_{ml}) \partial_{x_m}$$

and hence

$$\mathbf{d}_h \text{div}_\nabla(h)(\alpha, \beta) = \sum_{i,m=1}^n (\partial_{x_i}(h_{ml}) \partial_{x_m}(h_{ik}) - \partial_{x_i}(h_{mk}) \partial_{x_m}(h_{il})) = 0.$$

□

Let  $(M, \nabla, h)$  be an orientable contravariant pseudo-Hessian manifold and  $\Omega$  a volume form on  $M$ . For any  $f$  we denote by  $X_f = h_\#(df)$  and we define  $\mathbf{M}_\Omega : C^\infty(M, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R})$  by putting for any  $f \in C^\infty(M, \mathbb{R})$ ,

$$\nabla_{X_f} \Omega = \mathbf{M}_\Omega(f) \Omega.$$

It is obvious that  $\mathbf{M}_\Omega$  is a derivation and hence a vector field and  $\mathbf{M}_{e^f\Omega} = X_f + \mathbf{M}_\Omega$ . Moreover, if  $(x_1, \dots, x_n)$  is an affine coordinates system and  $\mu = \Omega(\partial_{x_1}, \dots, \partial_{x_n})$  then

$$\nabla_{X_f}\Omega(\partial_{x_1}, \dots, \partial_{x_n}) = X_f(\mu) = X_{\ln|\mu|}(f)\mu.$$

So in the coordinates system  $(x_1, \dots, x_n)$ , we have  $\mathbf{M}_\Omega = X_{\ln|\mu|}$ . This implies  $\mathbf{d}_h\mathbf{M}_\Omega = 0$ . The cohomology class of  $\mathbf{M}_\Omega$  doesn't depend on  $\Omega$  and we call it the *modular class* of  $(M, \nabla, h)$ .

**Proposition 2.15.** *The modular class of  $(M, \nabla, h)$  vanishes if and only if there exists a volume form  $\Omega$  such that  $\nabla_{X_f}\Omega = 0$  for any  $f \in C^\infty(M, \mathbb{R})$ .*

By analogy with the case of Poisson manifolds, one can ask if it is possible to find a volume form  $\Omega$  such that  $\mathcal{L}_{X_f}\Omega = 0$  for any  $f \in C^\infty(M, \mathbb{R})$ . The following proposition gives a negative answer to this question unless  $h = 0$ .

**Proposition 2.16.** *Let  $(M, \nabla, h)$  be an orientable contravariant pseudo-Hessian manifold. Then:*

1. *For any volume form  $\Omega$  and any  $f \in C^\infty(M, \mathbb{R})$ ,*

$$\mathcal{L}_{X_f}\Omega = [\mathbf{M}_\Omega(f) + \text{div}_\nabla(h)(f) + \langle h, \text{Hess}(f) \rangle] \Omega,$$

where  $\text{Hess}(f)(X, Y) = \nabla_X(df)(Y)$  and  $\langle h, \text{Hess}(f) \rangle$  is the pairing between the bivector field  $h$  and the 2-form  $\text{Hess}(f)$ .

2. *There exists a volume form  $\Omega$  such that  $\mathcal{L}_{X_f}\Omega = 0$  for any  $f \in C^\infty(M, \mathbb{R})$  if and only if  $h = 0$ .*

*Proof.* 1. Let  $(x_1, \dots, x_n)$  be an affine coordinates system. Then:

$$\begin{aligned} [X_f, \partial_{x_i}] &= \sum_{l,j=1}^n [\partial_{x_j}(f)h_{jl}\partial_{x_l}, \partial_{x_i}] \\ &= - \sum_{l,j=1}^n (h_{jl}\partial_{x_l}\partial_{x_j}(f) + \partial_{x_j}(f)\partial_{x_l}(h_{ji}))\partial_{x_i}, \\ \mathcal{L}_{X_f}\Omega(\partial_{x_1}, \dots, \partial_{x_n}) &= (\nabla_{X_f}\Omega)(\partial_{x_1}, \dots, \partial_{x_n}) - \sum_{i=1}^n \Omega((\partial_{x_1}, \dots, [X_f, \partial_{x_i}], \dots, \partial_{x_n})) \\ &= (\nabla_{X_f}\Omega)(\partial_{x_1}, \dots, \partial_{x_n}) + \sum_{i,j=1}^n (h_{ji}\partial_{x_l}\partial_{x_j}(f) + \partial_{x_j}(f)\partial_{x_l}(h_{ji}))\Omega(\partial_{x_1}, \dots, \partial_{x_n}) \end{aligned}$$

and the formula follows since  $\text{div}_\nabla(h) = \sum_{i,j=1}^n \partial_{x_i}(h_{ji})\partial_{x_j}$ .

2. This is a consequence of the fact that  $\mathbf{M}_\Omega$  and  $\text{div}_\nabla(h)$  are derivation and

$$\langle h, \text{Hess}(fg) \rangle = f \langle h, \text{Hess}(g) \rangle + g \langle h, \text{Hess}(f) \rangle + \langle h, df \odot dg \rangle.$$

□

### 3. The tangent bundle of a contravariant pseudo-Hessian manifold

In this section, we define and study the associated Poisson tensor on the tangent bundle of a contravariant pseudo-Hessian manifold. One can see [7] for the classical properties of the tangent bundle of a manifold endowed with a linear connection.

Let  $(M, \nabla)$  be an affine manifold,  $p : TM \rightarrow M$  the canonical projection and  $K : TTM \rightarrow TTM$  the connection map of  $\nabla$  locally given by

$$K \left( \sum_{i=1}^n b_i \partial_{x_i} + \sum_{j=1}^n Z_j \partial_{\mu_j} \right) = \sum_{l=1}^n \left( Z_l + \sum_{i=1}^n \sum_{j=1}^n b_i \mu_j \Gamma_{ij}^l \right) \partial_{x_l},$$

where  $(x_1, \dots, x_n)$  is a system of local coordinates,  $(x_1, \dots, x_n, \mu_1, \dots, \mu_n)$  the associated system of coordinates on  $TM$  and  $\nabla_{\partial_{x_i}} \partial_{x_j} = \sum_{l=1}^n \Gamma_{ij}^l \partial_{x_l}$ . Then

$$TTM = \ker Tp \oplus \ker K.$$

For  $X \in \Gamma(TM)$ , we denote by  $X^h$  its horizontal lift and by  $X^v$  its vertical lift. The flow of  $X^v$  is given by  $\Phi^X(t, (x, u)) = (x, u + tX(x))$  and  $X^h(x, u) = h^{(x,u)}(X(x))$ , where  $h^{(x,u)} : T_x M \rightarrow \ker K(x, u)$  is the inverse of the restriction of  $dp$  to  $\ker K(x, u)$ . Since the curvature of  $\nabla$  vanishes, we have

$$[X^h, Y^h] = [X, Y]^h, \quad [X^h, Y^v] = (\nabla_X Y)^v \quad \text{and} \quad [X^v, Y^v] = 0, \quad (3.1)$$

for any  $X, Y \in \Gamma(TM)$ . For any  $\alpha \in \Omega^1(M)$ , we define  $\alpha^v, \alpha^h \in \Omega^1(TM)$  by

$$\begin{cases} \alpha^v(X^v) = \alpha(X) \circ p, \\ \alpha^v(X^h) = 0, \end{cases} \quad \text{and} \quad \begin{cases} \alpha^h(X^h) = \alpha(X) \circ p, \\ \alpha^h(X^v) = 0. \end{cases}$$

The following proposition is well-known [7] and can be proved easily.

**Proposition 3.1.** *The connection  $\bar{\nabla}$  on  $TM$  given by*

$$\bar{\nabla}_{X^h} Y^h = (\nabla_X Y)^h, \quad \bar{\nabla}_{X^h} Y^v = (\nabla_X Y)^v \quad \text{and} \quad \bar{\nabla}_{X^v} Y^h = \bar{\nabla}_{X^v} Y^v = 0, \quad (3.2)$$

where  $X, Y \in \Gamma(TM)$ , defines an affine structure on  $TM$ . Moreover, the endomorphism vector field  $J : TTM \rightarrow TTM$  given by  $JX^h = X^v$  and  $JX^v = -X^h$  satisfies  $J^2 = -\text{Id}_{TTM}$ , is parallel with respect to  $\bar{\nabla}$  and hence defines a complex structure on  $TM$ .

Let  $h$  be a symmetric bivector field on  $M$ . We associate to  $h$  a skew-symmetric bivector field  $\Pi$  on  $TM$  by putting

$$\Pi(\alpha^v, \beta^v) = \Pi(\alpha^h, \beta^h) = 0 \quad \text{and} \quad \Pi(\alpha^h, \beta^v) = -\Pi(\beta^v, \alpha^h) = h(\alpha, \beta) \circ p,$$

for any  $\alpha, \beta \in \Omega^1(M)$ . For any  $\alpha \in \Omega^1(M)$ ,

$$\Pi_{\#}(\alpha^v) = -h_{\#}(\alpha)^h \quad \text{and} \quad \Pi_{\#}(\alpha^h) = h_{\#}(\alpha)^v. \quad (3.3)$$

To prove one of our main result in this section, we need the following proposition which is a part of the folklore.

**Proposition 3.2.** *Let  $(P, \nabla)$  be a manifold endowed with a torsionless connection and  $\pi$  is a bivector field on  $P$ . Then the Nijenhuis-Schouten bracket  $[\pi, \pi]$  is given by*

$$[\pi, \pi](\alpha, \beta, \gamma) = 2 \left( \nabla_{\pi_{\#}(\alpha)} \pi(\beta, \gamma) + \nabla_{\pi_{\#}(\beta)} \pi(\gamma, \alpha) + \nabla_{\pi_{\#}(\gamma)} \pi(\alpha, \beta) \right).$$

**Theorem 3.3.** *The following assertions are equivalent:*

- (i)  $(M, \nabla, h)$  is a contravariant pseudo-Hessian manifold.
- (ii)  $(TM, \Pi)$  is a Poisson manifold.

*In this case, if  $L$  is a leaf of  $\text{Im} h_{\#}$  then  $TL \subset TM$  is a symplectic leaf of  $\Pi$  which is also a complex submanifold of  $TM$ . Moreover, if  $\omega_L$  is the symplectic form of  $TL$  induced by  $\Pi$  and  $g_L$  is the pseudo-Riemannian metric given by  $g_L(U, V) = \omega(JU, V)$  then  $(TL, g_L, \omega_L, J)$  is a pseudo-Kähler manifold.*

*Proof.* We will use Proposition 3.2 to prove the equivalence. Indeed, by a direct computation one can establish easily, for any  $\alpha, \beta, \gamma \in \Omega^1(M)$ , the following relations

$$\begin{aligned} \bar{\nabla}_{\Pi_{\#}(\alpha^v)} \Pi(\beta^v, \gamma^v) &= \bar{\nabla}_{\Pi_{\#}(\alpha^v)} \Pi(\beta^h, \gamma^h) = \bar{\nabla}_{\Pi_{\#}(\alpha^h)} \Pi(\beta^v, \gamma^v) = \bar{\nabla}_{\Pi_{\#}(\alpha^h)} \Pi(\beta^h, \gamma^h) = \bar{\nabla}_{\Pi_{\#}(\alpha^h)} \Pi(\beta^h, \gamma^v) = 0, \\ \bar{\nabla}_{\Pi_{\#}(\alpha^v)} \Pi(\beta^h, \gamma^v) &= \nabla_{h_{\#}(\alpha)}(h)(\beta, \gamma) \circ p, \end{aligned}$$

and the equivalence follows. The second part of the theorem is obvious and the only point which need to be checked is that  $g_L$  is nondegenerate.  $\square$

**Remark 3.4.** 1. *The total space of the dual of a Lie algebroid carries a Poisson tensor (see [12]). If  $(M, \nabla, h)$  is a contravariant pseudo-Hessian manifold then, according to Theorem 2.4,  $T^*M$  carries a Lie algebroid structure and one can see easily that  $\Pi$  is the corresponding Poisson tensor on  $TM$ .*

2. *The equivalence of (i) and (ii) in Theorem 3.3 deserves to be stated explicitly in the case of  $\mathbb{R}^n$  endowed with its canonical affine structure  $\nabla$ . Indeed, let  $(h_{ij})_{1 \leq i, j \leq n}$  be a symmetric matrix where  $h_{ij} \in C^\infty(\mathbb{R}^n, \mathbb{R})$  and  $h$  the associated symmetric bivector field on  $\mathbb{R}^n$ . The associated bivector field  $\Pi_h$  on  $T\mathbb{R}^n = \mathbb{C}^n$  is*

$$\Pi_h = \sum_{i,j=1}^n h_{ij}(x) \partial_{x_i} \wedge \partial_{x_j},$$

*where  $(x_1 + iy_1, \dots, x_n + iy_n)$  are the canonical coordinates of  $\mathbb{C}^n$ . Then, according to Theorem 2.4,  $(\mathbb{R}^n, \nabla, h)$  is a contravariant pseudo-Hessian manifold if and only if  $(\mathbb{C}^n, \Pi_h)$  is a Poisson manifold.*

We explore now some relations between some invariants of  $(M, \nabla, h)$  and some invariants of  $(TM, \Pi)$ .

**Proposition 3.5.** *Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold. Then  $(\text{div}_{\nabla} h)^v = \text{div}_{\bar{\nabla}} \Pi$ .*

*Proof.* Fix  $(x, u) \in TM$  and choose a basis  $(e_1, \dots, e_n)$  of  $T_x M$ . Then  $(e_1^v, \dots, e_n^v, e_1^h, \dots, e_n^h)$  is a basis of  $T_{(x,u)} TM$  with  $((e_1^*)^v, \dots, (e_n^*)^v, (e_1^*)^h, \dots, (e_n^*)^h)$  as a dual basis. For any  $\alpha \in T_x^* M$ , we have

$$\begin{aligned} \langle \alpha^v, \text{div}_{\bar{\nabla}} \Pi \rangle &= \sum_{i=1}^n \left( \bar{\nabla}_{e_i^v} (\Pi)((e_i^*)^v, \alpha^v) + \bar{\nabla}_{e_i^h} (\Pi)((e_i^*)^h, \alpha^v) \right) \\ &\stackrel{(3.2)}{=} \langle \alpha, \text{div}_{\nabla} (h) \rangle \circ p = \langle \alpha^v, (\text{div}_{\nabla} (h))^v \rangle. \end{aligned}$$

In the same way we get that  $\langle \alpha^h, \text{div}_{\overline{\nabla}} \Pi \rangle = 0$  and the result follows.  $\square$

Let  $(M, \nabla, h)$  be a contravariant pseudo-Hessian manifold. For any multivector field  $Q$  on  $M$  we define its vertical lift  $Q^v$  on  $TM$  by

$$i_{\alpha^h} Q^v = 0 \quad \text{and} \quad Q^v(\alpha_1^v, \dots, \alpha_q^v) = Q(\alpha_1, \dots, \alpha_q) \circ p.$$

Recall that  $h$  defines a Lie algebroid structure on  $T^*M$  whose anchor is  $h_{\#}$  and the Lie bracket is given by (2.3). The Poisson tensor  $\Pi$  defines a Lie algebroid structure on  $T^*TM$  whose anchor is  $\Pi_{\#}$  and the Lie bracket is the Koszul bracket

$$[\phi_1, \phi_2]_{\Pi} = \mathcal{L}_{\Pi_{\#}(\phi_1)} \phi_2 - \mathcal{L}_{\Pi_{\#}(\phi_2)} \phi_1 - d\Pi(\phi_1, \phi_2), \quad \phi_1, \phi_2 \in \Omega^1(TM).$$

We denote by  $\mathbf{d}_h$  (resp.  $\mathbf{d}_{\Pi}$ ) the differential associated to the Lie algebroid structure on  $T^*M$  (resp.  $T^*TM$ ) defined by  $h$  (resp.  $\Pi$ ).

**Proposition 3.6.** (i) For any  $\alpha, \beta \in \Omega^1(M)$  and  $X \in \Gamma(TM)$ ,

$$\begin{cases} \mathcal{L}_{X^h} \alpha^h = (\mathcal{L}_X \alpha)^h, \quad \mathcal{L}_{X^h} \alpha^v = (\nabla_X \alpha)^v, \quad \mathcal{L}_{X^v} \alpha^h = 0 \quad \text{and} \quad \mathcal{L}_{X^v} \alpha^v = (\mathcal{L}_X \alpha)^h - (\nabla_X \alpha)^h, \\ [\alpha^h, \beta^h]_{\Pi} = 0, \quad [\alpha^v, \beta^v]_{\Pi} = -[\alpha, \beta]_h^v \quad \text{and} \quad [\alpha^h, \beta^v]_{\Pi} = (\mathcal{D}_{\beta} \alpha)^h, \end{cases}$$

where  $\mathcal{D}$  is the connection given by (2.4).

(ii)  $(\mathbf{d}_h Q)^v = -\mathbf{d}_{\Pi}(Q^v)$ .

*Proof.* The relations in (i) can be established by a straightforward computation. From these relations and the fact that  $\Pi_{\#}(\alpha^h) = (h_{\#}(\alpha))^v$  one can deduce easily that  $i_{\alpha^h} \mathbf{d}_{\Pi}(Q^v) = 0$ . On the other hand, since  $\Pi_{\#}(\alpha^v) = -(h_{\#}(\alpha))^h$  and  $[\alpha^v, \beta^v]_{\Pi} = -[\alpha, \beta]^v$  we can conclude.  $\square$

**Remark 3.7.** From Propositions 2.14 and Proposition 3.6, we can deduce that  $\mathbf{d}_{\Pi}(\text{div}_{\overline{\nabla}} \Pi) = 0$ . This is not a surprising result because  $\overline{\nabla}$  is flat and  $\text{div}_{\overline{\nabla}} \Pi$  is a representative of the modular class of  $\Pi$ .

As a consequence of Proposition 3.6 we can define a linear map from the cohomology of  $(T^*M, h_{\#}, [\cdot, \cdot]_h)$  to the cohomology of  $(T^*TM, \Pi_{\#}, [\cdot, \cdot]_{\Pi})$  by

$$V : H^*(M, h) \longrightarrow H^*(TM, \Pi), \quad [Q] \mapsto [Q^v].$$

**Proposition 3.8.**  $V$  is injective.

*Proof.* An element  $P \in \Gamma(\wedge^d TTM)$  is of type  $(r, d-r)$  if for any  $q \neq r$

$$P(\alpha_1^v, \dots, \alpha_q^v, \beta_1^h, \dots, \beta_{d-q}^h) = 0,$$

for any  $\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_{d-q} \in \Omega^1(M)$ . We have

$$\begin{cases} \Gamma(\wedge^d TTM) = \bigoplus_{r=0}^d \Gamma_{(r, d-r)}(\wedge^d TTM), \\ \mathbf{d}_{\Pi}(\Gamma_{(r, d-r)}(\wedge^d TTM)) \subset \Gamma_{(r+1, d-r)}(\wedge^{d+1} TTM) \oplus \Gamma_{(r, d+1-r)}(\wedge^{d+1} TTM). \end{cases}$$

Let  $Q \in \Gamma(\wedge^d TTM)$  such that  $\mathbf{d}_h Q = 0$  and there exists  $P \in \Gamma(\wedge^{d-1} TTM)$  such that  $\mathbf{d}_{\Pi} P = Q^v$ . Since  $Q^v \in \Gamma_{(d, 0)}(\wedge^d TTM)$  then  $P \in \Gamma_{(d-1, 0)}(\wedge^{d-1} TTM)$ . Let us show that  $P = T^v$ . For  $\alpha_1, \dots, \alpha_{d-1}, \beta \in \Omega^1(M)$ , we have

$$0 = \mathbf{d}_{\Pi} P(\beta^h, \alpha_1^v, \dots, \alpha_{d-1}^v) = (h_{\#}(\beta))^v \cdot P(\alpha_1^v, \dots, \alpha_{d-1}^v).$$

So the function  $P(\alpha_1^v, \dots, \alpha_{d-1}^v)$  is constant on the fibers of  $TM$  and hence there exists  $T \in \Gamma(\wedge^{d-1} TM)$  such that  $P(\alpha_1^v, \dots, \alpha_{d-1}^v) = T(\alpha_1, \dots, \alpha_{d-1}) \circ p$ . So  $[Q] = 0$  which completes the proof.  $\square$

#### 4. Linear, affine and multiplicative contravariant pseudo-Hessian structures

##### 4.1. Linear and affine contravariant pseudo-Hessian structures

As in the Poisson geometry context, we have the notions of linear and affine contravariant pseudo-Hessian structures. One can see [11] for the notion of cocycle in associative algebras.

Let  $(V, \nabla)$  be a finite dimensional real vector space endowed with its canonical affine structure. A symmetric bivector field  $h$  on  $V$  is called affine if there exists a commutative product  $\bullet$  on  $V^*$  and a symmetric bilinear form  $B$  on  $V^*$  such that, for any  $\alpha, \beta \in V^* \subset \Omega^1(V)$  and  $u \in V$ ,

$$h(\alpha, \beta)(u) = \langle \alpha \bullet \beta, u \rangle + B(\alpha, \beta).$$

One can see easily that if  $\alpha, \beta \in \Omega^1(V) = C^\infty(V, V^*)$  then

$$h(\alpha, \beta)(u) = \langle \alpha(u) \bullet \beta(u), u \rangle + B(\alpha(u), \beta(u)).$$

If  $B = 0$ ,  $h$  is called linear.

If  $(x_1, \dots, x_n)$  is a linear coordinates system on  $V^*$  associated to a basis  $(e_1, \dots, e_n)$  then

$$h(dx_i, dx_j) = b_{ij} + \sum_{k=1}^n C_{ij}^k x_k,$$

where  $e_i \bullet e_j = \sum_{k=1}^n C_{ij}^k e_k$  and  $b_{ij} = B(e_i, e_j)$ .

**Proposition 4.1.**  $(V, \nabla, h)$  is a contravariant pseudo-Hessian manifold if and only if  $\bullet$  is associative and  $B$  is a scalar 2-cocycle of  $(V^*, \bullet)$ , i.e.,

$$B(\alpha \bullet \beta, \gamma) = B(\alpha, \beta \bullet \gamma)$$

for any  $\alpha, \beta, \gamma \in V^*$ .

*Proof.* For any  $\alpha \in V^*$  and  $u \in V$ ,  $h_\#(\alpha)(u) = L_\alpha^* u + i_\alpha B$  where  $L_\alpha(\beta) = \alpha \bullet \beta$  and  $i_\alpha B \in V^{**} = V$ . We denote by  $\phi^{h_\#(\alpha)}$  the flow of the vector field  $h_\#(\alpha)$ . Then, for any  $\alpha, \beta, \gamma \in V^*$ ,

$$\begin{aligned} \nabla_{h_\#(\alpha)}(h)(\beta, \gamma)(u) &= \frac{d}{dt}\bigg|_{t=0} (\langle \beta \bullet \gamma, \phi^{h_\#(\alpha)}(t, u) \rangle + B(\beta, \gamma)) \\ &= \langle \beta \bullet \gamma, L_\alpha^* u + i_\alpha B \rangle \\ &= \langle \alpha \bullet (\beta \bullet \gamma), u \rangle + B(\alpha, \beta \bullet \gamma) \end{aligned}$$

and the result follows.  $\square$

Conversely, we have the following result.

**Proposition 4.2.** Let  $(\mathcal{A}, \bullet, B)$  be a commutative and associative algebra endowed with a symmetric scalar 2-cocycle. Then:

1.  $\mathcal{A}^*$  carries a structure of a contravariant pseudo-Hessian structure  $(\nabla, h)$  where  $\nabla$  is the canonical affine structure of  $\mathcal{A}^*$  and  $h$  is given by

$$h(u, v)(\alpha) = \langle \alpha, u(\alpha) \bullet v(\alpha) \rangle + B(u(\alpha), v(\alpha)), \quad \alpha \in \mathcal{A}^*, u, v \in \Omega^1(\mathcal{A}^*).$$

2. When  $B = 0$ , the leaves of the affine foliation associated to  $\text{Im}h_{\#}$  are the orbits of the action  $\Phi$  of  $(\mathcal{A}, +)$  on  $\mathcal{A}^*$  given by  $\Phi(u, \alpha) = \exp(L_u^*)(\alpha)$
3. The associated Poisson tensor  $\Pi$  on  $T\mathcal{A}^* = \mathcal{A}^* \times \mathcal{A}^*$  is the affine Poisson tensor dual associated to the Lie algebra  $(\mathcal{A} \times \mathcal{A}, [\ , \ ])$  endowed with the 2-cocycle  $B_0$  where

$$[(a, b), (c, d)] = (a \bullet d - b \bullet c, 0) \quad \text{and} \quad B_0((a, b), (c, d)) = B(a, d) - B(c, b).$$

*Proof.* It is only the third point which need to be checked. One can see easily that  $[\ , \ ]$  is a Lie bracket on  $\mathcal{A} \times \mathcal{A}$  and  $B_0$  is a scalar 2-cocycle for this Lie bracket. For any  $a \in \mathcal{A} \subset \Omega^1(\mathcal{A}^*)$ ,  $a^v = (0, a) \in \mathcal{A} \times \mathcal{A} \subset \Omega^1(\mathcal{A}^* \times \mathcal{A}^*)$  and  $a^h = (a, 0)$ . So

$$\Pi(a^h, b^v)(\alpha, \beta) = h(a, b)(\alpha) = \langle \alpha, a \bullet b \rangle + B(a, b).$$

On the other hand, if  $\Pi^*$  is the Poisson tensor dual, then

$$\begin{aligned} \Pi^*(a^h, b^v)(\alpha, \beta) &= \Pi^*((a, 0), (0, b))(\alpha, \beta) \\ &= \langle \alpha, \beta \rangle, [(a, 0), (0, b)] \rangle + B_0((a, 0), (0, b)) \\ &= \langle \alpha, a \bullet b \rangle + B(a, b) \\ &= \Pi(a^h, b^v)(\alpha, \beta). \end{aligned}$$

In the same way one can check the others equalities.  $\square$

This proposition can be used as a machinery to build examples of pseudo-Hessian manifolds. Indeed, by virtue of Proposition 2.5, any orbit  $L$  of the action  $\Phi$  has an affine structure  $\nabla_L$  and a pseudo-Riemannian metric  $g_L$  such that  $(L, \nabla_L, g_L)$  is a pseudo-Hessian manifold.

**Example 4.3.** We take  $\mathcal{A} = \mathbb{R}^4$  with its canonical basis  $(e_i)_{i=1}^4$  and  $(e_i^*)_{i=1}^4$  is the dual basis. We endow  $\mathcal{A}$  with the commutative associative product given by

$$e_1 \bullet e_1 = e_2, \quad e_1 \bullet e_2 = e_3, \quad e_1 \bullet e_3 = e_2 \bullet e_2 = e_4,$$

the others products are zero and we endow  $\mathcal{A}^*$  with the linear contravariant pseudo-Hessian structure associated to  $\bullet$ . We denote by  $(a, b, c, d)$  the linear coordinates on  $\mathcal{A}$  and  $(x, y, z, t)$  the dual coordinates on  $\mathcal{A}^*$ . We have

$$\Phi(ae_1 + be_2 + ce_3 + de_4, xe_1^* + ye_2^* + ze_3^* + te_4^*) = (x + ay + (\frac{1}{2}a^2 + b)z + (\frac{1}{6}a^3 + ab + c)t, y + az + (\frac{1}{2}a^2 + b)t, z + at, t)$$

and

$$X_{e_1} = y\partial_x + z\partial_y + t\partial_z, \quad X_{e_2} = z\partial_x + t\partial_y, \quad X_{e_3} = t\partial_x \quad \text{and} \quad X_{e_4} = 0.$$

Let us describe the pseudo-Hessian structure of the hyperplane  $M_c = \{t = c, c \neq 0\}$  endowed with the coordinates  $(x, y, z)$ . We denote by  $g_c$  the pseudo-Riemannian of  $M_c$ . We have, for instance,

$$g_c(X_{e_1}, X_{e_1})(x, y, z, c) = h(e_1, e_1)(x, y, z, c) = \langle e_1 \bullet e_1, (x, y, z, c) \rangle = y.$$

So, one can see that the matrix of  $g_c$  in  $(X_{e_1}, X_{e_2}, X_{e_3})$  is the passage matrix  $P$  from  $(X_{e_1}, X_{e_2}, X_{e_3})$  to  $(\partial_x, \partial_y, \partial_z)$  and hence

$$g_c = \frac{1}{c} \left( 2dx dz + dy^2 - \frac{2z}{c} dy dz + \frac{(z^2 - yc)}{c^2} dz^2 \right).$$

The signature of this metric is  $(+, +, -)$  if  $c > 0$  and  $(+, -, -)$  if  $c < 0$ . One can check easily that  $g_c$  is the restriction of  $\nabla d\phi$  to  $M_c$ , where

$$\phi(x, y, z, t) = \frac{z^4}{12t^3} + \frac{y^2}{2t} - \frac{z^2 y}{2t} + \frac{xz}{t}.$$

#### 4.2. Multiplicative contravariant pseudo-Hessian structures

A contravariant pseudo-Hessian structure  $(\nabla, h)$  on a Lie group  $G$  is called multiplicative if the multiplication  $m : (G \times G, \nabla \oplus \nabla, h \oplus h) \longrightarrow (G, \nabla, h)$  is affine and sends  $h \oplus h$  to  $h$ .

**Lemma 4.4.** *Let  $G$  be a connected Lie group and  $\nabla$  a connection on  $G$  such that the multiplication  $m : (G \times G, \nabla \oplus \nabla) \longrightarrow (G, \nabla)$  preserves the connections. Then  $G$  is abelian and  $\nabla$  is bi-invariant.*

*Proof.* We will denote by  $\chi^r(G)$  (resp.  $\chi^l(G)$ ) the space of left invariant vector fields (resp. the right invariant vector fields) on  $G$ . It is clear that for any  $X \in \chi^r(G)$  and  $Y \in \chi^l(G)$ , the vector field  $(X, Y)$  on  $G \times G$  is  $m$ -related to the vector field  $X + Y$  on  $G$ :

$$Tm(X_a, Y_b) = X_a \cdot b + a \cdot Y_b = X_{ab} + Y_{ab} = (X + Y)_{ab}$$

It follows that for any  $X_1, X_2 \in \chi^r(G)$  and  $Y_1, Y_2 \in \chi^l(G)$ , the vector field  $(\nabla \oplus \nabla)_{(X_1, Y_1)}(X_2, Y_2)$  is  $m$ -related to  $\nabla_{X_1+Y_1}(X_2 + Y_2)$ , hence:

$$Tm((\nabla_{X_1} X_2)_a, (\nabla_{Y_1} Y_2)_b) = (\nabla_{(X_1+Y_1)}(X_2 + Y_2))_{ab}$$

So we get

$$(\nabla_{X_1} X_2)_a \cdot b + a \cdot (\nabla_{Y_1} Y_2)_b = (\nabla_{X_1} X_2 + \nabla_{X_1} Y_2 + \nabla_{Y_1} X_2 + \nabla_{Y_1} Y_2)_{ab} \quad (4.1)$$

If we take  $Y_1 = 0 = Y_2$  we obtain that  $\nabla$  is right invariant. In the same way we get that  $\nabla$  is left invariant. Now, if we return back to the equation 4.1 we obtain that for any  $X \in \chi^r(G)$  and  $Y \in \chi^l(G)$  we have  $\nabla X = 0 = \nabla Y$ . This implies that any left invariant vector field is also right invariant ; indeed, if  $Y = \sum_{i=1}^n f_i X_i$  with  $Y \in \chi^l(G)$  and  $X_i \in \chi^r(G)$  then  $X_j f_i = 0$  for all  $i, j = 1, \dots, n$ . Hence the adjoint representation is trivial and hence  $G$  must be abelian.  $\square$

At the end of the paper, we give another proof of this Lemma based on parallel transport.

**Corollary 4.5.** *Let  $(\nabla, h)$  be multiplicative contravariant pseudo-Hessian structure on a simply connected Lie group  $G$ . Then  $G$  is a vector space,  $\nabla$  its canonical affine connection and  $h$  is linear.*

**Example 4.6.** *Based on the classification of complex associative commutative algebras given in [14], we can give a list of examples of affine contravariant pseudo-Hessian structures up to dimension 4.*

1. On  $\mathbb{R}^2$ :

$$h_1 = \begin{pmatrix} x_2 & 0 \\ 0 & 0 \end{pmatrix}, h_2 = \begin{pmatrix} x_1 & x_2 \\ x_2 & 0 \end{pmatrix} \quad \text{and} \quad h_3 = \begin{pmatrix} x_2 & 1 \\ 1 & 0 \end{pmatrix}.$$

2. On  $\mathbb{R}^3$ :

$$\begin{aligned} h_1 &= \begin{pmatrix} a & 0 & x_2 \\ 0 & 0 & 0 \\ x_2 & 0 & b \end{pmatrix}, h_2 = \begin{pmatrix} x_2 & x_3 & a \\ x_3 & a & 0 \\ a & 0 & 0 \end{pmatrix}, h_3 = \begin{pmatrix} a & 0 & x_1 \\ 0 & 0 & x_2 \\ x_1 & x_2 & x_3 \end{pmatrix}, \\ h_4 &= \begin{pmatrix} x_2 & 0 & x_2 \\ 0 & 0 & x_2 + a \\ x_2 & x_2 + a & x_3 \end{pmatrix} \quad \text{and} \quad h_5 = \begin{pmatrix} x_2 & 0 & x_1 \\ 0 & 0 & x_2 \\ x_1 & x_2 & x_3 \end{pmatrix}. \end{aligned}$$



3. On  $\mathbb{R}^4$  :

$$h_1 = \begin{pmatrix} x_3 & a & x_4 + b & 0 \\ a & -x_4 + c & 0 & 0 \\ x_4 + b & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad h_2 = \begin{pmatrix} x_2 & x_3 & x_4 & a \\ x_3 & x_4 & a & 0 \\ x_4 & a & 0 & 0 \\ a & 0 & 0 & 0 \end{pmatrix}, \quad h_3 = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 \\ x_2 & 0 & 0 & 0 \\ x_3 & 0 & 0 & 0 \\ x_4 & 0 & 0 & 0 \end{pmatrix},$$

$$h_4 = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 \\ x_2 & x_4 & 0 & 0 \\ x_3 & 0 & 0 & 0 \\ x_4 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad h_5 = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 \\ x_2 & x_3 & x_4 & 0 \\ x_3 & x_4 & 0 & 0 \\ x_4 & 0 & 0 & 0 \end{pmatrix}.$$

### 5. Quadratic contravariant pseudo-Hessian structures

Let  $V$  be a vector space of dimension  $n$ . Denote by  $\nabla$  its canonical affine connection. A symmetric bivector field  $h$  on  $V$  is quadratic if there exists a basis  $\mathbb{B}$  of  $V$  such that, for any  $i, j = 1, \dots, n$ ,

$$h(dx_i, dx_j) = \sum_{l,k=1}^n a_{l,k}^{i,j} x_l x_k,$$

where the  $a_{k,l}^{i,j}$  are real constants and  $(x_1, \dots, x_n)$  are the linear coordinates associated to  $\mathbb{B}$ .

For any linear endomorphism  $A$  on  $V$  we denote by  $\tilde{A}$  the associated linear vector field on  $V$ .

The key point is that if  $h$  is a quadratic contravariant pseudo-Hessian bivector field on  $V$  then its divergence is a linear vector field, i.e.,  $\text{div}_{\nabla}(h) = \tilde{L}^h$  where  $L^h$  is a linear endomorphism of  $V$ . Moreover, if  $F = (A, u)$  is an affine transformation of  $V$  then  $\text{div}_{\nabla}(F_*h) = \tilde{A^{-1}L^hA}$ . So the Jordan form of  $L_h$  is an invariant of the quadratic contravariant pseudo-Hessian structure. By using Maple we can classify quadratic contravariant pseudo-Hessian structures on  $\mathbb{R}^2$ . The same approach has been used by [9] to classify quadratic Poisson structures on  $\mathbb{R}^4$ . Note that if  $h$  is a quadratic contravariant pseudo-Hessian tensor on  $\mathbb{R}^n$  then its associated Poisson tensor on  $\mathbb{C}^n$  is also quadratic.

**Theorem 5.1.** 1. Up to an affine isomorphism, there is two quadratic contravariant pseudo-Hessian structures on  $\mathbb{R}^2$  which are divergence free

$$h_1 = \begin{pmatrix} 0 & 0 \\ 0 & ux^2 \end{pmatrix} \quad \text{and} \quad h_2 = \begin{pmatrix} \frac{r^2x^2}{c} - 2rxy + cy^2 & \frac{r^3x^2}{c^2} - \frac{2r^2xy}{c} + ry^2 \\ \frac{r^3x^2}{c^2} - \frac{2r^2xy}{c} + ry^2 & -\frac{2r^3xy}{c^2} + \frac{r^4x^2}{c^3} + \frac{r^2y^2}{c} \end{pmatrix}.$$

2. Up to an affine isomorphism, there is two quadratic contravariant pseudo-Hessian structures on  $\mathbb{R}^2$  with the divergence equivalent to the Jordan form  $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$ ,

$$h_1 = \begin{pmatrix} cy^2 + xy & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad h_2 = \begin{pmatrix} \frac{1}{2}xy + cy^2 & \frac{y^2}{4} \\ \frac{y^2}{4} & 0 \end{pmatrix}.$$

3. Up to an affine isomorphism, there is five quadratic contravariant pseudo-Hessian structures on  $\mathbb{R}^2$  with diagonalizable divergence

$$h_1 = \begin{pmatrix} ax^2 & 0 \\ 0 & by^2 \end{pmatrix}, \quad h_2 = \begin{pmatrix} ax^2 + by^2 & 0 \\ 0 & 0 \end{pmatrix}, \quad h_3 = \begin{pmatrix} ax^2 & axy \\ axy & ay^2 \end{pmatrix},$$

$$h_4 = \begin{pmatrix} \frac{2r^2x^2}{c} - 2rxy + cy^2 & ry^2 \\ ry^2 & \frac{2r^2y^2}{c} \end{pmatrix} \quad \text{and} \quad h_5 = \begin{pmatrix} (\frac{2p^2}{u} + \frac{q}{2})x^2 + \frac{pqxy}{u} + \frac{q^2y^2}{4u} & px^2 + qxy - \frac{pqy^2}{2u} \\ px^2 + qxy - \frac{pqy^2}{2u} & (\frac{2p^2}{u} + \frac{q}{2})y^2 + ux^2 - 2pxy \end{pmatrix}.$$

4. Up to an affine isomorphism, there is a unique quadratic pseudo-Hessian structure on  $\mathbb{R}^2$  with the divergence having non real eigenvalues

$$h = \begin{pmatrix} -2pxy - ux^2 + uy^2 & px^2 - py^2 - 2uxy \\ px^2 - py^2 - 2uxy & 2pxy + ux^2 - uy^2 \end{pmatrix}.$$

**Example 5.2.** The study of quadratic contravariant pseudo-Hessian structures on  $\mathbb{R}^3$  is more complicated and we give here a class of quadratic pseudo-Hessian structures on  $\mathbb{R}^3$  of the form  $\tilde{A} \odot \tilde{I}_3$  where  $\tilde{A}$  is linear.

1.  $A$  is diagonal:

$$h_1 = \begin{pmatrix} x^2 & xy & xz \\ xy & y^2 & yz \\ xz & yz & z^2 \end{pmatrix} \quad \text{and} \quad h_2 = \begin{pmatrix} x^2 & xy & 0 \\ xy & y^2 & 0 \\ 0 & 0 & -z^2 \end{pmatrix}.$$

2.  $A = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 0 \\ 0 & 0 & b \end{pmatrix}$ :

$$h_3 = \begin{pmatrix} 2x(y - px) & (y - px)y + pyx & pxz + (y - px)z \\ (y - px)y + pyx & 2py^2 & 2pyz \\ pxz + (y - px)z & 2pyz & 2pz^2 \end{pmatrix},$$

$$h_4 = \begin{pmatrix} 2x(y + px) & (y + px)y - pyx & pxz + (y + px)z \\ (y + px)y + pyx & -2py^2 & 0 \\ pxz + (y + px)z & 0 & 2pz^2 \end{pmatrix}.$$

## 6. Right-invariant contravariant pseudo-Hessian structures on Lie groups

Let  $(\mathfrak{g}, \bullet)$  be a left symmetric algebra, i.e., for any  $u, v, w \in \mathfrak{g}$ ,

$$\text{ass}(u, v, w) = \text{ass}(v, u, w) \quad \text{and} \quad \text{ass}(u, v, w) = (u \bullet v) \bullet w - u \bullet (v \bullet w).$$

This implies that  $[u, v] = u \bullet v - v \bullet u$  is a Lie bracket on  $\mathfrak{g}$  and  $L : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$ ,  $u \mapsto L_u$  is a representation of the Lie algebra  $(\mathfrak{g}, [\ , \ ])$ . We denote by  $L_u$  the left multiplication by  $u$ .

We consider a connected Lie group  $G$  whose Lie algebra is  $(\mathfrak{g}, [\ , \ ])$  and we define on  $G$  a right invariant connection by

$$\nabla_{u^-} v^- = -(u \bullet v)^-, \quad (6.1)$$

where  $u^-$  is the right vector field associated to  $u \in \mathfrak{g}$ . This connection is torsionless and without curvature and hence  $(G, \nabla)$  is an affine manifold. Let  $r \in \mathfrak{g} \otimes \mathfrak{g}$  which is symmetric and let  $r^-$  be the associated right invariant symmetric bivector field.

**Proposition 6.1.**  $(G, \nabla, r^-)$  is a contravariant pseudo-Hessian manifold if and only if, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$[[r, r]](\alpha, \beta, \gamma) := \langle \gamma, r_{\#}([\alpha, \beta]_r) - [r_{\#}(\alpha), r_{\#}(\beta)] \rangle = 0, \quad (6.2)$$

where

$$[\alpha, \beta]_r = L_{r_\#(\alpha)}^* \beta - L_{r_\#(\beta)}^* \alpha \quad \text{and} \quad \langle L_u^* \alpha, v \rangle = - \langle \alpha, u \bullet v \rangle.$$

In this case, the product on  $\mathfrak{g}$  given by  $\alpha \beta = L_{r_\#(\alpha)}^* \beta$  is left symmetric,  $[\ , \ ]_r$  is a Lie bracket and  $r_\#$  is a morphism of Lie algebras.

*Proof.* Note first that for any  $\alpha \in \mathfrak{g}^*$ ,  $r_\#^-(\alpha^-) = (r_\#(\alpha))^-$  and  $\nabla_u \alpha^- = -(L_u^* \alpha)^-$  and hence, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$\nabla_{r_\#^-(\alpha^-)}(r^-)(\beta^-, \gamma^-) = r(L_{r_\#(\alpha)}^* \beta, \gamma) + r(\beta, L_{r_\#(\alpha)}^* \gamma).$$

So,  $(G, \nabla, r^-)$  is a contravariant pseudo-Hessian manifold if and only if, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$\begin{aligned} 0 &= r(L_{r_\#(\alpha)}^* \beta, \gamma) + r(\beta, L_{r_\#(\alpha)}^* \gamma) - r(L_{r_\#(\beta)}^* \alpha, \gamma) - r(\alpha, L_{r_\#(\beta)}^* \gamma) \\ &= \langle \gamma, r_\#([\alpha, \beta]_r) - r_\#(\alpha) \bullet r_\#(\beta) + r_\#(\beta) \bullet r_\#(\alpha) \rangle \\ &= \langle \gamma, r_\#([\alpha, \beta]_r) - [r_\#(\alpha), r_\#(\beta)] \rangle \end{aligned}$$

and the first part of the proposition follows. Suppose now that  $r_\#([\alpha, \beta]_r) = [r_\#(\alpha), r_\#(\beta)]$  for any  $\alpha, \beta \in \mathfrak{g}^*$ . Then, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$\text{ass}(\alpha, \beta, \gamma) - \text{ass}(\beta, \alpha, \gamma) = L_{r_\#([\alpha, \beta]_r)}^* \gamma - L_{r_\#(\alpha)}^* L_{r_\#(\beta)}^* \gamma + L_{r_\#(\beta)}^* L_{r_\#(\alpha)}^* \gamma = 0.$$

This completes the proof.  $\square$

**Definition 6.2.** 1. Let  $(\mathfrak{g}, \bullet)$  be a left symmetric algebra. A symmetric bivector  $r \in \mathfrak{g} \otimes \mathfrak{g}$  satisfying  $[[r, r]] = 0$  is called a *S-matrix*.  
2. A left symmetric algebra  $(\mathfrak{g}, \bullet, r)$  endowed with a S-matrix is called a *contravariant pseudo-Hessian algebra*.

Let  $(\mathfrak{g}, \bullet, r)$  be a contravariant pseudo-Hessian algebra,  $[u, v] = u \bullet v - v \bullet u$  and  $G$  a connected Lie group with  $(\mathfrak{g}, [\ , \ ])$  as a Lie algebra. We have shown that  $G$  carries a right invariant contravariant pseudo-Hessian structure  $(\nabla, r^-)$ . On the other hand, in Section 3, we have associated to  $(\nabla, r^-)$  a flat connection  $\overline{\nabla}$ , a complex structure  $J$  and a Poisson tensor  $\Pi$  on  $TG$ . Now we will show that  $TG$  carries a structure of Lie group and the triple  $(\overline{\nabla}, J, \Pi)$  is right invariant. This structure of Lie group on  $TG$  is different from the usual one defined by the adjoint action of  $G$  on  $\mathfrak{g}$ .

Let us start with a general algebraic construction which is interesting on its own. Let  $(\mathfrak{g}, \bullet)$  be a left symmetric algebra, put  $\Phi(\mathfrak{g}) = \mathfrak{g} \times \mathfrak{g}$  and define a product  $\star$  and a bracket on  $\Phi(\mathfrak{g})$  by

$$(a, b) \star (c, d) = (a \bullet c, a \bullet d) \quad \text{and} \quad [(a, b), (c, d)] = ([a, c], a \bullet d - c \bullet b),$$

for any  $(a, b), (c, d) \in \Phi(\mathfrak{g})$ . It is easy to check that  $\star$  is left symmetric,  $[\ , \ ]$  is the commutator of  $\star$  and hence is a Lie bracket. We define also  $J_0 : \Phi(\mathfrak{g}) \rightarrow \Phi(\mathfrak{g})$  by  $J_0(a, b) = (b, -a)$ . It is also a straightforward computation to check that

$$N_{J_0}((a, b), (c, d)) = [J_0(a, b), J_0(c, d)] - J_0([a, b], J_0(c, d)) - J_0[J_0(a, b), (c, d)] - [(a, b), (c, d)] = 0.$$

For  $r \in \otimes^2 \mathfrak{g}$  symmetric, we define  $R \in \otimes^2 \Phi(\mathfrak{g})$  by

$$R((\alpha_1, \beta_1), (\alpha_2, \beta_2)) = r(\alpha_1, \beta_2) - r(\alpha_2, \beta_1), \quad (6.3)$$

for any  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathfrak{g}^*$ . We have obviously that  $R_\#(\alpha_1, \beta_1) = (-r_\#(\beta_1), r_\#(\alpha_1))$ .

**Proposition 6.3.**  $[[r, r]] = 0$  if and only if  $[R, R] = 0$ , where  $[R, R]$  is the Schouten bracket associated to the Lie algebra structure of  $\Phi(\mathfrak{g})$  and given by

$$[R, R](\alpha, \beta, \gamma) = \oint_{\alpha, \beta, \gamma} \langle \gamma, [R_{\#}(\alpha), R_{\#}(\beta)] \rangle, \quad \alpha, \beta, \gamma \in \Phi^*(\mathfrak{g}).$$

*Proof.* For any  $\alpha = (\alpha_1, \alpha_2), \beta = (\beta_1, \beta_2), \gamma = (\gamma_1, \gamma_2) \in \Phi(\mathfrak{g})^*$ ,

$$\begin{aligned} \langle \gamma, [R_{\#}(\alpha), R_{\#}(\beta)] \rangle &= \langle (\gamma_1, \gamma_2), [(-r_{\#}(\alpha_2), r_{\#}(\alpha_1)), (-r_{\#}(\beta_2), r_{\#}(\beta_1))] \rangle \\ &= \langle \gamma_1, [r_{\#}(\alpha_2), r_{\#}(\beta_2)] \rangle - \langle \gamma_2, r_{\#}(\alpha_2) \bullet r_{\#}(\beta_1) \rangle + \langle \gamma_2, r_{\#}(\beta_2) \bullet r_{\#}(\alpha_1) \rangle \\ &= \langle \gamma_1, [r_{\#}(\alpha_2), r_{\#}(\beta_2)] \rangle + \langle \beta_1, r_{\#}(\mathbb{L}_{r_{\#}(\alpha_2)}^* \gamma_2) \rangle - \langle \alpha_1, r_{\#}(\mathbb{L}_{r_{\#}(\beta_2)}^* \gamma_2) \rangle, \\ \langle \beta, [R_{\#}(\gamma), R_{\#}(\alpha)] \rangle &= \langle \beta_1, [r_{\#}(\gamma_2), r_{\#}(\alpha_2)] \rangle + \langle \alpha_1, r_{\#}(\mathbb{L}_{r_{\#}(\gamma_2)}^* \beta_2) \rangle - \langle \gamma_1, r_{\#}(\mathbb{L}_{r_{\#}(\alpha_2)}^* \beta_2) \rangle \\ \langle \alpha, [R_{\#}(\beta), R_{\#}(\gamma)] \rangle &= \langle \alpha_1, [r_{\#}(\beta_2), r_{\#}(\gamma_2)] \rangle + \langle \gamma_1, r_{\#}(\mathbb{L}_{r_{\#}(\beta_2)}^* \alpha_2) \rangle - \langle \beta_1, r_{\#}(\mathbb{L}_{r_{\#}(\gamma_2)}^* \alpha_2) \rangle. \end{aligned}$$

So

$$[R, R](\alpha, \beta, \gamma) = -[[r, r]](\beta_2, \gamma_2, \alpha_1) - [[r, r]](\gamma_2, \alpha_2, \beta_1) - [[r, r]](\alpha_2, \beta_2, \gamma_1)$$

and the result follows.  $\square$

Let  $G$  be a Lie group whose Lie algebra is  $(\mathfrak{g}, [\cdot, \cdot])$  and let  $\rho : G \rightarrow \text{GL}(\mathfrak{g})$  be the homomorphism of groups such that  $d_e \rho = L$  where  $L : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$  is the representation associated to  $\bullet$ . Then the product

$$(g, u) \cdot (h, v) = (gh, u + \rho(g)(v)), \quad g, h \in G, u, v \in \mathfrak{g}$$

induces a Lie group structure on  $G \times \mathfrak{g}$  whose Lie algebra is  $(\Phi(\mathfrak{g}), [\cdot, \cdot])$ . The complex endomorphism  $J_0$  and the left symmetric product  $\star$  induce a right invariant complex tensor  $J_0^-$  and a right invariant connection  $\widetilde{\nabla}$  given by

$$J_0^-(a, b)^- = (b, -a)^- \quad \text{and} \quad \widetilde{\nabla}_{(a, b)^-}(c, d)^- = -((a, b) \star (c, d))^-.$$

Let  $r \in \otimes^2 \mathfrak{g}$  symmetric such that  $[[r, r]] = 0$ ,  $r^-$  the associated right invariant symmetric bivector field and  $\nabla$  the affine connection given by (6.1). Then  $(G, \nabla, r^-)$  is a contravariant pseudo-Hessian manifold and let  $\overline{\nabla}$ ,  $J$  and  $\Pi$  be the associated structure on  $TG$  defined in Section 3.

**Theorem 6.4.** *If we identify  $TG$  with  $G \times \mathfrak{g}$  by  $u_g \rightarrow (g, T_g R_{g^{-1}} u_g)$  we denote also by  $\Pi$ ,  $\overline{\nabla}$  and  $J$  the images of  $\Pi$ ,  $\overline{\nabla}$  and  $J$  under this identification then  $\Pi = R^-$ ,  $\overline{\nabla} = \widetilde{\nabla}$  and  $J = J_0^-$ .*

To prove this theorem, we need some preparation.

**Proposition 6.5.** *Let  $(G, \nabla)$  be a Lie group endowed with a right invariant connection and  $\gamma : [0, 1] \rightarrow G$  a curve. Let  $V : [0, 1] \rightarrow TG$  be a vector field along  $\gamma$ . We define  $\mu : [0, 1] \rightarrow \mathfrak{g}$  and  $W : [0, 1] \rightarrow \mathfrak{g}$  by*

$$\mu(t) = T_{\gamma(t)} R_{\gamma(t)^{-1}}(\gamma'(t)) \quad \text{and} \quad W(t) = T_{\gamma(t)} R_{\gamma(t)^{-1}}(V(t)).$$

*Then  $V$  is parallel along  $\gamma$  with respect  $\nabla$  if and only if*

$$W'(t) - \mu(t) \bullet W(t) = 0,$$

*where  $u \bullet v = -(\nabla_u v^-)(e)$ .*

*Proof.* We consider  $(u_1, \dots, u_n)$  a basis of  $\mathfrak{g}$  and  $(X_1, \dots, X_n)$  the corresponding right invariant vector fields. Then

$$\begin{cases} \mu(t) = \sum_{i=1}^n \mu_i(t) u_i, & W(t) = \sum_{i=1}^n W_i(t) u_i, \\ \gamma'(t) = \sum_{i=1}^n \mu_i(t) X_i, & V(t) = \sum_{i=1}^n W_i(t) X_i. \end{cases}$$

Then

$$\begin{aligned} \nabla_t V(t) &= \sum_{i=1}^n W'_i(t) X_i + \sum_{i=1}^n W_i(t) \nabla_{\gamma'(t)} X_i \\ &= \sum_{i=1}^n W'_i(t) X_i + \sum_{i,j=1}^n W_i(t) \mu_j(t) \nabla_{X_j} X_i \\ &= \sum_{i=1}^n W'_i(t) X_i - \sum_{i,j=1}^n W_i(t) \mu_j(t) (u_j \bullet u_i)^- \\ &= (W'(t) - \mu(t) \bullet W(t))^- \end{aligned}$$

and the result follows having in mind that  $u^-$  is the right invariant vector field associated to  $u \in \mathfrak{g}$ .  $\square$

Let  $(G, \nabla)$  be a Lie group endowed with a right invariant connection. Then  $\nabla$  induces a splitting of  $TTG = \ker dp \oplus \mathcal{H}$ . For any tangent vector  $X \in T_g G$ , we denote by  $X^v, X^h \in T_{(g,u)} TG$  the vertical and the horizontal lift of  $X$ .

**Proposition 6.6.** *If we identify  $TG$  to  $G \times \mathfrak{g}$  by  $X_g \mapsto (g, T_g R_{g^{-1}}(X_g))$  then for any  $X \in T_g G$ ,*

$$X^v(g, u) = (0, T_g R_{g^{-1}}(X)) \quad \text{and} \quad X^h(g, u) = (X, T_g R_{g^{-1}}(X) \bullet u).$$

*Proof.* The first relation is obvious. Recall that the horizontal lift of  $X$  at  $u_g \in TG$  is given by:

$$X^h(u_g) = \frac{d}{dt} \Big|_{t=0} V(t)$$

where  $V : [0, 1] \rightarrow TG$  is the parallel vector field along  $\gamma : [0, 1] \rightarrow G$  a curve such that  $\gamma(0) = g$  and  $\gamma'(0) = X$ . If we denote by  $\Theta_R : TG \rightarrow G \times \mathfrak{g}$  the identification  $u_g \mapsto (g, T_g R_{g^{-1}}(u_g))$  then, by virtue of Proposition 6.5,

$$T_{u_g} \Theta_R(X^h) = \frac{d}{dt} \Big|_{t=0} (\gamma(t), W(t)) = (X, T_g R_{g^{-1}}(X) \bullet u).$$

$\square$

We consider now a left symmetric algebra  $(\mathfrak{g}, \bullet)$ ,  $G$  a connected Lie group associated to  $(\mathfrak{g}, [\ , \ ])$ ,  $\nabla$  the right invariant affine connection associated to  $\bullet$ . We have seen that  $G \times \mathfrak{g}$  has a structure of Lie group. We identify  $TG$  to  $G \times \mathfrak{g}$  and, for any vector field  $X$  on  $G$ , we denote by  $X^v$  and  $X^h$  the vector fields on  $G \times \mathfrak{g}$  obtained by the identification from the horizontal and the vertical lift of  $X$ . For  $a, b \in \mathfrak{g}$ ,  $\alpha, \beta \in \mathfrak{g}^*$ ,  $\alpha^-$  (resp.  $\alpha^+$ ) is the right invariant vector field (resp. 1-form) on  $G$  associated to  $a$  (resp.  $\alpha$ ),  $(a, b)^-$  (resp.  $(\alpha, \beta)^-$ ) the right invariant vector field (resp. 1-form) on  $G \times \mathfrak{g}$  associated to  $(a, b)$  (resp.  $(\alpha, \beta)$ ).

**Proposition 6.7.** For any  $(a, b) \in \mathfrak{g} \times \mathfrak{g}$  and  $(\alpha, \beta) \in \mathfrak{g}^* \times \mathfrak{g}^*$ ,

$$(a, b)^- = (a^-)^h + (b^-)^v \quad \text{and} \quad (\alpha, \beta)^- = (\alpha^-)^h + (\beta^-)^v.$$

*Proof.* We have

$$\begin{aligned} (a, b)^-(g, u) &= T_{(e,0)}R_{(g,u)}(a, b) \\ &= \frac{d}{dt}\bigg|_{t=0} (\exp(ta), tb)(g, u) \\ &= \frac{d}{dt}\bigg|_{t=0} (\exp(ta)g, tb + \rho(\exp(ta))(u)) \\ &= (a^-(g), b + a \bullet u) \\ &= (a^-(g), T_g R_{g^{-1}}(a^-(g)) \bullet u) + (0, T_g R_{g^{-1}}(b^-(g))) \\ &= (a^-)^h(g, u) + (b^-)^v(g, u). \quad (\text{Proposition 6.5}) \end{aligned}$$

The second relation can be deduced easily from the first one. □

*Proof of Theorem 6.4 .*

*Proof.* Let  $\Pi$  be the Poisson tensor on  $G \times \mathfrak{g}$  associated to  $r^-$ . Then, by using the precedent proposition,

$$\begin{aligned} \Pi((\alpha_1, \beta_1)^-, (\alpha_2, \beta_2)^-) &= \Pi((\alpha_1^-)^h + (\beta_1^-)^v, (\alpha_2^-)^h + (\beta_2^-)^v) \\ &= r^-(\alpha_1^-, \beta_2^-) - r^-(\alpha_2^-, \beta_1^-) \\ &= r(\alpha_1, \beta_2) - r(\alpha_2, \beta_1) \\ &= R^-((\alpha_1, \beta_1)^-, (\alpha_2, \beta_2)^-). \end{aligned}$$

In the same way,

$$\begin{aligned} J_0^-(a, b)^- &= (b, -a)^- = (b^-)^h - (a^-)^v, \\ J(a, b)^- &= (b^-)^h - (a^-)^v, \\ \bar{\nabla}_{(a,b)^-}(c, d)^- &= (\nabla_a c^-)^h + (\nabla_a d^-)^v = -((a \bullet c)^-)^h - ((a \bullet d)^-)^v = -((a, b).(c, d))^+ = \widetilde{\nabla}_{(a,b)^-}(c, d)^-. \end{aligned}$$

□

Let  $(\mathfrak{g}, \bullet)$  be a left symmetric algebra,  $(M, \nabla)$  and affine manifold and  $\rho : \mathfrak{g} \rightarrow \Gamma(TM)$  a linear map such that  $\rho(u \bullet v) = \nabla_{\rho(u)}\rho(v)$ . Then  $\rho$  defines an action on  $M$  of the Lie algebra  $(\mathfrak{g}, [\ , \ ])$ . We consider  $\rho^l : \Phi(\mathfrak{g}) \rightarrow \Gamma(TTM)$ ,  $(u, v) \rightarrow \rho(u)^h + \rho^v(v)$ . It is easy to check that

$$\rho^l([a, b]) = [\rho^l(a), \rho^l(b)].$$

Let  $r \in \otimes^2 \mathfrak{g}$  satisfying  $[[r, r]] = 0$  and  $R \in \otimes^2 \Phi(\mathfrak{g})$  given by (6.3).

**Theorem 6.8.** The bivector field on  $TM$  associated to  $\rho(r)$  is  $\rho^l(R)$  which is a Poisson tensor and  $(M, \nabla, \rho(r))$  is a contravariant pseudo-Hessian manifold.

*Proof.* Let  $(e_1, \dots, e_n)$  a basis of  $\mathfrak{g}$  and  $E_i = (e_i, 0)$  and  $F_i = (0, e_i)$ . Then  $(E_1, \dots, E_n, F_1, \dots, F_n)$  is a basis of  $\Phi(\mathfrak{g})$ . Then

$$r = \sum_{i,j} r_{i,j} e_i \otimes e_j \quad \text{and} \quad R = \sum_{i,j} r_{i,j} (E_i \otimes F_j - F_i \otimes E_j).$$

So

$$\rho(r) = \sum_{i,j=1}^n r_{i,j} \rho(e_i) \otimes \rho(e_j) \quad \text{and} \quad \rho^l(R) = \sum_{i,j=1}^n r_{i,j} (\rho(e_i)^h \otimes \rho(e_j)^v - \rho(e_i)^v \otimes \rho(e_j)^h).$$

Then for any  $\alpha, \beta \in \Omega^1(M)$

$$\rho^l(R)(\alpha^v, \beta^v) = \rho^l(R)(\alpha^h, \beta^h) = 0 \quad \text{and} \quad \rho^l(R)(\alpha^h, \beta^v) = \rho(r)(\alpha, \beta) \circ p.$$

According to Proposition 6.3,  $R$  is a solution of the classical Yang-Baxter equation and hence  $\rho^l(R)$  is a Poisson tensor. By using Theorem 3.3, we get that  $(M, \nabla, \rho(r))$  is a contravariant pseudo-Hessian manifold.  $\square$

- Example 6.9.** 1. Let  $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{R})$  be the Lie algebra of  $n$ -square matrices. It has a structure of left symmetric algebra given by  $A \bullet B = BA$ . Let  $\rho : \mathfrak{g} \rightarrow \Gamma(T\mathbb{R}^n)$  given by  $\rho(A) = A$ . Then  $\rho(A \bullet B) = \nabla_A B$ , where  $\nabla$  is the canonical connection of  $\mathbb{R}$ . According to Theorem 6.8, any  $S$ -matrix on  $\mathfrak{g}$  gives rise to a quadratic contravariant pseudo-Hessian structure on  $\mathbb{R}^n$ .
2. More generally, let  $(M, \nabla)$  be an affine manifold and  $\mathfrak{g}$  the finite dimensional Lie algebra of affine vector fields. Recall that  $X \in \mathfrak{g}$  if for any  $Y, Z \in \Gamma(TM)$ ,

$$[X, \nabla_Y Z] = \nabla_{[X,Y]} Z + \nabla_Y [X, Z].$$

Since the curvature and the torsion of  $\nabla$  vanish this is equivalent to

$$\nabla_{\nabla_Y Z} X = \nabla_Y \nabla_Z X.$$

From this relation, one can see easily that, for any  $X, Y \in \mathfrak{g}$ ,  $X \bullet Y := \nabla_X Y \in \mathfrak{g}$  and  $(\mathfrak{g}, \bullet)$  is an associative finite dimensional Lie algebra which acts on  $M$  by  $\rho(X) = X$ . Moreover,  $\rho(X \bullet Y) = \nabla_X Y$ . According to Theorem 6.8, any  $S$ -matrix on  $\mathfrak{g}$  gives rise to a contravariant pseudo-Hessian structure on  $M$ .

### Classification of two-dimensional contravariant pseudo-Hessian algebras

Using the classification of two-dimensional non-abelian left symmetric algebras given in [5] and the classification of abelian left symmetric algebras given in [14], we give a classification (over the field  $\mathbb{R}$ ) of 2-dimensional contravariant pseudo-Hessian algebras. We proceed as follows:

1. For any left symmetric 2-dimensional algebra  $\mathfrak{g}$ , we determine its automorphism group  $\text{Aut}(\mathfrak{g})$  and the space of  $S$ -matrices on  $\mathfrak{g}$ , we denote by  $\mathcal{A}(\mathfrak{g})$ .
2. We give the quotient  $\mathcal{A}(\mathfrak{g}) / \sim$  where  $\sim$  is the equivalence relation:

$$r^1 \sim r^2 \iff \exists A \in \text{Aut}(\mathfrak{g}) \text{ or } \exists \lambda \in \mathbb{R} \text{ such that } r_{\sharp}^2 = A \circ r_{\sharp}^1 \circ A^t \text{ or } r^2 = \lambda r^1.$$

$(\mathfrak{g}, \cdot)$	$\text{Aut}(\mathfrak{g})$	$\mathcal{A}(\mathfrak{g})/\sim$
$b_{1,\alpha \neq -1,1}$ $e_2.e_1 = e_1, e_2.e_2 = \alpha e_2$	$\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$b_{1,\alpha=-1}$ $e_2.e_1 = e_1, e_2.e_2 = -e_2$	$\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} b & 1 \\ 1 & 0 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix};$ $r_{\#}^3 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^4 = 0$
$b_{1,\alpha=1}$ $e_2.e_1 = e_1, e_2.e_2 = e_2$	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, ab \neq 0$	$r_{\#}^1 = \begin{pmatrix} 1 & c \\ c & c^2 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$b_{2,\beta \neq 0,1,2}$ $e_1.e_2 = \beta e_1, e_2.e_1 = (\beta - 1)e_1, e_2.e_2 = \beta e_2$	$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$b_{2,\beta=1}$ $e_1.e_2 = e_1, e_2.e_2 = e_2$	$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 1 & c \\ c & c^2 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$b_{2,\beta=2}$ $e_1.e_2 = 2e_1,$ $e_2.e_1 = e_1, e_2.e_2 = 2e_2$	$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 1 & 0 \\ 0 & c \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$b_3$ $e_2.e_1 = e_1, e_2.e_2 = e_1 + e_2$	$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$	$r_{\#}^1 = \begin{pmatrix} 1/2 & 1 \\ 1 & 1 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^3 = 0$
$b_4$ $e_1.e_1 = 2e_1, e_1.e_2 = e_2,$ $e_2.e_2 = e_1$	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}; \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$r_{\#}^1 = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^3 = 0$
$b_5$ $e_1.e_2 = e_1, e_2.e_2 = e_1 + e_2$	$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$	$r_{\#}^1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^2 = 0$
$As_2^1$ $e_1.e_1 = e_2$	$\begin{pmatrix} a & 0 \\ b & a^2 \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; r_{\#}^3 = 0$
$As_2^4$ $e_1.e_1 = e_1, e_1.e_2 = e_2,$ $e_2.e_2 = e_2$	$\begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}, a \neq 0$	$r_{\#}^1 = \begin{pmatrix} 0 & 1 \\ 1 & c \end{pmatrix}; r_{\#}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix};$ $r_{\#}^3 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; r_{\#}^4 = 0$

We end this paper by giving another proof to Lemma 4.4.

*Proof.* For any  $\gamma : [0, 1] \longrightarrow G \times G, t \mapsto (\gamma_1(t), \gamma_2(t))$  with  $\gamma(0) = (a, b)$  and  $\gamma(1) = (c, d)$ ,

$$\tau_{m(\gamma)}(T_{(a,b)}m(u, v)) = T_{(c,d)}m(\tau_{\gamma}(u, v)),$$

where  $\tau_{\gamma} : T_{(a,b)}(G \times G) \longrightarrow T_{(c,d)}(G \times G)$  and  $\tau_{m\gamma} : T_{ab}G \longrightarrow T_{cd}G$  are the parallel transports.

But

$$T_{(a,b)}m(u, v) = T_a R_b(u) + T_b L_a(v) \quad \text{and} \quad \tau_{\gamma}(u, v) = (\tau_{\gamma_1}(u), \tau_{\gamma_2}(v)).$$



So we get

$$\tau_{\gamma_1\gamma_2}(T_a R_b(u)) + \tau_{\gamma_1\gamma_2}(T_b L_a(v)) = T_c R_d(\tau_{\gamma_1}(u)) + T_d L_c(\tau_{\gamma_2}(v)).$$

If we take  $v = 0$  and  $\gamma_2(t) = b = d$ . We get

$$\tau_{\gamma_1 b}(T_a R_b(u)) = T_c R_b(\tau_{\gamma_1}(u))$$

and hence  $\nabla$  is right invariant. In the same way we get that  $\nabla$  is left invariant. And finally

$$\tau_{\gamma_1\gamma_2}(T_a R_b(u)) = T_c R_d(\tau_{\gamma_1}(u)) \quad \text{and} \quad \tau_{\gamma_1\gamma_2}(T_b L_a(v)) = T_d L_c(\tau_{\gamma_2}(v)).$$

If we take  $\gamma_2 = \gamma_1^{-1}$  we get that

$$\tau_{\gamma_1}(u) = T_a R_{a^{-1}c}(u) = T_a L_{ca^{-1}}(u).$$

This implies that the adjoint representation is trivial and hence  $G$  must be abelian.  $\square$

## References

- [1] C. Bai. *Bijective 1-cocycles and classification of 3-dimensional left-symmetric algebras*, Communications in Algebra Volume **37**, 1016-1057 (2009).
- [2] Bai, C., *Left-Symmetric Bialgebras and an Analogue of the Classical Yang-Baxter Equation*, Communication in Contemporary Mathematics, 2008, Vol. **10**; Numb. 2, 221-260.
- [3] S. Benayadi and M. Boucetta. *On para-Kähler Lie algebroids and contravariant pseudo-Hessian structures*. Mathematische Nachrichten. (2019);1-26.
- [4] S. Benayadi and M. Boucetta. *On Para-Kähler and Hyper-Kähler Lie algebras*. Journal of Algebra **436** (2015) 61-101.
- [5] D. Burde *Simple left-symmetric algebras with solvable Lie algebra*, manuscripta math. **95**, 397 - 411 (1998).
- [6] C. Calvin and L. M. C. Schochet, *Global analysis on foliated space*. MISRP (2006).
- [7] Dombrowski P., *On the geometry of the tangent bundle*. J. Reme Angew. Math. 210 (1962), 73-78.
- [8] Dufour, Jean-Paul, and Nguyen Tien Zung. *Poisson structures and their normal forms*. Vol. **242**. Springer Science and Business Media, 2006.
- [9] J. P. Dufour and A. Haraki, *Rotationnels et structures de Poisson quadratiques*. C. R. Acad. Sci. Paris Série. I Math., 312(1):137-140, 1991.
- [10] R. L. FERNANDES, *Lie algebroids, holonomy and characteristic classes*, Adv. in Math. **170** (2002), 119-179.
- [11] Christian Kassel, *Homology and cohomology of associative algebras. A concise introduction to cyclic homology*. Thematic school . August 2004 at ICTP, Trieste (Italy), 2006. jcel-00119891.
- [12] K. MACKENZIE, *Lie Groupoids and Lie Algebroids in Differential Geometry*, London Math. Soc. Lecture Notes Ser. **124**, Cambridge Univ. Press, Cambridge, 1987.
- [13] H. Shima. *The geometry of Hessian structures*. World Scientific Publishing (2007).
- [14] I.S. Rakhimov, I.M. Rikhsiboev, W. Basri. *Complete lists of low dimensional complex associative algebras*, arXiv:0910.0932v2
- [15] A. WEINSTEIN, *The local structure of Poisson manifolds*, J. Differential Geom. Volume **18**, Number 3 (1983), 523-557.