

# An approach to the Herzog-Schönheim conjecture using automata

Fabienne Chouraqui

## Abstract

Let  $G$  be a group and  $H_1, \dots, H_s$  be subgroups of  $G$  of indices  $d_1, \dots, d_s$  respectively. In 1974, M. Herzog and J. Schönheim conjectured that if  $\{H_i \alpha_i\}_{i=1}^{i=s}$ ,  $\alpha_i \in G$ , is a coset partition of  $G$ , then  $d_1, \dots, d_s$  cannot be distinct. In this paper, we present a new approach to the Herzog-Schönheim conjecture based on automata and present a translation of the conjecture as a problem on automata.

## 1 Introduction

Let  $G$  be a group and  $H_1, \dots, H_s$  be subgroups of  $G$ . If there exist  $\alpha_i \in G$  such that  $G = \bigcup_{i=1}^{i=s} H_i \alpha_i$ , and the sets  $H_i \alpha_i$ ,  $1 \leq i \leq s$ , are pairwise disjoint, then  $\{H_i \alpha_i\}_{i=1}^{i=s}$  is a *coset partition of  $G$*  (or a *disjoint cover of  $G$* ). In this case, all the subgroups  $H_1, \dots, H_s$  can be assumed to be of finite index in  $G$  [21, 18]. We denote by  $d_1, \dots, d_s$  the indices of  $H_1, \dots, H_s$  respectively. The coset partition  $\{H_i \alpha_i\}_{i=1}^{i=s}$  has *multiplicity* if  $d_i = d_j$  for some  $i \neq j$ .

If  $G$  is the infinite cyclic group  $\mathbb{Z}$ , a coset partition of  $\mathbb{Z}$  is  $\{d_i \mathbb{Z} + r_i\}_{i=1}^{i=s}$ ,  $r_i \in \mathbb{Z}$ , with each  $d_i \mathbb{Z} + r_i$  the residue class of  $r_i$  modulo  $d_i$ . These coset partitions of  $\mathbb{Z}$  were first introduced by P. Erdős [12] and he conjectured that if  $\{d_i \mathbb{Z} + r_i\}_{i=1}^{i=s}$ ,  $r_i \in \mathbb{Z}$ , is a coset partition of  $\mathbb{Z}$ , then the largest index  $d_s$  appears at least twice. Erdős' conjecture was proved independently by H. Davenport with R. Rado and L. Mirsky with D. Newman using analysis of complex function [13, 21, 22]. Furthermore, it was proved that the largest index  $d_s$  appears at least  $p$  times, where  $p$  is the smallest prime dividing  $d_s$  [21, 22, 34], that each index  $d_i$  divides another index  $d_j$ ,  $j \neq i$ , and that each index  $d_k$  that does not properly divide any other index appears at least twice [22]. We refer also to [25, 26, 27, 28, 35] for more details on coset partitions of  $\mathbb{Z}$  (also called covers of  $\mathbb{Z}$  by arithmetic progres-

sions) and to [14] for a proof of the Erdős' conjecture using group representations.

In 1974, M. Herzog and J. Schönheim extended Erdős' conjecture for arbitrary groups and conjectured that if  $\{H_i\alpha_i\}_{i=1}^s$ ,  $\alpha_i \in G$ , is a coset partition of  $G$ , then  $d_1, \dots, d_s$  cannot be distinct. In the 1980's, in a series of papers, M.A. Berger, A. Felzenbaum and A.S. Fraenkel studied the Herzog-Schönheim conjecture [2, 3, 4] and in [5] they proved the conjecture is true for the pyramidal groups, a subclass of the finite solvable groups. Coset partitions of finite groups with additional assumptions on the subgroups of the partition have been extensively studied. We refer to [6, 36, 37, 33]. In [19], the authors very recently proved that the conjecture is true for all groups of order less than 1440.

The common approach to the Herzog-Schönheim (HS) conjecture is to study it in finite groups. Indeed, given any group  $G$ , every coset partition of  $G$  induces a coset partition of a finite quotient group of  $G$  with the same indices [18]. In this paper, we present a completely different approach to the HS conjecture. The idea is to study it in free groups of finite rank and from there to provide answers for every group. This is possible since any finite or finitely generated group is a quotient group of a free group of finite rank and any coset partition of a quotient group  $F/N$  induces a coset partition of  $F$  with the same indices [7].

In order to study the Herzog-Schönheim conjecture in free groups of finite rank, we use the machinery of covering spaces. A pair  $(\tilde{X}, p)$  is a *covering space* of a topological space  $X$  if  $\tilde{X}$  is a path connected space,  $p : \tilde{X} \rightarrow X$  is an open continuous surjection and every  $x \in X$  has an open neighborhood  $U_x$  such that  $p^{-1}(U_x)$  is a disjoint union of open sets in  $\tilde{X}$ , each of which is mapped homeomorphically onto  $U_x$  by  $p$ . For each  $x \in X$ , the non-empty set  $Y_x = p^{-1}(x)$  is called *the fiber over  $x$*  and for all  $x, x' \in X$ ,  $|Y_x| = |Y_{x'}|$ . If the cardinal of a fiber is  $m$ , one says that  $(\tilde{X}, p)$  is a  *$m$ -sheeted covering* ( *$m$ -fold cover*) of  $X$  [16, 29].

The fundamental group of the bouquet with  $n$  leaves (or the wedge sum of  $n$  circles),  $X$ , is  $F_n$ , the free group of finite rank  $n$  and for any subgroup  $H$  of  $F_n$  of finite index  $d$ , there exists a  $d$ -sheeted covering space  $(\tilde{X}_H, p)$  with a fixed basepoint. The underlying graph of  $\tilde{X}_H$  is a directed labelled graph, with  $d$  vertices, called *the Schreier graph* and it can be seen as a finite complete bi-deterministic automaton; fixing the start and the end state at the basepoint, it recognises the set of elements in  $H$ . It is called *the Schreier coset diagram for  $F_n$  relative to the subgroup  $H$*  [32, p.107] or *the Schreier automaton for  $F_n$  relative to the subgroup  $H$*  [30, p.102]. The  $d$  vertices (or states) correspond to the  $d$  right cosets of  $H$ , any edge

(or transition) has the form  $Hg \xrightarrow{a} Hga$ ,  $g \in F_n$ ,  $a$  a generator of  $F_n$ , and it describes the right action of  $F_n$  on the right cosets of  $H$ . If we fix the start state at the basepoint  $(H)$ , and the end state at another vertex  $H\alpha$ , where  $\alpha$  denotes the label of some path from the start state to the end state, then this automaton recognises the set of elements in  $H\alpha$  and we call it *the Schreier automaton of  $H\alpha$*  and denote it by  $\tilde{X}_{H\alpha}$ .

In general, for any automaton  $M$ , with alphabet  $\Sigma$ , and  $d$  states, there exists a square matrix  $A$  of order  $d \times d$ , with  $a_{ij}$  equal to the number of directed edges from vertex  $i$  to vertex  $j$ ,  $1 \leq i, j \leq d$ . This matrix is non-negative and it is called *the transition matrix* [11]. If for every  $1 \leq i, j \leq d$ , there exists  $m \in \mathbb{Z}^+$  such that  $(A^m)_{ij} > 0$ , the matrix is *irreducible*. For  $A$  an irreducible non-negative matrix, *the period of  $A$*  is the gcd of all  $m \in \mathbb{Z}^+$  such that  $(A^m)_{ii} > 0$  (for any  $i$ ). If  $i$  and  $j$  denote respectively the start and end states of  $M$ , then the number of words of length  $k$  (in the alphabet  $\Sigma$ ) accepted by  $M$  is  $a_k = (A^k)_{ij}$ . *The generating function of  $M$*  is defined by  $p(z) = \sum_{k=0}^{k=\infty} a_k z^k$ . It is a rational function: the fraction of two polynomials in  $z$  with integer coefficients [11], [31, p.575].

In [9], we study the properties of the transition matrices and generating functions of the Schreier automata in the context of coset partitions of the free group. Let  $F_n = \langle \Sigma \rangle$ , and  $\Sigma^*$  the free monoid generated by  $\Sigma$ . Let  $\{H_i\alpha_i\}_{i=1}^{i=s}$  be a coset partition of  $F_n$  with  $H_i < F_n$  of index  $d_i > 1$ ,  $\alpha_i \in F_n$ ,  $1 \leq i \leq s$ . Let  $\tilde{X}_i$  denote the Schreier graph of  $H_i$ , with transition matrix  $A_i$  of period  $h_i \geq 1$  and  $\tilde{X}_{H_i\alpha_i}$  the Schreier automaton of  $H_i\alpha_i$ , with generating function  $p_i(z)$ ,  $1 \leq i \leq s$ . For each  $\tilde{X}_i$ ,  $A_i$  is a non-negative irreducible matrix and  $(A_i^k)_{bf}$ ,  $k \geq 0$ , counts the number of words of length  $k$  that belong to  $H_i\alpha_i \cap \Sigma^*$  (with  $b$  and  $f$  denoting the start and end state of  $H_i\alpha_i$  respectively). Since  $F_n$  is the disjoint union of the sets  $\{H_i\alpha_i\}_{i=1}^{i=s}$ , each element in  $\Sigma^*$  belongs to one and exactly one such set, so  $n^k$ , the number of words of length  $k$  in  $\Sigma^*$ , satisfies  $n^k = \sum_{i=1}^{i=s} a_{i,k}$ , for every

$k \geq 0$ , and moreover  $\sum_{k=0}^{k=\infty} n^k z^k = \sum_{i=1}^{i=s} p_i(z)$ . By using this kind of counting argument and studying the behaviour of the generating functions at their poles, we prove that if  $h = \max\{h_i \mid 1 \leq i \leq s\}$  is greater than 1, then there is a repetition of the maximal period  $h > 1$  and that, under certain conditions, the coset partition has multiplicity. Furthermore, we recover the Davenport-Rado result (or Mirsky-Newman result) for the Erdős' conjecture and some of its consequences.

In this paper, we deepen further our study of the transition matrices of the Schreier automata in the context of coset partitions of  $F_n$  and give some new conditions that ensure a coset partition of  $F_n$  has multiplicity.

**Theorem 1.** *Let  $F_n$  be the free group on  $n \geq 2$  generators. Let  $\{H_i\alpha_i\}_{i=1}^{i=s}$  be a coset partition of  $F_n$  with  $H_i < F_n$  of index  $d_i$ ,  $\alpha_i \in F_n$ ,  $1 \leq i \leq s$ , and  $1 < d_1 \leq \dots \leq d_s$ . Let  $\tilde{X}_i$  denote the Schreier graph of  $H_i$ , with transition matrix  $A_i$ , and period  $h_i \geq 1$ ,  $1 \leq i \leq s$ . Let  $H = \{h > 1 \mid \exists 1 \leq j \leq s, h_j = h\}$ . Assume  $H \neq \emptyset$  and different elements in  $H$  are pairwise coprime. Let  $r_h = |\{1 \leq j \leq s, \mid h_j = h\}|$ , the number of repetitions of  $h$ . If for some  $h \in H$ ,  $r_h = h$  or  $h < r_h \leq 2(h-1)$ , then  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity.*

Furthermore, we show the Herzog-Schönheim conjecture in free groups can be translated into a conjecture on automata.

**Theorem 2.** *If the following conjecture on automata is true:*

**Conjecture 1.** *Let  $\Sigma$  be a finite alphabet, and  $\Sigma^*$  be the free monoid generated by  $\Sigma$ . For every  $1 \leq i \leq s$ , let  $M_i$  be a finite, bi-deterministic and complete automaton with strongly-connected underlying graph. Let  $d_i$  be the number of states of  $M_i$  ( $d_i > 1$ ), and  $L_i \subsetneq \Sigma^*$  be the accepted language of  $M_i$ . If  $\Sigma^*$  is equal to the disjoint union of the  $s$  languages  $L_1, L_2, \dots, L_s$ , then there are  $1 \leq j, k \leq s$ ,  $j \neq k$ , such that  $d_j = d_k$ .*

*Then the Herzog-Schönheim conjecture is true.*

The paper is organized as follows. In Section 2, we give some preliminaries on automata and on irreducible non-negative matrices. In Section 3, we present a particular class of automata adapted to the study of the Herzog-Schönheim conjecture in free groups and describe some of their properties. In Section 4, we prove Theorem 1 and Theorem 2. The last section is an appendix with the proof of Lemma 4.5. We refer to [7] for more preliminaries and examples: Section 2, for free groups and covering spaces and Section 3.1, for graphs.

## 2 Automata, Non-negative irreducible matrices

### 2.1 Automata

We refer the reader to [30, p.96], [10, p.7], [23, 24], [11]. A *finite state automaton* is a quintuple  $(S, \Sigma, \mu, Y, s_0)$ , where  $S$  is a finite set, called the

state set,  $\Sigma$  is a finite set, called the *alphabet*,  $\mu : S \times \Sigma \rightarrow S$  is a function, called the *transition function*,  $Y$  is a (possibly empty) subset of  $S$  called the *accept (or end) states*, and  $s_0$  is called the *start state*. It is a directed graph with vertices the states and each transition  $s \xrightarrow{a} s'$  between states  $s$  and  $s'$  is an edge with label  $a \in \Sigma$ . The *label of a path*  $p$  of length  $n$  is the product  $a_1 a_2 \dots a_n$  of the labels of the edges of  $p$ . The finite state automaton  $M = (S, \Sigma, \mu, Y, s_0)$  is *deterministic* if there is only one initial state and each state is the source of exactly one arrow with any given label from  $\Sigma$ . In a deterministic automaton, a path is determined by its starting point and its label [30, p.105]. It is *co-deterministic* if there is only one final state and each state is the target of exactly one arrow with any given label from  $\Sigma$ . The automaton  $M = (S, \Sigma, \mu, Y, s_0)$  is *bi-deterministic* if it is both deterministic and co-deterministic. An automaton  $M$  is *complete* if for each state  $s \in S$  and for each  $a \in \Sigma$ , there is exactly one edge from  $s$  labelled  $a$ . We say that an automaton  $M$  is *strongly-connected* if there is a directed path from any state to any other state.

**Definition 2.1.** Let  $M = (S, \Sigma, \mu, Y, s_0)$  be a finite state automaton. Let  $\Sigma^*$  be the free monoid generated by  $\Sigma$ . Let  $\text{Map}(S, S)$  be the monoid consisting of all maps from  $S$  to  $S$ . The map  $\phi : \Sigma \rightarrow \text{Map}(S, S)$  given by  $\mu$  can be extended in a unique way to a monoid homomorphism  $\phi : \Sigma^* \rightarrow \text{Map}(S, S)$ . The range of this map is a monoid called *the transition monoid of  $M$* , which is generated by  $\{\phi(a) \mid a \in \Sigma\}$ . An element  $w \in \Sigma^*$  is *accepted* by  $M$  if the corresponding element of  $\text{Map}(S, S)$ ,  $\phi(w)$ , takes  $s_0$  to an element of the accept states set  $Y$ . The set  $L \subseteq \Sigma^*$  recognized by  $M$  is called *the language accepted by  $M$* , denoted by  $L(M)$ .

For any directed graph with  $d$  vertices or any finite state automaton  $M$ , with alphabet  $\Sigma$ , and  $d$  states, there exists a square matrix  $A$  of order  $d \times d$ , with  $a_{ij}$  equal to the number of directed edges from vertex  $i$  to vertex  $j$ ,  $1 \leq i, j \leq d$ . This matrix is non-negative (i.e  $a_{ij} \geq 0$ ) and it is called *the transition matrix* (as in [11]) or *the adjacency matrix* (as in [31, p.575]). For any  $k \geq 1$ ,  $(A^k)_{ij}$  is equal to the number of directed paths of length  $k$  from vertex  $i$  to vertex  $j$ . So, if  $M$  is a bi-deterministic automaton with alphabet  $\Sigma$ ,  $d$  states, start state  $i$ , accept state  $j$  and transition matrix  $A$ , then  $(A^k)_{ij}$  is the number of words of length  $k$  in the free monoid  $\Sigma^*$  accepted by  $M$ .

## 2.2 Irreducible non-negative matrices

We refer to [1, Ch.16], [20, Ch.8]. There is a vast literature on the topic. Let  $A$  be a transition matrix of order  $d \times d$  of a directed graph or an automaton with  $d$  states, as defined in Section 2.2. If for every  $1 \leq i, j \leq d$ , there exists  $m_{ij} \in \mathbb{Z}^+$  such that  $(A^{m_{ij}})_{ij} > 0$ , the matrix is *irreducible* and this is equivalent to the graph being strongly-connected. For  $A$  an irreducible non-negative matrix, *the period of  $A$*  is the gcd of all  $m \in \mathbb{Z}^+$  such that  $(A^m)_{ii} > 0$  (for any  $i$ ). If the period is 1,  $A$  is called *aperiodic*. In [20], an irreducible and aperiodic matrix  $A$  is called *primitive* and the period  $h$  is called the *index of imprimitivity*.

Let  $A$  be an irreducible non-negative matrix of order  $d \times d$  with period  $h \geq 1$  and spectral radius  $r$ . Then the Perron-Frobenius theorem states that  $r$  is a positive real number and it is a simple eigenvalue of  $A$ ,  $\lambda_{PF}$ , called the *Perron-Frobenius (PF) eigenvalue*. It satisfies  $\sum_i a_{ij} \leq \lambda_{PF} \leq \sum_j a_{ij}$ . The matrix  $A$  has a right eigenvector  $v_R$  with eigenvalue  $\lambda_{PF}$  whose components are all positive and likewise, a left eigenvector  $v_L$  with eigenvalue  $\lambda_{PF}$  whose components are all positive. Both right and left eigenspaces associated with  $\lambda_{PF}$  are one-dimensional. The behaviour of irreducible non-negative matrices depends strongly on whether the matrix is aperiodic or not.

**Theorem 2.2.** [20, Ch.8] *Let  $A$  be a  $d \times d$  irreducible non-negative matrix of period  $h \geq 1$ , with PF eigenvalue  $\lambda_{PF}$ . Let  $v_L$  and  $v_R$  be left and right eigenvectors of  $\lambda_{PF}$  whose components are all positive, with  $v_L v_R = 1$ .*

*If  $h = 1$ ,  $\lim_{k \rightarrow \infty} \frac{A^k}{\lambda_{PF}^k} = P$ , and if  $h > 1$ ,  $\lim_{k \rightarrow \infty} \frac{1}{k} \sum_{m=0}^{m=k-1} \frac{A^m}{\lambda_{PF}^m} = P$ ;  $P = v_R v_L$ .*

## 3 A particular class of automaton adapted to the study of the HS conjecture

### 3.1 The Schreier automaton of a coset of a subgroup

We now introduce the particular class of automata we are interested in, that is *the Schreier automaton for  $F_n$  relative to the subgroup  $H$*  [30, p.102], [32, p.107]. We refer to [7] for concrete examples.

**Definition 3.1.** Let  $F_n = \langle \Sigma \rangle$  and  $\Sigma^*$  the free monoid generated by  $\Sigma$ . Let  $H < F_n$  of index  $d$ . Let  $(\tilde{X}_H, p)$  be the covering of the  $n$ -leaves

bouquet with basepoint  $\tilde{x}_1$  and vertices  $\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_d$ . Let  $t_i \in \Sigma^*$  denote the label of a directed path of minimal length from  $\tilde{x}_1$  to  $\tilde{x}_i$ . Let  $\mathcal{T} = \{1, t_i \mid 1 \leq i \leq d\}$ . Let  $\tilde{X}_H$  be the Schreier coset diagram for  $F_n$  relative to the subgroup  $H$ , with  $\tilde{x}_1$  representing the subgroup  $H$  and the other vertices  $\tilde{x}_2, \dots, \tilde{x}_d$  representing the cosets  $Ht_i$  accordingly. We call  $\tilde{X}_H$  the *Schreier graph of  $H$* , with this correspondence between the vertices  $\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_d$  and the cosets  $Ht_i$  accordingly.

From its definition,  $\tilde{X}_H$  is a strongly-connected  $n$ -regular graph. So, its transition matrix  $A$  is non-negative and irreducible, with PF eigenvalue  $n$  (the sum of the elements at each row and at each column is equal to  $n$ ).

**Definition 3.2.** Let  $F_n = \langle \Sigma \rangle$  and  $\Sigma^*$  the free monoid generated by  $\Sigma$ . Let  $H < F_n$  of index  $d$ . Let  $\tilde{X}_H$  be the Schreier graph of  $H$ . Using the notation from Defn. 3.1, let  $\tilde{x}_1$  be the start state and  $\tilde{x}_f$  be the end state for some  $1 \leq f \leq d$ . We call the automaton obtained the *Schreier automaton of  $Ht_f$*  and denote it by  $\tilde{X}_{Ht_f}$ . The language accepted by  $\tilde{X}_{Ht_f}$  is the set of elements in  $\Sigma^*$  that belong to  $Ht_f$ . We call the elements in  $\Sigma^* \cap Ht_f$ , the *positive words in  $Ht_f$* . The identity may belong to this set.

**Example 3.3.** Let  $\Sigma = \{a, b\}$ ;  $F_2 = \langle a, b \rangle$ . Let  $K \leq F_2$ , of index 4.




Figure 3.3.1: The Schreier graph  $\tilde{X}_K$  of  $K = \langle a^4, b^4, ab^{-1}, a^2b^{-2}, a^3b^{-3} \rangle$ .

The transition matrix of  $\tilde{X}_K$  is  $\begin{pmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 2 & 0 & 0 & 0 \end{pmatrix}$  with period 4. If  $K$

and  $Ka$  are the start and end states,  $L$  is the set of positive words in  $Ka$ .

### 3.2 Properties of the Schreier automata in coset partitions

We recall here some results proved in [9].

**Theorem 3.4.** [9] Let  $F_n$  be the free group on  $n \geq 1$  generators. Let  $\{H_i \alpha_i\}_{i=1}^s$  be a coset partition of  $F_n$  with  $H_i < F_n$  of index  $d_i$ ,  $\alpha_i \in F_n$ ,  $1 \leq i \leq s$ , and  $1 < d_1 \leq \dots \leq d_s$ . Let  $\tilde{X}_i$  denote the Schreier graph of  $H_i$ , with transition matrix  $A_i$ , and period  $h_i \geq 1$ ,  $1 \leq i \leq s$ . Let  $1 \leq k, m \leq s$ .

- (i) Assume  $h_k > 1$ , where  $h_k = \max\{h_i \mid 1 \leq i \leq s\}$ . Then there exists  $j \neq k$  such that  $h_j = h_k$ .
- (ii) Let  $h_\ell > 1$ , such that  $h_\ell$  does not properly divide any other period  $h_i$ ,  $1 \leq i \leq s$ . Then there exists  $j \neq \ell$  such that  $h_j = h_\ell$ .
- (iii) For every  $h_i$ , there exists  $j \neq i$  such that either  $h_i = h_j$  or  $h_i \mid h_j$ .

If  $n = 1$  in Theorem 3.4,  $\{H_i\alpha_i\}_{i=1}^s$  is a coset partition of  $\mathbb{Z}$  and we recover the Davenport-Rado result (or Mirsky-Newman result) for the Erdős' conjecture and some of its consequences. Indeed, for every index  $d$ , the Schreier graph of  $d\mathbb{Z}$  has a transition matrix with period equal to  $d$ , so a repetition of the period is equivalent to a repetition of the index. For the unique subgroup  $H$  of  $\mathbb{Z}$  of index  $d$ , its Schreier graph  $\tilde{X}_H$  is a closed directed path of length  $d$  (with each edge labelled 1). So, its transition matrix  $A$  is the permutation matrix corresponding to the  $d$ -cycle  $(1, 2, \dots, d)$ , and it has period  $d$ . In particular, the period of  $A_s$  is  $d_s$ , and there exists  $j \neq s$  such that  $d_j = d_s$ . Also, if the period (index)  $d_k$  of  $A_k$  does not properly divide any other period (index), then there exists  $j \neq k$  such that  $d_j = d_k$ . For the free groups in general, we prove that in some cases, the repetition of the period implies the repetition of the index (see [9]).

## 4 Proof of the main results

### 4.1 Properties of the transition matrix of the Schreier graph

We study the properties of the transition matrix of a Schreier graph.

**Lemma 4.1.** *Let  $H < F_n$  of index  $d$ , with Schreier graph  $\tilde{X}_H$  and transition matrix  $A$  with period  $h \geq 1$ . Then the following properties hold:*

- (i) *The vectors  $v_L = \frac{1}{d}(1, 1, \dots, 1)$ ,  $v_R = (1, 1, \dots, 1)^T$  are left and right eigenvectors of  $n$  whose components are all positive, with  $v_L v_R = 1$ .*
- (ii) *The matrix  $P = v_R v_L$  is of order  $d \times d$  with all entries equal  $\frac{1}{d}$ .*
- (iii) *If  $h = 1$ , then  $\lim_{k \rightarrow \infty} \frac{A^k}{n^k} = P$  and if  $h > 1$ , then  $\lim_{k \rightarrow \infty} \frac{1}{k} \sum_{j=1}^{j=k} \frac{A^j}{n^j} = P$ .*

*Proof.* (i), (ii), (iii) As the sum of every row and every column in  $A$  is equal to  $n$ ,  $\lambda_{PF} = n$  with right eigenvalue  $v_R = (1, 1, \dots, 1)^T$  and left eigenvalue  $(1, 1, \dots, 1)$ . Since  $(1, 1, \dots, 1)v_R = d$ ,  $v_L = \frac{1}{d}(1, 1, \dots, 1)$  is a left eigenvector that satisfies  $v_L v_R = 1$ . Computing  $v_R v_L$  gives the matrix  $P$  of order  $d \times d$  with all entries equal  $\frac{1}{d}$ . (iii) results from Theorem 2.2.  $\square$



The behaviour of exponents of an aperiodic  $d \times d$  matrix of a Schreier graph  $\tilde{X}_H$  is well known: for every  $1 \leq i, j \leq d$ ,  $\lim_{k \rightarrow \infty} \frac{(A^k)_{ij}}{n^k} = \frac{1}{d}$ , from Lemma 4.1. It means that the proportion of positive words of every length  $k$  ( $k$  large enough) that belong to any coset of  $H$  tends to the fixed value  $\frac{1}{d}$ . We turn now to the study of  $\lim_{k \rightarrow \infty} \frac{(A^k)_{ij}}{n^k}$ , where  $A$  is the transition matrix of a Schreier graph  $\tilde{X}_H$  of period  $h > 1$ .

**Definition 4.2.** For  $1 \leq k, l \leq d$ , we define  $m_{ij}$ ,  $0 \leq m_{ij} \leq h$ , to be the minimal natural number such that  $(A^{m_{ij}})_{ij} \neq 0$ .

By definition, if  $i \neq j$ , then  $m_{ij}$  is the minimal length of a directed path from  $i$  to  $j$  in  $\tilde{X}_H$  and if  $i = j$ , then  $m_{ij} = 0$ . Whenever  $h > 1$ , only for the exponents  $m_{ij} + kh$ ,  $k \geq 0$ ,  $(A^{m_{ij}+kh})_{ij} \neq 0$ , that is only positive words of length  $m_{ij} + kh$  are accepted by the Schreier automaton, with  $i$  and  $j$  the start and end states respectively. Note that if  $H$  is a subgroup of  $\mathbb{Z} = \langle 1 \rangle$  of index  $d$ , its transition matrix  $A$  is a permutation matrix with period  $d$  and  $m_{ij} = r$ , where  $d\mathbb{Z} + r$  is the coset with  $i$  and  $j$  the start and end states respectively.

**Lemma 4.3.** Let  $H < F_n$  of index  $d$ , with Schreier graph  $\tilde{X}_H$  and transition matrix  $A$  with period  $h > 1$ . Then, the following properties hold:

- (i)  $\lim_{k \rightarrow \infty} \frac{(A^k)_{ij}}{n^k} = 0$ , whenever  $k \not\equiv m_{ij} \pmod{h}$ ,  $1 \leq i, j \leq d$ .
- (ii)  $\lim_{k \rightarrow \infty} \frac{(A^k)_{ij}}{n^k} = \frac{h}{d}$ , whenever  $k \equiv m_{ij} \pmod{h}$ ,  $1 \leq i, j \leq d$ .
- (iii) for every  $0 \leq m \leq h - 1$ , there is  $i$  such that  $m_{1i} \equiv m \pmod{h}$ .
- (iv)  $h$  divides  $d$ .

*Proof.* (i) By definition, whenever  $k \not\equiv m_{ij} \pmod{h}$ ,  $(A^k)_{ij} = 0$ .

(ii), (iii), (iv) We define a  $d \times h$  matrix  $B$  in the following way. Each row  $i$  is labelled by a right coset of  $H$  in the same order as they appear in the rows and columns of  $A$  and each column by  $m = 0, 1, 2, \dots, h - 1$ , and  $(B)_{ij} = \lim_{k \rightarrow \infty} \frac{(A^{j+kh})_{1i}}{n^{j+kh}}$ . Roughly,  $(B)_{ij}$  is the proportion of positive words of very large length that belong to the corresponding coset of  $H$ . From (i):

$$(B)_{ij} = \begin{cases} 0 & \text{if } m_{1i} \not\equiv j \pmod{h}, \\ \lim_{k \rightarrow \infty} \frac{(A^{m_{1i}+kh})_{1i}}{n^{m_{1i}+kh}} & \text{if } m_{1i} \equiv j \pmod{h}. \end{cases}$$

So, at each row of  $B$ , there is a single non-zero entry. As  $F_n$  is partitioned by the  $d$  cosets of  $H$ , all the non-zero elements in  $B$  are equal and for every  $k \geq 0$  and every  $1 \leq i \leq d$ ,  $\sum_{f=1}^{f=d} (A^k)_{if} = n^k$ , in particular  $\sum_{f=1}^{f=d} \frac{(A^k)_{1f}}{n^k} =$

1. So,  $\sum_{i=1}^{i=d} (B)_{ij} = \sum_{i=1}^{i=d} \lim_{k \rightarrow \infty} \frac{(A^{j+kh})_{1i}}{n^{j+kh}} = \lim_{k \rightarrow \infty} \sum_{i=1}^{i=d} \frac{(A^{j+kh})_{1i}}{n^{j+kh}} = 1$ , that is the sum of elements in each column of  $B$  is equal to 1. If  $h = d$ ,  $B$  is a square matrix and the right cosets can be arranged such that their  $m$  is in growing order and we have necessarily a diagonal matrix (otherwise there would be a column of zeroes). So,  $\lim_{k \rightarrow \infty} \frac{(A^{m_{1i}+kh})_{1i}}{n^{m_{1i}+kh}} = 1$  and  $(ii), (iii), (iv)$  hold. Now, assume  $d > h$ . At each column, there is at least one non-zero entry, so  $(iv)$  holds. Furthermore, the number of non-zero entries in each column needs to be the same, so  $h$  divides  $d$  and for any  $i$ ,  $\frac{d}{h} * (\lim_{k \rightarrow \infty} \frac{(A^{m_{1i}+kh})_{1i}}{n^{m_{1i}+kh}}) = 1$ .

That is,  $\lim_{k \rightarrow \infty} \frac{(A^{m_{1i}+kh})_{1i}}{n^{m_{1i}+kh}} = \frac{h}{d}$ . Furthermore,  $\lim_{k \rightarrow \infty} \frac{1}{h} \sum_{j=0}^{j=h-1} \frac{(A^{j+kh})_{1i}}{n^{j+kh}} = \frac{1}{d}$ .  $\square$

## 4.2 Conditions that ensure multiplicity in a coset partition

Let  $F_n$  be the free group on  $n \geq 1$  generators. Let  $\{H_i \alpha_i\}_{i=1}^{i=s}$  be a coset partition of  $F_n$  with  $H_i < F_n$  of index  $d_i$ ,  $\alpha_i \in F_n$ ,  $1 \leq i \leq s$ , and  $1 < d_1 \leq \dots \leq d_s$ . Let  $\tilde{X}_i$  denote the Schreier graph of  $H_i$ , with transition matrix  $A_i$ , and period  $h_i \geq 1$ ,  $1 \leq i \leq s$ . In the following lemmas, we prove, under these assumptions, there exist conditions that ensure multiplicity.

**Lemma 4.4.** *Assume there exists a unique  $h > 1$ . Let  $r$  denote the number of repetitions of  $h$ . Then,  $r \geq h$ . Furthermore, if  $r = h$  or if  $h < r \leq 2(h-1)$ , then  $\{H_i \alpha_i\}_{i=1}^{i=s}$  has multiplicity.*

*Proof.* For every  $1 \leq i \leq s$ ,  $(A_i^k)_{1f_i}$  denotes the number of positive words of length  $k$  that belong to the coset  $H_i \alpha_i$ . Let  $I = \{1 \leq i \leq s \mid h_i = h\}$  and for every  $i \in I$ ,  $m_i$  the minimal natural number ( $\text{mod } h$ ) such that  $(A_i^{m_i})_{1f_i} \neq 0$ . We define a  $r \times h$  matrix  $C$  in the following way. Each row  $i$  is labelled by a right coset  $H_i \alpha_i$ , where  $i \in I$  and each column by  $m = 0, 1, 2, \dots, h-1$ , and:

$$(C)_{ij} = \begin{cases} 0 & \text{if } m_i \not\equiv j(\text{mod } h), \\ \lim_{k \rightarrow \infty} \frac{(A_i^{m_i+hk})_{1f_i}}{n^{m_i+hk}} & \text{if } m_i \equiv j(\text{mod } h). \end{cases}$$

Roughly,  $(C)_{ij}$  is the proportion of positive words of very large length that belong to  $H_i \alpha_i$ , where  $i \in I$ . At each row of  $C$  there is a unique non-zero

entry. Since  $\{H_i\alpha_i\}_{i=1}^{i=s}$  is a coset partition of  $F_n$ , for every  $k$ ,  $\sum_{i=1}^{i=s} (A_i^k)_{1f_i} = n^k$ , that is  $\sum_{i=1}^{i=s} \frac{(A_i^k)_{1f_i}}{n^k} = 1$ . If  $A_i$  is aperiodic, then  $\lim_{k \rightarrow \infty} \frac{(A_i^k)_{1f_i}}{n^k} = \frac{1}{d_i}$  from Lemma 4.1. So,  $1 = \lim_{k \rightarrow \infty} \sum_{i=1}^{i=s} \frac{(A_i^k)_{1f_i}}{n^k} = \sum_{i=1}^{i=s} \lim_{k \rightarrow \infty} \frac{(A_i^k)_{1f_i}}{n^k} = \sum_{i \notin I} \frac{1}{d_i} + \sum_{i \in I} \lim_{k \rightarrow \infty} \frac{(A_i^k)_{1f_i}}{n^k}$ . That is,  $\sum_{i \in I} \lim_{k \rightarrow \infty} \frac{(A_i^k)_{1f_i}}{n^k} = 1 - \sum_{i \notin I} \frac{1}{d_i} = \sum_{i \in I} \frac{1}{d_i}$ , since  $\sum_{i=1}^{i=s} \frac{1}{d_i} = 1$ . So, the sum of elements in each column of  $C$  is equal to  $\sum_{i \in I} \frac{1}{d_i}$  and from Lemma 4.3, the non-zero entries in  $C$  have the form  $\frac{h}{d_i}$ . If  $r < h$ , then there is necessarily a column of zeroes, so  $r \geq h$ . If  $r = h$ , then  $C$  is a square matrix and the right cosets can be arranged such that their  $m$  is in growing order and we have necessarily a diagonal matrix (otherwise there would be a column of zeroes). So, for every  $i \in I$ ,  $\frac{h}{d_i} = \sum_{i \in I} \frac{1}{d_i}$ . That is, the coset partition  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity with all the  $d_i$  equal for  $i \in I$ . Now, assume  $r > h$ . At each column, there is at least one non-zero entry and there are necessarily columns with several non-zero entries. By some simple combinatorics,  $n_0$ , the number of columns with a single non-zero entry satisfies  $h - (r - h) \leq n_0 \leq h - 1$ , that is  $2h - r \leq n_0 \leq h - 1$ . If we assume  $r \leq 2(h - 1)$ , then  $n_0 \geq 2h - r - 2(h - 1) \geq 2$ , that is the number of columns with a single non-zero entry is at least 2, so there are at least two  $i \in I$ , such that  $\frac{h}{d_i} = \sum_{i \in I} \frac{1}{d_i}$ , that is the coset partition  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity. Note that for every  $0 \leq m \leq h - 1$ , there is  $i$  such that  $m_i \equiv m \pmod{h}$ .  $\square$

**Lemma 4.5.** *Assume there exists two  $h, h' > 1$  and  $h$  and  $h'$  are coprime. Let  $I = \{1 \leq i \leq s \mid h_i = h\}$ ,  $r = |I|$ ;  $I' = \{1 \leq i \leq s \mid h_i = h'\}$ ,  $r' = |I'|$ . If  $r = h$  or  $r \leq 2(h - 1)$  or  $r' = h'$  or  $r' \leq 2(h' - 1)$ , then  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity.*

The proof of Lemma 4.5 appears in the appendix. We prove there that coprime periods can be considered independently, that is each period can be assigned its own matrix  $C$  as defined in the proof of Lemma 4.4. The situation is different if  $h$  and  $h'$  are not coprime. Indeed, consider the following coset partition of  $F_2$ :  $F_2 = H \cup Ka \cup Ka^3$ , where  $K$  is the subgroup described in Example 3.3 and  $H = \langle a^2, b^2, ab \rangle < F_2$  of index 2. The period of  $\tilde{X}_H$  is  $h' = 2$  and the period of  $\tilde{X}_K$  is  $h = 4$  and the corresponding

matrix  $D$  as defined in the proof of Lemma 4.5 is  $D = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ ,

with the first row labelled  $H$ , the second row  $Ka$ , the third row  $Ka^3$  and at each column  $0 \leq m \leq 3$ . So, if  $h'$  divides  $h$ , each period cannot have its own matrix  $C$ . Yet, using the same kind of arguments as before, it is not difficult to prove that  $r \geq h - \frac{h}{h'}r'$  and that if  $r \leq 2(h - \frac{h}{h'}r' - 1)$  then the coset partition has multiplicity. We now turn to the proof of Theorem 1.

*Proof of Theorem 1.* We assume  $H = \{h > 1 \mid \exists 1 \leq j \leq s, h_j = h\} \neq \emptyset$  and different elements in  $H$  are pairwise coprime. Let  $r_h = |\{1 \leq j \leq s, h_j = h\}|$ , the number of repetitions of  $h$ . From the proof of Lemma 4.5, each period has its own matrix  $C$  and we apply the results of Lemma 4.3. That is, if for some  $h \in H$ ,  $r_h = h$  or  $h < r_h \leq 2(h - 1)$ , then  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity.  $\square$

### 4.3 Translation of the HS conjecture in terms of automata

Let  $F_n = \langle \Sigma \rangle$ , and  $\Sigma^*$  the free monoid generated by  $\Sigma$ . Let  $\{H_i\alpha_i\}_{i=1}^{i=s}$  be a coset partition of  $F_n$  with  $H_i < F_n$  of index  $d_i > 1$ ,  $\alpha_i \in F_n$ ,  $1 \leq i \leq s$ . Let  $\tilde{X}_i$  be the Schreier automaton of  $H_i\alpha_i$ , with language  $L_i = \Sigma^* \cap H_i\alpha_i$ .

*Proof of Theorem 2.* Assume Conjecture 1 is true. For every  $1 \leq i \leq s$ , the Schreier automaton  $\tilde{X}_i$  is a finite, bi-deterministic and complete automaton with strongly-connected underlying graph and alphabet  $\Sigma$ . Since  $F_n$  is the disjoint union of the sets  $\{H_i\alpha_i\}_{i=1}^{i=s}$ , each word in  $\Sigma^*$  belongs to one and exactly one such language, so  $\Sigma^*$  is the disjoint union of the  $s$  languages  $L_1, L_2, \dots, L_s$ . Since Conjecture 1 is true, there is a repetition of the number of states and this implies the coset partition  $\{H_i\alpha_i\}_{i=1}^{i=s}$  has multiplicity, that is the HS conjecture in free groups of finite rank is true. From [7, Thm.6], this implies the HS conjecture is true for all the finitely generated groups, in particular for all the finite groups. So, the HS conjecture is true for all the groups.  $\square$

Note that these two conjectures would have been equivalent if the conditions of Conjecture 1 implied the existence of a coset partition of the free group of rank  $|\Sigma|$ , which does not seem to be true. Nevertheless, any finite, bi-deterministic, complete and strongly-connected automaton  $M$  with  $d$  states can be considered as the Schreier automaton of a subgroup  $H$  of index  $d$  in  $F_{|\Sigma|}$ .

## References

- [1] R. Bellman, *Matrix Analysis*, S.I.A.M. Press, 1997.
- [2] M.A. Berger, A. Felzenbaum, A.S. Fraenkel, *Improvements to two results concerning systems of residue sets*, Ars. Combin. **20** (1985), 69-82.
- [3] M.A. Berger, A. Felzenbaum, A.S. Fraenkel, *The Herzog-Schönheim conjecture for finite nilpotent groups*, Canad. Math. Bull. **29**(1986),329-333.
- [4] M.A. Berger, A. Felzenbaum, A.S. Fraenkel, *Lattice parallelotopes and disjoint covering systems*, Discrete Math. **65** (1987), 23-44.
- [5] M.A. Berger, A. Felzenbaum, A.S. Fraenkel, *Remark on the multiplicity of a partition of a group into cosets*, Fund. Math. **128** (1987), 139-144.
- [6] M.A. Brodie, R.F. Chamberlain, L.C Kappe, *Finite coverings by normal subgroups*, Proc. Amer. Math. Soc. **104** (1988), 669-674.
- [7] F. Chouraqui, *The Herzog-Schönheim conjecture for finitely generated groups*, International J. of Alg. and Comp. (2019), vol.29, n.6, p. 1083-1112.
- [8] F. Chouraqui, *The space of coset partitions of  $F_n$  and Herzog-Schönheim conjecture*, ArXiv 1804.11103.
- [9] F. Chouraqui, *About an extension of the Davenport-Rado result to the Herzog-Schönheim conjecture for free groups*, ArXiv 1901.09898.
- [10] D.B.A. Epstein, J.W. Cannon, D.F. Holt, S.V.F. Levy, M.S. Paterson, W.P. Thurston, *Word Processing in Groups*, Jones and Bartlett Publishers (1992).
- [11] D.B.A. Epstein, A.R. Iano-Fletcher, U.Zwick, *Growth functions and automatic groups*, Experimental Math. **5** (1996), n.4.
- [12] P. Erdős, *On integers of the form  $2^k + p$  and some related problems*, Summa Brasil. Math. **2** (1950), 113-123.
- [13] P. Erdős, *Problems and results in Number theory*, Recent Progress in Analytic Number Theory, vol. 1, Academic Press, London-New York, 1981, 1-13.

- [14] Y. Ginosar, *Tile the group*, Elem. Math. **72**, Swiss Math. Society.
- [15] Y. Ginosar, O. Schnabel, *Prime factorization conditions providing multiplicities in coset partitions of groups*, J. Comb. Number Theory, **3** (2011), n.2, 75-86.
- [16] A. Hatcher, *Algebraic Topology*, Cambridge University Press (2002).
- [17] M. Herzog, J. Schönheim, *Research problem no. 9*, Canad. Math. Bull., **17** (1974), 150.
- [18] I. Korec, Š. Znám, *On disjoint covering of groups by their cosets*, Math. Slovaca, **27** (1977), 3-7.
- [19] L. Margolis, O. Schnabel, *The Herzog-Schonheim conjecture for small groups and harmonic subgroups*, ArXiv 1803.03569.
- [20] C.D. Meier, *Matrix analysis and applied linear algebra*, SIAM: Society for Industrial and Applied Mathematics (2010).
- [21] M. Newman, *Roots of unity and covering sets*, Math. Ann. **191** (1971), 279-282.
- [22] B. Novák, Š. Znám, *Disjoint covering systems*, Amer. Math. Monthly, **81** (1974), 42-45.
- [23] J.E. Pin, *On reversible automata*, Lecture Notes in Computer Science **583**, Springer 1992, p. 401-416.
- [24] J.E. Pin, *Mathematical foundations of automata theory*, <https://www.irif.fr/~jep/PDF/MPRI/MPRI.pdf>
- [25] Š. Porubský, *Natural exactly covering systems of congruences*, Czechoslovak Math. J. **24** (1974), 598-606.
- [26] Š. Porubský, *Covering systems and generating functions*, Acta Arith. **26** (1975), n.3, 223-231.
- [27] Š. Porubský, *Results and problems on covering systems of residue classes*, Mitt. Math. Sem. Giessen, **150** (1981).
- [28] Š. Porubský, J. Schönheim, *Covering systems of Paul Erdős. Past, present and future. Paul Erdős and his mathematics*, János Bolyai Math. Soc., **11** (2002), 581-627.

- [29] J.J. Rotman, *An Introduction to Algebraic Topology*, Graduate Texts in Mathematics **119**, Springer-Verlag, Berlin, Heidelberg, New York (1988).
- [30] C.C. Sims, *Computation with finitely presented groups*, Encyclopedia of Mathematics and its Applications **48** , Cambridge University Press (1994).
- [31] R.P. Stanley, *Enumerative Combinatorics*, Wadsworth and Brooks/Cole, Monterey, CA, 1986.
- [32] J. Stillwell, *Classical Topology and Combinatorial Group Theory*, Graduate Texts in Mathematics **72**, Springer-Verlag, Berlin, Heidelberg, New York (1980).
- [33] Z.W. Sun, *Finite covers of groups by cosets or subgroups*, Internat. J. Math. **17** (2006), n.9, 1047-1064.
- [34] Z.W. Sun, *An improvement of the Znám-Newman result*, Chinese Quart. J. Math. **6** (1991), n.3, 90-96.
- [35] Z.W. Sun, *Covering the integers by arithmetic sequences II* Trans. Amer. Math. Soc. **348** (1996), 4279-4320.
- [36] M.J. Tomkinson, *Groups covered by abelian subgroups*, London Math. Soc. Lecture Note Ser. **121**, Cambridge Univ. Press (1986).
- [37] M.J. Tomkinson, *Groups covered by finitely many cosets or subgroups*, Comm. Algebra **15**(1987), 845-859.

## Appendix: Proof of Lemma 4.5

**Lemma.** Assume there exists two  $h, h' > 1$  and  $h$  and  $h'$  are coprime. Let  $I = \{1 \leq i \leq s \mid h_i = h\}$ ,  $r = |I|$ ;  $I' = \{1 \leq i \leq s \mid h_i = h'\}$ ,  $r' = |I'|$ . If  $r = h$  or  $r \leq 2(h-1)$  or  $r' = h'$  or  $r' \leq 2(h'-1)$ , then  $\{H_i \alpha_i\}_{i=1}^{i=s}$  has multiplicity.

*Proof of Lemma 4.5.* Assume with no loss of generality that  $h' < h$ . From the same argument as in the proof of Lemma 4.4,  $\sum_{i \in I \cup I'} \lim_{k \rightarrow \infty} \frac{(A_i^k)_{1f_i}}{n^k} = \sum_{i \in I \cup I'} \frac{1}{d_i}$ . We show that each period can be considered independently, that is each period has its own matrix  $C$  as defined in the proof of Lemma 4.4. We define a  $(r' + r) \times L$  matrix  $D$ , where  $L = 2hh'$ , in the following way. The first  $r'$  rows are labelled by right cosets  $H_i \alpha_i$ , where  $i \in I'$ , the last  $r$  rows are labelled by right cosets  $H_i \alpha_i$ , where  $i \in I$  and each column by  $m = 0, 1, 2, \dots, h' - 1, \dots, h - 1, h, \dots, L - 1$ , and:

$$(D)_{ij} = \begin{cases} 0 & \text{if } i \in I', m_i \not\equiv j \pmod{h'}, \\ \lim_{k \rightarrow \infty} \frac{(A_i^{m_i + h'k})_{1f_i}}{n^{m_i + h'k}} & \text{if } i \in I', m_i \equiv j \pmod{h'} \\ 0 & \text{if } i \in I, m_i \not\equiv j \pmod{h}, \\ \lim_{k \rightarrow \infty} \frac{(A_i^{m_i + hk})_{1f_i}}{n^{m_i + hk}} & \text{if } i \in I, m_i \equiv j \pmod{h}. \end{cases}$$

So, the sum of elements in each column of  $D$  is equal to  $\sum_{i \in I \cup I'} \frac{1}{d_i}$  and from

Lemma 4.3, the non-zero entries in  $D$  have the form  $\frac{h}{d_i}$  for  $i \in I$  and  $\frac{h'}{d_i}$ , for  $i \in I'$ . Let  $0 \leq m \leq h' - 1$ , the minimal number such that the sum of the entries of the  $m$ -th column is  $\sum_{i \in J_0} \frac{h}{d_i} + \sum_{i \in J'_0} \frac{h'}{d_i}$ , where  $\emptyset \neq J_0 \subset I$

and  $\emptyset \neq J'_0 \subset I'$ . So, for every  $0 \leq k \leq h' - 1$ , the sum of the entries of the  $(m + kh)$ -th column is  $\sum_{i \in J_0} \frac{h}{d_i} + \sum_{i \in J'_k} \frac{h'}{d_i}$ , and this implies necessarily

$$\sum_{i \in J'_0} \frac{h'}{d_i} = \sum_{i \in J'_1} \frac{h'}{d_i} = \dots = \sum_{i \in J'_{h'-1}} \frac{h'}{d_i}. \text{ We show that } \{\frac{h'}{d_i} \mid i \in J'_0\}, \{\frac{h'}{d_i} \mid i \in$$

$J'_1\}, \dots, \{\frac{h'}{d_i} \mid i \in J'_{h'-1}\}$  appear in the first  $h'$  columns of  $D$  (not necessarily in this order). Let  $0 \leq k, l \leq h' - 1$ ,  $k \neq l$ . Assume by contradiction that  $m + kh \equiv m + lh \pmod{h'}$ . So,  $h'$  divides  $h(k - l)$ . As  $h$  and  $h'$  are coprime,  $h'$  divides  $k - l$ , a contradiction. So, for every  $0 \leq k, l \leq h' - 1$ ,  $k \neq l$ ,  $m + kh \not\equiv m + lh \pmod{h'}$ . As there are exactly  $h'$  values, these correspond to  $0, 1, \dots, h' - 1 \pmod{h'}$ , and  $\{\frac{h'}{d_i} \mid i \in J'_0\}, \{\frac{h'}{d_i} \mid i \in J'_1\}, \dots,$



$\{\frac{h'}{d_i} \mid i \in J'_{h'-1}\}$  appear in the first  $h'$  columns of  $D$  with  $\sum_{i \in J'_0} \frac{h'}{d_i} = \dots = \sum_{i \in J'_{h'-1}} \frac{h'}{d_i}$ . Furthermore,  $\sum_{i \in J'_0} \frac{h'}{d_i} = \dots = \sum_{i \in J'_{h'-1}} \frac{h'}{d_i} = \sum_{i \in I'} \frac{1}{d_i}$ . Indeed, on one hand, the sum of elements in the first  $r'$  rows and  $h'$  columns is equal to  $h' \sum_{i \in J'_0} \frac{h'}{d_i}$  and on the second hand, it is equal  $\sum_{i \in I'} \frac{h'}{d_i}$ . Using the same argument, for every  $0 \leq k \leq h-1$ , the sum of the entries of the  $(m + kh')$ -th column is  $\sum_{i \in J_k} \frac{h}{d_i} + \sum_{i \in J'_0} \frac{h'}{d_i}$ , and this implies necessarily  $\sum_{i \in J_0} \frac{h}{d_i} = \sum_{i \in J_1} \frac{h}{d_i} = \dots = \sum_{i \in J_{h-1}} \frac{h}{d_i}$ . We show that  $\{\frac{h}{d_i} \mid i \in J_0\}, \{\frac{h}{d_i} \mid i \in J_1\}, \dots, \{\frac{h}{d_i} \mid i \in J_{h-1}\}$  appear in the first  $h$  columns of  $D$  (not necessarily in this order). Let  $0 \leq k, l \leq h-1, k \neq l$ . Assume by contradiction that  $m + kh' \equiv m + lh' \pmod{h}$ . So,  $h$  divides  $h'(k-l)$ . As  $h$  and  $h'$  are coprime,  $h$  divides  $k-l$ , a contradiction. So, for every  $0 \leq k, l \leq h-1, k \neq l$ ,  $m + kh' \not\equiv m + lh' \pmod{h}$ . As there are exactly  $h$  values, these correspond to  $0, 1, \dots, h-1 \pmod{h}$ , and  $\{\frac{h}{d_i} \mid i \in J_0\}, \{\frac{h}{d_i} \mid i \in J_1\}, \dots, \{\frac{h}{d_i} \mid i \in J_{h-1}\}$  appear in the first  $h$  columns of  $D$ , with  $\sum_{i \in J_0} \frac{h}{d_i} = \sum_{i \in J_1} \frac{h}{d_i} = \dots = \sum_{i \in J_{h-1}} \frac{h}{d_i}$ . Furthermore,  $\sum_{i \in J_0} \frac{h}{d_i} = \dots = \sum_{i \in J_{h-1}} \frac{h}{d_i} = \sum_{i \in I} \frac{1}{d_i}$ . So, each period has its own matrix  $C$  and we apply the results of Lemma 4.3.  $\square$