

COMPLEMENTS OF SCHUBERT POLYNOMIALS

NEIL J.Y. FAN, PETER L. GUO, NICOLAS Y. LIU

ABSTRACT. Let $\mathfrak{S}_w(x)$ be the Schubert polynomial for a permutation w of $\{1, 2, \dots, n\}$. For any given composition μ , we say that $x^\mu \mathfrak{S}_w(x^{-1})$ is the complement of $\mathfrak{S}_w(x)$ with respect to μ . When each part of μ is equal to $n - 1$, Huh, Matherne, Mészáros and St. Dizier proved that the normalization of $x^\mu \mathfrak{S}_w(x^{-1})$ is a Lorentzian polynomial. They further conjectured that the normalization of $\mathfrak{S}_w(x)$ is Lorentzian. It can be shown that if there exists a composition μ such that $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then the normalization of $\mathfrak{S}_w(x)$ will be Lorentzian. This motivates us to investigate the problem of when $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial. We show that if $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then μ must be a partition. We also consider the case when μ is the staircase partition $\delta_n = (n - 1, \dots, 1, 0)$, and obtain that $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial if and only if w avoids the patterns 132 and 312. A conjectured characterization of when $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial is proposed.

Keywords: Schubert polynomial, Lorentzian polynomial, RC-graph

AMS Classifications: 05E05, 14N15

1 Introduction

The motivation of this paper is the Lorentzian property of Schubert polynomials studied by Huh, Matherne, Mészáros and St. Dizier [7]. For the beautiful theory developed around Lorentzian polynomials as well as its powerful applications, see the work of Brändén and Huh [3].

Given a polynomial $f = f(x_1, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$, the normalization of f , denoted $N(f)$, is the polynomial obtained from f after applying a linear operator N which is defined by

$$N(x^\alpha) = \frac{x^\alpha}{\alpha_1! \cdots \alpha_n!}, \quad \text{for each monomial } x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}.$$

Let S_n be the symmetric group of permutations of $[n] = \{1, 2, \dots, n\}$. For $w \in S_n$, let $\mathfrak{S}_w(x) = \mathfrak{S}_w(x_1, \dots, x_n)$ denote the Schubert polynomial indexed by w . Schubert polynomials were introduced by Lascoux and Schützenberger [8] as the polynomial representatives of Schubert classes in the cohomology ring of the flag variety.

Using the Lorentzian property of volume polynomials of irreducible complex projective varieties, Huh, Matherne, Mészáros and St. Dizier [7, Theorem 6] showed that

$$N(x_1^{n-1} \cdots x_n^{n-1} \mathfrak{S}_w(x^{-1})) := N(x_1^{n-1} \cdots x_n^{n-1} \mathfrak{S}_w(x_1^{-1}, \dots, x_n^{-1}))$$

is Lorentzian, see [3, 7] for the precise definition of Lorentzian polynomials. As a consequence, the normalization $N(s_\lambda(x))$ of a Schur polynomial is Lorentzian [7, Theorem

3]. This can be used to verify Okounkov's log-concavity conjecture [10, Conjecture 1] for the Littlewood–Richardson coefficients in the case of Kostka numbers [7, Theorem 2].

Huh, Matherne, Mészáros and St. Dizier [7, Conjecture 15] further conjectured that the normalization $N(\mathfrak{S}_w(x))$ of each Schubert polynomial is Lorentzian. Motivated by this conjecture, we investigate the following problem:

When $x^\mu \mathfrak{S}_w(x^{-1})$ is still a Schubert polynomial?

We call $x^\mu \mathfrak{S}_w(x^{-1})$ the complement of $\mathfrak{S}_w(x)$ with respect to a composition μ . As will be seen in the following remark, if there exists a μ such that $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then $N(\mathfrak{S}_w(x))$ will satisfy the Lorentzian property, and thus in this case we can verify the conjecture of Huh, Matherne, Mészáros and St. Dizier.

Remark 1. We explain that if there exists a μ such that $x^\mu \mathfrak{S}_w(x^{-1})$ for $w \in S_n$ is a Schubert polynomial, then $N(\mathfrak{S}_w(x))$ is Lorentzian. First, we claim that $N(x^\mu \mathfrak{S}_w(x^{-1}))$ is Lorentzian. This can be seen as follows. It was observed in [7, Lemma 7] that for any $\nu \in \mathbb{Z}_{\geq 0}^n$ and a polynomial $f = f(x_1, \dots, x_n)$, $N(f)$ is Lorentzian if and only if $N(x^\nu f)$ is Lorentzian. Without loss of generality, we may assume that $\max\{\mu_1, \dots, \mu_n\} \leq n-1$ since otherwise we can embed w into S_m with $m > n$. Let $\nu_i = n-1-\mu_i \in \mathbb{Z}_{\geq 0}$, so that

$$x_1^{n-1} \cdots x_n^{n-1} \mathfrak{S}_w(x^{-1}) = x^\nu \cdot x^\mu \mathfrak{S}_w(x^{-1}),$$

and hence, the Lorentzian property of $N(x_1^{n-1} \cdots x_n^{n-1} \mathfrak{S}_w(x^{-1}))$ [7, Theorem 6] implies that of $N(x^\mu \mathfrak{S}_w(x^{-1}))$. This verifies the claim. Suppose now that $x^\mu \mathfrak{S}_w(x^{-1})$ equals a Schubert polynomial, say $\mathfrak{S}_{w'}(x)$. Then we have $\mathfrak{S}_w(x) = x^\mu \mathfrak{S}_{w'}(x^{-1})$. By the above claim, $N(x^\mu \mathfrak{S}_{w'}(x^{-1}))$ is Lorentzian, and so it follows that $N(\mathfrak{S}_w(x))$ is Lorentzian.

Our first result shows that if $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then μ must be a partition, namely, the parts of μ are weakly decreasing.

Theorem 1.1. *Let $w \in S_n$. If there exists a composition μ such that $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then μ is a partition.*

Let $\delta_n = (n-1, \dots, 1, 0)$ denote the staircase partition. For $w \in S_n$, it is well known that if a monomial x^α appears in $\mathfrak{S}_w(x)$, then $x^\alpha | x^{\delta_n}$, that is, $\alpha_i \leq n-i$ for $1 \leq i \leq n$. In fact, the set $\{\mathfrak{S}_w(x) : w \in S_n\}$ of Schubert polynomials forms a \mathbb{Z} -basis of

$$V_n = \text{span}_{\mathbb{Z}}\{x^\alpha : 0 \leq \alpha_i \leq n-i\},$$

see for example [1, 9]. Notice that $x^\alpha \mapsto x^{\delta_n - \alpha}$ induces an involutive automorphism on V_n . Hence the set $\{x^{\delta_n} \mathfrak{S}_w(x^{-1}) : w \in S_n\}$ is a basis of V_n . So it is natural to investigate the typical case for $\mu = \delta_n$.

Theorem 1.2. *Let $w \in S_n$. Then $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial if and only if w avoids the patterns 132 and 312.*

The sufficiency of Theorem 1.2 can be easily deduced since the Schubert polynomial for a 132-avoiding permutation is a single monomial. However, the proof of the necessity of Theorem 1.2 turns out to be quite technical.

Remark 2. The complements $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ are related to the padded Schubert polynomials defined by Gaetz and Gao [5, 6]. The padded Schubert polynomial $\tilde{\mathfrak{S}}_w(x; y)$ is obtained from $\mathfrak{S}_w(x)$ by replacing each monomial x^α in $\mathfrak{S}_w(x)$ by $x^\alpha y^{\delta_n - \alpha}$. Setting $x_i = 1$ in $\tilde{\mathfrak{S}}_w(x; y)$, we see that $\tilde{\mathfrak{S}}_w(x; y)|_{x_i=1} = y^{\delta_n} \mathfrak{S}_w(y^{-1})$. Thus Theorem 1.2 is equivalent to saying that $\tilde{\mathfrak{S}}_w(x; y)|_{x_i=1}$ is a Schubert polynomial if and only if w avoids the patterns 132 and 312.

Computational verification suggests the following conjecture.

Conjecture 1.3. *Let $w \in S_n$. Then there exists a partition μ such that $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial if and only if w avoids the patterns 1432, 13254, 14253, 24153, 31524, 361452.*

This paper is organized as follows. In Section 2, we give a proof of Theorem 1.1. The arguments rely on the RC-graph (also called pipedream) model of Schubert polynomials. In Section 3, we present a proof of Theorem 1.2. Remarks about Conjecture 1.3 are given in Section 4.

Acknowledgments. We would like to thank the anonymous referee for valuable comments and suggestions. This work was supported by the National Science Foundation of China (11971250, 12071320, 12371329) and Sichuan Science and Technology Program (2023ZYD0012).

2 Proof of Theorem 1.1

To give a proof of Theorem 1.1, we require the RC-graph model of Schubert polynomials investigated by Bergeron and Billey [1].

Schubert polynomials are defined based on divided difference operators. For a polynomial $f(x) \in \mathbb{Z}[x_1, \dots, x_n]$, the divided difference operator ∂_i acts on $f(x)$ by letting

$$\partial_i f(x) = \frac{f(x) - s_i f(x)}{x_i - x_{i+1}},$$

where $s_i f(x)$ is obtained from $f(x)$ by exchanging x_i and x_{i+1} . It is easily checked that $\partial_i f(x)$ is symmetric in x_i and x_{i+1} .

For $w_0 = n \cdots 21$, set

$$\mathfrak{S}_{w_0}(x) = x_1^{n-1} x_2^{n-2} \cdots x_{n-1}.$$

For $w = w_1 w_2 \cdots w_n \neq w_0$, choose a position i such that $w_i < w_{i+1}$ and define

$$\mathfrak{S}_w(x) = \partial_i \mathfrak{S}_{ws_i}(x),$$

where ws_i is obtained from w by interchanging w_i and w_{i+1} . The above definition is independent of the choice of i since the operators ∂_i satisfy the Coxeter relations: $\partial_i \partial_j = \partial_j \partial_i$ for $|i - j| > 1$, and $\partial_i \partial_{i+1} \partial_i = \partial_{i+1} \partial_i \partial_{i+1}$.

Billey, Jockusch and Stanley [2] showed that $\mathfrak{S}_w(x)$ can be combinatorially generated in terms of reduced-word compatible sequences. Each permutation w can be expressed

as a product $s_{a_1}s_{a_2}\cdots s_{a_p}$ of adjacent transpositions. If p is minimal, then we write $\ell(w) = p$, which is the length of w , and in this case $a = (a_1, \dots, a_p)$ is called a reduced word of w . A sequence $\alpha = (\alpha_1, \dots, \alpha_p)$ of positive integers is said to be a compatible sequence of a if (i) $\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_p$, (ii) $\alpha_i < \alpha_{i+1}$ if $a_i < a_{i+1}$, and (iii) $1 \leq \alpha_i \leq a_i$. Let $\text{Red}(w)$ denote the set of reduced words of w . For a reduced word $a \in \text{Red}(w)$, $C(a)$ denotes the set of its compatible sequences. It was proved in [2] that for $w \in S_n$,

$$\mathfrak{S}_w(x) = \sum_{a \in \text{Red}(w)} \sum_{\alpha \in C(a)} x_{\alpha_1} x_{\alpha_2} \cdots x_{\alpha_p}. \quad (2.1)$$

The RC-graph corresponding to a compatible pair (a, α) can be defined as follows. Let Δ_n denote the array of left-justified boxes with $n+1-i$ boxes in row i for $1 \leq i \leq n$. The RC-graph corresponding to (a, α) is the subset of Δ_n consisting of boxes in row α_i and column $a_i - \alpha_i + 1$, where $1 \leq i \leq p$. For example, let $a = (4, 2, 3, 2, 4)$ be a reduced word of $w = 15342$, and $\alpha = (1, 1, 2, 2, 4)$ be a compatible sequence of a . Then the corresponding RC-graph is illustrated in Figure 2.1, where we use a cross to signify a box belonging to the RC-graph.




Figure 2.1: An RC-graph of $w = 15342$.

For an RC-graph D of w , let

$$x^D = \prod_{i \in [n]} x_i^{|\{\text{boxes in the } i\text{-th row of } D\}|}.$$

In this notation, formula (2.1) can be rewritten as

$$\mathfrak{S}_w(x) = \sum_D x^D,$$

where the sum runs over all the RC-graphs of w .

For the purpose of this paper, we pay attention to two specific RC-graphs: the bottom RC-graph and the top RC-graph. Let $d(w) = (d_1, d_2, \dots, d_n)$ denote the inversion code of w , that is, for $1 \leq i \leq n$,

$$d_i = |\{j : j > i, w_j < w_i\}|.$$

Clearly, $0 \leq d_i \leq n - i$. The bottom RC-graph of w is the RC-graph consisting of the first d_i boxes in row i . For example, Figure 2.2(a) is the bottom RC-graph of $w = 25143$.

Bergeron and Billey [1] showed that any RC-graph of w can be obtained from the bottom RC-graph of w by applying a sequence of ladder moves. Let D be an RC-graph




Figure 2.2: The bottom and top RC-graphs of $w = 25143$.

of w , and let (i, j) denote a box of D in row i and j . The ladder move $L_{i,j}$ is a local change of the crosses as illustrated in Figure 2.3. Formally, the resulting diagram after applying the ladder move $L_{i,j}$ is

$$L_{i,j}(D) = D \setminus \{(i, j)\} \cup \{(h, j + 1)\}.$$

It can be verified that $L_{i,j}(D)$ is still an RC-graph of w . Since the ladder move operation




Figure 2.3: A ladder move.

moves a cross upwards, the bottom RC-graph of w corresponds to the leading monomial of $\mathfrak{S}_w(x)$ in the reverse lexicographic order.

Proposition 2.1 (Bergeron–Billey [1]). *Let $w \in S_n$, and $d(w) = (d_1, \dots, d_n)$ be the inversion code of w . Then the monomial $x^{d(w)} = x_1^{d_1} \cdots x_n^{d_n}$ is the leading term of $\mathfrak{S}_w(x)$ in the reverse lexicographic order.*

It was shown in [1] that the transpose of an RC-graph of w is an RC-graph of the inverse w^{-1} of w . The top RC-graph of w is defined as the transpose of the bottom RC-graph of w^{-1} . For example, the inverse of $w = 25143$ is $w^{-1} = 31542$, and so the top RC-graph of w is as depicted in Figure 2.2(b).

Dual to the ladder moves, Bergeron and Billey [1] defined chute moves on RC-graphs, and showed that any RC-graph of w can be obtained from the top RC-graph of w by applying a sequence of chute moves. The chute move operation moves a cross downwards, and thus the top RC-graph of w corresponds to the smallest monomial of $\mathfrak{S}_w(x)$ in the reverse lexicographic order.

For a vector $v = (v_1, v_2, \dots, v_n)$ with $0 \leq v_i \leq n - i$, let

$$v^t = (v'_1, v'_2, \dots, v'_n) \tag{2.2}$$

denote the transpose of v , namely, for $1 \leq i \leq n$,

$$v'_i = |\{1 \leq j \leq n : v_j \geq i\}|.$$

It is easy to see that v^t is a partition for any v .

Proposition 2.2 (Bergeron–Billiey [1]). *For $w \in S_n$, let $d^t(w^{-1}) = (d'_1, \dots, d'_n)$ denote the transpose of the inversion code of w^{-1} . Then the monomial $x^{d^t(w^{-1})} = x_1^{d'_1} \cdots x_n^{d'_n}$ is the smallest term of $\mathfrak{S}_w(x)$ in the reverse lexicographic order.*

Using Propositions 2.1 and 2.2, we can now provide a proof of Theorem 1.1.

Proof of Theorem 1.1. Suppose that

$$x^\mu \mathfrak{S}_w(x^{-1}) = \mathfrak{S}_{w'}(x). \quad (2.3)$$

Equivalently,

$$x^\mu \mathfrak{S}_{w'}(x^{-1}) = \mathfrak{S}_w(x). \quad (2.4)$$

By Propositions 2.1 and 2.2, it follows that

$$d(w') = \mu - d^t(w^{-1}) \quad \text{and} \quad d(w) = \mu - d^t(w'^{-1}),$$

and so we have

$$d(w') + d^t(w^{-1}) = d(w) + d^t(w'^{-1}) = \mu. \quad (2.5)$$

Suppose to the contrary that μ is not a partition. Locate an index i such that $\mu_i < \mu_{i+1}$. Write $d(w) = (d_1, d_2, \dots, d_n)$ and $d(w') = (d'_1, d'_2, \dots, d'_n)$. Keep in mind that both $d^t(w^{-1})$ and $d^t(w'^{-1})$ are partitions. In view of (2.5) together with the assumption $\mu_i < \mu_{i+1}$, we must have $d_i < d_{i+1}$ and $d'_i < d'_{i+1}$. By the definition of inversion code, we obtain that $w_i < w_{i+1}$ and $w'_i < w'_{i+1}$. Therefore,

$$\mathfrak{S}_w(x) = \partial_i \mathfrak{S}_{ws_i} \quad \text{and} \quad \mathfrak{S}_{w'}(x) = \partial_i \mathfrak{S}_{w's_i},$$

implying that both $\mathfrak{S}_w(x)$ and $\mathfrak{S}_{w'}(x)$ are symmetric in x_i and x_{i+1} . Since $\mu_i < \mu_{i+1}$, both $x^\mu \mathfrak{S}_w(x^{-1})$ and $x^\mu \mathfrak{S}_{w'}(x^{-1})$ could not be symmetric in x_i and x_{i+1} , which clearly contradicts (2.3) and (2.4). This completes the proof. \blacksquare

3 Proof of Theorem 1.2

As mentioned in Introduction, the main difficulty is to prove the necessity of Theorem 1.2, that is, if $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then w must avoid the patterns 132 and 312.

3.1 Permutations avoiding 132 and 312

In this subsection, we prove a relationship concerning permutations avoiding the patterns 132 and 312, see Proposition 3.1. This proposition will be used in the proof of Lemma 3.4, which is crucial to prove the necessity of Theorem 1.2.

Given a permutation $w = w_1 w_2 \cdots w_n \in S_n$, we say that w is 132-avoiding if there do not exist $i < j < k$ such that $w_i < w_k < w_j$, and w is 312-avoiding if there do not exist $i < j < k$ such that $w_j < w_k < w_i$. For $1 \leq i \leq n$, define

$$l_i(w) = |\{j : j < i, w_j < w_i\}| \quad \text{and} \quad r_i(w) = |\{j : j < i, w_j > w_i\}|. \quad (3.1)$$

Clearly, $l_i(w) + r_i(w) = i - 1$. Let

$$L(w) = (l_1(w), \dots, l_n(w)) \quad \text{and} \quad R(w) = (r_1(w), \dots, r_n(w)). \quad (3.2)$$

We say that a vector $v = (v_1, \dots, v_n)$ is a rearrangement of a vector $v' = (v'_1, \dots, v'_n)$ if there exists a permutation $\pi \in S_n$ such that $v = (v'_{\pi_1}, \dots, v'_{\pi_n})$.

Proposition 3.1. *Let $w \in S_n$ be a permutation avoiding 312, and $u \in S_n$ be a permutation avoiding 132. If $L(w)$ is a rearrangement of $L(u)$ and $R(w)$ is a rearrangement of $R(u)$, then we have $w = u$.*

To prove Proposition 3.1, we need the matrix representation of w . Consider an $n \times n$ box grid, where the rows (respectively, the columns) are numbered $1, 2, \dots, n$ from top to bottom (respectively, from left to right). A box in row i and column j is denoted (i, j) . The matrix representation of w is obtained by putting a dot in the box (i, w_i) for $1 \leq i \leq n$. Figure 3.4 is the matrix representation of $w = 426315$. Obviously, for




Figure 3.4: The matrix representation of $w = 426315$.

$1 \leq i \leq n$, the value $l_i(w)$ (respectively, $r_i(w)$) is equal to the number of dots lying in the region to the strictly upper left (respectively, strictly upper right) of the box (i, w_i) . For $w = 426315$, it can be seen from Figure 3.4 that $L(w) = (0, 0, 2, 1, 0, 4)$ and $R(w) = (0, 1, 0, 2, 4, 1)$.

Proof of Proposition 3.1. We first prove that either $w_n = u_n = n$ or $w_n = u_n = 1$. Since w is 312-avoiding, we see that for $1 \leq j, k \leq n - 1$, if $w_j < w_n$ and $w_k > w_n$, then we must have $j < k$. This implies that the matrix representation of w is as illustrated in Figure 3.5, where the region L contains all the $l_n(w)$ dots that are on the upper left of (n, w_n) , and the region R contains all the $r_n(w) = n - 1 - l_n(w)$ dots that are on the upper right of (n, w_n) . We have the following observation:

(O1). For $1 \leq j \leq l_n(w)$, we have $l_j(w) < l_n(w)$, and for $l_n(w) < j \leq n$, we have $l_j(w) \geq l_n(w)$. Therefore, there are exactly

$$n - l_n(w) = r_n(w) + 1$$

indices j such that $l_j(w) \geq l_n(w)$.

Since $l_n(w) + r_n(w) = n - 1$, we have the following two cases: $l_n(w) \geq \frac{n-1}{2}$ and $r_n(w) > \frac{n-1}{2}$.




Figure 3.5: The matrix representation of w .

Case 1. $l_n(w) \geq \frac{n-1}{2}$. Since $L(w)$ is a rearrangement of $L(u)$, there exists some $1 \leq i \leq n$ such that $l_i(u) = l_n(w)$, that is, in the matrix representation of u , there are $l_n(w)$ dots lying in the region to the upper left of the box (i, u_i) . Since u is 132-avoiding, we see that for $1 \leq j, k < i$, if $u_j < u_i$ and $u_k > u_i$, then $j > k$. So the matrix representation of u is as illustrated in Figure 3.6, where the box marked with a dot is (i, u_i) , the region L' contains $l_i(u) = l_n(w)$ dots, and the region A contains $r_i(u)$ dots.




Figure 3.6: The matrix representation of u in Case 1.

We next show that there are no dots in the regions A, B, C of u in Figure 3.6. To proceed, we list the following straightforward observations.

- (O2). Since there are $l_n(w) \geq \frac{n-1}{2}$ dots in L' , there are at most $\frac{n-1}{2}$ dots in A . So, if (j, u_j) is a box in A , then $l_j(u) < l_n(w)$.
- (O3). Since u is 132-avoiding, every dot (if any) in B lies to the left of every dot in L' . Again, since there are $l_n(w) \geq \frac{n-1}{2}$ dots in L' , there are at most $\frac{n-1}{2}$ dots in B . So if (j, u_j) is a box in B , then $l_j(u) < l_n(w)$.
- (O4). If (j, u_j) is a box in C , then $l_j(u) \geq l_n(w)$.

By (O2), (O3) and (O4), we see that the set of indices j such that $l_j(u) \geq l_n(w)$ is

$$\{i\} \cup \{k : (k, u_k) \text{ is a box in } C\}.$$

Keep in mind that $L(w)$ is a rearrangement of $L(u)$. By (O1), we have

$$|\{i\} \cup \{k : (k, u_k) \text{ is a box in } C\}| = 1 + r_n(w),$$

and so

$$|\{\text{dots in } C\}| = r_n(w). \tag{3.3}$$

Notice that the total number of dots in A, B and C is equal to $n - 1 - l_n(w) = r_n(w)$. This together with (3.3) forces that there are no dots in A and B .

It remains to show that there are no dots in C . Suppose otherwise there is at least one dot in C . Note that the dot in (i, u_i) appears to the upper left of each dot in C . Since u is 132-avoiding, the dots in C must be listed from upper left to bottom right increasingly. This, along with the fact that there are no dots in A and B , implies that $u_n = n$. So we have $l_n(u) = n - 1$. Since $L(w)$ is a rearrangement of $L(u)$, there exists some index, say k , such that $l_k(w) = n - 1$. By the definition of $l_k(w)$, we must have $k = n$ and $w_n = n$. Recalling that $l_i(u) = l_n(w)$, we have $l_i(u) = l_n(w) = n - 1$, and thus $i = n$. However, from the assumption that there is at least one dot in C , it follows that $i < n$, leading to a contradiction. Hence the region C does not contain any dots.

Because there are no dots in B and C , we have $i = n$. By (3.3), we see that $r_n(w) = 0$, and so we have $w_n = n$. Moreover, since $l_n(u) = l_n(w) = n - 1 - r_n(w) = n - 1$, we see that $u_n = n$. Therefore, we have $w_n = u_n = n$.

Case 2. $r_n(w) > \frac{n-1}{2}$. In this case, from Figure 3.5, we see that $r_n(w)$ is the unique maximum value in $R(w)$. Since $R(u)$ is a rearrangement of $R(w)$, there is a unique i such that $r_i(u) = r_n(w)$. The matrix representation of u is illustrated in Figure 3.7, where the square marked with a dot is (i, u_i) and the region R' contains $r_i(u) = r_n(w)$ dots.

	u_i	
		R'
	D	
i		•
	E	F

Figure 3.7: The matrix representation of u in Case 2.

Since $r_i(u) = r_n(w) > \frac{n-1}{2}$ is the unique maximum value in $R(u)$, we see that there are no dots in the regions D and E . We next show that the region F does not contain any dots either. By the fact that $l_n(w) + r_n(w) = n - 1$, as well as the fact that D and E do not contain dots, it follows that there are $l_n(w)$ dots in F :

$$|\{\text{dots in } F\}| = l_n(w). \quad (3.4)$$

We aim to verify $l_n(w) = 0$. Suppose otherwise that $l_n(w) > 0$. By (O1), there are exactly $r_n(w) + 1$ indices j such that $l_j(w) \geq l_n(w)$. Since $L(u)$ is a rearrangement of $L(w)$, there are exactly $r_n(w) + 1$ indices j in u such that $l_j(u) \geq l_n(w)$. Since there are no dots in D , we have $l_i(u) = 0$. Along with the fact that there are no dots in E , we see that if j is an index such that $l_j(u) \geq l_n(w) > 0$, then the box (j, u_j) lies in either F or R' . Since $r_n(w) + l_n(w) = n - 1$, by the assumption $r_n(w) > \frac{n-1}{2}$, we see that $|\{\text{dots in } R'\}| = r_n(w) > l_n(w)$, and so, for each $1 \leq k \leq l_n(w)$, the box (k, u_k) is in R' . Clearly, for $1 \leq k \leq l_n(w)$, we have $l_k(u) < l_n(w)$. Hence the maximal possible number of indices j such that $l_j(u) \geq l_n(w)$ is

$$|\{\text{dots in } F\}| + |\{\text{dots in } R'\}| - l_n(w),$$

which, together with (3.4), becomes

$$|\{\text{dots in } R'\}| = r_i(u) = r_n(w),$$

contrary to the fact that there are exactly $r_n(w) + 1$ indices j such that $l_j(u) \geq l_n(w)$. This verifies $l_n(w) = 0$, and hence there are no dots in F .

Since $l_n(w) = 0$, we obtain that $w_n = 1$. As there are no dots in E and F , we have $i = n$, and so $r_n(u) = r_n(w) = n - 1$, leading to $u_n = 1$. Hence we have $w_n = u_n = 1$.

Now we have proved that either $w_n = u_n = n$ or $w_n = u_n = 1$. This allows us to finish the proof of the proposition by induction on n . To be more specific, if $w_n = u_n = n$, let $w' \in S_{n-1}$ (respectively, $u' \in S_{n-1}$) be the permutation obtained from w (respectively, u) by ignoring w_n (respectively, u_n), while if $w_n = u_n = 1$, let $w' \in S_{n-1}$ (respectively, $u' \in S_{n-1}$) be the permutation obtained from w (respectively, u) by ignoring w_n (respectively, u_n) and then decreasing each of the remaining elements by 1. It is easily seen that w' and u' satisfy the assumptions in the proposition. So, by induction, we have $w' = u'$, which yields $w = u$. This completes the proof. \blacksquare

3.2 Proof of Theorem 1.2

To present a proof of Theorem 1.2, we still need several lemmas. Recall that $d(w)$ denotes the inversion code of a permutation w . View

$$d: w \mapsto d(w)$$

as a bijection from S_n to the set $\{(v_1, \dots, v_n) : 0 \leq v_i \leq n-i\}$. With this notation, for any vector $v = (v_1, \dots, v_n)$ with $0 \leq v_i \leq n-i$, $d^{-1}(v)$ is the corresponding permutation in S_n . For $w \in S_n$, write $w^c = w_1^c w_2^c \cdots w_n^c$ for the complement of w , that is, $w_i^c = n+1-w_i$ for $1 \leq i \leq n$. It is routine to check that

$$d(w^c) = \delta_n - d(w),$$

and so we have

$$w^c = d^{-1}(\delta_n - d(w)). \quad (3.5)$$

Lemma 3.2. *Let $w \in S_n$ be a permutation. Assume that $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is equal to a Schubert polynomial $\mathfrak{S}_{w^*}(x)$. Then*

$$w^* = (d^{-1}(d^t(w^{-1})))^c, \quad (3.6)$$

which is a permutation in S_n .

Proof. By Proposition 2.2, the smallest term of $\mathfrak{S}_w(x)$ in the reverse lexicographic order is $x^{d^t(w^{-1})}$. Hence the leading term of $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is $x^{\delta_n - d^t(w^{-1})}$. On the other hand, by Proposition 2.1, the leading term of $\mathfrak{S}_{w^*}(x)$ is $x^{d(w^*)}$. Since $x^{\delta_n} \mathfrak{S}_w(x^{-1}) = \mathfrak{S}_{w^*}(x)$, we have

$$d(w^*) = \delta_n - d^t(w^{-1}). \quad (3.7)$$

Write $d(w^{-1}) = (a_1, a_2, \dots, a_n)$ and $d^t(w^{-1}) = (b_1, b_2, \dots, b_n)$. For $1 \leq i \leq n$, since $0 \leq a_i \leq n-i$, we have $0 \leq b_i \leq n-i$. This, together with (3.7), implies that w^* is a permutation in S_n . Moreover, by (3.5) and (3.7), we are led to

$$w^* = d^{-1}(\delta_n - d^t(w^{-1})) = (d^{-1}(d^t(w^{-1})))^c,$$

as desired. ■

The second lemma is a well-known characterization of 132-avoiding permutations, see for example [11, Chapter 1].

Lemma 3.3. *A permutation $w \in S_n$ is a 132-avoiding permutation if and only if its inversion code $d(w)$ is a partition. Moreover, the inverse w^{-1} of w is also a 132-avoiding permutation with $d(w^{-1}) = d^t(w)$.*

Based on Proposition 3.1 and Lemma 3.3, we are ready to prove the final lemma.

Lemma 3.4. *Let w be a permutation in S_n , and let w^* be as defined in (3.6). Then $(w^*)^* = w$ if and only if w avoids the patterns 132 and 312.*

Proof. We first show that if w avoids 132 and 312, then $(w^*)^* = w$. Since w is 132-avoiding, by Lemma 3.3, w^{-1} is also 132-avoiding. By (3.7) and Lemma 3.3, we have

$$d(w^*) = \delta_n - d^t(w^{-1}) = \delta_n - d(w),$$

yielding that $w^* = w^c$ by (3.5). As w is 312-avoiding, w^c is 132-avoiding, and hence w^* is 132-avoiding. Thus we obtain that

$$(w^*)^* = (w^c)^* = (w^c)^c = w,$$

where the second equality follows by applying (3.7) and Lemma 3.3, this time to the 132-avoiding permutation w^c .

Let us proceed to prove the reverse direction. By definition,

$$w = (w^*)^* = (d^{-1}(d^t((w^*)^{-1})))^c.$$

So we have $w^c = d^{-1}(d^t((w^*)^{-1}))$, and thus

$$d(w^c) = d^t((w^*)^{-1}). \tag{3.8}$$

Notice that for any vector v of nonnegative integers, its transpose v^t as defined in (2.2) is a partition. Thus, by (3.8) we see that $d(w^c)$ is a partition, and so it follows from Lemma 3.3 that w^c is 132-avoiding, or equivalently, w is 312-avoiding.

We still need to show that w is 132-avoiding. Since $d^t(w^{-1})$ is a partition, by Lemma 3.3, there exists a 132-avoiding permutation $u \in S_n$ such that

$$d(u) = d^t(w^{-1}). \tag{3.9}$$

So we have

$$w^* = (d^{-1}(d^t(w^{-1})))^c = (d^{-1}(d(u)))^c = u^c. \tag{3.10}$$

We now consider $(u^c)^*$. Since $d^t((u^c)^{-1})$ is a partition, there exists a 132-avoiding permutation $v \in S_n$ such that

$$d(v) = d^t((u^c)^{-1}), \tag{3.11}$$

and so

$$(u^c)^* = (d^{-1}(d^t((u^c)^{-1})))^c = (d^{-1}(d(v)))^c = v^c,$$

which along with (3.10) gives $(w^*)^* = v^c$. By the assumption $(w^*)^* = w$, we obtain that

$$w = v^c. \quad (3.12)$$

Using (3.9), (3.11) and (3.12), we can prove the following two claims.

Claim 1: $R(w)$ is a rearrangement of $R(u)$. Recall that $R(w) = (r_1(w), \dots, r_n(w))$, where $r_i(w)$ is equal to the number of dots lying to the upper right of the box (i, w_i) . Notice that in the inversion code $d(w) = (d_1, \dots, d_n)$, the entry d_i equals the number of dots lying to the lower left of the box (i, w_i) . Moreover, the matrix representation of w^{-1} is the transpose of the matrix representation of w . Therefore, $R(w)$ is a rearrangement of $d(w^{-1})$. For the same reason, $R(u)$ is a rearrangement of $d(u^{-1})$.

On the other hand, since u is 132-avoiding, it follows from Lemma 3.3 that $d(u) = d^t(u^{-1})$. Combined with (3.9), we have

$$d^t(u^{-1}) = d^t(w^{-1}).$$

It is easy to check that for any two vectors $\alpha, \beta \in \mathbb{Z}_{\geq 0}^n$, if $\alpha^t = \beta^t$, then α is a rearrangement of β . So $d(w^{-1})$ is a rearrangement of $d(u^{-1})$. We have explained that $R(w)$ is a rearrangement of $d(w^{-1})$ and $R(u)$ is a rearrangement of $d(u^{-1})$. Hence $R(w)$ is a rearrangement of $R(u)$. This proves Claim 1.

Claim 2: $L(w)$ is a rearrangement of $L(u)$. Since v is 132-avoiding, by Lemma 3.3 we have $d(v) = d^t(v^{-1})$. Moreover, by (3.12) we see that $v^{-1} = (w^c)^{-1}$. Hence,

$$d(v) = d^t(v^{-1}) = d^t((w^c)^{-1}).$$

In view of (3.11), we get

$$d^t((w^c)^{-1}) = d^t((u^c)^{-1}).$$

By the same arguments as in Claim 1, $R(w^c)$ is a rearrangement of $R(u^c)$. Noticing that $L(w) = R(w^c)$ and $L(u) = R(u^c)$, we conclude that $L(w)$ is a rearrangement of $L(u)$. This verifies Claim 2.

Since w is 312-avoiding and u is 132-avoiding, combining Proposition 3.1, Claim 1 and Claim 2, we obtain that $w = u$, and so w is 132-avoiding. This finishes the proof. ■

We are now in the position to provide a proof of Theorem 1.2.

Proof of Theorem 1.2. We first prove the sufficiency. Assume that $w \in S_n$ avoids the patterns 132 and 312. Since w is 132-avoiding, it follows from [9, Chapter IV] that $\mathfrak{S}_w(x) = x^{d(w)}$. So we have

$$x^{\delta_n} \mathfrak{S}_w(x^{-1}) = x^{\delta_n - d(w)}. \quad (3.13)$$

As $\delta_n - d(w)$ is the inversion code of w^c , we see that $x^{\delta_n - d(w)} = x^{d(w^c)}$. Since w is 312-avoiding, w^c is 132-avoiding, and so we have

$$\mathfrak{S}_{w^c}(x) = x^{d(w^c)} = x^{\delta_n - d(w)}.$$

This together with (3.13) yields that $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ is the Schubert polynomial $\mathfrak{S}_{w^c}(x)$.

It remains to prove the necessity. Assume that $x^{\delta_n} \mathfrak{S}_w(x^{-1})$ equals the Schubert polynomial $\mathfrak{S}_{w^*}(x)$. Equivalently, we have $\mathfrak{S}_w(x) = x^{\delta_n} \mathfrak{S}_{w^*}(x^{-1})$, which along with Lemma 3.2 leads to $w = (w^*)^*$. Invoking Lemma 3.4, we conclude that w avoids 132 and 312. This completes the proof. ■

4 Concluding remarks on Conjecture 1.3

The necessity of Conjecture 1.3 predicts that if $x^\mu \mathfrak{S}_w(x^{-1})$ is a Schubert polynomial, then w avoids the patterns 1432, 13254, 14253, 24153, 31524, 361452. In the case when w is the pattern 1432, we explain that $x^\mu \mathfrak{S}_{1432}(x^{-1})$ cannot be a Schubert polynomial. Notice that

$$x^\mu \mathfrak{S}_{1432}(x^{-1}) = \frac{x^\mu}{x_1^2 x_2^2 x_3} (x_1^2 + x_1 x_2 + x_1 x_3 + x_2^2 + x_2 x_3). \quad (4.1)$$

Suppose otherwise that $x^\mu \mathfrak{S}_{1432}(x^{-1})$ is a Schubert polynomial, say, $x^\mu \mathfrak{S}_{1432}(x^{-1}) = \mathfrak{S}_{w'}(x)$ for some permutation $w' \in S_n$. In the reverse lexicographic order, it follows from (4.1) that $\frac{x^\mu}{x_1^2 x_2^2 x_3} x_2 x_3$ is the leading monomial of $\mathfrak{S}_{w'}(x)$. Let $d(w') = (d_1, \dots, d_n)$ be the inversion code of w' . By Proposition 2.1,

$$d_1 = \mu_1 - 2, \quad d_2 = \mu_2 - 1, \quad d_3 = \mu_3, \quad d_4 = \mu_4, \dots, d_n = \mu_n. \quad (4.2)$$

By Theorem 1.1, μ is a partition. So we have

$$d_1 + 2 \geq d_2 + 1 \geq d_3 \geq d_4 \geq \dots \geq d_n. \quad (4.3)$$

On the other hand, we see from (4.1) that $\frac{x^\mu}{x_1^2 x_2^2 x_3} x_1^2$ is the smallest monomial of $\mathfrak{S}_w(x)$ in the reverse lexicographic order. By (4.2),

$$\frac{x^\mu}{x_1^2 x_2^2 x_3} x_1^2 = x_1^{d_1+2} x_2^{d_2-1} x_3^{d_3-1} x_4^{d_4} \dots x_n^{d_n}. \quad (4.4)$$

Denote by D_{bot} the bottom RC-graph of w' , which contains the first d_i boxes in row i . From Section 2, we know that the monomial (4.4) can be generated from D_{bot} by applying ladder moves. To be specific, we need to move two crosses from D_{bot} to the first row: one from the second row and one from the third row. This, in view of (4.3), ensures that $d_2 = d_1 + 1$ and $d_3 = d_2 + 1$. However, in this case we can apply ladders moves to generate a monomial

$$x_1^{d_1+2} x_2^{d_2} x_3^{d_3-2} x_4^{d_4} \dots x_n^{d_n},$$

which is smaller than the monomial in (4.4), see Figure 4.8 for an illustration. Hence,




Figure 4.8: Ladder moves in the case when $d_2 = d_1 + 1$ and $d_3 = d_2 + 1$.

the assumption that $x^\mu \mathfrak{S}_{1432}(x^{-1})$ is a Schubert polynomial is false.

From the above arguments, we see that even in the special case of $w = 1432$, the proof is not trivial. We would like to mention that Fink, Mészáros and St. Dizier [4] showed that if σ is a pattern of w , then $\mathfrak{S}_w(x)$ can be expressed as a monomial times $\mathfrak{S}_\sigma(x)$ (in reindexed variables) plus a polynomial with nonnegative coefficients. It seems that this result might be helpful in the investigation of Conjecture 1.3.

References

- [1] N. Bergeron and S. Billey, RC-graphs and Schubert polynomials, *Experiment. Math.* 2 (1993), 257–269.
- [2] S. Billey, W. Jockusch and R.P. Stanley, Some combinatorial properties of Schubert polynomials, *J. Algebraic Combin.* 2 (1993), 345–374.
- [3] P. Brändén and J. Huh, Lorentzian polynomials, *Ann. Math.* 192 (2020), 821–891.
- [4] A. Fink, K. Mészáros and A. St. Dizier, Zero-one Schubert polynomials, *Math. Z.* 297 (2021), 1023–1042.
- [5] C. Gaetz and Y. Gao, A combinatorial duality between the weak and strong Bruhat orders, *J. Combin. Theory Ser. A* 171 (2020), 105178.
- [6] C. Gaetz and Y. Gao, Padded Schubert polynomials and weighted enumeration of Bruhat chains, *arXiv:1905.00047*.
- [7] J. Huh, J.P. Matherne, K. Mészáros and A. St. Dizier, Logarithmic concavity of Schur and related polynomials, *Trans. Amer. Math. Soc.* 375 (2022), 4411–4427.
- [8] A. Lascoux and M.-P. Schützenberger, Polynômes de Schubert, *C. R. Acad. Sci. Paris Sér. I Math.* 294 (1982), 447–450.
- [9] I.G. Macdonald, Notes on Schubert Polynomials, *Laboratoire de combinatoire et d’informatique mathématique (LACIM), Université du Québec à Montréal, Montréal, 1991*.
- [10] A. Okounkov, Why would multiplicities be log-concave? The orbit method in geometry and physics (Marseille, 2000), *Progr. Math.*, Vol. 213, Birkhäuser Boston, Boston, MA, 2003, pp. 329–347.
- [11] R.P. Stanley, *Enumerative Combinatorics, Vol. 1, Second Edition*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2012.

NEIL J.Y. FAN, DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA. Email address: fan@scu.edu.cn

PETER L. GUO, CENTER FOR COMBINATORICS, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA. Email address: lguo@nankai.edu.cn

NICOLAS Y. LIU, CENTER FOR COMBINATORICS, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA. Email address: yiliu@mail.nankai.edu.cn